

Inflamm-Aging: Why Older Men Are the Most Susceptible to SARS-Cov-2 Complicated Outcomes

Massimiliano Bonafè¹, Francesco Prattichizzo², Angelica Giuliani³, Gianluca Storci¹, Jacopo Sabbatinelli^{3*}, Fabiola Olivieri^{3,4*}

¹ Department of Experimental, Diagnostic and Specialty Medicine, AlmaMater Studiorum, Università di Bologna, Bologna, Italy

² IRCCS MultiMedica, Milano, Italy

³ Department of Clinical and Molecular Sciences, DISCLIMO, Università Politecnica delle Marche, Ancona, Italy

⁴ Center of Clinical Pathology and Innovative Therapy, IRCCS INRCA, Ancona, Italy

* These authors contributed equally to the manuscript

Corresponding authors

Francesco Prattichizzo, PhD

IRCCS MultiMedica, Polo Scientifico e Tecnologico, Via Fantoli 16/15, Milano, Italy

E-mail: francesco.prattichizzo@multimedica.it

Angelica Giuliani, PhD

Department of Clinical and Molecular Sciences, Università Politecnica delle Marche, Via Tronto 10/A, Ancona, Italy

E-mail: angelica.giuliani@staff.univpm.it

ABSTRACT

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection is characterized by a high mortality of elderly men with age-related comorbidities. In most of these patients, uncontrolled local and systemic hyperinflammation induces severe and often lethal outcomes. The aging process is characterized by the gradual development of a chronic subclinical systemic inflammation (inflamm-aging) and by acquired immune system impairment (immune senescence). Here, we advance the hypothesis that some key features of aging contribute to the disproportionate SARS-CoV-2 mortality suffered by elderly men. At least four well-recognized aging-related characteristics that are strongly expressed in older men go some way towards explaining why these patients account for the vast majority of fatalities: i. the presence of subclinical systemic inflammation without overt disease, ii. a blunted acquired immune system and type I interferon response due to the chronic inflammation; iii. the downregulation of ACE2 (SARS-CoV-2 receptor), which triggers inflammation, particularly in patients with age-related comorbid diseases such as type II diabetes; and iv. accelerated biological aging, as measured by epigenetic and senescence markers (*e.g.* telomere shortening) associated to the chronic inflammatory state. Though typical of the aged, especially of elderly men, it is conceivable that these features are also shared by some subsets of the younger population. The high mortality rate of SARS-CoV-2 infection suggests that clarification of the mechanisms of inflamm-aging and immune senescence can help combat not only age-related disorders but also SARS-CoV-2 infection.

Keywords: SARS-CoV-2; COVID-19; IL-6; inflamm-aging; immune senescence; host-directed therapies

Introduction

A novel coronavirus, Severe Acute Respiratory Syndrome Corona Virus 2 (SARS-CoV-2), has been spreading across the world since December 2019. SARS-CoV-2 causes an atypical pneumonia associated with acute respiratory distress syndrome, acute respiratory failure, and other potentially lethal complications (Wu et al., 2020) including a deadly "cytokine storm", *i.e.* a strong increase in the plasma concentration of multiple cytokines (Shimabukuro-Vornhagen et al., 2018). In March 2020 Corona Virus Disease 2019 (COVID-19), the disorder caused by SARS-CoV-2, was declared a pandemic by the World Health Organization (World Health Organization, 2020). Its high infectivity seems to be mostly due to intrinsic characteristics of the virus (van Doremalen et al., 2020) and to the lack of previous exposure of the population to the strain. The elderly and patients with pre-existing comorbidities are bearing the brunt of the high case-fatality rate (CFR) of the disease, which is affecting the frailest groups of the population (Istituto Superiore di Sanità, 2020b). Data released by the Chinese Center for Disease Control and Prevention (Novel Coronavirus Pneumonia Emergency Response Epidemiology Team, 2020) suggest that the overall CFR of COVID-19 in China was 2.3%. In particular, whereas there were no fatalities among patients aged up to 9 years, the CFR of those aged 70-79 years and of those aged ≥ 80 years was 8.0% and 14.8%, respectively. Critically, pre-existing comorbidities were associated with a CFR of 10.5% (cardiovascular disease), 7.3% (diabetes), 6.3 % (chronic respiratory disease), 6.0% (hypertension), and 5.6% (cancer) (Young et al., 2020). Moreover, men were more likely to die (2.8%) than women (1.7%) (Novel Coronavirus Pneumonia Emergency Response Epidemiology Team, 2020). Thus, old age and male gender were among the main risk factors for an adverse outcome (Zhou et al., 2020).

These data are similar to those of Italy, where on March 30th infections were 97,780 (Istituto Superiore di Sanità, 2020a). The median age of the deceased was 80 years (interquartile range, 30-103); only 1.1% of those who died were aged less than 50 years. Notably, 70.9% of fatalities were men, whose mean age was 78 years compared to the 82 years of women; men were also 78.5% of deceased patients aged less than 50 years. About 1.4% of the patients who died in Italy suffered from no pre-existing

condition, whereas 51.2% had three or more age-related diseases (ARDs), such as cardiac ischemia, hypertension, type II diabetes mellitus, and chronic obstructive pulmonary disease. Similar mortality patterns have been described for severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome outbreaks, both of which are due to viral strains of the same family (Wu and McGoogan, 2020). These data indicate that advanced age and male gender are risk factors for an adverse outcome.

Two Chinese studies comparing the extreme patient phenotypes, *i.e.* discharged and deceased individuals, found that the most powerful clinical predictors of mortality COVID-19 were the levels of two markers of heart damage, myoglobin and cardiac troponin, and of three major proinflammatory mediators, high-sensitivity C-reactive protein (CRP), interleukin (IL)-6 (Ruan et al., 2020), and d-dimer (Zhou et al., 2020). In most patients with severe disease the infection was associated with a cytokine storm (Channappanavar and Perlman, 2017; Huang et al., 2020; Zumla et al., 2020). In particular, higher levels of circulating IL-6 have been reported in patients with more severe disease (Chen et al., 2020; Zhou et al., 2020).

Aging is characterized by the gradual development of a chronic subclinical systemic inflammation, which has been designated inflamm-aging (Franceschi et al., 2000), and by acquired immune system impairment, *i.e.* immune senescence (Fulop et al., 2018b). The rate of inflamm-aging is higher in men (Bonafè et al., 2001). IL-6 elevation is typical of aging (Maggio et al., 2006). Persistent IL-6 elevation can promote lung tissue inflammation and injury (Yu et al., 2002) and foster viral replication (Velazquez-Salinas et al., 2019). Targeting IL-6, the “cytokine for gerontologists” (Ershler, 1993), helps attenuate the cytokine storm (Lee et al., 2014). Tocilizumab, a biological drug approved for rheumatoid arthritis, is currently being evaluated for its efficacy against the effects of systemic IL-6 elevation (ClinicalTrial.gov, NCT04317092, NCT04320615, NCT04306705).

The evidence reviewed above suggested to us that the two key features of the aging process – inflamm-aging and immune senescence – and their implications can explain why older men with ARDs are the most prone to the adverse outcomes of SARS-COV2 infection.

I. In older men, accelerated inflamm-aging worsens COVID-19 infection outcomes

Inflamm-aging affects all individuals irrespective of their health status (Franceschi et al., 2000); inflammation is also a key pathogenic mechanism of COVID-19 disease (**Figure 1**).




Figure 1. Overview of the framework where inflamm-aging is the main phenomenon contributing to the high COVID-19 mortality rate seen in male, elderly, and frail patients. In these individuals, acute SARS-CoV-2 infection compounds their chronic, subclinical, aging-related proinflammatory state (inflamm-aging) which, together with immune senescence and the age- and gender-specific distribution of ACE2 in the airway epithelia, could blunt the antiviral response to inflammation. This model could explain the delayed viral clearance and the high rate of adverse outcomes observed in older patients in the late disease stages. The assessment of selected biological and immunological aging markers could be a valuable strategy for COVID-19 patient risk stratification in the earliest disease stage irrespective of their chronological age. Figure based on the classification of COVID-19 disease states proposed by Siddiqi and Mehra and adapted from the accompanying paper (Siddiqi and Mehra, 2020). ACE2, angiotensin converting enzyme 2; cfDNA, cell-free DNA; IFN, interferon.

In the elderly, especially men, IL-6 is chronically upregulated (Bonafè et al., 2001; Franceschi et al., 2000); its elevation also predicts mortality due to SARS-CoV-2 (Ruan et al., 2020; Zhou et al., 2020). The gender bias has also been characterized at the molecular level. The stronger age-dependent activation of the innate proinflammatory pathways demonstrated in men compared to women

(Marquez et al., 2020) is consistent with men's higher rate of inflamm-aging (Bonafè et al., 2001). Inflamm-aging is regarded as a major risk factor for the common ARDs, in line with the mounting evidence that an inflammatory pathogenesis is shared by most common ARDs (Franceschi et al., 2000). The situation is different in centenarians, who are characterized by specific longevity traits and anti-inflammatory markers that delay ARD onset, which seem to protect them against the adverse outcomes of sustained inflammation (Franceschi, 2007; Storci et al., 2019). In older men with ARDs, who have the highest rate of inflamm-aging, the inflammatory response induced by SARS-CoV2 infection seems to favor a poorer outcome.

II. In older men, immune senescence worsens COVID-19 infection outcomes

The elderly population is the most vulnerable to infection with influenza virus (IFV) and varicella-zoster virus (VZV), two major pathogens that are responsible for the most common infectious diseases worldwide (Oh et al., 2019). The high mortality from seasonal flu affecting the elderly, which most countries address with extensive vaccine campaigns (World Health Organization, 2019), is a consequence of immune senescence, which seems to be the result of lifelong immunogenic stimulation of the immune system by exogenous subclinical infections and endogenous agents (Fulop et al., 2018a). Endogenous factors such as misplaced nucleic acids (Franceschi et al., 2017; Storci et al., 2020) induce a shift of T and B lymphocytes from naïve to memory effector cells and impair the ability of the adaptive immune system to fight pathogens (Fulop et al., 2018b); since the impairment includes the proliferative exhaustion of memory cells, it involves both unencountered and previously encountered antigens (Ciabattini et al., 2018). A recent study has found that the depletion of B lymphocyte-driven acquired immunity affects predominantly men (Marquez et al., 2020). Thus, in older men the chronic inflammatory status is coupled with a dramatic depletion of B lymphocyte-driven acquired immunity. Notably, aging also attenuates the upregulation of co-stimulatory molecules critical for T cell priming and reduces antiviral interferon (IFN) production by alveolar macrophages and dendritic cells (DCs) in response to IFV (Kovacs et al., 2017). Consistent with this

finding, the ability of DCs and macrophages to elicit CD8⁺ T cell response and proliferation and to release antiviral cytokines is impaired in elderly individuals (Montgomery, 2017); in parallel, these subjects are characterized by a reduced activity of plasmacytoid DCs, the main sources of type I IFNs, which underpin the antiviral response and provide the first-line sentinels in immune surveillance, also in the lung (Abb et al., 1984; Cervantes-Barragan et al., 2007). An important animal study has found that aged cynomolgus macaques infected with SARS-CoV1 (which was responsible for the SARS-CoV1 epidemic in 2003-2004) developed more severe disease than their young adult counterparts and showed a blunted type I IFN response coupled with an elevated production of proinflammatory factors including IL-6 (Smits et al., 2010). Notably, as in other acquired immune system compartments, the impairment of the plasmacytoid DC type I IFN pathway is more pronounced in men compared to women (Griesbeck et al., 2015). Both responses help clear the virus: the former is aimed at providing a local/contained response, whereas the latter enhances the local and systemic response also through the recruitment of large numbers of leukocytes. The elevated production of proinflammatory factors may ultimately induce local and systemic damage by unleashing a cytokine storm (Tisoncik et al., 2012). In SARS-CoV-1-infected pneumocytes, the antiviral response triggered by plasmacytoid DCs is blunted by proinflammatory cytokines released from pneumocytes (Smits et al., 2010). Notably, the gender-biased association of inflamm-aging and immune-senescence is not simply correlative. Indeed, the two phenomena are recognized as being closely coupled in patients suffering from acute critical conditions such as sepsis and cytokine release syndrome (CRS), which is characterized by simultaneous systemic inflammation and immune paralysis (Bruse et al., 2019). A reciprocal inhibition between antiviral type-I IFN response and inflammation has been demonstrated in viral infections (Aman et al., 1996; Kindler and Thiel, 2014; Menachery et al., 2014; Nagata et al., 2007; Pauli et al., 2008; Reboldi et al., 2014; Wei et al., 2006; Zielecki et al., 2013). Therefore, in older men the fine balance between the antiviral response supported by T lymphocytes, B lymphocytes, and plasmacytoid DCs and inflammation is tilted in favor of inflammation.

III. In older men with age-related diseases, the aging-dependent reduction in ACE2 activity worsens SARS-CoV-2 infection outcomes

Angiotensin-converting enzyme (ACE)2, the main SARS-CoV2 host cell receptor, plays a crucial role in virus entry into the cell, as previously demonstrated in SARS and NL63 human coronaviruses (Mathewson et al., 2008). ACE cleaves angiotensin I to generate angiotensin II, whereas ACE2 is a terminal carboxypeptidase that reduces angiotensin II levels, thus playing important functions in the negative regulation of the renin-angiotensin system (Donoghue et al., 2000). ACE2 is expressed in the lung, oral mucosa (epithelial cells of the tongue), intestine (enterocytes), and endothelial cells (Hamming et al., 2004; Xu et al., 2020). Mounting evidence indicates that it plays a protective anti-inflammatory role on endothelial and lung function. Notably, ACE2 underexpression in the lung is implicated in respiratory diseases associated to severe infection (Kuba et al., 2005). In the mouse lung, attenuation of ACE2 activity in the setting of endotoxin inhalation enhanced neutrophil infiltration, exaggerated lung inflammation-induced injury by activating the des-Arg9 bradykinin (DABK)/bradykinin receptor B1 (BKB1R) axis (Sodhi et al., 2018), and elicited the release of proinflammatory chemokines (CXCL5, C-X-C motif chemokine 1), macrophage inflammatory protein-2 (MIP2), and TNF- α from airway epithelia (Sodhi et al., 2018). Intriguingly, the spike protein of SARS-CoV induces downregulation of ACE2 expression through an increased release of proinflammatory chemokines and cytokines, thus enhancing inflammation and vascular permeability and promoting neutrophil recruitment to the lung (Oh et al., 2019). Accordingly, lung ACE2 underexpression and the consequent impaired inhibition of DABK/BKB1R signaling result in abundant neutrophil infiltration and severe lung inflammation. In mice, *in vivo* injection of SARS-CoV spike protein worsens acute lung failure, which can be attenuated by blocking the renin-angiotensin pathway (Imai et al., 2008). As regards ACE2 expression in endothelial cells, vasculitis has been described during viral infection (Ding et al., 2003). Several studies suggest that ACE2 is downregulated in aging (Jia et al., 2019; Xie et al., 2006), explaining, at least in part, the increased susceptibility to vascular injury and cardiovascular disease affecting the elderly (Jia et al., 2019; Xie

et al., 2006). Interestingly, ACE2 overexpression has been reported in Asian females compared to males as well as in other ethnic groups (Jiawei Chen 2020). In contrast, ACE2 is significantly downregulated in patients with type II diabetes as well as in most aged diabetic men with severe COVID-19 outcomes (Cao et al., 2020).

IV. Older men show accelerated biological aging

Several aspects of inflamm-aging are still being investigated (Franceschi et al., 2018; Storci et al., 2020). The accumulation of senescent cells during aging is one of the most likely contributors to inflamm-aging, due to their acquisition of the senescence-associated secretory phenotype (SASP) (Coppe et al., 2008). Senescence affects most cell types including macrophages, which serve as sentinels against infection (Franceschi et al., 2000). Inflamm-aging has also been designated as macroph-aging, due to the important role played by this cell type in secreting proinflammatory mediators at the local and systemic level (Prattichizzo et al., 2016). Notably, studies of the modulation of circulating monocyte/macrophage polarization during aging suggest different age-related trends for the M2 subset (Costantini et al., 2018).

Macrophages are recruited to the lung by SARS-CoV-2-infected cells expressing ACE2 (Qi et al., 2020). The stronger inflammatory state described in circulating monocytes from men compared to women (Marquez et al., 2020) suggests that senescent-like macrophages compound the inflammation induced by COVID-19 through the production of proinflammatory cytokines like IL-6 (Hall et al., 2017; Liu et al., 2019; Rana et al., 2020), in a combination that in older men is especially lethal.

Telomere shortening, *i.e.* the loss of chromosome ends (Lopez-Otin et al., 2013; Mensà et al., 2019), is a hallmark of aging. Older men, particularly those suffering from ARDs, show reduced telomere length (TL) compared to age-matched women (Barrett and Richardson, 2011). Interestingly, telomeres released outside cells may exert an anti-inflammatory action, while shortened telomeres may favor the production of proinflammatory molecular species (Bonafè et al., 2020; Storci et al., 2018). Telomere shortening is seen in ARDs and is associated to an impaired response to infective

agents, suggesting that leukocyte TL may be a marker of general immune competence (Helby et al., 2017). Telomere research supports the notion that TL may play a role in inflamm-aging, which in older men is commonly accelerated. Other measures of biological aging, such as changes in epigenetic patterns, also exhibit a different expression in older men compared to age-matched women (Barrett and Richardson, 2011; Horvath, 2013). The data reviewed above support the notion that the aging process is associated to inflamm-aging and immune senescence and that biological aging is accelerated in men compared to women.

Potential implications for SARS-CoV-2 host-directed therapies

The mechanisms underpinning the chronic inflammatory state characterizing older men are the subject of intense research. We speculate that these mechanisms provide crucial insights into COVID-19 pathobiology and natural history and that their further investigation can lead to novel therapeutic interventions.

A variety of repurposed antiviral drugs (*e.g.* remdesivir, lopinavir/ritonavir) and anti-inflammatory agents, including chloroquine (ClinicalTrial.gov, NCT04323527, NCT04324463, NCT04303507) and tocilizumab (NCT04317092, NCT04320615, NCT04306705) are being tested to treat COVID-19 infection (FiercePharma, 2020; Walker, 2020). It is conceivable that drugs that modulate inflamm-aging could help treat elderly and/or frail patients with SARS-CoV-2 infection (Justice et al., 2019). In particular, “senolytic” compounds, which selectively promote senescent cell clearance, might help reduce systemic as well as local inflammation in such patients. Pilot trials are already in progress (Hickson et al., 2019; Justice et al., 2019), but a substantial number of potential senolytics with a low toxicity profile have been identified and can safely be explored (Gurau et al., 2018). Another approach could rely on harnessing the anti-inflammatory ability of free telomeric end oligonucleotides (Gursel et al., 2003; Yazar et al., 2020), whose airway administration in aerosol form has been studied in a model of mouse lung injury (Scheiermann and Klinman, 2016). This approach would not be limited to elderly COVID-19 patients.

The rate of biological aging (including inflamm-aging) varies among subjects as a result of interactions among genetic make-up, epigenetic modulation, environment, and random factors and can be traced and measured by biomarkers such as TL, circulating nucleic acids, epigenetic changes involving extracellular vesicles, and the expression of inflammatory cytokines such as IL-6 (Ferrucci et al., 2020; Jylhava et al., 2017; Maggio et al., 2006; Mensà et al., 2020; Mensà et al., 2019; Sabbatinelli et al., 2019; Storci et al., 2019). The ability to measure “real” biological age would help determine a person’s aging rate. Attaining this goal clearly requires more accurate and possibly more numerous biomarkers of biological aging capable of predicting functional capacity at a later age than chronological age. Since the variance of aging phenotypes widens over time, owing to different individual trajectories of aging (Ferrucci et al., 2020; Olivieri et al., 2017), immunological/biological age rather than chronological age appears to be a more sensitive approach also to assess individual susceptibility to the adverse outcomes of COVID-19 infection. Consequently, a panel of biomarkers of immunological/biological aging could help identify subjects at risk of adverse COVID-19 outcomes also among the younger population. Thus, young people whose inflamm-aging signature is not consistent with their chronological age – who may therefore be particularly prone to develop not only ARDs, but also a deadly communicable disease such as COVID-19 infection – would also benefit from aging research.

Acknowledgments

We are grateful to Word Designs for text editing (www.silviamodena.com).

Author Contributions

MB, FP, JS, and FO conceived the idea and wrote the manuscript. AG collected the literature and prepared the figure. AG and GS provided critical advice and reviewed the manuscript. All authors approve the final version of the manuscript.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

This work was supported by Italian Ministry of Health (Ricerca Corrente) grants to IRCCS MultiMedica and by Università Politecnica delle Marche (UNIVPM) grants to FO.

REFERENCES

- Abb, J., Abb, H., Deinhardt, F., 1984. Age-related decline of human interferon alpha and interferon gamma production. *Blut* 48, 285-289.
- Aman, M.J., Tretter, T., Eisenbeis, I., Bug, G., Decker, T., Aulitzky, W.E., Tilg, H., Huber, C., Peschel, C., 1996. Interferon-alpha stimulates production of interleukin-10 in activated CD4+ T cells and monocytes. *Blood* 87, 4731-4736.
- Barrett, E.L., Richardson, D.S., 2011. Sex differences in telomeres and lifespan. *Aging Cell* 10, 913-921.
- Bonafè, M., Olivieri, F., Cavallone, L., Giovagnetti, S., Marchegiani, F., Cardelli, M., Pieri, C., Marra, M., Antonicelli, R., Lisa, R., Rizzo, M.R., Paolisso, G., Monti, D., Franceschi, C., 2001. A gender-dependent genetic predisposition to produce high levels of IL-6 is detrimental for longevity. *European Journal of Immunology* 31, 2357-2361.
- Bonafè, M., Sabbatinelli, J., Olivieri, F., 2020. Exploiting the telomere machinery to put the brakes on inflamm-aging. *Ageing Research Reviews* 59, 101027.
- Bruse, N., Leijte, G.P., Pickkers, P., Kox, M., 2019. New frontiers in precision medicine for sepsis-induced immunoparalysis. *Expert Rev Clin Immunol* 15, 251-263.
- Cao, Y., Li, L., Feng, Z., Wan, S., Huang, P., Sun, X., Wen, F., Huang, X., Ning, G., Wang, W., 2020. Comparative genetic analysis of the novel coronavirus (2019-nCoV/SARS-CoV-2) receptor ACE2 in different populations. *Cell Discov* 6, 11.
- Cervantes-Barragan, L., Zust, R., Weber, F., Spiegel, M., Lang, K.S., Akira, S., Thiel, V., Ludewig, B., 2007. Control of coronavirus infection through plasmacytoid dendritic-cell-derived type I interferon. *Blood* 109, 1131-1137.
- Channappanavar, R., Perlman, S., 2017. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin Immunopathol* 39, 529-539.
- Chen, L., Liu, H.G., Liu, W., Liu, J., Liu, K., Shang, J., Deng, Y., Wei, S., 2020. [Analysis of clinical features of 29 patients with 2019 novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi* 43, 203-208.
- Ciabattini, A., Nardini, C., Santoro, F., Garagnani, P., Franceschi, C., Medagliani, D., 2018. Vaccination in the elderly: The challenge of immune changes with aging. *Semin Immunol* 40, 83-94.
- Coppe, J.P., Patil, C.K., Rodier, F., Sun, Y., Munoz, D.P., Goldstein, J., Nelson, P.S., Desprez, P.Y., Campisi, J., 2008. Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol* 6, 2853-2868.
- Costantini, A., Viola, N., Berretta, A., Galeazzi, R., Maccacchione, G., Sabbatinelli, J., Storci, G., De Matteis, S., Butini, L., Rippo, M.R., Procopio, A.D., Caraceni, D., Antonicelli, R., Olivieri, F., Bonafè, M., 2018. Age-related M1/M2 phenotype changes in circulating monocytes from healthy/unhealthy individuals. *Aging (Albany NY)* 10, 1268-1280.
- Ding, Y., Wang, H., Shen, H., Li, Z., Geng, J., Han, H., Cai, J., Li, X., Kang, W., Weng, D., Lu, Y., Wu, D., He, L., Yao, K., 2003. The clinical pathology of severe acute respiratory syndrome (SARS): a report from China. *J Pathol* 200, 282-289.
- Donoghue, M., Hsieh, F., Baronas, E., Godbout, K., Gosselin, M., Stagliano, N., Donovan, M., Woolf, B., Robison, K., Jeyaseelan, R., Breitbart, R.E., Acton, S., 2000. A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. *Circ Res* 87, E1-9.

- Ershler, W.B., 1993. Interleukin-6: a cytokine for gerontologists. *J Am Geriatr Soc* 41, 176-181.
- Ferrucci, L., Gonzalez-Freire, M., Fabbri, E., Simonsick, E., Tanaka, T., Moore, Z., Salimi, S., Sierra, F., de Cabo, R., 2020. Measuring biological aging in humans: A quest. *Aging Cell* 19, e13080.
- FiercePharma, 2020. China turns Roche arthritis drug Actemra against COVID-19 in new treatment guidelines.
- Franceschi, C., 2007. Inflammaging as a major characteristic of old people: can it be prevented or cured? *Nutr Rev* 65, S173-176.
- Franceschi, C., Bonafe, M., Valensin, S., Olivieri, F., De Luca, M., Ottaviani, E., De Benedictis, G., 2000. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* 908, 244-254.
- Franceschi, C., Garagnani, P., Vitale, G., Capri, M., Salvioli, S., 2017. Inflammaging and 'Garb-aging'. *Trends Endocrinol Metab* 28, 199-212.
- Franceschi, C., Zaikin, A., Gordleeva, S., Ivanchenko, M., Bonifazi, F., Storci, G., Bonafe, M., 2018. Inflammaging 2018: An update and a model. *Semin Immunol* 40, 1-5.
- Fulop, T., Larbi, A., Dupuis, G., Le Page, A., Frost, E.H., Cohen, A.A., Witkowski, J.M., Franceschi, C., 2018a. Immunosenescence and Inflamm-Aging As Two Sides of the Same Coin: Friends or Foes? *Frontiers in immunology* 8, 1960-1960.
- Fulop, T., Witkowski, J.M., Olivieri, F., Larbi, A., 2018b. The integration of inflammaging in age-related diseases. *Semin Immunol* 40, 17-35.
- Griesbeck, M., Ziegler, S., Laffont, S., Smith, N., Chauveau, L., Tomezsko, P., Sharei, A., Kourjian, G., Porichis, F., Hart, M., Palmer, C.D., Sirignano, M., Beisel, C., Hildebrandt, H., Cenac, C., Villani, A.C., Diefenbach, T.J., Le Gall, S., Schwartz, O., Herbeuval, J.P., Autran, B., Guery, J.C., Chang, J.J., Altfeld, M., 2015. Sex Differences in Plasmacytoid Dendritic Cell Levels of IRF5 Drive Higher IFN-alpha Production in Women. *J Immunol* 195, 5327-5336.
- Gurau, F., Baldoni, S., Prattichizzo, F., Espinosa, E., Amenta, F., Procopio, A.D., Albertini, M.C., Bonafe, M., Olivieri, F., 2018. Anti-senescence compounds: A potential nutraceutical approach to healthy aging. *Ageing Res Rev* 46, 14-31.
- Gursel, I., Gursel, M., Yamada, H., Ishii, K.J., Takeshita, F., Klinman, D.M., 2003. Repetitive elements in mammalian telomeres suppress bacterial DNA-induced immune activation. *J Immunol* 171, 1393-1400.
- Hall, B.M., Balan, V., Gleiberman, A.S., Strom, E., Krasnov, P., Virtuoso, L.P., Rydkina, E., Vujcic, S., Balan, K., Gitlin, II, Leonova, K.I., Consiglio, C.R., Gollnick, S.O., Chernova, O.B., Gudkov, A.V., 2017. p16(Ink4a) and senescence-associated beta-galactosidase can be induced in macrophages as part of a reversible response to physiological stimuli. *Aging (Albany NY)* 9, 1867-1884.
- Hamming, I., Timens, W., Bulthuis, M.L., Lely, A.T., Navis, G., van Goor, H., 2004. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol* 203, 631-637.
- Helby, J., Nordestgaard, B.G., Benfield, T., Bojesen, S.E., 2017. Shorter leukocyte telomere length is associated with higher risk of infections: a prospective study of 75,309 individuals from the general population. *Haematologica* 102, 1457-1465.
- Hickson, L.J., Langhi Prata, L.G.P., Bobart, S.A., Evans, T.K., Giorgadze, N., Hashmi, S.K., Herrmann, S.M., Jensen, M.D., Jia, Q., Jordan, K.L., Kellogg, T.A., Khosla, S., Koerber, D.M., Lagnado, A.B.,

- Lawson, D.K., LeBrasseur, N.K., Lerman, L.O., McDonald, K.M., McKenzie, T.J., Passos, J.F., Pignolo, R.J., Pirtskhalava, T., Saadiq, I.M., Schaefer, K.K., Textor, S.C., Victorelli, S.G., Volkman, T.L., Xue, A., Wentworth, M.A., Wissler Gerdes, E.O., Zhu, Y., Tchkonina, T., Kirkland, J.L., 2019. Senolytics decrease senescent cells in humans: Preliminary report from a clinical trial of Dasatinib plus Quercetin in individuals with diabetic kidney disease. *EBioMedicine* 47, 446-456.
- Horvath, S., 2013. DNA methylation age of human tissues and cell types. *Genome Biol* 14, R115.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., Xiao, Y., Gao, H., Guo, L., Xie, J., Wang, G., Jiang, R., Gao, Z., Jin, Q., Wang, J., Cao, B., 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497-506.
- Imai, Y., Kuba, K., Penninger, J.M., 2008. The discovery of angiotensin-converting enzyme 2 and its role in acute lung injury in mice. *Exp Physiol* 93, 543-548.
- Istituto Superiore di Sanità, 2020a. Characteristics of COVID-19 patients dying in Italy Report based on available data on March 24th, 2020.
- Istituto Superiore di Sanità, 2020b. Sorveglianza integrata COVID-19: i principali dati nazionali.
- Jia, G., Aroor, A.R., Jia, C., Sowers, J.R., 2019. Endothelial cell senescence in aging-related vascular dysfunction. *Biochim Biophys Acta Mol Basis Dis* 1865, 1802-1809.
- Jiawei Chen , Q.J., Xian Xia , Kangping Liu , Zhengqing Yu , Wanyu Tao , Wenxuan Gong , Jing-Dong J. Han, 2020. Individual Variation of the SARS-CoV2 Receptor ACE2 Gene Expression and Regulation. Preprints, 2020030191.
- Justice, J.N., Nambiar, A.M., Tchkonina, T., LeBrasseur, N.K., Pascual, R., Hashmi, S.K., Prata, L., Masternak, M.M., Kritchevsky, S.B., Musi, N., Kirkland, J.L., 2019. Senolytics in idiopathic pulmonary fibrosis: Results from a first-in-human, open-label, pilot study. *EBioMedicine* 40, 554-563.
- Jylhava, J., Pedersen, N.L., Hagg, S., 2017. Biological Age Predictors. *EBioMedicine* 21, 29-36.
- Kindler, E., Thiel, V., 2014. To sense or not to sense viral RNA--essentials of coronavirus innate immune evasion. *Curr Opin Microbiol* 20, 69-75.
- Kovacs, E.J., Boe, D.M., Boule, L.A., Curtis, B.J., 2017. Inflammaging and the Lung. *Clin Geriatr Med* 33, 459-471.
- Kuba, K., Imai, Y., Rao, S., Gao, H., Guo, F., Guan, B., Huan, Y., Yang, P., Zhang, Y., Deng, W., Bao, L., Zhang, B., Liu, G., Wang, Z., Chappell, M., Liu, Y., Zheng, D., Leibbrandt, A., Wada, T., Slutsky, A.S., Liu, D., Qin, C., Jiang, C., Penninger, J.M., 2005. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat Med* 11, 875-879.
- Lee, D.W., Gardner, R., Porter, D.L., Louis, C.U., Ahmed, N., Jensen, M., Grupp, S.A., Mackall, C.L., 2014. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood* 124, 188-195.
- Liu, J.Y., Souroullas, G.P., Diekman, B.O., Krishnamurthy, J., Hall, B.M., Sorrentino, J.A., Parker, J.S., Sessions, G.A., Gudkov, A.V., Sharpless, N.E., 2019. Cells exhibiting strong p16 (INK4a) promoter activation in vivo display features of senescence. *Proc Natl Acad Sci U S A* 116, 2603-2611.
- Lopez-Otin, C., Blasco, M.A., Partridge, L., Serrano, M., Kroemer, G., 2013. The hallmarks of aging. *Cell* 153, 1194-1217.

- Maggio, M., Guralnik, J.M., Longo, D.L., Ferrucci, L., 2006. Interleukin-6 in aging and chronic disease: a magnificent pathway. *J Gerontol A Biol Sci Med Sci* 61, 575-584.
- Marquez, E.J., Chung, C.H., Marches, R., Rossi, R.J., Nehar-Belaid, D., Eroglu, A., Mellert, D.J., Kuchel, G.A., Banchereau, J., Ucar, D., 2020. Sexual-dimorphism in human immune system aging. *Nat Commun* 11, 751.
- Mathewson, A.C., Bishop, A., Yao, Y., Kemp, F., Ren, J., Chen, H., Xu, X., Berkhout, B., van der Hoek, L., Jones, I.M., 2008. Interaction of severe acute respiratory syndrome-coronavirus and NL63 coronavirus spike proteins with angiotensin converting enzyme-2. *J Gen Virol* 89, 2741-2745.
- Menachery, V.D., Eisfeld, A.J., Schafer, A., Josset, L., Sims, A.C., Proll, S., Fan, S., Li, C., Neumann, G., Tilton, S.C., Chang, J., Gralinski, L.E., Long, C., Green, R., Williams, C.M., Weiss, J., Matzke, M.M., Webb-Robertson, B.J., Schepmoes, A.A., Shukla, A.K., Metz, T.O., Smith, R.D., Waters, K.M., Katze, M.G., Kawaoka, Y., Baric, R.S., 2014. Pathogenic influenza viruses and coronaviruses utilize similar and contrasting approaches to control interferon-stimulated gene responses. *mBio* 5, e01174-01114.
- Mensà, E., Guescini, M., Giuliani, A., Bacalini, M.G., Ramini, D., Corleone, G., Ferracin, M., Fulgenzi, G., Graciotti, L., Prattichizzo, F., Sorci, L., Battistelli, M., Monsurrò, V., Bonfigli, A.R., Cardelli, M., Recchioni, R., Marcheselli, F., Latini, S., Maggio, S., Fanelli, M., Amatori, S., Storci, G., Ceriello, A., Stocchi, V., De Luca, M., Magnani, L., Rippo, M.R., Procopio, A.D., Sala, C., Budimir, I., Bassi, C., Negrini, M., Garagnani, P., Franceschi, C., Sabbatinelli, J., Bonafè, M., Olivieri, F., 2020. Small extracellular vesicles deliver miR-21 and miR-217 as pro-senescence effectors to endothelial cells. *Journal of Extracellular Vesicles* 9, 1725285.
- Mensà, E., Latini, S., Ramini, D., Storci, G., Bonafè, M., Olivieri, F., 2019. The telomere world and aging: Analytical challenges and future perspectives. *Ageing Research Reviews* 50, 27-42.
- Montgomery, R.R., 2017. Age-related alterations in immune responses to West Nile virus infection. *Clin Exp Immunol* 187, 26-34.
- Nagata, N., Iwata, N., Hasegawa, H., Fukushi, S., Yokoyama, M., Harashima, A., Sato, Y., Saijo, M., Morikawa, S., Sata, T., 2007. Participation of both host and virus factors in induction of severe acute respiratory syndrome (SARS) in F344 rats infected with SARS coronavirus. *J Virol* 81, 1848-1857.
- Novel Coronavirus Pneumonia Emergency Response Epidemiology Team, 2020. Vital surveillances: the epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19)—China, 2020.
- Oh, S.J., Lee, J.K., Shin, O.S., 2019. Aging and the Immune System: the Impact of Immunosenescence on Viral Infection, Immunity and Vaccine Immunogenicity. *Immune Netw* 19, e37.
- Olivieri, F., Capri, M., Bonafe, M., Morsiani, C., Jung, H.J., Spazzafumo, L., Vina, J., Suh, Y., 2017. Circulating miRNAs and miRNA shuttles as biomarkers: Perspective trajectories of healthy and unhealthy aging. *Mech Ageing Dev* 165, 162-170.
- Pauli, E.K., Schmolke, M., Wolff, T., Viemann, D., Roth, J., Bode, J.G., Ludwig, S., 2008. Influenza A virus inhibits type I IFN signaling via NF-kappaB-dependent induction of SOCS-3 expression. *PLoS Pathog* 4, e1000196.
- Prattichizzo, F., Bonafè, M., Olivieri, F., Franceschi, C., 2016. Senescence associated macrophages and "macroph-aging": Are they pieces of the same puzzle? *Aging* 8, 3159-3160.
- Qi, F., Qian, S., Zhang, S., Zhang, Z., 2020. Single cell RNA sequencing of 13 human tissues identify cell types and receptors of human coronaviruses. *Biochem Biophys Res Commun*.

- Rana, T., Jiang, C., Liu, G., Miyata, T., Antony, V., Thannickal, V.J., Liu, R.M., 2020. PAI-1 Regulation of TGF-beta1-induced Alveolar Type II Cell Senescence, SASP Secretion, and SASP-mediated Activation of Alveolar Macrophages. *Am J Respir Cell Mol Biol* 62, 319-330.
- Reboldi, A., Dang, E.V., McDonald, J.G., Liang, G., Russell, D.W., Cyster, J.G., 2014. Inflammation. 25-Hydroxycholesterol suppresses interleukin-1-driven inflammation downstream of type I interferon. *Science* 345, 679-684.
- Ruan, Q., Yang, K., Wang, W., Jiang, L., Song, J., 2020. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med*.
- Sabbatinelli, J., Prattichizzo, F., Olivieri, F., Procopio, A.D., Rippo, M.R., Giuliani, A., 2019. Where Metabolism Meets Senescence: Focus on Endothelial Cells. *Front Physiol* 10, 1523.
- Scheiermann, J., Klinman, D.M., 2016. Suppressive oligonucleotides inhibit inflammation in a murine model of mechanical ventilator induced lung injury. *J Thorac Dis* 8, 2434-2443.
- Shimabukuro-Vornhagen, A., Godel, P., Subklewe, M., Stemmler, H.J., Schlosser, H.A., Schlaak, M., Kochanek, M., Boll, B., von Bergwelt-Baildon, M.S., 2018. Cytokine release syndrome. *J Immunother Cancer* 6, 56.
- Siddiqi, H.K., Mehra, M.R., 2020. COVID-19 Illness in Native and Immunosuppressed States: A Clinical-Therapeutic Staging Proposal. *The Journal of Heart and Lung Transplantation*.
- Smits, S.L., de Lang, A., van den Brand, J.M., Leijten, L.M., van, I.W.F., Eijkemans, M.J., van Amerongen, G., Kuiken, T., Andeweg, A.C., Osterhaus, A.D., Haagmans, B.L., 2010. Exacerbated innate host response to SARS-CoV in aged non-human primates. *PLoS Pathog* 6, e1000756.
- Sodhi, C.P., Wohlford-Lenane, C., Yamaguchi, Y., Prindle, T., Fulton, W.B., Wang, S., McCray, P.B., Jr., Chappell, M., Hackam, D.J., Jia, H., 2018. Attenuation of pulmonary ACE2 activity impairs inactivation of des-Arg(9) bradykinin/BKB1R axis and facilitates LPS-induced neutrophil infiltration. *Am J Physiol Lung Cell Mol Physiol* 314, L17-L31.
- Storci, G., Bacalini, M.G., Bonifazi, F., Garagnani, P., De Carolis, S., Salvioli, S., Olivieri, F., Bonafe, M., 2020. Ribosomal DNA instability: An evolutionary conserved fuel for inflammaging. *Ageing Res Rev* 58, 101018.
- Storci, G., De Carolis, S., Olivieri, F., Bonafe, M., 2018. Changes in the biochemical taste of cytoplasmic and cell-free DNA are major fuels for inflammaging. *Semin Immunol* 40, 6-16.
- Storci, G., De Carolis, S., Papi, A., Bacalini, M.G., Gensous, N., Marasco, E., Tesei, A., Fabbri, F., Arienti, C., Zanoni, M., Sarnelli, A., Santi, S., Olivieri, F., Mensa, E., Latini, S., Ferracin, M., Salvioli, S., Garagnani, P., Franceschi, C., Bonafe, M., 2019. Genomic stability, anti-inflammatory phenotype, and up-regulation of the RNaseH2 in cells from centenarians. *Cell Death Differ* 26, 1845-1858.
- Tisoncik, J.R., Korth, M.J., Simmons, C.P., Farrar, J., Martin, T.R., Katze, M.G., 2012. Into the eye of the cytokine storm. *Microbiol Mol Biol Rev* 76, 16-32.
- van Doremalen, N., Bushmaker, T., Morris, D.H., Holbrook, M.G., Gamble, A., Williamson, B.N., Tamin, A., Harcourt, J.L., Thornburg, N.J., Gerber, S.I., Lloyd-Smith, J.O., de Wit, E., Munster, V.J., 2020. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*.
- Velazquez-Salinas, L., Verdugo-Rodriguez, A., Rodriguez, L.L., Borca, M.V., 2019. The Role of Interleukin 6 During Viral Infections. *Front Microbiol* 10, 1057.
- Walker, J., 2020. Regeneron, Sanofi to Test Arthritis Drug as Coronavirus Treatment, *The Wall Street Journal*.

- Wei, L., Sandbulte, M.R., Thomas, P.G., Webby, R.J., Homayouni, R., Pfeffer, L.M., 2006. NFkappaB negatively regulates interferon-induced gene expression and anti-influenza activity. *J Biol Chem* 281, 11678-11684.
- World Health Organization, 2019. Recommendations on influenza vaccination during the 2019–2020 winter season.
- World Health Organization, 2020. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020.
- Wu, Y., Ho, W., Huang, Y., Jin, D.Y., Li, S., Liu, S.L., Liu, X., Qiu, J., Sang, Y., Wang, Q., Yuen, K.Y., Zheng, Z.M., 2020. SARS-CoV-2 is an appropriate name for the new coronavirus. *Lancet* 395, 949-950.
- Wu, Z., McGoogan, J.M., 2020. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*.
- Xie, X., Chen, J., Wang, X., Zhang, F., Liu, Y., 2006. Age- and gender-related difference of ACE2 expression in rat lung. *Life Sci* 78, 2166-2171.
- Xu, H., Zhong, L., Deng, J., Peng, J., Dan, H., Zeng, X., Li, T., Chen, Q., 2020. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci* 12, 8.
- Yazar, V., Kilic, G., Bulut, O., Canavar Yildirim, T., Yagci, F.C., Aykut, G., Klinman, D.M., Gursel, M., Gursel, I., 2020. A suppressive oligodeoxynucleotide expressing TTAGGG motifs modulates cellular energetics through the mTOR signaling pathway. *Int Immunol* 32, 39-48.
- Young, B.E., Ong, S.W.X., Kalimuddin, S., Low, J.G., Tan, S.Y., Loh, J., Ng, O.T., Marimuthu, K., Ang, L.W., Mak, T.M., Lau, S.K., Anderson, D.E., Chan, K.S., Tan, T.Y., Ng, T.Y., Cui, L., Said, Z., Kurupatham, L., Chen, M.I., Chan, M., Vasoo, S., Wang, L.F., Tan, B.H., Lin, R.T.P., Lee, V.J.M., Leo, Y.S., Lye, D.C., Singapore Novel Coronavirus Outbreak Research, T., 2020. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA*.
- Yu, M., Zheng, X., Witschi, H., Pinkerton, K.E., 2002. The role of interleukin-6 in pulmonary inflammation and injury induced by exposure to environmental air pollutants. *Toxicol Sci* 68, 488-497.
- Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., Xiang, J., Wang, Y., Song, B., Gu, X., Guan, L., Wei, Y., Li, H., Wu, X., Xu, J., Tu, S., Zhang, Y., Chen, H., Cao, B., 2020. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*.
- Zielecki, F., Weber, M., Eickmann, M., Spiegelberg, L., Zaki, A.M., Matrosovich, M., Becker, S., Weber, F., 2013. Human cell tropism and innate immune system interactions of human respiratory coronavirus EMC compared to those of severe acute respiratory syndrome coronavirus. *J Virol* 87, 5300-5304.
- Zumla, A., Hui, D.S., Azhar, E.I., Memish, Z.A., Maeurer, M., 2020. Reducing mortality from 2019-nCoV: host-directed therapies should be an option. *Lancet* 395, e35-e36.