

Gravity as the Global Standard Model

Ahmed Farag Ali

Dept. of Physics, Faculty of Sciences, Benha University, Benha 13518, Egypt.

ahmed.ali@fsc.bu.edu.eg

Abstract

We study the localization of gravity through the matching point between non-inertial frames and local inertial frames. This localization of gravity lead to an emergence of a timeless state of the universe in a mathematically consistent way. We find a geometric interpretation of the speed of light and mass. The experimental evidence of the timeless state of the universe is the quantum entanglement and internal symmetries that are independent of time. Since the spin measurement is the manifestation of quantum entanglement measurement. Therefore, the spin of quantum particles is correlated with the relative gravitational red-shift at two different points. The same can be applied to all types of internal symmetries that are independent of time. Therefore gravity represents all measurements independent of time including quantum entanglement. We conclude that the gravity is the global $SU(3) \times SU(2) \times U(1)$ symmetry that produces gauge fields such as Electromagnetism, weak and strong nuclear force through localization with their internal symmetries correlated with the varying of relative gravitational red-shift. We also introduce a gravitational or geometric interpretation of spin-0, spin-1 and spin-1/2 states. We answered the question why do we measure matter and not anti-matter. We Introduce a solution for the Cosmological Constant Problem Value.

1 Localization of gravity

We assume an existence of Schwarzschild black hole with event horizon. We investigate the gravitational red-shift which is a property of general covariance. To localize gravity, we consider two points (A and R) in the gravitational field of black hole as shown in the following Fig. (1). Notice here these two points form a triangle that follow a geodesics geometry of the considered black hole when connecting the two points with black hole center. If R and A are far enough from K , the triangle become approximately Euclidean triangle. Between these two points A and R , there are two possible local gravitational measurements as follows:

1. Relative gravitational red-shift which is represented by the ratio at two different points

$$\frac{z_A}{z_R} = \frac{(1 - \frac{r_s}{r_A})^{-1/2} - 1}{(1 - \frac{r_s}{r_R})^{-1/2} - 1} \quad (1.1)$$




Figure 1: Black hole

2. The difference in gravitational red-shift at two different points.

$$\Delta z = z_A - z_R = \left(1 - \frac{r_s}{r_A}\right)^{-1/2} - \left(1 - \frac{r_s}{r_R}\right)^{-1/2} \quad (1.2)$$

2 Matching point between local gravity measurement and local Inertial frames

2.1 Relative gravitational red-shift

At the moment of change of inertial frames into non-inertial frames, it is mathematically consistent to match the local gravitational measurements with the measurement in local inertial frames. Therefore, we consider the weak gravitational approximations, $r_s \ll r_K$ and $r_s \ll r_R$. The gravitational red-shift for both A and R can be approximated as follows

$$z_A = \left(1 - \frac{r_s}{r_A}\right)^{-1/2} - 1 \approx \frac{r_s}{2r_A} \quad (2.1)$$

$$z_R = \left(1 - \frac{r_s}{r_R}\right)^{-1/2} - 1 \approx \frac{r_s}{2r_R}. \quad (2.2)$$

We compute the relative gravitational red-shift using Eq(1.1). We express it in terms of all lengths measured at R including the distance between A and R (r_{AR}).

$$\frac{z_R}{z_A} = \frac{1}{\sqrt{1 - \frac{r_{AR}^2}{r_A^2} + 2\frac{r_R r_{AR}}{r_A^2} \cos \alpha}} = \delta \quad (2.3)$$

Notice the value of α can be $0 \leq \alpha \leq \pi/2$. This equation represents the relative gravitational-red-shift between two points A and R in a weak gravitational field. For the case $\alpha = \pi/2$. The relative gravitational red-shift will be given by

$$\frac{z_R}{z_A} = \frac{1}{\sqrt{1 - \left(\frac{r_{AR}}{r_A}\right)^2}} = \delta \quad (2.4)$$

On the other side, the measurement in local inertial frames are determined in terms of relative time dilation as follows

$$\frac{t_R}{t_A} = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (2.5)$$

where v is the relative speed between the two points A and R in the local inertial frames and c is the speed of light.

At the moment of changing inertial frames into non-inertial frames which is matching connecting point between them, the relative gravitational red-shift can set to equal to definition of time dilation in special relativity ($1/\sqrt{1 - v^2/c^2}$) which implies that the ratio r_{AR}^2/r_A^2 can be replaced by ratio v^2/c^2 . This matching connecting point is legitimate and mathematically consistent since the relative gravitational red-shift introduces a local measurement which should match with special relativity that holds only in "local" inertial frames. Therefore, for mathematical consistency of general relativity, its local measurement should be equivalent to measurement special relativity that hold only in local inertial frames. This means when $\alpha = \pi/2$, the gamma factor of special relativity emerges as a ratio between the gravitational red-shift at A and R . Local gravity measurements depends only on "**one variable**"; the distance from the gravitational source, which is the reason for velocity ratios turned to be lengths ratios in this delta factor in Eq. (2.4). The ratio r_{AR}^2/r_A^2 can be considered as a geometric or gravitational interpretation of the ratio v^2/c^2 . This comparison can be written as

$$\frac{r_{AR}}{r_A} = \frac{r_{AR}/t}{r_A/t} = \frac{v}{c} \quad (2.6)$$

This would support the approach of time varying speed of light as a solution of cosmological puzzles that was suggested in [1]. It may support also the experimental findings of changing physical constants such as fine structure constant in gravitational field as shown recently in [2]. In our case, the ratio v/c is varying depending on the distance from the gravitational source.

We note that time can be inserted easily in the previous equation as a "redundant variable" which can be interpreted that the state of matching local gravity measurements with local inertial frames would correspond to a possible timeless state which is consistent mathematically through the matching process that we performed. It has been intuited timeless universe is possible as well [3]

To realize the effect of other values of angle α in weak gravitational field, we consider an approximation which is $r_{AR} \ll r_A$, $r_{AR} \ll r_R$. In that case, the delta factor in Eq. (2.3) is approximated as following

$$\delta \approx 1 - \frac{r_R r_{AR}}{r_A^2} \cos \alpha \quad (2.7)$$

It is found that this equation matches with the derivative of Kepler equation.

$$\frac{dm}{dE} = 1 - e \cos E \quad (2.8)$$

where m is the mean anomaly, E is the eccentric anomaly, and e is the eccentricity. In our approximation, the eccentricity e is approximately equal to $r_R r_{AR} / r_A^2$, and E refers to the angle α . This gives a geometric interpretation of Kepler equation from the relative gravitational red-shift.

2.2 Difference in Gravitational red-shift

In this section, we compute local measurement as difference in gravitational red-shift. For weak gravitational approximation, we get

$$\Delta z = z_R - z_A = \frac{r_s}{2r_R} - \frac{r_s}{2r_A} \quad (2.9)$$

Let us make an approximation as following $r_A = r_R + x$, where $x \ll r_A$ and $x \ll r_R$. In that case, Eq. (2.9) will be rewritten as follows. We use the value of Schwarzschild radius $r_s = 2GM/c^2$

$$\Delta z M c^2 = GM^2 \frac{x}{r_R^2} \quad (2.10)$$

where G is the gravitational constant, M is the black hole mass and c is the speed of light. From Eq.(2.6), c can be set to equal to r_A if we take t to be unity since we agree that t is a redundant factor through matching local gravity measurement with local inertial frames. We find that Eq. (2.10) can be arranged to take the following form

$$\Delta z M = \Delta M = GM^2 \frac{x}{r_A^2 r_R^2} \quad (2.11)$$

where $\Delta M = \Delta z M$. ΔM represents relative relation of mass between any two different points in the gravitational field. This would give a geometric representation for relativistic relation mc^2 in terms of the difference between different points to the black hole. We want to understand the physical meaning of the factor GM^2 in r.h.s of Eq. (2.11). When we look at Bekenstein-Hawking entropy equation

$$S_{BH} = \frac{c^3 A}{4G\hbar} = \frac{4\pi}{c\hbar} GM^2 \quad (2.12)$$

where $A = 16\pi(GM/c^2)^2$ stands for surface area of a black hole. We found that the factor GM^2 in r.h.s of Eq. (2.11) between any two different points can be expressed in terms of black hole entropy as follows

$$\Delta z M = \Delta M = \frac{\hbar}{4\pi} \frac{x}{r_R^2 r_A} S_{BH} \quad (2.13)$$

We assumed that time is a unity. let us consider this unit as the Planck time. This means that the Planck constant in previous equation can be replaced through the following process

$$t_p = \sqrt{\frac{\hbar G}{c^5}} = 1 \quad (2.14)$$

Since the Planck time is our unity, then c can set to be r_A . Therefore, the Planck constant in this geometric picture will be given by

$$\hbar G = r_A^5 \quad (2.15)$$

This equation gives a geometric or gravitational interpretation of Planck constant when matching local gravity measurement with local inertial frames. The relative mass between any two different points is therefore given by

$$\Delta zM = \Delta M = \frac{1}{4\pi G} \frac{x r_A^4}{r_R^2} S_{BH} = \frac{1}{16\pi G} \frac{x r_A^2}{r_R^2} A \quad (2.16)$$

We notice that the difference in gravitational red-shift gives an emergence of relative mass. The previous equation gives purely a geometric expression for the relative mass in terms of the gravitational source area of its full entropy. It is experimentally proved that the difference in gravitational potential has an effect on the apparent weight of the 14.4-keV ray of Iron (Fe) [4, 5]. This may be an experimental support for the derived relation that connect the difference in gravitational red-shift and emergence of mass in this section.

3 Experimental quantum entanglement evidence of timeless state

The experimental evidence of the timeless state is the spin measurement in quantum entanglement at two different points in the space [7]. Besides, any internal symmetries of the elementary particles that are measured independent of time. Therefore, it is logically consistent to consider the quantum spin of elementary particles correlated with relative gravitational red-shift between two different points in the space. The same can be applied to all internal symmetries. In that sense, gravity may explain the origin of quantum spin for elementary particles in three different cases as follows:

- The relative gravitational red-shift $z_R/z_A=1$, would introduce the spin 1 excitation. In that case, $r_{AR} = r_A = r_R$ which corresponds to equilateral triangle.
- The measurement between two different points at different distances from the gravitational source would result the spin-1/2 excitation. In Local inertial frames, this ratio is perfectly 1/2, but it is expected to be varying in strong gravity field as we take gravitational red-shift into consideration of computing this ratio.
- The measurement on the same point, would result spin 0 state. This state is represented by a straight line that has an arbitrary length.

We interpret that as the quantum spin/internal symmetries as a non-local completeness of local quantum theory which is related to gravitational source. The spin/internal symmetries in that sense may be the gravity effect or gravitational degrees of freedom on every quantum particle. This may open a door for "gravity technology" through the suggested correlation between the quantum spin/internal symmetries and a relative gravitational red-shift.

4 Gravity and Uncertainty

In previous sections, we have shown that the concept of velocity is replaced with the relative distance between any any two different points in the space time when we match local gravitational measurements with the local inertial frames. This would generate a timeless state that is mathematically consistent in connecting non-inertial frames with local inertial frames or in other words matching non-locality with locality at their connecting point. In that state, the gravitational measurements happens in terms of only one variable which is the distance from the gravitational source. Time variable at the matching process appear to be a redundant variable. Since time, and therefore velocity dissolve when the matching process happens, therefore there is no meaning to define uncertainty in this timeless state. We conclude that the distance from the gravitational source may form the hidden variable of quantum mechanics, which may complete the connection between quantum mechanics and gravity in one unified theory, which is the timeless state. This may complete the picture that was introduced in EPR [6]. This implies that the uncertainty amount would decrease as the measurement happens closer to the gravitational source. The uncertainty emerges, once we start varying the distance from the black hole. The uncertainty emerges due to the difference in information between point A and point R without knowing the distance to the source. This difference is encoded in Eq. 2.16. The difference in information (uncertainty) would be given by

$$\frac{4\pi\Delta Mr_R^2 r_A}{x} = \hbar \ S_{BH} \quad (4.1)$$

Notice that the difference in information between Point A and point R depends on the distances r_A and r_R . If we do not know these values, this difference will be hidden in our local measurements, and therefore uncertainty emerges. Notice that the variables on the left hand side are greater than or equal to the Planck constant. This relation represent the hidden variables which is reason for emergence of uncertainty principle inequality in local measurements.

5 Strong Gravity Case

In strong gravity case, we can use Eqs. 1.1 and 1.2 without any approximation. These relations can be computed for any two points, and it gives a wide spectrum of measurements of relative gravitational red-shifts and masses in strong gravity field. In strong gravity field, the triangle will not be perfectly Euclidean but can be computed for every kind of measurement by knowing the length of this triangle.

6 Least computations as a guidance principle

The computations of gravitational measurements will use less computations if the relative gravitational red-shift equal to the difference in gravitational red-shift. In that case, what can represent the ratio will certainly represent the difference, with one variable. To achieve

the least computations, the local gravitational measurements should satisfy the following condition

$$z_A - z_R = \frac{z_A}{z_R} \quad (6.1)$$

For weak gravitational approximation; this condition can be written as

$$-\frac{r_s}{2} = \frac{r_R^2}{r_A - r_R} \quad (6.2)$$

Or it can be expressed in terms of black hole mass as

$$GM = -\frac{r_A^2 r_R^2}{r_A - r_R} \quad (6.3)$$

This equation introduces matter-gravity equation which is noted to be inverse proportionality between Matter and Gravity.

7 Standard model as Black hole global Symmetry

The experimental measurements would follow the standard model that has a symmetry $SU(3) \times SU(2) \times U(1)$ where its measurements have been confirmed in local inertial frames [8]. Therefore, we conjecture that our universe emerges from a black hole that has internal global symmetry $SU(3) \times SU(2) \times U(1)$. Or in other words, Gravity is a global symmetry of $SU(3) \times SU(2) \times U(1)$. Or in other words, Gravity is the timeless state of the universe. This group has 12 ($8 + 3 + 1$) free parameters. Through localization of this symmetry in the gravitational field between any two different points, the gauge fields are generated. We agree that localization in the gravitational field will be represented by the relative gravitational red-shift which explain why the internal symmetries such as spin of these gauge fields does not depend on time. Their relative ratio will be related by relative gravitational red-shift, and in local inertial frames, they will remain the same at the different points.

7.1 geometric interpretation of Spontaneous Symmetry Breaking in the timeless state

The spontaneous symmetry breaking would correspond to the emergence of mass in the timeless state of the universe. As we have shown in previous sections, the mass emerges when there is a change in gravitational red-shift between two different points at two different distances from the black hole. The black hole has a global symmetry $SU(3) \times SU(2) \times U(1)$. When we localize this symmetry between two different points, a mass emerges. This localization process would generate a two different spherical symmetry groups $SO(3)$ at the two different points. Using isomorphism, $SU(2)/\mathbb{Z}_2 \sim SO(3)$. This fundamental geometry relation would imply that the $SU(2)$ symmetry part of the gravity global symmetry has been broken into two different $SO(3)$ with two different radii, and the mass emerges as a result of difference in value between these two different radii. This

means $SU(3)$ and $U(1)$ symmetry will remain unbroken which explain why gluons and photon remain mass-less in the standard model and their internal symmetries are measured independent of time such as color and hyper charge. The Higgs modes in that sense can be understood as the gravitational red-shift at every point which formulate the spin-0 modes.

8 Event Horizon as Mirror Symmetry breaking

The localization process for gravity outside the black hole is only possible out side the event Horizon. Therefore, we can realize the black hole horizon as the edge of the black hole at which its global symmetry can start breaking. This mirror symmetry breaking at the event Horizon between “global” $SU(3) \times SU(2) \times U(1)$ and “Local” $SU(3) \times SU(2) \times U(1)$ can explain why do we have matter rather than anti-matter. If the creation of particle-antiparticle happens at the neighbourhood of event Horizon, then we can call the energy modes that are detected in “Local” $SU(3) \times SU(2) \times U(1)$ as a Matter or particles, and we can call the Matter or particles that enter the black hole global $SU(3) \times SU(2) \times U(1)$ as anti-matter or anti-particles. This would solve the fundamental question, why do we measure matter and not anti-matter!

9 Solution for the Cosmological Constant Problem

We have shown that Spin 0 states are represented by the gravitational red-shift at every point. This is interpreted as the scalar field that go through the space which is the Higgs field. Therefore, the Higgs mass would be related to the gravitational red-shift as follows:

$$z \propto m_H \approx 10^{-26} \text{Kg} \quad (9.1)$$

The number 10^{-26} is related the local measurement scale we have in the Laboratory. On the other side, if the cosmological constant can be represented by the vacuum energy which can be represented by the integration of gravitational red-shift through the space around the black hole. Therefore, the full integration of z over the whole space will be given by

$$\Lambda = \int z dz = \frac{z^2}{2} \quad (9.2)$$

Since Higgs modes are related to the gravitational red-shift in the scale of measurement that we realize, therefore, the total vacuum energy will be given by

$$\Lambda = \frac{z^2}{2} \approx \frac{m_H^2}{2} \approx 10^{-52} \quad (9.3)$$

This is around the cosmological constant value derived from Higgs mass modes when we interpret it as gravitational red-shift.

10 Origin of Conservation of Energy

Based on gravitational red-shift computations and the timeless state of the universe, a principle can be added for the emergence of conservation of energy

- For every action and a reaction, there is always a gravitational source to which both events exist simultaneously in a timeless state, which explains the origin of conservation of energy.

11 Conclusion

We match non-inertial frames with local inertial frames at their connecting point through equating the relative gravitational red-shift with the gamma factor in local inertial frames. We got a timeless state of the universe in which we found a geometric interpretations of speed of light, mass and spin. We found that the measured spin of quantum particles in quantum entanglement is an experimental evidence for the timeless state of the universe, in which relative gravitational red-shift is measured. The spin in that sense is the relative gravitational red-shift between any two different points of measurement. We found gravity would correspond to all measurements that are independent of time such as internal symmetries. We conjecture that the gravity is the global symmetry $SU(3) \times SU(2) \times U(1)$, and would correspond to all internal symmetries measurements that are independent of time. Gravity in that sense can be understood as the timeless state of the universe. We answered the question why do we measure matter and not anti-matter, and we introduced a solution for explaining the value of cosmological constant problem. The correlation that we introduce between internal symmetries and gravity would open the door for new gravitational technology.

Acknowledgment The author is deeply grateful for enlightening discussions with Ebte-sam Nafie, Klee Irwin, Raymond Aschiem, Hassan Alshal, Bahaa Hany Sorial, Ahmed Al, Fang Fang, Richard Clawson, Marcelo Amaral, David Chester, Haley Irwin, Alexandra Von Waden, Dugan Hammock, and Stephanie Nadanarajah. This work is supported by the quantum gravity research grant, Los Angeles, California.

References

- [1] A. Albrecht and J. Magueijo, Phys. Rev. D **59**, 043516 (1999) doi:10.1103/PhysRevD.59.043516 [astro-ph/9811018].
- [2] M. R. Wilczynska, J. K. Webb, M. Bainbridge, S. E. Bosman, J. D. Barrow, R. F. Carswell, M. P. Dabrowski, V. Dumont, A. C. Leite, C. Lee, K. Leszczynska, J. Liske, K. Marosek, C. J. Martins, D. Milakovic, P. Molaro and L. Pasquini, [arXiv:2003.07627 [astro-ph.CO]].

- [3] Gisin, N. Mathematical languages shape our understanding of time in physics. *Nat. Phys.* **16**, 114–116 (2020).
- [4] R. V. Pound and G. A. Rebka, Jr., *Phys. Rev. Lett.* **4**, 337 (1960).
doi:10.1103/PhysRevLett.4.337
- [5] R. V. Pound and J. L. Snider, *Phys. Rev. Lett.* **13**, 539 (1964).
doi:10.1103/PhysRevLett.13.539
- [6] A. Einstein, B. Podolsky and N. Rosen, *Phys. Rev.* **47**, 777-780 (1935)
doi:10.1103/PhysRev.47.777
- [7] Kwiat, Paul G., et al. "New high-intensity source of polarization-entangled photon pairs." *Physical Review Letters* **75**.24 (1995): 4337.
- [8] G. Aad *et al.* [ATLAS], *Phys. Lett. B* **716**, 1-29 (2012)
doi:10.1016/j.physletb.2012.08.020 [arXiv:1207.7214 [hep-ex]].