

A Study On Sums of Cubes of Generalized Fibonacci Numbers: Closed Formulas of

$$\sum_{k=0}^n x^k W_k^3 \text{ and } \sum_{k=1}^n x^k W_{-k}^3$$

Yüksel Soykan

*Department of Mathematics,
Art and Science Faculty,
Zonguldak Bülent Ecevit University,
67100, Zonguldak, Turkey
e-mail: yuksel_soykan@hotmail.com*

Abstract. In this paper, closed forms of the sum formulas $\sum_{k=0}^n x^k W_k^3$ and $\sum_{k=1}^n x^k W_{-k}^3$ for the cubes of generalized Fibonacci numbers are presented. As special cases, we give sum formulas of Fibonacci, Lucas, Pell, Pell-Lucas, Jacobsthal, Jacobsthal-Lucas numbers.

2010 Mathematics Subject Classification. 11B37, 11B39, 11B83.

Keywords : Fibonacci numbers; Lucas numbers; Pell numbers; Jacobsthal numbers; sum formulas

1. Introduction

There are so many studies in the literature that concern about special second order recurrence sequences such as Fibonacci and Lucas. The sequence of Fibonacci numbers $\{F_n\}$ is defined by

$$F_n = F_{n-1} + F_{n-2}, \quad n \geq 2, \quad F_0 = 0, \quad F_1 = 1.$$

and the sequence of Lucas numbers $\{L_n\}$ is defined by

$$L_n = L_{n-1} + L_{n-2}, \quad n \geq 2, \quad L_0 = 2, \quad L_1 = 1.$$

The Fibonacci numbers, Lucas numbers and their generalizations have many interesting properties and applications to almost every field. Horadam [8] defined a generalization of Fibonacci sequence, that is, he defined a second-order linear recurrence sequence $\{W_n(W_0, W_1; r, s)\}$, or simply $\{W_n\}$, as follows:

$$W_n = rW_{n-1} + sW_{n-2}; \quad W_0 = a, \quad W_1 = b, \quad (n \geq 2) \quad (1.1)$$

where W_0, W_1 are arbitrary complex numbers and r, s are real numbers, see also Horadam [7], [9] and [10]. Now these generalized Fibonacci numbers $\{W_n(a, b; r, s)\}$ are also called Horadam numbers. The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = -\frac{r}{s}W_{-(n-1)} + \frac{1}{s}W_{-(n-2)}$$

for $n = 1, 2, 3, \dots$ when $s \neq 0$. Therefore, recurrence (1.1) holds for all integer n .

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

For some specific values of a, b, r and s , it is worth presenting these special Horadam numbers in a table as a specific name. In literature, for example, the following names and notations (see Table 1) are used for the special cases of r, s and initial values.

Table 1. A few special case of generalized Fibonacci sequences.

Name of sequence	Notation: $W_n(a, b; r, s)$	OEIS: [17]
Fibonacci	$F_n = W_n(0, 1; 1, 1)$	A000045
Lucas	$L_n = W_n(2, 1; 1, 1)$	A000032
Pell	$P_n = W_n(0, 1; 2, 1)$	A000129
Pell-Lucas	$Q_n = W_n(2, 2; 2, 1)$	A002203
Jacobsthal	$J_n = W_n(0, 1; 1, 2)$	A001045
Jacobsthal-Lucas	$j_n = W_n(2, 1; 1, 2)$	A014551

The evaluation of sums of powers of these sequences is a challenging issue. Two pretty examples are

$$\sum_{k=0}^n (-1)^k F_k^3 = \frac{1}{2} ((-1)^n (F_{n+2}^3 + F_{n+1}^3 - 3F_{n+1}^2 F_{n+2}) + 1)$$

and

$$\sum_{k=1}^n (-1)^k L_{-k}^3 = \frac{1}{2} ((-1)^n (L_{-n+1}^3 + L_{-n}^3 - 3L_{-n}^2 L_{-n+1}) + 3).$$

In this work, we derive expressions for sums of second powers of generalized Fibonacci numbers. We present some works on sum formulas of powers of the numbers in the following Table 2.

Table 2. A few special study on sum formulas of second, third and arbitrary powers.

Name of sequence	sums of second powers	sums of third powers	sums of powers
Generalized Fibonacci	[1,2,6,11,12,18]	[5,19,21]	[3,4,13]
Generalized Tribonacci	[15,20]		
Generalized Tetranacci	[14,16]		

2. Sum Formulas of Generalized Fibonacci Numbers with Positive Subscripts

The following theorem presents some summing formulas of generalized Fibonacci numbers with positive subscripts.

THEOREM 2.1. *Let x be a complex number. For $n \geq 0$ we have the following formulas: If $(-s^3 x^2 + rsx + 1)(r^3 x + s^3 x^2 + 3rsx - 1) \neq 0$ then*

(a):

$$\sum_{k=0}^n x^k W_k^3 = \frac{\Lambda_1}{(-s^3 x^2 + rsx + 1)(r^3 x + s^3 x^2 + 3rsx - 1)}$$

where

$$\begin{aligned} \Lambda_1 = & -x^{n+2}(s^3 x^2 + 2rsx - 1)W_{n+2}^3 - x^{n+1}(r^3 x + s^3 x^2 + 3r^2 s^2 x^2 - r^3 s^3 x^3 + r^4 s x^2 + 2rsx - 1)W_{n+1}^3 \\ & + 3rsx^{n+3}(r + s^2 x)W_{n+2}^2 W_{n+1} - 3rs^2 x^{n+3}(rsx - 1)W_{n+1}^2 W_{n+2} + x(s^3 x^2 + 2rsx - 1)W_1^3 \\ & + (r^3 x + s^3 x^2 + 3r^2 s^2 x^2 - r^3 s^3 x^3 + r^4 s x^2 + 2rsx - 1)W_0^3 \\ & - 3rsx^2(r + s^2 x)W_1^2 W_0 + 3rs^2 x^2(rsx - 1)W_0^2 W_1 \end{aligned}$$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

(b):

$$\sum_{k=0}^n x^k W_k^2 W_{k+1} = \frac{\Lambda_2}{(-s^3 x^2 + rsx + 1)(r^3 x + s^3 x^2 + 3rsx - 1)}$$

where

$$\begin{aligned} \Lambda_2 = & -rx^{n+2}(rsx - 1)W_{n+2}^3 - rs^3x^{n+3}(rsx - 1)W_{n+1}^3 + sx^{n+2}(2r^3x - s^3x^2 + 1)W_{n+2}^2W_{n+1} \\ & -x^{n+1}(r^3x + s^3x^2 + r^4sx^2 - 2rs^4x^3 + 2rsx - 1)W_{n+1}^2W_{n+2} + rx(rsx - 1)W_1^3 \\ & +rs^3x^2(rsx - 1)W_0^3 - sx(2r^3x - s^3x^2 + 1)W_1^2W_0 \\ & +(r^3x + s^3x^2 + r^4sx^2 - 2rs^4x^3 + 2rsx - 1)W_0^2W_1 \end{aligned}$$

(c):

$$\sum_{k=0}^n x^k W_{k+1}^2 W_k = \frac{\Lambda_3}{(-s^3 x^2 + rsx + 1)(r^3 x + s^3 x^2 + 3rsx - 1)}$$

where

$$\begin{aligned} \Lambda_3 = & rx^{n+2}(r + s^2x)W_{n+2}^3 + rs^3x^{n+3}(r + s^2x)W_{n+1}^3 - x^{n+1}(r^3x + s^3x^2 + 3r^2s^2x^2 - 1)W_{n+2}^2W_{n+1} \\ & +s^2x^{n+2}(2r^3x - s^3x^2 + 1)W_{n+1}^2W_{n+2} - rx(r + s^2x)W_1^3 - rs^3x^2(r + s^2x)W_0^3 \\ & +(r^3x + s^3x^2 + 3r^2s^2x^2 - 1)W_1^2W_0 - s^2x(2r^3x - s^3x^2 + 1)W_0^2W_1 \end{aligned}$$

Proof. Using the recurrence relation

$$W_{n+2} = rW_{n+1} + sW_n$$

i.e.

$$sW_n = W_{n+2} - rW_{n+1}$$

we obtain

$$s^3W_n^3 = W_{n+2}^3 - 3rW_{n+2}^2W_{n+1} + 3r^2W_{n+1}^2W_{n+2} - r^3W_{n+1}^3$$

and so

$$\begin{aligned} s^3x^nW_n^3 &= x^nW_{n+2}^3 - 3rx^nW_{n+2}^2W_{n+1} + 3r^2x^nW_{n+1}^2W_{n+2} - r^3x^nW_{n+1}^3 \\ s^3x^{n-1}W_{n-1}^3 &= x^{n-1}W_{n+1}^3 - 3rx^{n-1}W_{n+1}^2W_n + 3r^2x^{n-1}W_n^2W_{n+1} - r^3x^{n-1}W_n^3 \\ s^3x^{n-2}W_{n-2}^3 &= x^{n-2}W_n^3 - 3rx^{n-2}W_n^2W_{n-1} + 3r^2x^{n-2}W_{n-1}^2W_n - r^3x^{n-2}W_{n-1}^3 \\ &\vdots \\ s^3x^2W_2^3 &= x^2W_4^3 - 3rx^2W_4^2W_3 + 3r^2x^2W_3^2W_4 - r^3x^2W_3^3 \\ s^3x^1W_1^3 &= x^1W_3^3 - 3rx^1W_3^2W_2 + 3r^2x^1W_2^2W_3 - r^3x^1W_2^3 \\ s^3x^0W_0^3 &= x^0W_2^3 - 3rx^0W_2^2W_1 + 3r^2x^0W_1^2W_2 - r^3x^0W_1^3 \end{aligned}$$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

If we add the above equations by side by, we get

$$\begin{aligned} s^3 \sum_{k=0}^n x^k W_k^3 &= (x^n W_{n+2}^3 + x^{n-1} W_{n+1}^3 - x^{-1} W_1^3 - x^{-2} W_0^3 + x^{-2} \sum_{k=0}^n x^k W_k^3) \\ &\quad - 3r(x^n W_{n+2}^2 W_{n+1} - x^{-1} W_1^2 W_0 + x^{-1} \sum_{k=0}^n x^k W_{k+1}^2 W_k) \\ &\quad + 3r^2(x^n W_{n+1}^2 W_{n+2} - x^{-1} W_0^2 W_1 + x^{-1} \sum_{k=0}^n x^k W_k^2 W_{k+1}) \\ &\quad - r^3(x^n W_{n+1}^3 - x^{-1} W_0^3 + x^{-1} \sum_{k=0}^n x^k W_k^3). \end{aligned} \quad (2.1)$$

Next we calculate $\sum_{k=0}^n x^k W_{k+1}^2 W_k$. Again, using the recurrence relation

$$W_{n+2} = rW_{n+1} + sW_n$$

i.e.

$$sW_n = W_{n+2} - rW_{n+1}$$

we obtain

$$sW_{n+1}^2 W_n = W_{n+1}^2 W_{n+2} - rW_{n+1}^3$$

and so

$$\begin{aligned} sx^n W_{n+1}^2 W_n &= x^n W_{n+1}^2 W_{n+2} - rx^n W_{n+1}^3 \\ sx^{n-1} W_n^2 W_{n-1} &= x^{n-1} W_n^2 W_{n+1} - rx^{n-1} W_n^3 \\ sx^{n-2} W_{n-1}^2 W_{n-2} &= x^{n-2} W_{n-1}^2 W_n - rx^{n-2} W_{n-1}^3 \\ &\vdots \\ sx^2 W_3^2 W_2 &= x^2 W_3^2 W_4 - rx^2 W_3^3 \\ sx^1 W_2^2 W_1 &= x^1 W_2^2 W_3 - rx^1 W_2^3 \\ sx^0 W_1^2 W_0 &= x^0 W_1^2 W_2 - rx^0 W_1^3 \end{aligned}$$

If we add the above equations by side by, we get

$$\begin{aligned} s \sum_{k=0}^n x^k W_{k+1}^2 W_k &= (x^n W_{n+1}^2 W_{n+2} - x^{-1} W_0^2 W_1 + x^{-1} \sum_{k=0}^n x^k W_k^2 W_{k+1}) \\ &\quad - r(x^n W_{n+1}^3 - x^{-1} W_0^3 + x^{-1} \sum_{k=0}^n x^k W_k^3) \end{aligned} \quad (2.2)$$

Next we calculate $\sum_{k=0}^n x^k W_k^2 W_{k+1}$. Again, using the recurrence relation

$$W_{n+2} = rW_{n+1} + sW_n$$

i.e.

$$sW_n = W_{n+2} - rW_{n+1} \Rightarrow s^2 W_n^2 = W_{n+2}^2 + r^2 W_{n+1}^2 - 2rW_{n+2} W_{n+1}$$

we obtain

$$s^2 W_n^2 W_{n+1} = W_{n+2}^2 W_{n+1} + r^2 W_{n+1}^3 - 2rW_{n+1}^2 W_{n+2}$$

and so

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

$$\begin{aligned}
 s^2 x^n W_n^2 W_{n+1} &= x^n W_{n+2}^2 W_{n+1} + r^2 x^n W_{n+1}^3 - 2rx^n W_{n+1}^2 W_{n+2} \\
 s^2 x^{n-1} W_{n-1}^2 W_n &= x^{n-1} W_{n+1}^2 W_n + r^2 x^{n-1} W_n^3 - 2rx^{n-1} W_n^2 W_{n+1} \\
 s^2 x^{n-2} W_{n-2}^2 W_{n-1} &= x^{n-2} W_n^2 W_{n-1} + r^2 x^{n-2} W_{n-1}^3 - 2rx^{n-2} W_{n-1}^2 W_n \\
 &\vdots \\
 s^2 x^2 W_2^2 W_3 &= x^2 W_4^2 W_3 + r^2 x^2 W_3^3 - 2rx^2 W_3^2 W_4 \\
 s^2 x^1 W_1^2 W_2 &= x^1 W_3^2 W_2 + r^2 x^1 W_2^3 - 2rx^1 W_2^2 W_3 \\
 s^2 x^0 W_0^2 W_1 &= x^0 W_2^2 W_1 + r^2 x^0 W_1^3 - 2rx^0 W_1^2 W_2
 \end{aligned}$$

If we add the above equations by side by, we get

$$\begin{aligned}
 s^2 \sum_{k=0}^n x^k W_k^2 W_{k+1} &= (x^n W_{n+2}^2 W_{n+1} - x^{-1} W_1^2 W_0 + x^{-1} \sum_{k=0}^n x^k W_{k+1}^2 W_k) \\
 &\quad + r^2 (x^n W_{n+1}^3 - x^{-1} W_0^3 + x^{-1} \sum_{k=0}^n x^k W_k^3) \\
 &\quad - 2r(x^n W_{n+1}^2 W_{n+2} - x^{-1} W_0^2 W_1 + x^{-1} \sum_{k=0}^n x^k W_k^2 W_{k+1})
 \end{aligned} \tag{2.3}$$

Solving the system (2.1)-(2.2)-(2.3), the required results of (a),(b) and (c) follow.

2.1. The case $x = 1$. See [19] for the case $x = 1$.

2.2. The case $x = -1$. In this subsection we consider the special case $x = -1$.

Taking $x = -1, r = s = 1$ in Theorem 2.1 (a),(b) and (c), we obtain the following proposition.

PROPOSITION 2.2. *If $x = -1, r = s = 1$ then for $n \geq 0$ we have the following formulas:*

- (a): $\sum_{k=0}^n (-1)^k W_k^3 = \frac{1}{2}((-1)^n (W_{n+2}^3 + W_{n+1}^3 - 3W_{n+1}^2 W_{n+2}) + W_1^3 + W_0^3 - 3W_0^2 W_1).$
- (b): $\sum_{k=0}^n (-1)^k W_k^2 W_{k+1} = \frac{1}{2}((-1)^n (W_{n+2}^3 - W_{n+1}^3 - W_{n+2}^2 W_{n+1}) + W_1^3 - W_0^3 - W_1^2 W_0).$
- (c): $\sum_{k=0}^n (-1)^k W_{k+1}^2 W_k = \frac{1}{2}((-1)^n (W_{n+2}^2 W_{n+1} - W_{n+1}^2 W_{n+2}) + W_1^2 W_0 - W_0^2 W_1).$

From the above proposition, we have the following corollary which gives sum formulas of Fibonacci numbers (take $W_n = F_n$ with $F_0 = 0, F_1 = 1$).

COROLLARY 2.3. *For $n \geq 0$, Fibonacci numbers have the following properties:*

- (a): $\sum_{k=0}^n (-1)^k F_k^3 = \frac{1}{2}((-1)^n (F_{n+2}^3 + F_{n+1}^3 - 3F_{n+1}^2 F_{n+2}) + 1).$
- (b): $\sum_{k=0}^n (-1)^k F_k^2 F_{k+1} = \frac{1}{2}((-1)^n (F_{n+2}^3 - F_{n+1}^3 - F_{n+2}^2 F_{n+1}) + 1).$
- (c): $\sum_{k=0}^n (-1)^k F_{k+1}^2 F_k = \frac{1}{2}((-1)^n (F_{n+2}^2 F_{n+1} - F_{n+1}^2 F_{n+2})).$

Taking $W_n = L_n$ with $L_0 = 2, L_1 = 1$ in the last proposition, we have the following corollary which presents sum formulas of Lucas numbers.

COROLLARY 2.4. *For $n \geq 0$, Lucas numbers have the following properties:*

- (a): $\sum_{k=0}^n (-1)^k L_k^3 = \frac{1}{2}((-1)^n (L_{n+2}^3 + L_{n+1}^3 - 3L_{n+1}^2 L_{n+2}) - 3).$
- (b): $\sum_{k=0}^n (-1)^k L_k^2 L_{k+1} = \frac{1}{2}((-1)^n (L_{n+2}^3 - L_{n+1}^3 - L_{n+2}^2 L_{n+1}) - 9).$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

$$(c): \sum_{k=0}^n (-1)^k L_{k+1}^2 L_k = \frac{1}{2}((-1)^n (L_{n+2}^2 L_{n+1} - L_{n+1}^2 L_{n+2}) - 2).$$

Taking $x = -1, r = 2, s = 1$ in Theorem 2.1 (a),(b) and (c), we obtain the following proposition.

PROPOSITION 2.5. *If $x = -1, r = 2, s = 1$ then for $n \geq 0$ we have the following formulas:*

$$\begin{aligned} (a): \sum_{k=0}^n (-1)^k W_k^3 &= \frac{1}{14}((-1)^n (2W_{n+2}^3 + 12W_{n+1}^3 - 3W_{n+2}^2 W_{n+1} - 9W_{n+1}^2 W_{n+2}) + 2W_1^3 + 12W_0^3 - 3W_1^2 W_0 - 9W_0^2 W_1). \\ (b): \sum_{k=0}^n (-1)^k W_k^2 W_{k+1} &= \frac{1}{14}((-1)^n (3W_{n+2}^3 - 3W_{n+1}^3 - 8W_{n+2}^2 W_{n+1} + 4W_{n+1}^2 W_{n+2}) + 3W_1^3 - 3W_0^3 - 8W_1^2 W_0 + 4W_0^2 W_1). \\ (c): \sum_{k=0}^n (-1)^k W_{k+1}^2 W_k &= \frac{1}{14}((-1)^n (W_{n+2}^3 - W_{n+1}^3 + 2W_{n+2}^2 W_{n+1} - 8W_{n+1}^2 W_{n+2}) + W_1^3 - W_0^3 + 2W_1^2 W_0 - 8W_0^2 W_1). \end{aligned}$$

From the last proposition, we have the following corollary which gives sum formulas of Pell numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1$).

COROLLARY 2.6. *For $n \geq 0$, Pell numbers have the following properties:*

$$\begin{aligned} (a): \sum_{k=0}^n (-1)^k P_k^3 &= \frac{1}{14}((-1)^n (2P_{n+2}^3 + 12P_{n+1}^3 - 3P_{n+2}^2 P_{n+1} - 9P_{n+1}^2 P_{n+2}) + 2). \\ (b): \sum_{k=0}^n (-1)^k P_k^2 P_{k+1} &= \frac{1}{14}((-1)^n (3P_{n+2}^3 - 3P_{n+1}^3 - 8P_{n+2}^2 P_{n+1} + 4P_{n+1}^2 P_{n+2}) + 3). \\ (c): \sum_{k=0}^n (-1)^k P_{k+1}^2 P_k &= \frac{1}{14}((-1)^n (P_{n+2}^3 - P_{n+1}^3 + 2P_{n+2}^2 P_{n+1} - 8P_{n+1}^2 P_{n+2}) + 1). \end{aligned}$$

Taking $W_n = Q_n$ with $Q_0 = 2, Q_1 = 2$ in the last proposition, we have the following corollary which presents sum formulas of Pell-Lucas numbers.

COROLLARY 2.7. *For $n \geq 0$, Pell-Lucas numbers have the following properties:*

$$\begin{aligned} (a): \sum_{k=0}^n (-1)^k Q_k^3 &= \frac{1}{14}((-1)^n (2Q_{n+2}^3 + 12Q_{n+1}^3 - 3Q_{n+2}^2 Q_{n+1} - 9Q_{n+1}^2 Q_{n+2}) + 16). \\ (b): \sum_{k=0}^n (-1)^k Q_k^2 Q_{k+1} &= \frac{1}{14}((-1)^n (3Q_{n+2}^3 - 3Q_{n+1}^3 - 8Q_{n+2}^2 Q_{n+1} + 4Q_{n+1}^2 Q_{n+2}) - 32). \\ (c): \sum_{k=0}^n (-1)^k Q_{k+1}^2 Q_k &= \frac{1}{14}((-1)^n (Q_{n+2}^3 - Q_{n+1}^3 + 2Q_{n+2}^2 Q_{n+1} - 8Q_{n+1}^2 Q_{n+2}) - 48). \end{aligned}$$

If $x = -1, r = 1, s = 2$ then $(-s^3 x^2 + rsx + 1)(r^3 x + s^3 x^2 + 3rsx - 1) = 0$ so we can't use Theorem 2.1 directly. But we can give another method to find $\sum_{k=0}^n (-1)^k W_k^3$, $\sum_{k=0}^n (-1)^k W_k^2 W_{k+1}$ and $\sum_{k=0}^n (-1)^k W_{k+1}^2 W_k$ by using Theorem 2.1.

THEOREM 2.8. *If $x = -1, r = 1, s = 2$ then for $n \geq 0$ we have the following formulas:*

$$\begin{aligned} (a): \sum_{k=0}^n (-1)^k W_k^3 &= \frac{1}{81}((-1)^n ((3n+18)W_{n+2}^3 - (24n+87)W_{n+1}^3 - 6(3n+13)W_{n+2}^2 W_{n+1} + 12(3n+11)W_{n+1}^2 W_{n+2} + 15W_1^3 - 63W_0^3 - 60W_1^2 W_0 + 96W_0^2 W_1). \\ (b): \sum_{k=0}^n (-1)^k W_k^2 W_{k+1} &= \frac{1}{81}((-1)^n (-(3n+8)W_{n+2}^3 + 8(3n+11)W_{n+1}^3 + 2(9n+36)W_{n+2}^2 W_{n+1} - (36n+147)W_{n+1}^2 W_{n+2} - 5W_1^3 + 64W_0^3 + 54W_1^2 W_0 - 111W_0^2 W_1). \\ (c): \sum_{k=0}^n (-1)^k W_{k+1}^2 W_k &= \frac{1}{81}((-1)^n ((3n+10)W_{n+2}^3 - 8(3n+13)W_{n+1}^3 - (18n+57)W_{n+2}^2 W_{n+1} + 4(9n+36)W_{n+1}^2 W_{n+2} + 7W_1^3 - 80W_0^3 - 39W_1^2 W_0 + 108W_0^2 W_1). \end{aligned}$$

Proof.

(a): We use Theorem 2.1 (a). If we set $r = 1, s = 2$ in Theorem 2.1 (a) then we have

$$\sum_{k=0}^n x^k W_k^3 = \frac{f_1(x)}{(-8x^2 + 2x + 1)(8x^2 + 7x - 1)}$$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

where

$$\begin{aligned} f_1(x) &= -x^{n+2}(8x^2 + 4x - 1)W_{n+2}^3 + x^{n+1}(8x^3 - 22x^2 - 5x + 1)W_{n+1}^3 \\ &\quad + 6x^{n+3}(4x + 1)W_{n+2}^2 W_{n+1} - 12x^{n+3}(2x - 1)W_{n+1}^2 W_{n+2} + x(8x^2 + 4x - 1)W_1^3 \\ &\quad - (8x^3 - 22x^2 - 5x + 1)W_0^3 - 6x^2(4x + 1)W_1^2 W_0 + 12x^2(2x - 1)W_0^2 W_1. \end{aligned}$$

For $x = -1$, the right hand side of the above sum formulas is an indeterminate form. Now, we can use L'Hospital rule. Then we get

$$\begin{aligned} \sum_{k=0}^n (-1)^k W_k^2 W_{k+1} &= \left. \frac{\frac{d}{dx}(f_1(x))}{\frac{d}{dx}((-8x^2 + 2x + 1)(8x^2 + 7x - 1))} \right|_{x=-1} \\ &= \frac{1}{81} ((-1)^n ((3n + 18) W_{n+2}^3 - (24n + 87) W_{n+1}^3 - 6(3n + 13) W_{n+2}^2 W_{n+1} \\ &\quad + 12(3n + 11) W_{n+1}^2 W_{n+2}) + 15W_1^3 - 63W_0^3 - 60W_1^2 W_0 + 96W_0^2 W_1). \end{aligned}$$

(b): We use Theorem 2.1 (b). If we set $r = 1, s = 2$ in Theorem 2.1 (b) then we have

$$\sum_{k=0}^n x^k W_k^2 W_{k+1} = \frac{f_2(x)}{(-8x^2 + 2x + 1)(8x^2 + 7x - 1)}$$

where

$$\begin{aligned} f_2(x) &= -x^{n+2}(2x - 1)W_{n+2}^3 - 8x^{n+3}(2x - 1)W_{n+1}^3 + 2x^{n+2}(-8x^2 + 2x + 1)W_{n+2}^2 W_{n+1} \\ &\quad - x^{n+1}(-32x^3 + 10x^2 + 5x - 1)W_{n+1}^2 W_{n+2} + x(2x - 1)W_1^3 + 8x^2(2x - 1)W_0^3 \\ &\quad + (-32x^3 + 10x^2 + 5x - 1)W_0^2 W_1 - 2x(-8x^2 + 2x + 1)W_1^2 W_0. \end{aligned}$$

For $x = -1$, the right hand side of the above sum formulas is an indeterminate form. Now, we can use L'Hospital rule. Then we get

$$\begin{aligned} \sum_{k=0}^n (-1)^k W_k^2 W_{k+1} &= \left. \frac{\frac{d}{dx}(f_2(x))}{\frac{d}{dx}((-8x^2 + 2x + 1)(8x^2 + 7x - 1))} \right|_{x=-1} \\ &= \frac{1}{81} ((-1)^n (-(3n + 8) W_{n+2}^3 + 8(3n + 11) W_{n+1}^3 + 2(9n + 36) W_{n+2}^2 W_{n+1} \\ &\quad - (36n + 147) W_{n+1}^2 W_{n+2}) - 5W_1^3 + 64W_0^3 + 54W_1^2 W_0 - 111W_0^2 W_1). \end{aligned}$$

(c): We use Theorem 2.1 (c). If we set $r = 1, s = 2$ in Theorem 2.1 (c) then we have

$$\sum_{k=0}^n x^k W_{k+1}^2 W_k = \frac{f_3(x)}{(-8x^2 + 2x + 1)(8x^2 + 7x - 1)}$$

where

$$\begin{aligned} f_3(x) &= x^{n+2}(4x + 1)W_{n+2}^3 + 8x^{n+3}(4x + 1)W_{n+1}^3 - x^{n+1}(20x^2 + x - 1)W_{n+2}^2 W_{n+1} \\ &\quad + 4x^{n+2}(-8x^2 + 2x + 1)W_{n+1}^2 W_{n+2} - x(4x + 1)W_1^3 - 8x^2(4x + 1)W_0^3 \\ &\quad + (20x^2 + x - 1)W_1^2 W_0 - 4x(-8x^2 + 2x + 1)W_0^2 W_1. \end{aligned}$$

For $x = -1$, the right hand side of the above sum formulas is an indeterminate form. Now, we can use L'Hospital rule. Then we get

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

$$\begin{aligned} \sum_{k=0}^n (-1)^k W_{k+1}^2 W_k &= \left. \frac{\frac{d}{dx}(f_3(x))}{\frac{d}{dx}((-8x^2 + 2x + 1)(8x^2 + 7x - 1))} \right|_{x=-1} \\ &= \frac{1}{81} ((-1)^n ((3n + 10) W_{n+2}^3 - 8(3n + 13) W_{n+1}^3 - (18n + 57) W_{n+2}^2 W_{n+1} \\ &\quad + 4(9n + 36) W_{n+1}^2 W_{n+2}) + 7W_1^3 - 80W_0^3 - 39W_1^2 W_0 + 108W_0^2 W_1). \end{aligned}$$

From the last theorem we have the following corollary which gives sum formulas of Jacobsthal numbers (take $W_n = J_n$ with $J_0 = 0, J_1 = 1$).

COROLLARY 2.9. For $n \geq 0$, Jacobsthal numbers have the following properties:

- (a): $\sum_{k=0}^n (-1)^k J_k^3 = \frac{1}{81} ((-1)^n ((3n + 18) J_{n+2}^3 - (24n + 87) J_{n+1}^3 - 6(3n + 13) J_{n+2}^2 J_{n+1} + 12(3n + 11) J_{n+1}^2 J_{n+2}) + 15)$.
- (b): $\sum_{k=0}^n (-1)^k J_k^2 J_{k+1} = \frac{1}{81} ((-1)^n (-(3n + 8) J_{n+2}^3 + 8(3n + 11) J_{n+1}^3 + 2(9n + 36) J_{n+2}^2 J_{n+1} - (36n + 147) J_{n+1}^2 J_{n+2}) - 5)$.
- (c): $\sum_{k=0}^n (-1)^k J_{k+1}^2 J_k = \frac{1}{81} ((-1)^n ((3n + 10) J_{n+2}^3 - 8(3n + 13) J_{n+1}^3 - (18n + 57) J_{n+2}^2 J_{n+1} + 4(9n + 36) J_{n+1}^2 J_{n+2}) + 7)$.

Taking $W_n = j_n$ with $j_0 = 2, j_1 = 1$ in the last theorem, we have the following corollary which presents sum formulas of Jacobsthal-Lucas numbers.

COROLLARY 2.10. For $n \geq 0$, Jacobsthal-Lucas numbers have the following properties:

- (a): $\sum_{k=0}^n (-1)^k j_k^3 = \frac{1}{81} ((-1)^n ((3n + 18) j_{n+2}^3 - (24n + 87) j_{n+1}^3 - 6(3n + 13) j_{n+2}^2 j_{n+1} + 12(3n + 11) j_{n+1}^2 j_{n+2}) - 225)$.
- (b): $\sum_{k=0}^n (-1)^k j_k^2 j_{k+1} = \frac{1}{81} ((-1)^n (-(3n + 8) j_{n+2}^3 + 8(3n + 11) j_{n+1}^3 + 2(9n + 36) j_{n+2}^2 j_{n+1} - (36n + 147) j_{n+1}^2 j_{n+2}) + 171)$.
- (c): $\sum_{k=0}^n (-1)^k j_{k+1}^2 j_k = \frac{1}{81} ((-1)^n ((3n + 10) j_{n+2}^3 - 8(3n + 13) j_{n+1}^3 - (18n + 57) j_{n+2}^2 j_{n+1} + 4(9n + 36) j_{n+1}^2 j_{n+2}) - 279)$.

2.3. The case $x = 1 + i$. In this subsection we consider the special case $x = 1 + i$.

Taking $x = 1 + i, r = s = 1$ in Theorem 2.1 (a), (b) and (c), we obtain the following proposition.

PROPOSITION 2.11. If $x = 1 + i, r = s = 1$ then for $n \geq 0$ we have the following formulas:

- (a): $\sum_{k=0}^n (1 + i)^k W_k^3 = \frac{1}{12+9i} ((1 + i)^n ((7 - 15i) W_{n+1}^3 + (6 + 6i) W_{n+1}^2 W_{n+2} + (-18 + 6i) W_{n+1} W_{n+2}^2 + (8 - 2i) W_{n+2}^3) - (3 - 5i) W_1^3 + (4 + 11i) W_0^3 + (6 - 12i) W_1^2 W_0 - 6W_0^2 W_1)$.
- (b): $\sum_{k=0}^n (1 + i)^k W_k^2 W_{k+1} = \frac{1}{12+9i} ((1 + i)^n ((2 + 2i) W_{n+1}^3 - (3 + 9i) W_{n+1}^2 W_{n+2} + 6i W_{n+1} W_{n+2}^2 + 2W_{n+2}^3) - (1 - i) W_1^3 - 2W_0^3 - (3 + 3i) W_1^2 W_0 + (6 + 3i) W_0^2 W_1)$.
- (c): $\sum_{k=0}^n (1 + i)^k W_{k+1}^2 W_k = \frac{1}{12+9i} ((1 + i)^n ((-6 + 2i) W_{n+1}^3 + 6i W_{n+1}^2 W_{n+2} + (9 - 9i) W_{n+1} W_{n+2}^2 + (-2 + 4i) W_{n+2}^3) - (1 + 3i) W_1^3 + (2 - 4i) W_0^3 + 9i W_1^2 W_0 - (3 + 3i) W_0^2 W_1)$.

From the above proposition, we have the following corollary which gives sum formulas of Fibonacci numbers (take $W_n = F_n$ with $F_0 = 0, F_1 = 1$).

COROLLARY 2.12. For $n \geq 0$, Fibonacci numbers have the following properties:

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{-k}^3$

$$\begin{aligned} \text{(a): } \sum_{k=0}^n (1+i)^k F_k^3 &= \frac{1}{12+9i} ((1+i)^n ((7-15i)F_{n+1}^3 + (6+6i)F_{n+1}^2 F_{n+2} + (-18+6i)F_{n+1} F_{n+2}^2 + (8-2i)F_{n+2}^3) - \\ &\quad 3 + 5i). \\ \text{(b): } \sum_{k=0}^n (1+i)^k F_k^2 F_{k+1} &= \frac{1}{12+9i} ((1+i)^n ((2+2i)F_{n+1}^3 - (3+9i)F_{n+1}^2 F_{n+2} + 6iF_{n+1} F_{n+2}^2 + 2F_{n+2}^3) - 1 + i). \\ \text{(c): } \sum_{k=0}^n (1+i)^k F_{k+1}^2 F_k &= \frac{1}{12+9i} ((1+i)^n ((-6+2i)F_{n+1}^3 + 6iF_{n+1}^2 F_{n+2} + (9-9i)F_{n+1} F_{n+2}^2 + (-2+4i)F_{n+2}^3) - \\ &\quad 1 - 3i). \end{aligned}$$

Taking $W_n = L_n$ with $L_0 = 2, L_1 = 1$ in the last proposition, we have the following corollary which presents sum formulas of Lucas numbers.

COROLLARY 2.13. For $n \geq 0$, Lucas numbers have the following properties:

$$\begin{aligned} \text{(a): } \sum_{k=0}^n (1+i)^k L_k^3 &= \frac{1}{12+9i} ((1+i)^n ((7-15i)L_{n+1}^3 + (6+6i)L_{n+1}^2 L_{n+2} + (-18+6i)L_{n+1} L_{n+2}^2 + (8- \\ &\quad 2i)L_{n+2}^3) + 17 + 69i). \\ \text{(b): } \sum_{k=0}^n (1+i)^k L_k^2 L_{k+1} &= \frac{1}{12+9i} ((1+i)^n ((2+2i)L_{n+1}^3 - (3+9i)L_{n+1}^2 L_{n+2} + 6iL_{n+1} L_{n+2}^2 + 2L_{n+2}^3) + 1 + 7i). \\ \text{(c): } \sum_{k=0}^n (1+i)^k L_{k+1}^2 L_k &= \frac{1}{12+9i} ((1+i)^n ((-6+2i)L_{n+1}^3 + 6iL_{n+1}^2 L_{n+2} + (9-9i)L_{n+1} L_{n+2}^2 + (-2+4i)L_{n+2}^3) + \\ &\quad 3 - 29i). \end{aligned}$$

Corresponding sums of the other second order linear sequences can be calculated similarly when $x = 1 + i$.

3. Sum Formulas of Generalized Fibonacci Numbers with Negative Subscripts

The following theorem presents some summing formulas of generalized Fibonacci numbers with negative subscripts.

THEOREM 3.1. Let x be a complex number. For $n \geq 1$ we have the following formulas: If $(-s^3 + x^2 + rsx)(r^3x + s^3 - x^2 + 3rsx) \neq 0$ then

$$\text{(a): } \sum_{k=1}^n x^k W_{-k}^3 = \frac{\Lambda_4}{(-s^3 + x^2 + rsx)(r^3x + s^3 - x^2 + 3rsx)}$$

where

$$\begin{aligned} \Lambda_4 &= x^{n+1}(s^3 - x^2 + 2rsx)W_{-n+1}^3 + x^{n+1}(s^3x - r^3s^3 + r^3x^2 - x^3 + 2rsx^2 + r^4sx + 3r^2s^2x)W_{-n}^3 \\ &\quad - 3rsx^{n+1}(rx + s^2)W_{-n+1}^2W_{-n} + 3rs^2x^{n+1}(-x + rs)W_{-n}^2W_{-n+1} - x(s^3 - x^2 + 2rsx)W_1^3 \\ &\quad + x(-s^3x + r^3s^3 - r^3x^2 + x^3 - 2rsx^2 - r^4sx - 3r^2s^2x)W_0^3 \\ &\quad + 3rsx(rx + s^2)W_1^2W_0 - 3rs^2x(-x + rs)W_0^2W_1 \end{aligned}$$

$$\text{(b): } \sum_{k=1}^n x^k W_{-k+1}^2 W_{-k} = \frac{\Lambda_5}{(-s^3 + x^2 + rsx)(r^3x + s^3 - x^2 + 3rsx)}$$

where

$$\begin{aligned} \Lambda_5 &= -rx^{n+2}(rx + s^2)W_{-n+1}^3 - rs^3x^{n+1}(rx + s^2)W_{-n}^3 \\ &\quad + x^{n+2}(r^3x + 3r^2s^2 + s^3 - x^2)W_{-n+1}^2W_{-n} + s^2x^{n+1}(-2r^3x + s^3 - x^2)W_{-n}^2W_{-n+1} \\ &\quad + rx^2(rx + s^2)W_1^3 + rs^3x(rx + s^2)W_0^3 - x^2(r^3x + 3r^2s^2 + s^3 - x^2)W_1^2W_0 \\ &\quad + s^2x(2r^3x - s^3 + x^2)W_0^2W_1 \end{aligned}$$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{k+1}^3$

(c):

$$\sum_{k=1}^n x^k W_{-k}^2 W_{-k+1} = \frac{\Lambda_6}{(-s^3 + x^2 + rsx)(r^3 x + s^3 - x^2 + 3rsx)}$$

$$\begin{aligned} \Lambda_6 &= rx^{n+2}(-x+rs)W_{-n+1}^3 + rs^3x^{n+1}(-x+rs)W_{-n}^3 + sx^{n+1}(-2r^3x+s^3-x^2)W_{-n+1}^2W_{-n} \\ &\quad + x^{n+1}(-2rs^4+s^3x+r^3x^2-x^3+2rsx^2+r^4sx)W_{-n}^2W_{-n+1} + rx^2(x-rs)W_1^3 \\ &\quad + rs^3x(x-rs)W_0^3 + sx(2r^3x-s^3+x^2)W_1^2W_0 \\ &\quad + x(2rs^4-s^3x-r^3x^2+x^3-2rsx^2-r^4sx)W_0^2W_1 \end{aligned}$$

Proof. Using the recurrence relation

$$W_{-n+2} = rW_{-n+1} + sW_{-n} \Rightarrow W_{-n} = -\frac{r}{s}W_{-n+1} + \frac{1}{s}W_{-n+2}$$

i.e.

$$sW_{-n} = W_{-n+2} - rW_{-n+1}$$

we obtain

$$s^3W_{-n}^3 = W_{-n+2}^3 - 3rW_{-n+2}^2W_{-n+1} + 3r^2W_{-n+1}^2W_{-n+2} - r^3W_{-n+1}^3$$

and so

$$\begin{aligned} s^3x^nW_{-n}^3 &= x^nW_{-n+2}^3 - 3rx^nW_{-n+2}^2W_{-n+1} + 3r^2x^nW_{-n+1}^2W_{-n+2} - r^3x^nW_{-n+1}^3 \\ s^3x^{n-1}W_{-n+1}^3 &= x^{n-1}W_{-n+3}^3 - 3rx^{n-1}W_{-n+3}^2W_{-n+2} + 3r^2x^{n-1}W_{-n+2}^2W_{-n+3} - r^3x^{n-1}W_{-n+2}^3 \\ s^3x^{n-2}W_{-n+2}^3 &= x^{n-2}W_{-n+4}^3 - 3rx^{n-2}W_{-n+4}^2W_{-n+3} + 3r^2x^{n-2}W_{-n+3}^2W_{-n+4} - r^3x^{n-2}W_{-n+3}^3 \\ &\vdots \\ s^3x^3W_{-3}^3 &= x^3W_{-1}^3 - 3rx^3W_{-1}^2W_{-2} + 3r^2x^3W_{-2}^2W_{-1} - r^3x^3W_{-2}^3 \\ s^3x^2W_{-2}^3 &= x^2W_0^3 - 3rx^2W_0^2W_{-1} + 3r^2x^2W_{-1}^2W_0 - r^3x^2W_{-1}^3 \\ s^3x^1W_{-1}^3 &= x^1W_1^3 - 3rx^1W_1^2W_0 + 3r^2x^1W_0^2W_1 - r^3x^1W_0^3 \end{aligned}$$

If we add the above equations by side by, we get

$$\begin{aligned} s^3\left(\sum_{k=1}^n x^k W_{-k}^3\right) &= (-x^{n+1}W_{-n+1}^3 - x^{n+2}W_{-n}^3 + x^1W_1^3 + x^2W_0^3 + x^2\sum_{k=1}^n x^k W_{-k}^3) \\ &\quad - 3r(-x^{n+1}W_{-n+1}^2W_{-n} + x^1W_1^2W_0 + x^1\sum_{k=1}^n x^k W_{-k+1}^2W_{-k}) \\ &\quad + 3r^2(-x^{n+1}W_{-n}^2W_{-n+1} + x^1W_0^2W_1 + x^1\sum_{k=1}^n x^k W_{-k}^2W_{-k+1}) \\ &\quad - r^3(-x^{n+1}W_{-n}^3 + x^1W_0^3 + x^1\sum_{k=1}^n x^k W_{-k}^3) \end{aligned} \quad (3.1)$$

Next we calculate $\sum_{k=1}^n x^k W_{-k+1}^2 W_{-k}$. Again using the recurrence relation

$$W_{-n+2} = rW_{-n+1} + sW_{-n} \Rightarrow W_{-n} = -\frac{r}{s}W_{-n+1} + \frac{1}{s}W_{-n+2}$$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{k+1}^3$

i.e.

$$sW_{-n} = W_{-n+2} - rW_{-n+1}$$

we obtain

$$sW_{-n+1}^2 W_{-n} = W_{-n+1}^2 W_{-n+2} - rW_{-n+1}^3$$

and so

$$\begin{aligned} sx^n W_{-n+1}^2 W_{-n} &= x^n W_{-n+1}^2 W_{-n+2} - rx^n W_{-n+1}^3 \\ sx^{n-1} W_{-n+2}^2 W_{-n+1} &= x^{n-1} W_{-n+2}^2 W_{-n+3} - rx^{n-1} W_{-n+2}^3 \\ sx^{n-2} W_{-n+3}^2 W_{-n+2} &= x^{n-2} W_{-n+3}^2 W_{-n+4} - rx^{n-2} W_{-n+3}^3 \\ &\vdots \\ sx^3 W_{-2}^2 W_{-3} &= x^3 W_{-2}^2 W_{-1} - rx^3 W_{-2}^3 \\ sx^2 W_{-1}^2 W_{-2} &= x^2 W_{-1}^2 W_0 - rx^2 W_{-1}^3 \\ sx^1 W_0^2 W_{-1} &= x^1 W_0^2 W_1 - rx^1 W_0^3 \end{aligned}$$

If we add the above equations by side by, we get

$$\begin{aligned} s \sum_{k=1}^n x^k W_{-k+1}^2 W_{-k} &= (-x^{n+1} W_{-n}^2 W_{-n+1} + x^1 W_0^2 W_1 + x^1 \sum_{k=1}^n x^k W_{-k}^2 W_{-k+1}) \\ &\quad - r(-x^{n+1} W_{-n}^3 + x^1 W_0^3 + x^1 \sum_{k=1}^n x^k W_{-k}^3) \end{aligned} \quad (3.2)$$

Next we calculate $\sum_{k=1}^n x^k W_{-k+1}^2 W_{-k}$. Again using the recurrence relation

$$W_{-n+2} = rW_{-n+1} + sW_{-n}$$

i.e.

$$sW_{-n} = W_{-n+2} - rW_{-n+1}$$

we obtain

$$\begin{aligned} s^2 W_{-n}^2 &= W_{-n+2}^2 - 2rW_{-n+2}W_{-n+1} + r^2 W_{-n+1}^2 \\ \Rightarrow s^2 W_{-n}^2 W_{-n+1} &= W_{-n+2}^2 W_{-n+1} - 2rW_{-n+1}^2 W_{-n+2} + r^2 W_{-n+1}^3 \end{aligned}$$

and so

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_k^3$

$$\begin{aligned}
 s^2 x^n W_{-n}^2 W_{-n+1} &= x^n W_{-n+2}^2 W_{-n+1} - 2rx^n W_{-n+1}^2 W_{-n+2} + r^2 x^n W_{-n+1}^3 \\
 s^2 x^{n-1} W_{-n+1}^2 W_{-n+2} &= x^{n-1} W_{-n+3}^2 W_{-n+2} - 2rx^{n-1} W_{-n+2}^2 W_{-n+3} + r^2 x^{n-1} W_{-n+2}^3 \\
 s^2 x^{n-2} W_{-n+2}^2 W_{-n+3} &= x^{n-2} W_{-n+4}^2 W_{-n+3} - 2rx^{n-2} W_{-n+3}^2 W_{-n+4} + r^2 x^{n-2} W_{-n+3}^3 \\
 &\vdots \\
 s^2 x^3 W_{-3}^2 W_{-2} &= x^3 W_{-1}^2 W_{-2} - 2rx^3 W_{-2}^2 W_{-1} + r^2 x^3 W_{-2}^3 \\
 s^2 x^2 W_{-2}^2 W_{-1} &= x^2 W_0^2 W_{-1} - 2rx^2 W_{-1}^2 W_0 + r^2 x^2 W_{-1}^3 \\
 s^2 x^1 W_{-1}^2 W_0 &= x^1 W_1^2 W_0 - 2rx^1 W_0^2 W_1 + r^2 x^1 W_0^3
 \end{aligned}$$

If we add the above equations by side by, we get

$$\begin{aligned}
 s^2 \sum_{k=1}^n x^k W_{-k}^2 W_{-k+1} &= (-x^{n+1} W_{-n+1}^2 W_{-n} + x^1 W_1^2 W_0 + x^1 \sum_{k=1}^n x^k W_{-k+1}^2 W_{-k}) \\
 &\quad - 2r(-x^{n+1} W_{-n}^2 W_{-n+1} + x^1 W_0^2 W_1 + x^1 \sum_{k=1}^n x^k W_{-k}^2 W_{-k+1}) \\
 &\quad + r^2(-x^{n+1} W_{-n}^3 + x^1 W_0^3 + x^1 \sum_{k=1}^n x^k W_{-k}^3)
 \end{aligned} \tag{3.3}$$

Then, solving the system (3.1)-(3.2)-(3.3), the required results of (a),(b) and (c) follow.

3.1. The case $x = 1$. See [19] for the case $x = 1$.

3.2. The case $x = -1$. In this subsection we consider the special case $x = -1$.

Taking $x = -1, r = s = 1$ in Theorem 3.1 (a),(b) and (c), we obtain the following proposition.

PROPOSITION 3.2. *If $x = -1, r = s = 1$ then for $n \geq 1$ we have the following formulas:*

- (a): $\sum_{k=1}^n (-1)^k W_{-k}^3 = \frac{1}{2}((-1)^n (W_{-n+1}^3 + W_{-n}^3 - 3W_{-n}^2 W_{-n+1}) - W_1^3 - W_0^3 + 3W_0^2 W_1).$
- (b): $\sum_{k=1}^n (-1)^k W_{-k+1}^2 W_{-k} = \frac{1}{2}((-1)^n (W_{-n+1}^2 W_{-n} - W_{-n}^2 W_{-n+1}) - W_1^2 W_0 + W_0^2 W_1).$
- (c): $\sum_{k=1}^n (-1)^k W_{-k}^2 W_{-k+1} = \frac{1}{2}((-1)^n (W_{-n+1}^3 - W_{-n}^3 - W_{-n+1}^2 W_{-n}) - W_1^3 + W_0^3 + W_1^2 W_0).$

From the above proposition, we have the following corollary which gives sum formulas of Fibonacci numbers (take $W_n = F_n$ with $F_0 = 0, F_1 = 1$).

COROLLARY 3.3. *For $n \geq 1$, Fibonacci numbers have the following properties.*

- (a): $\sum_{k=1}^n (-1)^k F_{-k}^3 = \frac{1}{2}((-1)^n (F_{-n+1}^3 + F_{-n}^3 - 3F_{-n}^2 F_{-n+1}) - 1).$
- (b): $\sum_{k=1}^n (-1)^k F_{-k+1}^2 F_{-k} = \frac{1}{2}((-1)^n (F_{-n+1}^2 F_{-n} - F_{-n}^2 F_{-n+1})).$
- (c): $\sum_{k=1}^n (-1)^k F_{-k}^2 F_{-k+1} = \frac{1}{2}((-1)^n (F_{-n+1}^3 - F_{-n}^3 - F_{-n+1}^2 F_{-n}) - 1).$

Taking $W_n = L_n$ with $L_0 = 2, L_1 = 1$ in the last proposition, we have the following corollary which presents sum formulas of Lucas numbers.

COROLLARY 3.4. *For $n \geq 1$, Lucas numbers have the following properties.*

- (a): $\sum_{k=1}^n (-1)^k L_{-k}^3 = \frac{1}{2}((-1)^n (L_{-n+1}^3 + L_{-n}^3 - 3L_{-n}^2 L_{-n+1}) + 3).$
- (b): $\sum_{k=1}^n (-1)^k L_{-k+1}^2 L_{-k} = \frac{1}{2}((-1)^n (L_{-n+1}^2 L_{-n} - L_{-n}^2 L_{-n+1}) + 2).$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_k^3$

$$(c): \sum_{k=1}^n (-1)^k L_{-k}^2 L_{-k+1} = \frac{1}{2}((-1)^n (L_{-n+1}^3 - L_{-n}^3 - L_{-n+1}^2 L_{-n}) + 9).$$

Taking $x = -1, r = 2, s = 1$ in Theorem 3.1 (a),(b) and (c), we obtain the following proposition.

PROPOSITION 3.5. *If $x = -1, r = 2, s = 1$ then for $n \geq 1$ we have the following formulas:*

$$\begin{aligned} (a): \sum_{k=1}^n (-1)^k W_{-k}^3 &= \frac{1}{14}((-1)^n (2W_{-n+1}^3 + 12W_{-n}^3 - 3W_{-n+1}^2 W_{-n} - 9W_{-n}^2 W_{-n+1}) - 2W_1^3 - 12W_0^3 + 3W_1^2 W_0 + 9W_0^2 W_1). \\ (b): \sum_{k=1}^n (-1)^k W_{-k+1}^2 W_{-k} &= \frac{1}{14}((-1)^n (W_{-n+1}^3 - W_{-n}^3 + 2W_{-n+1}^2 W_{-n} - 8W_{-n}^2 W_{-n+1}) - W_1^3 + W_0^3 - 2W_1^2 W_0 + 8W_0^2 W_1). \\ (c): \sum_{k=1}^n (-1)^k W_{-k}^2 W_{-k+1} &= \frac{1}{14}((-1)^n (3W_{-n+1}^3 - 3W_{-n}^3 + 4W_{-n}^2 W_{-n+1} - 8W_{-n+1}^2 W_{-n}) - 3W_1^3 + 3W_0^3 + 8W_1^2 W_0 - 4W_0^2 W_1). \end{aligned}$$

From the last proposition, we have the following corollary which gives sum formulas of Pell numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1$).

COROLLARY 3.6. *For $n \geq 1$, Pell numbers have the following properties.*

$$\begin{aligned} (a): \sum_{k=1}^n (-1)^k P_{-k}^3 &= \frac{1}{14}((-1)^n (2P_{-n+1}^3 + 12P_{-n}^3 - 3P_{-n+1}^2 P_{-n} - 9P_{-n}^2 P_{-n+1}) - 2). \\ (b): \sum_{k=1}^n (-1)^k P_{-k+1}^2 P_{-k} &= \frac{1}{14}((-1)^n (P_{-n+1}^3 - P_{-n}^3 + 2P_{-n+1}^2 P_{-n} - 8P_{-n}^2 P_{-n+1}) - 1). \\ (c): \sum_{k=1}^n (-1)^k P_{-k}^2 P_{-k+1} &= \frac{1}{14}((-1)^n (3P_{-n+1}^3 - 3P_{-n}^3 + 4P_{-n}^2 P_{-n+1} - 8P_{-n+1}^2 P_{-n}) - 3). \end{aligned}$$

Taking $P_n = Q_n$ with $Q_0 = 2, Q_1 = 2$ in the last proposition, we have the following corollary which presents sum formulas of Pell-Lucas numbers.

COROLLARY 3.7. *For $n \geq 1$, Pell-Lucas numbers have the following properties.*

$$\begin{aligned} (a): \sum_{k=1}^n (-1)^k Q_{-k}^3 &= \frac{1}{14}((-1)^n (2Q_{-n+1}^3 + 12Q_{-n}^3 - 3Q_{-n+1}^2 Q_{-n} - 9Q_{-n}^2 Q_{-n+1}) - 16). \\ (b): \sum_{k=1}^n (-1)^k Q_{-k+1}^2 Q_{-k} &= \frac{1}{14}((-1)^n (Q_{-n+1}^3 - Q_{-n}^3 + 2Q_{-n+1}^2 Q_{-n} - 8Q_{-n}^2 Q_{-n+1}) + 48). \\ (c): \sum_{k=1}^n (-1)^k Q_{-k}^2 Q_{-k+1} &= \frac{1}{14}((-1)^n (3Q_{-n+1}^3 - 3Q_{-n}^3 + 4Q_{-n}^2 Q_{-n+1} - 8Q_{-n+1}^2 Q_{-n}) + 32). \end{aligned}$$

If $x = -1, r = 1, s = 2$ then $(-s^3 + x^2 + rsx)(r^3 x + s^3 - x^2 + 3rsx) = 0$ so we can't use Theorem 3.1 directly. But we can give another method to find $\sum_{k=1}^n (-1)^k W_{-k}^3$, $\sum_{k=1}^n (-1)^k W_{-k+1}^2 W_{-k}$ and $\sum_{k=1}^n (-1)^k W_{-k}^2 W_{-k+1}$ by using Theorem 3.1.

THEOREM 3.8. *If $x = -1, r = 1, s = 2$ then for $n \geq 1$ we have the following formulas:*

$$\begin{aligned} (a): \sum_{k=1}^n (-1)^k W_{-k}^3 &= \frac{1}{27}((-1)^n (-(n-1)W_{-n+1}^3 + (8n+11)W_{-n}^3 + 2(3n+2)W_{-n+1}^2 W_{-n} - 4(3n+4)W_{-n}^2 W_{-n+1}) - W_1^3 - 11W_0^3 - 4W_1^2 W_0 + 16W_0^2 W_1). \\ (b): \sum_{k=1}^n (-1)^k W_{-k+1}^2 W_{-k} &= \frac{1}{81}((-1)^n (-(3n+5)W_{-n+1}^3 + 8(3n+2)W_{-n}^3 + 3(6n+11)W_{-n+1}^2 W_{-n} - 36(n+1)W_{-n}^2 W_{-n+1}) + 5W_1^3 - 16W_0^3 - 33W_1^2 W_0 + 36W_0^2 W_1). \\ (c): \sum_{k=1}^n (-1)^k W_{-k}^2 W_{-k+1} &= \frac{1}{81}((-1)^n ((3n+7)W_{-n+1}^3 - 8(3n+4)W_{-n}^3 - 18(n+1)W_{-n+1}^2 W_{-n} + 3(12n+11)W_{-n}^2 W_{-n+1}) - 7W_1^3 + 32W_0^3 - 33W_0^2 W_1 + 18W_1^2 W_0). \end{aligned}$$

Proof.

(a): We use Theorem 3.1 (a). If we set $r = 1, s = 2$ in Theorem 3.1 (a) then we have

$$\sum_{k=1}^n x^k W_{-k}^3 = \frac{f_4(x)}{(-x^2 + 7x + 8)(x^2 + 2x - 8)}$$

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_k^3$

where

$$f_4(x) = x^{n+1}(-x^2+4x+8)W_{-n+1}^3 + x^{n+1}(-x^3+5x^2+22x-8)W_{-n}^3 - 6x^{n+1}(x+4)W_{-n+1}^2W_{-n} - 12x^{n+1}(x-2)W_{-n}^2W_{-n+1} - x(-x^2+4x+8)W_1^3 - x(-x^3+5x^2+22x-8)W_0^3 + 6x(x+4)W_1^2W_0 + 12x(x-2)W_0^2W_1.$$

For $x = -1$, the right hand side of the above sum formulas is an indeterminate form. Now, we can use L'Hospital rule. Then we get

$$\begin{aligned} \sum_{k=1}^n (-1)^k W_{-k}^3 &= \frac{\frac{d}{dx}(f_4(x))}{\frac{d}{dx}((-x^2+7x+8)(x^2+2x-8))} \Big|_{x=-1} \\ &= \frac{1}{27}((-1)^n(-(n-1)W_{-n+1}^3 + (8n+11)W_{-n}^3 + 2(3n+2)W_{-n+1}^2W_{-n} \\ &\quad - 4(3n+4)W_{-n}^2W_{-n+1}) - W_1^3 - 11W_0^3 - 4W_1^2W_0 + 16W_0^2W_1). \end{aligned}$$

(b): We use Theorem 3.1 (b). If we set $r = 1, s = 2$ in Theorem 3.1 (b) then we have

$$\sum_{k=1}^n x^k W_{-k+1}^2 W_{-k} = \frac{f_5(x)}{(-x^2+7x+8)(x^2+2x-8)}$$

where

$$f_5(x) = -x^{n+2}(x+4)W_{-n+1}^3 - 8x^{n+1}(x+4)W_{-n}^3 + x^{n+2}(-x^2+x+20)W_{-n+1}^2W_{-n} - 4x^{n+1}(x^2+2x-8)W_{-n}^2W_{-n+1} + x^2(x+4)W_1^3 + 8x(x+4)W_0^3 - x^2(-x^2+x+20)W_1^2W_0 + 4x(x^2+2x-8)W_0^2W_1.$$

For $x = -1$, the right hand side of the above sum formulas is an indeterminate form. Now, we can use L'Hospital rule. Then we get

$$\begin{aligned} \sum_{k=1}^n (-1)^k W_{-k+1}^2 W_{-k} &= \frac{\frac{d}{dx}(f_5(x))}{\frac{d}{dx}((-x^2+7x+8)(x^2+2x-8))} \Big|_{x=-1} \\ &= \frac{1}{81}((-1)^n(-(3n+5)W_{-n+1}^3 + 8(3n+2)W_{-n}^3 + 3(6n+11)W_{-n+1}^2W_{-n} \\ &\quad - 36(n+1)W_{-n}^2W_{-n+1}) + 5W_1^3 - 16W_0^3 - 33W_1^2W_0 + 36W_0^2W_1) \end{aligned}$$

(c): We use Theorem 3.1 (c). If we set $r = 1, s = 2$ in Theorem 3.1 (c) then we have

$$\sum_{k=1}^n x^k W_{-k}^2 W_{-k+1} = \frac{f_6(x)}{(-x^2+7x+8)(x^2+2x-8)}$$

where

$$f_6(x) = (-x^{n+2}(x-2)W_{-n+1}^3 - 8x^{n+1}(x-2)W_{-n}^3 - 2x^{n+1}(x^2+2x-8)W_{-n+1}^2W_{-n} + x^{n+1}(-x^3+5x^2+10x-32)W_{-n}^2W_{-n+1} + x^2(x-2)W_1^3 + 8x(x-2)W_0^3 + 2x(x^2+2x-8)W_1^2W_0 - x(-x^3+5x^2+10x-32)W_0^2W_1).$$

For $x = -1$, the right hand side of the above sum formulas is an indeterminate form. Now, we can use L'Hospital rule. Then we get

$$\begin{aligned} \sum_{k=1}^n (-1)^k W_{-k}^2 W_{-k+1} &= \frac{\frac{d}{dx}(f_6(x))}{\frac{d}{dx}((-x^2+7x+8)(x^2+2x-8))} \Big|_{x=-1} \\ &= \frac{1}{81}((-1)^n((3n+7)W_{-n+1}^3 - 8(3n+4)W_{-n}^3 - 18(n+1)W_{-n+1}^2W_{-n} \\ &\quad + 3(12n+11)W_{-n}^2W_{-n+1}) - 7W_1^3 + 32W_0^3 - 33W_0^2W_1 + 18W_1^2W_0) \end{aligned}$$

From the last theorem, we have the following corollary which gives sum formula of Jacobsthal numbers (take $W_n = J_n$ with $J_0 = 0, J_1 = 1$).

COROLLARY 3.9. For $n \geq 1$, Jacobsthal numbers have the following properties:

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_{\tilde{k}}^3$

- (a): $\sum_{k=1}^n (-1)^k J_{-k}^3 = \frac{1}{27}((-1)^n(-(n-1)J_{-n+1}^3 + (8n+11)J_{-n}^3 + 2(3n+2)J_{-n+1}^2 J_{-n} - 4(3n+4)J_{-n}^2 J_{-n+1}) - 1)$.
- (b): $\sum_{k=1}^n (-1)^k J_{-k+1}^2 J_{-k} = \frac{1}{81}((-1)^n(-(3n+5)J_{-n+1}^3 + 8(3n+2)J_{-n}^3 + 3(6n+11)J_{-n+1}^2 J_{-n} - 36(n+1)J_{-n}^2 J_{-n+1}) + 5)$.
- (c): $\sum_{k=1}^n (-1)^k J_{-k}^2 J_{-k+1} = \frac{1}{81}((-1)^n((3n+7)J_{-n+1}^3 - 8(3n+4)J_{-n}^3 - 18(n+1)J_{-n+1}^2 J_{-n} + 3(12n+11)J_{-n}^2 J_{-n+1}) - 7)$.

Taking $J_n = j_n$ with $j_0 = 2, j_1 = 1$ in the last theorem, we have the following corollary which presents sum formulas of Jacobsthal-Lucas numbers.

COROLLARY 3.10. For $n \geq 1$, Jacobsthal-Lucas numbers have the following properties:

- (a): $\sum_{k=1}^n (-1)^k j_{-k}^3 = \frac{1}{27}((-1)^n(-(n-1)j_{-n+1}^3 + (8n+11)j_{-n}^3 + 2(3n+2)j_{-n+1}^2 j_{-n} - 4(3n+4)j_{-n}^2 j_{-n+1}) - 33)$.
- (b): $\sum_{k=1}^n (-1)^k j_{-k+1}^2 j_{-k} = \frac{1}{81}((-1)^n(-(3n+5)j_{-n+1}^3 + 8(3n+2)j_{-n}^3 + 3(6n+11)j_{-n+1}^2 j_{-n} - 36(n+1)j_{-n}^2 j_{-n+1}) - 45)$.
- (c): $\sum_{k=1}^n (-1)^k j_{-k}^2 j_{-k+1} = \frac{1}{81}((-1)^n((3n+7)j_{-n+1}^3 - 8(3n+4)j_{-n}^3 - 18(n+1)j_{-n+1}^2 j_{-n} + 3(12n+11)j_{-n}^2 j_{-n+1}) + 153)$.

3.3. The case $x = 1 + i$. In this subsection we consider the special case $x = 1 + i$.

Taking $x = 1 + i, r = s = 1$ in Theorem 3.1 (a),(b) and (c), we obtain the following proposition.

PROPOSITION 3.11. If $x = 1 + i, r = s = 1$ then for $n \geq 1$ we have the following formulas:

- (a): $\sum_{k=1}^n (1+i)^k F_{-k}^3 = \frac{1}{-6+15i}((1+i)^n(3(1+i)F_{-n+1}^3 + (-3+15i)F_{-n}^3 - (3+9i)F_{-n+1}^2 F_{-n} + (3-3i)F_{-n}^2 F_{-n+1}) - (3+3i)F_1^3 + (3-15i)F_0^3 + (3+9i)F_1^2 F_0 - (3-3i)F_0^2 F_1)$.
- (b): $\sum_{k=1}^n (1+i)^k F_{-k+1}^2 F_{-k} = \frac{1}{-6+15i}((1+i)^n((2-4i)F_{-n+1}^3 - (1+3i)F_{-n}^3 + (2+10i)F_{-n+1}^2 F_{-n} + (3-5i)F_{-n}^2 F_{-n+1}) - (2-4i)F_1^3 + (1+3i)F_0^3 - (2+10i)F_1^2 F_0 - (3-5i)F_0^2 F_1)$.
- (c): $\sum_{k=1}^n (1+i)^k F_{-k}^2 F_{-k+1} = \frac{1}{-6+15i}((1+i)^n(2F_{-n+1}^3 + (1-i)F_{-n}^3 + (3-5i)F_{-n+1}^2 F_{-n} + (-4+8i)F_{-n}^2 F_{-n+1}) - 2F_1^3 - (1-i)F_0^3 - (3-5i)F_1^2 F_0 + (4-8i)F_0^2 F_1)$.

From the above proposition, we have the following corollary which gives sum formulas of Fibonacci numbers (take $W_n = F_n$ with $F_0 = 0, F_1 = 1$).

COROLLARY 3.12. For $n \geq 1$, Fibonacci numbers have the following properties.

- (a): $\sum_{k=1}^n (1+i)^k F_{-k}^3 = \frac{1}{-6+15i}((1+i)^n(3(1+i)F_{-n+1}^3 + (-3+15i)F_{-n}^3 - (3+9i)F_{-n+1}^2 F_{-n} + (3-3i)F_{-n}^2 F_{-n+1}) - 3-3i)$.
- (b): $\sum_{k=1}^n (1+i)^k F_{-k+1}^2 F_{-k} = \frac{1}{-6+15i}((1+i)^n((2-4i)F_{-n+1}^3 - (1+3i)F_{-n}^3 + (2+10i)F_{-n+1}^2 F_{-n} + (3-5i)F_{-n}^2 F_{-n+1}) - 2+4i)$.
- (c): $\sum_{k=1}^n (1+i)^k F_{-k}^2 F_{-k+1} = \frac{1}{-6+15i}((1+i)^n(2F_{-n+1}^3 + (1-i)F_{-n}^3 + (3-5i)F_{-n+1}^2 F_{-n} + (-4+8i)F_{-n}^2 F_{-n+1}) - 2)$.

Taking $W_n = L_n$ with $L_0 = 2, L_1 = 1$ in the last proposition, we have the following corollary which presents sum formulas of Lucas numbers.

A STUDY ON SUMS OF CUBES OF GENERALIZED FIBONACCI NUMBERS: CLOSED FORMULAS OF $\sum_{k=0}^n x^k W_k^3$ AND $\sum_{k=1}^n x^k W_k^3$

COROLLARY 3.13. For $n \geq 1$, Lucas numbers have the following properties.

- (a): $\sum_{k=1}^n (1+i)^k L_{-k}^3 = \frac{1}{-6+15i} ((1+i)^n (3(1+i)L_{-n+1}^3 + (-3+15i)L_{-n}^3 - (3+9i)L_{-n+1}^2 L_{-n} + (3-3i)L_{-n}^2 L_{-n+1}) + 15 - 93i).$
- (b): $\sum_{k=1}^n (1+i)^k L_{-k+1}^2 L_{-k} = \frac{1}{-6+15i} ((1+i)^n ((2-4i)L_{-n+1}^3 - (1+3i)L_{-n}^3 + (2+10i)L_{-n+1}^2 L_{-n} + (3-5i)L_{-n}^2 L_{-n+1}) - 10 + 28i).$
- (c): $\sum_{k=1}^n (1+i)^k L_{-k}^2 L_{-k+1} = \frac{1}{-6+15i} ((1+i)^n (2L_{-n+1}^3 + (1-i)L_{-n}^3 + (3-5i)L_{-n+1}^2 L_{-n} + (-4+8i)L_{-n}^2 L_{-n+1}) - 14i).$

Corresponding sums of the other second order linear sequences can be calculated similarly when $x = 1 + i$.

References

- [1] Čerin, Z., Formulae for sums of Jacobsthal–Lucas numbers, *Int. Math. Forum*, 2(40), 1969–1984, 2007.
- [2] Čerin, Z., Sums of Squares and Products of Jacobsthal Numbers. *Journal of Integer Sequences*, 10, Article 07.2.5, 2007.
- [3] Chen, L., Wang, X., The Power Sums Involving Fibonacci Polynomials and Their Applications, *Symmetry*, 11, 2019, doi.org/10.3390/sym11050635.
- [4] Frontczak, R., Sums of powers of Fibonacci and Lucas numbers: A new bottom-up approach, *Notes on Number Theory and Discrete Mathematics*, 24(2), 94–103, 2018.
- [5] Frontczak, R., Sums of Cubes Over Odd-Index Fibonacci Numbers, *Integers*, 18, 2018.
- [6] Gnanam, A., Anitha, B., Sums of Squares Jacobsthal Numbers. *IOSR Journal of Mathematics*, 11(6), 62-64. 2015.
- [7] Horadam, A.F., A Generalized Fibonacci Sequence, *American Mathematical Monthly*, Vol. 68, pp. 455-459, 1961.
- [8] Horadam, A. F., Basic Properties of a Certain Generalized Sequence of Numbers, *Fibonacci Quarterly* 3.3, 161-176, 1965.
- [9] Horadam, A. F., Special Properties of The Sequence $w_n(a, b; p, q)$, *Fibonacci Quarterly*, Vol. 5, No. 5, pp. 424-434, 1967.
- [10] Horadam, A. F., Generating functions for powers of a certain generalized sequence of numbers. *Duke Math. J* 32, 437-446, 1965.
- [11] Kiliç, E., Taşçi, D., The Linear Algebra of The Pell Matrix, *Boletín de la Sociedad Matemática Mexicana*, 3(11), 2005.
- [12] Kılıc, E., Sums of the squares of terms of sequence $\{u_n\}$, *Proc. Indian Acad. Sci. (Math. Sci.)* 118(1), 27–41, 2008.
- [13] Prodinger, H., Sums of Powers of Fibonacci Polynomials, *Proc. Indian Acad. Sci. (Math. Sci.)*, 119(5), 567-570, 2009.
- [14] Prodinger, H., Selkirk, S.J., Sums of Squares of Tetranacci Numbers: A Generating Function Approach, 2019, <http://arxiv.org/abs/1906.08336v1>.
- [15] Raza, Z., Riaz, M., Ali, M.A., Some Inequalities on the Norms of Special Matrices with Generalized Tribonacci and Generalized Pell-Padovan Sequences, *arXiv*, 2015, <http://arxiv.org/abs/1407.1369v2>
- [16] Schumacher, R., How to sum the squares of the Tetranacci numbers and the Fibonacci m-step numbers. *Fibonacci Quarterly*, 57:168–175, 2019.
- [17] Sloane, N.J.A., The on-line encyclopedia of integer sequences. Available: <http://oeis.org/>
- [18] Soykan, Y. Closed Formulas for the Sums of Squares of Generalized Fibonacci Numbers, *Asian Journal of Advanced Research and Reports*, 9(1), 23-39, 2020. <https://doi.org/10.9734/ajarr/2020/v9i130212>
- [19] Soykan Y., Closed Formulas for the Sums of Cubes of Generalized Fibonacci Numbers: Closed Formulas of $\sum_{k=0}^n W_k^3$ and $\sum_{k=1}^n W_k^3$, *Archives of Current Research International*, 20(2), 58-69, 2020. DOI: 10.9734/ACRI/2020/v20i230177
- [20] Soykan, Y., A Closed Formula for the Sums of Squares of Generalized Tribonacci numbers, *Journal of Progressive Research in Mathematics*, Accepted.
- [21] Wamiliana., Suharsono., Kristanto, P. E., Counting the sum of cubes for Lucas and Gibonacci Numbers, *Science and Technology Indonesia*, 4(2), 31-35, 2019.