

The Forrest Gump approach to prevent severe COVID-19 – reverse the predisposing pro-inflammatory state with exercise

Chris Kenyon^{1,2*}

¹HIV/STI Unit, Institute of Tropical Medicine, Antwerp, Belgium; ²Division of Infectious Diseases and HIV Medicine, University of Cape Town, Anzio Road, Observatory 7700, South Africa;

**Corresponding author. HIV/STI Unit, Institute of Tropical Medicine, Antwerp, 2000, Belgium. Tel: +32 3 2480796; Fax: +32 3 2480831; E-mail: ckenyon@itg.be*

Word Count: 914

Keywords: Exercise; Angiotensin-converting enzyme; SARS-CoV-2; COVID-19; Ecology

25 **Abstract**

26 We hypothesize that the reason severe COVID-19 is closely linked to a limited number of
27 conditions such as obesity, hypertension, older age and diabetes is that these conditions are
28 characterized by a pro-inflammatory state in general and an angiotensin converting enzyme-
29 1-bias in particular. We review recent evidence that this pro-inflammatory state can be
30 reversed via regular exercise and a healthy diet. Studies are suggested that could test this
31 hypothesis.

Why do only a small proportion of those infected with SARS-CoV-2 get severe disease? Most infections are pauci- or asymptomatic [1]. Studies show that the vast majority of those who get severe disease have predisposing conditions such as hypertension, obesity, cardiovascular disease, diabetes or older age [2-5]. For example, in New York, of the 5700 patients sick enough from coronavirus to need hospitalization, 57% had high blood pressure, 42% were obese and 34% had diabetes [2]. However, only a small set of chronic diseases predispose to severe disease. Diseases like asthma and HIV do not [2, 5]. Why should hypertension and cardiovascular disease be such prominent risk factors for a severe infectious disease? Increasing evidence suggests that the reason why only certain chronic diseases predispose to severe disease is that they are associated with a pro-inflammatory state and an imbalance in their ACE1/ACE2 axes in particular [6, 7]. These are linked enzyme pathways that in health are in a state of balance between the pro- and anti-inflammatory effects of ACE1 and ACE2, respectively [7]. The chronic diseases associated with COVID-19 are typified by a dominance of the ACE1 axis which promotes vasoconstriction as well as inflammatory, oxidative and fibrotic tissue damage [6, 7]. A sedentary, excess-calorie-lifestyle results in the same proinflammatory ACE1 bias [8]. So does aging without staying fit [9].

This low-grade tissue inflammation may be critical in determining who gets severe COVID-19 [6, 7]. As shown on the right side of Fig. 1, an unfit, overweight individual with a high blood pressure has an overactive ACE1 and an underactive ACE2 axis [6, 7]. This generates a proinflammatory state in their lungs [7]. SARS-CoV-2 infection in this setting may precipitate an excessive inflammatory reaction. Part of the reason the inflammatory response is so excessive is that SARS-CoV-2 uses ACE2 as its cellular receptor and in so doing dramatically reduces the expression of ACE2 [7]. The resulting exuberant inflammation fills the alveoli in

fluid (which is the key clinicopathological feature of severe COVID-19) before the immune system has a chance to mount an effective immune response to clear the virus [7]. The fit healthy individual on the left has a balanced ACE1/2 system with sufficient ACE2 and no tissue inflammation at baseline. Following infection, the virus depletes ACE2, but this is from a high baseline, and sufficient ACE2 remains to ensure that the inflammatory response is proportionate. The immune cells are not overwhelmed by non-specific inflammation and are able to effectively eliminate the virus [7].

The ACE-1 bias and excess inflammation can be reversed by exercise/diet

Whilst this is an over-simplification of a complex process it goes some way to explaining the association between severe COVID-19 and age, obesity, and hypertension. As one ages, becomes more obese or hypertensive one's ACE axis is biased in the ACE1 direction which then predisposes one to severe COVID-19 [6, 7, 9]. Other pro-inflammatory pathways are also activated [9]. These changes are however not inevitable with aging or irreversible. A large number of studies in humans and other animals have shown that aerobic exercise can fairly rapidly reverse this bias, regardless of one's age, sex or abdominal girth [10, 11]. Whilst the initial reversal of this bias occurs within minutes to hours of exercise, more profound reversal requires sustaining a regular program of exercise [10-12]. Exercise has also consistently been shown to be an effective way to reduce blood pressure in hypertensives [13]. A healthy, tailored diet can also play an important complementary role. As an example, randomized controlled trials have shown that certain diets can lead to remission of diabetes in up to 50% of participants [14, 15].

Testing the exercise/diet hypothesis

If this theory is correct, then regular exercise and a healthy diet would not reduce one's risk of getting infected with SARS-CoV-2, but they would reduce one's risk of getting severe disease. We must acknowledge that randomized clinical trials in high risk populations would need to be conducted before we can conclude that exercise and diet have this protective effect. As an alternative, the protective-effect-of-exercise/diet could be tested in animal models of severe COVID-19 [16]. Unfortunately, no such trials are currently registered on www.clinicaltrials.gov.

Whilst we wait for these studies it is worth recalling the good evidence for a broad array of other health benefits from exercise. One study for example found that every hour of running prolongs one's life by 7 hours [12]. If exercise protects against severe COVID-19 too, then this would be the cherry on the cake. The advice "run, Forrest, run!" saved the handicapped Forrest Gump from nasty bullies, cured his physical disability and paved the way for a new healthy lifestyle. In this time of widespread COVID-19 induced anxiety and sense of helplessness, advising people to exercise regularly will likewise accomplish more than preventing severe COVID-19. At a macro level, a number of authors have concluded that the current COVID-19 pandemic was ultimately caused by excessive human consumption that has resulted in extensive habitat destruction and spillover of SARS-CoV-2 into humans [17-19]. This is mirrored at a micro level by an imbalance between excessive consumption, massive ecological footprints and insufficient exercise [18, 20]. This imbalance extends to the extraordinary contemporary inequalities in wealth and opportunities between individuals and populations [20, 21]. These insights suggest that dealing with the current pandemic and preventing future pandemics and other health disasters will ultimately require rebalancing

103 each of these levels [18, 19, 21]. Each of us could start this process by donning our running
104 shoes more regularly.

105 **Figure 1.** Schematic representation of hypothesized link between being physically
106 unfit/normotensive and the risk of severe COVID-19, contrasting SARS-CoV-2 infection in the
107 alveolus of a fit, normotensive individual with balanced ACE1/2 axes (left) with an unfit,
108 hypertensive individual with an ACE1-biased proinflammatory alveolus (right). Inflammation
109 is represented by red and the pulmonary exudative fluid characteristic of severe COVID-19 is
110 depicted by yellow-green fluid.

References

1. Sutton D, Fuchs K, D'Alton M, Goffman D. Universal Screening for SARS-CoV-2 in Women Admitted for Delivery. *N Engl J Med*. 2020. Epub 2020/04/14. doi: 10.1056/NEJMc2009316. PubMed PMID: 32283004; PubMed Central PMCID: PMC67175422.

2. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA*. 2020.

3. Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *Jama*. 2020.

4. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama*. 2020;323(13):1239-42.

5. Mehra MR, Desai SS, Kuy S, Patel AN. Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *NEJM*. 2020. doi: 10.1056/NEJMoa2007621.

6. Henry BM, Vikse J, Benoit S, Favaloro EJ, Lippi G. Hyperinflammation and Derangement of Renin-Angiotensin-Aldosterone System in COVID-19: a novel hypothesis for clinically suspected hypercoagulopathy and microvascular immunothrombosis. *Clinica Chimica Acta*. 2020.

7. Sriram K, Insel PA. A hypothesis for pathobiology and treatment of COVID-19: the centrality of ACE1/ACE2 imbalance. *Br J Pharmacol*. 2020. Epub 2020/04/26. doi: 10.1111/bph.15082. PubMed PMID: 32333398.

8. Migdalis IN, Iliopoulou V, Kalogeropoulou K, Koutoulidis K, Samartzis M. Elevated serum levels of angiotensin-converting enzyme in patients with diabetic retinopathy. *South Med J*. 1990;83(4):425-7. Epub 1990/04/01. PubMed PMID: 2157294.

9. Ferrucci L, Fabbri E. Inflammageing: chronic inflammation in ageing, cardiovascular disease, and frailty. *Nat Rev Cardiol*. 2018;15(9):505-22. Epub 2018/08/02. doi: 10.1038/s41569-018-0064-2. PubMed PMID: 30065258; PubMed Central PMCID: PMC6146930.

10. Echeverria-Rodriguez O, Gallardo-Ortiz IA, Del Valle-Mondragon L, Villalobos-Molina R. Angiotensin-(1-7) Participates in Enhanced Skeletal Muscle Insulin Sensitivity After a Bout of Exercise. *J Endocr Soc*. 2020;4(2):bvaa007. Epub 2020/02/28. doi: 10.1210/jendso/bvaa007. PubMed PMID: 32104748; PubMed Central PMCID: PMC67039407.

11. Magalhaes DM, Nunes-Silva A, Rocha GC, Vaz LN, de Faria MHS, Vieira ELM, et al. Two protocols of aerobic exercise modulate the counter-regulatory axis of the renin-angiotensin system. *Heliyon*. 2020;6(1):e03208. Epub 2020/01/29. doi: 10.1016/j.heliyon.2020.e03208. PubMed PMID: 31989052; PubMed Central PMCID: PMC6970173.

12. Lee D-c, Brellenthin AG, Thompson PD, Sui X, Lee I-M, Lavie CJ. Running as a key lifestyle medicine for longevity. *Progress in cardiovascular diseases*. 2017;60(1):45-55.

13. Hegde SM, Solomon SD. Influence of Physical Activity on Hypertension and Cardiac Structure and Function. *Curr Hypertens Rep*. 2015;17(10):77. Epub 2015/08/19. doi: 10.1007/s11906-015-0588-3. PubMed PMID: 26277725; PubMed Central PMCID: PMC64624627.

14. Avenell A, Brown TJ, McGee M, Campbell MK, Grant AM, Broom J, et al. What interventions should we add to weight reducing diets in adults with obesity? A systematic

- 157 review of randomized controlled trials of adding drug therapy, exercise, behaviour therapy or
158 combinations of these interventions. *Journal of Human Nutrition and Dietetics*.
159 2004;17(4):293-316.
- 160 15. Lean ME, Leslie WS, Barnes AC, Brosnahan N, Thom G, McCombie L, et al. Primary
161 care-led weight management for remission of type 2 diabetes (DiRECT): an open-label,
162 cluster-randomised trial. *Lancet*. 2018;391(10120):541-51. Epub 2017/12/10. doi:
163 10.1016/S0140-6736(17)33102-1. PubMed PMID: 29221645.
- 164 16. Chan JF, Zhang AJ, Yuan S, Poon VK, Chan CC, Lee AC, et al. Simulation of the clinical
165 and pathological manifestations of Coronavirus Disease 2019 (COVID-19) in golden Syrian
166 hamster model: implications for disease pathogenesis and transmissibility. *Clin Infect Dis*.
167 2020. Epub 2020/03/28. doi: 10.1093/cid/ciaa325. PubMed PMID: 32215622.
- 168 17. Nabi G, Siddique R, Ali A, Khan S. Preventing bat-born viral outbreaks in future using
169 ecological interventions. *Environmental Research*. 2020.
- 170 18. Dunk JH, Jones DS, Capon A, Anderson WH. Human Health on an Ailing Planet -
171 Historical Perspectives on Our Future. *N Engl J Med*. 2019;381(8):778-82. Epub 2019/08/23.
172 doi: 10.1056/NEJMms1907455. PubMed PMID: 31433929.
- 173 19. Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, et al. Impacts of
174 biodiversity on the emergence and transmission of infectious diseases. *Nature*.
175 2010;468(7324):647-52. Epub 2010/12/03. doi: 10.1038/nature09575. PubMed PMID:
176 21124449.
- 177 20. Pickett KE, Wilkinson RG. Income inequality and health: a causal review. *Social science*
178 *& medicine*. 2015;128:316-26.
- 179 21. Ahmed F, Ahmed Ne, Pissarides C, Stiglitz J. Why inequality could spread COVID-19.
180 *The Lancet Public Health*. 2020.
- 181