

Communication

Expression pattern of the SARS-CoV-2 entry genes *ACE2* and *TMPRSS2* in the respiratory tract

Yichuan Liu^{1*}, Hui-Qi Qu^{1*}, Jingchun Qu¹, Lifeng Tian¹, Hakon Hakonarson^{1,2†}

¹ Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA

² Divisions of Human Genetics and Pulmonary Medicine, Department of Pediatrics, The Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

* Both authors contributed equally to this work.

† Corresponding authors:

Dr. Hakon Hakonarson

Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, United States of America, Department of Pediatrics, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America

Email: Hakonarson@chop.edu

Abstract: To address the expression pattern of the SARS-CoV-2 receptor *ACE2* and the viral priming protease, *TMPRSS2*, in the respiratory tract, this study investigated RNA sequencing transcriptome profiling of samples of airway and oral mucosa. As shown, *ACE2* has medium levels of expression in both small airway epithelium and masticatory mucosa, and high levels of expression in nasal epithelium. The expression of *ACE2* is low in mucosal associated invariant T (MAIT) cells, and can't be detected in alveolar macrophages. *TMPRSS2* is highly expressed in small airway epithelium and nasal epithelium, and has lower expression in masticatory mucosa. Our results provide the molecular basis that the nasal mucosa is the most susceptible locus in the respiratory tract for SARS-CoV-2 infection and consequently for subsequent droplet transmission and should be the focus for protection against SARS-CoV-2 infection.

Keywords: *ACE2*; COVID-19; SARS-CoV-2; *TMPRSS2*

1. Introduction

Concerning the pandemic of Coronavirus Disease 2019 (COVID-19), on Sep 26th, 2020, it has been diagnosed in 32.6 million people globally, causing 990,000 deaths. COVID-19 is caused by the infection of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). For this highly infectious and deadly disease, there is no effective anti-viral treatment, and the mortality rate is high, or 2-3%(1). The human angiotensin I converting enzyme 2 (ACE2) has been suggested to serve as the receptor for the cell entry of SARS-CoV-2 to cause infection(2). ACE2 is a member of the renin-angiotensin system (RAS), with the function of converting angiotensin II to angiotensin-(1-7) (with 7 amino acids), and converting angiotensin I to angiotensin-(1-9)(3), thereby negatively regulating the effects of angiotensin I converting enzyme (ACE) and the RAS system. In addition to its critical roles in RAS, ACE2 binds the S1 domain of the SARS-CoV Spike (S) protein as the viral receptor, and accounts for the infection of SARS-CoV and syncytia formation(4). The genome sequence of SARS-CoV-2 shows significant similarity (79%) to that of SARS-CoV, while its receptor-binding domain shows even higher similarity to that of SARS-CoV(2), further supporting ACE2 as the receptor of SARS-CoV-2. COVID-19 is a highly infectious respiratory disease with a basic reproduction number R_0 (95% CI) of 3.28 (1.4, 6.49)(5). After binding with ACE2, SARS-CoV-2 priming by the serine protease encoded by the transmembrane serine protease 2 gene (*TMPRSS2*) is also required for the viral entry into host cells(6; 7). In addition, three other genes that may be involved in the SARS-CoV-2 infection were highlighted by a recent study, i.e. the furin, paired basic amino acid cleaving enzyme gene (*FURIN*), encoding the proprotein convertase subtilisin/kexin 3 (PCSK3), the dipeptidyl peptidase 4 gene (*DPP4*), and the basigin (Ok blood group) gene (*BSG*)(8). PCSK3 may activate the SARS-CoV-2 S protein when *TMPRSS2* has low expression(9). *DPP4* and *BSG* have the possibility of serving as the alternative receptor of SARS-CoV-2 invasion(8).

Knowledge about the expression of *ACE2* and *TMPRSS2*, as well as other potential entry genes of SARS-CoV-2 encoded by *FURIN*, *DPP4*, and *BSG*, is extremely important to understand the infection of SARS-CoV-2, and to find ways to prevent the infection. For this purpose, we investigated RNA sequencing transcriptome profiling of samples of airway and oral mucosa, including small airway epithelium, alveolar macrophages, nasal epithelium, and masticatory mucosa. In addition, considering the critical role of mucosal associated invariant T (MAIT) cells in mucosal immune defense against viral infection(10), transcriptome profiling of MAIT was also examined in this study.

2. Materials and Methods

Five datasets of transcriptome profiling by RNA sequencing (RNAseq) were acquired from the NCBI Gene Expression Omnibus (GEO) database (Table 1). We mapped and quantified the trimmed RNA-SEQ reads using HISAT2 (<https://ccb.jhu.edu/software/hisat2/index.shtml>) to hg19 refSeq for each sample at default thresholds. The expression matrix was generated based on Cuffnorm functions in Cufflink package version 2.2.1(11). Library sizes (i.e. sequencing depths) are normalized by the classic-fpkms method. Comparisons of the levels of *ACE*, *ACE2*, and *TMPRSS2*, across different samples were based on the control or pre-exposure samples in each dataset, i.e.

small airway epithelium of 10 healthy never-smokers before smoking E-cigarette; alveolar macrophages of 10 healthy never-smokers before smoking E-cigarette; nasal epithelium of 4 non-smoker females before exposure to third hand smoke; masticatory mucosa of 21 never smokers; and MAIT of 5 healthy bodyweight donors. The relative levels of the target genes were presented as the Fragments Per Kilobase of transcript per Million mapped reads (FPKM). Different types of human tissues are compared in this study. The expression of a single house-keeping genes (HKG) may not be constant in different cell types or different stages of the cell cycle. Instead, we chose 6 most stable HKGs with different essential functions for cell survival, to correct the expression levels of target genes. All values of the target genes were corrected by the average of relative levels of 6 HKGs, i.e. *ACTB*, *GAPDH*, *HMBS*, *HPRT1*, *RPL13A*, and *TBP*.

Table 1 Five datasets of RNAseq profiling analyzed in this study

GEO accession	Sample	Data description	Library Prep	Sequencing	Spots (M)	Bases	Size	GC content	Reference
GSE85121	small airway epithelium(12)	10 healthy never-smokers before and after smoking E-cigarette	TruSeq v2	Illumina HiSeq2500	39.1	9.8Gbp	3.7G	44.27%	hg19
GSE85121	alveolar macrophages(12)	10 healthy never-smokers before and after smoking E-cigarette	TruSeq v2	Illumina HiSeq2500	37.5	9.4Gbp	3.6G	45.62%	hg19
GSE129959	nasal epithelium	4 non-smoker females before and after exposure to thirdhand smoke	Nextera XT	Illumina NextSeq500	51.1	4.2Gbp	1.6G	43.39%	hg19
GSE136262	masticatory mucosa(13)	21 never smokers; 17 current smokers	TruSeq	Illumina HiSeq3000	16.5	840.4M bp	305.7 M	51.43%	hg19
GSE126169	MAIT(10)	5 healthy bodyweight donors; 4 morbidly obese donors	SMAR T-Seq v4	Illumina NextSeq 500	25.2	2.1Gbp	794.4 M	47.29%	hg19

Results and Discussion

Gene expression patterns in five types of normal tissues are shown in Fig.1. As the major counterpart of *ACE2* expression, the expression of *ACE* is also investigated. *ACE* and *ACE2* have medium and comparable levels of expression in both small airway epithelium and masticatory mucosa. These findings suggest that SARS-CoV-2 can infect both small airway epithelium and oral mucosa. The *ACE2* expression with the highest level of *TMPRSS2* expression in small airway epithelium provides explanation for the vulnerability infected individuals have for the characteristic pneumonia of COVID-19. The *ACE2* expression in masticatory mucosa helps explain the high level of infectivity via droplet transmission from SARS-CoV-2 infection residing in the oral mucous membrane. In addition to this study, the analysis on single-cell RNA-seq (scRNA-seq) data of human tissues showed high expression of *ACE2* in pulmonary type II alveolar cells (AT2) and respiratory epithelial cells(14). Liao et al. further demonstrated expression of *ACE2* and its binding with the SARS-CoV-2 S protein in human bronchial epithelial cells(15). In addition, a number of studies demonstrated

the expression of *ACE2* in airway epithelial cells and its correlation with risk factors of severe COVID-19(16; 17).

Interestingly, the expression level of both *ACE2* and *TMPRSS2* in nasal epithelium is much higher than the levels of *ACE* expression, which is consistent with the studies by Sungnak et al(18) and Bunyavanich et al(19). These results provide mechanistic evidence that the SARS-CoV-2 virus resides in both the oral and nasal mucosa of the upper respiratory tract where it is able to bind to the *ACE2* receptor and serving as the principal locus of infections. These results also provide explanation for the high level of viral load in the oral and nasal mucosa and resulting high level of droplets transmission, with the lower airways being responsible for the severe form of pneumonia as well as the aerosol transmission of the virus. With *ACE2* and *TMPRSS2* as the major entry genes, however, other potential entry genes of SARS-CoV-2 encoded by *FURIN*, *DPP4*, and *BSG*, in the respiratory tract should not be neglected for the possibility of serving as the alternative pathway of SARS-CoV-2 infection.




Figure 1. The expression of SARS-CoV-2 receptor genes. a. Established SARS-CoV-2 receptor genes, *ACE*, *ACE2*, and *TMPRSS2* in five different types of samples. b. Possible SARS-CoV-2 alternative receptor genes, *BSG*, *DPP4*, and *FURIN*. Y-axis represents FPKM

values. The boxplot produced by the IBM SPSS Statistics Version 23 shows the mean, the first quartile and the third quartile, and 95% confidence interval (CI).

The expressions of *ACE*, *ACE2*, and *TMPRSS2* are low in MAIT cells. The expression of *ACE* is high in alveolar macrophages, but the expression of *ACE2* can't be detected with a low level of *TMPRSS2* in alveolar macrophages. These patterns of gene expression in the two types of innate immune cells suggest that SARS-CoV-2 has no or little direct impact on these two components of the innate immune system. In addition, these findings suggest that the infection of SARS-CoV-2 can be limited to the respiratory tract, which explains the absent of viremia in many COVID-19 patients(20; 21).

In parallel, we examined whether the exposure factor in each RNAseq dataset affected the expression levels of *ACE*, *ACE2*, and *TMPRSS2*, to assess whether those common factors (i.e. smoking E-cigarette, third hand smoke, smoking, and obesity) are associated with the susceptibility of SARS-CoV-2 infection. FPKM values of *ACE*, *ACE2*, and *TMPRSS2*, within each dataset were compared by paired T test (for GSE85121 and GSE129959) or independent T test (for GSE136262 and GSE126169). However, there was no difference observed ($P>0.05$) between the dataset for *ACE*, *ACE2*, or *TMPRSS2*. Accordingly, while the SARS-CoV-2 virus is highly infectious, our results do not suggest significant change of infectious susceptibility of SARS-CoV-2 infection by these factors. However, because of the modest sample size for each datasets, we must acknowledge that we are underpowered to identify minor effects.

In summary, this study highlights that the nasal mucosa is the most susceptible locus in the respiratory tract for SARS-CoV-2 infection and replication, responsible for the subsequent high level of droplet transmission and should be the focus for protection against SARS-CoV-2 infection, in line with a recently virological analysis(22). Accordingly, local interventions with *ACE2* inhibitor(23) or *TMPRSS2* inhibitor (e.g. camostat mesylate)(7) may represent novel interventions to block SARS-CoV-2 cell entry and treat COVID-19.

Author Contributions: YL, HQ, and JQ, analyzed and interpreted the RNAseq data. YL, HQ, LT, and HH wrote the manuscript. All authors read and approved the final manuscript.

Funding: The study was supported by Institutional Development Funds from the Children's Hospital of Philadelphia to the Center for Applied Genomics, The Children's Hospital of Philadelphia Endowed Chair in Genomic Research to HH.

Conflicts of Interest: The authors declare that they have no competing interests

References

1. Wu Z, McGoogan JM: Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. JAMA 2020;
2. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, Wang W, Song H, Huang B, Zhu N: Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. The Lancet 2020;
3. Ferrario CM, Ahmad S, Nagata S, Simington SW, Varagic J, Kon N, Dell'italia LJ: An evolving story of angiotensin-II-forming pathways in rodents and humans. Clinical science (London, England : 1979) 2014;126:461-469

4. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC: Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* 2003;426:450-454
5. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J: The reproductive number of COVID-19 is higher compared to SARS coronavirus. *Journal of travel medicine* 2020;
6. Lukassen S, Chua RL, Trefzer T, Kahn NC, Schneider MA, Muley T, Winter H, Meister M, Veith C, Boots AW: SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in bronchial transient secretory cells. *The EMBO Journal* 2020;
7. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu N-H, Nitsche A: SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;
8. Latini A, Agolini E, Novelli A, Borgiani P, Giannini R, Gravina P, Smarrazzo A, Dauri M, Andreoni M, Rogliani P, Bernardini S, Helmer-Citterich M, Biancolella M, Novelli G: COVID-19 and Genetic Variants of Protein Involved in the SARS-CoV-2 Entry into the Host Cells. *Genes* 2020;11
9. Millet JK, Whittaker GR: Host cell proteases: Critical determinants of coronavirus tropism and pathogenesis. *Virus research* 2015;202:120-134
10. O'Brien A, Loftus RM, Pisarska MM, Tobin LM, Bergin R, Wood NAW, Foley C, Mat A, Tinley FC, Bannan C, Sommerville G, Veerapen N, Besra GS, Sinclair LV, Moynagh PN, Lynch L, Finlay DK, O'Shea D, Hogan AE: Obesity Reduces mTORC1 Activity in Mucosal-Associated Invariant T Cells, Driving Defective Metabolic and Functional Responses. *Journal of immunology (Baltimore, Md : 1950)* 2019;202:3404-3411
11. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L: Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol* 2010;28:511-515
12. Staudt MR, Salit J, Kaner RJ, Hollmann C, Crystal RG: Altered lung biology of healthy never smokers following acute inhalation of E-cigarettes. *Respir Res* 2018;19:78
13. Richter GM, Kruppa J, Munz M, Wiehe R, Hasler R, Franke A, Martins O, Jockel-Schneider Y, Bruckmann C, Dommisch H, Schaefer AS: A combined epigenome- and transcriptome-wide association study of the oral masticatory mucosa assigns CYP1B1 a central role for epithelial health in smokers. *Clin Epigenetics* 2019;11:105
14. Zou X, Chen K, Zou J, Han P, Hao J, Han Z: Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Frontiers of medicine* 2020:1-8
15. Liao Y, Li X, Mou T, Zhou X, Li D, Wang L, Zhang Y, Dong X, Zheng H, Guo L, Liang Y, Jiang G, Fan S, Xu X, Xie Z, Chen H, Liu L, Li Q: Distinct infection process of SARS-CoV-2 in human bronchial epithelial cells line. *J Med Virol* 2020;
16. Ziegler CG, Allon SJ, Nyquist SK, Mbano IM, Miao VN, Tzouanas CN, Cao Y, Yousif AS, Bals J, Hauser BM: SARS-CoV-2 receptor ACE2 is an interferon-stimulated gene in human airway epithelial cells and is detected in specific cell subsets across tissues. *Cell* 2020;
17. Cai G, Bossé Y, Xiao F, Kheradmand F, Amos CI: Tobacco smoking increases the lung gene expression of ACE2, the receptor of SARS-CoV-2. *American journal of respiratory and critical care medicine* 2020;

18. Sungnak W, Huang N, Bécavin C, Berg M, Queen R, Litvinukova M, Talavera-López C, Maatz H, Reichart D, Sampaziotis F: SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nature medicine* 2020;26:681-687
19. Bunyavanich S, Do A, Vicencio A: Nasal gene expression of angiotensin-converting enzyme 2 in children and adults. *Jama* 2020;
20. Lescure F-X, Bouadma L, Nguyen D, Parisey M, Wicky P-H, Behillil S, Gaymard A, Bouscambert-Duchamp M, Donati F, Le Hingrat Q: Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *The Lancet Infectious Diseases* 2020;
21. Chang L, Yan Y, Wang L: Coronavirus Disease 2019: Coronaviruses and Blood Safety. *Transfusion Medicine Reviews* 2020;
22. Ehmann KZ, Drosten C, Wendtner C, Zange M, Vollmar P, Rosina Ehmann D, Zwirgmaier K, Guggemos M, Seilmaier M, Niemeyer D: Virological assessment of hospitalized cases of coronavirus disease 2019.
23. Huentelman MJ, Zubcevic J, Hernandez Prada JA, Xiao X, Dimitrov DS, Raizada MK, Ostrov DA: Structure-based discovery of a novel angiotensin-converting enzyme 2 inhibitor. *Hypertension* 2004;44:903-906