

Retention in care and viral suppression in differentiated service delivery models for HIV treatment in sub-Saharan Africa: a rapid systematic review

Lawrence Long^{ab††}, Salome Kuchukhidze^{a*}, Sophie Pascoe^b, Brooke E Nichols^{ab}, Matthew P Fox^{ab}, Refiloe Cele^b, Caroline Govathson^b, Amy N Huber^b, David Flynn^c, Sydney Rosen^{ab}

^aDepartment of Global Health, Boston University School of Public Health, Boston, MA, USA

^bHealth Economics and Epidemiology Research Office, Department of Internal Medicine, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

^cAlumni Medical Library, Boston University, Boston, MA, USA

*Contributed equally

†Author for correspondence: Lawrence Long, Boston University School of Public Health, 801 Massachusetts Ave, Boston, MA 02118 USA, lclong@bu.edu

Lawrence Long, lclong@bu.edu

Salome Kuchukhidze, skuchukh@bu.edu

Sophie Pascoe, spascoe@heroza.org

Brooke E Nichols, brooken@bu.edu

Matthew P Fox, mfox@bu.edu

Refiloe Cele, rcele@heroza.org

Caroline Govathson, cgovathson@heroza.org

Amy Huber, ahuber@heroza.org

David Flynn, dflynn@bu.edu

Sydney Rosen, sbrosen@bu.edu

Keywords: Antiretroviral therapy, Differentiated service delivery, Retention, Suppression, Africa, Systematic review

Word count: 3429

Abstract

(350 word limit) 340 words

Introduction:

Differentiated service delivery (DSD) models for antiretroviral treatment (ART) for HIV are being scaled up in the expectation that they will improve the quality and efficiency of treatment delivery and reduce costs while maintaining at least equivalent clinical outcomes. Even this minimum requirement of equivalent clinical outcomes is poorly documented for most models and settings, however. We reviewed the recent literature on DSD models to describe what is known about clinical outcomes.

Methods:

We conducted a rapid systematic review of peer-reviewed publications in PubMed, Embase, and the Web of Science and major international conference abstracts that reported outcomes of DSD models for the provision of ART in sub-Saharan Africa from January 1, 2016 to September 12, 2019. Sources reporting standard clinical HIV treatment metrics, primarily retention in care and viral load suppression, were reviewed and categorized by DSD model and source quality assessed.

Results and discussion:

Twenty-nine papers and abstracts describing 37 DSD models and reporting 52 discrete outcomes met search inclusion criteria. Of the 37 models, 7 (19%) were facility-based individual models, 12 (32%) out-of-facility based individual models, 5 (14%) client-led groups, and 13 (35%) healthcare worker-led groups. Retention was reported for 73% of the models and viral suppression for 57%. Where a comparison with conventional care was provided, retention in most DSD models was within 5% of that for conventional care; where no comparison was provided, retention generally exceeded 80%. For viral suppression, all those with a comparison to conventional care reported a small increase in suppression in the DSD model; reported suppression exceeded 90% in 11/21 models. Analysis was limited by the extensive heterogeneity of study designs, outcomes, models, and populations. Most sources did not provide comparisons with conventional care, and metrics for assessing outcomes varied widely and were in many cases poorly defined.

Conclusion:

Existing evidence on the clinical outcomes of DSD models for HIV treatment in sub-Saharan Africa is limited in both quantity and quality but suggests that retention in care and viral suppression are roughly equivalent to those in conventional models of care.

Prospero number: CRD42019118230

Introduction

Throughout sub-Saharan Africa, most national HIV programs are striving to achieve the 95-95-95 targets for HIV diagnosis, treatment, and viral suppression¹. The rapid expansion of antiretroviral therapy (ART) programs to reach these targets has created shortfalls in health system capacity and quality.² In response, many countries are scaling up alternative service delivery approaches, or differentiated service delivery (DSD) models. DSD models differ from conventional HIV care in the location and frequency of interactions with the healthcare system, cadre of provider involved, and/or types of services provided.³ Grimsrud and colleagues broadly categorize DSD models as individual or group models, with service delivery at a facility or in the community.⁴ DSD models aim to achieve a wide range of potential benefits to both providers and patients. The attractiveness of DSD models is generally considered to be conditional on maintaining at least equivalent clinical outcomes to conventional care; assuming no deterioration in clinical outcomes, DSD models are hoped to generate greater patient satisfaction, lower cost to both providers and patients, and create efficient and convenient service delivery.

Despite the large-scale rollout of DSD models in various formats across multiple countries, there is a dearth of evidence to document the purported benefits of the new models in routine implementation. Even the minimum requirement of equivalent clinical outcomes is poorly documented for most models and settings. The studies and evaluations available are widely inconsistent in their designs, methods, and outcomes, making it difficult to draw an overall picture of the impact of the models. Monitoring and evaluation systems have not kept up with DSD model implementation, and DSD participation is poorly captured in routine records, making it challenging to compare outcomes in DSD models with those in conventional care.⁵ The information available to policy makers, funders, and program implementers is thus incomplete and difficult to interpret.

To help fill this gap and create a baseline to guide future research, we conducted a comprehensive, rapid review of the most recent peer-reviewed reports of the outcomes of DSD model implementation in sub-Saharan Africa. In view of the importance of achieving non-inferior clinical outcomes as a condition for adopting DSD models, we report here the results of our search for retention in care, viral suppression, and related clinical outcomes.

Methods

Following World Health Organization guidance for rapid reviews,⁶ we conducted a rapid systematic review of peer-reviewed publications and conference abstracts that reported outcomes of differentiated service delivery (DSD) models for the provision of antiretroviral treatment (ART) in sub-Saharan Africa since 2016.⁷ The search protocol was previously presented,⁷ and the review was registered on the International Prospective Register of Systematic Reviews (PROSPERO CRD42019118230).

Although the full review included a wide range of outcomes for both providers and patients, the most widely available information pertained to patient-level clinical outcomes, specifically retention in care and viral suppression. In this report we focus on these outcomes only, to allow for a more detailed examination and discussion of consistently-defined indicators. The full report of the review is available online.⁸

Search strategy and study selection

For this review, we adopted and modified the widely-cited frameworks put forward by Grimsrud et al⁴ and Duncombe et al³ and defined as a “differentiated model of service delivery” any approach to providing ART that focused on a specific population, the location of service delivery, the frequency of patient interaction with the healthcare system, or the cadre of health care provider involved. We did not consider a change in services provided, without adjustment of any other characteristics, to constitute a DSD model. Inclusion and exclusion criteria for the review are shown in Supplementary Table 1.

We searched the PubMed, Embase, and Web of Science databases with a search string developed to identify publications which reported on HIV treatment delivery models in sub-Saharan Africa from 1 January 2016 until 12 September 2019. The final search was conducted on 12 September 2019. We supplemented the peer reviewed publications by manually searching peer-reviewed abstracts from major conferences for the same period. Search strings and a full list of conferences included can be found in Supplementary Table 2. Limiting eligible articles and abstracts to those published or presented since 2016 was intended to ensure that results come as close as possible to reflecting the current state of DSD model implementation and to avoid repeating the efforts of previous reviews.^{2,9-12} If a source reported patient follow up data collected both before and after January 1, 2016, we included it only if the majority of follow-up time, as stated in the source or estimated by the authors, occurred after that date.

We excluded sources that reported interventions aimed at improving conventional care that we judged did not in themselves comprise DSD models, such as adherence interventions that strengthened existing counseling or offered incentives for retention within the conventional model of care. We also excluded cross-sectional surveys of patients or providers who were asked to comment on DSD models but did not have personal experience with it. If two source documents described what we determined to be the same cohort of patients enrolled in the same instance of the model, we counted only one model but cited both references for it. If one source document superseded another, e.g. by reporting more complete data or longer-term outcomes, we kept only the more informative source.

All peer reviewed references identified using the respective search strings from PubMed, Embase and Web of Science were imported into an EndNote™ library, where deduplication occurred. An initial, independent, blinded review (reviewers were not aware of each other’s decisions) of the titles and abstracts was conducted by three study team members (SK, RC, CG) using Rayyan QCRI.¹³ A full text review was then conducted for all publications remaining after the initial review by two study team members (SK, CG). Reasons for excluding publications were recorded during the full text review. As a quality check, another author (LL) also checked a sample (10%) of the excluded sources against exclusion criteria. At each stage of the review process any conflicts between reviewers were assessed and resolved through consensus of two authors (LL, SR). The results of the search were documented in accordance with the PRISMA-P reporting checklist (Supplementary Text 1).^{6,14}

Data extraction

The data extraction tool was designed to capture each DSD model separately, regardless of whether the source publication described one or many models. In addition to standard bibliographic descriptors, we collected two types of data: a) a detailed description of the model of service delivery; and b) the outcomes that were reported for the model. We categorized each model according to the taxonomy described by Grimsrud⁴, with four categories: facility based individual models, out of facility based

individual models, healthcare worker led groups, and client led groups. We then used the adapted Duncombe³ schema to describe the model in terms of population, provider, location, frequency, and services provided as well as and its outcomes. Where a comparison was provided with the pre- or non-differentiated standard of care, we also extracted data about these comparison models, henceforth referred to as conventional care.

Outcomes

We report here standard clinical HIV treatment metrics, including retention in care, viral load suppression, adherence, and pharmacy refill rates. We used each source's own definition and timing of these outcomes, accepting that definitions for "retention in care" vary widely, as do thresholds for determining viral suppression. Retention usually referred to the proportion of patients enrolled in a DSD model and retained in the ART program at a specific time point after enrollment in the study. The point at which a patient was considered no longer in care (i.e. not retained) varied by study or country. Where a loss to follow up (LTFU) proportion was reported, we converted it to a retention rate (as 100-LTFU%). Most sources defined viral suppression as <1000 copies/mL. Adherence and prescription refill frequency were uncommon outcomes but are included in this analysis when reported. Other outcomes from the full review, such as costs to providers and patients, can be found in the previously cited report.¹⁵

Analysis

To structure the results, we first divided the models into the four categories mentioned above: facility based individual models (FBIM), out of facility based individual models (OFBIM), client led groups (CLG), and healthcare worker led groups (HCWLG). In publications where more than one model was described, we reported each model separately. We report outcomes as stated in the original publications, adjusted where possible to utilize uniform metrics (for example, by converting a reported percentage of patients lost to follow up to the percentage of patients retained). As explained in the search protocol⁷, we feared that it would be misleading to conduct aggregate analyses due to the heterogeneity of model designs, participating populations, and study settings, even where outcomes themselves were similar. We thus report only the disaggregated results.

We assessed the quality of the cohort studies using the Newcastle-Ottawa scale.^{16,17} The quality rating covered a review of selection, comparability, and outcome domains and generated a score out of 9. There are no standardized quality rating categories, but to simplify interpretation of scores, those studies that scored 7 or above were categorized as high quality, those scoring between 4-6 were of moderate quality, and those scoring below 4 were considered low quality, as done in previous studies.¹⁷ Randomized controlled trials were assessed using the Cochrane Collaboration's tool for assessing risk of bias in cluster randomized controlled trials.¹⁸ We assessed sequence generation, participant recruitment with respect to randomization timing, deviation from intended intervention, completeness of outcome data for each main outcomes, bias in measurement of outcome, bias in selection of the reported result. Risk of bias assessment for the one remaining cross-sectional study was not conducted.¹⁹

Results


Sources identified

The results of the systematic search are shown in Figure 1. A total of 3,498 non-duplicate abstracts of peer reviewed journal articles and 12,822 abstracts from the selected conferences were screened. After

the initial title and abstract review, 16,092 articles and abstracts were excluded, leaving 228 documents for full review. During the full review, an additional 181 were excluded. Reasons for exclusions are reported in Supplementary Table 3. The primary reason (60%) for excluding articles was date: most or all of the underlying data were collected prior to 2016. The main reason for excluding conference abstracts (33%) was insufficient information to adequately describe the model and at least one of the outcomes of interest.

Nine peer reviewed articles and 38 conference abstracts (47 total) were retained in the final data set for the full review. Of these, 29 included one or more clinical outcomes and were included in the analysis reported here. Three quarters of these sources (76%) reported observational cohort studies; most of the rest (21%) were randomized trials. South Africa (27%) and Zambia (22%) jointly accounted for nearly half the sample (Supplementary Table 3).

Figure 1. PRISMA flow chart



Differentiated models included in the review

The 29 sources described outcomes for a total of 37 discrete differentiated service delivery models, excluding conventional care models for comparison. Models are described briefly in Table 1 below and in full in Supplementary Table 4. In the tables, each model is assigned a model identifier (ID), which is

used to reference that model throughout the review. If a source document (article or abstract) reports on more than one DSD model, multiple model IDs will be associated with it in Table 2. Each model identifier contains an acronym for the model category (FBIM, OFBIM, CLG, or HCWLG) followed by a number. For example, client led groups have model IDs CLG1 through CLG5, indicating that there were 5 distinct CLG models identified. In one instance (HCWLG11), the same model is referred to in more than one source document.^{20,21}

Table 1. Models included in the review

ID	Source	Country	Model	Eligibility for model (condition, age*)	Sample size§	Study period
Facility based individual models (FBIM)						
FBIM1 ²²	Kamerhe 2018	Democratic Republic of Congo (DRC)	Fast-track ART refill circuit	Stable†, ≥18	974	10/2016-03/2018
FBIM2 ²³	Havliir 2019¥	Kenya and Uganda	Patient-centered streamlined care	No restriction, all ages	186,354	2013-2017
FBIM3 ²⁴	Mdala 2018	Namibia	Nurse-initiated management of ART	No restriction, all ages	18,367	09/2015-09/2017
FBIM4 ²⁵	Cassidy 2018	South Africa	Quick pick-up model	Stable, not specified	4,753	10/2015-10/2017
FBIM5 ¹⁹	Kaimal 2017	Uganda	Pharmacy refill plus program	Stable, not specified	624	08/2015-09/2016
FBIM6 ²⁶	Nsumba 2019	Uganda	Treatment failure management service	Unstable, not specified	862	03/2017-
FBIM7 ²⁷	Mwila 2018	Zambia	FBO-based community ART delivery	No restriction, all ages	5,014	12/2015-04/2018
Out of facility based individual models (OFBIM)						
OFBIM1 ²⁸	Tchissambou 2018	DRC	Community ART distribution centers	Stable, ≥15	2,027	01/2016-12/2017
OFBIM2 ²²	Kamerhe 2018	DRC	Community-based point of ART distribution	Stable, ≥18	974	10/2016-03/2018
OFBIM3 ²⁹	Mothibi 2018	DRC	Community-based individual drug distribution	Stable, ≥18	1,484	10/2016-09/2017
OFBIM4 ³⁰	Pasipamire 2019¥	Eswatini	Comprehensive outreach	Stable, ≥16	918	02/2015-08/2016
OFBIM5 ³¹	Avong 2018	Nigeria	Community pharmacy ART dispensing	Stable, ≥18	295	02/2016-05/2017
OFBIM6 ³²	Nwabueze 2018	Nigeria	Out-of-facility ARV delivery	Stable, not specified	283	03/2016-02/2017
OFBIM7 ²¹	Fox 2019	South Africa	Decentralized medication delivery	Stable, ≥18	578	03/2016-09/2018
OFBIM8 ³³	Geldsetzer 2018	Tanzania	ARV community delivery	Stable, not specified	2,172	03/2016-10/2017
OFBIM9 ³⁴	Mulenga 2019	Zambia	Community pharmacy dispensing	Stable, not specified	237	08/2018-12/2018
OFBIM10 ³⁵	Mwanda 2018	Zambia	DSD model for prisoners	Stable, not specified	241	10/2016-03/2018
OFBIM11 ³⁶	Chibesakunda 2019	Zambia	Community ART centers	No restriction, all ages	7037	10/2016-01/2019
OFBIM12 ³⁷	Limbada 2019	Zambia	Home-based delivery	Stable, ≥18	2,493	05/2017-12/2017
Client led groups (CLG)						
CLG1 ³⁰	Pasipamire 2019¥	Eswatini	Community ART groups	Stable, ≥16	918	02/2015-08/2016

ID	Source	Country	Model	Eligibility for model (condition, age*)	Sample size§	Study period
CLG2 ³⁸	Van Rompaey 2019	Mozambique	Community adherence groups	Stable, not specified	15,457	08/2016–12/2017
CLG3 ³⁹	Okechukwu 2018	Tanzania	Community ART refill groups	Stable, not specified	13,372	05/2017–11/2017
CLG4 ⁴⁰	Kagimu 2018	Uganda	Community client led ART delivery	Stable, not specified	14	10/2017–09/2018
CLG5 ⁴¹	Mwamba 2018	Zambia	Community adherence groups and viral load clinic	Unstable, ≥14	386	10/2017–11/2018
Healthcare worker led groups (HCWLG)						
HCWLG1 ²²	Kamerhe 2018	DRC	ART support groups	No restriction, all ages	974	10/2016–03/2018
HCWLG2 ³⁰	Pasipamire 2019	Eswatini	Treatment clubs	Stable, ≥16	918	02/2015–08/2016
HCWLG3 ⁴²	Finci 2018	Mozambique	Adherence clubs	Stable, not specified	687	10/2015–03/2018
HCWLG4 ⁴³	Manganye 2018	South Africa	Adherence clubs	Stable, ≥18	141,269	10/2016–06/2018
HCWLG5 ⁴⁴	Meehan 2018	South Africa	Community based adherence clubs	Stable, ≥18	465	08/2017–11/2017
HCWLG6 ⁴⁵	Pahad 2018	South Africa	Youth care clubs	No restriction, 12–24	589	08/2016–12/2017
HCWLG7 ⁴⁶	Lebelo 2019¥	South Africa	6 month-refill adherence clubs	Stable, not specified	2,150	10/2017–02/2019
HCWLG8 ⁴⁶	Lebelo 2019¥	South Africa	2 month-refill adherence clubs	Stable, not specified	2,150	10/2017–02/2019
HCWLG9 ²⁵	Cassidy 2018	South Africa	Community clubs	Stable, not specified	4,753	10/2015–10/2017
HCWLG10 ²⁵	Cassidy 2018	South Africa	Facility clubs	Stable, not specified	4,753	10/2015–10/2017
HCWLG11 ^{20,21}	Fox 2018; Fox 2019	South Africa	Adherence clubs	Stable, ≥18	569; 578	Pre-/–post 01/2015
HCWLG12 ⁴⁷	Roy 2018	Zambia	Urban adherence groups	Not specified, ≥14	1,096; 15 FGD	05/2016–10/2017
HCWLG13 ³⁷	Limbada 2019	Zambia	Adherence clubs	Stable, ≥18	5,252; 869	05/2017–12/2017

*Most models where age was not specified appeared to be limited to adults.

¥The authors used associated documents (e.g. published study protocols, unpublished reports) relevant to these source documents to supplement the DSD model description, if insufficient detail was provided in the publication itself.

§Sample sizes pertain to the entire study population rather than for a specific DSD model. For publications that evaluated different DSD models in each arm, we report the total N for the study cohort rather than the N in each study arm.

‡For most models, stable was defined per national guidelines, though clinicians used clinical criteria to define stability when necessary laboratory tests were not available.

FBO, faith-based organization; FGD, focus group discussion

In addition to the models listed in Table 1, 11 source documents reported comparative results for a conventional care model, creating a total of 48 model-instances with clinical outcomes included in this review (37 DSD + 11 conventional models). Out of facility based individual models (32%) and healthcare worker led group models (35%) were the most commonly reported categories (Supplementary Table 5).

Three quarters (76%) of the models were limited to clinically stable patients, and most (59%) were for adults (Table 1). Definitions of stability varied. Some models required prior evidence of viral suppression while others relied on clinical condition, for example, and minimum duration on ART prior to model entry also differed. Details of how a stable patient is defined are presented elsewhere.⁴⁸

Additional model characteristics are described in Supplementary Table 6. Most models provided basic clinical care, antiretroviral medications (ARVs), and laboratory monitoring only (78%). Almost half (46%) included services delivered both in the clinic and in the community, rather than solely one or the other. For those that identified clinical care and pharmacy refill providers, nearly all clinical care (96%) was provided by trained clinicians, though few sources specified the clinical cadre involved; more than two thirds of medication refills (70%) were provided by non-clinician staff (community health workers, designated patients, or lay counselors). More than half the models (57%) required patients to have a total of 4-8 clinic visits or DSD model interactions per year; most of the rest required more than 8 visits or interactions per year, though a few (18%) were structured for 3 or fewer per year (Supplementary Table 6).

Outcomes

A total of 52 outcomes were reported for the 37 models included in the review (Table 2). Retention in care was the most common, reported for 73% of the models. Just over half the models (57%) reported viral suppression.

Table 2. Frequency of DSD model outcomes reported, by model category*

Outcome reported (n, %)	Facility based individual (FBIM) (n=7)	Out of facility individual (OFBIM) (n=12)	Client led group (CLG) (n=5)	Healthcare worker led group (HCWLG) (n=13)	Total (n=37)
Retention	4 (57%)	10 (83%)	1 (20%)	12 (92%)	27 (73%)
Viral load suppression	6 (86%)	6 (50%)	3 (60%)	6 (46%)	21 (57%)
Adherence	0 (0%)	0 (0%)	1 (20%)	0 (0%)	1 (3%)
Prescription refill	0 (0%)	1 (8%)	1 (20%)	1 (8%)	3 (8%)

*Most models reported more than one outcome, resulting in column totals that are greater than the number of models in each category.

Quantitative results for each study are shown in Table 3. Some studies included effect sizes in comparison with conventional care, while others did not provide comparison values at all, but simply reported the outcomes of the DSD models. Table 4 provides additional information, including effect sizes, for studies that did report these measures. More detailed versions of both tables, including any estimates or calculations by the authors, can be found in Supplementary Table 7.

Retention in care

Although retention in care was the most commonly reported outcome, only a few sources provided a comparison to conventional care. For those that did, retention in the DSD model was generally within 5% of that in conventional care, with the exception of a healthcare worker led group model in the Democratic Republic of Congo, which greatly improved retention.²² Among those not providing a comparison, retention generally exceeded 80%. For the few sources (n=4) which reported retention outcomes with an effect size, effects varied widely, from much better than conventional care to somewhat worse.

Viral load suppression

Among the 21 models that reported viral load suppression, eleven included a comparison with conventional care (including one that reported only an effect estimate and not actual values). All those with a comparison reported a small increase in suppression in the DSD model. Reported suppression exceeded 90% in 11/21 models. Five models reported viral suppression with an effect size estimate. Three of these found no difference in suppression when adjusting for baseline differences. Streamlined care in Uganda and Kenya²³ and CAGs in Mozambique³⁸ both reported approximately 15% (prevalence ratio=1.15 and unadjusted odds ratio=1.16, respectively) improvements in suppression.

Adherence and prescription refill rates

Few sources (n=4) used adherence to ARVs or prescription refill rates as outcomes; results are shown in Table 3. Rates of adherence (n=1) and prescription refill (n=3) were >90% across the models. Only two reported a comparison with conventional care and the DSD model outperformed conventional care in both instances. No effect sizes were reported for adherence or prescription refill measures.

Quality of evidence

Among the three quarters of the sources included that were cohort studies and thus evaluated on the Newcastle-Ottawa scale, the quality of the evidence was generally low to moderate (Supplementary Table 8). Only two of the 22 cohort studies received a score of 7 points (high quality) on the 9-point scale. The relatively low quality of evidence among cohort studies was due mainly to the absence of comparators in many of the studies and the scarcity of detail found in conference abstracts. Most of the remaining studies (n=6) were randomized controlled trials, for which we assessed quality using the Cochrane Collaboration's tool for assessing risk of bias cluster randomized trials (Supplementary Table 9).¹⁸ All three full-length articles (four models) were at low risk for bias^{21,23,33} but a concern about bias applied to the two abstracts, driven mainly by the fact that the conference abstracts did not contain full information on study methodology.^{37,46}

Table 3. Clinical outcomes as reported for DSD and comparison models

Source	Model ID	Country	Model name	Timing of outcome	Differentiated model outcome (%)	Conventional care model outcome (%)	Difference (=DSD minus conventional)
Retention in care							
<i>Facility based individual model</i>							
Kamerhe 2018 ²²	FBIM1	DRC	Fast-track ART refill	≤12 months	96.0%	60.0%	+33.7%
Cassidy 2018 ²⁵	FBIM4	South Africa	Quick pick-up model	≤12 months	91.0%	86.0%	+5%
Kaimal 2018 ¹⁹	FBIM5	Uganda	Pharmacy refill plus program	≤12 months	99.3%		
Nsumba 2019 ²⁶	FBIM6	Uganda	Treatment failure management service	Unknown	47.3%		
<i>Out of facility based individual model</i>							
Kamerhe 2018 ²²	OFBIM2	DRC	Community-based point of ART distribution	≤12 months	96.0%	60.0%	+33.7%
Mothibi 2018 ²⁹	OFBIM3	DRC	Community-based individual drug distribution	≤12 months	96.5%		
Tchissambou 2018 ²⁸	OFBIM1	DRC	Community ART distribution center	≤12 months	86.1%		
Pasipamire 2019 ³⁰	OFBIM4	Eswatini	Comprehensive outreach	≤12 months	90.8%		
Avong 2018 ³¹	OFBIM5	Nigeria	Community pharmacy ART dispensing	≤12 months	97.2%**		
Nwabueze 2018 ³²	OFBIM6	Nigeria	Out-of-facility ART delivery	≤12 months	92.5%		
Fox 2019* ²¹	OFBIM7	South Africa	Decentralized medication delivery	≤12 months	81.5%	87.2%	-5.7%
Chibesakunda 2019 ³⁶	OFBIM11	Zambia	Community ART centers	≤12 months	83%		
Limbada 2019 ³⁷	OFBIM12	Zambia	Home-based delivery	12-24 months	84.9%		
Mulenga 2019 ³⁴	OFBIM9	Zambia	Community pharmacy dispensation	≤12 months	100%		
<i>Healthcare worker led group</i>							
Kamerhe 2018 ²²	HCWLG1	DRC	ART support groups	≤12 months	93.7%	60.0%	+33.7%
Pasipamire 2019 ³⁰	HCWLG2	Eswatini	Comprehensive outreach	≤12 months	90.8%		
Finci 2018 ⁴²	HCWLG3	Mozambique	Adherence clubs	12-24 months	84.4%		
Cassidy 2018 ⁴²⁵	HCWLG9	South Africa	Community clubs	≤12 months	89.9%	86.0%	+3.9%
Cassidy 2018 ⁴²⁵	HCWLG10	South Africa	Facility clubs	≤12 months	85.1%	86.0%	+0.9%
Fox 2019* ²¹	HCWLG11	South Africa	Adherence clubs	≤12 months	89.5%	81.6%	+7.9%
Lebelo 2019* ⁵⁴⁶	HCWLG7 and HCWLG8	South Africa	6 month-refill adherence clubs and 2 month-refill adherence clubs (comparison)	≤12 months	97%	98%	-1%
Manganye 2018 ⁴³	HCWLG4	South Africa	Adherence clubs	12-24 months	94.9%		

Source	Model ID	Country	Model name	Timing of outcome	Differentiated model outcome (%)	Conventional care model outcome (%)	Difference (=DSD minus conventional)
Meehan 2018 ⁴⁴	HCWLG5	South Africa	Community based adherence clubs	12-24 months	82.5%		
Pahad 2018 ⁴⁵	HCWLG6	South Africa	Youth care clubs	≤12 months	80.9%	84.0%	-3.1%
Limbada 2019 ³⁷	HCWLG13	Zambia	Adherence clubs	12-24 months	92.6%		
Roy 2018 ⁴⁷	HCWLG12	Zambia	Urban adherence groups	≤12 months	71.0%	42.0%	29.0%
Client led group							
Pasipamire 2019 ³⁰	CLG1	Eswatini	Community ART groups	≤12 months	94.4%		
Kagimu 2018 ⁴⁰	CLG4	Uganda	Community client led ART delivery	≤12 months	100.0%		
Viral load<1000 copies/mm³							
Facility based individual model							
Havilir 2019 ²³	FBIM2	Kenya and Uganda	Patient-centered streamlined care	Unknown	79%	68%	+9%
Mdala 2018 ²⁴	FBIM3	Namibia	Nurse-initiated management of ART	Unknown	86.0%		
Cassidy 2018 ²⁵	FBIM4	South Africa	Quick pick-up model	12-24 months	96.0%	91.0%	+5%
Kaimal 2018 ¹⁹	FBIM5	Uganda	Pharmacy refill plus program	≤12 months	98.8%		
Nsumba 2019 ²⁶	FBIM6	Uganda	Treatment failure management service	Unknown	79.3%		
Mwila 2018 ²⁷	FBIM7	Zambia	FBO based community ART delivery	12-24 months	89.1%	83.8%***	+5.3%
Out of facility based individual model							
Mothibi 2018 ²⁹	OFBIM3	DRC	Community-based individual drug distribution	≤12 months	98.5%		
Nwabueze 2018 ³²	OFBIM6	Nigeria	Out-of-facility ART delivery	≤12 months	100.0%		
Fox 2019* ²¹	OFBIM7	South Africa	Decentralized medication delivery	≤12 months	77.2%	74.3%	+2.9%
Geldsetzer 2018* ³³	OFBIM8	Tanzania	ARV community delivery	≤12 months	90.3%	89.1%	+1.2%
Chibesakunda 2019 ³⁶	OFBIM11	Zambia	Community ART centres	≤12 months	90.4%	84.8%	+5.6%
Mwanda 2018 ³⁵	OFBIM10	Zambia	DSD model for prisoners	Unknown	91.7%		
Health care worker led group							
Finci 2018 ⁴²	HCWLG3	Mozambique	Adherence club	12-24 months	81.0%		
Cassidy 2018 ²⁵	HCWLG9	South Africa	Community clubs	12-24 months	98.0%	91.0%	+7.0%
Cassidy 2018 ²⁵	HCWLG10	South Africa	Facility clubs	12-24 months	94.9%	91.0%	+3.9%
Fox 2019* ²¹		South Africa	Adherence clubs	≤12 months	80.0%	79.6%	+0.4%

Source	Model ID	Country	Model name	Timing of outcome	Differentiated model outcome (%)	Conventional care model outcome (%)	Difference (=DSD minus conventional)
Lebelo 2019* ^{§46}	HCWLG7 and HCWLG8	South Africa	6 month-refill adherence clubs and 2 month-refill adherence clubs (comparison)	≤12 months	97.8%	96.5%	+1.3%
Pahad 2018 ⁴⁵	HCWLG6	South Africa	Youth care clubs	≤12 months	75.0%		
Client led group							
Kagimu 2018 ⁴⁰	CLG4	Uganda	Community client led ART delivery	≤12 months	100.0%		
Mwamba 2018 ⁴¹	CLG5	Zambia	Community adherence groups and dedicated VL clinic for unstable patients	Unknown	27.8%		
Van Rompaey ^{¥38}	CLG2	Mozambique	Community adherence groups	Unknown	uOR=1.16*		
Adherence							
Client led group							
Kagimu 2018 ⁴⁰	CLG4	Uganda	Community client led ART delivery	≤12 months	95.0%		
Prescription refill							
Avong 2018 ³¹	OFBIM5	Nigeria	Community pharmacy ART dispensing	≤12 months	100.0%		
Fox 2018 ²⁰	HCWLG11	South Africa	Adherence clubs	≤12 months	92.0%	88.0%	+4%
Okechukwu 2018 ³⁹	CLG3	Tanzania	Community ART refill groups	≤12 months	97.9%	87.0%	+10.9%

*Cluster randomized trial

**Rate calculated by authors

***National average

§ The comparator is a 2-month pick-up model in adherence clubs which is counted as a DSD model, rather than conventional care. This source document was not reported as including the standard of care in Supplementary table 5.

¥ Only the effect estimates were reported by the source.

Table 4. Clinical outcomes of DSD models with effect size estimates compared to conventional care

Source	Model ID	Country	Model Name	N	Outcome		Effect size		Outcome definition
					DSD model	Conventional model	Crude	Adjusted	
Retention									
Pasipamire 2019 ³⁰	OFBIM4	Eswatini	Comprehensive outreach	918	90.8	-	HR=0.68** (0.30 to 1.51)	HR=0.58** (0.25 to 1.35)	Patient loss to follow up (LTFU) was defined as not having a recorded visit for 120 days or more before database closure. LTFU from care was time from enrolment to the composite endpoint of LTFU and death, regardless whether the outcome occurred while enrolled in the care model or in routine facility-based ART care. Retention was measured at 12 months
	CLG1		Community ART groups	918	94.4	-	HR=1.0 (reference)	HR=1.0 (reference)	
	HCWLG2		Treatment clubs	918	94.4	-	HR=1.07** (0.54 to 2.08)	HR=1.09** (0.54 to 2.22)	
Fox 2019* ²¹	OFBIM7	South Africa	Decentralized medication delivery	1,147	81.5	87.2	DID= -6.0% (-10.6% to -1.5%)	DID=-5.9% (-12.5% to 0.8%) ^a	Retention in care at 12 months after eligibility defined as 100% - % attrition, with attrition as the sum of reported deaths, loss to follow-up, and transfers. Loss to follow-up was defined as failure to attend the clinic within 90 days of a scheduled appointment.
	HCWLG11		Adherence clubs	1,147	89.5	81.6	DID=7.4% (2.9% to 11.9%)	DID=8.3% (1.1% to 15.6%) ^a	
Lebelo 2019* ⁴⁶	HCWLG7 HCWLG8	South Africa	6 month-refill adherence clubs and 2 month-refill adherence clubs (comparison)	2,150	97	98	-	HR=1.47 (0.79 to 2.7) [§]	Retention was defined as any visit within 3-months of the 12-month scheduled appointment
Roy 2018 ⁴⁷	HCWLG12	Zambia	Urban adherence groups	1,096	71.0	42.0	RR=1.7 (1.32 to 2.22)**	-	Cumulative incidence of the first late drug pick-up at 12 months used as a proxy for retention**
Viral load<1000 copies/mm3									
Havlir 2019* ²³	FBIM2	Kenya and Uganda	Patient-centered streamlined care	186,354	79	68	-	PR (prevalence ratio) =1.15 (1.11 to 1.20) ^c	Viral suppression defined as HIV RNA <500 copies per milliliter at 3 years (cross-sectional measurement)

Source	Model ID	Country	Model Name	N	Outcome		Effect size		Outcome definition
					DSD model	Conventional model	Crude	Adjusted	
Van Rompaey ³⁸	CLG2	Mozambique	Community adherence groups	15,457	-	-	uOR=1.16 [‡]		Viral suppression defined as HIV RNA <75 copies per milliliter. Time of follow-up is unclear.
Fox 2019 ^{*21}	HCWLG11	South Africa	Adherence clubs	1,147	80	79.6	DID= 3.1% (-3.8% to 10.0%)	DID= 3.8% (-6.9% to 14.4%) ^a	Viral suppression (<400 copies/mL) at 12 months after eligibility
	OFBIM7		Decentralized medication delivery	1,266	77.2	74.3	DID= -0.5% (-7.5% to 6.6%)	DID= -1.0% (-12.2% to 10.1%) ^a	
Geldsetzer 2018 ^{*33}	OFBIM8	Tanzania	ARV community delivery	2,172	90.3	89.1	RR=1.12 (0.80 to 1.59)**	RR=1.00 (0.74 to 1.35)** ^b	Viral load ≥1000 copies/mL** at 6 months
Prescription refill									
Fox 2018 ²⁰	HCWLG11	South Africa	Adherence clubs	579	92.0	88.0	DiD=6.7% (3.4% to 10.4%)	DiD=7.5% (-1.3% to 16.2%) ^a	Proportion of patients who picked up medication within four months of enrolling in the intervention.

*Cluster randomized trial

**Author calculated

§ The model compares 6-month dispensing adherence clubs to 2-month dispensing adherence clubs, not the SOC

‡ Only the effect estimates were reported

^a Cluster and covariate adjusted

^b Adjusted for baseline viral load, CD4 count, age and sex

^c Adjusted for incomplete measures of HIV serostatus and HIV RNA with the use of individual-level targeted maximum likelihood estimation (adjustment variables included sex, age group, marital status, educational level, occupation, alcohol use, household wealth, mobility, previous HIV testing, and care status).

Discussion

We systematically reviewed and synthesized the current evidence related to clinical outcomes of differentiated service delivery models for HIV treatment in sub-Saharan Africa between 2016 and 2019. While we identified 29 sources that described one or more clinical outcomes of 37 DSD models in 11 countries, only a minority (28%) compared the alternative models to conventional care or to one another, making it difficult to draw strong conclusions about the overall impact of DSD models on clinical outcomes. Because of the heterogeneity of outcome definitions and timing and the highly variable quality, size, and scope of the studies included, we opted to present outcomes individually for each model, stratified by model category and outcome, rather than to estimate aggregate statistics.

For those models that did provide a comparison with conventional care, retention in care in DSD models was generally within 5% of that in conventional care, with a few exceptions that reported much better retention. Similarly, viral suppression was generally equivalent or slightly higher in the DSD models. We did not expect to see a marked improvement in clinical impact (retention or viral suppression) because most DSD models are limited to already-stable patients, for whom outcomes can be sustained but cannot improve. Where comparisons with conventional care were provided and effect sizes reported, effects on retention and suppression varied widely, from slightly worse than conventional care to moderately better. In general, DSD models were not associated with a meaningful deterioration in patient outcomes, despite in many cases having fewer interactions with patients or relying on lower cadres of clinicians than did conventional care.

As is evident from the discussion above, this review had many limitations. While we believe that our search of the peer-reviewed, published literature and abstracts was thorough, the lack of standard terminology for describing DSD models hampered the creation of precise search strings, and it is possible that some sources were missed. Most sources did not describe procedures for recruiting patients into DSD models, but it is possible that self- and provider-selection biased participation toward the most motivated and empowered patients, among all those who met formal eligibility criteria. More important, the extreme heterogeneity of the sources that did meet inclusion criteria rendered any attempt to aggregate results or produce summary statistics misleading. This heterogeneity manifested itself in multiple ways. The topic of DSD models is highly diverse in itself. Evaluation methods ranged from single-site, single-arm observational cohorts to large randomized trials. The majority of sources did not provide comparisons with conventional care, and metrics for assessing outcomes varied widely and were in many cases poorly defined. The underlying patient populations were often poorly described or differed by design between models even within countries.

Stemming from these limitations, the search reported here identifies gaps in the evidence base and research priorities for DSD model implementation in the coming years. In particular, rigorous evaluation of clinical outcomes, with relevant comparisons, is needed if we are to fully understand the implications of DSD models for HIV control. Longer-term follow-up under routine care settings, beyond the first 12 or 24 months, should be undertaken, as it is critical to know what happens to retention and viral suppression three, five, or ten years after entry into a DSD model. This is especially important when DSD models are focused on stable patients and large changes in treatment outcomes are unlikely in the short term. Evaluation reports on the outcomes of DSD models should consistently include a description of the population served, as models limited to already-stable patients are likely to have different outcomes from those that enroll a cross-section of the ART patient population. There is also a need for electronic medical record systems to evolve to capture data on DSD model participation, as this is an essential step towards understanding the true clinical and other impacts of DSD models.

Conclusions

We note that there is a difference between the clinical outcomes of the patient enrolled in DSD models and the “impact” of implementing DSD models as part of national HIV programs. In many of the studies included in this review, only a small proportion of eligible patients were enrolled in a DSD model, and only those patients’ outcomes reported. The effect of those patients’ outcomes on the overall, aggregate outcomes of the healthcare facilities at which the DSD models were implemented may have been modest, or even trivial, if large numbers of other patients remained in conventional care. Future evaluations of the outcomes of DSD models would be of greater value if they considered the entire, relevant patient population—for example, all the ART patients served by a facility, or all the ART patients in a catchment area—as the denominator for assessing success.

Conflicts of interest

The authors declare that they have no conflicting interests.

Authors’ contributions

LL, SK, and SR conceived of and designed the study. SK, SP, BEN, RC, CG and AH identified and reviewed sources and extracted data. LL, SK, and SR analyzed the data and drafted the manuscript. DF contributed to design and data collection. MF contributed to design and data analysis. All authors reviewed and edited the manuscript.

Acknowledgements

Funding for the study was provided the Bill & Melinda Gates Foundation through OPP1192640 to Boston University. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Additional files

Supplementary text 1. PRISMA checklist

Supplementary table 1. Inclusion/exclusion criteria for publications and abstracts

Supplementary table 2. Search strategy

Supplementary table 3. Reasons for exclusion after full-text review

Supplementary table 4. Original source documents and models

Supplementary table 5. Characteristics of source documents

Supplementary table 6. Characteristics of service delivery models

Supplementary table 7. Details of treatment outcomes reported in the source documents

Supplementary table 8. Risk of bias assessment for cohort studies

Supplementary table 9. Risk of bias assessment for randomised trials

References

1. UNAIDS. Fast-track: ending the AIDS epidemic by 2030. 2014. Available from: https://www.unaids.org/sites/default/files/media_asset/JC2686_WAD2014report_en.pdf
2. Decroo T, Rasschaert F, Telfer B, Remartinez D, Laga M, Ford N. Community-based antiretroviral therapy programs can overcome barriers to retention of patients and decongest health services in sub-saharan africa: A systematic review. *Int Health*. 2013;5(3):169–79.
3. Duncombe C, Rosenblum S, Hellmann N, Holmes C, Wilkinson L, Biot M, et al. Reframing HIV care: Putting people at the centre of antiretroviral delivery. *Trop Med Int Heal*. 2015 Apr 1;20(4):430–47.
4. Grimsrud A, Barnabas R V., Ehrenkranz P, Ford N. Evidence for scale up: The differentiated care research agenda. *J Int AIDS Soc*. 2017;20(00):1–6.
5. Reidy WJ, Rabkin M, Syowai M, Schaaf A, El-Sadr WM. Patient-level and program-level monitoring and evaluation of differentiated service delivery for HIV: A pragmatic and parsimonious approach is needed. *AIDS*. 2018;32(3):399–401.
6. Stevens A, Garritty C, Hersi M, Moher D. Developing PRISMA-RR, a reporting guideline for rapid reviews of primary studies (Protocol). 2018. Available from: <https://www.equator-network.org/wp-content/uploads/2018/02/PRISMA-RR-protocol.pdf>
7. Long L, Kuchukhidze S, Pascoe S, Nichols B, Cele R, Govathson C, et al. Differentiated models of service delivery for antiretroviral treatment of HIV in sub-Saharan Africa: a rapid review protocol. *Syst Rev*. 2019;8(1):314.
8. Long L, Kuchukhidze S, Pascoe S, Nichols B, Cele R, Govathson C, et al. Differentiated Service Delivery Models for Anitretroviral Treatment of HIV in Sub-Saharan Africa: A Rapid Systematic Review. 2020. Available from: <http://www.heroza.org/wp-content/uploads/2020/02/AMBIT-report-04-systematic-review-v-1.0-2020-02-06.pdf>
9. Hagey A, Li JM, Barr-Walker X. Differentiated HIV care in sub-Saharan Africa: a scoping review to inform antiretroviral therapy provision for stable HIV-infected individuals in Kenya. *AIDS Care*. 2018;30(12):1477–87.
10. Murray KR, Dulli LS, Ridgeway K, Dal Santo L, De Mora DD, Olsen P, et al. Improving retention in HIV care among adolescents and adults in low- and middle-income countries: A systematic review of the literature. *PLoS One*. 2017;12(9).
11. Mukumbang FC, Van Belle S, Marchal B, van Wyk B. An exploration of group-based HIV/AIDS treatment and care models in Sub-Saharan Africa using a realist evaluation (Intervention-Context-Actor-Mechanism-Outcome) heuristic tool: A systematic review. *Implement Sci*. 2017;12(1).
12. Wouters E, Van Damme W, Van Rensburg D, Masquillier C, Meulemans H. Impact of community-based support services on antiretroviral treatment programme delivery and outcomes in resource-limited countries: A synthetic review. *BMC Health Serv Res*. 2012;12(194).
13. Ouzzani M, Hammady H, Elmagarmid A, Fedorowicz Z, Ryman QCRI. *Syst Rev*. 2016;5(210).
14. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Rev Esp Nutr Humana y Diet*. 2016;20(2):148–60.
15. Long L, Kuchukhidze S, Pascoe SJ, Nichols B, Cele R, Govathson C, et al. Differentiated service delivery models for antiretroviral treatment of HIV in sub-Saharan Africa: a rapid systematic review. Boston; 2020. (AMBIT Project Report). Report No.: 04. Available from: <http://www.heroza.org/wp-content/uploads/2020/02/AMBIT-report-04-systematic-review-v-1.0-2020-02-06.pdf>
16. Wells G, Shea B, O'Connell D. Newcastle-Ottawa Quality Assessment Scale for Cohort Studies. 2019. Available from: http://www.ohri.ca/programs/clinical_epidemiology/nosgen.pdf

17. Brennan AT, Bonawitz R, Gill CJ, Thea DM, Kleinman M, Long L, et al. A Meta-analysis Assessing Diarrhea and Pneumonia in HIV-Exposed Uninfected Compared With HIV-Unexposed Uninfected Infants and Children. *J Acquir Immune Defic Syndr*. 2019;82(1).
18. Eldridge S, Campbell M, Campbell M, Dahota A, Giraudeau B, Higgins J, et al. Revised Cochrane risk of bias tool for randomized trials (RoB 2.0) Additional considerations for cluster-randomized trials. 2016. Available from: <https://sites.google.com/site/riskofbiastool/welcome/rob-2-0-tool/archive-rob-2-0-cluster-randomized-trials-2016?authuser=0>
19. Kaimal A, Castelnuovo B, Atwiine M, Musomba R, Nabaggala MS, Ratanshi RP, et al. Experiences with retention in care and viral suppression in a pharmacy refill program. Abstract 1001. In Seattle, USA: Conference on Retroviruses and Opportunistic Infections; 2017.
20. Fox MP, Pascoe S, Huber A, Murphy J, Phokojoe M, Gorgens M, et al. Short-term Effects of Alternative Medication Refill Strategies in South Africa. Abstract 1102. In Boston, MA: Conference on Retroviruses and Opportunistic Infections; 2018.
21. Fox MP, Pascoe S, Huber AN, Murphy J, Phokojoe M, Gorgens M, et al. Adherence clubs and decentralized medication delivery to support patient retention and sustained viral suppression in care: Results from a cluster-randomized evaluation of differentiated ART delivery models in South Africa. *PLoS Med*. 2019;16(7).
22. Kamerhe D, Tendo C, Mwamba R, Thior I, Canagasabay D, Kiluba J, et al. Impact of offering differentiated HIV care on treatment retention and health facility workload: Results from Kenya health zone in the Democratic Republic of the Congo. Abstract WEPEE749. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
23. Havlir D V., Balzer LB, Charlebois ED, Clark TD, Kwarisiima D, Ayieko J, et al. HIV testing and treatment with the use of a community health approach in rural Africa. *N Engl J Med*. 2019;381(3):219–29.
24. Mdala J., Wabmba S., Kalimugogo P, McQuide P, Indongo R. National Policy on Nurse-Initiated Management of ART Supports Decentralization of HIV Service in Namibia. Abstract THPED556. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
25. Cassidy T, O’Connell R, Euvrard J, Ndlovu S, Duran LT, Steele SJ, et al. A “Quick Pick-Up” Differentiated Model of ART Delivery Shows Good Retention in Care. Abstract WEPEE751. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
26. Nsumba S, Musomba R, Lumu I, Namitala F, Nakiyingi W, Tibakabikoba H, et al. Implementation and Outcomes of a Dedicated Treatment Failure Management Service in an Urban HIV Treatment Clinic in Uganda. Abstract 286. In Accra, Ghana: The 13th INTEREST Conference; 2019.
27. Mwila E, Mwanda K, Chikuba-Mcleod M, Chungulo P, O’bra H, Waddle I, et al. Differentiated service delivery: Harnessing the reach of faith-based organizations to deliver quality HIV services in Zambia. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
28. Tchissambou T, Jha A, Dane S, Dibulundu D, Matsumaini S, Madevu-Matson C, et al. Community ART distribution center (PODI) as a differentiated service delivery measure in The Democratic Republic of the Congo. Abstract WEPEE735. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
29. Mothibi E, Mangoma-Chaurura J, Kabamba D, Kabongo B, Kanyinda M, Makumbe B, et al. Postes de Distribution Communautaire (PODIs) in the Democratic Republic of Congo (DRC). Abstract WEPEE754. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
30. Pasipamire L, Nesbitt RC, Ndlovu S, Sibanda G, Mamba S, Lukhele N, et al. Retention on ART and predictors of disengagement from care in several alternative community-centred ART refill models in rural Swaziland. *J Int AIDS Soc*. 2018;21(9).
31. Avong YK, Aliyu GG, Jatau B, Gurumnaan R, Danat N, Kayode GA, et al. Integrating community pharmacy into community based anti-retroviral therapy program: A pilot implementation in

- Abuja, Nigeria. PLoS One. 2018;13(1).
32. Nwabueze E, Bukar A, Ezekiel J, Makumbi M, Ndulue N, Waibale P. Optimizing clients' retention on antiretroviral treatment using an "Out-of-facility health care worker group differentiated model" in Nigeria: Experiences and challenges. Abstract TUPEE603. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
 33. Geldsetzer P, Francis JM, Sando D, Asmus G, Lema IA, Mboggo E, et al. Community delivery of antiretroviral drugs: A non-inferiority cluster-randomized pragmatic trial in Dar es Salaam, Tanzania. PLoS Med. 2018 Sep;15(9).
 34. Mulenga, Helen B;Mubiana,Muhau;Kateule K. Differentiated service delivery using a "Community pharmacy ART drug pickup" model: an evaluation from Lusaka, Zambia. Abstract MOPED623. In Mexico City, Mexico: 10th IAS Conference on HIV Science; 2019.
 35. Mwanda K, Chikuba-McLeod M, Mwila E, Lungu Y, Chungulo P, Phiri D, et al. ART behind bars: Differentiated service delivery results in high viral load suppression among prisoners in Zambia. Abstract WEPED530. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
 36. Chibesakunda K, Chikuba M, Mwila E, Chungulo P, Kawanga L. Community ART centres – are they filling the gap and meeting the grade? A comparison of quality and numbers between community based ART centres and conventional hospitals in Zambia. Abstract WEPED853. In Mexico City, Mexico: 10th IAS Conference on HIV Science; 2019.
 37. Limbada M, Macleod D, Fidler S, Schaap A, Shibwela O, Chiti B, et al. Retention in non-facility-based ART delivery models among stable ART patients in Lusaka, Zambia – Findings from the HPTN 071 (PopART) trial. Abstract MOPED530. In Mexico City, Mexico: 10th IAS Conference on HIV Science; 2019.
 38. Van Rompaey S, Melo M, Bilhete F, Tancredo I, Wester C, De Schacht C. Improved viral suppression rates among HIV-positive adults receiving antiretroviral therapy (ART) via community adherence group (CAG) support in Zambézia province, Mozambique. Abstract 305. In Accra, Ghana: The 13th INTEREST Conference; 2019.
 39. Okechukwu E. Decentralization of ARV refill in Njombe region, Tanzania; USAID Boresha Afya Southern Zone strategy towards reaching the 'hard-to-reach' patients. Abstract THPEE730. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
 40. Kagimu D, Egessa J, Ocul L SE. Overcoming barriers to access of HIV/AIDS services among female sex workers through differentiated service delivery models, TASO Entebbe experience. Abstract WEPEE759. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
 41. Mwamba D, Thulani R, Herce M. Community adherence group (CAG) for HIV viremic patients: early lessons learnt from Lusaka, Zambia. Abstract 184. In Kigali, Rwanda: 12th INTEREST Conference; 2018.
 42. Finci I, Flores A, Gutierrez Zamudio AG, De Abreu E, Issufo S, Gaspar I, et al. The first experience with ART adherence clubs in Maputo, Mozambique: An analysis of a second-line ART subcohort. Abstract WEPEE757. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
 43. Manganye M, Berrada S, Wagner A, Gombarume S, Make A, Wylie C. Increasing retention in care through community systems strengthening: lessons learned from 15 districts in South Africa. Abstract WEA502. In 22nd International AIDS Conference; 2018.
 44. Meehan A, Gunst C, Wademan D, Hodinott G, Holtman R, Grobbelaar N, et al. Evaluating the impact of community-based adherence clubs on the clinical outcomes amongst ART patients in the Cape Winelands district of Western Cape Province of South Africa. Abstract THPEE770. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
 45. Pahad S, Beery M, Martin CE, Henwood R, Dondolo E, Ngobeni M, et al. Youth care clubs: Optimising clinic time, fostering peer support, improving adherence. Abstract 89. In Kigali, Rwanda: 12th INTEREST Conference; 2018.

46. Lebelo K, Cassidy T, Grimsrud A, Keene C, Ndlovu S, Hayes H, et al. Twelve-month retention and viral load outcomes from a non-inferiority cluster randomized trial extending adherence club ART refill dispensing intervals to 6-monthly. Abstract 201. In Mexico City, Mexico: 10th IAS Conference on HIV Science; 2019.
47. Roy M, Bolton C, Sikazwe I, Mukumbwa-Mwenechanya M, Efronson E, Somwe P, et al. Implementation and effectiveness of urban adherence clubs in Zambia. Abstract FRAE0104. In Amsterdam, Netherlands: 22nd International AIDS Conference; 2018.
48. Long L, Kuchukhidze S, Pascoe S, Nichols B, Huber A, Rosen S. Eligibility criteria for differentiated service delivery models of HIV treatment in sub-Saharan Africa: Evidence from a literature review 2016-2019. Boston, MA; 2020. Available from: <https://sites.bu.edu/ambit/files/2020/03/Brief-1-DSD-eligibility-from-AMBIT-review-2020-02-28-final.pdf>