

*Review*

# Factors Affecting Vitamin C Status and Prevalence of Deficiency: A Global Health Perspective

Anitra C. Carr <sup>1,\*</sup> and Sam Rowe <sup>2</sup><sup>1</sup> Nutrition in Medicine Research Group, Department of Pathology & Biomedical Science, University of Otago, Christchurch, 8011, New Zealand<sup>2</sup> Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, L35QA, UK  
sam.rowe@lstmed.ac.uk\* Correspondence: [anitra.carr@otago.ac.nz](mailto:anitra.carr@otago.ac.nz); Tel.: +64-3364-0649

**Abstract:** A recent review of global vitamin C status indicated a high prevalence of deficiency, particularly in low- and middle-income countries, as well as in specific subgroups within high-income countries. Here we provide a narrative of potential factors influencing vitamin C status. The in vivo status of vitamin C is primarily affected by dietary intake and supplement use. Dietary intake can be influenced by cultural aspects such as staple foods and traditional cooking practices. Environmental factors can also affect vitamin C intake and status; these include geographic region, season and climate, as well as pollution. Demographic factors such as sex, age, and race are known to affect vitamin C status, as do socioeconomic factors such as deprivation, education and social class, and institutionalization. Various health aspects affect vitamin C status; these include body weight, pregnancy and lactation, genetic variants, smoking, and disease states, including severe infections as well as various non-communicable diseases such as cardiovascular disease and cancer. Some of these factors have changed over time, therefore we also explore if vitamin C status has shown temporal changes. Overall, there are numerous factors that can affect vitamin C status to different extents in various regions of the world. Many of these factors are not taken into consideration during the setting of global recommended dietary intakes for vitamin C.

**Keywords:** vitamin C; vitamin C status; vitamin C deficiency; global health; dietary intake; obesity; smoking; communicable disease; infection; non-communicable disease

## 1. Introduction

Due to random genetic mutations that have occurred during our evolution, humans have lost the ability to synthesize ascorbate in our livers [1]. Therefore, instead of being a normal liver metabolite, as is the case for most other animal species, ascorbate has instead become a vitamin and must be obtained through the diet to prevent hypovitaminosis C and outright deficiency [2]. In a recent review we described the global status of vitamin C and prevalence of deficiency as assessed by plasma levels [3]. Whilst vitamin C levels are normally tightly controlled in humans with adequate intake a number of studies suggest that hypovitaminosis C and deficiency are not uncommon worldwide and may be very common in some settings. Vitamin C is well absorbed in small quantities. Uptake occurs primarily through the sodium-dependent vitamin C transporter-1 (SVCT-1) which is rapidly saturated allowing relatively limited absorption of vitamin C per serving [4]. Vitamin C is then concentrated at varying levels in different body tissues [5]. The vitamin is excreted unchanged in the urine, but re-uptake occurs in the renal tubules [6]. Further loss of vitamin C occurs through the oxidation of ascorbic acid to dehydroascorbic acid which may then be recycled to ascorbic acid or undergo further oxidative decomposition [7]. Thus, in vivo vitamin C levels are affected by a range of factors that include dietary intake, absorption, distribution, oxidative decomposition and elimination.

Vitamin C acts as an important antioxidant and plays a myriad of functions in optimal health and prevention of disease [8, 9]. Even mild insufficiency or hypovitaminosis C can be associated with symptoms such as low mood [10]. More severe deficiency can be associated with a range of clinical presentations [11]. Prolonged severe deficiency results in the clinical syndrome of scurvy a condition that continues to be diagnosed in individuals and occasionally in public health outbreaks to this day [12]; if left untreated scurvy is fatal disease. The optimum intake and levels of vitamin C, however, have been subject to debate and recommended values are not universal [13]. Due to a range of benefits, a number of international authorities have increased their dietary recommendations for vitamin C from those previously recommended on the basis of preventing scurvy [13]. Although these recommendations generally take into account variations in requirements based on age, sex, pregnancy and lactation, and sometimes smoking status, there are many other factors that affect vitamin C status that are not taken into consideration by these global authorities. In this review we highlight the numerous factors that impact on vitamin C status and prevalence of deficiency globally.


2. Dietary factors determining vitamin C status

2.1. Dietary intake

Dietary intake of vitamin C is a key determinant of body status, with the amount consumed and the frequency of consumption correlating with plasma status and prevalence of deficiency (Table 1) [14, 15]. Fresh fruit and vegetables are the major dietary source of vitamin C, with fruit intake correlating more strongly with plasma vitamin C status [16, 17]. Foods rich in vitamin C include oranges and other citrus, guava, kiwifruit, cranberries, strawberries, papaya, mango, melon, cantaloupe, spinach, Swiss chard, tomato, asparagus, and Brussels sprouts (Figure 1) [18]. In contrast, cereals (e.g. rice, millet, wheat/couscous, corn), some starchy roots and tubers, meat (other than liver), eggs and dairy contain very little vitamin C.

Table 1. Dietary factors determining vitamin C status.

Factor	Summary and comments	References
Dietary intake	Dietary intake, particularly fruit intake, correlates with improved vitamin C status and decreased prevalence of deficiency. Depends on amount consumed, frequency of consumption, and type of food consumed as vitamin C content of food varies. High dietary fat and sugar intake is associated with reduced vitamin C intake and status.	[14-17, 19-24]
Staple foods	Staple foods such as cereals (e.g. rice, millet, wheat/couscous, corn) and some starchy roots and tubers are low in vitamin C; populations who consume these staples can have lower overall vitamin C intake.	[18, 25]
Traditional cooking practices	Through boiling or steaming water soluble vitamins may be leached from food and prolonged cooking of food can destroy vitamin C; this could lead to decreased vitamin C status in certain social or ethnic group. Drying of leafy vegetables reduces water soluble vitamins.	[26, 27]
Supplement use	Supplement users have significantly higher vitamin C status and negligible prevalence of deficiency. Non-users have a 2-3 fold odds ratio of insufficient and deficient vitamin C status.	[14, 15, 19, 28-30]



**Figure 1.** Typical vitamin C content of selected fruit and vegetables. Data derived primarily from the United States Department of Agriculture (<https://fdc.nal.usda.gov/>). Note that vitamin C content can vary depending on the plant variety and cooking may decrease the vitamin C content to variable extents. Pulses include kidney beans, chickpeas, mung beans, pinto beans, soy beans, lentils, peanuts, split peas; nuts include hazelnuts, pistachios, macadamia nuts, pecans, walnuts, brazil nuts, cashew nuts; seeds include chia, flax seeds, pumpkin seeds, sunflower seeds, sesame seeds; grains include rice, millet, wheat/couscous, cornmeal. Animal products, such as meat (other than liver), eggs and milk contain negligible vitamin C. Dotted lines: lower line indicates daily intake (10 mg/d) to prevent scurvy; upper line indicates daily intake (200 mg/d) for optimal health.

Suboptimal diet is an important preventable risk factor for non-communicable diseases and low intake of fruit was found to be the third leading dietary risk for deaths (2 million deaths) and disability-adjusted life-years (65 million DALYs) globally [31]. Historically, scurvy has been reported in refugee camps in Ethiopia; the relief food distributed to the refugees was almost completely deficient in vitamin C and the environment where the camps were located precluded an adequate supply of fresh food [32]. However, vitamin C deficiency may even be common in fertile areas of Africa such as Uganda [33]. There have been a number of outbreaks of clinical scurvy in recent years, including in tribesmen in Kenya and during the Afghanistan winter [34, 35], and cases of clinical scurvy continue to be diagnosed related to poor intake even in high income countries [12, 36].

## 2.2. Cultural aspects: staple foods and traditional cooking practices

Food consumption is influenced by a number of factors, such as geographical, economical, social and cultural [25]. It is well known that economic factors, such as poverty, can limit the consumption of healthy foods. Furthermore, tradition and social customs may influence the consumption of a variety of foods. For example a study in Nepal highlighted cultural beliefs around menstruation, pregnancy and lactation that reduced micronutrient intake and intrahousehold disfavouritism towards women in distribution of micronutrient rich foods [27]. In LMICs, geographic considerations and agricultural practices influence the production and consumption of staple foods, which can greatly influence vitamin C intake. For example, countries whose populations consume cereals, such as rice and millet, as staple foods (e.g. in Asia and parts of Africa) tend to have lower intakes of vitamin C [18, 25]. In contrast, some areas of Africa and Latin America where yams, cassava and sweet potato are staple foods, vitamin C intakes may be higher. However, the vitamin C content of food can also vary depending on time of harvest, transportation and storage conditions, and food preparation conditions. For example, the vitamin C content of some staple foods, e.g. cassava, is significantly depleted by different processing methods, which could lead to inaccuracies in estimated vitamin C intake [37]. Vitamin C is heat-labile and can be destroyed by cooking; the lower vitamin C status of Indians and Malays is thought to be partly due to its destruction by more prolonged cooking [26].

## 2.3. Supplement use

Although it is possible to obtain an optimal vitamin C intake (i.e. 200 mg/d) from five plus daily servings of fresh fruit and vegetables, providing that at least one or two servings are high vitamin C foods, this is not always possible for many people. Therefore, taking supplements in addition to dietary intake can help to maintain optimal vitamin C status. There are many different forms of vitamin C supplements, however, research indicates that these generally have the same bioavailability as food-derived vitamin C [38]. Numerous studies have found that even in developed countries, where there is no shortage of fresh fruit and vegetables, those who consume supplements have significantly higher vitamin C status and/or lower prevalence of deficiency (Table 4). Two large health surveys carried out in Canada (CHMS) and the USA (NHANES IV) reported 20% supplement use in Canada and 37 to 47% supplement use, for men and women respectively, in the USA [14, 15]. It should be noted, however, that in these surveys supplement use was considered to be as little as once in the previous month. Nevertheless, both studies found vitamin C status was at least 20  $\mu\text{mol/L}$  higher in supplement users, with a low prevalence of deficiency (0-2%) in those who supplemented [14, 15]. Two studies in the UK indicated that people who did not supplement had a two-fold odds ratio of having vitamin C status < 28  $\mu\text{mol/L}$  and a three-fold odds ratio of vitamin C status being < 11  $\mu\text{mol/L}$  relative to >28  $\mu\text{mol/L}$  [19, 28].

## 3. Environmental factors affecting vitamin C status

### 3.1. Geographic region

Geographic differences in vitamin C status have been reported (Table 2); there are likely many factors that contribute to the observed differences. For example, significant differences in vitamin C status and prevalence of deficiency were observed between Finland and neighboring Russia (where strikingly low plasma levels were observed). Geographical differences in the consumption of fresh fruit and vegetables were apparent which were associated with deficiency. Low educational status was also noted to be associated with deficiency on the Russian side of the border [17]. Another study carried out in five countries across Europe indicated a difference in the vitamin C status of women, with the biggest differences being observed between Northern Ireland and the Republic of Ireland [39]. No differences in vitamin C status were observed for men from these different countries, whilst women had higher vitamin C status than the men in three of the five countries. Mosdol et al. [19] reported that low income participants living in Scotland and Northern Ireland had a lower prevalence

of deficiency than those living in England, whilst those living in Wales had a higher prevalence of deficiency. As part of the SU.VI.MAX trial, France was divided into seven regions and vitamin C status was found to be significantly lower in the Northern region. The reasons for these regional differences are unknown [40]. In a large study of elderly people in India, a higher prevalence of deficiency was observed in the North (74% deficiency) compared with the South (46% deficiency) [24]. Similarly, in Mexico, a higher prevalence of deficiency was observed in children in the North and South regions (~26% deficiency) compared with those living in Mexico City (12% deficiency) [41]. This could reflect differences in the socioeconomic status between urban and rural environments.

**Table 2.** Environmental factors affecting vitamin C status.

Factor	Summary and comments	References
Geographical region	Vitamin C status varies by geographical region, both within and between countries; this could partly reflect differences in socioeconomic status and available foods.	[17, 19, 24, 39-43]
Season	Vitamin C status varies seasonally between countries, likely reflecting different crops and thus the types and/or amounts of vitamin C-rich foods consumed.	[24, 30, 44, 45]
Climate	Drought and harsh winter climates have been associated with recent outbreaks of clinical scurvy.	[35]
Pollution	Exposure to environmental pollutants, e.g. smoke, can deplete vitamin C status; this is partly due to enhanced oxidative stress.	[24, 46-51]

3.2. *Season and climate*

Since fresh fruit and vegetables are the major source of vitamin C for most people, it is perhaps not surprising that there have been reports of seasonal differences in vitamin C status (Table 2). Surprisingly, however, studies carried out in England and China have indicated that vitamin C status tends to be highest in winter and lowest in autumn, with up to 10 µmol/L difference between these seasons [30, 44]. Another study in China indicated much higher vitamin C status in winter relative to spring [45]. This likely reflects the types or amounts of vitamin C-rich foods being consumed in winter. Similar trends were observed in Northern India, with less deficiency being observed in the winter months, however, in Southern India, the winter months were associated with higher prevalence of deficiency [24]. This likely reflects the different climatic and agricultural patterns across the sub-continent. It is possible that seasonal variances in plasma vitamin C levels are affected by baseline vitamin C status, i.e. whether the individual is already saturated or not. Bates et al. [52] reported seasonal variations in participants with higher intakes and blood status at baseline compared with little variation in those with low intakes and status, the latter likely due to a depleted body pool and preferential uptake of the vitamin by tissues. Changes in climate, such as drought, are also likely to impact on the vitamin C status of the populations reliant on local foods in the affected region [35]. This also leads to increased reliance on staple crops like cassava that lose their vitamin content prior to consumption.

3.3. *Pollution*

Currently, over half of the world’s population lives in urban areas and WHO data indicate that more than 80% of people living in urban areas are exposed to air quality levels that exceed WHO guideline limits, with LMIC suffering from the highest exposures [53, 54]. Air pollution causes an estimated seven million premature deaths worldwide every year, primarily resulting from increased mortality from strokes, heart disease, chronic obstructive pulmonary disease, lung cancer and acute respiratory infections [54]. Environmental air pollution, such as smoke derived from burning biomass, can comprise reactive species that potentially affect in vivo antioxidant status [24]. Another example

is environmental tobacco smoke, which is an underestimated pollutant in many parts of the world. Exposure to environmental tobacco smoke is associated with depleted vitamin C status in both non-smoking adults and children [46-51]. Tribble et al. [46] found that the vitamin C status of passive smokers was significantly lower than that of nonexposed nonsmokers, despite comparable dietary intakes of the vitamin. Hypovitaminosis C was observed in 12% of passive smokers, but not in nonexposed nonsmokers. Analysis of the data of children from the NHANES III and IV surveys revealed a dose-response relationship between levels of tobacco exposure and serum vitamin C concentrations [48, 49]. Not all studies in adults, however, have shown effects of passive smoking on vitamin C status [55-58]. Nevertheless, supplementation with vitamin C and other antioxidants was found to decrease oxidative biomarkers in participants exposed to environmental tobacco smoke [55, 59].

4. Effect of demographic factors on vitamin C status

4.1. Sex

In High income settings females appear to have higher vitamin C status and a lower prevalence of deficiency than men (ref: vitC status paper or CRFSN rev). According to McCall et al. [28], UK males have a four-fold odds ratio of deficiency compared with females (Table 3). The difference in vitamin C status between males and females is thought to be partly a result of a volumetric dilution effect due to the higher fat free mass of males [29]. There are also differences in dietary intakes between men and women, with women generally having comparable or higher intakes than men in high income countries, although comparable data on this is limited from low income settings [3]. It should also be noted that pregnancy and lactation typically lower women’s vitamin C status due to hemodilution and the needs of the developing fetus and growing infant (see below). Many countries have higher dietary recommendations for men, to take into account the differences in body mass, and pregnant and lactating women [13].

Table 3. Effect of demographic factors on vitamin C status.

Factor	Summary and comments	References
Sex	Males generally have lower vitamin C status, and a higher prevalence of deficiency, than females; this is partly a result of a volumetric dilution effect due to the higher fat free mass of males. This difference is less apparent in some low- and middle-income countries.	[14, 15, 17, 19, 22, 24, 26, 28-30, 39, 40, 45, 47, 60-64]
Age	Both children and elderly tend to have higher vitamin C status in high-income settings; this could partly be due to lower body weight. Elderly can have a higher prevalence of vitamin C deficiency in some settings; this could be due to lower intake and/or comorbidities.	[14, 15, 20, 24, 28, 29, 40, 43, 65, 66]
Race	In the US and UK, African-Caribbean’s and South Asians had lower status than Caucasians. In South Asia, Malays and Indians had lower status than Chinese; this is thought to be partly due to differences in culinary practices. Differences are more apparent between women of different races.	[15, 26, 30, 67, 68]

4.2. Age

Some studies carried out in people aged >60 years (e.g. the French POLA study and the British National Diet and Nutrition Survey) have shown lower vitamin C status than other studies carried out in the same countries with younger age groups (e.g. the French SU.VI.MAX study and the European EPIC-Norfolk study), however, it is less than ideal to compare values between different

studies due to potential methodological differences. A number of studies have indicated that older age within the same study population is associated with an increased prevalence of vitamin C deficiency, particularly in men, and older men also tend to have a lower vitamin C intake than older women [20, 24, 28, 65]. However, a counterpoint to this is that overall vitamin C status can be higher in older people [15, 29, 40], although not in all cases as was found in men in Finland who had lower vitamin C status with older age [66]. Schleicher et al. found a U-shaped curve for vitamin C status over the age range of 6 to 60+ years [15]. It is possible that the lower body mass of children and elderly, due to age-related frailty, could contribute to their higher vitamin C status. However, clearly there are subpopulations within the aging population who have increased deficiency, likely due to lower intake and/or comorbidities.

#### 4.3. Race

A number of studies have indicated that vitamin C status varies by race (Table 3; reviewed in [68]). The prevalence of vitamin C deficiency appeared to be highest amongst South Asians. This is thought to be partly due to traditional cooking practices. It was also suggested that the high proportion of individuals with low concentrations of vitamin C in South Asian populations might contribute to high rates of cardiovascular disease. Only a few studies have assessed different ethnic groups within the same study [15, 26, 30, 67]. In 1980 Koh et al. [67] assessed vitamin C status in a cohort of black and white participants in Mississippi, USA. Black males and females had significantly lower vitamin C status than white males and females (up to 10  $\mu\text{mol/L}$  difference for females). The more recent US NHANES IV survey also found significantly lower vitamin C status in black females compared with white females [15]. In London, the vitamin C status of black and South Asians was significantly lower than white participants (up to 15  $\mu\text{mol/L}$  lower for South Asian females) [30]. Similarly, in Singapore Malays and Asian Indians were found to have significantly lower vitamin C status than Chinese (up to 12  $\mu\text{mol/L}$  for females) [26]. Studies in Uganda, South Africa and Nigeria all showed high rates of deficiency in predominately black Africans (Table 2).

### 5. Effect of socioeconomic factors on vitamin C status

#### 5.1. Socioeconomic status/deprivation

Socioeconomic status affects diet quality as foods of higher quality and higher nutritional value generally cost more [69]. Therefore, it is not surprising that numerous studies have shown that lower socioeconomic status and deprivation negatively impact on vitamin C status (Table 4). In a UK study, McCall et al. [28] reported that those who were most deprived (based on the Townsend deprivation index) had a two-fold odds ratio of vitamin C deficiency relative to those with plasma values above 28  $\mu\text{mol/L}$ . This finding was supported by Mosdol et al. [19] who specifically focused on the low-income population within the UK, and showed a higher prevalence of deficiency and insufficiency than has been reported for the general UK population (Table 1). Bates et al. [43] reported a three-fold difference in plasma vitamin C status for people of low socioeconomic status in the UK relative to those of higher status. A two-fold difference in vitamin C intake between the two socioeconomic groups likely contributed to the observed differences in vitamin C status. Other more recent surveys have confirmed the association of low vitamin C status with low socioeconomic status [15, 61].v

#### 5.2. Education and social class

Similar findings have been reported for level of education, with the lowest level of education associated with the lowest vitamin C status [14, 17]. McCall et al. [28] reported that those with the lowest level of education in the UK had a 4.5 fold odds ratio of deficiency relative vitamin C status >28  $\mu\text{mol/L}$ . The same authors investigated social class and found that those having manual occupations had a three-fold odd ratio of deficiency compared with those having non-manual occupations [28]. Similar findings were reported by Wreiden et al. [20], with two- to three-fold higher prevalence of vitamin C deficiency for females and males, respectively, with manual occupations.

**Table 4.** Effect of socioeconomic factors on vitamin C status.

Factor	Summary and comments	References
Socioeconomic status/deprivation	Individuals with lower socioeconomic status or higher deprivation have lower vitamin C status and a higher prevalence of deficiency; this is partly due to the higher cost of good quality, nutrient dense food.	[15, 19, 21, 24, 28, 41, 43, 47, 61, 70]
Education and social class	Similarly, individuals with lower education, and manual occupations, have lower vitamin C status.	[14, 17, 20, 28, 47]
Institutionalized	Institutionalized elderly, and other institutionalized individuals (e.g. priests, prisoners, boarding school children), have lower vitamin C status and higher prevalence of deficiency; this is partly due to lower dietary intake.	[43, 71-73]

5.3. Institutionalization

One survey of elderly people assessed institutionalized individuals relative to free-living individuals [43]. The authors found significantly lower vitamin C status (25 vs 44  $\mu\text{mol/L}$ ) and a higher prevalence of deficiency (40 vs 14%) in institutionalized elderly compared with free-living elderly, respectively. Much of this is likely due to a higher proportion of individuals consuming less than the recommended nutrient intake for vitamin C in institutional settings [43]. This will become a growing concern with an increasingly ageing population. Other institutionalized individuals, e.g. priests, prisoners, and boarding school children, have also been found to have lower vitamin C status, higher hypovitaminosis C and severe deficiency, compared to control groups [71-73].

6. Health aspects that affect vitamin C status

6.1. Body weight and body mass index (BMI)

Body weight, and the related BMI, waist circumference or waist-to-hip ratio, are well known to have a significant association with vitamin C status (Table 5). Numerous studies have shown an inverse association between body weight or BMI and vitamin C status, with obese individuals having the lowest vitamin C status [14-16, 29, 30, 60, 74]. Schleicher et al. [15] reported up to a 15  $\mu\text{mol/L}$  difference in the vitamin C status of obese women compared with women of healthy weight (i.e. 45 versus 60  $\mu\text{mol/L}$ , respectively). Pearson et al. [61] found that individuals with hypovitaminosis C (i.e. < 23  $\mu\text{mol/L}$ ) had significantly higher weight, BMI and waist circumference. Lower vitamin C status in those of highest weight could be partly due to lower dietary intake of the vitamin [74, 75]. In support of this, lower vitamin C status was found in those with highest fat intake [23]. It has also been suggested that the observed differences in vitamin C status could be a result of volumetric dilution due to differences in fat-free mass [29]. This premise is supported by supplementation studies that have indicated that individuals with higher body weight do not replete as readily as those of normal body weight [76, 77].

Based on these observations, Block et al. [76] proposed that recommended vitamin C intakes should be based on a dose per kg body weight or in terms of an ideal plasma concentration. This is a prudent recommendation considering the ongoing increase in body weight globally, with the prevalence of obesity exceeding 50% in some countries [78, 79]. Obesity is a risk factor for numerous diseases, particularly cardiometabolic diseases such diabetes and cardiovascular disease, which are also associated with lower vitamin C status and higher prevalence of deficiency [9]. Enhanced dietary fat and sugar, which are risk factors for cardiometabolic diseases, are also associated with decreased vitamin C intake and status [22, 23].

**Table 5.** Health aspects that affect vitamin C status.

Factor	Summary and comments	References
Body weight, BMI	Individuals with higher body weight or BMI have lower vitamin C status; this is partly due to a volumetric dilution effect.	[14-16, 24, 29, 30, 60, 61, 74, 75, 80, 81]
Physical activity	Physical activity level positively correlates with vitamin C status, with inactive individuals having a 3-fold odds ratio of deficiency; this is partly due to associated lifestyle factors such as diet and body weight.	[28, 29]
Pregnancy and lactation	Pregnancy is associated with lower vitamin C status; this is partly due to hemodilution and active transfer of vitamin C to the developing fetus and growing infant via breastmilk.	[33]
Genetic variants	Genetic polymorphisms in the genes for the vitamin C transporter (SVCT1) and haptoglobin (Hp2-2) are associated with lower vitamin C status; the latter is thought to be due to enhanced oxidative stress.	[6, 70, 82-85]
Smoking	Smokers have lower vitamin C status, and a higher prevalence of deficiency, than non-smokers; this is partly due enhanced oxidative stress.	[14-16, 19, 20, 24, 28-30, 40, 45, 46, 60, 61, 65, 70, 86-89]
Disease states	Various communicable and non-communicable diseases are associated with lower vitamin C status; this is partly due to inflammatory processes and enhanced oxidative stress.	[9, 90-92]

### 6.2. Pregnancy and lactation

Pregnant women typically have lower vitamin C status compared with non-pregnant women [33]. This is most likely due to hemodilution as well as active transfer of the vitamin to the developing fetus [93]. Women with complications of pregnancy can have even lower vitamin C status [33, 94]. Supplementation of pregnant women with vitamin C can potentially decrease the risk of some pregnancy-related complications [95]. Studies in high-income countries, which included women with normal vitamin C status, have failed to show benefit from supplementation [96]. However, studies in low income settings with high rates of vitamin deficiency show potential benefits including decreased rates of low birth weight, decreased hospital admissions and possible decreased rates of pre-eclampsia [97]. It is noteworthy that the recently discovered epigenetic regulatory activities of vitamin C could have important roles to play in fetal development [98]. It is likely that lactating women also have lower vitamin C status due to transfer of vitamin C to the growing infant via breastmilk. Many global authorities have taken the enhanced requirements of pregnant and lactating women into consideration with recommendations above their standard dietary recommendations of +10-20 mg/d for pregnant women and +20-60 mg/d for lactating women [13].

### 6.3. Genetic variants

Vitamin C status can potentially be influenced by genetic variants. The SLC23A1 gene encodes the sodium-dependent vitamin C transporter-1 (SVCT1) which is responsible for active uptake of dietary vitamin C through the intestinal epithelium and reuptake of filtered vitamin C via the kidney tubules, and is essential in vitamin C-requiring animals [4, 6]. A number of single nucleotide variants have been identified in the SLC23A1 gene and modelling of in vitro data for four of these indicate 40 to 75% decreases in ascorbate uptake [6, 99]. These variants are relatively common in those of African descent (6-16%), and the variant with the largest decrease in vitamin C transport has a high

prevalence in African Americans [6]. Meta-analysis has indicated a strong association of another of these variants with decreased vitamin C status in five cohorts in the UK (frequency of ~3%) [70]. In the British Women's Heart and Health Study a further two variants, with relatively common frequencies of ~30%, were also found to be associated with vitamin C status [70]. Cahill et al. [82] reported that variants of SVCT1 can also modify the strength of the correlation between dietary vitamin C and serum vitamin C.

Vitamin C status could be further influenced by genetic variants affecting the metabolism of the vitamin [100]. A common variant of the hemoglobin-binding protein haptoglobin (Hp2-2) has a decreased ability to bind hemoglobin and results in increased oxidation of vitamin C in vitro [83]. Several studies have shown that this variant is associated with lower circulating vitamin C concentrations [83-85]. The Hp2-2 variant appeared to have a greater effect in individuals with dietary vitamin C intakes <90 mg/day [82]. It is noteworthy that genetic variants can show marked geographical differences; e.g. Hp2-2 is present in ~36% Northwest Europeans, 51% Iranians, 55% Thais and Chinese, and 84% Indians (where it is thought to have originated) [101]. Other genetic variants, such as those of the detoxifying enzyme glutathione S-transferase, may also have associations with vitamin C status, particularly in individuals with lower intakes, thereby resulting in an increased risk of deficiency [102]. High-dose vitamin supplementation has been shown to ameliorate certain gene variant defects [103], and is hypothesized to occur with vitamin C-related variants [104]. Therefore, individuals with genetic variants affecting vitamin C status may require higher dietary intakes.

#### 6.4. Smoking

Smoking is a well known source of oxidants and in vivo oxidative stress [105]. Multiple studies have shown depleted vitamin C status and a higher prevalence of deficiency in smokers compared with non-smokers (Table 4). McCall et al. [28] reported that current smokers had over a seven-fold odds ratio of deficiency compared with non-smokers, and Wrieden et al. [20] showed two- to three-fold more deficiency in male and female smokers, respectively. Smokers tended to have dietary intakes that were lower in vitamin C e.g. lower fruit and vegetable intake and higher fat intake [86, 87, 106, 107]. However, when differences in dietary intake were taken into account, smokers still exhibited lower vitamin C status and higher requirements than non-smokers [86, 87, 108, 109]. Kallner et al. [110] demonstrated over a 40% increase in vitamin C turnover in smokers compared with nonsmokers.

A few international regulatory authorities have taken the enhanced requirements of smokers into consideration with additional intake recommendations of 20 to 80 mg/d above their normal adult recommendations [13]. However, it is likely that these additional intakes are insufficient to compensate for the enhanced requirements of smokers [88]. Furthermore, some countries continue to show an increased trend in smoking rates which could potentially impact on their population vitamin C status and requirements [111]. Smoking cessation results in an increase in vitamin C status [112].

#### 6.5. Disease states

Vitamin C status can be depleted by various disease states due to inflammatory processes and enhanced oxidative stress (reviewed in [9, 90, 91]). A number of studies of hospitalized patients showed a high prevalence of depleted plasma vitamin C status, and concentrations were inversely correlated with inflammatory markers [113-116]. A wide range of medical conditions are associated with vitamin C insufficiency; these include non-communicable diseases such as cardiovascular diseases (e.g. strokes, coronary artery disease and hypertension), congestive heart failure, malignancy, chronic inflammatory states (e.g. rheumatoid arthritis), metabolic disorders (e.g. diabetes), and cataracts [9, 90, 91]. Acute infectious diseases leading to enhanced inflammation are also associated with depleted plasma vitamin C concentrations in plasma and immune cells, as are a range of chronic infections such as HIV, *Helicobacter pylori* and tuberculosis, which are prevalent in many LMICs [92, 117]. It should also be noted that requirements for vitamin C during infections increase with the

severity of the infection, requiring significantly enhanced intakes to reach normal plasma status [92]. This is particularly pertinent to the current global coronavirus (SARS-CoV-2) pandemic which causes severe pneumonia and sepsis, conditions known to be associated with depleted vitamin C status [118, 119]. It should also be noted that people with marginal vitamin C status are at higher risk of developing vitamin C deficiency [120], and once people have become depleted, higher than recommended intakes of the vitamin are required to fully replete them again [76, 77].

## 7. Has vitamin C status changed over time?

Many factors have changed over time that could potentially affect vitamin C status. These include population demographics and health status, e.g. obesity and smoking rates. Changing climate also affects what crops can be grown in different regions of the world. Public health policies, such as recommended dietary intakes, have changed in many countries [13]. Therefore, have these temporal changes been reflected in changing vitamin C status over time? There have been relatively few studies that have assessed vitamin C status in the same populations or regions over time. The findings can also be complicated by changes in analytical methodologies over time. Two surveys have been carried out in elderly people in Britain; one in 1979 by the Department of Health and Social Security, and the National Diet and Nutrition Survey in 1998 [43]. There was an increase in vitamin C status from 29 to 44  $\mu\text{mol/L}$  between the two surveys. However, it should be noted that participants in the first survey were older, and analytical methodologies may have varied between the two surveys.

Schleicher et al. [15] carried out an in-depth comparison of US vitamin C status in NHANES IV (2003–2004) relative to NHANES III that was undertaken 10–15 years earlier (i.e. 1988–1994) [121]. For those aged  $\geq 20$  years, overall vitamin C deficiency decreased by about 44% (from 15 to 8%) between the two surveys. Of note, there was an increase in the recommended dietary intake for vitamin C in the US in 2000, from 60 mg/d to 90 mg/d for men and 75 mg/d for women [122]. However, it is unlikely that this increase in dietary recommendations fully accounts for the decrease in vitamin C deficiency over time as there were also some differences in methodology between the two surveys, with the more recent survey likely being more accurate [15]. Looking back even earlier to the NHANES II survey (carried out between 1976 and 1980), vitamin C status was reported to be higher for both men and women (i.e.  $\geq 50 \mu\text{mol/L}$  for men and  $\geq 60 \mu\text{mol/L}$  for women), with 5% deficiency reported, despite lower dietary intakes compared to the later NHANES III survey [123]. It should be noted, however, that vitamin C in the NHANES II samples was analysed using the 2,4-dinitrophenylhydrazine method, which can be prone to interference by other components in plasma and thus potentially overestimate vitamin C concentrations [124].

Other time course data is available from studies carried out in Finland and neighboring Russia between 1992 and 2002 [17, 42]. In North Karelia, Finland, the vitamin C status of men decreased from 54 to 27  $\mu\text{mol/L}$  over 5 years, then rose slightly to 37  $\mu\text{mol/L}$  over the next 5 year period [17]. Nyyssonen et al. reported a small increase in baseline vitamin C values in Finnish men over 6 years [66]. In Pitkäranta, Russia, the vitamin C status of women decreased from 22 to 13  $\mu\text{mol/L}$  over a 5 year period, while the vitamin C status of Russian men remained in the deficiency range over a 10 year period [17]. The samples from the more recent survey (2002) were from fasting participants, however, it is not clear whether they were fasting in the earlier surveys (1992 and 1997).

Another set of studies was carried out in Java, Indonesia between 2005 and 2011 [125, 126]. Mean vitamin C status decreased in the subjects from 45 to 29  $\mu\text{mol/L}$  over 6 years, although the percentage of deficiency decreased from 15 to 11% over this time. The decrease in mean vitamin C status could partly be explained by the earlier study assessing non-fasting plasma (as demonstrated by a higher percentage of vitamin C status  $>23 \mu\text{mol/L}$ , i.e. 72 versus 54%). However, there are possibly other environmental, lifestyle or health-related issues that could be responsible for the changes in vitamin C status over time in Indonesia, Finland and Russia.

Not all studies have shown changes in vitamin C status over time. A survey carried out in Shanghai, China, over a five year period (between 1995 and 2000) showed no change in vitamin C status over this time period, although there was a dip in status at the midpoint (from 43 to 34  $\mu\text{mol/L}$ )

[44]. Thus, there is currently no clear trend in vitamin C status over time. However, much of the vitamin C data is very dated, and thus may not accurately reflect current status, particularly in countries where there have been major changes in factors that can affect vitamin C status.

## 8. Conclusions and future directions

This narrative review describes the findings of numerous studies which highlight the various factors that can impact on vitamin C status and prevalence of deficiency. The studies were of variable quality; ideally multivariate analysis of the various factors highlighted in these studies would be performed to further examine the relationships discussed.

Global vitamin C status appears to be a cause for concern and we believe this is primarily due to poor diet, particularly in low- and middle-income countries, although also apparent in subgroups within high-income settings [3]. Vitamin C levels provide a useful biomarker for a healthy diet, unfortunately accurate analysis is costly, technically challenging and time consuming [127]. Thus, there is a need for low cost accurate commercially available methods to assess plasma levels. Equally, current assessment of intake is also limited and fails to take into account that the vitamin may be almost entirely eliminated prior to consumption in some globally important crops, such as cassava [37]. Future amendments to nutrient reference tables should take this aspect into account.

Clearly, there is a global need for increased consumption of vitamin C rich fresh fruit and vegetables. From our previous review, this appears to be most notable in LMIC's, however it is also a concern in other at risk groups [3]. Further education is required globally in how to meet the daily vitamin C requirements with locally available crops year round. Some staple foods such as cassava require prolonged cooking to remove toxins (in this case cyanide); efforts have been made to reduce this content in genetically modified cassava, which could allow higher vitamin C contents from this vitamin rich plant at the time of consumption [37]. Given the high rates of deficiency in some diets, cooking practices may also require modification to help increase intake.

A number of studies have assessed the potential for vitamin C supplements to reduce a range of non-communicable disease and infections [9, 92]. However, there remains much controversy as to the efficacy of supplementation, primarily due to badly designed studies that do not take into account the baseline vitamin C status of the participants [96]. Therefore, knowledge of the high-risk groups for deficiency should be utilised for future studies as these groups are likely to gain more benefit.

In the meantime, global and local policy makers should consider the local data available on deficiency in an attempt to modify dietary intake and other modifiable risk factors. More of these risk factors should be taken into account during review and update of global recommended dietary intakes for the vitamin. Clinicians worldwide should also remain vigilant to detect and correct hypovitaminosis C and deficiency in the at risk groups highlighted.

**Author Contributions:** Both authors contributed equally to the review. All authors have read and agreed to the published version of the manuscript.

**Acknowledgments:** A.C. is a recipient of a Sir Charles Hercus Health Research Fellowship from the Health Research Council of New Zealand.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Nishikimi M, Yagi K. Molecular basis for the deficiency in humans of gulonolactone oxidase, a key enzyme for ascorbic acid biosynthesis. *Am J Clin Nutr.* 1991;54(6 Suppl):1203S-8S.
2. Stone I. Homo sapiens ascorbicus, a biochemically corrected robust human mutant. *Med Hypotheses.* 1979;5(6):711-21.

3. Rowe S, Carr A C. Global vitamin C status and prevalence of deficiency: A cause for concern? Preprints. 2020;202005.0394.v1.
4. Savini I, Rossi A, Pierro C, Avigliano L, Catani MV. SVCT1 and SVCT2: key proteins for vitamin C uptake. *Amino Acids*. 2008;34(3):347-55.
5. Hornig D. Distribution of ascorbic acid, metabolites and analogues in man and animals. *Ann N Y Acad Sci*. 1975;258:103-18.
6. Corpe CP, Tu H, Eck P, Wang J, Faulhaber-Walter R, Schnermann J, Margolis S, Padayatty S, Sun H, Wang Y, Nussbaum RL, Espey MG, Levine M. Vitamin C transporter Slc23a1 links renal reabsorption, vitamin C tissue accumulation, and perinatal survival in mice. *J Clin Invest*. 2010;120(4):1069-83.
7. Washko PW, Wang Y, Levine M. Ascorbic acid recycling in human neutrophils. *J Biol Chem*. 1993;268(21):15531-5.
8. Carr A, Frei B. Does vitamin C act as a pro-oxidant under physiological conditions? *FASEB J*. 1999;13(9):1007-24.
9. Carr AC, Frei B. Toward a new recommended dietary allowance for vitamin C based on antioxidant and health effects in humans. *Am J Clin Nutr*. 1999;69(6):1086-107.
10. Levine M, Conry-Cantilena C, Wang Y, Welch RW, Washko PW, Dhariwal KR, Park JB, Lazarev A, Graumlich JF, King J, Cantilena LR. Vitamin C pharmacokinetics in healthy volunteers: evidence for a recommended dietary allowance. *Proc Natl Acad Sci U S A*. 1996;93(8):3704-9.
11. Deirawan H, Fakhoury JW, Zarka M, Bluth MH, Moossavi M. Revisiting the pathobiology of scurvy: a review of the literature in the context of a challenging case. *Int J Dermatol*. 2020.
12. Ceglie G, Macchiarulo G, Marchili MR, Marchesi A, Rotondi Aufiero L, Di Camillo C, Villani A. Scurvy: still a threat in the well-fed first world? *Arch Dis Child*. 2019;104(4):381-3.
13. Carr AC, Lykkesfeldt J. Discrepancies in global vitamin C recommendations: A review of RDA criteria and underlying health perspectives. *Crit Rev Food Sci Nutr*. 2020;1-14.
14. Langlois K, Cooper M, Colapinto CK. Vitamin C status of Canadian adults: Findings from the 2012/2013 Canadian Health Measures Survey. *Health Rep*. 2016;27(5):3-10.
15. Schleicher RL, Carroll MD, Ford ES, Lacher DA. Serum vitamin C and the prevalence of vitamin C deficiency in the United States: 2003-2004 National Health and Nutrition Examination Survey (NHANES). *Am J Clin Nutr*. 2009;90(5):1252-63.
16. Drewnowski A, Rock CL, Henderson SA, Shore AB, Fischler C, Galan P, Preziosi P, Hercberg S. Serum beta-carotene and vitamin C as biomarkers of vegetable and fruit intakes in a community-based sample of French adults. *Am J Clin Nutr*. 1997;65(6):1796-802.
17. Paalanen L, Prattala R, Alfthan G, Salminen I, Laatikainen T. Vegetable and fruit consumption, education and plasma vitamin C concentration in Russian and Finnish Karelia, 1992-2002. *Public Health Nutr*. 2014;17(10):2278-86.
18. Joint FAO/WHO expert group. Human Vitamin and Mineral Requirements. Rome: Food and Agriculture Organization; 2001.
19. Mosdol A, Erens B, Brunner EJ. Estimated prevalence and predictors of vitamin C deficiency within UK's low-income population. *Journal of public health (Oxford, England)*. 2008;30(4):456-60.
20. Wrieden WL, Hannah MK, Bolton-Smith C, Tavendale R, Morrison C, Tunstall-Pedoe H. Plasma vitamin C and food choice in the third Glasgow MONICA population survey. *J Epidemiol Community Health*. 2000;54(5):355-60.

21. Madruga de Oliveira A, Rondo PH, Mastroeni SS, Oliveira JM. Plasma concentrations of ascorbic acid in parturients from a hospital in Southeast Brazil. *Clin Nutr.* 2008;27(2):228-32.
22. Charlton KE, Kolbe-Alexander TL, Nel JH. Micronutrient dilution associated with added sugar intake in elderly black South African women. *Eur J Clin Nutr.* 2005;59(9):1030-42.
23. Carter B, Monsivais P, Drewnowski A. Absorption of folic acid and ascorbic acid from nutrient comparable beverages. *J Food Sci.* 2010;75(9):H289-93.
24. Ravindran RD, Vashist P, Gupta SK, Young IS, Maraini G, Camparini M, Jayanthi R, John N, Fitzpatrick KE, Chakravarthy U, Ravilla TD, Fletcher AE. Prevalence and risk factors for vitamin C deficiency in north and south India: a two centre population based study in people aged 60 years and over. *PLoS One.* 2011;6(12):e28588.
25. Joint FAO-WHO Expert Group. Requirements of ascorbic acid, vitamin D, vitamin B12, folate, and iron. . Switzerland: World Health Organization; 1970. Report No.: 0512-3054 (Print)  
0512-3054 Contract No.: 452.
26. Hughes K, Ong CN. Vitamins, selenium, iron, and coronary heart disease risk in Indians, Malays, and Chinese in Singapore. *J Epidemiol Community Health.* 1998;52(3):181-5.
27. Gittelsohn J, Thapa M, Landman LT. Cultural factors, caloric intake and micronutrient sufficiency in rural Nepali households. *Soc Sci Med.* 1997;44(11):1739-49.
28. McCall SJ, Clark AB, Luben RN, Wareham NJ, Khaw KT, Myint PK. Plasma vitamin C levels: risk factors for deficiency and association with self-reported functional health in the European Prospective Investigation into Cancer-Norfolk. *Nutrients.* 2019;11(7).
29. Jungert A, Neuhauser-Berthold M. The lower vitamin C plasma concentrations in elderly men compared with elderly women can partly be attributed to a volumetric dilution effect due to differences in fat-free mass. *Br J Nutr.* 2015;113(5):859-64.
30. Ness AR, Cappuccio FP, Atkinson RW, Khaw KT, Cook DG. Plasma vitamin C levels in men and women from different ethnic backgrounds living in England. *Int J Epidemiol.* 1999;28(3):450-5.
31. GBD 2017 Diet Collaborators. Health effects of dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2019;393(10184):1958-72.
32. Desenclos JC, Berry AM, Padt R, Farah B, Segala C, Nabil AM. Epidemiological patterns of scurvy among Ethiopian refugees. *Bull World Health Organ.* 1989;67(3):309-16.
33. Kiondo P, Tumwesigye NM, Wandabwa J, Wamuyu-Maina G, Bimenya GS, Okong P. Plasma vitamin C assay in women of reproductive age in Kampala, Uganda, using a colorimetric method. *Trop Med Int Health.* 2012;17(2):191-6.
34. Halestrap P, Scheenstra S. Outbreak of scurvy in Tana River County, Kenya: A case report. *African journal of primary health care & family medicine.* 2018;10(1):e1-e3.
35. Cheung E, Mutahar R, Assefa F, Ververs MT, Nasiri SM, Borrel A, Salama P. An epidemic of scurvy in Afghanistan: assessment and response. *Food Nutr Bull.* 2003;24(3):247-55.
36. Khalife R, Grieco A, Khamisa K, Tinmouh A, McCudden C, Saidenberg E. Scurvy, an old story in a new time: The hematologist's experience. *Blood Cells Mol Dis.* 2019.
37. Montagnac JA, Davis CR, Tanumihardjo SA. Nutritional value of cassava for use as a staple food and recent advances for improvement. *Comprehensive Reviews in Food Science and Food Safety.* 2009;8:181-94.
38. Carr AC, Vissers MC. Synthetic or food-derived vitamin C--are they equally bioavailable? *Nutrients.* 2013;5(11):4284-304.

39. Olmedilla B, Granado F, Southon S, Wright AJ, Blanco I, Gil-Martinez E, Berg H, Corridan B, Roussel AM, Chopra M, Thurnham DI. Serum concentrations of carotenoids and vitamins A, E, and C in control subjects from five European countries. *Br J Nutr.* 2001;85(2):227-38.
40. Faure H, Preziosi P, Roussel AM, Bertrais S, Galan P, Hercberg S, Favier A. Factors influencing blood concentration of retinol, alpha-tocopherol, vitamin C, and beta-carotene in the French participants of the SU.VI.MAX trial. *Eur J Clin Nutr.* 2006;60(6):706-17.
41. Villalpando S, Montalvo-Velarde I, Zambrano N, Garcia-Guerra A, Ramirez-Silva CI, Shamah-Levy T, Rivera JA. Vitamins A, and C and folate status in Mexican children under 12 years and women 12-49 years: a probabilistic national survey. *Salud Publica Mex.* 2003;45 Suppl 4:S508-19.
42. Matilainen T, Vartiainen E, Puska P, Alfthan G, Pokusajeva S, Moisejeva N, Uhanov M. Plasma ascorbic acid concentrations in the Republic of Karelia, Russia and in North Karelia, Finland. *Eur J Clin Nutr.* 1996;50(2):115-20.
43. Bates CJ, Prentice A, Cole TJ, van der Pols JC, Doyle W, Finch S, Smithers G, Clarke PC. Micronutrients: highlights and research challenges from the 1994-5 National Diet and Nutrition Survey of people aged 65 years and over. *Br J Nutr.* 1999;82(1):7-15.
44. Frankenfeld CL, Lampe JW, Shannon J, Gao DL, Li W, Ray RM, Chen C, King IB, Thomas DB. Fruit and vegetable intakes in relation to plasma nutrient concentrations in women in Shanghai, China. *Public Health Nutr.* 2012;15(1):167-75.
45. Lam TK, Freedman ND, Fan JH, Qiao YL, Dawsey SM, Taylor PR, Abnet CC. Prediagnostic plasma vitamin C and risk of gastric adenocarcinoma and esophageal squamous cell carcinoma in a Chinese population. *Am J Clin Nutr.* 2013;98(5):1289-97.
46. Tribble DL, Giuliano LJ, Fortmann SP. Reduced plasma ascorbic acid concentrations in nonsmokers regularly exposed to environmental tobacco smoke. *Am J Clin Nutr.* 1993;58(6):886-90.
47. Chiplonkar SA, Agte VV, Mengale SS, Tarwadi KV. Are lifestyle factors good predictors of retinol and vitamin C deficiency in apparently healthy adults? *Eur J Clin Nutr.* 2002;56(2):96-104.
48. Strauss RS. Environmental tobacco smoke and serum vitamin C levels in children. *Pediatrics.* 2001;107(3):540-2.
49. Wilson KM, Finkelstein JN, Blumkin AK, Best D, Klein JD. Micronutrient levels in children exposed to secondhand tobacco smoke. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco.* 2011;13(9):800-8.
50. Preston AM, Rodriguez C, Rivera CE. Plasma ascorbate in a population of children: influence of age, gender, vitamin C intake, BMI and smoke exposure. *P R Health Sci J.* 2006;25(2):137-42.
51. Preston AM, Rodriguez C, Rivera CE, Sahai H. Influence of environmental tobacco smoke on vitamin C status in children. *Am J Clin Nutr.* 2003;77(1):167-72.
52. Bates CJ, Rutishauser IH, Black AE, Paul AA, Mandal AR, Patnaik BK. Long-term vitamin status and dietary intake of healthy elderly subjects. 2. Vitamin C. *Br J Nutr.* 1979;42(1):43-56.
53. United Nations. 68% of the world population projected to live in urban areas by 2050, says UN 2018 [Available from: <https://www.un.org/development/desa/en/news/population/2018-revision-of-world-urbanization-prospects.html>].
54. World Health Organization. Air pollution [Available from: [https://www.who.int/health-topics/air-pollution#tab=tab\\_1](https://www.who.int/health-topics/air-pollution#tab=tab_1)].

55. Dietrich M, Block G, Benowitz NL, Morrow JD, Hudes M, Jacob P, 3rd, Norkus EP, Packer L. Vitamin C supplementation decreases oxidative stress biomarker f2-isoprostanes in plasma of nonsmokers exposed to environmental tobacco smoke. *Nutr Cancer*. 2003;45(2):176-84.
56. Farchi S, Forastiere F, Pistelli R, Baldacci S, Simoni M, Perucci CA, Viegi G. Exposure to environmental tobacco smoke is associated with lower plasma beta-carotene levels among nonsmoking women married to a smoker. *Cancer Epidemiol Biomarkers Prev*. 2001;10(8):907-9.
57. Forastiere F, Mallone S, Lo Presti E, Baldacci S, Pistelli F, Simoni M, Scalera A, Pedreschi M, Pistelli R, Corbo G, Rapiti E, Agabiti N, Farchi S, Basso S, Chiaffi L, Matteelli G, Di Pede F, Carrozzi L, Viegi G. Characteristics of nonsmoking women exposed to spouses who smoke: epidemiologic study on environment and health in women from four Italian areas. *Environ Health Perspect*. 2000;108(12):1171-7.
58. Ayaori M, Hisada T, Suzukawa M, Yoshida H, Nishiwaki M, Ito T, Nakajima K, Higashi K, Yonemura A, Ohsuzu F, Ishikawa T, Nakamura H. Plasma levels and redox status of ascorbic acid and levels of lipid peroxidation products in active and passive smokers. *Environ Health Perspect*. 2000;108(2):105-8.
59. Howard DJ, Ota RB, Briggs LA, Hampton M, Pritsos CA. Oxidative stress induced by environmental tobacco smoke in the workplace is mitigated by antioxidant supplementation. *Cancer Epidemiol Biomarkers Prev*. 1998;7(11):981-8.
60. Galan P, Viteri FE, Bertrais S, Czernichow S, Faure H, Arnaud J, Ruffieux D, Chenal S, Arnault N, Favier A, Roussel AM, Hercberg S. Serum concentrations of beta-carotene, vitamins C and E, zinc and selenium are influenced by sex, age, diet, smoking status, alcohol consumption and corpulence in a general French adult population. *Eur J Clin Nutr*. 2005;59(10):1181-90.
61. Pearson JF, Pullar JM, Wilson R, Spittlehouse JK, Vissers MCM, Skidmore PML, Willis J, Cameron VA, Carr AC. Vitamin C status correlates with markers of metabolic and cognitive health in 50-year-olds: Findings of the CHALICE cohort study. *Nutrients*. 2017;9(8):E831.
62. Itoh R, Yamada K, Oka J, Echizen H, Murakami K. Sex as a factor in levels of serum ascorbic acid in a healthy elderly population. *Int J Vitam Nutr Res*. 1989;59(4):365-72.
63. Hughes K, New AL, Lee BL, Ong CN. Plasma vitamins A, C and E in the general population of Singapore, 1993 to 1995. *Ann Acad Med Singapore*. 1998;27(2):149-53.
64. Dherani M, Murthy GV, Gupta SK, Young IS, Maraini G, Camparini M, Price GM, John N, Chakravarthy U, Fletcher AE. Blood levels of vitamin C, carotenoids and retinol are inversely associated with cataract in a North Indian population. *Invest Ophthalmol Vis Sci*. 2008;49(8):3328-35.
65. Birlouez-Aragon I, Delcourt C, Tessier F, Papoz L. Associations of age, smoking habits and diabetes with plasma vitamin C of elderly of the POLA study. *Int J Vitam Nutr Res*. 2001;71(1):53-9.
66. Nyyssonen K, Parviainen MT, Salonen R, Tuomilehto J, Salonen JT. Vitamin C deficiency and risk of myocardial infarction: prospective population study of men from eastern Finland. *BMJ*. 1997;314(7081):634-8.
67. Koh ET, Chi MS, Lowenstein FW. Comparison of selected blood components by race, sex, and age. *Am J Clin Nutr*. 1980;33(8):1828-35.
68. Khan RM, Iqbal MP. Deficiency of Vitamin C in South Asia. *Pakistan journal of medical sciences*. 2006;22(3):347-55.
69. Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev*. 2015;73(10):643-60.
70. Timpson NJ, Forouhi NG, Brion MJ, Harbord RM, Cook DG, Johnson P, McConnachie A, Morris RW, Rodriguez S, Luan J, Ebrahim S, Padmanabhan S, Watt G, Bruckdorfer KR, Wareham NJ, Whincup PH,

- Chanock S, Sattar N, Lawlor DA, Davey Smith G. Genetic variation at the SLC23A1 locus is associated with circulating concentrations of L-ascorbic acid (vitamin C): evidence from 5 independent studies with >15,000 participants. *Am J Clin Nutr.* 2010;92(2):375-82.
71. Viroonudomphol D, Mahaisiriyodom A, Mingkhawn R, Sodomthian P, Korchasri N, Jittngamkhum S, Sangcakul A. Relationship between serum antioxidant vitamins A, E, and C and lipid profiles in priest subjects at the Priest Hospital. *Southeast Asian J Trop Med Public Health.* 2005;36 Suppl 4:246-53.
  72. Gould C, Tousignant B, Brian G, McKay R, Gibson R, Bailey K, Venn BJ. Cross-sectional dietary deficiencies among a prison population in Papua New Guinea. *BMC Int Health Hum Rights.* 2013;13:21.
  73. Ene-Obong HN, Odoh IF, Ikwuagwu OE. Plasma vitamin A and C status of in-school adolescents and associated factors in Enugu State, Nigeria. *J Health Popul Nutr.* 2003;21(1):18-25.
  74. Canoy D, Wareham N, Welch A, Bingham S, Luben R, Day N, Khaw KT. Plasma ascorbic acid concentrations and fat distribution in 19,068 British men and women in the European Prospective Investigation into Cancer and Nutrition Norfolk cohort study. *Am J Clin Nutr.* 2005;82(6):1203-9.
  75. Garcia OP, Ronquillo D, Caamano Mdel C, Camacho M, Long KZ, Rosado JL. Zinc, vitamin A, and vitamin C status are associated with leptin concentrations and obesity in Mexican women: results from a cross-sectional study. *Nutr Metab (Lond).* 2012;9(1):59.
  76. Block G, Mangels AR, Patterson BH, Levander OA, Norkus EP, Taylor PR. Body weight and prior depletion affect plasma ascorbate levels attained on identical vitamin C intake: a controlled-diet study. *J Am Coll Nutr.* 1999;18(6):628-37.
  77. Carr AC, Pullar JM, Bozonet SM, Vissers MC. Marginal ascorbate status (hypovitaminosis C) results in an attenuated response to vitamin C supplementation. *Nutrients.* 2016;8(6):E341.
  78. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, Mullany EC, Biryukov S, Abbafati C, Abera SF, Abraham JP, Abu-Rmeileh NM, Achoki T, AlBuhairan FS, Alemu ZA, Alfonso R, Ali MK, Ali R, Guzman NA, Ammar W, Anwari P, Banerjee A, Barquera S, Basu S, Bennett DA, Bhutta Z, Blore J, Cabral N, Nonato IC, Chang JC, Chowdhury R, Courville KJ, Criqui MH, Cundiff DK, Dabhadkar KC, Dandona L, Davis A, Dayama A, Dharmaratne SD, Ding EL, Durrani AM, Esteghamati A, Farzadfar F, Fay DF, Feigin VL, Flaxman A, Forouzanfar MH, Goto A, Green MA, Gupta R, Hafezi-Nejad N, Hankey GJ, Harewood HC, Havmoeller R, Hay S, Hernandez L, Hussein A, Idrisov BT, Ikeda N, Islami F, Jahangir E, Jassal SK, Jee SH, Jeffreys M, Jonas JB, Kabagambe EK, Khalifa SE, Kengne AP, Khader YS, Khang YH, Kim D, Kimokoti RW, Kinge JM, Kokubo Y, Kosen S, Kwan G, Lai T, Leinsalu M, Li Y, Liang X, Liu S, Logroscino G, Lotufo PA, Lu Y, Ma J, Mainoo NK, Mensah GA, Merriman TR, Mokdad AH, Moschandreas J, Naghavi M, Naheed A, Nand D, Narayan KM, Nelson EL, Neuhauser ML, Nisar MI, Ohkubo T, Oti SO, Pedroza A, Prabhakaran D, Roy N, Sampson U, Seo H, Sepanlou SG, Shibuya K, Shiri R, Shiue I, Singh GM, Singh JA, Skirbekk V, Stapelberg NJ, Sturua L, Sykes BL, Tobias M, Tran BX, Trasande L, Toyoshima H, van de Vijver S, Vasankari TJ, Veerman JL, Velasquez-Melendez G, Vlassov VV, Vollset SE, Vos T, Wang C, Wang X, Weiderpass E, Werdecker A, Wright JL, Yang YC, Yatsuya H, Yoon J, Yoon SJ, Zhao Y, Zhou M, Zhu S, Lopez AD, Murray CJ, Gakidou E. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2014;384(9945):766-81.
  79. Yatsuya H, Li Y, Hilawe EH, Ota A, Wang C, Chiang C, Zhang Y, Uemura M, Osako A, Ozaki Y, Aoyama A. Global trend in overweight and obesity and its association with cardiovascular disease incidence. *Circ J.* 2014;78(12):2807-18.

80. Saito K, Yokoyama T, Yoshida H, Kim H, Shimada H, Yoshida Y, Iwasa H, Shimizu Y, Yoshitaka K, Handa S, Maruyama N, Ishigami A, Suzuki T. A significant relationship between plasma vitamin C concentration and physical performance among Japanese elderly women. *J Gerontol A Biol Sci Med Sci*. 2011.
81. Cahill L, Corey PN, El-Sohemy A. Vitamin C deficiency in a population of young Canadian adults. *Am J Epidemiol*. 2009;170(4):464-71.
82. Cahill LE, El-Sohemy A. Vitamin C transporter gene polymorphisms, dietary vitamin C and serum ascorbic acid. *J Nutrigenet Nutrigenomics*. 2009;2(6):292-301.
83. Langlois MR, Delanghe JR, De Buyzere ML, Bernard DR, Ouyang J. Effect of haptoglobin on the metabolism of vitamin C. *Am J Clin Nutr*. 1997;66(3):606-10.
84. Na N, Delanghe JR, Taes YE, Torck M, Baeyens WR, Ouyang J. Serum vitamin C concentration is influenced by haptoglobin polymorphism and iron status in Chinese. *Clin Chim Acta*. 2006;365(1-2):319-24.
85. Cahill LE, El-Sohemy A. Haptoglobin genotype modifies the association between dietary vitamin C and serum ascorbic acid deficiency. *Am J Clin Nutr*. 2010;92(6):1494-500.
86. Marangon K, Herbeth B, Lecomte E, Paul-Dauphin A, Grolier P, Chancerelle Y, Artur Y, Siest G. Diet, antioxidant status, and smoking habits in French men. *Am J Clin Nutr*. 1998;67(2):231-9.
87. Schectman G, Byrd JC, Gruchow HW. The influence of smoking on vitamin C status in adults. *Am J Public Health*. 1989;79(2):158-62.
88. Schectman G, Byrd JC, Hoffmann R. Ascorbic acid requirements for smokers: analysis of a population survey. *Am J Clin Nutr*. 1991;53(6):1466-70.
89. Boonpangrak S, Tantimongcolwat T, Treeratanapiboon L, Leelahakul P, Prachayasittikul V. Lifestyle behaviors and serum vitamin C in the Thai population in Bangkok Metropolitan. *EXCLI journal*. 2018;17:452-66.
90. McGregor GP, Biesalski HK. Rationale and impact of vitamin C in clinical nutrition. *Curr Opin Clin Nutr Metab Care*. 2006;9(6):697-703.
91. Carr AC, Cook J. Intravenous vitamin C for cancer therapy - identifying the current gaps in our knowledge. *Front Physiol*. 2018;9:1182.
92. Carr AC. Vitamin C in pneumonia and sepsis. In: Chen Q, Vissers M, editors. *Vitamin C: New Biochemical and Functional Insights. Oxidative Stress and Disease*. Boca Raton, FL CRC Press/Taylor & Francis; 2020. p. 115-35.
93. Juhl B, Lauszus FF, Lykkesfeldt J. Is Diabetes Associated with Lower Vitamin C Status in Pregnant Women? A Prospective Study. *Int J Vitam Nutr Res*. 2017;1-5.
94. Juhl B, Lauszus FF, Lykkesfeldt J. Poor Vitamin C Status Late in Pregnancy Is Associated with Increased Risk of Complications in Type 1 Diabetic Women: A Cross-Sectional Study. *Nutrients*. 2017;9(3).
95. Rumbold A, Ota E, Nagata C, Shahrook S, Crowther CA. Vitamin C supplementation in pregnancy. The Cochrane database of systematic reviews. 2015(9):Cd004072.
96. Lykkesfeldt J, Poulsen HE. Is vitamin C supplementation beneficial? Lessons learned from randomised controlled trials. *Br J Nutr*. 2010;103(9):1251-9.
97. Ugwa E. Vitamin C supplementation in pregnancy: A review of current literature. *Nigerian Journal of Basic and Clinical Sciences*. 2015;12(1):1-5.
98. Camarena V, Wang G. The epigenetic role of vitamin C in health and disease. *Cell Mol Life Sci*. 2016;73(8):1645-58.
99. Michels AJ, Hagen TM, Frei B. Human genetic variation influences vitamin C homeostasis by altering vitamin C transport and antioxidant enzyme function. *Annu Rev Nutr*. 2013;33:45-70.

100. Delanghe JR, Langlois MR, De Buyzere ML, Torck MA. Vitamin C deficiency and scurvy are not only a dietary problem but are codetermined by the haptoglobin polymorphism. *Clin Chem*. 2007;53(8):1397-400.
101. Langlois MR, Delanghe JR. Biological and clinical significance of haptoglobin polymorphism in humans. *Clin Chem*. 1996;42(10):1589-600.
102. Cahill LE, Fontaine-Bisson B, El-Sohemy A. Functional genetic variants of glutathione S-transferase protect against serum ascorbic acid deficiency. *Am J Clin Nutr*. 2009;90(5):1411-7.
103. Ames BN, Elson-Schwab I, Silver EA. High-dose vitamin therapy stimulates variant enzymes with decreased coenzyme binding affinity (increased K(m)). *Am J Clin Nutr*. 2002;75(4):616-58.
104. Levine M, Eck P. Vitamin C: working on the x-axis. *Am J Clin Nutr*. 2009;90(5):1121-3.
105. Lykkesfeldt J, Viscovich M, Poulsen HE. Plasma malondialdehyde is induced by smoking: a study with balanced antioxidant profiles. *Br J Nutr*. 2004;92(2):203-6.
106. Zondervan KT, Ocke MC, Smit HA, Seidell JC. Do dietary and supplementary intakes of antioxidants differ with smoking status? *Int J Epidemiol*. 1996;25(1):70-9.
107. Larkin FA, Basiotis PP, Riddick HA, Sykes KE, Pao EM. Dietary patterns of women smokers and non-smokers. *J Am Diet Assoc*. 1990;90(2):230-7.
108. Dietrich M, Block G, Norkus EP, Hudes M, Traber MG, Cross CE, Packer L. Smoking and exposure to environmental tobacco smoke decrease some plasma antioxidants and increase gamma-tocopherol in vivo after adjustment for dietary antioxidant intakes. *Am J Clin Nutr*. 2003;77(1):160-6.
109. Lykkesfeldt J, Christen S, Wallock LM, Chang HH, Jacob RA, Ames BN. Ascorbate is depleted by smoking and repleted by moderate supplementation: a study in male smokers and nonsmokers with matched dietary antioxidant intakes. *Am J Clin Nutr*. 2000;71(2):530-6.
110. Kallner AB, Hartmann D, Hornig DH. On the requirements of ascorbic acid in man: steady-state turnover and body pool in smokers. *Am J Clin Nutr*. 1981;34(7):1347-55.
111. GBD 2015 Tobacco Collaborators. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990-2015: a systematic analysis from the Global Burden of Disease Study 2015. *Lancet*. 2017;389(10082):1885-906.
112. Lykkesfeldt J, Prieme H, Loft S, Poulsen HE. Effect of smoking cessation on plasma ascorbic acid concentration. *BMJ*. 1996;313(7049):91.
113. Sharma Y, Miller M, Shahi R, Doyle A, Horwood C, Hakendorf P, Thompson C. Vitamin C deficiency in Australian hospitalized patients: an observational study. *Intern Med J*. 2018.
114. Gan R, Eintracht S, Hoffer LJ. Vitamin C deficiency in a university teaching hospital. *J Am Coll Nutr*. 2008;27(3):428-33.
115. Gariballa S, Forster S. Effects of acute-phase response on nutritional status and clinical outcome of hospitalized patients. *Nutrition*. 2006;22(7-8):750-7.
116. Fain O, Paries J, Jacquart B, Le Moel G, Kettaneh A, Stirnemann J, Heron C, Sitbon M, Taleb C, Letellier E, Betari B, Gattegno L, Thomas M. Hypovitaminosis C in hospitalized patients. *Eur J Intern Med*. 2003;14(7):419-25.
117. Hemilä H. Vitamin C and infections. *Nutrients*. 2017;9(4):E339.
118. Carr AC, Spencer E, Dixon L, Chambers ST. Patients with community acquired pneumonia exhibit depleted vitamin C status and elevated oxidative stress. *Nutrients*. 2020;12, ; (1318):10 pages.
119. Carr AC, Rosengrave PC, Bayer S, Chambers S, Mehrtens J, Shaw GM. Hypovitaminosis C and vitamin C deficiency in critically ill patients despite recommended enteral and parenteral intakes. *Crit Care*. 2017;21:300.

120. Johnston CS, Corte C. People with marginal vitamin C status are at high risk of developing vitamin C deficiency. *J Am Diet Assoc.* 1999;99(7):854-6.
121. Hampl JS, Taylor CA, Johnston CS. Vitamin C deficiency and depletion in the United States: the Third National Health and Nutrition Examination Survey, 1988 to 1994. *Am J Public Health.* 2004;94(5):870-5.
122. Institute of Medicine Panel on Dietary Antioxidants and Related Compounds. Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids. Washington: National Academies Press; 2000. 529 p.
123. Loria CM, Whelton PK, Caulfield LE, Szklo M, Klag MJ. Agreement among indicators of vitamin C status. *Am J Epidemiol.* 1998;147(6):587-96.
124. Washko PW, Welch RW, Dhariwal KR, Wang Y, Levine M. Ascorbic acid and dehydroascorbic acid analyses in biological samples. *Anal Biochem.* 1992;204(1):1-14.
125. Amaliya, Timmerman MF, Abbas F, Loos BG, Van der Weijden GA, Van Winkelhoff AJ, Winkel EG, Van der Velden U. Java project on periodontal diseases: the relationship between vitamin C and the severity of periodontitis. *J Clin Periodontol.* 2007;34(4):299-304.
126. Amaliya A, Laine ML, Delanghe JR, Loos BG, Van Wijk AJ, Van der Velden U. Java project on periodontal diseases: periodontal bone loss in relation to environmental and systemic conditions. *J Clin Periodontol.* 2015;42(4):325-32.
127. Pullar JM, Bayer S, Carr AC. Appropriate handling, processing and analysis of blood samples is essential to avoid oxidation of vitamin C to dehydroascorbic acid. *Antioxidants.* 2018;7(2):E29.