

Consciousness as an emergent phenomenon: a tale of different levels of description

Ramón Guevara^{1,2}, Diego Mateos³, and José Luis Pérez Velázquez⁴

¹Integrative Neuroscience and Cognition Centre (INCC UMR8002), University of Paris and CNRS, Paris, France

²Department of Physics and Astronomy, University of Padova, Padova, Italy

³Universidad Autónoma de Entre Ríos, Parana, Argentina.

⁴Instituto de Matemática Aplicada del Litoral (IMAL-CONICET-UNL), Santa Fé, Argentina

⁵The Ronin Institute, Montclair, NJ, United States

Abstract

One of the biggest queries in cognitive sciences is the emergence of consciousness from matter. Modern neurobiological theories of consciousness propose that conscious experience is the result of interactions between large-scale neuronal networks in the brain, traditionally described within the realm of classical physics. Here, we propose a generalized connectionist framework in which the emergence of “conscious networks” is not exclusive of large brain areas, but can be identified in sub-cellular networks exhibiting non-trivial quantum phenomena. The essential feature of such networks is the existence of strong correlations in the system (classical or quantum coherence) and the presence of an optimal point at which the system’s complexity is maximized, expressed either by maximization of the information content in large scale functional networks or by achieving optimal efficiency through the quantum Goldilock effect.

Introduction

The relation between behavior and brain function is the main task of cognitive neuroscience. In particular, one of the biggest queries in cognitive sciences and philosophy is the emergence of consciousness from matter (Crick and Clark (1994), Edelman and Tononi (2003)). A natural question to ask is at what level is consciousness generated, that is, what is the neural substrate of conscious experience and where it is located. Different possibilities have been explored, ranging from the cellular level to the level of the whole or at least large parts of the brain (Koch *et al.* (2016)).

One reason that makes this question hard to answer is that, as with other primordial questions in science, there is not a precise definition of consciousness, one that is accepted by all the scientific community (Tassi and Muzet (2001)) (but see some proposal about the advantages of a less strict definition of consciousness in Velazquez (2020)). Similar conundrums can be found all across biological sciences. Even the concept of life itself leads to several possible definitions and diverse proposals to define life had been advanced during the years, more or less emphasizing particular attributes of biological systems (Luisi (1998)). Furthermore, if one were asked where in an organism resides a property called “life”, in other words, what part of the organism is alive, one would probably consider the question misleading or simply wrong. One would eventually conclude that life is a global property that has no specific scale. And in fact, considering that basic aspects of life are exchange of energy, reproduction and compartmentalization, inorganic compounds which have these properties could be considered “alive” to some extent (Monnard and Walde (2015)), as similar principles of organization apply to the living and the nonliving (Velazquez (2009)). The organism is a complex system at all possible scales, from systems made up of organs down to the molecular scale, and each part contributes to the whole in an irreducible way. Failure at any scale can lead to the collapse of the organism; consider, for example, damage to the DNA structure by ionizing radiation at the molecular level, or heart failure at the organ level.

Where in the nervous system (or more generally, in the organism) consciousness resides may constitute a question of a similar nature. One possible answer could be that consciousness does not reside within any specific structure and that it cannot be clearly associated with a particular spatial scale. Another possibility is that consciousness appears at a specific scale, associated with a structure or group of structures. This is the question we address here. We start by acknowledging the overwhelming evidence supporting the idea that integration in large-scale neural networks may constitute the underlying mechanism of conscious phenomena, at least at the mesoscopic or macroscopic level of description.

But we also address the possibility that conscious phenomena can result from, or at least be enhanced by, non-trivial quantum effects. Does quantum physics play a role in the emergence of conscious phenomena? The answer to this question has been negative in the neuroscientific community at large (Tegmark (2000), Koch and Hepp (2006), Baars and Edelman (2012)). However, alternative explanations to prevalent paradigms do exist. In particular, several authors have proposed during the course of the last few decades, the possibility that consciousness could arise as a result of non-trivial quantum mechanical processes, such as quantum coherence and entanglement (for a magnificent recent review see Adams and Petruccione (2020)). Here we focus on such approaches to enquire about the possibility that these can be incorporated into a connectionist approach to the emergence of consciousness. We propose that consciousness is a multi-scale phenomena, the result of both large-scale interactions in neural networks and non-trivial quantum phenomena within cells.

Connectionists theories of consciousness

Several theories of consciousness are strongly based on a connectionist approach, that is, on the idea that large webs of interacting neurons are fundamental to understand brain functionality, cognition and behavior (O'Brien and Opie (1999), Maia and Cleeremans (2005)). These are "global" theories of consciousness, in that typically large brain areas are activated and necessary for the emergence of consciousness (McGovern and Baars (2007), Seth (2007)). These theories provide a point of view according to which it is possible to properly characterize conscious phenomenon in terms of certain properties common to all of them, and they have offered means to quantify them by identifying concomitant brain processes and possible underlying mechanisms. In the words of the authors of two influential neurobiological theories of consciousness, the dynamic core hypothesis (Tononi (1998), Edelman and Tononi (2003)) and the Integrated Information Theory (IIT) (Tononi (2004), Tononi (2008)), "instead of arguing whether a particular brain area or group of neurons contributes to consciousness or not, our strategy is to characterize the kinds of neural processes that might account for key properties of conscious experience" (Tononi (1998)).

For example, the Baars' global workspace theory (Baars (1993)) has used the traditional Cartesian argument of consciousness, an influential idea in cognitive sciences, in association with specific neural hypothesis based on electrophysiological evidence (Baars (1998)). In the Cartesian theater metaphor consciousness is an illuminated part of a stage representing immediate memory. This spotlight is shifted across the stage, directed by attention under executive action, whereas the rest of the theater is in darkness and unconscious. The bright spot correlates with the activation of a specific sensory projection cortical area (e.g., visual, auditory), whereas other cortical areas are simultaneously inhibited as they correspond to non-illuminated parts of the theater. Contents of conscious states are then sent to specialized networks all around the brain, corresponding to the audience sitting in darkness; this may happen in the brain through corticocortical or corticotalamalamic pathways. Specifically, the spotlight region content is chosen by attentional mechanisms, using bottom-up limbic and thalamic pathways or from executive top-down prefrontal cortex input (Baars (2002)). Therefore, the most important role of consciousness is that of integrating and coordinating a large number of specialized networks that would otherwise operate autonomously (Mountcastle (1978)).

Similarly, in the implementation of the global workspace theory due to Dehaene, Changeux and colleagues, a pattern of global activity is generated in the workspace when sensory stimuli activate excitatory neurons with long-range cortico-cortical axons (Dehaene *et al.* (2003)). Such patterns inhibit other possible patterns of activity that therefore remain unconscious, preventing the processing of alternative stimuli.

According to some of the "global" theories of consciousness, it is possible to identify two general properties of conscious experiences: integration and differentiation (Tononi (1998), Tononi (2004)).

Integration reflects the unity of conscious experiences, in the sense that each such experience is unique, so it cannot be decomposed into independent components. Differentiation refers to the enormous amount of potential conscious experiences that are simultaneous, that is, the extremely large repertoire of possible conscious experiences, of which one is selected.

In correspondence with these general properties of conscious phenomena, IIT, the dynamic core hypothesis and other related theories of consciousness identify neural processes that underlay conscious experience (Tononi (2004)). The first such process is the activation and deactivation of widely distributed brain areas, particularly in the thalamocortical system (segregation). The second is the integration through strong and rapid reentrant interactions between differentiated neuronal groups. In simple words, differentiation is achieved by the simultaneous activation of distinct and widely distributed neuronal groups, and integration refers to the process of connection of these groups through reentrant interactions to achieve a unique conscious experience.

A crucial element in connectionists approaches is that consciousness appears as an emergent property of large neuronal networks, requiring the interaction of neurons to form a web sufficiently complex such as to sustain conscious experiences. In such view, neurons are crucial - that is, necessary- but not sufficient to generate consciousness. This is not an exceptional perspective in cognitive neuroscience, but rather representative of contemporary ideas. In the dominant computing argot, neurons constitute the computing units within large networks, and computation is performed at the synaptic level. We can abstract out the crucial point: the interaction (integration) of a large ensemble of processing units (segregation) leads to emergent phenomena, among them consciousness. Thus, conscious experience reflects such underlying substrate expressed in the neurophysiological properties of integration and segregation.

But what is the mechanism of brain integration underlying conscious phenomena? Temporal correlations are known to play a crucial role in the functioning of the brain, cognition and conscious experience. Indeed, it has for long been recognized that temporal correlations of neural activity is crucial for communication between brain areas and between neurons (Singer (1999), Engel *et al.* (2001), Varela *et al.* (2001), Fries (2005), Velazquez and Frantseva (2011)).

Temporal correlation in the brain is often associated to the concept of neural synchronization. It is well known that nonlinear coupled autonomous oscillators (self-sustained oscillators) show synchronization phenomena (Pikovsky *et al.* (2003)). Synchronized oscillators show temporal correlations, specifically correlations in the phase of the corresponding signals. Neurons are not an exception, and synchronized neuronal activity (phase synchronization) has been found at multiple scales within the nervous system (Buzsáki (2009)), contributing to the so-called "brainweb" (Varela *et al.* (2001)). Not only individual neurons synchronize their activities in local regions, but also large neuronal networks (Perez Velazquez *et al.* (2015)).

It is currently believed that neural synchronization is the mechanism at the basis of integration in the brain (Varela *et al.* (2001)). Neural correlations are therefore of crucial importance in connectionist theories of brain activity and cognition, and in particular in connectionist theories of consciousness, such as IIT, because they underlay the property of integration in conscious experience.

Synchronization is also crucial for the development of brain rhythms, that appear to have a huge relevance in cognitive processes, including conscious experience (Buzsáki (2009)). Indeed, synchronization is required for neurons to produce an activity that is quasi-periodic across large brain areas, such as the rhythms observed in electroencephalographic (EEG) recordings (Nunez and Srinivasan (1981)). They appear as a result of feedback activity across large neuronal loops, and they have been associated to a myriad of cognitive functions (Buzsáki and Draguhn (2004)). Of crucial importance in the current context, the gamma band activity (around 40 Hz) has been associated with high cognitive functions, including consciousness; however, the generation of gamma activity may be a consequence of the natural activity of neurons during conscious experience: gamma frequencies reflect the emergence and dissolution of communication among many neuronal ensembles, fluctuations providing the variability needed in brain dynamics to process information leading to adaptive behaviours (Velazquez *et al.* (2019)).

Several studies have emphasized the importance of the thalamocortical system and gamma band synchronization in the generation of conscious experience (Llinás *et al.* (1998)). In particular, it seems plausible that synchronous oscillations in thalamocortical loops creates conscious states.

It is also important to notice that the breakdown of neural synchronization may lead to pathological brain dynamics. This is exemplified in diseases such as epilepsy and Parkinson disease, characterized by abnormal synchronization of brain activity (Jiruska *et al.* (2013)). In the case

of epilepsy, over-synchronization in extended brain areas leads to the large, periodic discharges observed in EEG recordings during seizures and the concomitant loss of responsiveness. Some authors suggest that during epileptic absent seizures, where consciousness is temporarily lost, there is a disruption of the thalamocortical "resonance" and gamma band synchronization which leads to disorders of consciousness (Llinás *et al.* (1998)).

A crucial point in connectionist theories of consciousness is the relationship between consciousness and complexity. According to the dynamic core hypothesis, complexity means that the dynamic core must be highly differentiated. The dynamic core also correlates with the conscious state of the subject (Tononi (1998)). Therefore it is expected that complexity is higher for "more conscious" states. For example, complexity of deep sleep stages should be lower than complexity during awake states.

Based on a statistical mechanics approach, some features of brain organization optimal for sensory processing and that correlate with consciousness were recently identified. Neurophysiological recordings taken during conscious and unconscious states (epilepsy, sleep and coma) were analyzed. Normal wakeful states were characterized by greatest number of possible configurations of interactions between brain networks, representing highest Shannon entropy values, suggesting that the information content is larger in networks associated with conscious awareness and that consciousness could be the result of an optimization of information processing (Guevara Erra *et al.* (2016), Mateos *et al.* (2017), Mateos *et al.* (2018)). These observations also suggest a fine-tuning of parameters in order for brain dynamics to achieve maximal efficiency. Conscious states are typically near the maximum of the curve of complexity vs synchronization (Fig.1), implying that networks are more efficient in information processing and consciousness emerges at intermediate values of synchronization: too high or too low values of synchronization were associated to low information content in functional networks, or equivalently, low network complexity (or low network global connectivity). Indeed, low values of complexity are observed in deep sleep stages and during generalized epileptic seizures, where consciousness is lost.

Another way to look at this fine-tuning is to describe it in terms of a trade off between noise and phase correlation in the system. Too much noise typically indicates a very low level of correlation, a system that is too random. On the other end of the spectrum, if the system has too little noise and too much correlation (for example, during a seizure) it is excessively ordered and cannot convey information. So, it seems that consciousness appear within a region of brain dynamics where noise and correlation values are intermediate (no too low nor too high). This is in agreement with ideas going back for decades about the emergence of complexity in biological systems, such as the "edge of chaos" concept of Stuart Kauffman (Kauffman (1993)). According to these ideas, complexity arises in the border between order (as in a crystalline lattice) and disorder (as in a gas).

Summarizing, we identify three properties that are common for the emergence of consciousness in a global connectionist approach:

1. The existence of functional networks that are activated and deactivated in accordance with concomitant conscious phenomena. Such networks are global because they expand across large areas of the brain.
2. The presence of temporal correlations (neural synchronization) within networks relevant to consciousness.
3. A fine-tuning of network parameters is associated with the emergence of consciousness. Consciousness appear at intermediate levels of noise and network correlation. The complexity of the networks is close to maximal for conscious states and decreases on altered states of consciousness.

Towards a multiple-scale theory of consciousness

When characterizing networks underlying cognitive phenomena, the identification of the constituting units of the network is fundamental. According to the influential ideas put forward by global connectionist theories of consciousness, such entities are restricted to either individual neurons or neuronal networks. However, extremely complex networks of interacting units do exist within cells and in particular within neurons. Could it be that a connectionist approach to consciousness can be also established at the infra-neuronal level, with sub cellular mechanisms contributing to the

emergence of consciousness? Interestingly, brain activity at the molecular level is very rich, and networks of all sorts can be identified within cells (consider, for example, metabolic networks).

More generally, a purely connectionist approach leads naturally to the following question: why consciousness does not emerge from any network (not necessarily neuronal) fulfilling the necessary ingredients (integration and differentiation) for its emergence? Complex networks are widespread in biological systems, and they can be identified at multiples spatio-temporal scales. Consider an extreme case: is a sufficiently complex ecological network conscious? This is a very difficult question, that is beyond the scope of our text. Instead, we focus on networks of interactions within the nervous system. We have to consider that the nervous system has evolved to process information and to organize the responses of organisms to environmental changes in a timely fashion, contrary to other networks that are not specific in this sense. Since conscious experiences, at least at the psychological level, are intimately related to perception and cognition, it makes sense to analyze networks within the brain or within neurons.

An important property of the nervous system is that its architecture is extremely complex and there is an intricate connection between structures at different scales. For example, the detection of a few photons at a single retinal neuron may produce a cascade of neural processes leading to the perception of light. In humans and other mammals, it is argued that the first steps of this cascade are actually unconscious, and only when large brain areas are activated this percept effectively enters into the realm of conscious awareness. In much simpler animals with a very small nervous system, the cascade is finished much sooner. Some could argue that such animals are not conscious. However, several authors have suggested that consciousness may be a general property of vertebrates, including very early forms (Feinberg and Mallatt (2018)).

Of course these perspectives depend on the definition of consciousness one adopts, but we have to admit there is no single definition that satisfies all, and we will not delve into this point here, rather only will mention that, as commented above, there are advantages if less strict definitions of consciousness are adopted (Velazquez (2020)). To wit, just like the notion of life, consciousness can be defined by enumerating characteristics: features like self-reproduction, compartmentalization, etc. can be used to “define” life and consciousness can be described by sensorimotor actions, self-awareness, memory etc. In this manner, “primitive” creatures and even microorganisms can be endowed with some sort of primordial consciousness, as these entities possess some features of consciousness like response to environmental changes, memory, and others. But few numbers of neurons and networks in worms do not allow them, presumably, to display self-awareness—even though we note that worms have a withdrawal reflex in that these animals respond to external cutaneous touch but not to self-induced touch so they seem to distinguish between inputs from the outside world and the self-induced sensory input, phenomenon that occurs because of the corollary discharge, the transmission of a copy of motor commands to sensory areas where the expected sensation is generated (Feinberg (1978)). In the final analysis these behaviours represent a continuum in cognition, the essentials being present in primordial nervous systems, and it should always be kept in mind that cognition and consciousness developed in a graded, continuous fashion, these are not static properties but a dynamic processes (Haken (2006)).

In any case, complex structure and dynamics has been identified at many levels within the nervous system. One important property of complex systems and processes is the apparent lack of a specific, dominant scale, a property that also pervades brain architecture and function (Turner *et al.* (2018)). Indeed, scale-free properties are common in neuronal networks. Just to mention an example, the electric signal detected in electroencephalographic recording exhibit scale free-dynamics, characterized by the so-called $1/f$ property of the power spectrum (Nunez and Srinivasan (2005)). It is then arguable that phenomena at the cellular level may have either a very important impact on conscious experience, an assertion that may be speculative and controversial and yet not ruled out by experimental evidence.

So, what criteria do we use in order to identify networks correlates of consciousness or self-awareness, apart from those already known - large neuronal networks in large brains - that constitute the central theme of modern neuroscience? Above, we proposed that we can follow a connectionist approach, and identified three properties that such networks should satisfy. We also propose that a fundamental property of such networks is the existence of synchronization phenomena, or resonance phenomena leading to important levels of temporal correlation in the network. We emphasize that synchronization is a widespread phenomenon in nonlinear dynamics, and therefore is commonplace among complex interconnected networks, with abundant examples in physics, chemistry and biology. Mathematically speaking, synchronization is the adjustment of rhythms

of weakly nonlinear oscillators (Pikovsky *et al.* (2003)). However, "neural synchronization", as understood in neuroscience, is an umbrella term compressing all sorts of phenomena in the nervous system where temporal correlations are present (Guevara Erra *et al.* (2017)); for example, there may be resonant phenomena in neurons but also coherency of the mean field of large neuronal networks. We propose that is neural synchronization - both in a strict mathematical sense and in the neuroscientific broad sense - that underlies conscious phenomena, because it leads to temporal coordination and correlations within neuronal networks, a crucial ingredient for conscious phenomena, that is, integration of information.

The question is then if temporal coordinated activity has been identified in networks within cells, and to verify that such networks correlate to conscious phenomena. Curiously, an interesting and unexpected candidate has recently appeared in approaches to the problem of consciousness, based on quantum physics rather than classical physics, and associated to sub-cellular or even molecular structures. We refer here to this approach as "quantum consciousness". Temporal coordination in that case comes in the form of quantum coherence and entanglement. We will dedicate the rest of the article to articulate this idea.

Quantum effects in biology

Is there any relevance of quantum physics for living organisms? One possible answer is that ultimately any kind of matter is described by quantum mechanics. Indeed, there is one obvious way in which quantum mechanics enters the realm of biological phenomena: at the biochemical level, molecular structure and interactions are affected by quantum phenomena, because quantum effects are important at the scale of atoms and molecules. This has important consequences, for example, in the enzymatic reactions occurring in biological membranes within cells. However, we do not refer here to this kind of quantum phenomena, which is rather trivial, in the sense that they can be equally described by approximations using classical physics. We refer, instead, to irreducible quantum phenomena that require in one way or another superposition of quantum states, typically at a larger scale, compared to the molecular scale, and involving large polymers and complex chemical reactions. Non-trivial quantum effects are related to quantum coherence. Coherence appears because in quantum mechanics particles have wave-like properties, in the sense that their probability distributions are given by a wave-like equation (Schrödinger equation) (Griffiths and Schroeter (2018)), and therefore, as is typical for other waves, are subject to interference phenomena. A very well-known example is the double-slit Young experiment. Individual particles pass through each of the slits, but the ensemble is described by the probability distribution of the particles, which has wave-like properties including interference. Under certain conditions, quantum coherence can appear at the macroscopic scale, as in Bose-Einstein condensates. The destruction of coherence by interaction with the environment is called decoherence. Although more difficult to prove in a biological context, some authors also refer to the possible physiological effect of a purely quantum correlation, called entanglement (Bes (2012)), impossible to describe in terms of classical physics (Brookes (2017)). Entanglement is a phenomenon that results when the wavefunction describing the state of a group of particles cannot be factored out as the product of the states of individual particles. Entanglement is a simple consequence of the superposition principle applied to multi-particle systems. It implies that the properties of certain configurations of particles are intrinsically intermingled (entangled) in such a way that results of experiments on different particles are correlated in a non-trivial way, impossible to explain in terms of classical physics. Finally, quantum tunneling, another quantum effect in which particles have non-zero probability of passing through a barrier potential, have also been associated with biological phenomena such as olfaction and neuroreception (Adams and Petruccione (2020)).

Of great importance for the argument we develop here, quantum coherence and entanglement lead to temporal correlation between particles, in the same manner as synchronization lead to correlation between nonlinear oscillators. This is the crucial analogy we emphasize in this article. The analogy is of course not complete. Quantum objects obey the Schrödinger equation, that is linear, whereas synchronization phenomena occur in nonlinear systems. So, it should be perhaps more correct to say that such correlation arrives from resonances at the microscopic level. Another similarity of quantum correlations with their classical counterparts is that both are connected to phase transitions and the emergence of new properties. As for the brain, phase transitions between synchronized and desynchronized states in networks are commonplace, and they may even lead to pathological states in epilepsy and Parkinson disease, or to altered states of consciousness. Such

phase transitions are important because they constitute the basis for "emergence phenomena", a class to which consciousness seems to belong. Apart from neuroscience and many other areas of biology, emergency has been also traditionally studied by condense matter physicist, in the context of large systems containing interacting units. Indeed, the title of the seminal paper by Anderson "More is different" (Anderson (1972)), succinctly summarize what emergency is: new, unexpected properties that appear in large systems of interacting particles. Or great importance in this context, some of these phenomena are quantum in nature, as is the case of superconductivity and superfluidity. Indeed, they constitute "macroscopic quantum phenomena", leading to emergent properties and phase transitions. For years the possibility of such macroscopic quantum phenomena in the brain were excluded, because they normally appear at very low temperatures, far from the physiologically relevant temperatures at which an organism function normally.

However, the last decade has witnessed the development of a the emerging field of quantum biology (Lambert *et al.* (2013), Fleming *et al.* (2011)), and several unexpected quantum effects have been discovered in soft, biological tissues at room temperature, at times having crucial physiological implications, such as the discovery of quantum coherence in the photosynthesis process (Engel *et al.* (2007), Panitchayangkoon *et al.* (2010), Ishizaki and Fleming (2012)). Other examples of relevancy of quantum effects in biology include enzyme catalysis, olfaction, the avian compass and DNA replication (McFadden and Al-Khalili (2016), Marais *et al.* (2018)). Such advances demonstrate that decoherence may fail to destroy quantum correlations in biological tissue at physiological temperature (in the "warm, wet and messy" biological tissue (Marais *et al.* (2018))), and it actually seems that at least in some cases molecular architecture and function has been designed in such a way as to avoid the deleterious effects of noise, in order to preserve quantum coherence and in a way as to benefit living organisms. As well, it has been speculated that evolution may have taken advantage of such processes to maintain ordered states in crucial biological functions that could be otherwise difficult to preserve.

To illustrate this point, consider photosynthesis, a paradigmatic system of study in the emerging field of quantum biology (Marais *et al.* (2018)), that sheds light on the importance of non-trivial quantum effects in living matter. In the photochemical reactions occurring during photosynthesis within plants and algae chloroplasts, light quanta is absorbed by chlorophyll, and by a chain of chemical reactions involving electron transport, solar energy is stored in the NADPH and ATP molecules, for subsequent use in metabolism (Alberts *et al.* (2002)). Chlorophyll and other similar molecules used in photosynthesis by diverse organisms are chromophores (literally a "carrier of color"), molecules that constitute pigments in that they absorb light in certain wavelenghts and are transparent in others (Rüdiger (1986)). In the case of chlorophyll, the molecule is transparent in the wavelength corresponding to green light. One end of the chlorophyll molecule is a flat, antenna-like structure with a magnesium atom at its center. When a quantum of light hits the magnesium atom, an electron is emitted, carrying with it the energy delivered by the photon. After this initial stage, the excited electron has to be transferred to a reaction center, where charge separation occurs and free energy can be stocked within more stable molecules, for subsequent use. These first stages in photosynthesis, including the transfer of energy to the reaction center, are very efficient. This is difficult to explain if it is assumed that excited electrons classically diffuse across chromophores to reach the reaction center. If this was the case, most energy would be lost and efficiency should be very low. Indeed, a search strategy based on classical Brownian motion is extremely inefficient and could not account to the almost 100% efficiency found in the first stages in photosynthesis (Lloyd *et al.* (2011), Fleming *et al.* (2011)).

Here is where quantum mechanics comes at rescue. Once an electron is excited by a quantum of light the lost negative charge can be considered as a positive "hole" in the magnesium atom, as is typically done in condense matter and field theory, a typical example being the electrons and holes used to describe semiconductors (Knox (1963)). The ensemble of the electron and the hole can be treated quantum-mechanically as an elementary excitation, or exciton (Novoderezhkin and van Grondelle (2010)). As it turns out, in photosynthetic light harvesting complexes these collective optical excitations lead to correlations of electron's motions in different chromophores (Mohseni *et al.* (2008), Fassioli *et al.* (2014)). In other words, there is a superposition of the electronic excitations of individual chlorophyll molecules, leading to much more efficient transfer of energy from chlorophyll molecules to the reaction center. This possibility was already pointed out by pioneer work of Franck and Teller (Franck and Teller (1938)) but it was not corroborated experimentally until recently. Indeed, nowadays, there seems to exist strong evidence of wave-like energy transfer through quantum coherence in photosynthetic systems (van Grondelle and

Novoderezhkin (2011), Marais *et al.* (2018), but also see Wilkins and Dattani (2015) and Duan *et al.* (2017) for an opposing view). This was possible originally with femtosecond transient absorption spectroscopy (Chachisvilis *et al.* (1994), Vos *et al.* (1994), Kumble *et al.* (1996), Agarwal *et al.* (2000), Novoderezhkin *et al.* (2000), Novoderezhkin *et al.* (2004)) and more recently with two dimensional electronic spectroscopy (Brixner *et al.* (2005)). The first observations were done on the Fenna–Matthews–Olson (FMO) complex of (photosynthetic) green sulfur bacteria at cryogenic temperatures (Engel *et al.* (2007)), but subsequent studies extended the original results to cover physiological temperatures and other organisms including plants (Schlau-Cohen *et al.* (2009), Panitchayangkoon *et al.* (2010), Collini *et al.* (2010)).

Connectionist approach to quantum consciousness

In spite of the growing evidence of quantum effects in biology, at first sight it seems improbable that quantum mechanics could play any significant role in any brain process involved in the psychological features associated with conscious awareness. Quantum effects are tiny compared to the scale of these physiological or cognitive events, and the consensus is that the mechanisms underlying brain function are well understood and can be described in terms of classical physics. Indeed, the attitude towards such proposals from the neuroscientific community has been typically negative. One crucial argument against the existence of quantum effects in the human brain is that decoherence should destroy such effects in the high temperature, noisy biological tissue that constitutes the nervous system. Indeed, macroscopic quantum phenomena such as superconductivity and superfluidity occur in a cryogenic regime. For many years, this argument has been a fundamental limitation to the notion of consciousness arising from purely quantum effects. However, as outlined in the last section, recent experimental results in the new field of quantum biology have made this argument much less stringent, as it is now clear that important physiological processes such as photosynthesis would benefit from non trivial quantum effects such as long-lived quantum coherence.

Another argument against a possible role of quantum physics in brain function is that there is no need for it as an explanatory aid: why use sophisticated and more mathematically involved ideas from quantum physics when neuroscience can be expressed in terms of very simple concepts requiring at most high school physics? Nevertheless, we should be cautious about this point: as we discuss later, some authors have suggested that consciousness could be irreducibly quantum (for reviews see Jedlicka (2017) and Adams and Petruccione (2020)).

Under this new light, a question can be naturally posed: at what level does consciousness appear? At first sight it seems that either consciousness is the result of the classical, chemical and electrical interactions between neurons, and in particular is the result of the integration of information on large neuronal networks, or it arises within cellular structures inside neurons via quantum coherence.

We propose that these two perspectives have actually a lot in common and that perhaps consciousness is an emergent phenomena at several spatio-temporal levels within the nervous system. As explained above, we propose a general connectionist approach, in which consciousness naturally arises as an emergent phenomena in brain networks in which an equilibrium is reached between noise and correlation. In the following we very briefly examine several possible intra-cellular candidates to such types of networks within the realm of quantum theories of consciousness, and we put them in the context of a general connectionist theory of consciousness.

Perhaps the most important candidate for a sub-cellular neural correlates of consciousness are microtubules (Hameroff (2006), Tuszynski (2006)). Microtubules are polymers of the protein called tubulin, and constitute the main part of cytoskeleton in eukaryotic and some prokaryotic cells (Alberts *et al.* (2002)). They form, together with microtubules associated proteins (MAP), acting and intermediate filaments, a dense network inside those cells. Microtubules have important physiological roles: they constitute the "skeleton" of cells (contributing to its shape and structure and connecting organelles) and are crucial for molecular transport within cells, by means of kinesin motor proteins. They are also central to cell division. Microtubules are constantly created within the cytoplasm, by polymerization of tubulin dimmers (α and β tubulin proteins).

According to one of the most outstanding theories of "quantum consciousness" in the brain, the orchestrated objective reduction theory (Orch OR), quantum coherence in networks of microtubules within neurons constitute a neural correlate of consciousness (Hameroff (1998), Hameroff and Penrose (2014)). Orch OR focuses on neurons, and in particular in the neuronal soma (cell body) and dendrites. The reason is that microtubules within dendrites and neural soma are interrupted

and of mixed polarity, and are connected to MAPs to form recursive networks that prevent them to disassemble (Hameroff and Penrose (2014), Adams and Petruccione (2020)). On the contrary, microtubules in axons are unipolar and continuous, forming radial, regular arrangements. In cells other than neurons, microtubules are unstable, disassembling in various ways (Adams and Petruccione (2020)). Therefore, the authors of Orch OR suggest that microtubules networks within neurons body and dendrites (contrary to those in axons and within other types of cells) could better support information processing.

According to Orch OR there are several reasons why microtubules networks are associated to consciousness (Hameroff and Penrose (2014)). The most important argument relates to the action of anesthetics. Indeed, anesthetics constitute a probe into consciousness. According to Turing, "the only thing we are sure about consciousness is that it is soluble in chloroform" (Rinaldi (2014)). Crucially, during anesthesia consciousness is suppressed, whereas neuronal activity can still be measured in the brain (Hameroff (2006)). Furthermore, certain anesthetics that influence cognition and conscious experience seem to involve microtubules (Adams and Petruccione (2020)). Relatedly, anesthetics are effective on any organism (Kelz and Mashour (2019)). How can anesthetic inhibit some responses to the environment in unicellular organisms such as slime molds that do not even have a nervous system (Keller and Segel (1970), Perouansky (2012))? This cannot be explained by a general anesthetic action on synaptic effects (Richards (2002), Voss *et al.* (2019), Adams and Petruccione (2020)). Relatedly, complex responses in simple organisms such as the paramecium seem to be connected to the cytoskeleton. So, it could be hypothesised that cognition in general and in particular consciousness (or loss of it through anesthesia effects) are linked to the cytoskeleton and in particular microtubules. They may constitute, for unicellular organisms but also for larger organism, what neurons constitute for the brain: basic information processing units.

Furthermore, the authors of Orch OR argue that, while action potentials (neuronal firing) represent a main mean of communication between neurons giving rise to synaptic potentials, it is the integration of these synaptic activity in the dendrites and cell bodies that is fundamental for neuronal information transmission. Also, the authors point out that gamma synchronization is related to somatic and dendritic integration of potentials (Hameroff and Penrose (2014)). And, as pointed out above, microtubule networks seem to be more suitable for information processing in bodies and dendrites of neurons. Therefore, they identify gamma synchronization with microtubules networks.

In the original version of Orch OR, Hameroff and Penrose explored the possibility of a quantum superposition of mechanical conformations coupled to London force electric dipoles. In later versions of the theory, they locate the quantum effects within the constituent aromatic rings (phenylalanine, tyrosine, and tryptophan) that make up the tubulin proteins (Adams and Petruccione (2020)), mediated via magnetic dipoles. In any case, they propose the existence of temporal correlation between microtubules, sustained by quantum coherence. The authors even go further and reason that a network of microtubules connected by quantum coherence can develop not only inside neurons but, via their connection with electrical and chemical synapses, they could extend across entire brain areas (Hameroff and Penrose (2014)). In this sense, it would be very difficult to distinguish a network of microtubules connected by quantum coherence or even entanglement, from a network of neurons connected by more familiar, synaptically-related connections and neural synchronization. One crucial distinction could be that if microtubules constitute a network where information is processed, as in Orch OR, the potential possibilities for the brain are enormously enhanced, because they constitute a dense network inside neurons.

However, it remains to answer for how long can quantum coherence be maintained in microtubules at physiological temperature. According to Tegmark, not for too long to be important in a neurophysiological context, because decoherence destroy quantum correlation between microtubules after approximately 10^{-12} s, too short a time lapse as compared with the scale of neuron firing (milliseconds, 10^{-3} s) (Tegmark (2000)). However, a re-calculation due to Hagan *et al.*, more in accordance to Orch OR theory, produced decoherence times of the order of 10 to 100 μ s (10^{-5} s - 10^{-4} s), much closer to the time scales of neural activity (Hagan *et al.* (2002)).

Going back to anesthetics, there are quantum effects that seem to be implicated in the action of general anesthetics (Hameroff (2006), Hameroff and Penrose (2014)). These are better understood by reasoning in analogy with photosynthesis. Indeed, as we sketched above, photosynthesis energy transfer and charge separation is enhanced by quantum effects. Quantum betas are present in light-harvesting complexes in plants and bacteria (Engel *et al.* (2007)). Similarly, tryptophan residues in tubulin proteins are also capable of quantum coherence transfer, at least according to theoretical models (Craddock *et al.* (2014)). Craddock and co-workers point out that a possible action of

anesthetics is the inhibition of quantum coherence between microtubules (Craddock *et al.* (2015)). More specifically, the authors argued that anesthetics act on "quantum channels" (hydrophobic pathways of tryptophan rings in tubulin). Other, very recent evidence comes from spin changes in anesthetized fruit flies. Anesthetic action seems to generally disrupt electronic activity, suggesting that nuclear spin of anesthetic molecules may influence their efficiency (Turin *et al.* (2014)).

Apart from research related with microtubules, the possibility of spin (which is an intrinsically quantum property) having an influence on conscious phenomena has been worked out by several authors. An important line of research in this sense examines quantum entanglement between atoms associated with crucial neural phenomena such as action potentials. Wu and Hu proposed a pioneer theory of spin-mediated consciousness, based on entanglement between different types of atoms within cell membranes (Hu and Wu (2004)). More recently, Fisher built a theory of quantum cognition by processing information with entangled phosphorus atoms. Phosphorus is a crucial substance for organisms; it can be found, for example, within metabolically important molecules such as ATP. In particular, it is found within Posner molecules $\text{Ca}_9(\text{PO}_4)_6$, believed to be important for bone formation (Posner and Betts (1975)). According to Fisher and co-workers, entanglement of phosphorus atoms within Posner molecules is "protected" from the environment, in the sense that decoherence times are very long (of the order of hours or even days (Fisher (2015), Swift *et al.* (2018))). Although other authors estimated a much shorter decoherence time (37 minutes or even much shorter), it is in any case much longer than those estimated in quantum coherence in microtubules. In Fisher's proposal, entangled Posner molecules could elicit the simultaneous firing of neurons, by a complex set of events leading to the release of neurotransmitters in the corresponding neurons. The mechanism through which entangled Posner molecules release neurotransmitters remains hypothetical, but the result would be that of temporal correlation of the activities of neurons containing entangled Posner molecules.

Finally, other studies point towards the possibility of quantum entanglement of photons in the brain (for a review of such ideas see Adams and Petruccione (2020)). For example, it has been hypothesized using a theoretical model, that photons can be used for communication between neurons, by traveling through myelinated axons, that could serve as waveguides, in analogy with the way photons travel in optical fibers (Kumar *et al.* (2016)). Of course, such ideas remain speculative.

Whatever the details of such quantum entanglement mechanism, the relevance for the current paper is that entangled atoms or photons within neurons can be correlated across the whole brain, effectively constituting a network of correlations, or functional network, exactly as those described in neuronal network theories of consciousness.

Summarizing, we identify two types of quantum correlations in sub-cellular neurons that could in principle be related to cognition and conscious experience. The first is quantum coherence between adjacent structures, such as microtubules or other cytoskeleton structures, leading to coordinated activity within neurons. The second is whole brain (or at least macroscopic) correlations between entangled atoms within molecules, in turn leading to a cascade of events synchronizing neural activity. To our knowledge, there is little if no experimental evidence that any of these processes are directly related to cognition or consciousness, and for now they rely on hypothetical thinking and indirect evidence. One such indirect evidence comes from the field of quantum biology, and in particular for photosynthetic light harvesting complexes, where both quantum coherence and entanglement has been demonstrated.

The quantum Goldilock effect

An important clue to understand the possible importance of quantum effects in cognition comes from the effect of noise in biological tissue, and in particular the effect of noise on information processing. As we explained above, conscious states tend to appear at intermediate levels of noise and correlation in functional networks (Guevara Erra *et al.* (2016)). At first sight, it seems paradoxical that consciousness is achieved at intermediate levels of noise. Indeed, normally, noise is considered as a hindrance for information-processing systems and devices. However, there are instances where noise can be beneficial in biological systems, specially in information processing in the nervous system, where it can even enhance neural processing (Ward (2003), Faisal *et al.* (2008), McDonnell and Ward (2011)). A paradigmatic example is stochastic resonance, where a sub-threshold input can be detected at appropriate levels of noise, and not at lower levels (Gammaitoni *et al.* (1998)). Strikingly, in the phenomenon of psychophysical stochastic resonance, an image




Figure 1: Shannon entropy (S) of brain networks (a measure of network complexity) as a function of network connectivity (a measure of correlation in brain activity) for conscious and non-conscious states. Each data point is a measure of brain activity using either electroencephalography or magnetoencephalography. baseline = brain activity before a seizure. sws 3-4 = slow wave 3-4, a phase of deep sleep. During awake and "baseline" (non-seizure) states (conscious states), entropy is large, as compared with coma, seizure and sws 3-4 states (non-conscious). Conscious states appear at intermediate values (not too large, nor too low) of correlations in brain networks (C), also corresponding to intermediate values of noise in brain activity. The inverted U curve represents the dependency of S with C , calculated using a statistical mechanics model (Guevara Erra *et al.* (2016))




Figure 2: Excitation transfer in a fully connected network modeling a light harvesting complex. The probability of excitation transfer to the reaction center, or "sink" (p_{sink}) is a function of the noise in the environment. Noise is quantified by the local dissipation rate Γ_{N+1} , the transfer of energy from the exciton to individual nodes in the network, and a pure dephasing noise rate γ , randomizing the phase of the exciton and therefore destroying quantum coherence. Maximal exciton transfer efficiency is achieved at intermediate levels of noise and coherence. The network is constituted by $N = 5$ sites. Reproduced from [Caruso *et al.* \(2009\)](#), with the permission of AIP Publishing.).

embedded in noise is not visible until sufficient noise is added (Velazquez *et al.* (2007)). There is also experimental evidence that stochastic resonance enhance transport in biological systems (Douglass *et al.* (1993), Shang *et al.* (2007)). Interesting, converging ideas appear in apparently unrelated studies of transport in quantum systems. Indeed, one question that puzzled researchers for decades (starting with Schrödinger in his masterpiece "What is life?" Schrodinger and Schrödinger (1992)) is how can certain biological processes be possible in spite of molecular noise. New developments in quantum biology points toward the possibility that an effect similar to stochastic resonance, that is, maximal efficiency of certain processes at intermediate levels of noise, also operates at the quantum level of description, and can be applied to important physiological processes related to charge and energy transport, such as photosynthesis and respiration (Plenio and Huelga (2008), Rebentrost *et al.* (2009), Caruso *et al.* (2009)).

A particularly striking example comes from models of photosynthetic complexes in bacteria: the Fenna-Matthews-Olsen (FMO) complex already mentioned above, that has been thoroughly studied by means of X-ray crystallography and spectroscopy (Collini *et al.* (2010), Panitchayangkoon *et al.* (2010)). This system is so well known that a model can be constructed with a single free-parameter: temperature (Lloyd *et al.* (2011)). As it turns out, the passage of excitons from chlorophyll molecules to the reaction center can be modeled as a "quantum Brownian motion", explaining the large efficiency of the transport processes (Brookes (2017)).

The efficiency is however, temperature dependent (Lloyd *et al.* (2011)). At very low temperatures, interference of the different possible paths taken by the exciton are mainly destructive, leading to a localization of the exciton after a few molecular steps. At higher temperatures, the environment induce a stronger decoherence of the exciton paths, decreasing the destructive interference and therefore increasing the distance achieved by the exciton, that can eventually arrive to the reaction center. At even higher temperatures, decoherence is so strong that that the exciton essentially does not move. Maximal efficiency was reached at around 290 K, and maintained for several degrees Kelvin in both directions (Lloyd *et al.* (2011)). This is the range of temperatures of the water within which these bacteria live. Within this range, efficiency of transport was very close to 100%. We emphasize that the only parameter changed in the model was temperature, which is a measure of molecular disorder and can be easily interpreted as "noise". In other words, at large and low levels of "molecular noise" the exciton does not move very far, whereas at intermediate levels maximal efficiency is achieved.

Interestingly, this kind of effect should be present in any kind of partially coherent quantum transport process in a disordered system, not only in photosynthesis and was baptized by the authors of the model as Environmentally Assisted Quantum Transport (ENAQT). Furthermore, the authors suggest a "quantum Goldilock effect", in that natural selection drives certain biological systems into a level of quantum coherence at which optimal efficiency is achieved. This speculation is sustained by the observation of a convergence of temporal scales in photosynthesis (Lloyd *et al.* (2011), Novo *et al.* (2016)).

A concrete example of this kind of effect is shown in Fig 2, that shows the influence of dephasing noise on the transfer rate of excitons between rings of light harvesting photosynthetic units (?). An optimal performance in excitonic transfer is achieved at an intermediate level of dephasing noise. Too much noise or too little are detrimental for conduction (Huelga and Plenio (2013)).

We speculate that the quantum Goldilock effect is the analogous, at the quantum level, to the fine tuning between noise and coherence resulting in conscious states in the brain, observed during epileptogenic activity and sleep and described above (Guevara Erra *et al.* (2016)). During epileptic seizures and deep stages of sleep, brain functionality diminishes as compared with normal states, and the information in brain networks also diminishes. We may say that the corresponding brain activity is not optimal, and the brain is less efficient at information processing. Comparing figures 1 and 2 may be illustrative in this sense. In both figures efficiency is achieved at intermediate levels of noise and coherence. Huelga and Plenio suggested that the optimal efficiency in transport is achieved at an intermediate level, located between the classical regime, where dephasing noise dominates, and the quantum one, dominated by coherence (Huelga and Plenio (2013)). Based on these observations, it is inviting to conjecture that a general principle, the self-organization of highly correlated networks, underlies the emergence of conscious phenomena.

Conclusions

The main question we have tried to address can be summarized as follows: what is the size of the smallest illuminated spots, the building blocks in the Cartesian theatre of consciousness mentioned above? We have presented plausible arguments in favor of a multi-scale connectionist approach to consciousness, with quantum effects playing a possible role in enhancing the generation of conscious phenomena. The experimental evidence in favor of such idea is not too clear, but there are certain puzzling questions about the mechanism of action of general anesthesia that may require the incorporation of quantum physics. Also, the last years had witnessed the development of the new field of quantum biology. Quantum coherence and entanglement seem to play a role in several physiological processes. Under this perspective, it is instructive to analyze the problem of the emergence of consciousness under a generalized connectionist point of view, where networks associated to conscious experience appear at different scales, including sub-cellular, and can be affected or enhanced by quantum correlations. We identify quantum coherence and entanglement as possible sources of temporal correlation, in analogy as neural synchronization supposedly underlies large neuronal network information integration. We also identify the quantum Goldilock effect to be analogue to the fine tuning of parameters observed in global neuronal networks during the emergency of consciousness. In both cases, it is the interplay between noise and correlation in the system that leads to an increase in system's efficiency. We speculate that is this constitute a general principle underlying the emergence of conscious awareness.

References

- Crick, F.; Clark, J. The astonishing hypothesis. *Journal of Consciousness Studies* **1994**, *1*, 10–16.
- Edelman, G.M.; Tononi, G. A universe of consciousness: how matter becomes imagination. *Contemporary Psychology* **2003**, *48*, 92.
- Koch, C.; Massimini, M.; Boly, M.; Tononi, G. Neural correlates of consciousness: progress and problems. *Nature Reviews Neuroscience* **2016**, *17*, 307.
- Tassi, P.; Muzet, A. Defining the states of consciousness. *Neuroscience & Biobehavioral Reviews* **2001**, *25*, 175–191.
- Velazquez, J.L.P. On the emergence of cognition: from catalytic closure to neuroglial closure. *Journal of Biological Physics* **2020**, pp. 1–25.
- Luisi, P.L. About various definitions of life. *Origins of Life and Evolution of the Biosphere* **1998**, *28*, 613–622.
- Monnard, P.A.; Walde, P. Current ideas about prebiological compartmentalization. *Life* **2015**, *5*, 1239–1263.
- Velazquez, J.L.P. Finding simplicity in complexity: general principles of biological and nonbiological organization. *Journal of biological physics* **2009**, *35*, 209–221.
- Tegmark, M. Importance of quantum decoherence in brain processes. *Physical review E* **2000**, *61*, 4194.
- Koch, C.; Hepp, K. Quantum mechanics in the brain. *Nature* **2006**, *440*, 611–611.
- Baars, B.J.; Edelman, D.B. Consciousness, biology and quantum hypotheses. *Physics of life reviews* **2012**, *9*, 285–294.
- Adams, B.; Petruccione, F. Quantum effects in the brain: A review. *AVS Quantum Science* **2020**, *2*, 22901. doi:black10.1116/1.5135170.
- O'Brien, G.; Opie, J. A connectionist theory of phenomenal experience. *Behavioral and Brain Sciences* **1999**, *22*, 127–148.
- Maia, T.V.; Cleeremans, A. Consciousness: Converging insights from connectionist modeling and neuroscience. *Trends in cognitive sciences* **2005**, *9*, 397–404.

- McGovern, K.; Baars, B.J. Cognitive theories of consciousness. *The Cambridge handbook of consciousness* **2007**, pp. 177–205.
- Seth, A. Models of consciousness. *Scholarpedia* **2007**, *2*, 1328.
- Tononi, G. Consciousness and Complexity. *Science* **1998**, *282*, 1846–1851. doi:10.1126/science.282.5395.1846.
- Tononi, G. An information integration theory of consciousness. *BMC neuroscience* **2004**, *5*, 42.
- Tononi, G. Consciousness as integrated information: a provisional manifesto. *The Biological Bulletin* **2008**, *215*, 216–242.
- Baars, B.J. *A cognitive theory of consciousness*; Cambridge University Press, 1993.
- Baars, B.J. Metaphors of consciousness and attention in the brain. *Trends in Neurosciences* **1998**, *21*, 58–62.
- Baars, B.J. The conscious access hypothesis: origins and recent evidence. *Trends in cognitive sciences* **2002**, *6*, 47–52.
- Mountcastle, V. An organizing principle for cerebral function: the unit module and the distributed system. *The mindful brain* **1978**.
- Dehaene, S.; Sergent, C.; Changeux, J.P. A neuronal network model linking subjective reports and objective physiological data during conscious perception. *Proceedings of the National Academy of Sciences* **2003**, *100*, 8520–8525.
- Singer, W. Neuronal synchrony: a versatile code for the definition of relations? *Neuron* **1999**, *24*, 49–65.
- Engel, A.K.; Fries, P.; Singer, W. Dynamic predictions: oscillations and synchrony in top-down processing. *Nature Reviews Neuroscience* **2001**, *2*, 704–716.
- Varela, F.; Lachaux, J.P.; Rodriguez, E.; Martinerie, J. The brainweb: phase synchronization and large-scale integration. *Nature reviews neuroscience* **2001**, *2*, 229–239.
- Fries, P. A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. *Trends in cognitive sciences* **2005**, *9*, 474–480.
- Velazquez, J.L.P.; Frantseva, M. *The Brain-Behaviour Continuum: The Subtle Transition Between Sanity and Insanity*; World Scientific, 2011.
- Pikovsky, A.; Rosenblum, M.; Kurths, J. *Synchronization: a universal concept in nonlinear sciences*; Vol. 12, Cambridge university press, 2003.
- Buzsáki, G. *Rhythms of the Brain*; Oxford University Press, 2009.
- Perez Velazquez, J.; Guevara Erra, R.; Rosenblum, M. The Epileptic Thalamocortical Network is a Macroscopic Self-Sustained Oscillator: Evidence from Frequency-Locking Experiments in Rat Brains. *Scientific reports* **2015**, *5*.
- Nunez, P.L.; Srinivasan, R. *Electric fields of the brain: The neurophysiology of EEG*; Oxford University Press, New York, 1981.
- Buzsáki, G.; Draguhn, A. Neuronal oscillations in cortical networks. *science* **2004**, *304*, 1926–1929.
- Velazquez, J.L.P.; Mateos, D.M.; Erra, R.G. On a simple general principle of brain organization. *Frontiers in Neuroscience* **2019**, *13*.
- Llinás, R.; Ribary, U.; Contreras, D.; Pedroarena, C. The neuronal basis for consciousness. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **1998**, *353*, 1841–1849.
- Jiruska, P.; De Curtis, M.; Jefferys, J.G.; Schevon, C.A.; Schiff, S.J.; Schindler, K. Synchronization and desynchronization in epilepsy: controversies and hypotheses. *The Journal of physiology* **2013**, *591*, 787–797.

- Guevara Erra, R.; Mateos, D.; Wennberg, R.; Perez Velazquez, J. Statistical mechanics of consciousness: Maximization of information content of network is associated with conscious awareness. *Physical Review E* **2016**, *94*.
- Mateos, D.; Wennberg, R.; Guevara Erra, R.; Velazquez, J.P. Consciousness as a global property of brain dynamic activity. *Physical Review E* **2017**, *96*, 062410.
- Mateos, D.; Guevara Erra, R.; Wennberg, R.; Velazquez, J.P. Measures of entropy and complexity in altered states of consciousness. *Cognitive neurodynamics* **2018**, *12*, 73–84.
- Kauffman, S.A. *The origins of order: Self-organization and selection in evolution*; OUP USA, 1993.
- Feinberg, T.E.; Mallatt, J.M. *Consciousness demystified*; MIT Press, 2018.
- Feinberg, I. Efference copy and corollary discharge: implications for thinking and its disorders. *Schizophrenia bulletin* **1978**, *4*, 636.
- Haken, H. *Information and self-organization: A macroscopic approach to complex systems*; Springer Science & Business Media, 2006.
- Turner, S.; Hanel, R.; Klimek, P. *Introduction to the theory of complex systems*; Oxford University Press, 2018.
- Nunez, P.L.; Srinivasan, R. *Electric Fields of the Brain: The Neurophysics of EEG* **2005**.
- Guevara Erra, R.; Perez Velazquez, J.L.; Rosenblum, M. Neural synchronization from the perspective of non-linear dynamics. *Frontiers in computational neuroscience* **2017**, *11*, 98.
- Griffiths, D.J.; Schroeter, D.F. *Introduction to quantum mechanics*; Cambridge University Press, 2018.
- Bes, D. *Quantum mechanics: a modern and concise introductory course*; Springer Science & Business Media, 2012.
- Brookes, J.C. Quantum effects in biology: golden rule in enzymes, olfaction, photosynthesis and magnetodetection. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **2017**, *473*, 20160822. doi:black10.1098/rspa.2016.0822.
- Anderson, P.W. More Is Different. *Science* **1972**, *177*, 393–396. doi:black10.1126/science.177.4047.393.
- Lambert, N.; Chen, Y.N.; Cheng, Y.C.; Li, C.M.; Chen, G.Y.; Nori, F. Quantum biology. *Nature Physics* **2013**, *9*, 10–18.
- Fleming, G.R.; Scholes, G.D.; Cheng, Y.C. Quantum effects in biology. *Procedia Chemistry* **2011**, *3*, 38–57.
- Engel, G.S.; Calhoun, T.R.; Read, E.L.; Ahn, T.K.; Mančal, T.; Cheng, Y.C.; Blankenship, R.E.; Fleming, G.R. Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature* **2007**, *446*, 782–786.
- Panitchayangkoon, G.; Hayes, D.; Fransted, K.A.; Caram, J.R.; Harel, E.; Wen, J.; Blankenship, R.E.; Engel, G.S. Long-lived quantum coherence in photosynthetic complexes at physiological temperature. *Proceedings of the National Academy of Sciences* **2010**, *107*, 12766–12770.
- Ishizaki, A.; Fleming, G.R. Quantum coherence in photosynthetic light harvesting. *Annu. Rev. Condens. Matter Phys.* **2012**, *3*, 333–361.
- McFadden, J.; Al-Khalili, J. *Life on the edge: the coming of age of quantum biology*; Broadway Books, 2016.
- Marais, A.; Adams, B.; Ringsmuth, A.K.; Ferretti, M.; Gruber, J.M.; Hendrikx, R.; Schuld, M.; Smith, S.L.; Sinayskiy, I.; Krüger, T.P.J.; Petruccione, F.; van Grondelle, R. The future of quantum biology. *Journal of The Royal Society Interface* **2018**, *15*, 20180640. doi:black10.1098/rsif.2018.0640.

- Alberts, B.; Bray, D.; Lewis, J.; Raff, M.; Roberts, K.; Watson, J. Molecular biology of the cell. Garland Pub. Inc., London **2002**.
- Rüdiger, W. The chromophore. In *Photomorphogenesis in plants*; Springer, 1986; pp. 17–33.
- Lloyd, S.; Mohseni, M.; Shabani, A.; Rabitz, H. The quantum Goldilocks effect: on the convergence of timescales in quantum transport. *arXiv preprint arXiv:1111.4982* **2011**.
- Knox, R.S. Theory of Excitons. *Solid State Phys.* **1963**, *5*.
- Novoderezhkin, V.I.; van Grondelle, R. Physical origins and models of energy transfer in photosynthetic light-harvesting. *Physical Chemistry Chemical Physics* **2010**, *12*, 7352–7365.
- Mohseni, M.; Rebentrost, P.; Lloyd, S.; Aspuru-Guzik, A. Environment-assisted quantum walks in photosynthetic energy transfer. *The Journal of chemical physics* **2008**, *129*, 11B603.
- Fassioli, F.; Dinshaw, R.; Arpin, P.C.; Scholes, G.D. Photosynthetic light harvesting: excitons and coherence. *Journal of The Royal Society Interface* **2014**, *11*, 20130901.
- Franck, J.; Teller, E. Migration and photochemical action of excitation energy in crystals. *The Journal of Chemical Physics* **1938**, *6*, 861–872.
- van Grondelle, R.; Novoderezhkin, V.I. Quantum effects in photosynthesis. *Procedia Chemistry* **2011**, *3*, 198–210.
- Wilkins, D.M.; Dattani, N.S. Why quantum coherence is not important in the Fenna–Matthews–Olsen complex. *Journal of chemical theory and computation* **2015**, *11*, 3411–3419.
- Duan, H.G.; Prokhorenko, V.I.; Cogdell, R.J.; Ashraf, K.; Stevens, A.L.; Thorwart, M.; Miller, R.J.D. Nature does not rely on long-lived electronic quantum coherence for photosynthetic energy transfer. *Proceedings of the National Academy of Sciences* **2017**, *114*, 8493–8498. doi:black10.1073/pnas.1702261114.
- Chachisvilis, M.; Pullerits, T.; Jones, M.; Hunter, C.; Sundström, V. Vibrational dynamics in the light-harvesting complexes of the photosynthetic bacterium *Rhodobacter sphaeroides*. *Chemical physics letters* **1994**, *224*, 345–354.
- Vos, M.H.; Jones, M.R.; Hunter, C.N.; Breton, J.; Martin, J.L. Coherent nuclear dynamics at room temperature in bacterial reaction centers. *Proceedings of the National Academy of Sciences* **1994**, *91*, 12701–12705.
- Kumble, R.; Palese, S.; Visschers, R.; Dutton, P.L.; Hochstrasser, R.M. Ultrafast dynamics within the B820 subunit from the core (LH-1) antenna complex of *Rs. rubrum*. *Chemical physics letters* **1996**, *261*, 396–404.
- Agarwal, R.; Krueger, B.P.; Scholes, G.D.; Yang, M.; Yom, J.; Mets, L.; Fleming, G.R. Ultrafast energy transfer in LHC-II revealed by three-pulse photon echo peak shift measurements. *The Journal of Physical Chemistry B* **2000**, *104*, 2908–2918.
- Novoderezhkin, V.; Monshouwer, R.; van Grondelle, R. Electronic and Vibrational Coherence in the Core Light-Harvesting Antenna of *Rhodospseudomonas v. iridis*. *The Journal of Physical Chemistry B* **2000**, *104*, 12056–12071.
- Novoderezhkin, V.I.; Yakovlev, A.G.; Van Grondelle, R.; Shuvalov, V.A. Coherent nuclear and electronic dynamics in primary charge separation in photosynthetic reaction centers: a Redfield theory approach. *The Journal of Physical Chemistry B* **2004**, *108*, 7445–7457.
- Brixner, T.; Stenger, J.; Vaswani, H.M.; Cho, M.; Blankenship, R.E.; Fleming, G.R. Two-dimensional spectroscopy of electronic couplings in photosynthesis. *Nature* **2005**, *434*, 625–628.
- Schlau-Cohen, G.S.; Calhoun, T.R.; Ginsberg, N.S.; Read, E.L.; Ballottari, M.; Bassi, R.; van Grondelle, R.; Fleming, G.R. Pathways of energy flow in LHCI from two-dimensional electronic spectroscopy. *The Journal of Physical Chemistry B* **2009**, *113*, 15352–15363.

- Collini, E.; Wong, C.Y.; Wilk, K.E.; Curmi, P.M.; Brumer, P.; Scholes, G.D. Coherently wired light-harvesting in photosynthetic marine algae at ambient temperature. *Nature* **2010**, *463*, 644–647.
- Jedlicka, P. Revisiting the Quantum Brain Hypothesis: Toward Quantum (Neuro)biology? *Frontiers in Molecular Neuroscience* **2017**, *10*. doi:black10.3389/fnmol.2017.00366.
- Hameroff, S.R. The Entwined Mysteries of Anesthesia and Consciousness. *Anesthesiology* **2006**, *105*, 400–412. doi:black10.1097/0000542-200608000-00024.
- Tuszynski, J.A. *The emerging physics of consciousness*; Springer Science & Business Media, 2006.
- Hameroff, S. Quantum computation in brain microtubules? The Penrose–Hameroff ‘Orch OR’ model of consciousness. *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* **1998**, *356*, 1869–1896.
- Hameroff, S.; Penrose, R. Consciousness in the universe: A review of the ‘Orch OR’ theory. *Physics of life reviews* **2014**, *11*, 39–78.
- Rinaldi, A. Reawakening anaesthesia research. *EMBO reports* **2014**, *15*, 1113–1118.
- Kelz, M.B.; Mashour, G.A. The biology of general anesthesia from Paramecium to primate. *Current Biology* **2019**, *29*, R1199–R1210.
- Keller, E.F.; Segel, L.A. Initiation of slime mold aggregation viewed as an instability. *Journal of theoretical biology* **1970**, *26*, 399–415.
- Perouansky, M. The quest for a unified model of anesthetic action. *Anesthesiology* **2012**, *117*, 465.
- Richards, C. Anaesthetic modulation of synaptic transmission in the mammalian CNS. *British journal of anaesthesia* **2002**, *89*, 79–90.
- Voss, L.J.; Garcia, P.S.; Hentschke, H.; Banks, M.I. Understanding the effects of general anesthetics on cortical network activity using ex vivo preparations. *Anesthesiology: The Journal of the American Society of Anesthesiologists* **2019**, *130*, 1049–1063.
- Hagan, S.; Hameroff, S.R.; Tuszyński, J.A. Quantum computation in brain microtubules: Decoherence and biological feasibility. *Physical Review E* **2002**, *65*, 061901.
- Craddock, T.J.A.; Friesen, D.; Mane, J.; Hameroff, S.; Tuszynski, J.A. The feasibility of coherent energy transfer in microtubules. *Journal of the Royal Society Interface* **2014**, *11*, 20140677.
- Craddock, T.J.A.; Hameroff, S.R.; Ayoub, A.T.; Klobukowski, M.; Tuszynski, J.A. Anesthetics Act in Quantum Channels in Brain Microtubules to Prevent Consciousness. *Current Topics in Medicinal Chemistry* **2015**, *15*, 523–533. doi:black10.2174/1568026615666150225104543.
- Turin, L.; Skoulakis, E.M.C.; Horsfield, A.P. Electron spin changes during general anesthesia in *Drosophila*. *Proceedings of the National Academy of Sciences* **2014**, *111*, E3524–E3533. doi:black10.1073/pnas.1404387111.
- Hu, H.; Wu, M. Spin-mediated consciousness theory: possible roles of neural membrane nuclear spin ensembles and paramagnetic oxygen. *Medical Hypotheses* **2004**, *63*, 633–646.
- Posner, A.S.; Betts, F. Synthetic amorphous calcium phosphate and its relation to bone mineral structure. *Accounts of Chemical Research* **1975**, *8*, 273–281.
- Fisher, M.P. Quantum cognition: The possibility of processing with nuclear spins in the brain. *Annals of Physics* **2015**, *362*, 593–602.
- Swift, M.W.; Van de Walle, C.G.; Fisher, M.P. Posner molecules: from atomic structure to nuclear spins. *Physical Chemistry Chemical Physics* **2018**, *20*, 12373–12380.
- Kumar, S.; Boone, K.; Tuszyński, J.; Barclay, P.; Simon, C. Possible existence of optical communication channels in the brain. *Scientific Reports* **2016**, *6*. doi:black10.1038/srep36508.

- Caruso, F.; Chin, A.W.; Datta, A.; Huelga, S.F.; Plenio, M.B. Highly efficient energy excitation transfer in light-harvesting complexes: The fundamental role of noise-assisted transport. *The Journal of Chemical Physics* **2009**, *131*, 09B612.
- Ward, L.M. Synchronous neural oscillations and cognitive processes. *Trends in cognitive sciences* **2003**, *7*, 553–559.
- Faisal, A.A.; Selen, L.P.J.; Wolpert, D.M. Noise in the nervous system. *Nature Reviews Neuroscience* **2008**, *9*, 292–303. doi:black10.1038/nrn2258.
- McDonnell, M.D.; Ward, L.M. The benefits of noise in neural systems: bridging theory and experiment. *Nature Reviews Neuroscience* **2011**, *12*, 415–426.
- Gammaitoni, L.; Hänggi, P.; Jung, P.; Marchesoni, F. Stochastic resonance. *Reviews of modern physics* **1998**, *70*, 223.
- Velazquez, J.L.P.; Dominguez, L.G.; Erra, R.G. Fluctuations in neuronal synchronization in brain activity correlate with the subjective experience of visual recognition. *Journal of biological physics* **2007**, *33*, 49–59.
- Douglass, J.K.; Wilkens, L.; Pantazelou, E.; Moss, F. Noise enhancement of information transfer in crayfish mechanoreceptors by stochastic resonance. *Nature* **1993**, *365*, 337–340.
- Shang, Y.; Claridge-Chang, A.; Sjulson, L.; Pypaert, M.; Miesenböck, G. Excitatory local circuits and their implications for olfactory processing in the fly antennal lobe. *Cell* **2007**, *128*, 601–612.
- Schrodinger, R.; Schrödinger, E. *What is life?: With mind and matter and autobiographical sketches*; Cambridge University Press, 1992.
- Plenio, M.B.; Huelga, S.F. Dephasing-assisted transport: quantum networks and biomolecules. *New Journal of Physics* **2008**, *10*, 113019. doi:black10.1088/1367-2630/10/11/113019.
- Rebentrost, P.; Mohseni, M.; Kassal, I.; Lloyd, S.; Aspuru-Guzik, A. Environment-assisted quantum transport. *New Journal of Physics* **2009**, *11*, 033003.
- Novo, L.; Mohseni, M.; Omar, Y. Disorder-assisted quantum transport in suboptimal decoherence regimes. *Scientific reports* **2016**, *6*, 18142.
- Huelga, S.F.; Plenio, M.B. Vibrations, quanta and biology. *Contemporary Physics* **2013**, *54*, 181–207. doi:black10.1080/00405000.2013.829687.
- Chin, A.W.; Prior, J.; Rosenbach, R.; Caycedo-Soler, F.; Huelga, S.F.; Plenio, M.B. The role of non-equilibrium vibrational structures in electronic coherence and recoherence in pigment-protein complexes. *Nature Physics* **2013**, *9*, 113–118. doi:black10.1038/nphys2515.