

## Semen quality as a potential susceptibility indicator to SARS-CoV-2 insults in polluted areas

Luigi Montano<sup>1\*</sup>, Ian Marc Bonapace<sup>2</sup>, Francesco Donato<sup>3</sup>, Pietro Massimiliano Bianco<sup>4</sup>, Antonino Guglielmino<sup>5</sup>, Marina Piscopo<sup>6</sup>

<sup>1</sup>Andrology Unit, Local Health Authority (ASL) Salerno, EcoFoodFertility Project, Coordination Unit, Oliveto Citra, Via M. Clemente, 84020, Oliveto Citra, (SA), Italy

<sup>2</sup>Department of Biotechnology and Life Sciences, University of Insubria (VA), Italy

<sup>3</sup>Department of Medical and Surgical Specialties Radiological Sciences and Public Health, Unit of Hygiene, Epidemiology, and Public Health, University of Brescia, Italy.

<sup>4</sup>ISPRA, Italian Institute for Environmental Protection and Research, Via Vitaliano Brancati 60, 00144 Roma, Italy

<sup>5</sup>Unit of Reproductive Medicine (UMR) Catania, Italy

<sup>6</sup>Marina Piscopo, Department of Biology, University of Naples Federico II, 80126, Napoli, Italy

\*Corresponding author:

Luigi Montano, MD, Andrology Unit, Local Health Authority (ASL) Salerno, EcoFoodFertility Project, Coordination Unit, Oliveto Citra, Via M. Clemente, 84020, Oliveto Citra, SA Italy. Tel: 00390828797263 email: [l.montano@aslsalerno.it](mailto:l.montano@aslsalerno.it) or [luigimontano@gmail.com](mailto:luigimontano@gmail.com)

ORCID ID: 0000-00015670-9439

### Abstract

High levels of air pollution can contribute to high rate of the COVID-19 outbreaks. Air pollutants induce oxidative stress, inflammatory process, immune imbalance and coagulation at systemic level, making the organism susceptible to complications caused by various pathogens, including viruses, resulting in a possible important damage co-factor. Sperm cells are highly sensitive to the pro-oxidant effects of environmental pollutants, and may represent an important alarm bell indicating that the burden of environmental pressure in a certain area is causing damage to humans. A comparison of the maps of COVID-19 case fatality rates, male infertility rates and air pollution may suggest a way to understand the dynamics of the virus impact. Semen quality may be considered as an early and sensitive environmental marker, and also a potential susceptibility indicator to viral insults (including SARS-CoV-2) in heavily polluted areas. Therefore, assessing

the burden of environmental exposure of a given population and its potential susceptibility to insults through early biological stress indicators may be helpful for predicting the risk of the adverse effects by the SARS-CoV-2 epidemic .

**Key words:** air pollution, COVID-19, semen quality, environmental marker, health marker, oxidative stress, SARS-CoV-2

### **Air pollution and COVID-19**

A recent review point out that chronic exposure to air pollutants in the most polluted areas of the world leads to more severe and lethal forms of COVID-19<sup>1</sup>. Indeed, a preprint study, carried out in 3,080 counties in the United States calculated an 8% increase in the COVID-19 death rate adjusted by 20 potential confounding factors for 1 microgram/m<sup>3</sup> elevation of fine particulate matter with a diameter of 2,5 µm or less (PM<sub>2,5</sub>)<sup>2</sup>. Furthermore, in 2003, during SARS infection in China, scientists reported that Case Fatality Rate (CFR) in the most polluted areas was twice as high as in the least polluted ones<sup>3</sup>. Another preprint study has reported a higher CFR of COVID-19 in Wuhan, China, with increasing of air particular matter with diameter of 10 µm or less (PM<sub>10</sub>) and PM<sub>2,5</sub> after adjusting for humidity and temperature<sup>4</sup>. Moreover, some authors pointed out that the regions of Northern Italy most affected by COVID-19, were among those with the highest levels of PM<sub>10</sub> and PM<sub>2,5</sub> in Europe<sup>5-7</sup>. A recent preprint found a significant correlation between high levels of PM<sub>2,5</sub>, carbon monoxide (CO), nitrogen dioxide (NO<sub>2</sub>) and COVID-19 spread and mortality in Italy, U.S.A. and China<sup>8</sup>.

COVID-19 spread after initial outbreaks (China, South Korea, Iran) occurred in Italy before that observed in the rest of Europe and Eastern U.S.A., between latitude 30° and 50° North, in Winter (December to April), when weather patterns, such as low temperature (between 5 and 11 °C) and low specific and absolute humidity of 3-6 g/kg and 4-7 g/m<sup>3</sup> respectively were favourable for the spread of a respiratory virus<sup>9</sup>. Furthermore, it is significant that in the same seasonal period (May for Brazilian winter) the highest local peak out of this belt is in Sao Paulo, Brazil, a heavily polluted city (figure 1)<sup>10</sup>. It's known that in the winter period the air pollution rates are higher and the cities where COVID-19 has hit hardest are those with PM<sub>10</sub>, PM<sub>2,5</sub> and NO<sub>2</sub> annual average above the WHO recommended values of 20 µg/m<sup>3</sup>, 10 µg/m<sup>3</sup> and 40 µg/m<sup>3</sup> respectively (figure 1)<sup>11-17</sup>. Stability of SARS-CoV-2 in different environmental conditions supports the hypothesis that air pollution may favour the human-to-human chain of transmission of the infection, particularly in elevated crowding situations, COVID-19 severity and risk of related death<sup>1,18</sup>.

## **Pollution and susceptibility to viral insults**

The World Health Organization (WHO) estimates that about a quarter of diseases is due to prolonged exposure to environmental pollutants<sup>19,20</sup>, including cardiovascular and chronic degenerative disease, premature deaths and reproduction dysfunctions<sup>19,21–23</sup> along with lifestyle, as reported in the European Code against Cancer<sup>24</sup>. Moreover, environmental pollutants can increase susceptibility to non-communicable diseases (NCDs) and determine a significant decline in body's defences, possibly due to also transgenerational effects that may partly explain the worldwide disease burden consisting not only in non-communicable but also communicable diseases (dengue, yellow fever, tuberculosis, etc.)<sup>24–36</sup>. Furthermore, it shows that this is transmitted to the following generations, by reducing the defence ability toward viral pathogens<sup>26</sup>.

In particular, the chronic exposure to the air-borne finest fraction of particulate matter (PM<sub>2.5</sub>), not only induces inflammation to alveolar district, but, passing the alveolar barrier, it reaches the blood and hence the peripheral tissues, inducing oxidative stress both directly and through the guest response to the chemical insult. This contributes to the activation of inflammasome and, particularly, NLRP3, influencing the maturation and secretion of Cytokine such as IL-1 beta and IL-18 both involved in the systemic inflammatory syndrome and in the conditions facilitating the pathogen agent virulence, including vascular leakage and coagulopathy<sup>37–40</sup>.

SARS-CoV-2 is able to unleash a fast process of autoimmune dysregulation, by inducing a significant Cytokine storm, mainly TNF- $\alpha$ , IL-6 e IL-1 $\beta$ , IL-17, IL-18, in genetically predisposed individuals<sup>41</sup>. Such a process could be even more important when pre-existing environmental factors have already altered regulatory mechanisms for Cytokine release and/or even when there are some polymorphisms for IL-6, such as in specific populations or ethnic groups who in fact make them more susceptible to virus complications<sup>42</sup>. Specifically, all populations are susceptible to COVID-19, but the elderly, individuals with chronic diseases or low immune defences, pregnant women and newborns are most exposed to complications<sup>43–45</sup>.

In addition, air pollutants induce oxidative stress, inflammatory process, immune imbalance and coagulation at systemic level, making the organism susceptible to complications arising from the pathogen agents, including SARS-CoV-2, resulting as a possible important damage co-factor<sup>46</sup>.

This is even more true in those areas of the world where poor air quality (figure 2) could, favour viral contagion and/or increased virulence, decreasing the antioxidant and immune defences of the organism.

## **Sperm decline in polluted areas**

Many data suggest a decline in sperm parameters in different areas of the world, particularly in developed ones or those undergoing strong industrial development with high levels of air pollution<sup>47</sup>. A systematic review and meta-regression analysis reported a decline of total sperm counts by 59.3% in Europe, U.S.A., Canada and New Zealand between 1971 and 2011<sup>48</sup>. In Asia, the infertility rate of Iranian men has increased by 20% over the last 20 years and in China, out of a total of 30,636 young donors, the sperm concentration and percentage of sperm with normal morphology decreased from  $68 \times 10^6/\text{mL}$  to  $47 \times 10^6/\text{mL}$  and from 31.8% to 10.8%, respectively<sup>49,50</sup>. In Brasil, in the past 23 years a median reduction of 0.24 million/mL of spermatozoa per year was reported<sup>51</sup>.

The male reproductive system is, indeed, extremely sensitive to environmental pollutants. In particular, chronic exposure to high levels of  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$  and individual air pollutants such as  $\text{NO}_2$  and sulfur dioxide ( $\text{SO}_2$ ) are negatively associated with sperm count, motility and testicular volume in infertile subjects<sup>52,53</sup>. The mechanisms of spermatogenesis damage by environmental agents are largely unknown, although radical oxygen species (ROS) imbalance and associated oxidative stress may be the common denominator through which pollutants alter the most sensitive parameters of seminal quality such as sperm count, motility, morphology and integrity of sperm DNA<sup>54-56</sup>. As a matter of fact, sperm cells are highly sensitive to the pro-oxidant effects of environmental pollutants, due to the limited volume of the cytoplasmic space, with less antioxidant defence, and sperm membrane lipids are target of ROS<sup>57</sup>. In this regard, we recently reported that male gametes are the most sensitive cells to the accumulation of damaged DNA and showed, through molecular investigations, that the sperm nuclear basic proteins from samples of men living in polluted areas have a new and unexpected behaviour, resulting involved in DNA oxidative damage<sup>58</sup>. The COVID-19 CFR map of the most important outbreaks in the world areas presenting, in recent decades, a negative trend in sperm quality, together with the average annual levels of  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$  and  $\text{NO}_2$ , shows a certain overlapping (figure 1). It could be argued that sperm decline may represent the earliest clinical sign of environmental pressure.

### **Human semen as environmental and health marker**

We and other authors indicate human semen as a “sentinel biomarker” of subclinical biological effect suitable for monitoring the impact of adverse environmental exposures<sup>59-63</sup>.

Semen quality has also been found to reflect individuals' general health condition, as recent studies showed an association between semen quality and the onset of chronic diseases, with male infertility as a predictor of future hospitalization and overall mortality<sup>64-68</sup>.

We speculate that this observation could help understanding the dynamics that may have facilitated the COVID-19 severity in polluted areas. As an early and sensitive environmental and health marker, semen quality could be considered as a potential susceptibility indicator of external insults to the health of the general population and be used for health risk management, innovative prevention programs and health surveillance, especially in heavily polluted areas<sup>62,63</sup>. However, although high population density (according for human-to-human transmission mechanisms), climatic characteristics, age, comorbidity, different capacity of health systems to face the pandemic and prevention policies adopted in the various countries currently play the central role, we should not overlook the possible facilitating contribution of pollution in increasing the risk for people living in areas with higher environmental pressure to the COVID-19 impact. Moreover, it must be considered that higher incidence of non-communicable diseases (NCDs) and male infertility are reported in these same areas because of a complex interaction between factors of chronic chemical and physical exposure, along with the contribution of lifestyle behavioural risk factors and individuals' genetic background. In particular, human sperm decline in the last decades, represent an important alarm bell indicating how the burden of environmental pressure is becoming increasingly unsustainable. In this perspective, the first signs of damage to organo-sentinel systems such as the male reproductive system, detectable by the reduction in semen quality over time can be an opportunity to know the health status of the population in a given environmental context, including susceptibility to the virus impact in that population, and predict the medium to long term adverse effects on human health.

However, in our opinion semen quality as an early environmental and health marker could help the policy makers to intervene promptly in areas with significant environmental criticalities in order to reduce air, water and soil pollution with an integrated approach in a One Health perspective, where the sharing of information between different professional figures (clinicians, biologists, chemists, virologists, veterinarians, economists, epidemiologists) can succeed to find a systemic approach that could be effective on a global scale<sup>69</sup>. Above all, it is necessary to estimate effectiveness of the measures adopted to safeguard the health of community and also its social and productive organization, with the aim to avoid, or at least reduce, the rapid and destructive spread of future viruses.

## Refecences

1. Domingo, J. L. & Rovira, J. Effects of air pollutants on the transmission and severity of respiratory viral infections. *Environmental Research* **187**, 109650 (2020).
2. Wu, X., Nethery, R., Benjamin, M., Braun, D. & Dominici, F. *Exposure to air pollution and COVID-19 mortality in the United States: A nationwide cross-sectional study. medRxiv : the preprint server for health sciences* (2020). doi:10.1101/2020.04.05.20054502.
3. Cui, Y. *et al.* Air pollution and case fatality of SARS in the People's Republic of China: an ecologic study. *Environ Health* **2**, 15 (2003).
4. Yao, Y. *et al.* Temporal Association Between Particulate Matter Pollution and Case Fatality Rate of COVID-19 in Wuhan, China. *medRxiv* 2020.04.09.20049924 (2020) doi:10.1101/2020.04.09.20049924.
5. Conticini, E., Frediani, B. & Caro, D. Can atmospheric pollution be considered a co-factor in extremely high level of SARS-CoV-2 lethality in Northern Italy? *Environ Pollut* **261**, 114465 (2020).
6. Martelletti, L. & Martelletti, P. Air Pollution and the Novel Covid-19 Disease: a Putative Disease Risk Factor. *SN Compr Clin Med* 1–5 (2020) doi:10.1007/s42399-020-00274-4.
7. Fattorini, D. & Regoli, F. Role of the chronic air pollution levels in the Covid-19 outbreak risk in Italy. *Environ Pollut* **264**, 114732 (2020).
8. Pansini, R. & Fornacca, D. Initial evidence of higher morbidity and mortality due to SARS-CoV-2 in regions with lower air quality. *medRxiv* 2020.04.04.20053595 (2020) doi:10.1101/2020.04.04.20053595.
9. Sajadi, M. M. *et al.* Temperature, Humidity, and Latitude Analysis to Estimate Potential Spread and Seasonality of Coronavirus Disease 2019 (COVID-19). *JAMA Netw Open* **3**, e2011834–e2011834 (2020).
10. Andrade, M. de F. *et al.* Air quality in the megacity of São Paulo: Evolution over the last 30 years and future perspectives. *Atmospheric Environment* **159**, 66–82 (2017).
11. Update of WHO Global Air Quality Guidelines. <https://www.euro.who.int/en/health-topics/environment-and-health/air-quality/activities/update-of-who-global-air-quality-guidelines>.
12. US EPA, O. Air Quality - Cities and Counties. *US EPA* <https://www.epa.gov/air-trends/air-quality-cities-and-counties> (2016).
13. Qualidade do Ar. <https://qualidadedoar.org.br/>.

14. Wu, X. PM<sub>2.5</sub>/PM<sub>10</sub> Ratio Prediction Based on a Long Short-term Memory Neural Network in Wuhan, China. **3**, (2019).
15. Beloconi, A., Chrysoulakis, N., Lyapustin, A., Utzinger, J. & Vounatsou, P. Bayesian geostatistical modelling of PM<sub>10</sub> and PM<sub>2.5</sub> surface level concentrations in Europe using high-resolution satellite-derived products. *Environment International* **121**, 57–70 (2018).
16. Pansini, R. & Fornacca, D. Higher virulence of COVID-19 in the air-polluted regions of eight severely affected countries. *medRxiv* 2020.04.30.20086496 (2020)  
doi:10.1101/2020.04.30.20086496.
17. QA 2019 conf stampa 08012020.pdf.
18. Two mechanisms for accelerated diffusion of COVID-19 outbreaks in regions with high intensity of population and polluting industrialization: the air pollution-to-human and human-to-human transmission dynamics | medRxiv.  
<https://www.medrxiv.org/content/10.1101/2020.04.06.20055657v1>.
19. Forouzanfar, M. H. *et al.* Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet* **388**, 1659–1724 (2016).
20. WHO | Almost a quarter of all disease caused by environmental exposure. *WHO*  
<https://www.who.int/mediacentre/news/releases/2006/pr32/en/>.
21. WHO | First WHO Global Conference on Air Pollution and Health, 30 October – 1 November 2018. *WHO* <http://www.who.int/airpollution/events/conference/en/>.
22. Ji, W. & Zhao, B. Estimating Mortality Derived from Indoor Exposure to Particles of Outdoor Origin. *PLoS One* **10**, (2015).
23. Cohen, A. J. *et al.* The global burden of disease due to outdoor air pollution. *J. Toxicol. Environ. Health Part A* **68**, 1301–1307 (2005).
24. Anderson, A. S. *et al.* European Code against Cancer 4th Edition: Obesity, body fatness and cancer. *Cancer Epidemiol* **39 Suppl 1**, S34-45 (2015).
25. Heindel, J. J. The developmental basis of disease: Update on environmental exposures and animal models. *Basic Clin. Pharmacol. Toxicol.* **125 Suppl 3**, 5–13 (2019).
26. Post, C. M. *et al.* The Ancestral Environment Shapes Antiviral CD8+ T cell Responses across Generations. *iScience* **20**, 168–183 (2019).
27. Horan, T. S., Marre, A., Hassold, T., Lawson, C. & Hunt, P. A. Germline and reproductive tract effects intensify in male mice with successive generations of estrogenic exposure. *PLoS Genet.* **13**, e1006885 (2017).

28. Suzuki, K. The developing world of DOHaD. *J Dev Orig Health Dis* **9**, 266–269 (2018).
29. Bianco-Miotto, T., Craig, J. M., Gasser, Y. P., van Dijk, S. J. & Ozanne, S. E. Epigenetics and DOHaD: from basics to birth and beyond. *J Dev Orig Health Dis* **8**, 513–519 (2017).
30. Soubry, A. POHaD: why we should study future fathers. *Environ Epigenet* **4**, (2018).
31. Soubry, A. Epigenetics as a Driver of Developmental Origins of Health and Disease: Did We Forget the Fathers? *Bioessays* **40**, (2018).
32. Curley, J. P., Mashoodh, R. & Champagne, F. A. Epigenetics and the origins of paternal effects. *Horm Behav* **59**, 306–314 (2011).
33. Soubry, A., Hoyo, C., Jirtle, R. L. & Murphy, S. K. A paternal environmental legacy: evidence for epigenetic inheritance through the male germ line. *Bioessays* **36**, 359–371 (2014).
34. Braun, K. & Champagne, F. A. Paternal influences on offspring development: behavioural and epigenetic pathways. *J. Neuroendocrinol.* **26**, 697–706 (2014).
35. Zhao, Z.-H., Schatten, H. & Sun, Q.-Y. Environmentally induced paternal epigenetic inheritance and its effects on offspring health. *Reproductive and Developmental Medicine* **1**, 89 (2017).
36. Xavier, M. J., Roman, S. D., Aitken, R. J. & Nixon, B. Transgenerational inheritance: how impacts to the epigenetic and genetic information of parents affect offspring health. *Hum. Reprod. Update* **25**, 518–540 (2019).
37. Ghio, A. J., Carraway, M. S. & Madden, M. C. Composition of air pollution particles and oxidative stress in cells, tissues, and living systems. *J Toxicol Environ Health B Crit Rev* **15**, 1–21 (2012).
38. Li, Y. *et al.* Indoor nanoscale particulate matter-induced coagulation abnormality based on a human 3D microvascular model on a microfluidic chip. *Journal of Nanobiotechnology* **17**, 20 (2019).
39. Tsai, D.-H. *et al.* Effects of short- and long-term exposures to particulate matter on inflammatory marker levels in the general population. *Environ Sci Pollut Res Int* **26**, 19697–19704 (2019).
40. Caso, F. *et al.* Could Sars-coronavirus-2 trigger autoimmune and/or autoinflammatory mechanisms in genetically predisposed subjects? *Autoimmun Rev* **19**, 102524 (2020).
41. Masters, S. L., Simon, A., Aksentjevich, I. & Kastner, D. L. Horror Autoinflammaticus: The Molecular Pathophysiology of Autoinflammatory Disease. *Annu Rev Immunol* **27**, 621–668 (2009).

42. El-Maadawy, E. A., Talaat, R. M., Ahmed, M. M. & El-Shenawy, S. Z. Interleukin-6 promotor gene polymorphisms and susceptibility to chronic hepatitis B virus in Egyptians. *Human Immunology* **80**, 208–214 (2019).
43. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* **395**, 497–506 (2020).
44. Chen, H. *et al.* Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *The Lancet* **395**, 809–815 (2020).
45. Working Group for the Prevention and Control of Neonatal 2019-nCoV Infection in the Perinatal Period of the Editorial Committee of Chinese Journal of Contemporary Pediatrics. [Perinatal and neonatal management plan for prevention and control of 2019 novel coronavirus infection (1st Edition)]. *Zhongguo Dang Dai Er Ke Za Zhi* **22**, 87–90 (2020).
46. Glencross, D. A., Ho, T.-R., Camiña, N., Hawrylowicz, C. M. & Pfeffer, P. E. Air pollution and its effects on the immune system. *Free Radic. Biol. Med.* **151**, 56–68 (2020).
47. Mann, U., Shiff, B. & Patel, P. Reasons for worldwide decline in male fertility. *Curr Opin Urol* **30**, 296–301 (2020).
48. Levine, H. *et al.* Temporal trends in sperm count: a systematic review and meta-regression analysis. *Hum Reprod Update* **23**, 646–659 (2017).
49. Safarinejad, M. R. Infertility among couples in a population-based study in Iran: prevalence and associated risk factors. *Int. J. Androl.* **31**, 303–314 (2008).
50. Huang, C. *et al.* Decline in semen quality among 30,636 young Chinese men from 2001 to 2015. *Fertil. Steril.* **107**, 83-88.e2 (2017).
51. Siqueira, S. *et al.* Changes in seminal parameters among Brazilian men between 1995 and 2018. *Scientific Reports* **10**, 6430 (2020).
52. Zhang, J., Cai, Z., Ma, C., Xiong, J. & Li, H. Impacts of Outdoor Air Pollution on Human Semen Quality: A Meta-Analysis and Systematic Review. *BioMed Research International* vol. 2020 e7528901 <https://www.hindawi.com/journals/bmri/2020/7528901/> (2020).
53. Ambient sulfur dioxide could have an impact on testicular volume from an observational study on a population of infertile male - Abstract - Europe PMC. <https://europepmc.org/article/ppr/ppr125689>.
54. Jurewicz, J., Dziewirska, E., Radwan, M. & Hanke, W. Air pollution from natural and anthropic sources and male fertility. *Reprod Biol Endocrinol* **16**, (2018).
55. Nowicka-Bauer, K. & Nixon, B. Molecular Changes Induced by Oxidative Stress that Impair Human Sperm Motility. *Antioxidants (Basel)* **9**, (2020).

56. Bosco, L. *et al.* Sperm DNA fragmentation: An early and reliable marker of air pollution. *Environ. Toxicol. Pharmacol.* **58**, 243–249 (2018).
57. Aitken, R. J., Gibb, Z., Baker, M. A., Drevet, J. & Gharagozloo, P. Causes and consequences of oxidative stress in spermatozoa. *Reprod. Fertil. Dev.* **28**, 1–10 (2016).
58. Lettieri, G. *et al.* Discovery of the Involvement in DNA Oxidative Damage of Human Sperm Nuclear Basic Proteins of Healthy Young Men Living in Polluted Areas. *International Journal of Molecular Sciences* **21**, 4198 (2020).
59. Nordkap, L., Joensen, U. N., Blomberg Jensen, M. & Jørgensen, N. Regional differences and temporal trends in male reproductive health disorders: semen quality may be a sensitive marker of environmental exposures. *Mol. Cell. Endocrinol.* **355**, 221–230 (2012).
60. Bergamo, P. *et al.* Human semen as an early, sensitive biomarker of highly polluted living environment in healthy men: A pilot biomonitoring study on trace elements in blood and semen and their relationship with sperm quality and RedOx status. *Reprod. Toxicol.* **66**, 1–9 (2016).
61. Montano, L. *et al.* The role of human semen for assessing environmental impact on human health in risk areas: Novels and early biomarkers of environmental pollution. EcoFoodFertility project. *Reproductive Toxicology* **72**, 44–45 (2017).
62. Montano, L., Bergamo, P., Lorenzetti, S., Andreassi, M. G. The Role of Human Semen as an Early and Reliable Tool of Environmental Impact Assessment on Human Health | IntechOpen. <https://www.intechopen.com/books/spermatozoa-facts-and-perspectives/the-role-of-human-semen-as-an-early-and-reliable-tool-of-environmental-impact-assessment-on-human-health> (2018).
63. Montano, L. Reproductive biomarkers as early indicators for assessing environmental health risk. in Marfe, G., and Di Stefano C. Toxic Waste Management and Health Risk, Bentham Science Publishers eBook eISBN 978-981-14-5474-5 (Forthcoming)
64. Eisenberg, M. L., Li, S., Behr, B., Pera, R. R. & Cullen, M. R. Relationship between semen production and medical comorbidity. *Fertility and Sterility* **103**, 66–71 (2015).
65. Choy, J. T. & Eisenberg, M. L. Male infertility as a window to health. *Fertil. Steril.* **110**, 810–814 (2018).
66. Pisarska, M. D. Fertility Status and Overall Health. *Semin. Reprod. Med.* **35**, 203–204 (2017).
67. Glazer, C. *et al.* Male Infertility and Risk of Nonmalignant Chronic Diseases: A Systematic Review of the Epidemiological Evidence. *Seminars in Reproductive Medicine* **35**, 282–290 (2017).
68. Jensen, T. K., Jacobsen, R., Christensen, K., Nielsen, N. C. & Bostofte, E. Good semen quality and life expectancy: a cohort study of 43,277 men. *Am. J. Epidemiol.* **170**, 559–565 (2009).

69. Mackenzie, J. S. & Jeggo, M. The One Health Approach—Why Is It So Important? *Trop Med Infect Dis* **4**, (2019).

### **Acknowledgments**


The authors thank Dr. Federico Marra for graphic elaboration of the figure 2.

### **Author contributions**


LM developed the concept and drafted the manuscript, PMB collected the data, IMB, AG, FD and MP edited and revised manuscript, LM, IMB, FD, PMB, AG and MP approved the final version of the manuscript.

### **Competing interests**

The authors declare no competing financial interest.



**Figure 1:** Overlapping between Case Fatality Rate maps (from pink to brown) of the most important COVID-19 outbreaks, air pollution and world areas presenting, in recent decades, a negative trend in sperm quality. Some cities of the same areas with  $PM_{10}$ ,  $PM_{2.5}$  and  $NO_2$  annual average above the WHO recommended values of  $20 \mu\text{g}/\text{m}^3$ ,  $10 \mu\text{g}/\text{m}^3$  and  $40 \mu\text{g}/\text{m}^3$  respectively.



**Figure 2:** World map of tropospheric  $NO_2$  concentrations from the Copernicus Sentinel-5P satellite (2019)