

Review

High Homogenization Pressures to Improve Food Quality, Functionality and Sustainability

Jose Mesa¹, Leidy I. Hinestroza^{1,2}, Cristina Barrera¹, Lucía Seguí¹, Ester Betoret^{3*}, Noelia Betoret^{1*}

¹ Institute of Food Engineering for Development, Universitat Politècnica de València, CP 46022 València, Spain; jomegu1@etsiamn.upv.es (J.M.); mbarpu@tal.upv.es (C.B.); lusegil@upvnet.upv.es (L.S.)

² Grupo de Valoración y Aprovechamiento de la Biodiversidad, Universidad Tecnológica del Chocó. AA.292, Calle 22 No. 18B-10, Quibdó-Chocó, CP. 270001 Colombia; leihicor@doctor.upv.es (L.H.)

³ Instituto de Agroquímica y Tecnología de Alimentos, Consejo Superior de Investigaciones Científicas, 46980 Paterna, Spain.

* Correspondence: noebeval@tal.upv.es (N.B.); ester_betoret@hotmail.es (E.B.). Tel: +34963877000 (83624)

Abstract: The interest in high homogenization pressure technology has grown over the years. It is a green technology with low energy consumption, not generating high CO₂ emissions or polluting effluents. The main food applications derive from its effect on particle size, causing a more homogeneous distribution of fluid elements (particles, globules, droplets, aggregates, etc.) and favouring the release of intracellular components; and its effect on the structure and configuration of chemical components such as polyphenols and macromolecules such as carbohydrates (fibres) and proteins (also microorganisms and enzymes). The challenges of the 21st century lead food industry processing towards obtaining food with high nutritional quality and taking advantage of waste to obtain ingredients with specific properties. For this purpose, soft and non-thermal technologies such as high pressures homogenization have a huge potential. The objective of this work is to review how the need to combine safety, functionality and sustainability in food industry has conditioned the last decade applications of high-pressure homogenization technology.

Keywords: High homogenization pressure; food functionality; bioactive components; agri-food waste; sustainability.

1. Introduction

In the homogenization process, a fluid is forced to pass through a gap, causing energy transformations that directly affect the dissolved, dispersed or emulsified components. The fluid undergoes mechanical (shear, hydrodynamic and cavitation effects) stress and an increase in temperature (thermal effect) of approximately 2-3 °C for every 10 MPa of homogenization pressure [1]. These affect fluid structure and properties and also those of its constituent elements (particles, molecules, globules, droplets, aggregates, granules, etc.). The particle size decreases and a more homogeneous distribution of the elements is achieved, facilitating operations such as mixing and emulsification.

Initially the homogenization operation was introduced as a manufacturing step in the dairy industry. This operation reduced the size of the fat globules, increasing the stability of the emulsion and thus the physical and chemical stability of milk. It had a great impact on the quality of dairy products such as condensed milk, curd or ice cream. The applied pressure was less than 30 MPa and it was applied in one or two steps. However, technological development increased greatly having an impact on the design and geometry of the homogenization valves, which allowed working at higher pressures and with very short processing times (some seconds) [2]. High homogenization pressures operation was established at the beginning of the 2000's as an alternative non-thermal treatment in the food industry and applications were extended to industries other than dairy and to other fields such as textile or biotechnologic.

Currently, the existence of valves of different geometries has given rise to different equipment able to work at pressures higher than 400 MPa. Thus, a distinction is made among standard

homogenization for pressures between 0 and 50 MPa, high pressure homogenization (HPH) for pressures between 50 and 300 MPa and ultra-high pressure homogenization (UHPH) for pressures equal to or greater than 400 MPa. Processing efficiency is modulated by applying various pressure ranges or combining a pressure value with a specific number of passes through the equipment [3,4]. In addition, the possibility of operating in continuous conditions for a great diversity of pumpable fluids has made it possible to extend the application to activation/inactivation of enzymes, reduction of microbial load, mixing, dispersion, emulsification or encapsulation processes, cell breakage processes and modification of proteins or macromolecules to obtain ingredients or additives with differentiated properties.

Nowadays, concern about food functionality and sustainability has triggered research for increasing the bioavailability and bioaccessibility of active components and probiotics and for the extraction of macro and micro-molecules from food by-products. Moreover, the challenge of increasing the nutritional characteristics of food must be combined with a reduction of the environmental impact and food security. In this context, alternative, soft and non-thermal technologies such as high pressures homogenization have a huge potential. The objective of this work is to review how the need to combine safety, functionality and sustainability has conditioned the applications of high pressure homogenization technology in food.

2. Last decade evolution and main applications

Publications in peer-reviewed journals show that the main applications of HPH in food have the following objectives:

- Conservation and safety by decreasing the microbial load and inactivating enzymes. It occurs as a consequence of thermal effect derived from mechanical stress or from structural changes in proteins.
- Recovery and extraction of proteins, fibrous materials and bioactive compounds of special interest (mainly polyphenols) and increase of the functionality considered in terms of technologic use (stabilization of emulsions and dispersions, flow capacity and viscosity modifications, emulsifying activity improvement...). Mechanical stresses and hydrodynamic effects induce cell disruption favouring the release of intracellular content or structural components of cell wall. Moreover, dispersed particles or fat droplets can be reduced in size and modified in structure.
- Increase of functionality in terms of health effect (increase bioaccessibility, bioavailability or probiotic effect). These effects result from favouring the release of bioactive compounds, modification of biopolymer structure and development of novel particle interactions and networking. Micro or nanocapsules can also be developed.

As it can be observed in Figure 1a, between 2000 and 2009 HPH were used mainly for the extraction of proteins, although a large number of research works were aimed to microorganisms and enzymes inactivation, contributing to food preservation and safety. The last decade (from 2010 until now) revealed a significant increase (74.87%) in the total number of scientific articles published. The main areas in which there has been an increase greater than the total value are related to the use of HPH for microorganisms inactivation, fibre extraction, and above all, that related to bioactive and probiotic components. Application of HPH to extract or increase functionality of food bioactive compounds and to improve probiotic effect grew by 89% and 87,9% respectively (figure 1a). The increasing interest of consumers and the food industry in improving the organoleptic and nutritional quality of foods, along with the concern for food waste valorization might explain this result.

Figure 1b. shows the evolution in the number of published research works related to HPH application with regard to the type of food. Although the majority of works belongs to the area of fruit juices, the largest growth has occurred in plant-based beverages and food waste. The huge increase in plant-based beverages consumption [5] and the general concern about food processing waste related issues are responsible for this fact. Moreover, HPH technology is identified as a green technology, due to short processing times, low energy consumption, low CO₂ emissions and the fact that it does not require polluting solvents.




Figure 1. Number of scientific articles published regarding the main food application areas of HPH (a) and the different types of food to which it applies (b). % INCREASE has been calculated as the difference in the number of articles published between the periods 2010-2020 and 2000-2009, divided by the number of articles published in the period 2000-2009, expressed as percentage. (Source: Science Direct. The following keywords and their combinations have been used as main search terms: high homogenization pressures, non-thermal technologies, food processing, encapsulation, functional food, bioactive components, probiotics, microbial load, enzyme inactivation, protein extraction, milk, fruit juice, food waste, plant-based beverages and alcoholic beverages).

This increase in research works based on HPH technology is also due to the development of new homogenization equipment that works at elevated pressures (up to 400 MPa) and supports specific conditions. Since the invention of adjustable valves in 1930 [6], the possibilities of homogenization technology have increased. The geometry and design of the valve determines the mechanical effects on the treated fluid. In 1982 the invention of the Gaulin Micro-Gap valve [7] greatly boosted the efficiency of the process, making high homogenization pressures possible in subsequent years and, more recently, the ultra-high homogenization pressures technology. In general, improvements have been obtained in all fields of application, making it an efficient tool with great potential for use in the food industry [8].

3. Preservation and safety

HPH treatment for enzyme and microbial inactivation is one of the technologies that have been used in recent years as an alternative to thermal processes that, in most cases, cause undesirable effects such as non-enzymatic browning, cooked flavour or valuable components degradation. It has been demonstrated that HPH treatment at processing pressures higher than 100 MPa contributes to microbial load reduction and enzyme inactivation. As previously stated the heating derived from homogenization (a rise of about 2.5 °C per 10 MPa) together with structural modifications of cell wall are the main effects responsible for microorganisms or enzymes inactivation. Final impact of HPH on microorganism viability or enzyme activity depend on several factors such as processing pressure, microbial strain or enzyme and food matrix.

In general, it has been verified that gram-negative bacteria exhibit a greater susceptibility to inactivation by HPH than gram-positive ones, due to the reduced content of peptidoglycan in the cell wall that makes it thinner and therefore easier to disrupt. Fungi and yeasts seem to have a susceptibility intermediate in between gram-positive and gram-negative bacteria, probably due to their wall structure, which is thicker but more complex than that from gram-positive bacteria [9]. For bacterial spores inactivation, pressures up to 400 MPa and additional hurdles are required [8].

Table 1. Research works evaluating the decrease in microbial load in different food products by HPH.

PRODUCT	TREATMENT	TERMS	MICROBIOLOGIC CONTROL	RESULTS	REFERENCE
Fruit juices (apricot and carrot)	HPH + rapid cooling	100 MPa (1-8 passes)	<i>Zygosaccharomyces bailii</i> 45	The juice type affected the yeast fate (growth or death) and viscosity change after HPH treatment	[8]
Mango nectar	HPH + thermal shock	200 MPa 10-20 s at 60-85 ° C	<i>A. niger</i> (COI 4573)	The combination of HPH with subsequent thermal shock was efficient in inactivating the heat resistant mould in the mango nectar	[10]
Banana juice	HPH + rapid cooling	0, 150, 200, 300 and 400 MPa	Total mesophilic bacteria	Pressures greater than 200 MPa were required to obtain a reduction of 4 logarithmic units	[11]
Apricot juice	HPH+ citral + rapid cooling	100 MPa (1,3,5 and 8 passes)	<i>Saccharomyces cerevisiae</i> SPA	Decrease of the viability of the yeasts following a linear tendency with the pressure. Improvement of the antimicrobial effect by adding citral	[12]
Mango juice (<i>Mangifera indica</i> L.)	HPH + heat treatment	40-190 MPa (1-5 passes)	Total plate count, moulds and yeasts	Complete inactivation of moulds and yeasts was achieved by 1 and 3 passes at 190 MPa and 60 °C, while the total plate count was less than 2.0 log CFU/mL	[13]
Mulberry juice (<i>Morus atropurpurea</i> Roxb.)	HPH + heat treatment + Addition of Dimethyl Dicarboxylate (DMDC)	200 MPa (1-3 passes)	Total count, yeast, mould and lactic acid bacteria	Combination treatment with 3 passes at 200 MPa and 250 mg DMDC/L, decreased total count to the level reached by heat treatment at 95 ° C	[14]
Lupine based drinks	HPH + refrigeration	50, 100 and 175 MPa (2,4,6 passes)	Total bacterial count, moulds and yeasts. <i>Bacillus cereuses</i> and coliform bacteria	At 175 MPa yeasts, moulds and coliforms were completely eliminated with 2 and 4 passes	[15]

Granada juice	HPH + low temperature pasteurization	100, 150 MPa (10 passes) 55 or 65°C for 15 s	<i>Escherichia coli</i> (ATCC 26) and <i>Saccharomyces pastorianus</i> (ATCC 42376)	HPH at 150 MPa followed by a low heat intensity at 65 °C for 15 s showed a reduction of 3 log CFU/ mL	[16]
Skim milk	Heat treatment + HPH	100-300 MPa	<i>Bacillus stearothermophilus</i> ATCC 7953 and <i>Clostridium sporogenes</i> PA 3679	The efficacy of HPH is similar to pasteurization and must be combined with other conservation technique	[17]
Milk	Heat treatment + HPH	300 MPa	Spores of <i>B. cereus</i> , <i>B. lincheniformis</i> , <i>B. sporothermodurans</i> , <i>B. coagulans</i> , <i>B. stearothermophilus</i> , and <i>B. subtilis</i>	Sterility at 300 MPa can be achieved with an initial milk temperature of 85 °C	[18]
Skim and whole milk concentrates	Heat treatment + HPH	Skim milk: 0,20,50,70, 100,120 and 150 MPa. Whole milk: 0,20,30,35 and 40 MPa.	Total count, coliforms, enterobacteriaceae, moulds and yeasts and <i>Staphylococco</i>	HPH at 120 MPa completely inactivates the microbial load of milk concentrates	[19]
Almond beveages	Heat treatment + HPH	200, 300 MPa (1,2 passes)	<i>Micrococcaceae</i> , <i>Bacillus cereus</i> and Mesophilic aerobic bacterias	Complete elimination of microbial growth when working with the highest pressure and with an inlet temperature of 65- 75 °C	[20]
Rice drink	HPH+ sonication	50-100 MPa (1-3 passes)	<i>Lactobacillus Plantarum</i> , <i>Lactobacillus Casei</i> , y <i>Bifidobacterium Animalis</i>	Reduction and elimination of post-acidification by lactic acid bacteria.	[21]
Tiger nuts' milk beverage	HPH + refrigeration	200 and 300 MPa	Psychrotrophic bacteria, Lactobacilli, Enterobacteriaceae and faecal coliforms	Improved shelf life and microbial inactivation compared to other heat treatments	[22]
Lagger beer	HPH + lysozyme addition	0-300 MPa	<i>Lactobacillus brevis</i> (CCT 3745)	The inhibitory concentration of lysozyme against <i>L. brevis</i> was 100 mg/ L. HPH at 100, 140, and 150 MPa promoted decimal reductions of 1, 3, and 6 in microbial counts	[23]

Pilsen beer	Heat treatment + HPH	100, 150, 200 and 250 MPa (1-3 passes)	<i>Lactobacillus del brueckii</i>	It is possible to inactivate the most common microorganisms of beer deterioration at 250 MPa. The effect increases with increasing number of passes. HPH at 150 MPa the best treatment by inducing yeast autolysis, also suitable to accelerate sur lie maturation	[24]
Wine	Chemical treatment + HPH	0, 50, 100 and 150 MPa	<i>Saccharomyces bayanus</i>		[25]

Significant work has been performed on the application of HPH to reduce microbial load in fruit juices. In this type of food, it has been observed that the presence of some aroma compounds and essential oils can greatly influence the effect of HPH treatment. Patrignani et al. [12] studied the effect of number of passes and citral addition on spoilage microbiota of apricot juice when subjected to HPH at 100 MPa. Their results showed that yeast cell viability decreased with the increase of passes and the relationship between both variables followed a linear trend. Moreover, the citral addition enhanced the effect of HPH, increasing the storage time by 6-8 days. To analyse the effect of the food matrix, the same authors compared the effect of HPH treatment at 100 MPa on viability loss of *S. cerevisiae* 635 inoculated at a level of about 6.0 Log₁₀ cfu/mL in apricot juice and carrot juice. In apricot juice a significant viability decreases (2.2 Log₁₀ cfu/ml) was obtained with only four repeated passes at 100 MPa. A further increase of the number of passes at 100 MPa did not significantly increase the effectiveness of HPH treatment. Concerning carrot juice, eight repeated passes at 100 MPa were unable to completely inactivate the inoculated cells. They concluded that because of the higher viscosity and sugar content, apricot juice required more passes in HPH treatment to reduce yeast load [26]. On the contrary, *Zygosaccharomyces bailii* 45 exhibited the same susceptibility to HPH treatment in both juices. Eight passes at 100 MPa allowed a yeast inactivation higher than 2.5 log CFU/mL regardless of the juice considered [27]. Nevertheless, [16] established that HPH treatment at 100 and 150 MPa was not sufficient to reduce total bacteria and yeast count in pomegranate juice and it needed to be combined with a thermal treatment at 65 °C for 15 s, to achieve the same effect than pasteurization at 75 °C.

Besides fruit juices, plant-based beverages are complex dispersions with suspended proteins and oil droplets that require a homogenization stage to stabilize them and extend its commercial life. HPH can be applied at pressures higher than 100 MPa using multiple passes for these purposes along with microbial cells destruction [21,22]. Valencia-Flores et al. [20] compared the effect on bacterial growth of HPH at 200-300 MPa and soft temperature inlet (55-75 °C) with conventional pasteurization treatment (90 °C, 90 s) in almond beverage. They showed that 200 MPa and an inlet temperature of 55 °C of inlet temperature was enough to improve conventional pasteurization effect on microbiological quality.

Moreover, beer is another beverage susceptible to be treated by HPH. Some research works established it was possible to completely inactivate the microorganisms in addition to improving the colour of the beer by HPH at pressures between 200-300 MPa and with 1 to 3 passes. The addition of antimicrobials as lysozyme enzyme (50 mg/L) had a synergistic effect, reducing the pressure needed to 100-150 MPa. However, HPH treatment could result in greater values for turbidity and it would be necessary to perform another stabilization treatments to minimize the negative effects [23,24].

HPH treatment has also been used to modulate the activity of various enzymes. This treatment can increase or decrease the enzyme activity depending on processing conditions (pressure and number of passes), homogenizing valve structure, specific enzyme, pH, temperature and food matrix.

Since enzymes are a complex type of globular protein, mechanical forces and cavitation effects associated to HPH treatment result in conformational and structural changes modifying enzyme activity and stability. The main modifications in the enzyme are linked to changes in the quaternary, tertiary and even secondary structure. Formation or interruption of hydrogen bonds, Van der Waals, hydrophobic and electrostatic interactions can occur increasing the number of hydrophobic sites, revealing amino acid and sulfhydryl groups and thus, accelerating, delaying or impeding enzyme-substrate interaction[24]. Furthermore, the magnitude of the changes induced by the HPH treatment will determine their reversibility or irreversibility. Aguilar et al. [28] mentioned that protein denaturation can be reversible at 100 MPa and irreversible above 200 MPa.

In the case of juices, the main alteration reactions are caused by polyphenoloxidase responsible for browning and oxidation reactions. It was possible to inactivate it with homogenization pressures of 80-150 MPa [29,30]. On the other hand, α -amylase, whose use in recent years has been increasing since it reduces the starch content of beverages to avoid turbidity and gelatinization, is resistant to HPH [31]. A similar resistance has been observed on *Pseudomonas fluorescens* protease when HPH at 100-150 MPa was applied to reduce its proteolytic rate [32].

Table 2. Research works evaluating enzyme activity modulation by HPH.

PRODUCT	ENZYMES	TREATMENT	EFFECT	REFERENCE
Commercial enzymes	Glucose oxidase	50, 100, 150 MPa	Decrease in enzyme activity at 50 MPa. Improvement in activity and stability at 100 and 150 MPa	[33]
Commercial enzymes	Amyloglucosidase, Glucose oxidase, Neutral protease	Amyloglucosidase, neutral protease: 150, 200 MPa (3 passes). Glucose oxidase: 100, 150 MPa (3 passes)	Improvement of enzymatic activity	[24]
Fruit juices	α -amilase	0, 40, 80, 120 and 150 MPa	Stability of the enzyme	[31]
Apple juice	Polyphenoloxidase	150 MPa (10 passes)	Inactivation	[29]
Lettuce waste juice	Polyphenoloxidase	80 MPa (1 pass) and 150 MPa (1-10 passes)	Inactivation	[30]
Peanut protein	Alcalase	0, 1, 40 and 80 MPa	Increased enzymatic hydrolysis	[34]
Chicken egg white	Lysozyme muramidase	40, 80, 120, 160 and 190 MPa	Activation and increase of enzymatic activity	[35]

Raw skim milk	Alkaline phosphatase and lactoperoxidase	100, 150, 200, 250 and 300 MPa	Decrease and inactivation of alkaline phosphatase. Increased activity of lactoperoxidase.	[17]
Milk	Protease <i>Pseudomonas fluorescens</i>	100 and 150 MPa	Decreased proteolytic rate	[32]

On the other hand, HPH can also be applied to enhance enzyme activity. Some authors have applied HPH to increase the activity of enzymes involved in the shelf life or processing of several food matrices. Lysozyme and lactoferrin have increased their antimicrobial activity against *L. monocytogenes* by HPH at 100 MPa in milk [8,36]. Pinho et al. [17] obtained a rise on enzymatic activity of lactoperoxidase in skim milk at pressures between 100 and 250 MPa. In contrast, if the homogenization pressure increased up to 300 MPa a reduction of around 30% in enzyme activity was detected. In another work, defatted peanut flour was dispersed in distilled water and pH adjusted, and further subjected to HPH treatment at 40 and 80 MPa. After that, the peanut protein was recovered from the dispersed solution by an acid precipitation and re-dispersed in distilled water. The HPH treatment increased the extraction yield and the hydrolysis of the peanut protein isolates by endogenous enzymes. DPPH radical scavenging and hydroxyl radical scavenging activities were also increased [34].

It is evidenced that costs, versatility and performance improvement of enzymatic processes can be achieved when activity of commercial enzymes is increased by HPH. In particular, Tribst et al. [24] improved the activity of amyloglucosidase, glucose oxidase and neutral protease, at HPH between 100-150 MPa and non- optimum temperatures. Commercial enzymes derived from fungi and available as powders were diluted in acetate buffer solutions and then subjected to HPH treatment. They observed an uneven effect of the number of passes. Only one pass was required to increase activity of amyloglucosidase and neutral protease while no effect was observed in subsequent passes, successive steps continued to increase the enzyme activity of glucose oxidase. The energy involved in the molecular changes associated to the increase in enzyme activity might be responsible for it.

3. Extraction and technological functionality improvement of proteins, fibrous materials and bioactive compounds

HPH is being used in the last years to contribute to food process sustainability [5]. In this area, HPH has been applied for valorisation of agrifood by-products with two different objectives: to increase extractability of intracellular or cell wall structural components and to improve technologic functionality of biomolecules from food by-products. Most agri-food wastes or by-products are rich in fibrous material and, in some cases, in proteins or bioactive compounds of interest for the food industry to be used as food ingredients [37], or either to be used as environmentally safe and sustainable packaging materials [38]. HPH induces cells disruption favouring the release of structural and intracellular content and consequently its valorisation.

The main kind of products in which HPH is applied to extract fibres, proteins or bioactive compounds are solid by-products such as pomace from fruit or vegetable juicing, fruits or vegetables peels, minimal processing waste and vegetal parts of plants or cereal seed hulls. In this cases, the solid wastes need to be fluidized by diluting them in water or another solvent. In other cases, an extraction method is applied and the extracted phase further subjected to HPH. [39] extracted pectin from potato peel by HPH at 200 MPa. They obtained improvements in the viscosity, emulsifying properties, degree of esterification and physicochemical characteristics. Therefore they recommend the use of HPH to obtain pectin that could be used as an ingredient acting as a stabilizing agent or a thickener in food manufacturing. Similarly, Fayaz et al. [40] showed that HPH favours the release of okara proteins and soluble fibre. Soy okara was dispersed in deionized water at 10 g/100 g

concentration and pre-homogenized with a high-speed blender. After that, an homogenization pressure of 150 MPa for 5 passes made it possible to extract proteins with a yield of 90%. [41] applied HPH to make edible and biodegradable films for food packaging from a type of edible fungus of the variety *Flammulina velutipes*. Wu et al. [42] demonstrated the possibility of using HPH treatment to make biodegradable biopolymer films from pomelo peel.

Otherwise, HPH reduces particle size and structure of macromolecules modifying solubility, interaction properties, viscosity, or other physic-chemical properties. Saricaoglu et al. [43] improved the functionality of proteins from hazelnut industry by HPH at 100 MPa and 1 pass. The homogenization pressure decreased particle size of proteins, increasing zeta potential and water solubility. Emulsifying and sparkling properties were improved too. Hua et al. [44] demonstrated a microstructural change of tomato waste fibres by applying HPH at 100 MPa and 10 passes. The authors transformed around 8% of the insoluble fibres into soluble ones. [45] indicated that for the preparation of soluble peach fibre from fresh peach marc it must be dispersed in 3 times the volume of deionized water, thus improving the efficiency of cellulose hydrolysis. For pectin extraction from milled dried lemon peel, variations in dilutions change the properties of the pectin extracted, resulting in residues with different pectic characteristics [46]. Discarded external lettuce leaves were dispersed in hydroalcoholic solutions and polyphenols extracted with ethanol to get good phenolic extraction yields [47].

Table 3. Research works aimed to the extraction and improvement of technological functionality of proteins, fibres or bioactive compounds from agri-food wastes by HPH.

SUSTRATE	COMPONENT	TREATMENT	OBJECTIVE	REFERENCE
Sweet potato leaves	Flavonoids	100 MPa (2 passes)	Strengthens the antioxidant activities of the flavonoid	[48]
Potato peel	Biopolymer film	150 MPa	Extraction	[49]
Peach pomace	Soluble fibres	140 MPa (4 passes)	Significantly improved the efficiency of cellulase hydrolysis in the preparation of soluble fibres and a high binding capacity for sodium cholate and cholesterol.	[45]
Potato peel	Phenolic acids	159 MPa (2 passes) + NaOH treatment	Improved extraction and release of total phenolic content and total flavonoid content	[37]
Desmodosmus sp. F51	Carotenoids	69 - 276 MPa (1-4 passes)	Extraction	[50]
Dry tomato residue waste	Fibres	100 MPa (10 passes)	Improved the soluble fibre content and its oil holding capacity	[44]
Citrus peel	Fibres	90, 160 MPa (2 passes)	Improvement of physical, chemical and functional properties including surface area, water holding capacity, texture and viscosity	[51]
Lemon peels fibre	Pectin	20 and 80 MPa	Extraction	[46]

Soybean	Protein	100 MPa	Extraction	[52]
Hazelnut oil industry by-products	Hazelnut meal proteins	0, 25, 50, 75, 100 and 150 MPa	Improves functional (solubility, emulsifying and foaming properties) and rheological properties of proteins	[53]
Black cherry tomato waste	Pectin	0, 40, 80, 120 and 160 MPa (2 passes)	Increase the esterification degree of pectins	[54]
Carrot processing waste	Biodegradable composite films were prepared	138 MPa (7 passes)	Extraction	[55]
Lettuce waste	Polyphenols	50, 100 MPa	Extraction	[47]
Potato peel	Pectin	200 MPa	Increased galacturonic acid content, viscosity and emulsifying properties. Decreased esterification degree and molecular weight	[39]
Broccoli seeds	Sulforaphane	20-160MPa (1-5 passes)	Increases the extraction yield	[56]
Agri-food waste (tomato peel, coffee beans)	Application for structuring peanut oil	70 MPa (3 passes)	Replacing part of the lipids with water and low calorie fibres.	[57]
Edible mushroom by-products	Biodegradable edible film	100 MPa (3 passes)	Improve tensile strength, elongation at break, water vapor permeability, oxygen barrier and thermal stability.	[41]
Grape seeds, tomato stem, walnut shells, coffee	Polyphenolic compounds and antioxidants	20, 50, 100, 120 MPa	Extraction	[58]
Soybean okara	Proteins and soluble fibers	50, 100, 150 MPa (1 pass) 150MPa (5 passes)	Extraction	[40]
Sugar palm	nanofibrillated cellulose	50 MPa (3 passes)	Extraction	[59]
Tomato peels	Bioactive compounds: proteins, polyphenols, lycopene	100 MPa (1-10 passes)	Increased release of intracellular compounds (proteins, sugars, antioxidants)	[60]

Pomelo peel	Biopolymer film	20, 40, 60 and 80 MPa (10 passes)	Improve mechanical properties, microstructure, optical and barrier properties	[42]
Soybean meal	Resins	20 MPa	Extraction	[61]

4. Increase of bioavailability and encapsulation of bioactive compounds

In the last decade, many studies have been carried out to demonstrate that the application of HPH in liquid foods can produce a modification in the bioaccessibility (i.e. fraction of an ingested nutrient that is released from the food matrix and made available for intestinal absorption) or bioavailability (i.e. fraction of an ingested nutrient that is absorbed in the intestine and incorporated into the bloodstream) of its bioactive compounds. In most studies, an increase in the bioaccessibility of phytochemicals is observed due to their release within the structure of the food in which they are found. In other cases a modification of its biological functionality occurs due to some change in its chemical structure. [62] carried out an interesting review that demonstrates these effects in three bioactive components: carotenoids, phenolic compounds and vitamin C. The review shows that fruit juices (carrot, tomato, orange, apple and berries) are the most common food in which HPH increase bioaccessibility of bioactive compounds. HPH decreases the particle size of suspended pulp, increases cloud stability and thus the availability of bioactive components. Treatment with HPH in mandarin juices increased the bioaccessibility of total carotenoids by five times, although in the case of flavonoids no such drastic changes were observed. Therefore, the HPH treatment was recommended for the production of tangerine juices that promote health, mainly through the improvement of the bioaccessibility of its carotenoids [63]. Quan et al. [64] established that bioaccessibility improvement could be conditioned by food matrix. They observed that HPH at 250 MPa favoured the release from cell walls and increased the content of total phenolic compounds in kiwi and pomelo juices, but it had a negative effect on its bioaccessibility (in vitro) as a consequence of a major degradation along the digestion process. Conversely, the addition of skimmed or whole milk to the juices had no significant effect on total phenolics content but increased phenolic bioaccessibility in kiwi juice and pomelo juice from 21.6% to 37.8% and 60.1% to 63.3%, respectively. Similarly, Alongi et al. [65] showed that chlorogenic acids bioaccessibility (in vitro) increased from nearly 25% to more than 50% by adding milk with different fat content to coffee and applying HPH (50-150 MPa). They observed the pressure required was lower the lower the fat percentage and they attributed the effect to chlorogenic acids micellarization, a phenomenon that reduced their susceptibility to degradation during digestion. Sometimes, positive effect of HPH could be observed after storage. Betoret et al. [66] found in low pulp mandarin juice that, despite the suspended pulp increased with HPH and trehalose addition, flavonoid hesperidin initially decreased but resulted in less flavonoid degradation during storage.

HPHs have also been applied, with a much smaller number of published articles, to non-dairy vegetable based beverages. Although in some cases no significant improvements in the nutritional characteristics were detected, in others, such as almond or soy beverages, a reduction in antinutrients was achieved [67]. Denaturation, aggregation and chemical modification of proteins may change their allergic potential. Toro- Funes et al. [68] demonstrated an increase of 40% in the extractability of phytosterols from almond milk subjected to a HPH of 300 MPa (6 passes). However, the content of tocopherol and polyamines such as spermidine were reduced by up to 90%. The application of HPH for kefir production from hazelnut beverage achieved improvements in the total content of phenolic compounds and antioxidant capacity, causing a reduction in the content of lactic and citric acid [69].

Improvement in the bioavailability of bioactive components by HPH is also possible in solid foods. HPH (10-20 cycles and 100- 200 MPa) was used to fabricate fermented soybean powder into an aqueous nanosuspension, favouring in-vitro release of isoflavones from nanosuspension [70].

The other way in which HPH can contribute to an improvement in the nutritional properties of foods is by using this technology for the encapsulation of bioactive components. In this way it is

possible to increase its stability, conservation and controlled delivery in the target site increasing food functionality. HPH produces intense disruptive forces that break up particles into smaller sizes favouring encapsulation of specific components in a suitable media. Mechanical stress, heating associated and emulsifier interactions can affect the effectiveness of the process and therefore the activity and bioaccessibility of the bioactive compound.

Many studies have investigated nanoencapsulation of curcumin by HPH at 40-100 MPa. Results showed that the emulsifier type had an influence in curcumin bioaccessibility [71]. Frank et al. [72] studied the degradation of anthocyanins from bilberry extract by subjecting them to temperature and mechanical stress similar to the process of emulsification and encapsulation by HPH. The HPH were applied with a simple pass and in the pressure range between 30 and 150 MPa. Thereafter the samples were immediately cooled to 298K. The results showed no significant influence of mechanical stresses associated to HPH on anthocyanin stability, even at high-pressure treatment up to 150 MPa. The combination temperature–time was the main parameter affecting anthocyanin degradation.

A great interest has been shown in applying HPH as encapsulation technique in bacteria with probiotic effect. Patrignani et al. [73] underlined the applicative potential of HPH microencapsulation of probiotic microorganisms to produce fermented milk with improved functionality and with enhanced sensory properties. They established 50 MPa and 5 passes as adequate conditions to produce stable microcapsules of *Lactobacilli* with high yield and viability during storage. Moreover, microencapsulation of adjunct bacteria reduced acidity of fermented milk. Calabuig-Jiménez et al. [74] microencapsulated *L. salivarius* spp. *salivarius* in alginate coatings by HPH at 70 MPa. A positive effect of microcapsules was observed when evaluating the survival of the probiotic strain on simulated gastrointestinal conditions.

Table 4. Research works in which HPH treatment was applied to encapsulate.

COMPONENT ENCAPSULATED	MATRIX	CONDITIONS	RESULTS	REFERENCE
<i>Lactobacillus paracasei</i> A13 and <i>Lactobacillus salivarius</i> subsp. <i>salivarius</i> CET 4063	Fermented milk	50 MPa (5 passes)	The microcapsules presented high yields in terms of trapped viable cells and acceptable sizes. Furthermore, microencapsulation caused the decrease in acidity in fermented milk.	[73]
Phenolic compounds and anthocyanins from blueberry pomace	-	50- 200 MPa	The encapsulation efficiency, size and charge characteristics of the emulsion droplets were affected by HPH.	[75]
<i>Lactobacillus salivarius</i> spp. <i>Salivarius</i>	Mandarin Juice	70 MPa (2 passes)	Improving the survival of probiotics with the use of alginate as a coating	[74]
Phenolic powder from strawberry pomace	-	50 and 70 MPa (3, 5, 7 passes)	High encapsulation efficiency	[76]
<i>L. salivarius</i> spp. <i>Salivarius</i>	Mandarin juice impregnated in apple	70 MPa (2 passes)	The final count of <i>L. salivarius</i> spp. <i>Salivarius</i> encapsulated was high enough to exert a potential beneficial effect	[77]

HPH treatments at pressure levels below 100 MPa, considered as sublethal pressures, are applied to microbial cultures: initiators, co-initiators, or probiotics and yeasts, in order to produce cultures with improved functional, technological and sensory properties. The use of strains belonging to the genus *Bifidobacterium* and *Lactobacillus* predominate as probiotics, and to a lesser extent *Enterococcus*, *Streptococcus* and *Saccharomyces* [78]. The bacterial cells response to mechanical stress induced by HPH modifying their metabolic activity and membrane composition. Therefore technological and functional properties such as fermentation kinetic, enzymatic activities, hydrophobicity or resistance to gastrointestinal digestion can be improved.

Table 5. Research works in which HPH treatment was applied to probiotic cells.

FOOD MATRIX	MICROBIAL STRAIN	CONDITIONS	RESULTS	REFERENCES
Yogurt	<i>L. Delbrueckii</i> ssp. <i>bulgaricus</i> LB-12, <i>S. Salivarius</i> ssp. <i>thermophilus</i> ST-M5 and <i>L. acidophilus</i> LA-K	0, 3.45, 6.90, 10.34 and 13.80 MPa	Improved tolerance to acid and bile	[79]
-	<i>L. acidophilus</i> Dru y <i>L. paracasei</i> A13	0.1 and 50 MPa	Increased probiotic characteristics in vivo, there was no modification in the interaction of lactobacilli with the small intestine	[80]
-	<i>Lactobacillus paracasei</i> A13, <i>Lactobacillus acidophilus</i> 08 and Dru, <i>Lactobacillus delbrueckii</i> spp. <i>lactis</i> 200	50 MPa	Increased functional characteristics depending on the type of strain	[12]
Fermented milks	<i>Lactobacillus rhamnosus</i> BFE5264, <i>L. delbrueckii</i> spp. <i>bulgaricus</i> FP1 and <i>Streptococcus thermophilus</i> LI3	60 MPa	Reduced product clotting time and increased viability of the probiotic strain	[81]
Cacciotta cheese	<i>Lactobacillus paracasei</i> A13	50 MPa	Increase in quality and decrease in cheese maturation time.	[82]
Mandarin juice	<i>L. salivarius</i> spp. <i>Salivarius</i>	0, 20 and 100 MPa	Improvement of cellular hydrophobicity	[66]
Clementine juice	<i>L. salivarius</i> spp. <i>Salivarius</i>	25, 50, 100 and 150 MPa	Improvement of the antioxidant properties of the juice.	[83]

Fresh Culture (1% v/v)	<i>Lactobacillus paracasei</i> A13	50, 150, 200 MPa	Increase the unsaturation in membrane fatty acids.	[84]
------------------------	------------------------------------	------------------	--	------

Siroli et al. [84] reported that the main regulatory mechanism that probiotic lactobacilli adopt to counteract pressure stress is the modification of the composition of membrane fatty acids. Specifically, they observed an increase of unsaturated fatty acids when HPH at 100 and 150 MPa was applied to *Lactobacillus paracasei* A13. Considering that the increase of unsaturation level is a key mechanism to compensate for the oxidative damages induced by physico-chemical stressors in microbial cells, they concluded that HPH at sublethal pressures is useful to improve activity of some *Lactobacillus* species.

Lanciotti et al. [85] studied the effect of HPH between 50 MPa and 100 MPa on fermentation kinetics, metabolic profile and enzymatic activity of four species of Lactobacilli involved in dairy product fermentation and ripening. Although the results varied according to the species, they documented no significant effect on cell viability, an increased proteolytic activity and positive changes in fermentation dynamics. The resistance to simulated gastric conditions, hydrophobicity and auto-aggregation capacity resulted also strain-dependent for *L. acidophilus* Dru and *L. paracasei* A13 when subjected to HPH at 50 MPa. HPH increased the three properties for *L. paracasei* A13 while reduced them for *L. acidophilus* Dru. Authors attribute the differences to the compositional and structural differences in the cellular outer structures, thus suggesting the HPH effects on macromolecules and their interactions with the gut immune cells have a key role in the probiotic effect. The same authors noted that HPH treated *L. paracasei* cells modified their interaction with the small intestine of mice, inducing a higher IgA response compared to untreated *L. paracasei* cells [12,80] Betoret et al. [66] demonstrated the improvement in the hydrophobicity of *Lactobacillus salivarius* spp. *salivarius* added to mandarin juice with trehalose when HPH at 0, 20 and 100 MPa were applied.

5. Conclusions

Although in the beginning the application of high homogenizing pressures was aimed at homogenizing and increasing the stability of emulsions such as milk, the advances in valve design and equipment allowed for an increase in working pressure extending the scope of application.

In the last decade, it has grown by more than 80% the number of research works related to the implementation of HPH in the process of extracting bioactive components from agri-food wastes, to improve bioavailability and probiotic properties of bioactive components and microorganisms and also as an encapsulation technique. At the same time, progress has been made in the application of HPH to reduce the microbial load or modulate the activity of some enzymes.

The general mechanisms responsible for the effect of HPH are known, but the final effect is largely conditioned by the type of valve, pressure applied, number of passes, nature of the components and macromolecules, and food matrix. For this reason research is needed for each specific application.

Results published in the last decade have shown HPH as a non-thermal technology suitable to accomplish the food industry's objectives of quality and safety, functionality and sustainability.

Author Contributions: Conceptualization, N.B. and E.B.; methodology and search, J.M., L.H. and N.B.; writing—original draft preparation, J.M. and N.B.; writing—review and editing, E.B., L.S. and C.B.; funding acquisition, N.B. All authors have read and agreed to the published version of the manuscript.

Funding: Please add: This research and APC were funded by Generalitat Valenciana, Project AICO/2017/049-

Acknowledgments: The authors thank the research project “Fortalecimiento de los Encadenamientos Productivos de las Subregiones del Chocó” BPIN 2013000100284 Tecnológica del Chocó (in Spanish) by financial support to Leidy Indira Hinestroza-Córdoba.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Augusto, P.E.D.; Tribst, A.A.L.; Cristianini, M. High Hydrostatic Pressure and High-Pressure Homogenization Processing of Fruit Juices. In *Fruit Juices: Extraction, Composition, Quality and Analysis*; Elsevier Inc., 2018; pp. 393–421 ISBN 9780128024911.
2. Osorio-Arias, J.C.; Vega-Castro, O.; Martínez-Monteaquedo, S.I. Fundamentals of High-Pressure Homogenization of Foods. In *Reference Module in Food Science*; Elsevier, 2020.
3. Barba, F.J.; Esteve, M.J.; Frígola, A. High Pressure Treatment Effect on Physicochemical and Nutritional Properties of Fluid Foods During Storage: A Review. *Compr. Rev. Food Sci. Food Saf.* **2012**, *11*, 307–322, doi:10.1111/j.1541-4337.2012.00185.x.
4. Bevilacqua, A.; Campaniello, D.; Speranza, B.; Altieri, C.; Sinigaglia, M.; Corbo, M.R. Two nonthermal technologies for food safety and quality—ultrasound and high pressure homogenization: Effects on microorganisms, advances, and possibilities: A review. *J. Food Prot.* **2019**, *82*, 2049–2064.
5. Picart-Palmade, L.; Cunault, C.; Chevalier-Lucia, D.; Belleville, M.P.; Marchesseau, S. Potentialities and limits of some non-thermal technologies to improve sustainability of food processing. *Front. Nutr.* **2019**, *5*, 130, doi:10.3389/fnut.2018.00130.
6. McClatchie, J.M. The Borden Company. Valve For homogenizing devices 1930, 28, 131–134.
7. Pandolfe, W.D. Development of the New Gaulin Micro-Gap™ Homogenizing Valve. *J. Dairy Sci.* **1982**, *65*, 2035–2044, doi:10.3168/jds.S0022-0302(82)82456-9.
8. Patrignani, F.; Siroli, L.; Braschi, G.; Lanciotti, R. Combined use of natural antimicrobial based nanoemulsions and ultra high pressure homogenization to increase safety and shelf-life of apple juice. *Food Control* **2020**, *111*, 107051, doi:10.1016/j.foodcont.2019.107051.
9. Balasubramaniam, V.M.; Barbosa-Cánovas, G.; Lelieveld, H. *High Pressure Processing of Food-Principles, Technology and Application*; 2016; ISBN 978-1-4939-3233-7.
10. Tribst, A.A.L.; Franchi, M.A.; de Massaguer, P.R.; Cristianini, M. Quality of Mango Nectar Processed by High-Pressure Homogenization with Optimized Heat Treatment. *J. Food Sci.* **2011**, *76*, 106–110, doi:10.1111/j.1750-3841.2010.02006.x.
11. Calligaris, S.; Foschia, M.; Bartolomeoli, I.; Maifreni, M.; Manzocco, L. Study on the applicability of high-pressure homogenization for the production of banana juices. *LWT - Food Sci. Technol.* **2012**, *45*, 117–121, doi:10.1016/j.lwt.2011.07.026.
12. Tabanelli, G.; Patrignani, F.; Vinderola, G.; Reinheimer, J.A.; Gardini, F.; Lanciotti, R. Effect of sub-lethal high pressure homogenization treatments on the invitro functional and biological properties of lactic acid bacteria. *LWT - Food Sci. Technol.* **2013**, *53*, 580–586, doi:10.1016/j.lwt.2013.03.013.
13. Guan, Y.; Zhou, L.; Bi, J.; Yi, J.; Liu, X.; Chen, Q.; Wu, X.; Zhou, M. Change of microbial and quality attributes of mango juice treated by high pressure homogenization combined with moderate inlet temperatures during storage. *Innov. Food Sci. Emerg. Technol.* **2016**, *36*, 320–329, doi:10.1016/j.ifset.2016.07.009.

14. Yu, Y.; Wu, J.; Xu, Y.; Xiao, G.; Zou, B. Effect of High Pressure Homogenization and Dimethyl Dicarboxylate (DMDC) on Microbial and Physicochemical Qualities of Mulberry Juice. *J. Food Sci.* **2016**, *81*, M702–M708, doi:10.1111/1750-3841.13213.
15. Xia, X.; Dai, Y.; Wu, H.; Liu, X.; Wang, Y.; Cao, J.; Zhou, J. Effects of pressure and multiple passes on the physicochemical and microbial characteristics of lupin-based beverage treated with high-pressure homogenization. *J. Food Process. Preserv.* **2019**, *43*, 1–12, doi:10.1111/jfpp.13912.
16. Benjamin, O.; Gamrasni, D. Microbial, nutritional, and organoleptic quality of pomegranate juice following high-pressure homogenization and low-temperature pasteurization. *J. Food Sci.* **2020**, *85*, 592–599, doi:10.1111/1750-3841.15032.
17. Pinho, C.R.G.; Franchi, M.A.; Tribst, A.A.L.; Cristianini, M. Effect of Ultra High Pressure Homogenization on Alkaline Phosphatase and Lactoperoxidase Activity in Raw Skim Milk. *Procedia Food Sci.* **2011**, *1*, 874–878, doi:10.1016/j.profoo.2011.09.132.
18. Amador Espejo, G.G.; Hernández-Herrero, M.M.; Juan, B.; Trujillo, A.J. Inactivation of *Bacillus* spores inoculated in milk by Ultra High Pressure Homogenization. *Food Microbiol.* **2014**, *44*, 204–210, doi:10.1016/j.fm.2014.06.010.
19. Mercan, E.; Sert, D.; Akin, N. Determination of powder flow properties of skim milk powder produced from high-pressure homogenization treated milk concentrates during storage. *LWT* **2018**, *97*, 279–288, doi:10.1016/j.lwt.2018.07.002.
20. Valencia-Flores, D.C.; Hernández-Herrero, M.; Guamis, B.; Ferragut, V. Comparing the Effects of Ultra-High-Pressure Homogenization and Conventional Thermal Treatments on the Microbiological, Physical, and Chemical Quality of Almond Beverages. *J. Food Sci.* **2013**, *78*, 199–205, doi:10.1111/1750-3841.12029.
21. Bevilacqua, A.; Casanova, F.P.; Petrucci, L.; Sinigaglia, M.; Corbo, M.R. Using physical approaches for the attenuation of lactic acid bacteria in an organic rice beverage. *Food Microbiol.* **2016**, *53*, 1–8, doi:10.1016/j.fm.2015.08.005.
22. Codina-Torrella, I.; Guamis, B.; Zamora, A.; Quevedo, J.M.; Trujillo, A.J. Microbiological stabilization of tiger nuts' milk beverage using ultra-high pressure homogenization. A preliminary study on microbial shelf-life extension. *Food Microbiol.* **2018**, *69*, 143–150, doi:10.1016/j.fm.2017.08.002.
23. Franchi, M.A.; Tribst, A.A.L.; Cristianini, M. Inactivation of *Lactobacillus brevis* in Beer Utilizing a Combination of High-Pressure Homogenization and Lysozyme Treatment. *J. Inst. Brew.* **2011**, *117*, 634–638, doi:10.1002/j.2050-0416.2011.tb00515.x.
24. Franchi, M.A.; Tribst, A.A.L.; Cristianini, M. High-pressure homogenization: A non-thermal process applied for inactivation of spoilage microorganisms in beer. *J. Inst. Brew.* **2013**, *119*, 237–241, doi:10.1002/jib.99.
25. Comuzzo, P.; Calligaris, S.; Iacumin, L.; Ginaldi, F.; Palacios Paz, A.E.; Zironi, R. Potential of high pressure homogenization to induce autolysis of wine yeasts. *Food Chem.* **2015**, *185*, 340–348, doi:10.1016/j.foodchem.2015.03.129.

26. Capra, M.L.; Patrignani, F.; Quiberoni, A. del L.; Reinheimer, J.A.; Lanciotti, R.; Guerzoni, M.E. Effect of high pressure homogenization on lactic acid bacteria phages and probiotic bacteria phages. *Int. Dairy J.* **2009**, *19*, 336–341, doi:10.1016/j.idairyj.2008.11.002.
27. Patrignani, F.; Vannini, L.; Kamdem, S.L.S.; Lanciotti, R.; Guerzoni, M.E. Potentialities of High-Pressure Homogenization to Inactivate *Zygosaccharomyces bailii* in Fruit Juices. *J. Food Sci.* **2010**, *75*, 116–120, doi:10.1111/j.1750-3841.2009.01508.x.
28. Dos Santos Aguilar, J.G.; Cristianini, M.; Sato, H.H. Modification of enzymes by use of high-pressure homogenization. *Food Res. Int.* **2018**, *109*, 120–125, doi:10.1016/j.foodres.2018.04.011.
29. Bot, F.; Calligaris, S.; Cortella, G.; Plazzotta, S.; Nocera, F.; Anese, M. Study on high pressure homogenization and high power ultrasound effectiveness in inhibiting polyphenoloxidase activity in apple juice. *J. Food Eng.* **2018**, *221*, 70–76, doi:10.1016/j.jfoodeng.2017.10.009.
30. Plazzotta, S.; Manzocco, L. High-pressure homogenisation combined with blanching to turn lettuce waste into a physically stable juice. *Innov. Food Sci. Emerg. Technol.* **2019**, *52*, 136–144, doi:10.1016/j.ifset.2018.11.008.
31. Tribst, A.A.L.; Cristianini, M. High pressure homogenization of a fungi α -amylase. *Innov. Food Sci. Emerg. Technol.* **2012b**, *13*, 107–111, doi:10.1016/j.ifset.2011.10.006.
32. Oliveira, M.M. de; Leite Júnior, B.R. de C.; Tribst, A.A.L.; Cristianini, M. Use of high pressure homogenization to reduce milk proteolysis caused by *Pseudomonas fluorescens* protease. *LWT - Food Sci. Technol.* **2018**, *92*, 272–275, doi:10.1016/j.lwt.2018.02.052.
33. Tribst, A.A.L.; Cristianini, M. Changes in commercial glucose oxidase activity by high pressure homogenization. *Innov. Food Sci. Emerg. Technol.* **2012a**, *16*, 355–360, doi:10.1016/j.ifset.2012.08.002.
34. Dong, X.; Zhao, M.; Shi, J.; Yang, B.; Li, J.; Luo, D.; Jiang, G.; Jiang, Y. Effects of combined high-pressure homogenization and enzymatic treatment on extraction yield, hydrolysis and function properties of peanut proteins. *Innov. Food Sci. Emerg. Technol.* **2011**, *12*, 478–483, doi:10.1016/j.ifset.2011.07.002.
35. Tribst, A.A.L.; Ribeiro, L.R.; Cristianini, M. Comparison of the effects of high pressure homogenization and high pressure processing on the enzyme activity and antimicrobial profile of lysozyme. *Innov. Food Sci. Emerg. Technol.* **2017**, *43*, 60–67, doi:10.1016/j.ifset.2017.07.026.
36. Iucci, L.; Patrignani, F.; Vallicelli, M.; Guerzoni, M.E.; Lanciotti, R. Effects of high pressure homogenization on the activity of lysozyme and lactoferrin against *Listeria monocytogenes*. *Food Control* **2007**, *18*, 558–565, doi:10.1016/j.foodcont.2006.01.005.
37. Zhu, X.; Cheng, Y.; Chen, P.; Peng, P.; Liu, S.; Li, D.; Ruan, R. Effect of alkaline and high-pressure homogenization on the extraction of phenolic acids from potato peels. *Innov. Food Sci. Emerg. Technol.* **2016**, *37*, 91–97, doi:10.1016/j.ifset.2016.08.006.
38. Flôres, S.H.; Rios, A. de O.; Iahnke, A.O.S.; de Campo, C.; Vargas, C.G.; Santos, C.D.M.; Caetano, K. dos S.; Stoll, L.; Crizel, T. de M. Films for Food From Ingredient Waste. In *Reference Module in Food Science*; Elsevier, 2017.

39. Xie, F.; Zhang, W.; Lan, X.; Gong, S.; Wu, J.; Wang, Z. Effects of high hydrostatic pressure and high pressure homogenization processing on characteristics of potato peel waste pectin. *Carbohydr. Polym.* **2018**, *196*, 474–482, doi:10.1016/j.carbpol.2018.05.061.
40. Fayaz, G.; Plazzotta, S.; Calligaris, S.; Manzocco, L.; Nicoli, M.C. Impact of high pressure homogenization on physical properties, extraction yield and biopolymer structure of soybean okara. *LWT* **2019**, *113*, 108324, doi:10.1016/j.lwt.2019.108324.
41. Wang, W.; Zhang, K.; Li, C.; Cheng, S.; Zhou, J.; Wu, Z. A novel biodegradable film from edible mushroom (*F. velutipes*) by product: Microstructure, mechanical and barrier properties associated with the fiber morphology. *Innov. Food Sci. Emerg. Technol.* **2018**, *47*, 153–160, doi:10.1016/j.ifset.2018.02.004.
42. Wu, H.; Xiao, D.; Lu, J.; Jiao, C.; Li, S.; Lei, Y.; Liu, D.; Wang, J.; Zhang, Z.; Liu, Y.; et al. Effect of high-pressure homogenization on microstructure and properties of pomelo peel flour film-forming dispersions and their resultant films. *Food Hydrocoll.* **2020**, *102*, 105628, doi:10.1016/j.foodhyd.2019.105628.
43. Saricaoglu, F.T.; Atalar, I.; Yilmaz, V.A.; Odabas, H.I.; Gul, O. Application of multi pass high pressure homogenization to improve stability, physical and bioactive properties of rosehip (*Rosa canina* L.) nectar. *Food Chem.* **2019**, *282*, 67–75, doi:10.1016/j.foodchem.2019.01.002.
44. Hua, X.; Xu, S.; Wang, M.; Chen, Y.; Yang, H.; Yang, R. Effects of high-speed homogenization and high-pressure homogenization on structure of tomato residue fibers. *Food Chem.* **2017**, *232*, 443–449, doi:10.1016/j.foodchem.2017.04.003.
45. Xu, H.; Jiao, Q.; Yuan, F.; Gao, Y. In vitro binding capacities and physicochemical properties of soluble fiber prepared by microfluidization pretreatment and cellulase hydrolysis of peach pomace. *LWT - Food Sci. Technol.* **2015**, *63*, 677–684, doi:10.1016/j.lwt.2015.03.033.
46. Willemsen, K.L.D.D.; Panozzo, A.; Moelants, K.; Debon, S.J.J.; Desmet, C.; Cardinaels, R.; Moldenaers, P.; Wallecan, J.; Hendrickx, M.E.G. Physico-chemical and viscoelastic properties of high pressure homogenized lemon peel fiber fraction suspensions obtained after sequential pectin extraction. *Food Hydrocoll.* **2017**, *72*, 358–371, doi:10.1016/j.foodhyd.2017.06.020.
47. Plazzotta, S.; Manzocco, L. Effect of ultrasounds and high pressure homogenization on the extraction of antioxidant polyphenols from lettuce waste. *Innov. Food Sci. Emerg. Technol.* **2018**, *50*, 11–19, doi:10.1016/j.ifset.2018.10.004.
48. Huang, X.; Tu, Z.; Xiao, H.; Li, Z.; Zhang, Q.; Wang, H.; Hu, Y.; Zhang, L. Dynamic high pressure microfluidization-assisted extraction and antioxidant activities of sweet potato (*Ipomoea batatas* L.) leaves flavonoid. *Food Bioprod. Process.* **2013**, *91*, 1–6, doi:10.1016/j.fbp.2012.07.006.
49. Rommi, K.; Rahikainen, J.; Vartiainen, J.; Holopainen, U.; Lahtinen, P.; Honkapää, K.; Lantto, R. Potato peeling costreams as raw materials for biopolymer film preparation. *J. Appl. Polym. Sci.* **2016**, *133*, doi:10.1002/app.42862.
50. Xie, Y.; Ho, S.H.; Chen, C.N.N.; Chen, C.Y.; Jing, K.; Ng, I.S.; Chen, J.; Chang, J.S.; Lu, Y. Disruption of thermo-tolerant *Desmodesmus* sp. F51 in high pressure homogenization as a prelude to carotenoids extraction. *Biochem. Eng. J.* **2016**, *109*, 243–251,

- doi:10.1016/j.bej.2016.01.003.
51. Zhu, X.; Lundberg, B.; Cheng, Y.; Shan, L.; Xing, J.; Peng, P.; Chen, P.; Huang, X.; Li, D.; Ruan, R. Effect of high-pressure homogenization on the flow properties of citrus peel fibers. *J. Food Process Eng.* **2018**, *41*, doi:10.1111/jfpe.12659.
 52. Preece, K.E.; Hooshyar, N.; Zuidam, N.J. Whole soybean protein extraction processes: A review. *Innov. Food Sci. Emerg. Technol.* **2017**, *43*, 163–172.
 53. Saricaoglu, F.T.; Gul, O.; Besir, A.; Atalar, I. Effect of high pressure homogenization (HPH) on functional and rheological properties of hazelnut meal proteins obtained from hazelnut oil industry by-products. *J. Food Eng.* **2018**, *233*, 98–108, doi:10.1016/j.jfoodeng.2018.04.003.
 54. Zhang, W.; Xie, F.; Lan, X.; Gong, S.; Wang, Z. Characteristics of pectin from black cherry tomato waste modified by dynamic high-pressure microfluidization. *J. Food Eng.* **2018**, *216*, 90–97, doi:10.1016/j.jfoodeng.2017.07.032.
 55. Otoni, C.G.; Lodi, B.D.; Lorevice, M. V.; Leitão, R.C.; Ferreira, M.D.; Moura, M.R. d.; Mattoso, L.H.C. Optimized and scaled-up production of cellulose-reinforced biodegradable composite films made up of carrot processing waste. *Ind. Crops Prod.* **2018**, *121*, 66–72, doi:10.1016/j.indcrop.2018.05.003.
 56. Xing, J. jie; Cheng, Y. ling; Chen, P.; Shan, L.; Ruan, R.; Li, D.; Wang, L. jun Effect of high-pressure homogenization on the extraction of sulforaphane from broccoli (*Brassica oleracea*) seeds. *Powder Technol.* **2019**, *358*, 103–109, doi:10.1016/j.powtec.2018.12.010.
 57. Mustafa, W.; Pataro, G.; Ferrari, G.; Donsì, F. Novel approaches to oil structuring via the addition of high-pressure homogenized agri-food residues and water forming capillary bridges. *J. Food Eng.* **2018**, *236*, 9–18.
 58. Griffin, S.; Sarfraz, M.; Farida, V.; Nasim, M.J.; Ebokaiwe, A.P.; Keck, C.M.; Jacob, C. No time to waste organic waste: Nanosizing converts remains of food processing into refined materials. *J. Environ. Manage.* **2018**, *210*, 114–121, doi:10.1016/j.jenvman.2017.12.084.
 59. Ilyas, R.A.; Sapuan, S.M.; Ishak, M.R.; Zainudin, E.S. Sugar palm nanofibrillated cellulose (*Arenga pinnata* (Wurmb.) Merr): Effect of cycles on their yield, physic-chemical, morphological and thermal behavior. *Int. J. Biol. Macromol.* **2019**, *123*, 379–388, doi:10.1016/j.ijbiomac.2018.11.124.
 60. Jurić, S.; Ferrari, G.; Velikov, K.P.; Donsì, F. High-pressure homogenization treatment to recover bioactive compounds from tomato peels. *J. Food Eng.* **2019**, *262*, 170–180, doi:10.1016/j.jfoodeng.2019.06.011.
 61. Zhang, Y.; Shi, R.; Xu, Y.; Chen, M.; Zhang, J.; Gao, Q.; Li, J. Developing a stable high-performance soybean meal-based adhesive using a simple high-pressure homogenization technology. *J. Clean. Prod.* **2020**, *256*, 120336, doi:10.1016/j.jclepro.2020.120336.
 62. Zhou, L. High-Pressure Homogenization Effect on the Stability and Bioaccessibility of Bioactive Phytochemicals and Vitamins in the Food Matrix. In *Reference Module in Food Science*; Elsevier, 2019.
 63. Sentandreu, E.; Stinco, C.M.; Vicario, I.M.; Mapelli-Brahm, P.; Navarro, J.L.; Meléndez-

- Martínez, A.J. High-pressure homogenization as compared to pasteurization as a sustainable approach to obtain mandarin juices with improved bioaccessibility of carotenoids and flavonoids. *J. Clean. Prod.* **2020**, *262*, 121325, doi:10.1016/j.jclepro.2020.121325.
64. Quan, W.; Tao, Y.; Qie, X.; Zeng, M.; Qin, F.; Chen, J.; He, Z. Effects of high-pressure homogenization, thermal processing, and milk matrix on the in vitro bioaccessibility of phenolic compounds in pomelo and kiwi juices. *J. Funct. Foods* **2020**, *64*, 103633, doi:10.1016/j.jff.2019.103633.
65. Alongi, M.; Calligaris, S.; Anese, M. Fat concentration and high-pressure homogenization affect chlorogenic acid bioaccessibility and α -glucosidase inhibitory capacity of milk-based coffee beverages. *J. Funct. Foods* **2019**, *58*, 130–137, doi:10.1016/j.jff.2019.04.057.
66. Betoret, E.; Calabuig-Jiménez, L.; Patrignani, F.; Lanciotti, R.; Dalla Rosa, M. Effect of high pressure processing and trehalose addition on functional properties of mandarin juice enriched with probiotic microorganisms. *LWT - Food Sci. Technol.* **2017**, *85*, 418–422, doi:10.1016/j.lwt.2016.10.036.
67. Munekata, P.E.S.; Domínguez, R.; Budaraju, S.; Roselló-Soto, E.; Barba, F.J.; Mallikarjunan, K.; Roohinejad, S.; Lorenzo, J.M. Effect of innovative food processing technologies on the physicochemical and nutritional properties and quality of non-dairy plant-based beverages. *Foods* **2020**, *9*.
68. Toro-Funes, N.; Bosch-Fusté, J.; Veciana-Nogués, M.T.; Vidal-Carou, M.C. Influence of ultra-high-pressure homogenization treatment on the phytosterols, tocopherols, and polyamines of almond beverage. *J. Agric. Food Chem.* **2014**, *62*, 9539–9543, doi:10.1021/jf503324f.
69. Atalar, I. Functional kefir production from high pressure homogenized hazelnut milk. *LWT* **2019**, *107*, 256–263, doi:10.1016/j.lwt.2019.03.013.
70. Kapoor, R.; Pathak, S.; Najmi, A.K.; Aeri, V.; Panda, B.P. Processing of soy functional food using high pressure homogenization for improved nutritional and therapeutic benefits. *Innov. Food Sci. Emerg. Technol.* **2014**, *26*, 490–497, doi:10.1016/j.ifset.2014.05.015.
71. Jiang, T.; Liao, W.; Charcosset, C. Recent advances in encapsulation of curcumin in nanoemulsions: A review of encapsulation technologies, bioaccessibility and applications. *Food Res. Int.* **2020**, *132*, 109035.
72. Frank, K.; Köhler, K.; Schuchmann, H.P. Stability of anthocyanins in high pressure homogenisation. *Food Chem.* **2012**, *130*, 716–719, doi:10.1016/j.foodchem.2011.07.086.
73. Patrignani, F.; Siroli, L.; Serrazanetti, D.I.; Braschi, G.; Betoret, E.; Reinheimer, J.A.; Lanciotti, R. Microencapsulation of functional strains by high pressure homogenization for a potential use in fermented milk. *Food Res. Int.* **2017**, *97*, 250–257, doi:10.1016/j.foodres.2017.04.020.
74. Calabuig-Jiménez, L.; Betoret, E.; Betoret, N.; Patrignani, F.; Barrera, C.; Seguí, L.; Lanciotti, R.; Dalla Rosa, M. High pressures homogenization (HPH) to microencapsulate *L. salivarius* spp. *salivarius* in mandarin juice. Probiotic survival and in vitro digestion. *J. Food Eng.* **2019**, *240*, 43–48.
75. Bamba, B.S.B.; Shi, J.; Tranchant, C.C.; Xue, S.J.; Forney, C.F.; Lim, L.T.; Xu, W.; Xu, G.

- Coencapsulation of polyphenols and anthocyanins from blueberry pomace by double emulsion stabilized by whey proteins: Effect of Homogenization parameters. *Molecules* **2018**, *23*, doi:10.3390/molecules23102525.
76. Cilek Tatar, B.; Sumnu, G.; Oztop, M. Microcapsule characterization of phenolic powder obtained from strawberry pomace. *J. Food Process. Preserv.* **2019**, *43*, 1–8, doi:10.1111/jfpp.13892.
77. Ester, B.; Noelia, B.; Laura, C.J.; Francesca, P.; Cristina, B.; Rosalba, L.; Marco, D.R. Probiotic survival and in vitro digestion of *L. salivarius* spp. *salivarius* encapsulated by high homogenization pressures and incorporated into a fruit matrix. *LWT* **2019**, *111*, 883–888, doi:10.1016/j.lwt.2019.05.088.
78. Siroli, L.; Braschi, G.; Rossi, S.; Gottardi, D.; Patrignani, F.; Lanciotti, R. Lactobacillus paracasei A13 and High-Pressure Homogenization Stress Response. *Microorganisms* **2020**, *8*, 439, doi:10.3390/microorganisms8030439.
79. Muramalla, T.; Aryana, K.J. Some low homogenization pressures improve certain probiotic characteristics of yogurt culture bacteria and *Lactobacillus acidophilus* LA-K1. *J. Dairy Sci.* **2011**, *94*, 3725–3738, doi:10.3168/jds.2010-3737.
80. Tabanelli, G.; Burns, P.; Patrignani, F.; Gardini, F.; Lanciotti, R.; Reinheimer, J.; Vinderola, G. Effect of a non-lethal High Pressure Homogenization treatment on the in vivo response of probiotic lactobacilli. *Food Microbiol.* **2012**, *32*, 302–307, doi:10.1016/j.fm.2012.07.004.
81. Patrignani, F.; Serrazanetti, D.I.; Mathara, J.M.; Siroli, L.; Gardini, F.; Holzapfel, W.H.; Lanciotti, R. Use of homogenisation pressure to improve quality and functionality of probiotic fermented milks containing *Lactobacillus rhamnosus* BFE 5264. *Int. J. Dairy Technol.* **2016**, *69*, 262–271, doi:10.1111/1471-0307.12251.
82. Burns, P.G.; Patrignani, F.; Tabanelli, G.; Vinderola, G.C.; Siroli, L.; Reinheimer, J.A.; Gardini, F.; Lanciotti, R. Potential of high pressure homogenisation on probiotic Caciotta cheese quality and functionality. *J. Funct. Foods* **2015**, *13*, 126–136, doi:10.1016/j.jff.2014.12.037.
83. Barrera, C.; Burca, C.; Betoret, E.; García-Hernández, J.; Hernández, M.; Betoret, N. Improving antioxidant properties and probiotic effect of clementine juice inoculated with *Lactobacillus salivarius* spp. *salivarius* (CECT 4063) by trehalose addition and/or sublethal homogenisation. *Int. J. Food Sci. Technol.* **2019**, *54*, 2109–2122, doi:10.1111/ijfs.14116.
84. Siroli, L.; Braschi, G.; Rossi, S.; Gottardi, D.; Patrignani, F.; Lanciotti, R. Lactobacillus paracasei A13 and high-pressure homogenization stress response. *Microorganisms* **2020**, *8*, doi:10.3390/microorganisms8030439.
85. Lanciotti, R.; Patrignani, F.; Iucci, L.; Saracino, P.; Guerzoni, M.E. Potential of high pressure homogenization in the control and enhancement of proteolytic and fermentative activities of some *Lactobacillus* species. *Food Chem.* **2007**, *102*, 542–550, doi:10.1016/j.foodchem.2006.06.043.