

Article

Targeting Hypoxia Sensitizes TNBC to Cisplatin and Promotes Inhibition of Both Bulk and Cancer Stem Cells

Andrew Sulaiman^{1,2,3,‡}, Sarah McGarry^{1,2,‡}, Jason Chambers¹, Emil Al-Kadi¹, Alexandra Phan¹, Li Li^{1,2}, Karan Mediratta¹, Jim Dimitroulakos^{1,4}, Christina Addison^{1,4}, Xuguang Li^{1,5}, Lisheng Wang^{1,2,6,*}

¹ Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine, University of Ottawa, 451 Smyth Road, Ottawa, Ontario K1H 8M5, Canada

² Ottawa Institute of Systems Biology, University of Ottawa, 451 Smyth Road, Ottawa, Ontario K1H 8M5, Canada

³ Department of Basic Science, Kansas City University of Medicine and Bioscience, 1750 Independence Ave, Kansas City, MO 64106, USA

⁴ Centre for Cancer Therapeutics, Ottawa Hospital Research Institute, Ottawa, Ontario K1H 8L6, Canada

⁵ Centre for Biologics Evaluation, Biologics and Genetic Therapies Directorate, Health Canada, Sir Frederick G. Banting Research Centre, Ottawa, Canada

⁶ Regenerative Medicine Program, Ottawa Hospital Research Institute, Ottawa, Ontario K1H 8L6, Canada

‡ These authors contributed equally

* Correspondence: lisheng.wang@uottawa.ca (L.W.); Tel.: +1-613-562-5624

Abstract: Development of targeted therapies for triple-negative breast cancer (TNBC) is an unmet medical need. Cisplatin has demonstrated its promising potential for the treatment of TNBC in clinical trials; however, cisplatin treatment is associated with hypoxia that in turn promotes cancer stem cell (CSC) enrichment and drug resistance. Therapeutic approaches to attenuate this may lead to increased cisplatin efficacy in the clinic for the treatment of TNBC. In this report, we analyzed clinical dataset of TNBC and found that TNBC patients possessed higher levels of EGFR and hypoxia gene expression. A similar expression pattern was also observed in cisplatin-resistant ovarian cancer cells. We thus developed a new therapeutic approach to inhibit EGFR and hypoxia by combination of metformin and gefitinib that sensitized TNBC cells to cisplatin and led to the inhibition of both CD44+/CD24- and ALDH+ CSCs. We demonstrated a similar inhibition efficacy on organotypic cultures of TNBC patient samples *ex vivo*. Since these drugs have already been used frequently in the clinic, this study illustrates a novel, clinically translatable therapeutic approach to treat patients with TNBC.

Keywords: Triple Negative Breast Cancer; Cancer Stem Cell; Hypoxia; EGFR; Cisplatin; PDX

1. Introduction

Breast cancer is one of the leading causes of cancer-related deaths in women throughout the world [1]. The triple-negative breast cancer (TNBC) subtype is characterized as negative for the estrogen receptor 1 (ESR1), progesterone receptor (PGR), and human epidermal growth factor receptor type 2 (HER2).

In contrast to other breast cancer subtypes, TNBC does not have specific targets and is reliant on chemotherapeutic regimens for systemic treatment and disproportionately accounts for the majority of breast cancer related deaths. Cisplatin is a platinum 2 complex capable of creating adducts, causing DNA damage, and subsequently inducing apoptosis in a multitude of cancers [2]. Recent clinical trials have showed efficacy of cisplatin in combinational chemotherapy in comparison to conventional chemotherapeutic approaches for the treatment of TNBC [3-5]. However, others have demonstrated lackluster outcomes, and underlying mechanisms remain convoluted [6,7].

Hypoxia following cisplatin treatment has been reported and associated with the onset of drug resistance and disease progression [8]. Additionally, while chemotherapy effectively inhibits bulk tumor populations, it enriches cancer stem cells (CSCs) [9]. CSCs retain stem cell-like characteristics and are highly associated with drug resistance, metastasis and tumorigenicity [10,11]. In breast cancer, two main populations of CSCs have been characterized according to the expression of CD44+/CD24- or ALDH+ [10,12,13]. Two populations of CSC exhibited plasticity and strong tumorigenic capacity, capable of generating new tumors and responsible for disease relapse. Thus, a new strategy to simultaneously target both CSC populations and suppress hypoxia will lead to the effective treatment of TNBC.

Gefitinib (an EGFR inhibitor) elicits anti-hypoxic properties and has been demonstrated to overcome cisplatin-induced hypoxia resistance in TNBC [14]. However, gefitinib is prone to resistance via stimulation of bypass signalling which in turn promotes Akt/mTORC1 and subsequent hypoxia, diminishing the efficacy of gefitinib for long-term treatment [15-17]. To circumvent this, metformin (an AMPK activator and first line treatment for type II diabetes) has been employed. Metformin exhibits anti-hypoxic effects via inhibition of mTORC1 activity and HIF-1 α protein stability [18,19]. Metformin has been shown to reverse gefitinib hypoxia resistance in head and neck squamous cell carcinoma and attenuate hypoxia-induced cisplatin resistance; moreover, metformin has been found to target CSCs in TNBC [20-22]. However, whether the combination of metformin and gefitinib can prevent cisplatin-induced hypoxia in TNBC, increase apoptosis and inhibit CSCs has not yet been reported.

In this report, we found that TNBC tumors from the patients expressed high levels of EGFR and hypoxia genes compared to normal mammary tissue in a clinical database. As such, we developed a combinational therapy consisting of metformin (AMPK activator), gefitinib (EGFR inhibitor) and cisplatin. We found that inhibition of EGFR and hypoxia sensitized TNBC cell lines to cisplatin treatment. Both epithelial and mesenchymal CSC populations could be inhibited. These findings were further verified using ex vivo organotypic cultures of TNBC clinical samples and TNBC patient-derived xenograft tumors. Together, these findings present a new strategy to prevent cisplatin-induced CSCs and hypoxia, which may lead to an efficacy treatment to reduce disease relapse in TNBC patients..

2. Results

2.1. Upregulated gene expressions of EGFR and hypoxia associate with cisplatin resistance, anti-apoptosis, and stemness

Clinical trials showed mixed results for the treatment of TNBC with cisplatin with convoluted mechanisms [3-7]. To identify potential targets associated with the cisplatin resistance (cisplatin sensitive and resistant) due to unavailable dataset for TNBC, we adopted an ovarian cancer bioinformatics model. We assessed gene overexpression in cisplatin resistant vs. naive ovarian cancer cells using the NCBI Gene Expression Omnibus (GEO2R). It was found that cisplatin resistance was associated with the increased expressions of EGFR, hypoxia, stemness and anti-apoptotic genes (Supplemental Figure 1A-D). We then analyzed NCBI Gene Expression Omnibus (GEO2R) to compare 30 TNBC with 13 normal mammary tissue samples and identified a similar trend where TNBC samples possessed high levels of EGFR, hypoxia, stemness and anti-apoptotic gene expressions compared to normal mammary tissues (Figure 1A-D). EGFR and hypoxia have both been associated with TNBC resistance to cisplatin [35,36], suggesting potentially clinical targets for further investigation.





A EGFR**B Hypoxia****C Apoptosis****D CSC Related Genes**

Figure 1. TNBC is associated with high levels of hypoxia, EGFR, stemness-related and anti-apoptosis related gene expression. (A-D) The relative expression levels (A.U arbitrary unit) of genes in 30 TNBC and 13 normal mammary tissue samples using the NCBI Gene Expression Omnibus (GEO2R). GSE38959 samples were analyzed using the GPL4133 Agilent-014850 Whole Human Genome Microarray 4x44K.

2.2. Combination of Hypoxia and EGFR Inhibitors with Cisplatin Effectively Suppress TNBC bulk and CSC populations

To determine whether hypoxia and EGFR inhibition could increase cisplatin efficacy, 25 μM of metformin (an AMPK activator used to treat type II diabetes with anti-hypoxic effects), 5 μM gefitinib (an EGFR inhibitor) and 5 μM of cisplatin at clinically relevant concentrations were used in different combinations (designated as CMG). As shown in Figure 2A-B, CMG treatment for 120 hours significantly reduced bulk cell viabilities in both SUM 149-PT and MDA-MB-231 TNBC cell lines.




Figure 2. Co-inhibition of hypoxia and EGFR in combination with cisplatin effectively suppress bulk tumor cells and CSC subpopulations in TNBC. (A) Relative viability (MTT assay) of SUM 149-PT TNBC cells after 120 hours of exposure to DMSO (vehicle control), cisplatin (Cis, 5 μ M), gefitinib (Gef, 5 μ M) and/or Metformin (Met, 25 μ M). (B) Relative viability (viable cell counting) of MDA-MB-231 TNBC cells after 120 hours of the same treatment as described in (A) (C-D) Living CSCs (CD44+/CD24-) in SUM 149-PT and MDA-MB-231 cells after 120 hours of exposure to the drugs as described in (A). (E-F) Living CSCs (ALDH+) in SUM 149-PT and MDA-MB-231 cells after 120 hours of exposure to the drugs as described in (A). (G-H) Representative flow cytometric plots of the CSC subpopulations (CD44 + /CD24- or ALDH+) in MDA-MB-231 cells after 120 hours of treatment with the drugs as described in (A). Data represent means \pm SD, n = 3 repeats; *P < 0.05, **P < 0.01, ***P < 0.001.

To assess whether the CMG combinational therapy could also target both epithelial-like and mesenchymal-like CSC populations, we performed FACs analysis on SUM 149-PT and MDA-MB-231 cells after 120 hours of treatment. CSCs have been closely associated with drug resistance, cancer

metastasis, and poor prognosis and chemotherapeutics enriched CSCs [9]. There are two main CSC populations characterized in TNBC: CD44+/CD24- mesenchymal and ALDH+ epithelial CSCs. Two CSC subpopulations exhibit plasticity and 100-fold greater tumor-initiating capacity than their non-CSC counterparts [11,37].

We found that CMG effectively inhibited CD44+/CD24- CSC population in both SUM 149-PT and MDA-MB-231 TNBC cell lines and reduced the living CSCs by ~90% (Figure 2C-D and G, Supplemental Figure 2A). In contrast, single and dual treatments exhibited weak or mixed results. Additionally, gefitinib alone did not inhibit CD44+/CD24- CSCs in MDA-MB-231 cells although it showed efficacy in SUM149-PT cells.

For the ALDH+ CSC subpopulation, we found that the combinations of cisplatin+getinib, metformin+getinib and CMG were all effective (Figure 2E-F and H, Supplemental Figure) with CMG reducing ALDH+ CSCs in both TNBC cell lines by ~90%. Together, these data demonstrate that the CMG combination can effectively inhibit TNBC bulk, CD44+/CD24- mesenchymal CSCs and ALDH+ epithelial CSC populations in both MDA-MB-231 and SUM149-PT cell lines at clinically relevant concentrations.

2.3. Combination of Metformin, Gefitinib and Cisplatin Effectively Repress Hypoxia and Promote Apoptosis in TNBC

As hypoxia has been closely associated with CSC enrichment and chemoresistance in breast cancer, we asked whether combinational CMG treatment suppressed hypoxia [38,39]. Hypoxia was assessed using a luciferase reporter plasmid containing three hypoxia response elements from the P_{gk}-1 gene upstream of firefly luciferase [26]. Using this system, active HIFs would bind to hypoxia response elements to mediate transcription and induce luciferase transcription [40]. We found that cisplatin increased the transcriptional activity of HIFs while the combination of metformin and gefitinib (MG) reduced HIF activity (Figure 3A). Moreover, the combination of CMG was able to antagonize cisplatin-upregulated HIF activity (Figure 3A). Similar findings were observed using a HIF1 α oxygen-dependent degradation sequence luciferase reporter (ODD-Luc) hypoxia reporter (Figure 3B) [25]. Additionally, RT-qPCR analysis of hypoxia-response-element containing genes showed that CMG treatment suppressed PDK1 and LDH1 gene expression (Figure 3C-D) [41,42].




Figure 3. Co-inhibition of hypoxia and EGFR suppresses cisplatin-upregulated hypoxia and promotes apoptosis in TNBC. (A-B) Relative luciferase reporter expression of MDA-MB-231 TNBC cells after 24 hours of drug treatment. Cells were transfected with HRE-Luciferase or ODD-Luciferase and normalized with the pRL-SV40P Renilla construct. The following day, cells were treated with DMSO (vehicle control), cisplatin (Cis, 5 μ M), Gefitinib (Gef, 5 μ M) and/or Metformin (Met, 25 μ M). (C-D) RT-qPCR analysis and comparison of relative mRNA expression levels of hypoxia genes (*PDK1* and *LDH1*) after treatment for 48 hours as described in (A-B). (E) Flow cytometric analysis of cell states (Dead, Alive, Early Apoptosis or Late Apoptosis) based on Annexin-V/7-AAD markers after 72 hours of exposure to DMSO (vehicle control), cisplatin (Cis, 5 μ M), gefitinib (Gef, 5 μ M) and/or Metformin (Met, 25 μ M). Data represent means \pm SE, n = 3 repeats; *P < 0.05, **P < 0.01, ***P < 0.001.

FACs analysis was performed to assess whether inhibition of hypoxia with CMG would correlate with a sensitization of cells to cisplatin-mediated apoptosis. Individual and dual combinations of cisplatin, metformin and gefitinib did not significantly induce early apoptosis (Annexin-V+/7-AAD-) or late apoptosis (Annexin-V+/7-AAD+) in TNBC cells (Figure 3E). Early apoptotic cells represent the cells undergoing apoptosis while retaining an intact cell membrane at the time of analysis. Notably, only the CMG combination significantly increased both early and late apoptosis in MDA-MB-231 cells (Figure 3E). Together, these data suggest that the CMG combinational treatment is able to inhibit hypoxia and promote TNBC apoptosis.

2.4. CMG Treatment Effectively Inhibits the Viability of TNBC Organotypic Cultures and Reduces Their CSC Subpopulations

To verify the above findings in a clinically relevant model, we obtained two clinical samples (CRDCA and SEM-1) from patients with TNBC and three TNBC patient-derived xenograft (PDX) tumors (HCI-001, HCI-002 and HCI-016). Samples were sliced into 2 × 1mm, cultured *ex vivo* and treated with the drug in different combinations at the clinically relevant concentrations for 144 hours. Alamar blue viability analysis demonstrated the efficacy of CMG treatment in all five clinically heterologous samples, showing a significant reduction in cell viability post-treatment (Figure 4A). Moreover, CMG combination also significantly suppressed both epithelial (ALDH+) and mesenchymal (CD44+/CD24-) CSC subpopulations in all five clinically heterologous samples (Figure 4B-E).




Figure 4. Co-inhibition of Hypoxia and EGFR in combination with Cisplatin effectively inhibits the viability of patient tumors and CSC populations in ex vivo organotypic cultures. (A) Alamar blue viability analysis of organotypic cultures from two primary patient TNBC tumors (CRDCA and SEM-1 samples) and three patient-derived xenograft samples (HCI-001, HCI-002, HCI-016) after 144 hours of exposure to DMSO (vehicle control), Cisplatin (Cis, 5 μ M), Gefitinib (Gef, 5 μ M) and/or Metformin (Met, 25 μ M). (B-C) Representative flow cytometric plots of the CSC subpopulations (CD44 + /CD24- or ALDH+) in SEM-1 and HCI-001 cells after 144 hours of treatment with the different drugs as described in (A). (D-E) Flow cytometric data of the relative living CSC (CD44 + /CD24- or ALDH+) populations in SEM-1, CRDCA, HCI-016 and HCI-001 patient TNBC tumors after 144 hours of exposure to the drugs as described in (A). Data represent means \pm SE, n = 3 repeats; *P < 0.05, **P < 0.01, ***P < 0.001.

Of note, the results from clinical TNBC samples (3 PDX and 2 clinical samples) amongst the single drug inhibitors are not same as the results from TNBC cell lines (as shown in Figure 3). Indeed, the gene expressions of EGFR, apoptotic resistance, stemness and hypoxia between TNBC patient tumors and TNBC cell line were different in the NCBI Gene Expression Omnibus (GEO2R) (Supplemental Figure 3A-C). These differences also highlight the importance to use clinically translatable models such as PDX for the development of therapeutic interventions.

3. Discussion

Cisplatin is a chemotherapeutic drug and has been widely used for the effective treatment of ovarian, cervical, bone, esophageal and lung cancers [43-48]. Renewed interest in cisplatin has led to more clinical trials, showing efficacious results in the treatment of other types of cancer, including TNBC [49-54]. Clinical trials using cisplatin for the treatment of TNBC discourage the use of cisplatin alone as adjuvant or neoadjuvant therapy [50]. However, cisplatin in combination with gemcitabine/docetaxel or bevacizumab in clinical trials has shown better outcomes [3,55,56]. Trial CBCSG006 found that cisplatin and gemcitabine combination displayed superior efficacy in comparison to the conventional paclitaxel and gemcitabine combination for the treatment of metastatic TNBC [49].

A drawback for cisplatin treatment is its modulation of an assortment of signalling pathways that bars its long-term efficacy. Cisplatin has been found to induce Interleukin-6 (IL-6) which promotes the accumulation of hypoxia inducible factors and CSC enrichment [57,58]. Cisplatin has also been demonstrated to induce STAT3 and NF- κ B expression following treatment. Both STAT3 and NF- κ B pathways are direct regulators to stimulate HIF-1 α , chemoresistance and CSC enrichment [59-61]. With the expanding role of cisplatin in TNBC treatment, suppressing cisplatin-induced hypoxia is critical for preventing cisplatin-resistance and CSC enrichment following treatment [57,62-64].

To prevent cisplatin-induced hypoxia, we employed metformin (first-line therapy for the treatment of type II diabetes), which was found to potently suppress hypoxia [19,65-67]. Metformin inhibits complex I of the mitochondria to indirectly stimulate AMPK, which in turn activates TSC1/TSC2 to inhibit mTORC1 and subsequent hypoxia signalling. Mechanistic analysis of metformin in an esophageal cancer cell model demonstrated a widespread downregulation of P13K/mTOR related proteins and stemness-related proteins, which correlated with a reduction in ALDH⁺ CSCs and increased apoptosis when combined with chemotherapy [68]. However, a study using AMPK α siRNA on HEP2G and Huh7 liver cancer cell lines showed that AMPK activation did not affect HIF-1 α . Interestingly, treatment with metformin was still able to potently suppress HIF-1 α protein levels and transcriptional activity via AMPK independent modulation of HIF1 α protein stability [19]. Thus, metformin exhibits anti-hypoxic properties through AMPK dependant and independent mechanisms.

Short-term Gefitinib treatment prevents EGFR activity and subsequently activate Akt/mTORC1, leading to inhibition of cisplatin-induced hypoxia [69,70]. Long-term gefitinib treatment promotes Akt/mTORC1 signalling via activation of STAT3, PLAU and/or TGF- α , which leads to hypoxia, CSC related gene expression and gefitinib resistance [16,71]. We found that even short-term treatment of TNBC cells with gefitinib led to the upregulation of hypoxia-related PDK1 gene, a potential early sign of resistance (Figure 3C).

Since gene expressions of hypoxia and EGFR were upregulated in TNBC patient tumors in comparison to normal mammary tissue (Figure 1A-B), we reasoned that their inhibition may increase cisplatin efficacy in TNBC. In addition, upregulation of hypoxia and EGFR is also associated with cisplatin resistance in ovarian cancer models (Supplemental Figure 1A-D), which further supports this possibility. Indeed, CMG combination effectively reducing TNBC cell viability and inhibiting both mesenchymal CD44⁺/CD24⁻ CSCs and epithelial ALDH⁺ CSCs (Figure 2A-H and Supplemental Figure 2A-B). Luciferase-gene reporter analysis demonstrated that cisplatin promoted TNBC hypoxia (Figure 3A-C), which may associate with drug resistance and CSC enrichment. However, CMG combination was able prevent cisplatin-induced hypoxia and promote apoptosis (Figure 3A-E).

Using organotypic cultures of five TNBC patient tumor fragments, we further verified our findings. Importantly, we revealed a differential pattern of gene expressions between TNBC patient tumors TNBC cell lines (Supplemental Figure 3A-C). This highlights the importance to use ex vivo patient tumors in experiments for clinical translation. Of note, the patient tumors from PDX samples were selected to represent a clinical setting, including paclitaxel-resistant HCI-001, paclitaxel-sensitive HCI-002 and naïve HCI-016 (obtained from patient without prior exposure to chemotherapy) [72,73]. Importantly, combination of CMG at clinically relevant dosages was capable

of inhibiting both bulk and CSC subpopulations in all five clinical tumors in ex vivo organotypic cultures (Figure 4A-E) [74-76]. As assessment of drug efficacy using short-term organotypic cultures of PDX have demonstrated clinical concordance [77-80], our data support the clinical translatability of the CMG treatment. As cisplatin, gefitinib and metformin are all currently used in clinic, our study may lead to a new avenue in TNBC treatment.

4. Materials and Methods

4.1 Cell culture and reagents

MDA-MB-231 breast cancer cells were purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA) and maintained in DMEM-F12 media supplemented with 10% Fetal bovine serum (FBS, HyClone, Logan, UT, USA) and 1% penicillin/streptomycin. SUM 149-PT breast cancer cells were obtained from Asterand (Detroit, MI, USA) and cultured in Hams F-12 media (Mediatech, Manassas, VA, USA) containing 5% FBS, 5 µg/mL insulin, 1 µg/mL hydrocortisone, 10 mm HEPES, and 1% penicillin/streptomycin. Cells were cultured at 37 °C in a 5% CO₂ incubator. Cisplatin was purchased from Caymen Chemical Company (Ann Arbor, Michigan, USA), gefitinib from LC labs- G-4408 (Woburn, Massachusetts, USA) and metformin from Caymen Chemical Company (Ann Arbor, Michigan, USA).

4.2 Organotypic cultures of patient TNBC breast tumor and patient-derived xenograft fragments

Tumor tissues from two TNBC patients undergoing routine surgical procedures were obtained following the protocol approved by The Ottawa Hospital Research Ethics Board (Protocol# 20120559-01H). Approximately 2 mm cores were obtained using a sterile biopsy punch that was further sliced with a scalpel to obtain approximately 2 × 1 mm tumor slices. The slices were randomized and placed into a well of 24-well plate and cultured in DMEM-F12 medium supplemented with 10% FBS, 1% penicillin/streptomycin, 1 µg/ml insulin, 0.5 ng/ml hydrocortisol and 3 ng/ml epidermal growth factor. The tissue fragments were treated with the same concentrations of inhibitors as described in the figures, followed by a viability assay and flow cytometric analysis as previously described [23,24]. The TNBC patient-derived xenograft samples HCI-001, HCI-002, HCI-016 was obtained from University of Utah, cut similarly to the patient tumor samples and cultured in the same conditions as the clinical samples.

4.3 Flow cytometry analysis

Dissociated cancer cells were filtered through a 4 µm strainer and suspended in PBS supplemented with 2% FBS and 2 mM EDTA (FACS buffer) as previously described [24]. One µL of mouse IgG (1 mg/mL) was added and incubated at 4 °C for 10 minutes. The cells were then re-suspended in 1× binding buffer and anti-CD44 (APC) in combination with anti-CD24 (PE) antibodies (BD, Mississauga, ON, Canada) according to the manufacturer's instructions for 30 min. The cells were washed twice with FACS buffer and 7-aminoactinomycin D (7-AAD, eBioscience, San Diego, CA) and Annexin-V/V450 (BD) was added and incubated for 15 minutes at room temperature to assess dead and apoptotic cells. Flow cytometry was performed on the BD LSRFortessa. Data were analyzed with FlowJo software (Ashland, OR, USA).

4.4 Quantitative real-time PCR

Total RNAs were extracted using RNeasy kit (QIAGEN) and real-time qPCR (RT-qPCR) analysis was performed using Bio-Rad MyiQ (Bio-Rad, Hercules, CA, USA) as described previously [23]. The conditions for RT-qPCR reactions were one cycle at 95 °C for 20 seconds followed by 45 cycles at 95 °C for three seconds and annealing at 60 °C for 30 seconds. Results were normalized to the housekeeping gene 18S ribosomal RNA (18S) or GAPDH. Relative expression level of genes from different groups were calculated with the $2^{-\Delta\Delta CT}$ method and compared with the expression levels of appropriate control cells. Specific primer sequences for individual genes are listed in Table 1.

Table 1. Primers used in qPCR.

Genes	Forward	Reverse
18S	AACCCGTTGAACCCATT	CCATCCAATCGGTAGTAGCG
GAPDH	ACAGTCAGCCGCATCTTCTT	GACAAGCTTCCCGTTCTCAG
PDK1	CAACAGAGGTGTTTACCCCC	ATTTTCCTCAAAGGAACGCC
LDH1	GGCCTGTGCCATCAGTATCT	GGAGATCCATCATCTCTCCC

4.5 Luciferase Reporter

MDA-MB-231 TNBC cells were seeded into 24-well plates and transfected with 500 ng of ODD-Luciferase-pcDNA3 (Addgene Plasmid # 18965, a gift from William Kaelin) [25], or HRE-luciferase (Addgene Plasmid # 26731, a gift from Navdeep Chandel) [26] in conjunction with 500 ng Renilla pRL-SV40P construct (Addgene Plasmid #27163, a gift from Dr. Ron Prywes)[27] using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. After 18 hours, cells were treated with either DMSO (vehicle), cisplatin (5uM), metformin (25uM), gefitinib (5 μ M) or different combinations for 24 hours, after which cells were lysed and both Firefly and Renilla luciferase activities were quantified using a Dual-Luciferase® Reporter Assay System (Promega) following the manufacturer's instructions.

4.6 Cell viability Assays

MDA-MB-231 cells were seeded into 24 well plates (5 \times 10⁴ cells/well). After 120 hours of treatment, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT, 5 mg/ml) was added to determine cellular viability. Absorbance was read at 570 nm. For clinical/PDX tumors, Alamar blue viability analysis was performed by incubation with 10% Alamar blue reagent (Thermo Fisher Scientific) for 4 hours. Florescence was measured at 560 nm excitation and 590 nm emission. Alternatively, cell counting was preformed after trypsinization of cells into single cell suspension followed by 1:10 dilution with trypan blue dye, followed by counting viable cells in haemocytometer under microscopy.

4.7 Clinical database analysis

Gene Expression Omnibus2R database [28,29] was used to analyze various datasets. Dataset: GSE15709 was used to compare cisplatin resistant and sensitive ovarian cancer cell lines [30]: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE15709> (Supplemental Figure 1). Dataset GSE38959 was used to compare 30 TNBC patient samples to 13 normal mammary tissue samples [31]: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE38959>. Dataset GSE65216 was used to compare 55 TNBC patient samples to 12 TNBC cell lines [32-34]: <https://www.ncbi.nlm.nih.gov/geo/geo2r/?acc=GSE65216&platform=GPL570>.

4.8 Statistical analysis

For all clinical databases, the log rank test was performed to determine the statistical differences between groups. Data are expressed as means +/-Standard Deviation (SD) or Standard Error (SE). Statistical significance was determined using ANOVA or Student's t test wherever appropriate. Results were considered significant when * p < 0.05, ** p < 0.01, or *** p < 0.001.

Supplementary Materials: Supplementary materials can be found at the end of the document.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used "Conceptualization, X.X. and Y.Y.; methodology, X.X.; software, X.X.; validation, X.X., Y.Y. and Z.Z.; formal analysis, X.X.; investigation, X.X.; resources, X.X.; data curation, X.X.; writing—original draft preparation, X.X.; writing—review and editing, X.X.; visualization, X.X.; supervision, X.X.; project administration, X.X.; funding acquisition, Y.Y. All authors have read and agreed to the published version of the manuscript.", please turn to the [CRediT taxonomy](#) for the term explanation. Authorship must be limited to those who have contributed substantially to the work reported.

Funding: This work was supported by operating grants Cancer Research Society/University of Ottawa (24064) and Natural Sciences and Engineering Research Council (NSERC, RGPIN-2019-0522) to LW; Alexander Graham Bell Canada Graduate Scholarship of NSERC and the Michael Smith Foreign Study Supplements to AS.

Acknowledgments: We thank the Huntsman Cancer Institute in Salt Lake City, UT for the use of the Preclinical Research Resource (PRR) to provide the PDX (HCI-001, HCI-002, and HCI-016) samples.

Conflicts of Interest: The authors declare no conflict of interest

Abbreviations

EGFR	epidermal growth factor receptor
CD44	cluster of differentiation 44 (hyaluronic acid receptor)
CD24	cluster of differentiation 24
CSC	cancer stem cell
ALDH	aldehyde dehydrogenase
ESR1	estrogen receptor 1 (alpha)
PGR	progesterone receptor
HER2	human epidermal growth factor receptor type 2
PDX	patient derived xenograft
TNBC	triple negative breast cancer
CMG	Cisplatin, Metformin and Gefitinib combination;
MG	Metformin and Gefitinib combination

References

1. Siegel, R.L.; Miller, K.D.; Jemal, A. Cancer statistics, 2017. *CA: a cancer journal for clinicians* 2017, *67*, 7-30.
2. Go, R.S.; Adjei, A.A. Review of the comparative pharmacology and clinical activity of cisplatin and carboplatin. *Journal of Clinical Oncology* 1999, *17*, 409-409.
3. Hu, X.-C.; Zhang, J.; Xu, B.-H.; Cai, L.; Ragaz, J.; Wang, Z.-H.; Wang, B.-Y.; Teng, Y.-E.; Tong, Z.-S.; Pan, Y.-Y. Cisplatin plus gemcitabine versus paclitaxel plus gemcitabine as first-line therapy for metastatic triple-negative breast cancer (CBCSG006): a randomised, open-label, multicentre, phase 3 trial. *The lancet oncology* 2015, *16*, 436-446.
4. Rodler, E.T.; Kurland, B.F.; Griffin, M.; Gralow, J.R.; Porter, P.; Yeh, R.F.; Gadi, V.K.; Guenthoer, J.; Beumer, J.H.; Korde, L. Phase I study of veliparib (ABT-888) combined with cisplatin and vinorelbine in advanced triple-negative breast cancer and/or BRCA mutation-associated breast cancer. *Clinical Cancer Research* 2016, *22*, 2855-2864.
5. Ferreira, A.R.; Metzger-Filho, O.; Sarmiento, R.; Bines, J. Neoadjuvant Treatment of Stage IIB/III Triple Negative Breast Cancer with Cyclophosphamide, Doxorubicin, and Cisplatin (CAP Regimen): A Single Arm, Single Center Phase II Study (GBECAM 2008/02). *Frontiers in oncology* 2018, *7*, 329.
6. Su, Y.-W.; Hung, C.-Y.; Lam, H.-B.; Chang, Y.-C.; Yang, P.-S.J.C.M.I.O. A single institution experience of incorporation of cisplatin into adjuvant chemotherapy for patients with triple-negative breast cancer of unknown BRCA mutation status. 2018, *12*, 1179554918794672.
7. Silver, D.P.; Richardson, A.L.; Eklund, A.C.; Wang, Z.C.; Szallasi, Z.; Li, Q.; Juul, N.; Leong, C.-O.; Calogrias, D.; Buraimoh, A.J.J.o.c.o. Efficacy of neoadjuvant Cisplatin in triple-negative breast cancer. 2010, *28*, 1145.
8. Zhang, F.; Duan, S.; Tsai, Y.; Keng, P.C.; Chen, Y.; Lee, S.O.; Chen, Y. Cisplatin treatment increases stemness through upregulation of hypoxia-inducible factors by interleukin-6 in non-small cell lung cancer. *Cancer science* 2016, *107*, 746-754, doi:10.1111/cas.12937.
9. Jia, D.; Tan, Y.; Liu, H.; Ooi, S.; Li, L.; Wright, K.; Bennett, S.; Addison, C.L.; Wang, L.J.O. Cardamonin reduces chemotherapy-enriched breast cancer stem-like cells in vitro and in vivo. 2016, *7*, 771.
10. Liu, S.; Cong, Y.; Wang, D.; Sun, Y.; Deng, L.; Liu, Y.; Martin-Trevino, R.; Shang, L.; McDermott, S.P.; Landis, M.D. Breast cancer stem cells transition between epithelial and mesenchymal states reflective of their normal counterparts. *Stem cell reports* 2014, *2*, 78-91.
11. Al-Hajj, M.; Wicha, M.S.; Benito-Hernandez, A.; Morrison, S.J.; Clarke, M.F. Prospective identification of tumorigenic breast cancer cells. *Proceedings of the National Academy of Sciences* 2003, *100*, 3983-3988.

12. Sulaiman, A.; McGarry, S.; Li, L.; Jia, D.; Ooi, S.; Addison, C.; Dimitroulakos, J.; Arnaout, A.; Nessim, C.; Yao, Z. Dual inhibition of Wnt and Yes-associated protein signaling retards the growth of triple-negative breast cancer in both mesenchymal and epithelial states. *Molecular oncology* 2018, 12, 423-440.
13. Li, W.; Ma, H.; Zhang, J.; Zhu, L.; Wang, C.; Yang, Y. Unraveling the roles of CD44/CD24 and ALDH1 as cancer stem cell markers in tumorigenesis and metastasis. *Scientific Reports* 2017, 7, 13856.
14. Rho, J.K.; Choi, Y.J.; Lee, J.K.; Ryoo, B.-Y.; Na, I.I.; Yang, S.H.; Kim, C.H.; Yoo, Y.D.; Lee, J.C. Gefitinib circumvents hypoxia-induced drug resistance by the modulation of HIF-1 α . *Oncology reports* 2009, 21, 801-807.
15. Liu, S.; Yang, H.; Ge, X.; Su, L.; Zhang, A.; Liang, L. Drug resistance analysis of gefitinib-targeted therapy in non-small cell lung cancer. *Oncology letters* 2016, 12, 3941-3943.
16. Minakata, K.; Takahashi, F.; Nara, T.; Hashimoto, M.; Tajima, K.; Murakami, A.; Nurwidya, F.; Yae, S.; Koizumi, F.; Moriyama, H. Hypoxia induces gefitinib resistance in non-small-cell lung cancer with both mutant and wild-type epidermal growth factor receptors. *Cancer science* 2012, 103, 1946-1954.
17. Huang, W.-C.; Chen, Y.-J.; Li, L.-Y.; Wei, Y.-L.; Hsu, S.-C.; Tsai, S.-L.; Chiu, P.-C.; Huang, W.-P.; Wang, Y.-N.; Chen, C.-H. Nuclear translocation of epidermal growth factor receptor by Akt-dependent phosphorylation enhances breast cancer-resistant protein expression in gefitinib-resistant cells. *Journal of Biological Chemistry* 2011, 286, 20558-20568.
18. Kalender, A.; Selvaraj, A.; Kim, S.Y.; Gulati, P.; Brûlé, S.; Viollet, B.; Kemp, B.E.; Bardeesy, N.; Dennis, P.; Schlager, J.J. Metformin, independent of AMPK, inhibits mTORC1 in a rag GTPase-dependent manner. *Cell metabolism* 2010, 11, 390-401.
19. Zhou, X.; Chen, J.; Yi, G.; Deng, M.; Liu, H.; Liang, M.; Shi, B.; Fu, X.; Chen, Y.; Chen, L. Metformin suppresses hypoxia-induced stabilization of HIF-1 α through reprogramming of oxygen metabolism in hepatocellular carcinoma. *Oncotarget* 2016, 7, 873.
20. Yin, X.; Wei, Z.; Song, C.; Tang, C.; Xu, W.; Wang, Y.; Xie, J.; Lin, Z.; Han, W. Metformin sensitizes hypoxia-induced gefitinib treatment resistance of HNSCC via cell cycle regulation and EMT reversal. *Cancer management and research* 2018, 10, 5785.
21. Fujita, H.; Hirose, K.; Sato, M.; Fujioka, I.; Fujita, T.; Aoki, M.; Takai, Y. Metformin attenuates hypoxia-induced resistance to cisplatin in the HepG2 cell line. *Oncology letters* 2019, 17, 2431-2440.
22. Shi, P.; Liu, W.; Wang, H.; Li, F.; Zhang, H.; Wu, Y.; Kong, Y.; Zhou, Z.; Wang, C.; Chen, W.J.C.d. Metformin suppresses triple-negative breast cancer stem cells by targeting KLF5 for degradation. 2017, 3, 1-13.
23. Sulaiman, A.; Sulaiman, B.; Khouri, L.; McGarry, S.; Nessim, C.; Arnaout, A.; Li, X.; Addison, C.; Dimitroulakos, J.; Wang, L. Both bulk and cancer stem cell subpopulations in triple-negative breast cancer are susceptible to Wnt, HDAC, and ER α coinhibition. *FEBS letters* 2016, 590, 4606-4616.
24. Sulaiman, A.; McGarry, S.; Lam, K.M.; El-Sahli, S.; Chambers, J.; Kaczmarek, S.; Li, L.; Addison, C.; Dimitroulakos, J.; Arnaout, A. Co-inhibition of mTORC1, HDAC and ESR1 α retards the growth of triple-negative breast cancer and suppresses cancer stem cells. *Cell death & disease* 2018, 9, 815.
25. Safran, M.; Kim, W.Y.; O'Connell, F.; Flippin, L.; Günzler, V.; Horner, J.W.; DePinho, R.A.; Kaelin, W.G. Mouse model for noninvasive imaging of HIF prolyl hydroxylase activity: assessment of an oral agent that stimulates erythropoietin production. *Proceedings of the National Academy of Sciences* 2006, 103, 105-110.
26. Emerling, B.M.; Weinberg, F.; Liu, J.-L.; Mak, T.W.; Chandel, N.S. PTEN regulates p300-dependent hypoxia-inducible factor 1 transcriptional activity through Forkhead transcription factor 3a (FOXO3a). *Proceedings of the National Academy of Sciences* 2008, 105, 2622-2627.
27. Chen, X.; Prywes, R. Serum-induced expression of the cdc25A gene by relief of E2F-mediated repression. *Molecular and Cellular Biology* 1999, 19, 4695-4702.
28. Edgar, R.; Domrachev, M.; Lash, A.E. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic acids research* 2002, 30, 207-210.
29. Barrett, T.; Wilhite, S.E.; Ledoux, P.; Evangelista, C.; Kim, I.F.; Tomashevsky, M.; Marshall, K.A.; Phillippy, K.H.; Sherman, P.M.; Holko, M. NCBI GEO: archive for functional genomics data sets—update. *Nucleic acids research* 2012, 41, D991-D995.
30. Li, M.; Balch, C.; Montgomery, J.S.; Jeong, M.; Chung, J.H.; Yan, P.; Huang, T.H.; Kim, S.; Nephew, K.P. Integrated analysis of DNA methylation and gene expression reveals specific signaling pathways associated with platinum resistance in ovarian cancer. *BMC medical genomics* 2009, 2, 34.
31. Komatsu, M.; Yoshimaru, T.; Matsuo, T.; Kiyotani, K.; Miyoshi, Y.; Tanahashi, T.; Rokutan, K.; Yamaguchi, R.; Saito, A.; Imoto, S. Molecular features of triple negative breast cancer cells by genome-wide gene expression profiling analysis. *International journal of oncology* 2013, 42, 478-506.

32. Maubant, S.; Tesson, B.; Maire, V.; Ye, M.; Rigai, G.; Gentien, D.; Cruzalegui, F.; Tucker, G.C.; Roman-Roman, S.; Dubois, T. Transcriptome analysis of Wnt3a-treated triple-negative breast cancer cells. *PLoS one* 2015, 10, e0122333.
33. Maire, V.; Baldeyron, C.; Richardson, M.; Tesson, B.; Vincent-Salomon, A.; Gravier, E.; Marty-Prouvost, B.; De Koning, L.; Rigai, G.; Dumont, A. TTK/hMPS1 is an attractive therapeutic target for triple-negative breast cancer. *PLoS one* 2013, 8, e63712.
34. Maire, V.; Némati, F.; Richardson, M.; Vincent-Salomon, A.; Tesson, B.; Rigai, G.; Gravier, E.; Marty-Prouvost, B.; De Koning, L.; Lang, G. Polo-like kinase 1: a potential therapeutic option in combination with conventional chemotherapy for the management of patients with triple-negative breast cancer. *Cancer research* 2013, 73, 813-823.
35. Soleymani Abyaneh, H.; Gupta, N.; Radziwon-Balicka, A.; Jurasz, P.; Seubert, J.; Lai, R.; Lavasanifar, A.J.C. STAT3 but not HIF-1 α is important in mediating Hypoxia-Induced chemoresistance in MDA-MB-231, a triple negative breast cancer cell line. 2017, 9, 137.
36. Kim, E.M.; Mueller, K.; Gartner, E.; Boerner, J.J.o.s.r. Dasatinib is synergistic with cetuximab and cisplatin in triple-negative breast cancer cells. 2013, 185, 231-239.
37. Ginestier, C.; Hur, M.H.; Charafe-Jauffret, E.; Monville, F.; Dutcher, J.; Brown, M.; Jacquemier, J.; Viens, P.; Kleer, C.G.; Liu, S. ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell stem cell* 2007, 1, 555-567.
38. Shiraishi, A.; Tachi, K.; Essid, N.; Tsuboi, I.; Nagano, M.; Kato, T.; Yamashita, T.; Bando, H.; Hara, H.; Ohneda, O. Hypoxia promotes the phenotypic change of aldehyde dehydrogenase activity of breast cancer stem cells. *Cancer science* 2017, 108, 362-372.
39. Crowder, S.W.; Balikov, D.A.; Hwang, Y.-S.; Sung, H.-J. Cancer stem cells under hypoxia as a chemoresistance factor in the breast and brain. *Current pathobiology reports* 2014, 2, 33-40.
40. Dengler, V.L.; Galbraith, M.D.; Espinosa, J.M. Transcriptional regulation by hypoxia inducible factors. *Critical reviews in biochemistry and molecular biology* 2014, 49, 1-15.
41. Semba, H.; Takeda, N.; Isagawa, T.; Sugiura, Y.; Honda, K.; Wake, M.; Miyazawa, H.; Yamaguchi, Y.; Miura, M.; Jenkins, D.M.J.N.c. HIF-1 α -PDK1 axis-induced active glycolysis plays an essential role in macrophage migratory capacity. 2016, 7, 11635.
42. McGinn, O.; Gupta, V.K.; Dauer, P.; Arora, N.; Sharma, N.; Nomura, A.; Dudeja, V.; Saluja, A.; Banerjee, S.J.S.r. Inhibition of hypoxic response decreases stemness and reduces tumorigenic signaling due to impaired assembly of HIF1 transcription complex in pancreatic cancer. 2017, 7, 1-11.
43. Dasari, S.; Tchounwou, P.B. Cisplatin in cancer therapy: molecular mechanisms of action. *European journal of pharmacology* 2014, 740, 364-378.
44. Group, I.A.L.C.T.C. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. *New England Journal of Medicine* 2004, 350, 351-360.
45. Collaboration, N.C.f.L.A.C.C.M.-a. Neoadjuvant chemotherapy for locally advanced cervical cancer: a systematic review and meta-analysis of individual patient data from 21 randomised trials. 2003.
46. McClay, E.F.; Howell, S.B. A review: intraperitoneal cisplatin in the management of patients with ovarian cancer. *Gynecologic oncology* 1990, 36, 1-6.
47. Andronescu, E.; Fica, A.; Albu, M.G.; Mitran, V.; Sonmez, M.; Fica, D.; Ion, R.; Cimpean, A. Collagen-hydroxyapatite/cisplatin drug delivery systems for locoregional treatment of bone cancer. *Technology in cancer research & treatment* 2013, 12, 275-284.
48. Walsh, T.N.; Noonan, N.; Hollywood, D.; Kelly, A.; Keeling, N.; Hennessy, T.P. A comparison of multimodal therapy and surgery for esophageal adenocarcinoma. *New England Journal of Medicine* 1996, 335, 462-467.
49. Zhang, J.; Lin, Y.; Sun, X.; Wang, B.; Wang, Z.; Luo, J.; Wang, L.; Zhang, S.; Cao, J.; Tao, Z. Biomarker assessment of the CBCSG006 trial: a randomized phase III trial of cisplatin plus gemcitabine compared with paclitaxel plus gemcitabine as first-line therapy for patients with metastatic triple-negative breast cancer. *Annals of Oncology* 2018, 29, 1741-1747.
50. Silver, D.P.; Richardson, A.L.; Eklund, A.C.; Wang, Z.C.; Szallasi, Z.; Li, Q.; Juul, N.; Leong, C.-O.; Calogrias, D.; Buraimoh, A. Efficacy of neoadjuvant Cisplatin in triple-negative breast cancer. *Journal of clinical oncology* 2010, 28, 1145.
51. Baselga, J.; Gómez, P.; Greil, R.; Braga, S.; Climent, M.A.; Wardley, A.M.; Kaufman, B.; Stemmer, S.M.; Pêgo, A.; Chan, A. Randomized phase II study of the anti-epidermal growth factor receptor monoclonal antibody

- cetuximab with cisplatin versus cisplatin alone in patients with metastatic triple-negative breast cancer. *Journal of clinical oncology* 2013, 31, 2586.
52. Jovanović, B.; Mayer, I.A.; Mayer, E.L.; Abramson, V.G.; Bardia, A.; Sanders, M.E.; Kuba, M.G.; Estrada, M.V.; Beeler, J.S.; Shaver, T.M. A randomized phase II neoadjuvant study of cisplatin, paclitaxel with or without everolimus in patients with stage II/III Triple-Negative Breast Cancer (TNBC): responses and long-term outcome correlated with increased frequency of DNA damage response gene mutations, TNBC Subtype, AR Status, and Ki67. *Clinical Cancer Research* 2017, 23, 4035-4045.
 53. Isakoff, S.J.; Mayer, E.L.; He, L.; Traina, T.A.; Carey, L.A.; Krag, K.J.; Rugo, H.S.; Liu, M.C.; Stearns, V.; Come, S.E. TBCRC009: a multicenter phase II clinical trial of platinum monotherapy with biomarker assessment in metastatic triple-negative breast cancer. *Journal of clinical oncology* 2015, 33, 1902.
 54. Baek, D.W.; Park, J.-Y.; Lee, S.J.; Chae, Y.S.J.Y.U.J.o.M. Impressive effect of cisplatin monotherapy on a patient with heavily pretreated triple-negative breast cancer with poor performance. 2020.
 55. Fan, Y.; Xu, B.; Yuan, P.; Ma, F.; Wang, J.; Ding, X.; Zhang, P.; Li, Q.; Cai, R. Docetaxel–cisplatin might be superior to docetaxel–capecitabine in the first-line treatment of metastatic triple-negative breast cancer. *Annals of oncology* 2012, 24, 1219-1225.
 56. Ryan, P.; Tung, N.; Isakoff, S.; Golshan, M.; Richardson, A.; Corben, A.; Smith, B.; Gelman, R.; Winer, E.; Garber, J. Neoadjuvant cisplatin and bevacizumab in triple negative breast cancer (TNBC): safety and efficacy. *Journal of Clinical Oncology* 2009, 27, 551-551.
 57. Zhang, F.; Duan, S.; Tsai, Y.; Keng, P.C.; Chen, Y.; Lee, S.O.; Chen, Y. Cisplatin treatment increases stemness through upregulation of hypoxia-inducible factors by interleukin-6 in non-small cell lung cancer. *Cancer Science* 2016, 107, 746-754.
 58. Liu, C.C.; Lin, J.H.; Hsu, T.W.; Su, K.; Li, A.F.Y.; Hsu, H.S.; Hung, S.C. IL-6 enriched lung cancer stem-like cell population by inhibition of cell cycle regulators via DNMT1 upregulation. *International Journal of Cancer* 2015, 136, 547-559.
 59. Kuo, W.-Y.; Hwu, L.; Wu, C.-Y.; Lee, J.-S.; Chang, C.-W.; Liu, R.-S. STAT3/NF- κ B-Regulated Lentiviral TK/GCV Suicide Gene Therapy for Cisplatin-Resistant Triple-Negative Breast Cancer. *Theranostics* 2017, 7, 647.
 60. Morais, C.; Gobe, G.; Johnson, D.W.; Healy, H. Inhibition of nuclear factor kappa B transcription activity drives a synergistic effect of pyrrolidine dithiocarbamate and cisplatin for treatment of renal cell carcinoma. *Apoptosis* 2010, 15, 412-425.
 61. Gu, F.; Ma, Y.; Zhang, Z.; Zhao, J.; Kobayashi, H.; Zhang, L.; Fu, L. Expression of Stat3 and Notch1 is associated with cisplatin resistance in head and neck squamous cell carcinoma. *Oncology reports* 2010, 23, 671-676.
 62. Shin, D.H.; Choi, Y.-J.; Park, J.-W. SIRT1 and AMPK mediate hypoxia-induced resistance of non-small cell lung cancers to cisplatin and doxorubicin. *Cancer research* 2014, 74, 298-308.
 63. Fischer, C.; Leithner, K.; Wohlkoenig, C.; Quehenberger, F.; Bertsch, A.; Olschewski, A.; Olschewski, H.; Hrzenjak, A. Panobinostat reduces hypoxia-induced cisplatin resistance of non-small cell lung carcinoma cells via HIF-1 α destabilization. *Molecular cancer* 2015, 14, 1.
 64. Samanta, D.; Park, Y.; Ni, X.; Li, H.; Zahnow, C.A.; Gabrielson, E.; Pan, F.; Semenza, G.L. Chemotherapy induces enrichment of CD47+/CD73+/PDL1+ immune evasive triple-negative breast cancer cells. *Proceedings of the National Academy of Sciences* 2018, 201718197.
 65. Ye, J.; Chen, K.; Qi, L.; Li, R.; Tang, H.; Zhou, C.; Zhai, W. Metformin suppresses hypoxia-induced migration via the HIF-1 α /VEGF pathway in gallbladder cancer in vitro and in vivo. *Oncology reports* 2018, 40, 3501-3510.
 66. Tadakawa, M.; Takeda, T.; Li, B.; Tsuiji, K.; Yaegashi, N. The anti-diabetic drug metformin inhibits vascular endothelial growth factor expression via the mammalian target of rapamycin complex 1/hypoxia-inducible factor-1 α signaling pathway in ELT-3 cells. *Molecular and cellular endocrinology* 2015, 399, 1-8.
 67. Saha, I.B.; Regazzetti, C.; Robert, G.; Laurent, K.; Le Marchand-Brustel, Y.; Auberger, P.; Tanti, J.-F.; Giorgetti-Peraldi, S.; Bost, F. Metformin, independent of AMPK, induces mTOR inhibition and cell-cycle arrest through REDD1. *Cancer research* 2011, 71, 4366-4372.
 68. Honjo, S.; Ajani, J.A.; Scott, A.W.; Chen, Q.; Skinner, H.D.; Stroehlein, J.; Johnson, R.L.; Song, S.J.I.j.o.o. Metformin sensitizes chemotherapy by targeting cancer stem cells and the mTOR pathway in esophageal cancer. 2014, 45, 567-574.

69. Solomon, B.; Binns, D.; Roselt, P.; Weibe, L.I.; McArthur, G.A.; Cullinane, C.; Hicks, R.J. Modulation of intratumoral hypoxia by the epidermal growth factor receptor inhibitor gefitinib detected using small animal PET imaging. *Molecular cancer therapeutics* 2005, 4, 1417-1422.
70. Rho, J.K.; Choi, Y.J.; Lee, J.K.; Ryoo, B.-Y.; Na, I.I.; Yang, S.H.; Kim, C.H.; Yoo, Y.D.; Lee, J.C.J.O.r. Gefitinib circumvents hypoxia-induced drug resistance by the modulation of HIF-1 α . 2009, 21, 801-807.
71. Murakami, A.; Takahashi, F.; Nurwidya, F.; Kobayashi, I.; Minakata, K.; Hashimoto, M.; Nara, T.; Kato, M.; Tajima, K.; Shimada, N. Hypoxia increases gefitinib-resistant lung cancer stem cells through the activation of insulin-like growth factor 1 receptor. *PloS one* 2014, 9, e86459.
72. PDXNet. PDX DEVELOPMENT AND TRIAL CENTERS RESEARCH NETWORK. <https://brandi-davis-7wsf.squarespace.com/hcibcm>.
73. DeRose, Y.S.; Wang, G.; Lin, Y.C.; Bernard, P.S.; Buys, S.S.; Ebbert, M.T.; Factor, R.; Matsen, C.; Milash, B.A.; Nelson, E., et al. Tumor grafts derived from women with breast cancer authentically reflect tumor pathology, growth, metastasis and disease outcomes. *Nat Med* 2011, 17, 1514-1520, doi:10.1038/nm.2454.
74. 74. Rajkumar, P.; Mathew, B.S.; Das, S.; Isaiah, R.; John, S.; Prabha, R.; Fleming, D.H.J.J.o.C.; JCDR, D.R. Cisplatin concentrations in long and short duration infusion: implications for the optimal time of radiation delivery. 2016, 10, XC01.
75. McCreight, L.J.; Stage, T.B.; Connelly, P.; Lonergan, M.; Nielsen, F.; Prehn, C.; Adamski, J.; Brøsen, K.; Pearson, E.R.J.D., Obesity; Metabolism. Pharmacokinetics of metformin in patients with gastrointestinal intolerance. 2018, 20, 1593-1601.
76. McKillop, D.; Partridge, E.A.; Kemp, J.V.; Spence, M.P.; Kendrew, J.; Barnett, S.; Wood, P.G.; Giles, P.B.; Patterson, A.B.; Bichat, F.J.M.c.t. Tumor penetration of gefitinib (Iressa), an epidermal growth factor receptor tyrosine kinase inhibitor. 2005, 4, 641-649.
77. Borodovsky, A.; McQuiston, T.J.; Stetson, D.; Ahmed, A.; Whitston, D.; Zhang, J.; Grondine, M.; Lawson, D.; Challberg, S.S.; Zinda, M. Generation of stable PDX derived cell lines using conditional reprogramming. *Molecular cancer* 2017, 16, 177.
78. Yuan, H.; Myers, S.; Wang, J.; Zhou, D.; Woo, J.A.; Kallakury, B.; Ju, A.; Bazylewicz, M.; Carter, Y.M.; Albanese, C. Use of reprogrammed cells to identify therapy for respiratory papillomatosis. *New England Journal of Medicine* 2012, 367, 1220-1227.
79. Fong, E.L.S.; Toh, T.B.; Lin, Q.X.X.; Liu, Z.; Hooi, L.; Rashid, M.B.M.A.; Benoukraf, T.; Chow, E.K.-H.; Huynh, T.H.; Yu, H. Generation of matched patient-derived xenograft in vitro-in vivo models using 3D macroporous hydrogels for the study of liver cancer. *biomaterials* 2018, 159, 229-240.
80. Roife, D.; Dai, B.; Kang, Y.a.; Perez, M.V.R.; Pratt, M.; Li, X.; Fleming, J.B. Ex vivo testing of patient-derived xenografts mirrors the clinical outcome of patients with pancreatic ductal adenocarcinoma. *Clinical Cancer Research* 2016, 22, 6021-6030.