

Article

Climate change and alpine scree: no future for glacial relict *Papaver occidentale* (Papaveraceae) in Western Prealps

Yann Fragnière¹, Loïc Pittet¹, Benoît Clément¹, Sébastien Bétrisey¹, Emanuel Gerber², Michał Ronikier³, Christian Parisod⁴, Gregor Kozłowski^{1,2,5*}

¹ Department of Biology and Botanic Garden, University of Fribourg, Chemin du Musée 10, CH-1700 Fribourg, Switzerland (yann.fragniere@unifr.ch, loic.pittet@unifr.ch, sebastien.betrisey@unifr.ch, benoit.clement@unifr.ch, gregor.kozlowski@unifr.ch)

² Natural History Museum Fribourg, Chemin du Musée 6, CH-1700 Fribourg, Switzerland (emanuel.gerber@fr.ch)

³ Institute of Botany, Polish Academy of Sciences, Lubicz 46, PL-31-512 Krakow, Poland (m.ronikier@botany.pl)

⁴ Institute of Plant Sciences, University of Bern, Altenbergrain 21, CH-3013 Bern, Switzerland (christian.parisod@ips.unibe.ch)

⁵ Plant Systematics and Evolutionary Biology, Shanghai Chenshan Plant Science Research Center, Chinese Academy of Sciences, 201602 Shanghai, China

* **Correspondence:** Gregor Kozłowski, Department of Biology and Botanic Garden, University of Fribourg, Chemin du Musée 10, CH-1700 Fribourg, Switzerland (gregor.kozlowski@unifr.ch)

Abstract: A single paragraph of about 200 words maximum. For research articles, abstracts should give a pertinent overview of the work. We strongly encourage authors to use the following style of structured abstracts, but without headings: (1) Background: Place the question addressed in a broad context and highlight the purpose of the study; (2) Methods: Describe briefly the main methods or treatments applied; (3) Results: Summarize the article's main findings; and (4) Conclusions: Indicate the main conclusions or interpretations. The abstract should be an objective representation of the article, it must not contain results which are not presented and substantiated in the main text and should not exaggerate the main conclusions. Glacial relicts, especially those with very narrow habitat requirements, are particularly affected by global warming and thus good models for studying the future biodiversity patterns of the Alps. We have used as a model taxon *Papaver occidentale*, a glacial relict endemic to the Western Prealps, belonging to the alpine poppy complex (*P. alpinum* aggr.). All known localities were visited, each population was georeferenced and the number of individuals was estimated. Species Distribution Modelling (SDM) was used to evaluate the present and future potential distribution range and habitat suitability, taking into account the specificity of its habitat (calcareous screes). According to our study, there are globally 19 natural populations of *P. occidentale*. The total number of individuals was estimated at 30756 (with 72% of all individuals in canton of Bern). The taxon is a highly specialized alpine plant growing preferentially between 1900 and 2100 m a.s.l. on north facing screes. Predictions for the end of the 21st century indicate that suitable area will significantly decrease, for both the entire studied area (0-30% remaining) and sites nearby current *P. occidentale* populations (0-17% remaining). Under the most severe scenario, the species risks complete extinction. The long-term *in situ* conservation of *P. occidentale*, and all other taxa of the *P. alpinum* complex, is unlikely to be achieved without slowing global climate change. More generally, our fine-scale study shows that local environmental buffering of large-scale climate change in high-mountain flora may be very limited in specialised taxa of such patchy environments as screes.

Keywords: Arctic-alpine flora, Extinction due to climate change, Narrow endemism, *Papaver alpinum* aggr., Scree species, Species distribution modelling

1. Introduction

Understanding how species respond to climate change is a major challenge of conservation biology [1]. This is especially the case for high mountain plants, and more specifically glacial relicts inhabiting European Alps [2-4]. Relicts are remnants of past populations that have become fragmented by climate-driven changes and habitat loss [5]. These remnants were left behind during past range shifts and can persist today only in enclaves with suitable environmental conditions in areas with inhospitable regional climates [6]. The Alps, along with the neighboring mountain ranges, played an important role in forming the biogeographical patterns in Europe and acted as a refugium for many taxa throughout several ice age cycles [7-12]. During recent decades, the glacial and postglacial histories of arctic-alpine plant species have been extensively investigated [13-18]. However, many relict taxa of the alpine region still await thorough analysis since only a few punctual studies have explored the conservation status of the glacial relicts in Europe [19-20]. Similarly, although glacial relicts are expected to respond dramatically to climate warming, very few studies exist on their reaction to global warming in their current refugia [1,21].

In our study, we investigated a typical glacial relict *Papaver occidentale* (Markgr.) H.E. Hess et al. (Papaveraceae) (Fig. 1). The species is a narrow endemic of the Western Prealps, growing in the alpine zone in plant associations typical of calcareous screes (*Thlaspion rotundifolii*) [22-25]. *Papaver occidentale* belongs to the monophyletic section *Meconella* including ca. 50 poppy species, that have mainly circumpolar arctic distribution with some geographically isolated species distributed in the alpine zone of temperate regions [26,27]. This group hosts some of the most northerly-growing plants in the world, like *Papaver radicum*, which grows at latitudes up to 83° North [28,29]. More specifically, *P. occidentale* belongs to the alpine poppy complex *Papaver alpinum* s.l., which is distributed in small isolated regions all across the Alps and neighbouring mountain ranges [24,30,31]. The *P. alpinum* group is a very complex taxon, perfectly illustrating complicated

evolutionary and biogeographic history of the Alps [31]. Because of the highly polymorphic character of *P. alpinum* complex, its taxonomic division is not fully resolved [31]. Nevertheless, it was demonstrated that *P. occidentale* forms a natural and probably monophyletic group with three other white-flowered and narrow-lobed alpine poppies, that are confined to calcareous bedrocks: *P. alpinum* s.str., *P. sendtneri* (both north-eastern Alps), and *P. tatricum* (Tatra mountains in the Carpathians) [30-34]. The recolonization history of the Prealps by this taxon after the last glaciation was complex and the present genetic structure of *P. occidentale* is a result of a mixture of local survival and immigration [34]. Despite of its priority status in many regions of Switzerland and France [35] and its symbolic value to some regions [36], the plant was never studied across its entire distribution area. Thus, besides of very few local and technical reports [37-39], many aspects of its biology and biogeography (e.g. chorology, phylogeography, population genetics, sensitivity to global change) have not been studied at all.

Papaver occidentale grows exclusively on steep screes and in inaccessible summit parts of the alpine zone [36]. Thus, its habitat is usually not disturbed by any direct human activity (Fig. 1). However, the plant is perennial and stenothermic, making it vulnerable to climate change [38]. Species Distribution Modelling (SDM) combines species current natural occurrence data and environmental variables to estimate the geographic range of suitable habitats in the past, present, and future [40-43]. It has been widely applied in conservation biology to predict potential shifts in geographic ranges in response to climate changes [44]. However, the predictions and using of SDM for alpine plants are challenging due to a complex topographic and edaphic pattern of the high-altitude areas [45]. This is even more the case for plants inhabiting very specific alpine habitats, such as calcareous screes, which is the case of *P. occidentale*.

The present study aims thus to answer the following questions: (1) What is the present global distribution of *Papaver occidentale*? (2) What are the population sizes in different regions across its distribution area? (3) What are the general

ecological characteristics of its populations? Furthermore, we aimed to apply the Species Distribution Modelling (SDM) based on the present occurrence data, as well as on climatic and ecological variables in order to (4) evaluate the present potential distribution range, and (5) predict the location of suitable habitats of *P. occidentale* in the future, considering not only various climate change scenarios but also its unique habitat specificity. Finally, based on our findings, we discuss the conservation implications of our results for the model taxon as well as for other relict plants, especially those highly specialized for alpine scree.

2. Materials and Methods

2.1. Study area and data collection

The studied area covered the whole distribution of *Papaver occidentale*, namely the Western Prealps between Switzerland and France. Information about all known locations of *P. occidentale* was obtained from Info Flora (www.infoflora.ch), as well as elucidated from the literature [32,36], from the unpublished reports [38,39] and from personal communications of local experts (B. Clément from the Botanical Garden of the University of Fribourg and D. Jordan from France). The populations from the Swiss canton of Lucerne were identified as belonging to *P. sendtneri* [34], and were not taken into consideration in the present study. All information was gathered, duplicates as well as non-precise locations were removed, and the remaining information was used to conduct the populations sampling.

All populations of *P. occidentale* were visited and investigated between July and September 2018 (Figures 1 and 2, Table S1, Suppl. Mat.). The location of each population was georeferenced directly using GPS. The number of individuals was estimated as follows: for large populations (> 100 individuals), several GPS locations were recorded (10 to 50 m distance between points) with each time the estimation of the number of individuals. For small populations (< 100 individuals) one GPS

location was measured and all individuals counted. Because of the field characteristics (moving screes), it was not possible to use a standard procedure (like a grid pattern) to collect the data. Proximity of high cliffs and strong slopes could have decreased the GPS precision, so all GPS locations were checked afterward on a GIS software with orthophotos, and corrected if necessary (final horizontal error < 20 m). For species distribution modelling, the number of estimated individuals was used as presence data (generation of “presence points” in a delimited area around each GPS location).

2.2. Environmental variables

The statistical analyses and species distribution modelling (SDM) was conducted in the Swiss Western Prealps, from the Lake of Geneva to the Lake of Thun (the centre of *P. occidentale* distribution). For the SDM, a selection of environmental variables, which were evaluated as the most likely to play a role in *P. occidentale* distribution, has been done (topographic, climatic and land cover). The most highly reliable data for the studied area, with the finest available resolution, were used. These data were gathered from different sources and the final resolution of environmental variables used for the SDM was 10 m (Table 1, Table S2, Suppl. Mat.).

Climatic and topographic variables are the basis of SDM, but landcover (e.g. soil, geology) plays also an essential role, mainly for plants [46]. As *P. occidentale* is exclusively linked to moving calcareous screes, this azonal factor plays an essential role to model its distribution. As no exhaustive data exists, we developed this layer ourselves, by delimiting area mainly using orthophotos and in-field knowledge.

The variable EASTNESS (west-east slope aspect) was excluded because it was not estimated as relevant. Moreover, it is biased as the mountain chains are mainly oriented from SW to NE in the studied area. Indeed, as *P. occidentale* occurs mainly in north exposed screes (Fig. 3), they are in the studied area mainly NW exposed and

EASTNESS appears to be significant. But we assumed that it is just a bias and that this variable can't play a role to explain the niche of *P. occidentale*.

The variable ELEVATION was not used in the model because it is an indirect variable [42,47]. It was substituted by the temperature (TEMP), which is a direct variable and more suitable to make predictions with future climatic scenarios. The original data for temperature (1 km resolution) was highly correlated with ELEVATION (R-squared value = 0.92, p-value < 0.001). TEMP was thus calculated as a function of ELEVATION, to downscale the resolution to 10 m.

We checked for multicollinearity problems between variables, which is assumed as an important source of model uncertainty. It is safer to avoid selecting highly correlated variables for SDM [47-50]. The variance inflation factor (VIF) was calculated and we looked at paired correlation plots and r-squared values [47,51,52]. With the retained variables, all VIF scores were under 5, (recommendation for value below 10, [47]) and paired correlation R-squared values didn't exceed 0.6 (Figures S1, S2, S3, Suppl. Mat.), which is below an acceptable threshold of 0.8 [53].

For SDM, pseudo-absence data (or background data) were generated to obtain the available environmental conditions in the studied area. The number of pseudo-absences was selected to equals the number of presences. Pseudo-absences were randomly generated, which is the strategy with the least assumptions that should be used by default if there are no strong argument in favour of a different approach [47,54,55]. Pseudo-absences have been first generated in the entire studied area. With this method, it appears in all models that SCREES was the most important variable, that explained above all the distribution of *P. occidentale* and largely hide other variables effects. As *P. occidentale* occurs only in calcareous screes, pseudo-absences were generated only inside the delimited area with screes (where SCREES = 1). Consequently, SCREES was removed from variables in the models. The pseudo-absences were randomly generated in this restricted area, to the same number as the presences.

For future prediction under different climate change scenarios, we gathered raster data from Meteoswiss ([56], Table S3, Suppl. Mat.). We used the data that includes evolution of temperature and precipitation (annual, summer, winter) for the end of the century (sometime between year 2070 and 2099, noted hereafter year 2085). Three different RCP (representative concentration pathway) scenarios were considered: RCP 2.6 (implies strong reduction of greenhouse gas emissions early in the 21st century), RCP 4.5 (emissions decline after 2050, stabilization of radiative forcing) and RCP 8.5 (continuously increasing radiative forcing) [56,57].

On average, the annual mean temperature is expected to increase in Switzerland [56], from + 1.3 °C (RCP 2.6) to + 4.4 °C (RCP 8.5). The actual mean precipitation in winter (PREC-Win data) was also replaced by the raster projections for 2085. On average, the mean winter precipitations are expected to increase in Switzerland from + 6 % (RCP 2.6) to + 13 % (RCP 8.5). On the contrary, summer precipitations are expected to decrease (but this variable was not retained in the final validated model).

2.3. Model selection and calibration

Two logistic regression techniques were used for SDM: the general linear models (GLM) and the general additive models (GAM). GLM are a generalization of ordinary linear regression. They allow the linear model to be related to the response variable via a link function [58]. For our purpose, the link function was the logistic function (logit) and we used quadratic functions of all environmental variables to allow for nonlinear relationships between the covariates and response variable. GAM are GLM in which the usual linear relationship between the response and covariates are replaced by several nonlinear smooth functions to model and capture the nonlinearities in the data [59-61].

All preselected environmental variables were first included in the models. Then, the significance of the variables in the model, the explained deviance, the AIC, Akaike information criterion and BIC, Bayesian information criterion [62,63] were

checked to remove some superfluous variables and build to the most parsimonious model.

2.4. Model validation

Different cross validation techniques have been used to validate the model and prevent overfitting, like 10-fold cross validation [47,64,65] and Monte-Carlo cross validation [66,67]. However, our data were strongly spatially autocorrelated. Consequently, the 10-fold and the Monte-Carlo cross validations were not suitable to validate our models (for example, the ROC curves of validation datasets were always quasi identical to the ROC curve of the full model, see Fig. S4, Suppl. Mat.). We therefore designed a spatial cross validation [68,69]. The studied area was split into two blocks on the W-E axis. The data within one block serves to train the model while the data in the second block serves to validate the model, and vice versa. Then the same exercise was done with two blocks on the N-S axis. Presences and pseudo-absences were each time equilibrated. This validation procedure serves us mainly to delete environmental variables from the model that conduct to overfitting.

The results of cross validation were analysed with a ROC curve [70]. The true positive rate was plotted against false positive rate for each validation dataset, next to the full model ROC curve, which permits to investigate for overfitting problems. With validation datasets, the area under the curve (AUC) and mean squared residuals (MSR) served to evaluate the model performance.

The spatial cross validation revealed a strong overfitting of the models (both GLM and GAM). The models were thus simplified step by step (suppression of some environmental variables, decreasing of the degree of smoothing in GAM or suppression of some quadratic terms in GLM), to optimize the dataset during the spatial cross validation. We lastly also tried to add interaction terms, but they never improved the model and were thus not included. All calculations and graphs were done in R ([71], Table S4, Suppl. Mat.). Maps and spatial analyses were created with QGIS [72].

3. Results

3.1. Distribution and population size of *Papaver occidentale*

The species could be confirmed in all sites indicated in the available sources (e.g., old and recent literature, reports, personal communication of local experts), and no new populations were discovered. *Papaver occidentale* possesses globally 21 populations (Figures 1 and 2, Table S1, Suppl. Mat.) growing in six administrative units: 6 populations in canton of Fribourg, 5 in Bern, 4 in Vaud, and 1 in Valais, as well as 5 populations in Haute-Savoie (France). Two populations of the canton of Vaud from the Jura Mountains are introduced (TEN and CUN, ancient introductions, [38]). Thus, the majority of 19 natural populations (74%) are growing in Switzerland.

The total population size of natural occurrences of *P. occidentale* was estimated at 30756 individuals (estimation from 153 GPS locations). The two introduced sites in the Jura Mountains are very poor in individuals (TEN: 6, CUN: 1). The largest population grows in canton of Bern (Hinderi Fromatt), with 9851 individuals. Generally, canton of Bern with 22117 (72%) of all individuals taken together, is the main distribution center of the taxon. Canton of Fribourg has 5167 (17%), Vaud 748 (2.5%) and Valais 141 (0.5%) of individuals. Haute-Savoie, and thus France, counts 2547 individuals, and thus merely 8% of all individuals of the species (Figures 1 and 2).

The species is growing preferentially between 1900 and 2100 m a.s.l. and on the northern slopes (Fig. 3, Table S1, Suppl. Mat.), usually between 100 and 300 meters from the highest summit. The highest mean elevation of *P. occidentale* occurrence (2400 m a.s.l.) was measured in the population on Pointe Blanche (PTB) in Haute-Savoie as well as in canton of Fribourg, with 2300 m a.s.l. on Pointe de Paray (PAR) and with 2235 m a.s.l. at Combe Vanil de l'Écri (COM). The natural populations with the lowest mean altitude occur in canton of Vaud, with 1760 m a.s.l. in Les Salaires (SAL) and with 1653 m a.s.l. in La Pierreuse (PIE). The two very small

and introduced populations of Jura are growing also at very low altitudes (TEN: 1658 m a.s.l and CUN: 1571 m a.s.l.) (Table S1, Suppl. Mat.).

3.2. Species distribution modelling

A total of 28209 individuals (presences) were comprised in the area considered for the SDM (representing 92% of all individuals and 74% of all natural populations). Thus, the number of pseudo-absences randomly generated (in the area with calcareous screes, see methods), was 28209 as well.

Some environmental variables have been directly removed to avoid multicollinearity problems. The eight remaining variables included in the first models were the following: SLOPE, NORTHNESS, TEMP, PREC_Sum, PREC_Win, SUNSHINE, IRRAD, HOURS_SUN. The eight included variables were significant in GLM and GAM, and were relevant according to AIC and BIC. Only HOURS_SUN was removed from GLM as it increased the AIC. The cross validation procedure revealed a strong overfitting of the model. Some environmental variables were removed consequently (see methods).

The final validated model chosen to predict *P. occidentale* habitat suitability was a GAM including the following four environmental variables: SLOPE, TEMP, NORTHNESS, PREC_Win. It explained 58.64 % of the total variance. The degree of smoothing (k) of the GAM was 2 for all variables. A GLM with the same environmental variables was very similar to the GAM, but the GAM was slightly more performant during the spatial cross validation, explaining why the GAM was finally selected. The mean AUC of the validation datasets during the spatial cross validation was 0.9. The four included variables were significant (P-value < 0.001). The best threshold calculated from the ROC curve was 0.55 (Fig. S1, Table S5, Suppl. Mat.).

3.3. Predicted habitat suitability and ecological preferences

The above model was used to predict the current habitat suitability for *P. occidentale* on the studied area (Fig. 4). In the studied area (2335 km²), the habitat can be considered as suitable (predicted value > 0.55) for *P. occidentale* on 33 km² (1.4 %). The species effectively occurs in area of about 1 km².

The most important environmental variables that influence the distribution of *P. occidentale* are those included in the final model; namely, the slope (SLOPE), the mean annual temperature (TEMP) and the slope orientation (NORTHNESS). More surprisingly, the mean precipitations during the winter (PREC_Win) also contributes to explain the species distribution. The response curve of these four variables can be found in Figure S5 (Suppl. Mat.). The ecological preferences of *P. occidentale* concerning some selected environmental variables are shown in Fig. 5.

A small percentage of *P. occidentale* individuals were growing on south facing slopes, however only above 1900 m a.s.l. Below 1900 m a.s.l. all individuals were only growing on north facing slopes. Above this elevation, the slope aspect becomes less important (Fig. 3).

The model was used to predict habitat suitability of *P. occidentale* under different climatic change scenarios (RCP 2.6, RCP 4.5 and RCP 8.5, see Methods), for the end of the century (year 2085, (Table 2, Fig. 6). For the calculations, the actual mean annual temperature (TEMP data) was replaced by the raster projections for 2085. Finally, the remaining suitable area at the end of the 21st century was calculated for each climatic scenario. The suitable area will be drastically reduced, both for all populations of *P. occidentale* in the study area, and for all climatic scenarios. For the entire studied area reduction to 0-30% is predicted, and to 0-17% for the sites nearby of the current *P. occidentale* populations. Under the most severe scenario (RCP 8.5), the species risks complete extinction, and this not only for small isolated population but also for all five individual-reach and large-surface populations of Spillgerte in the canton of Bern (Table 2, Fig. 6).

3.4. Figures and Tables

Figure 1. *Papaver occidentale*. (a) Individual plant forming numerous flowers; (b) Close up of the flower; (c) Typical population growing in a calcareous scree (Spillgerte, canton of Bern); (d) Typical habitat, north-facing screes and summits (Spillgerte, canton of Bern); (e) Each tuft is formed by one individual, anchored in the mobile scree with one large and long pivotal root. Plants reproduce exclusively sexually (no vegetative reproduction) and reach the maximum age of ca. 8 years.




Figure 2. Location of the 22 populations of *Papaver occidentale*, with their corresponding name and number of individuals (size of dots reflect the population size). The shaded zone indicates the area considered for SDM.




Figure 3. (a) Radial histogram indicating the number of *Papaver occidentale* individuals according to the slope aspect. The small surrounding ticks indicates single individuals and the black solid line gives a smooth approximation, calculated with a GAM. (b) Northness (N-S slope aspect, values from -1: south to 1: north) as a function of elevation (meters), for all *P. occidentale* individuals in the studied area (grey dots). The black line shows a smooth approximation of the mean (GAM fit, with 95% confidence interval).




Figure 4. Modeled current habitat suitability of *Papaver occidentale* in the Western Swiss Prealps. White areas are the less suitable, red and purple are the most suitable. Three areas are zoomed on the right panel. The black dots indicate the locations of *P. occidentale* (presences).




Figure 5. Ecological preferences of *Papaver occidentale* concerning some selected environmental variables. *P. occidentale* individuals are represented by red dots and background (pseudo-absences) are represented by grey dots. The box plots summarize the preference of *P. occidentale* concerning each variable. ELEVATION and IRRAD are shown but were not included in the final model.




Figure 6. Modeled future habitat suitability of *Papaver occidentale* in the Western Swiss Prealps, under different climatic change scenarios (RCP 2.6, RCP 4.5 and RCP 8.5), at the end of the century (year 2085). White areas are the less suitable, red and purple are the most suitable. Two areas are zoomed on the right panel. Black dots indicate location of present *P. occidentale* populations.




Table 1. Environmental variables initially selected for Species Distribution Model (SDM) of *Papaver occidentale*.

Abbreviation	Variable type	Unit	Source	Native resolution	Resolution used in SDM
ELEVATION	Digital elevation model (DEM)	Meters (m)	swissALTI3D, The high precision digital elevation model of Switzerland [90]	2 m, vertical uncertainty $\pm 0.5 \text{ m} < 2000 \text{ m}$, 1–3 m $> 2000 \text{ m}$	Upscaled to 10 m, median value
SLOPE	terrain inclination	Degrees (°)	Calculated from DEM	2 m	Upscaled to 10 m, median value
NORTHNESS	Slope aspect on the south-north axis	No unit, a real number between -1 (south) and 1 (north)	Calculated from DEM, cosinus of the slope aspect	2 m	Upscaled to 10 m, median value
EASTNESS	Slope aspect on the west-east axis	No unit, a real number between -1 (west) and 1 (east)	Calculated from DEM, sinus of the slope aspect	2 m	Upscaled to 10 m, median value
TEMP	Mean yearly mean temperature (norm, 1981–2010)	degrees °C	TnormY8110 [93]	1000 m	Downscaled to 10 m, using DEM to improve resolution (strong negative linear relationship between ELEVATION and TEMP, R-squared value = 0.92, p-value < 0.001 ***. TEMP was thus calculated as a function of ELEVATION, see Fig. S1, Suppl. Mat.)
TEMP_Sum	Mean summer mean temperature (norm, 1981–2010)	degrees °C	TnormM8110 mean for June, July and August [93]	1000 m	Downscaled to 10 m, same procedure as TEMP
TEMP_Win	Mean winter mean temperature (norm, 1981–2010)	degrees °C	TnormM8110, mean for December, January and February [93]	1000 m	Downscaled to 10 m, same procedure as TEMP
PREC	Mean yearly precipitation (norm, 1981–2010)	Milimeters (mm)	RnormY8110 [93]	1000 m	Downscaled to 10m, using bicubic spline interpolation to get a smooth result

PREC_Sum	Mean summer precipitation (norm, 1981-2010)	Milimeters (mm)	RnormM8110, mean for June, July and August [93]	1000 m	Downscaled to 10m, using bicubic spline interpolation to get a smooth result
PREC_Win	Mean winter precipitation (norm, 1981-2010)	Milimeters (mm)	RnormM8110, mean for December, January and February [93]	1000 m	Downscaled to 10m, using bicubic spline interpolation to get a smooth result
SUNSHINE	Mean yearly relative sunshine duration (norm, 1981-2010)	Percent (%)	SnormY8110 [93]	1000 m	Downscaled to 10m, using bicubic spline interpolation to get a smooth result
IRRAD	Total irradiance during one day at the solstice (21 of June)	Wh / m2 /day	Calculate from DEM with r.sun (Solar irradiance and irradiation model) in GRASS GIS [92]	10 m	10 m
HOURS_SU N	Hours of sun during one day at the solstice (21 of June)	Hours (h)	Calculate from DEM with r.sun (Solar irradiance and irradiation model) in GRASS GIS [92]	10 m	10 m
SCREES	Landcover : area with presence of Calcareous screes	Presence-absence (1-0)	Area delimited using GEOCOVER (Geological vector datasets for better subsurface management, [91]) and interpretation of orthophotos. A 100 m buffer was added to the delimited areas, as the precise limit of the scree is difficult to get.	NA, vector data.	Rasterized to10 m

Table 2. Remaining suitable area (predicted value > 0.55) under different climatic change scenarios (RCP 2.6, RCP 4.5 and RCP 8.5), at the end of the century for *Papaver occidentale*. The entire studied area was considered (first column) or only the areas nearby actual *P. occidentale* populations, within a radius of 5 km (second column). The remaining futures suitable areas are given as a percentage of the actual suitable predicted area.

Climatic change scenario	Remaining suitable area (entire studied area)	Remaining suitable area (nearby <i>P. occidentale</i> populations)
Actual	100 %	100 %
RCP 2.6, year 2085	30 %	17 %
RCP 4.5, year 2085	4 %	0.3 %
RCP 8.5, year 2085	0 %	0 %

4. Discussion

Our study demonstrates that *P. occidentale* is a highly specialized, cold-adapted alpine plant. It grows preferentially between 1900 and 2100 meters of altitude, and all populations and individuals that grow under 1900 meters a.s.l. are strictly confined to north-facing screes. Although some populations (e.g. in canton of Bern) show locally relatively high number of individuals, the present distribution pattern and the extremely small effectively occupied area, left no doubt: *P. occidentale* likely reaches today the strongest reduction of its distribution range in the evolutionary history of this taxon.

Papaver occidentale, along with all members of the *P. alpinum* complex, belongs to the section *Meconella*, and shows a large number of morphological and ecological similarities with other members of this taxonomic group [30]. The *Meconella* poppies (ca. 50 species) are an evolutionary young and taxonomically poorly understood branch, mainly due to a Pleistocene or perhaps even later speciation of the majority of its members [73,74]. All species of this section are specialized arctic-alpine poppies, adapted to survive in the most harsh (edaphically and climatically) ecosystems inhabited by vascular plants [75]. However, the *P. alpinum* group is the most basal in the sect. *Meconella* [26,27] and some authors proposed that it colonized the Alps rather from the east than from the north [31].

Nevertheless, *Meconella* poppies were among the very first immigrants during the late glacial period in the high Arctic [76]. Notably, *P. radicum* was probably among the earliest colonizers of deglaciated parts of northern Norway between the 13,000 and 14,000 years BP [28]. *Papaver radicum* is also today nearly the only vascular plant (along with *Saxifraga oppositifolia*) inhabiting the northernmost piece of land in the Northern Hemisphere (islet of Kaffeklubben, northern Greenland) [75]. Another species of the *Meconella* section, *P. dahlianum*, forms on Svalbard floristically poor communities and is inhabiting similar habitats to *P. occidentale* (scree-covered mountain slopes), but also gravel beds and moraines. The species is also among the very first immigrants growing on moraines left behind the retracting glaciers on

Svalbard [76]. Furthermore, *P. dahlianum* makes part of a restricted number of vascular plants capable of growing, flowering and yielding seeds at the average daily temperature slightly over 0°C [77], and reaching on Svalbard altitude of 940 m a.s.l. [78].

In northeastern and southeastern Alps, individual plants and even large populations of *P. alpinum* s.l. were observed in gravel river beds and road sides at low altitudes [31]. Therefore, it is highly probable, that during the last glacial period, ancestors and/or relatives of *P. occidentale* were frequent in the vast alluvial terraces surrounding the central glaciated part of the Alps. They then followed the retracting glaciers and survived only in very few isolated and extremely small areas [30]. However, recent molecular data indicate, that both local survival and immigration from edge alpine refugia played an important role during the recolonization of the Western Prealps by *P. occidentale*, leading to a complex genetic pattern [34]. *Papaver occidentale* is clearly in refugial situation today.

4.1. Current geographic range

According to SDM, the habitats that can be considered as suitable for *P. occidentale* in the studied area reach about 33 km². However, the effective area where the species occurs does not exceed 1 km². There are likely two main reasons why the realized niche is much more restricted than the fundamental niche. Firstly, the history of the recolonization of the Alps by the species after the last glacial maximum is complex. The species probably followed the retracting glaciers in some valleys on the moraines, but not in all of them. With the Holocene climate warming, the species was blocked in these valleys and was unable to reach other parallel valleys because of its limited dispersion potential. It moreover probably became extinct in some areas. The high genetic isolation of some populations, geographically close, could support such a scenario or a possible local in situ survival without postglacial expansion [1].

The second reason is the method of the delimitation of calcareous screes, which could lead to a certain bias. Some delimited screes on orthophotos were included in the studied area, but might be not suitable for the species (scree already too stabilized, concurrent vegetation, etc.). This lead to an overoptimistic suitable area, but the percentage of reduction of those suitable areas under future climatic change scenarios is proportional, and thus reliable.

4.2. No future for *P. occidentale*?

Thuiller et al. [79] estimated that up to 48.5% of mountain plant species in the European Alps are at risk of extinction due to the accelerated climate change. *Papaver occidentale* belongs without doubts to the group of alpine plants, which are the most exposed to the recent global warming. According to our results, the remaining suitable area will be drastically reduced, even by the weakest global warming scenario. By the RCP 8.5 climate change scenario *P. occidentale* will face a complete extinction by 2085 across its whole distribution area. The severe future range contraction predicted for *P. occidentale* reflects similar results for other alpine endemic species, e.g. *Berardia subacaulis* [80] or *Lilium pomponium* [81]. It was often concluded, that global warming may result in a significant upward shift in species optimum elevation [82], giving the threatened species an additional survival option. However, marginal Western Prealps have relatively low summit heights, making their alpine plant populations particularly vulnerable to extinctions [25,83]. Additionally, *P. occidentale* is confined to calcareous scree habitats, which will not enlarge their surface till the end of the 21st century. Although exceptionally at highest elevations, isolated individual plants can be found growing in weathered crevices of calcareous rocks, an altitudinal shift does not appear to be a viable option for this taxon.

Several studies demonstrated that alpine plants' response to climate warming might be much slower than predicted using standard species distribution models [84,85]. Additionally, even highly specialized and/or endemic plants can defy

unfavorable climatic conditions and persist for millennia in so-called topo-climatic microrefugia, as was demonstrated for *Saxifraga florulenta* in the Alps [86] or a relict lineage of *Ranunculus glacialis* in the Carpathians [16]. In extreme cases, exiguous spots can ensure a long-term survival of isolated relict populations [87]. Our study however, is a rare example of SDM specifically developed for plants inhabiting calcareous screes (e.g. not including other non-relevant landcover variables), enhancing thus its precision and reliability. It is very probable that plants inhabiting calcareous screes will react faster than others plant communities (e.g. subalpine grasslands, [88]).

4.3. Conservation implications

Only 19 populations grouped in six isolated regions form the entire native distribution area of *P. occidentale*, with a total number of individuals estimated at slightly more than 30000. Surprisingly, the region of Spillgerte (comprising all 5 populations of the canton of Bern), harbors more than 72%, and thus a huge portion of the total number of individuals. Such a domination in this region is an important discovery of the present study, since the only reports on *P. occidentale* available to date, attributed regions of Fribourg and Vaud as the main distribution centers of this taxon [38]. Furthermore, the five populations of the canton of Bern represent a unique genetic composition, being highly differentiate in comparison with all the other populations [34]. These results should be therefore taken into consideration in redistribution of the conservation responsibilities and priorities for the concerned regions and administrative units. The introduced plants in Jura Mountains grow in a habitat that is very different from the natural ones [38,88,89]. The very low number of individuals in the introduced region, detected in our survey, confirms that *P. occidentale* has very specific habitat requirements and cannot prosper in any other habitat. Thus, no conservation efforts should be undertaken for these non-natural populations.

Despite its quite narrow distribution and its very specialized environmental requirements, the *P. occidentale* populations are not threatened by any direct human activity in a short term. The steep and mobile calcareous screes are habitats which are not appropriate for any human activity. Merely few hiking trails get close to populations [38]. Wild animals do not put the species under extinction risk neither. Ibexes and chamois have been observed in the vicinity but only few individuals have been reported as grazed [39]. Thus, according to our study, global warming will be the main (if not the only) factor threatening the last remaining populations of *P. occidentale* in the near future.

Our results confront local nature conservation authorities with major challenges. The long-term *in situ* conservation of *P. occidentale*, and probably all other taxa of the *P. alpinum* complex, as well as other glacial relicts in the Alps, is unlikely to be achieved without slowing global climate change.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1,

Table S1. Populations of *P. occidentale* and estimated population sizes (number of individuals).

Table S2. Additional analyses related to SDM.

Table S3. Information on climate change scenarios data.

Table S4. R script.

Table S5. Cross validation of the final validated GAM: AUC and MSR of the validated model, with different cross validation techniques.

Figure S1. Series of maps representing environmental variables on the studied area.

Figure S2. Paired correlation plot with all environmental variables.

Figure S3. Paired correlation plot with the 8 selected (uncorrelated) environmental variable.

Figure S4. Roc curves for different cross validation techniques (red line, full model ROC curve with all the data, grey lines, ROC curves of validation datasets).

Figure S5. Response curves for the final validated GAM.

Author Contributions: GK and YF conceived the idea. LP, SB, BC and EG conducted the field work. YF conducted statistical and modelling analyses. GK, MR and CP edited the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: MR is financially supported by the statutory funds of the W. Szafer Institute of Botany, Polish Academy of Sciences.

Acknowledgments: We would like to thank Peter Wandeler from the Natural History Museum in Fribourg (Switzerland) for his advice and the financial support, Denis Jordan (France), Sarah Kasraian, David Giovannini and Beat Wolf. Finally, we thank Francesca Cheda from the Nature Protection Office of the Canton Fribourg (Switzerland), Brigitte Holzer from the Nature Protection Office of the Canton Bern (Switzerland), Olivier Guex from the Service des forêts, des cours d'eau et du paysage of the Canton Valais (Switzerland) and Franco Ciardo from the Direction générale de l'environnement of the Canton Vaud for the permissions.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1 Jiménez-Alfaro, B.; García-Calvo, L.; García, P.; Luis Acebes, J. Anticipating extinctions of glacial relict populations in mountain refugia. *Biol. Conserv.* **2016**, *201*, 243-251.
- 2 Arrigo, N.; Bétrisey, S.; Graf, L.; Bilat, J.; Gerber, E.; Kozłowski, G. Hybridization as a threat in climate relict *Nuphar pumila* (Nymphaeaceae). *Biodiv. Conserv.* **2016**, *25*, 1863-1877.
- 3 Zecca, G.; Casazza, G.; Piscopo, S.; Minuto, L.; Grassi, F. Are the responses of plant species to Quaternary climatic changes idiosyncratic? A demographic perspective from Western Alps. *Plant. Ecol. Divers.* **2017**, *10*, 273-281.
- 4 Schönswetter, P.; Schneeweiss, G.M. Is the incidence of survival in interior Pleistocene refugia (nunataks) underestimated? Phylogeography of the high mountain plant *Androsace alpina* (Primulaceae) in the European Alps revisited. *Ecol. Evol.* **2019**, *9*, 4078-4086.
- 5 Hampe, A.; Jump, A.S. Climate relicts: past, present, future. *Annu. Rev. Ecol. Evol.* **2011**, *42*, 313-333.
- 6 Woolbright, S.A.; Whitham, T.G.; Gering, C.A.; Allan, G.J.; Bailey, J.K. Climate relicts and their associated communities as natural ecology and evolution laboratories. *Trends Ecol. Evol.* **2014**, *29*, 406-416.
- 7 Taberlet, P.; Fumagalli, L.; Wust-Saucy, A.-G.; Cosson, J.-F. Comparative phylogeography and postglacial colonization routes in Europe. *Mol. Ecol.* **1998**, *7*, 453-464.
- 8 Brochmann, C.; Gabrielsen, T.M.; Nordal, I.; Landvik, J.Y.; Elven, R. Glacial survival or *tabula rasa*? The history of North Atlantic biota revisited. *Taxon* **2003**, *52*, 417-450.
- 9 Tribsch, A.; Schönswetter, P. (2003) Patterns of endemism and comparative phylogeography confirm palaeoenvironmental evidence for Pleistocene refugia in the eastern Alps. *Taxon* **2003**, *52*, 477-497.
- 10 Holderegger, R.; Thiel-Egenter, C. A discussion of different types of glacial refugia used in mountain biogeography and phylogeography. *J. Biogeogr.* **2009**, *36*, 476-480.

- 11 Berthouzoz, M.; Maendly, S.; Bétrisey, S.; Mangili, S.; Prunier, P.; Lexer, C.; Kozłowski, G. Some like it cold: distribution, ecology and phylogeny of *Arenaria bernensis* Favarger (Caryophyllaceae) from the Western Prealps in Switzerland. *Alp. Bot.* **2013**, *123*, 65-75.
- 12 Windmaisser, T.; Kattari, S.; Heubl, G.; Reisch, C. Glacial refugia and postglacial expansion of the alpine-prealpine plant species *Polygala chamaebuxus*. *Ecol. Evol.* **2016**, *6*, 7809-7819.
- 13 Stehlik, I.; Blattner, F.; Holderegger, R.; Bachmann, K. Nunatak survival of the high alpine plant *Eritrichium nanum* (L.) Gaudin in the Central Alps during the ice ages. *Mol. Ecol.* **2002**, *11*, 2027-2036.
- 14 Stehlik, I. Resistance or emigration? Response of alpine plants to the ice ages. *Taxon* **2003**, *52*, 499-510.
- 15 Paun, O.; Schönschwetter, P.; Winkler, M.; Consortium, I.; Tribsch, A. Historical divergence vs. contemporary gene flow: Evolutionary history of the calcicole *Ranunculus alpestris* group (Ranunculaceae) in the European Alps and the Carpathians. *Mol. Ecol.* **2008**, *17*, 4263-4275.
- 16 Ronikier, M.; Schneeweis, G.M.; Schönschwetter, P. The extreme disjunction between Beringia and Europe in *Ranunculus glacialis* sl. (Ranunculaceae) does not coincide with the deepest genetic split - A story of the importance of temperate mountain ranges in arctic-alpine phylogeography. *Mol. Ecol.* **2012**, *21*, 5561-5578.
- 17 Winkler, M.; Tribsch, A.; Schneeweis, G.M.; Brodbeck, S.; Gugerli, F.; Holderegger, R.; Abbott, R.J.; Schönschwetter, P. Tales of the unexpected: Phylogeography of the arctic-alpine model plant *Saxifraga oppositifolia* (Saxifragaceae) revisited. *Mol. Ecol.* **2012**, *21*, 4618-4630.
- 18 Eidesen, P.B.; Ehrich, D.; Bakkestuen, V.; Alsos, I.G.; Gilg, O.; Taberlet, P.; Brochmann, C. Genetic roadmap of the Arctic: Plant dispersal highways, traffic barriers and capitals of diversity. *New Phytol.* **2013**, *200*, 898-910.
- 19 Schmitt, T.; Muster, C.; Schönschwetter, P. Are disjunct alpine and arctic-alpine animal and plant species in the western Palearctic really "relics of the cold past"? In: Habel JC, Assman T (eds) Relict species: phylogeny and conservation biology. Springer, Berlin Heidelberg. 2010, pp. 239-252.
- 20 Zimmermann, M.; Vischer-Leopold, M.; Ellwanger, G.; Ssymak, A.; Schröder, E. The EU Habitat Directive and the German Natura 200 network of protected areas as tool for implementing the conservation of relict species. In: Habel JC, Assman T (eds) Relict species: phylogeny and conservation biology. Springer, Berlin Heidelberg. 2010, pp. 323-340.
- 21 Parisod, C.; Besnard, G. Glacial *in situ* survival in the Western Alps and polytopic autopolyploidy in *Biscutella laevigata* L. (Brassicaceae). *Mol. Ecol.* **2007**, *16*, 2755-2767.
- 22 Hess, H.E.; Landolt, E.; Hirzel, R. Flora der Schweiz und angrenzender Gebiete. Birkhäuser Verlag, Basel 2:955. 1977.
- 23 Richard, J.-L. La végétation du Vanil Noir et du Vallon des Morteys. *Bul. Soc. Frib. Sc. Nat.* **1977**, *66*, 1-52.

- 24 Aeschimann, D.; Lauber, K.; Moser, D.M.; Theurillat, J.-P. Flora Alpina. Band 1-3, Haupt, Bern, Switzerland, 2004.
- 25 Gerber, E.; Kozłowski, G.; Mariéthoz, A.-. La flore des Préalpes du lac de Thun au Léman. Rossolis, Bussigny. 2010.
- 26 Kadereit JW, Schwarzbach AE, Jork KB (1997) The phylogeny of *Papaver* s.l. (*Papaveraceae*): polyphyly or monophyly? *Pl Syst Evol* 204:75-98.
- 27 Carolan, J.C.; Hook, I.L.I.; Chase, M.W.; Kadereit, J.W.; Hodkinson, T.R. Phylogenetics of *Papaver* and related genera based on DNA sequences from ITS nuclear ribosomal DNA and plastid *trnL* intron and *trnL-F* intergenic spacers. *Ann. Bot.* **2006**, *98*, 141-155.
- 28 Nordal, I.; Hestmark, G.; Solstad, H. Reproductive biology and demography of *Papaver radicum* – A key species in Nordic plant geography. *Opera Bot.* **1997**, *132*, 77-87.
- 29 Solstad, H.; Elven, R.; Nordal, I. Isozyme variation among and within North Atlantic species of *Papaver* sect. *Meconella* (*Papaveraceae*) and taxonomic implications. *Biol. J. Linn. Soc.* **2003**, *143*, 255–269.
- 30 Kadereit, J.W. Notes on the taxonomy, distribution, phylogeny and ecology of *Papaver alpinum* L. (*Papaveraceae*) *Bot. Jahrb. Syst.* **1990**, *112*, 79-97.
- 31 Schönswetter, P.; Solstad, H.; Escobar, G.P. A combined molecular and morphological approach to the taxonomically intricate European mountain plant *Papaver alpinum* s.l. (*Papaveraceae*) - Taxa or informal phylogeographical groups? *Taxon* **2009**, *58*, 1326-1348.
- 32 Markgraf, F. *Papaveraceae*. In: Hegi G. *Illustrierte Flora von Mitteleuropa*, 2nd edition, vol. IV, 1/2, Hanser, München. 1958a, pp. 15-49.
- 33 Markgraf, F. Eine neue Gliederung der Alpenmohne. *Phyton* **1958b**, *7*, 302-314.
- 34 Pittet, L.; Fragnière, Y.; Grünig, S.; Bétrisey, S.; Clément, B.; Gerber, E.; Ronikier, R.; Kozłowski, G.; Parisod, C. Genetic structure of the narrow endemic *Papaver occidentale* indicates complex patterns of survival and immigration in the Western Prealps. *Alp. Bot.* **2020**, doi: [10.1007/s00035-020-00238-3](https://doi.org/10.1007/s00035-020-00238-3).
- 35 Bornand, C.; Gyax, A.; Juillerat, P.; Jutzi, M.; Möhl, A.; Rometsch, S.; Sager, L.; Santiago, H.; Eggenberg, S. Rote Liste Gefäßpflanzen. Gefährdete Arten der Schweiz. Bundesamt für Umwelt, Bern und Info Flora, Genf. Umwelt-Vollzug Nr. 1621. 2016.
- 36 Jordan, D. La flore rare ou menacée de Haute-Savoie. Naturalia Publications. 2015.
- 37 Bornand, C.; Hoffer-Massard, F. La Pierreuse ou le parfum du pavot, le 20 juillet 2002. *Bull. Cercle Vaud. Bot.* **2003**, *32*, 53-61.
- 38 Delarze, R. Plan d'action pour *Papaver occidentale* (Markgraf) H.E. Hess & Landolt (Pavot des Alpes occidentales). Coordination régionale pour la protection de la flore. 2012.
- 39 Bétrisey, S. Adaptation du plan d'action pour *Papaver occidentale*. Jardin botanique de l'Université de Fribourg, Suisse. 2014.
- 40 Guisan, A.; Thuiller, W. Predicting species distribution: Offering more than simple habitat models. *Ecol. Lett.* **2005**, *8*, 993–1009.

- 41 Elith, J.; Leathwick, J.R.J.R. Species distribution models: ecological explanation and prediction across space and time. *Annu. Rev. Ecol. Evol. Syst.* **2009**, *40*, 677-697.
- 42 Peterson, A.T.; Soberón, J.; Pearson, R.G.; Anderson, R.P.; Martínez-Meyer, E.; Nakamura, M.; Araújo, M.B. Ecological niches and geographic distributions (MPB-49) (Vol. 56). Princeton University Press. 2011.
- 43 Peterson, A.T.; Soberón, J. Species distribution modeling and ecological niche modeling: getting the concepts right. *Nat. Conserv.* **2012**, *10*, 102-107.
- 44 Song, Y.-G.; Petipierre, B.; Deng, M.; Wu, J.-P.; Kozłowski, G. Predicting climate change impacts on the threatened *Quercus arbutifolia* in montane cloud forests in southern China and Vietnam: Conservation implications. *Forest Ecol. Manag.* **2019**, *444*, 269-279.
- 45 Buri, A.; Grand, S.; Yashiro, E.; Adatte, T.; Spangenberg, J.E.; Pinto-Figueroa, E.; Verrecchia, E.; Guisan, A. What are the most crucial soil variables for predicting the distribution of mountain plant species? A comprehensive study in the Swiss Alps. *J Biogeogr.* **2020**, *47*, 1143-1153.
- 46 Hageer, Y.; Esperón-Rodríguez, M.; Baumgartner, J.B.; Beaumont, L.J. Climate, soil or both? Which variables are better predictors of the distributions of Australian shrub species? *PeerJ* **2017**, *5*, e3446.
- 47 Guisan, A.; Thuiller, W.; Zimmermann, N.E. Habitat suitability and distribution models: with applications in R. Cambridge University Press. 2017.
- 48 Graham, M.H. Confronting multicollinearity in ecological multiple regression. *Ecology* **2003**, *84*, 2809-2815.
- 49 Braunisch, V.; Coppes, J.; Arlettaz, R.; Suchant, R.; Schmid, H.; Bollmann, K. Selecting from correlated climate variables: a major source of uncertainty for predicting species distributions under climate change. *Ecography* **2013**, *36*, 971-983.
- 50 Júnior, P.D.M.; Nóbrega, C.C. Evaluating collinearity effects on species distribution models: An approach based on virtual species simulation. *PloS one* **2018**, *13*, e0202403.
- 51 Craney, T.A.; Surles, J.G. Model-dependent variance inflation factor cutoff values. *Qual. Eng.* **2002**, *14*, 391-403.
- 52 O'Brien, R.M. A caution regarding rules of thumb for variance inflation factors. *Qual. Quant.* **2007**, *41*, 673-690.
- 53 Menard, S. Applied logistic regression analysis (Vol. 106). Sage. 2002.
- 54 Wisz, M.S.; Guisan, A. Do pseudo-absence selection strategies influence species distribution models and their predictions? An information-theoretic approach based on simulated data. *BMC Ecol.* **2009**, *9*, 8.
- 55 Barbet-Massin, M.; Jiguet, F.; Albert, C.H.; Thuiller, W. Selecting pseudo-absences for species distribution models: how, where and how many? *Methods Ecol. Evol.* **2012**, *3*, 327-338.
- 56 CH2018 Climate Scenarios for Switzerland, Technical Report, National Centre for ClimateServices, Zurich, 2018, pp. 1-271.

- 57 Van Vuuren, D.P.; Edmonds, J.; Kainuma, M.; Riahi, K.; Thomson, A.; Hibbard, K.; Hurtt, G.C.; Kram, T.; Krey, V.; Lamarque, J.-F.; Masui, T.; Meinshausen, M.; Nakicenovic, N.; Smith, S.J.; Rose, S.K. The representative concentration pathways: an overview. *Climatic Change* **2011**, *109*, 5.31.
- 58 McCullagh, P. Generalized linear models. Routledge. 2019.
- 59 Guisan, A.; Edwards, Jr. T.C.; Hastie, T. Generalized linear and generalized additive models in studies of species distributions: setting the scene. *Ecol. Model.* **2002**, *157*, 89-100.
- 60 Hastie, T.J. Generalized additive models. In Statistical models in S. Routledge. 2017, pp. 249-307.
- 61 Wood, S.N. Generalized additive models: an introduction with R. Chapman and Hall/CRC. 2017.
- 62 Atkinson, A.C. A note on the generalized information criterion for choice of a model. *Biometrika* **1980**, *67*, 413-418.
- 63 Sakamoto, Y.; Ishiguro, M.; Kitagawa, G. Akaike information criterion statistics. Dordrecht, The Netherlands: D. Reidel 81. 1986.
- 64 Rodriguez, J.D.; Perez, A.; Lozano, J.A. (2009) Sensitivity analysis of k-fold cross validation in prediction error estimation. *IEEE Transactions on Pattern Analysis and Machine Intelligence* **2009**, *32*, 569-575.
- 65 Lever, J.; Krzywinski, M.; Altman, N. Points of significance: model selection and overfitting. *Nat. Methods* **2016**, *13*, 703–704.
- 66 Shao, J. Linear model selection by cross-validation. *J. Am. Stat. Assoc.* **1993**, *88*, 486-494.
- 67 Xu, Q.S.; Liang, Y.Z. (2001) Monte Carlo cross validation. *Chemomet. Intell. Lab. Syst.* **2001**, *56*, 1-11.
- 68 Pohjankukka, J.; Pahikkala, T.; Nevalainen, P.; Heikkonen, J. (2017) Estimating the prediction performance of spatial models via spatial k-fold cross validation. *Int. J. Geogr. Inf. Sci.* **2017**, *31*, 2001-2019.
- 69 Roberts, D.R.; Bahn, V.; Ciuti, S.; Boyce, M.S.; Elith, J.; Guillera-Aroita, G.; Hauenstein, S.; Lahoz-Monfort, J.J.; Schröder, B.; Thuiller, W.; Warton, D.I.; Wintle, B.A.; Hartig, F.; Dormann, C.F. (2017) Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography* **2017**, *40*, 913-929.
- 70 Hanley, J.A.; McNeil, B.J. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* **1982**, *143*, 29-36.
- 71 R Core Team. R: A language and environment for statistical computing. 2019.
- 72 QGIS Development Team. QGIS geographic information system. Open Source Geospatial Foundation Project. 2019.
- 73 Petrovsky, V.V. The problem of the species-width in poppies of Russian Arctic. *Skr. Norske Vidensk. Akad. Oslo* **1999**, *38*, 243–259.

- 74 PAF. Panarctic Flora Project – Annotated checklist of the Panarctic flora - vascular plants. (<http://panarcticflora.org>). Consulted on 30.01.2020. 2020.
- 75 Solstad, H.; Elven, R.; Nordal, I. Are there too many species and subspecies in the *Papaver radicum* complex? *Skr. Norske Vidensk. Akad. Oslo* **1999**, *38*, 281-294.
- 76 Dubiel, E.; Olech, M. Plant communities of NW Sörkapp Land (Spitsbergen). *Zeszyty Naukowe Uniwersytetu Jagiellońskiego* **1990**, *21*, 35-74.
- 76 Paus, A. Late Weichselian vegetation, climate, and floral migration at Sandvikvatn, North Rogaland, south-western Norway. *Boreas* **1988**, *17*, 113-139.
- 77 Dubiel, E. Ecological observations on vascular plants in the NW Sörkapp Land (Spitsbergen). *Zeszyty Naukowe Uniwersytetu Jagiellońskiego* **1991a**, *22*, 39-46.
- 78 Dubiel, E. Geobotanical problems of NW Sörkapp Land (Spitsbergen). *Zeszyty Naukowe Uniwersytetu Jagiellońskiego* **1991b**, *22*, 15-38.
- 79 Thuiller W, Lavorel S, Araujo MB, Sykes MT, Prentice IC. Climate change threats to plant diversity in Europe. *PNAS* **2005**, *102*, 8245– 8250.
- 80 Guerrina, M.; Conti, E.; Minuto, L.; Casazza, G. Knowing the past to forecast the future: a case study on a relictual, endemic species of the SW Alps, *Berardia subacaulis*. *Reg. Environ. Change* **2016**, *16*, 1035-1045.
- 81 Casazza, G.; Giordani, P.; Benesperi, R.; Foggi, B.; Viciani, D.; Filigheddu, R.; Farris, E.; Bagella, S.; Pisanu, S.; Mariotti, M.G. Climate change hastens the conservation urgency of range-restricted plant species in the central-northern Mediterranean region. *Biol. Conserv.* **2014**, *179*, 129-138.
- 82 Lenoir, J.; Gégout, J.C.; Marquet, P.A.; de Ruffray, P., Brisse, H. A significant upward shift in plant species optimum elevation during the 20th century. *Science* **2008**, *320*, 1768-1771.
- 83 Dulliger, S.; Gatringer, A.; Thuiller, W.; Moser, D.; Zimmermann, N.E.; Guisan, A.; Willner, W.; Plutzar, C.; Leitner, M.; Mang, T.; Caccianiga, M.; Dirnböck, T.; Ertl, S.; Fischer, A.; Lenoir, J.; Svenning, J.C.; Psomas, A.; Schmatz, D.R.; Silc, U.; Vittoz, P.; Hülber, K. Extinction debt of high-mountain plant under twenty-first-century climate change. *Nat. Clim. Change* **2012**, *2*, 619-622.
- 84 Cotto, O.; Wessely, J.; Georges, D.; Klonner, G.; Schmid, M.; Dulliger, S.; Thuiller, W.; Guillaume, F. A dynamic eco-evolutionary model predicts slow response of alpine plants to climate warming. *Nat. Commun.* **2017**, *8*, 15399.
- 85 Alexander, J.M.; Chalmandrier, L.; Lenoir, J.; Burgess, T.I.; Essl, F.; Haider, S.; Kueffer, C.; McDougall, K.; Milbau, A.; Nunez, M.A.; Pauchard, A.; Rabitsch, W.; Rew, L.J.; Sanders, N.; Pellissier, L. Lags in the response of mountain plant communities to climate change. *Glob. Change Biol.* **2017**, *24*, 563-579.
- 86 Patsiou, T.S.; Conti, E.; Zimmermann, N.E.; Theodoridis, S.; Randin, C. Topo-climatic microrefugia explain the persistence of a rare endemic plant in the Alps during the last 21 millennia. *Glob. Change Biol.* **2014**, *20*, 2286-2300.

- 87 Suchan, T.; Malicki, M.; Ronikier, M. Relict populations and Central European glacial refugia: The case of *Rhododendron ferrugineum* (Ericaceae). *J. Biogeogr.* **2019**, *46*, 392-404.
- 88 Aubert, S. La flore du Mont Tendre en raccourci. *Bull. Cercle Vaud. Bot.* **1952**, *2*, 25-26.
- 88 Vittoz, P.; Randin, C.; Dutoit, A.; Bonnet, F.; Hegg, O. (2009) Low impact of climate change on subalpine grassland in the Swiss Northern Alps. *Glob. Change Biol.* **2009**, *15*, 209-220.
- 89 Moret, J.-L. Les plantes introduites à la Vallée de Joux. *Bull. Cercle Vaud. Bot.* **2008**, *37*, 95-105.
- 90 Swisstopo. Bundesamt für Landestopografie swisstopo-swissALTI3D-Das hoch aufgelöste Terrainmodell der Schweiz. 2018.
- 91 Swisstopo. Metadata of the GeoCover V2 product. 2018.
- 92 Neteler, M.; Mitasova, H. Open source GIS: a GRASS GIS approach (Vol. 689). Springer Science & Business Media. 2013.
- 93 MeteoSwiss. Monthly and annual reports. Federal Office of Meteorology and Climatology. <https://www.meteoswiss.admin.ch>. 2018.