

# Automatic Segmentation of White Matter Hyperintensities from Brain Magnetic Resonance Images in the Era of Deep Learning and Big Data – A Systematic Review

Ramya Balakrishnan<sup>†1,3</sup>, Maria del C. Valdés Hernández<sup>†\*1,2</sup>, Andrew J. Farrall<sup>1,2</sup>

<sup>1</sup> Edinburgh Imaging Academy, University of Edinburgh, Edinburgh, UK

<sup>2</sup> Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

<sup>3</sup> International Institute of Health Sciences, Sri Lanka

\*Corresponding author

† Equal contribution

Correspondence to: Dr. Maria del C. Valdés Hernández

Centre for Clinical Brain Sciences

Dementia Research Institute at The University of Edinburgh, Room FU427

Chancellor's Building

49 Little France Crescent

Edinburgh, EH16 4SB, UK

e-mail: [M.Valdes-Hernan@ed.ac.uk](mailto:M.Valdes-Hernan@ed.ac.uk)

## ABSTRACT

**Background:** White matter hyperintensities (WMH), of presumed vascular origin, are visible and quantifiable neuroradiological markers of brain parenchymal change. These changes may range from damage secondary to inflammation and other neurological conditions, through to healthy ageing. Fully automatic WMH quantification methods are promising, but still, traditional semi-automatic methods seem to be preferred in clinical research. We systematically reviewed the literature for fully automatic methods developed in the last five years, to assess what are considered state-of-the-art techniques, as well as trends in the analysis of WMH of presumed vascular origin.

**Method:** We registered the systematic review protocol with the International Prospective Register of Systematic Reviews (PROSPERO), registration number - CRD42019132200. We conducted the search for fully automatic methods developed from 2015 to July 2020 on Medline, Science direct, IEE Explore, and Web of Science. We assessed risk of bias and applicability of the studies using QUADAS 2.

**Results:** The search yielded 2327 papers after removing 104 duplicates. After screening titles, abstracts and full text, 37 were selected for detailed analysis. Of these, 16 proposed a supervised segmentation method, 10 proposed an unsupervised segmentation method, and 11 proposed a deep learning segmentation method. Average DSC values ranged from 0.538 to 0.91, being the highest value obtained from an unsupervised segmentation method. Only four studies validated their method in longitudinal samples, and eight performed an additional validation using clinical parameters. Only 8/37 studies made available their methods in public repositories.

**Conclusions:** We found no evidence that favours deep learning methods over the more established k-NN, linear regression and unsupervised methods in this task. Data and code availability, bias in study design and ground truth generation influence the wider validation and applicability of these methods in clinical research.

**Keywords:** White matter lesions, white matter hyperintensities, supervised segmentation, unsupervised segmentation, deep learning, FLAIR hyperintensities

## 1. INTRODUCTION

In 1987, Hachinski, Potter, and Merskey (Hachinski, Potter & Merskey, 1987) first used the term leukoaraiosis to describe abnormal findings observed in subcortical white matter on a brain computed tomography (CT) scan. Leukoaraiosis has also been referred to as white matter lesions (WMLs) (Inzitari, 2003). With increasing use of magnetic resonance imaging (MRI) as a diagnostic tool, leukoaraiosis is increasingly referred to as white matter hyperintensities (WMH) (Wardlaw et al., 2013). WMHs are associated with reduced cognitive function, dementia, gait, balance, mobility, and mood disorders (Fazekas & Wardlaw, 2013; Zheng, Delbaere, Close, Sachdev & Lord, 2011). WMHs are also frequently observed in the asymptomatic aged and associated with common geriatric conditions such as cerebrovascular disease, cardiovascular disease, multiple sclerosis, other autoimmune diseases and psychiatric disorders such as depressive disorder, bipolar disorder and schizophrenia (Kim, MacFall & Payne, 2008; Rachmadi et al., 2018). WMH prevalence in the general population ranges from 11 – 21% in 64 year olds and increases with age to 94% in 82 year olds (DeBette & Markus, 2010). One study reported that amongst an elderly population aged 60 to 90 years, 90% have WMH (Hasan et al., 2019).

Detailed WMH evaluation for number, volume, location, and distribution on MRI may provide crucial information on aetiology, prognosis, and progression of diseases; accurate quantification may help measure treatment effectiveness (Manjón et al., 2018; Qin et al., 2018). WMH severity is considered an indirect marker of normal appearing white matter integrity and a surrogate marker of small vessel disease (SVD) (Maltais et al., 2019; Maniega et al., 2015). Advancing MRI technology means several methods have been developed to quantify WMH volumes through image segmentation: “a process which typically partitions the spatial domain of an image into mutually exclusive subsets called regions, each one of which is uniform and homogeneous with respect to some property such as tone or texture and whose property value

## White Matter Hyperintensities Segmentation Review

differs in some significant way from the property value of each neighbouring regions” (Haralick et al., 1991). However, WMH are not homogeneous, have ill-defined boundaries and their tone and texture may not significantly differ from neighbouring tissues. Biologically, they represent the “tip of the iceberg” of demyelinating, inflammatory processes which affect the whole brain: they accompany and sometimes coalesce with many neuroradiological features. Essential for digital image segmentation is recognition of edges which separate WMH from “background”. WMH identification subjectivity and boundary recognition, challenge WMH segmentation, leading to low agreement in studies of manual delineation of WMH ground truth segmentations (Akudjedu et al., 2018; Despotović, Goossens & Philips, 2015; Keller & Roberts, 2009).

Unlike normal tissues, for which validated fully automatic protocols exist and have become standard, a robust method for WMH segmentation has yet to emerge. Clinical research groups usually select a WMH segmentation method based on their own capabilities, existing methods’ specifications, availability and sustainability of the source code, and image acquisition protocols. Then, groups adapt these methods in-house and validate them for a specific study protocol. Normal tissue intensities follow a normal distribution, but abnormalities do not. In the specific case of WMH, signal intensity and spatial distributions vary, displaying unique signatures for each disease and cohort. Table 1 summarises some WMH signatures in normal ageing, SVD, Alzheimer’s disease (AD), multiple sclerosis (MS) and vanishing white matter disease (an autosomal recessive disorder) (Labauge et al., 2009). In addition to specific disease / neurological condition characteristics, WMH appearances vary widely in individuals from different disease groups (Figure 1).

## White Matter Hyperintensities Segmentation Review




Figure 1. T2-hyperintensities in MS (top row) and sporadic SVD (middle and bottom rows). Top row: Representative axial slice from two MS patients displayed, from left to right, in FLAIR, T1-weighted and T2-weighted MRI at 1.5T, showing pseudocavitated FLAIR hyperintense lesions (enclosed in rectangles). Middle row: From left to right, sagittal, coronal and axial views of a FLAIR 3T MRI scan from a patient with SVD and a high burden of WMH of presumed vascular origin. Bottom row: From left to right, sagittal, coronal and axial views of a FLAIR 3T MRI scan displaying a large confounding image artefact (enclosed in rectangles in coronal and axial views) from a patient with SVD with a low burden of WMH of presumed vascular origin.

White Matter Hyperintensities Segmentation Review

WMHs arising as a result of infections (e.g. viral, bacterial), can overlay those which already exist due to other processes (e.g., normal ageing) or comorbidities (e.g., SVD): this poses a challenge for their differential identification and segmentation. For example, in COVID-19 patients, in addition to large vessel strokes, WMHs have been reported bilaterally in the thalami, cerebellum and temporal lobes, and also in the corpus callosum, along with abnormal T2 signal in the olfactory bulb and microbleeds in the thalami (Imaging in COVID-19 complications - ESR Connect, 2020). In these brain regions, typical WMHs are uncommon. Symmetric frontal WMH and cortical hyperintensities have been reported in other COVID-19 patients with more severe respiratory disease status (MRI Shows Brain Abnormalities in Some COVID-19 Patients, 2020) along with punctate cortical blooming artefacts. But influence of treatment, comorbidities and disease severity make it difficult even for neuroradiologists to identify specific disease-related patterns that could differentially aid in diagnosis and patient stratification.

Table 1. Neuroradiological signatures of T2 WMHs in normal ageing, SVD, AD, MA and vanishing WMD.

|        | Normal ageing                                                  | SVD                                                              | Alzheimer’s disease                               | Multiple sclerosis                                                                                                                                          | Vanishing white matter disease                                                                                                   |
|--------|----------------------------------------------------------------|------------------------------------------------------------------|---------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|
| Extent | From thin peri-ventricular lining to confluent deep WM regions | From peri-ventricular with few deep WM foci to confluent regions | Thin peri-ventricular lining with foci in deep WM | From peri-ventricular with few deep WM foci to large confluent regions (may enclose “pseudo-cavities” of low T1 signal, also referred as “cavitary lesions” | Large confluent regions enclosing “pseudocavities” of low T1 signal, also referred as “cavitary lesions” (Ayrignac et al., 2016) |

## White Matter Hyperintensities Segmentation Review

|                                            |                                                                                                                                                                                                                                    |                                                                                                                                           |                                                                                           | (Ayrignac et al., 2016)                                                                                                                |                                                                                                 |
|--------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|
| <b>Characteristic brain regions</b>        | Non-specific. Disseminated throughout periventricular & deep WM & corpus striatum. Gradual progression extending from periventricular WM, from frontal & parietal regions.<br><br>Rare in temporal lobes, brain stem & cerebellum. | Non-specific. Disseminated throughout periventricular & deep WM, corpus striatum & thalami.<br><br>Rare in cerebellum.                    | Non-specific. Disseminated throughout WM & deep GM.<br><br>Rare in juxtacortical regions. | Disseminated throughout the whole brain, including midbrain, brainstem, juxtacortical regions, corpus callosum & cerebellar peduncles. | Disseminated throughout the whole brain. External capsule & corpus callosum diffusely involved. |
| <b>Symmetry between brain hemispheres</b>  | Symmetric distribution                                                                                                                                                                                                             | Symmetric distribution                                                                                                                    | Symmetric distribution                                                                    | Symmetric distribution in cerebrum, but not in cerebellum                                                                              | Symmetric distribution                                                                          |
| <b>Histogram distribution in FLAIR MRI</b> | Tail (from normal WM) fits.<br><br>Extreme Value distributions (e.g. Fréchet or Gumbel).                                                                                                                                           | Tail (from normal WM) fits.<br><br>Extreme Value distributions (e.g. Fréchet or Gumbel).<br><br>Laplacian distribution can be observed in | Very skewed independent of (i.e. separated from) that of normal WM.                       | Bimodal independent of (i.e. separated from) that of normal WM (considering cavitation).                                               | Bimodal independent of (i.e. separated from) that of normal WM (considering cavitation).        |

## White Matter Hyperintensities Segmentation Review

|                                       |                                                                                                                             |                                                                                                                                                  |                                                                       |                                                                                                                                                                                                                                                             |                                                                                                                                                                                   |
|---------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                       |                                                                                                                             | some cases<br>If/when strokes<br>are considered<br>part of the<br>WMHs                                                                           |                                                                       |                                                                                                                                                                                                                                                             |                                                                                                                                                                                   |
| <b>Signatures</b>                     | "Cotton-wool"<br>appearance.                                                                                                | "Cotton-wool"<br>appearance<br>combined with<br>subtle<br>hyperintense<br>large areas, or<br>overt &<br>confluent with<br>irregular<br>patterns. | "Cotton-wool"<br>appearance.<br>Confluent<br>regions are<br>uncommon. | Lesions<br>juxtacortically &<br>in corpus<br>callosum,<br>perpendicular to<br>ventricular<br>linings (known<br>as "Dawson<br>fingers").<br>Predominantly<br>periventricular<br>cavitary lesions.<br>Punctate<br>asymmetrical<br>posterior fossa<br>lesions. | Diffuse corpus<br>callosum<br>involvement.<br>Symmetric<br>involvement of<br>cerebellum &<br>middle<br>cerebellar<br>peduncles.<br>Predominantly<br>anterior cavitary<br>lesions. |
| <b>Presence of<br/>other features</b> | Perivascular<br>spaces (from<br>none to mild) &<br>possibly lacunar<br>lesions +/-<br>chronic mild<br>ischaemic<br>strokes. | Perivascular<br>spaces, lacunes,<br>chronic<br>ischaemic<br>strokes & few<br>microbleeds.                                                        | Marked brain<br>atrophy &<br>hippocampal<br>sclerosis.                | n/a                                                                                                                                                                                                                                                         | n/a                                                                                                                                                                               |
| <b>Borders</b>                        | Not well<br>defined.                                                                                                        | Not well<br>defined.                                                                                                                             | Better defined<br>than in SVD or<br>healthy<br>ageing.                | Better defined<br>than in SVD or<br>healthy ageing.<br>(Griffanti et<br>al.2016)                                                                                                                                                                            | Not well<br>defined.                                                                                                                                                              |



To help select WMH segmentation methods and discuss their applicability, other systematic literature reviews have been published, but on focused topics specific to diseases, e.g. MS lesion segmentation (García-Lorenzo et al., 2013; Llado et al., 2011, 2012; Miller et al., 1998; Mortazavi, Kouzani & Soltanian-Zadeh, 2012). Methods which work for MS may only perform moderately if applied to individuals with SVD or to the normal elderly (Table 1). Caligiuri et al. conducted a systematic review on fully automated methods for segmenting WMH in normal ageing and in patients with vascular pathology and risk factors, covering from 1980 – 2014 (Caligiuri et al., 2015). Two other non-overlapping reviews (Wardlaw et al., 2015; Blair et al., 2017) discussed different approaches published up to 2016, both for segmenting WMH, and also for assessing other neuroimaging markers of SVD. Another study which systematically reviewed machine-learning methods which differentiate healthy aging from different dementia types (Pellegrini et al., 2018) included studies (from 2006 to September 2016) aimed at detecting and segmenting WMH in ageing and dementia. The last five years (i.e., since 2015) have seen a boost in sample sizes, computational power and the introduction / application of deep learning in clinical research, in parallel with an increase in high-quality imaging acquisitions, facilitated by 3T MRI scanners.

We systematically reviewed the literature from 2015 to 2020 in order to assess and overview those fully automatic computational methods developed to segment WMH of presumed vascular origin.

## 2. METHODS

### 2.1. Literature Search

This systematic review protocol is registered on the International Prospective Register of Systematic Reviews (PROSPERO), registration number - CRD42019132200 (2020) to avoid

unintended duplication and to aid in transparent reporting. The search was conducted from January 2015 to July 2020 on Medline, Science direct, IEE Explore and Web of Science. For each database, we developed a search strategy to retrieve as many WMH segmentation method articles as possible. We identified keywords by expanding the subject components from the review question: white matter lesion, white matter hyperintensities, leukoaraiosis, aging, WMH, segmentation, supervised segmentation, unsupervised segmentation, machine learning, deep learning, parcellation, artificial neural network, pattern recognition, clustering, classification, magnetic resonance imaging, MRI. We applied language restriction and age limits (45 plus years) for Medline. We summarize search strategy details for each database in Appendix 1. We imported all articles retrieved into the reference manager Mendeley, and removed all duplicates. We then screened abstracts and titles to exclude studies outwith the scope of the review. Then we evaluated the full text of the remaining articles, applying inclusion and exclusion criteria (explained below). We also reviewed references of these articles for possible papers missed in the primary search.

Additionally, the following journals were hand-searched to identify articles which presented a method for segmenting WMH in the period covered by this review.

1. Neuroimage – keywords WMH and segmentation from January 2015 to July 2020: identified 6 articles which matched the search results (100% recall).
2. NeuroImage Clinical – from January 2015 to July 2020: identified 7 articles which matched the search results (100% recall).
3. Neuroinformatics – keywords WMH and segmentation from January 2015 to July 2020: identified 2 articles which matched the search results (100% recall).

## **2.2. Inclusion and Exclusion Criteria**

### *2.2.1. Inclusion criteria*

## White Matter Hyperintensities Segmentation Review

- Presentation and / or validation of a fully automated method for segmenting WMH of presumed vascular origin from human brain MR images from January 2015 to July 2020
- Studies published in English

### 2.2.2. *Exclusion criteria*

- Animal studies
- Segmentation methods solely for MS lesions or validated using only MS patients
- New fully automated segmentation methods not described
- Segmentation method accuracy evaluated on CT images
- Segmentation of brain regions, brain tumours or other pathologies which are not WMH of presumed vascular origin
- Semi-automated segmentation methods
- No information on segmentation method similarity metrics used or no evaluation against ground truth segmentations
- Focus on pre-processing or classification of MR images
- Methods designed for WMH evolution assessment (i.e. longitudinal analysis) without cross-sectional validation of their results
- Insufficient information to replicate or apply the segmentation method
- Studies published only as abstracts
- Conference proceedings
- No segmentation method description

### 2.3. **Assessment of Methodological Quality**

We evaluated methodological quality for each study using QUADAS 2: a tool to assess the risk of bias and the applicability of the methods / procedures (<https://www.bristol.ac.uk/media-library/sites/quadas/migrated/documents/quadas2.pdf>). QUADAS 2 contains four domains: 1) patient selection; 2) index test; 3) reference test; and 4) flow and timing. In our case, index test refers to WMH segmentation method / algorithm. Different from the original QUADAS 2

questionnaire, the evaluation of the index text consisted in assessing whether or not the reference standard was used in any way by the segmentation method. We completed the online form for each of the included studies. If a study were judged low in all four domains in relation to bias or applicability from answering the specific questions from each domain, then it was considered as “low risk of bias”. If a study were judged high or unclear for one or more domains, then it was considered as “risk of bias” or as having concerns regarding applicability.

## 2.4. Data Extraction

From the included papers, we extracted the following data:

- Title, year of publication, journal name, study design
- Number of subjects or images, age, gender
- Patient selection criteria, sample size
- Type of MRI sequences used (details about the scanner used)
- Information on imaging features used for investigation
- Details about pre-processing steps (registration, brain extraction, intensity inhomogeneity correction, noise reduction, intensity normalization)
- Method to remove false positives
- Reference standard(s)
- Segmentation method details
- Non-imaging features used for clinical correlation with WMH volume (e.g., cognitive test)
- Sensitivity, specificity, accuracy, dice similarity index, false negative ratio (FNR) and false positive ratio (FPR) of the proposed segmentation method
- Visual rating scale used for validating the segmentation method (if any)

Extracted data were tabulated, synthesized, and evaluated for methodological flaws and applicability of the proposed techniques.

### 3. RESULTS

#### 3.1. Search Results

The search yielded 2327 papers after removing 104 duplicate citations. We schematically represent the selection process in Figure 2; we conducted it according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).

##### 3.1.1. Exclusions

We removed 2268 papers after screening titles and abstracts, leaving 59 for full text screening. We excluded a further 19 for these reasons: validated the method using datasets with brain tumours (2); or MS patients only (6); method for segmenting lacunes (1); or perivascular spaces (3); or only small T2 hyperintensities (1); full text unavailable (1); sample size less than 20 (1); presented a tool for displaying but not segmenting WMH (1); modelling WMH distribution (1 study); or only quantifying longitudinal change (1). Also, three studies did not propose a new segmentation method of WMHs but compared the performance of existing machine learning based segmentation methods of WMHs (Dadar et al., 2017b; Kuijf et al., 2019; Rachmadi, Valdés-Hernández, Agan & Komura 2017), leaving 37 studies for full analysis.




Figure 2. PRISMA Flow diagram of the systematic literature search.

### 3.2. Studies Characteristics

#### 3.2.1. Individuals analysed

The 37 studies included analysed imaging data from approximately 7000 individuals. The exact number and quantitative breakdown of sample characteristics was not possible to ascertain because several studies used overlapping publicly available datasets lacking accompanying clinical or demographic metadata. One large-scale multicentre study involved 2781 subjects from 12 different sites (Schirmer et al., 2019); two studies analysed more than 500 individuals (Griffanti et al., 2016 (n=559); Jiang et al., 2018 (n=566)); eight studies used small relevant samples (i.e., fewer than 30 individuals with WMH of presumed vascular origin) (Ding et al.,

2020; Rachmadi et al., 2020; Rincón et al., 2017; Roy et al., 2015; Stone et al., 2016, Van Opbroek et al., 2015a, 2015b; Valverde et al., 2017); three additionally used data from MS patients (Rachmadi et al., 2020; Van Opbroek et al., 2015a, 2015b).

3.2.2. *Validity of the Data Extracted*

Fourteen studies validated their results in data provided by Medical Image Computing and Computer Assisted Intervention (MICCAI) segmentation challenges (Roy et al., 2015; Sudre et al., 2015; Van Obproek et al., 2015a, 2015b; Wang et al, 2015; Valverde et al., 2017; Zhan et al., 2017; Knight et al., 2018; Li et al., 2018; Manjón et al., 2018; Moeskops et al., 2018; Sundaresan et al., 2019; Wu, Zhang, Wang & Tang, 2019; Liu et al., 2020). Nine of them also used additional datasets or patient data from clinics. Out of 37, twelve studies reported using data from prospective studies or clinics (Atlason et al., 2019; Bowles et al., 2017; Guerrero et al., 2017; Hong et al., 2020; Moeskops et al., 2018; Ling et al., 2018; Park et al., 2018; Qin et al., 2018; Rincón et al., 2017; Roy et al., 2015; Sundaresan et al., 2019; Wang et al., 2015). Four studies used data from the Alzheimer’s Disease Neuroimaging Initiative (ADNI) (Sudre et al., 2017; Dadar et al., 2017a; Rachmadi et al., 2018, 2020), of which only two declared the subset used (Rachmadi et al., 2018, 2020). Two studies recruited patients diagnosed with migraine (Hong et al., 2020; Park et al., 2018). We summarize the population characteristics in Table 2.

Table 2: Sample characteristics, type of segmentation algorithm proposed and average spatial agreement with reference segmentations in the relevant dataset used, from the 37 studies reviewed. Studies appear in alphabetical order of the first author’s surname by year of publication.

| STUDY / CODE REPOSITORY | SAMPLE SIZE | SAMPLE CHARACTERISTICS                                                           | SEQUENCES AND FIELD STRENGTH | TYPE OF SEGMENTATION | AVERAGE SPATIAL AGREEMENT WITH REFERENCE SEGMENTATION |
|-------------------------|-------------|----------------------------------------------------------------------------------|------------------------------|----------------------|-------------------------------------------------------|
| Roy et al. (2015)       | 24          | Selected from a larger cohort of elderly subjects with hypertension based on the | T1W, FLAIR / 1.5T            | Supervised           | DSC 0.60 - 0.76                                       |

## White Matter Hyperintensities Segmentation Review

|                                                                                                                                      |                                    |                                                                                                                                                                                                                                                                                                                                                                  |                             |              |                                                                                                |
|--------------------------------------------------------------------------------------------------------------------------------------|------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|--------------|------------------------------------------------------------------------------------------------|
|                                                                                                                                      | Additional eval. in data from n=20 | burden of WMH. Further evaluation in data from 20 MS patients from the MICCAI MS Segmentation Challenge II 2008<br>( <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a> )                                                                                                                                                   |                             |              | In MS patients: DSC 0.42, TPR 0.55, PPV 0.38                                                   |
| Sudre et al. (2015)                                                                                                                  | Not specified                      | Simulated data from the BrainWeb project (Kwan, Evans & Pike, 1999), MS patient data from<br>( <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a> ) and data with age-related WMH from the MICCAI BrainS challenge                                                                                                          | T1W, FLAIR / 1.5T           | Unsupervised | DSC 0.46, FPR 0.1, TPR 0.38, FNR 0.62                                                          |
| Van Opbroek et al. (2015a)                                                                                                           | 40                                 | 20 healthy elderly subjects from the Rotterdam scan study and 20 MS patients from<br>( <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a> )                                                                                                                                                                                 | T1W, T2W/PD, FLAIR / 1.5T   | Supervised   | Total accuracy score 78%, TPR 42.3 & 52.7 %, FPR 73.2 & 70.0 %                                 |
| Van Opbroek et al. (2015b)                                                                                                           | 40                                 | 20 healthy elderly subjects from the Rotterdam scan study and 20 MS patients from<br>( <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a> )                                                                                                                                                                                 | T1W, T2W/PD, FLAIR / 1.5T   | Supervised   | -                                                                                              |
| Wang et al. (2015)                                                                                                                   | 60 +                               | Elderly patients scanned as part of normal patient care, aged 61- 86 years (mean age 68.2 years), with various degrees of vascular white matter abnormalities. Further evaluation used data from 10 MS patients from the MICCAI MS Segmentation Challenge II 2008<br>( <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a> ) | T1W, T2W, FLAIR / 1.5T & 3T | Unsupervised | DSC 0.81-0.84, FPR 0.13-0.24, FNR 0.14-0.2<br><br>In MS patients: TPR 0.38-0.40, PPV 0.36-0.48 |
| Zhan et al. (2015)                                                                                                                   | 40                                 | Patient selection process not provided, mean age $62.2 \pm 5.9$ years                                                                                                                                                                                                                                                                                            | T1W, T2W, FLAIR / 1.5T      | Unsupervised | DSC 0.75, TPF 0.72, FPF 0.46                                                                   |
| Damangir et al. (2016)                                                                                                               | 119                                | Patients from Kings Health Partners-Dementia Case Register with AD, MCI, and healthy controls, aged $76.4 \pm 7.4$ years, 56% Females                                                                                                                                                                                                                            | T1W, T2W, FLAIR, PD / 1.5T  | Unsupervised | DSC 0.85 – 0.91                                                                                |
| Griffanti et al. (2016)<br><a href="https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/BIANCA">https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/BIANCA</a> | 559                                | Selection criteria not provided. Dataset 1 (neurodegenerative cohort)- 85 patients Dataset 2 (vascular cohort)-MRI data from 474 consecutive patients                                                                                                                                                                                                            | T1W, FLAIR / 3T             | Supervised   | DSC 0.76, FNR 0.25, FPR 0.22                                                                   |
| Stone et al. (2016)                                                                                                                  | 24                                 | Traumatic brain injured patients aged 28-58 years (mean age $39.6 \pm 8.1$ years) with isolated traumatic lesions in the cerebral white matter, recruited based on potential concussive events                                                                                                                                                                   | T1W, T2W, FLAIR / 3T        | Supervised   | Sensitivity 0.68, PPV 0.51                                                                     |



## White Matter Hyperintensities Segmentation Review

|                                                                                                   |      |                                                                                                                                                                                                                                                            |                                          |               |                                        |
|---------------------------------------------------------------------------------------------------|------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------|---------------|----------------------------------------|
| Bowles et al. (2017)                                                                              | 127  | Selection criteria not provided. Heterogeneous dataset containing imaging data from three different acquisition protocols                                                                                                                                  | T1W, T2W, FLAIR / 1.5T                   | Unsupervised  | DSC 0.70                               |
| Dadar et al. (2017a)                                                                              | 130  | Dataset 1: 80 subjects aged 70-90 years old with either normal cognition, MCI or AD.<br>Dataset 2: 40 cognitively normal subjects at risk of AD aged 55-75 years<br>Dataset 3: 10 subjects from ADNI2/GO selected to have different WMH burden             | T1W, FLAIR / 1.5T & 3T                   | Supervised    | DSC 0.51 - 0.62                        |
| Ghafoorian et al. (2017)                                                                          | 420  | Subjects for the RUN DMC study aged between 50 and 85 years with cerebral SVD on neuroimaging (appearance of WMH and/or lacunes)                                                                                                                           | T1W, FLAIR / 1.5T                        | Deep learning | DSC 0.78-0.79                          |
| Rincón et al. (2017)                                                                              | 28   | 13 patients with cortical and lacunar ischemic infarctions, between 40 and 79 years of age, mini-mental state examination (MMSE) scores $\geq 23$ , no severe problems of language and visual/auditory neglect. 15 MCI patients meeting Petersen criteria. | T1W, FLAIR / 1.5T                        | Supervised    | DSC 0.64                               |
| Sudre et al. (2017)                                                                               | 85 + | Two datasets of unspecified size: one from ADNI with minimal to none WMH load and other with a range of WMH burden. Additional Synthetic data derived from a non-specified number of data was used.                                                        | T1W, FLAIR / 3T                          | Unsupervised  | DSC 0.2 – 0.7                          |
| Valverde et al. (2017)<br><a href="http://atc.udg.edu/nic/msseg">http://atc.udg.edu/nic/msseg</a> | 20   | MRBrainS public database, selected with varying atrophy degrees and WMH loads                                                                                                                                                                              | T1W, FLAIR / 3T                          | Unsupervised  | TPF 0.7- 0.8                           |
| Zhan et al. (2017)                                                                                | 104  | Dataset 1: 50 subjects from ACCORD-MIND mean age 62 years<br>Dataset2: 54 image datasets from <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a>                                                                      | T1W, T2W, PD, FLAIR / 1.5T               | Supervised    | DSC 0.76, TPR 0.83                     |
| Bandeira Diniz et al. (2018)                                                                      | 91   | Image data from a database with some of the volumes containing less than 10 lesions clusters while other volumes contains more than 100.                                                                                                                   | FLAIR / Acquisition details not provided | Deep learning | Sensitivity 78.79%, Specificity 98.77% |
| Guerrero et al. (2018)                                                                            | 167  | Patients with their first clinically evident non-disabling lacunar or mild cortical ischemic stroke                                                                                                                                                        | T1W, FLAIR / 1.5T                        | Deep learning | DSC 0.607                              |

## White Matter Hyperintensities Segmentation Review

|                                                                                                                                                                      |                                      |                                                                                                                                                                                                                                                                                 |                        |               |                                                           |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|---------------|-----------------------------------------------------------|
| Jiang et al. (2018)<br><a href="https://cheba.unsw.edu.au/research-groups/neuroimaging/pipeline">https://cheba.unsw.edu.au/research-groups/neuroimaging/pipeline</a> | 566                                  | Data from two sources: 1) Sydney Memory and Ageing Study (n=166 aged 70 - 90 years old provided longitudinal data in 3 time points).<br>2) Older Australian Twins Study (n=400 aged 65 and above).                                                                              | T1W, FLAIR / 1.5T & 3T | Supervised    | DSC 0.85, Sensitivity 0.91, Specificity 0.99              |
| Knight et al. (2018)                                                                                                                                                 | 96                                   | From 7 different sources, provided by different WMH and MS lesion segmentation challenges                                                                                                                                                                                       | FLAIR / 1.5T & 3T      | Supervised    | DSC 0.41-0.70                                             |
| Li et al. (2018)<br><a href="https://github.com/hongweilibran/wmh_ibbmTum">https://github.com/hongweilibran/wmh_ibbmTum</a>                                          | 170                                  | WMH segmentation challenge dataset ( <a href="https://wmh.isi.uu.nl/">https://wmh.isi.uu.nl/</a> ).                                                                                                                                                                             | T1W, FLAIR / 1.5T & 3T | Deep learning | DSC 0.8, Recall 0.84                                      |
| Ling et al. (2018)                                                                                                                                                   | 156                                  | CADASIL patients 18 years old and above. Mutation of gene NOTCH3 confirmed. 90 patients with 2D FLAIR images were aged 24–74 years, mean age $49 \pm 11$ years, F:M = 54:36.<br>66 patients with 3D FLAIR images were aged 35–81 years, mean age $57 \pm 11$ years, F:M = 46:20 | T1W, FLAIR / 1.5T & 3T | Supervised    | DSC 0.79 for 2D FLAIR images and 0.76 for 3D FLAIR images |
| Manjón et al. (2018)                                                                                                                                                 | 128 +                                | 128 subjects with a wide range of WMH load aged 38.6–92.1 years (M:F = 60:68) from Australian Imaging Biomarkers and Lifestyle (AIBL) study, and unspecified data from <a href="https://www.nitrc.org/projects/msseg">https://www.nitrc.org/projects/msseg</a>                  | T1W, FLAIR / 1.5T & 3T | Deep learning | DSC 0.78                                                  |
| Moeskops et al. (2018)                                                                                                                                               | 226                                  | Patient data from MRBrainS13 challenge and two other datasets: patients with type 2 diabetes mellitus, healthy controls and patients from memory clinic                                                                                                                         | T1W, FLAIR / 3T        | Deep learning | DSC 0.67                                                  |
| Park et al. (2018)<br><a href="https://github.com/bypark/DEWS">https://github.com/bypark/DEWS</a>                                                                    | 148                                  | Diagnosis of migraine confirmed by two headache specialists mean age 44.4 (SD 12.4) years                                                                                                                                                                                       | T1W, FLAIR / 3T        | Supervised    | TPR 0.7-0.91                                              |
| Qin et al. (2018)                                                                                                                                                    | 88 (uses WM lesion 3D atlas from 277 | Adults from mid to older ages with WMH without confounding radiological evidence of recent or old strokes                                                                                                                                                                       | T1W, FLAIR / 1.5T      | Supervised    | DSC 0.65 - 0.67, Precision 0.68, Recall 0.68 - 0.69       |

## White Matter Hyperintensities Segmentation Review

|                                                                                                                       | FLAIR images)                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                      |               |                                                                                                                     |
|-----------------------------------------------------------------------------------------------------------------------|------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------|---------------|---------------------------------------------------------------------------------------------------------------------|
| Rachmadi et al. (2018)<br><a href="https://github.com/deepmedic/deepmedic">https://github.com/deepmedic/deepmedic</a> | 288                          | ADNI subjects randomly selected. From the subsample with ground truth segmentations (n=20) 3 were cognitively normal, 12 had early MCI and 5 had late MCI                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | T1W, T2W, FLAIR / 3T                 | Deep learning | DSC 0.54                                                                                                            |
| Atlason et al. (2019)                                                                                                 | 170                          | AGES-Reykjavik study. Age 66–93 years at first visit, and WMH segmentation challenge dataset ( <a href="https://wmh.isi.uu.nl/">https://wmh.isi.uu.nl/</a> ).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            | T1W, T2W, FLAIR / 1.5T & 3T          | Unsupervised  | DSC 0.77, TPR 0.64 in AGES-Reykjavik dataset.<br>DSC 0.53-0.67, TPR 0.25-0.40 in WMH segmentation challenge dataset |
| Schirmer et al. (2019)                                                                                                | 2783 (WMH extracted in 2533) | Acute ischaemic stroke patients ages 63.28 (SD 14.70) years, 61% male, 10.6% had a prior stroke                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | FLAIR / Field strength not specified | Deep learning | (Provides ICC = 0.84, Pearson r = 0.86 with p < 0.001 (validation only used 144 images))                            |
| Sundaresan et al. (2019)                                                                                              | 133                          | Five datasets:<br>1) Neurodegenerative cohort (n=21, age range 63–86 years, mean age 77.1 ±5.8 years F:M = 10:11)<br>2) Patients that recently experienced a minor non-disabling stroke or transient ischemic attack (n=18, (age range 50–91 years, mean age 73.27 ±12.32 years, F:M = 7:11)<br>3) CADASIL patients (n=15, age range 33–70 years, mean age 53.73 ±11.31 years, F:M = 11:4)<br>4) Healthy controls (n=19, age range 29–70 years, mean age 54.58 ±11.25 years, F:M = 6:13)<br>5) MICCAI WMH segmentation challenge dataset (n=60, selected from 3 different primary cohorts, <a href="https://wmh.isi.uu.nl/">https://wmh.isi.uu.nl/</a> ) | T1W, T2W, FLAIR / 1.5T & 3T          | Supervised    | DSC 0.77<br>TPR 0.73 – 0.98                                                                                         |
| Wu et al. (2019)                                                                                                      | 135                          | Data from BIOCARD study. Subjects either cognitively normal (n=113) or MCI (n=22), based on the (NIA/AA) research diagnostic criteria                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | T1W, FLAIR / 3T                      | Supervised    | DSC 0.62, FPR 0.35, FNR 0.37                                                                                        |

## White Matter Hyperintensities Segmentation Review

|                                                                                                                                      |     |                                                                                                                                                                                      |                        |               |                                                      |
|--------------------------------------------------------------------------------------------------------------------------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|---------------|------------------------------------------------------|
| Wu, Zhang, Wang & Tang (2019)                                                                                                        | 60  | Data from a MICCAI WMH challenge ( <a href="https://wmh.isi.uu.nl/">https://wmh.isi.uu.nl/</a> ). Demographic data not provided                                                      | T1W, FLAIR / 1.5T & 3T | Deep learning | DSC 0.78, Recall 0.81                                |
| Ding et al. (2020)                                                                                                                   | 20  | Participants from ongoing normal aging study. Average age 81.2 (SD 7.15) years. 70% Females. 85% white and 15% African-Americans                                                     | T1W, FLAIR / 3T        | Supervised    | DSC 0.78, FPR 0.009 PPV 0.85, TPR 0.70               |
| Fiford et al. (2020)                                                                                                                 | 60  | 30 controls and 30 AD patients from ADNI. Mean age controls 73.4 (6.2) years Mean age AD patients 74.9 years                                                                         | T1W, FLAIR / 3T        | Unsupervised  | DSC 0.74 (F/T PR and NR given by anatomical regions) |
| Hong et al. (2020) <a href="https://github.com/jisu-hong/deepwmh">https://github.com/jisu-hong/deepwmh</a>                           | 148 | Patients with migraine without aura, migraine with typical aura, and chronic migraine. Mean age: 44.4 years, 82 Females                                                              | T1W, FLAIR / 3T        | Deep learning | TPR 0.87, FDR 0.10                                   |
| Liu et al. (2020)                                                                                                                    | 60  | Data from WMH segmentation challenge ( <a href="https://wmh.isi.uu.nl/">https://wmh.isi.uu.nl/</a> ), and SISS challenge 2015. Out of 60, 25 cases contained ischemic stroke lesions | T1W, FLAIR / 1.5T & 3T | Deep learning | DSC 0.829                                            |
| Rachmadi et al. (2020) <a href="https://github.com/febrianrachmadi/lots-iam-gpu">https://github.com/febrianrachmadi/lots-iam-gpu</a> | 60  | 20 subjects from ADNI and 40 MS patients with different loads of MS lesions.                                                                                                         | T1W, T2W, FLAIR / 3T   | Unsupervised  | DSC 0.47 - 0.56 PPV 0.59 TPR 0.47                    |

Legend: ADNI: Alzheimer's Disease Neuroimaging Initiative, AD: Alzheimer's disease, MCI: Mild Cognitive Impairment, M:F (or F:M): number of Male : number of Female (or viceversa) participants, CADASIL: Cerebral autosomal dominant arteriopathy with subcortical infarcts and leuko-encephalopathy, T1W: T1-weighted MRI sequence, T2W: T2-weighted MRI sequence, FLAIR: fluid attenuated inversion recovery MRI sequence, PD: proton density MRI sequence, 1.5T/3T: MRI scanner magnet strength 1.5 or 3 Teslas, DSC: Dice Similarity Coefficient, TPR: True Positive Rate (equivalent to Recall and Sensitivity), FNR: False Negative Rate, FPR: False Positive Rate, TNR: True Negative Rate (equivalent to Specificity), PPV: Positive Predictive Value (equivalent to Precision), WMH: white matter hyperintensities, MS: Multiple Sclerosis, ICC: Intra-class Correlation Coefficient

### 3.3. Risk of Bias Assessment within Studies

We observed four types of bias: spectrum bias, observer bias, verification bias and selection bias (Figure 3). Observer and data selection biases were common. Observer bias, found in 23/37 studies, mainly occurred in studies that proposed a supervised segmentation method.

White Matter Hyperintensities Segmentation Review

These “learned” from reference data generated by one or more observers, or used limited overlapping retrospective data. A study that reported consensus between observers in the generation of reference segmentation data proposed an unsupervised segmentation method (Sudre et al., 2015). Data selection bias was also observed in 25/37 studies.




Figure 3. Types of bias observed in the articles reviewed.

Lack of consideration of differences in disease severity (i.e. WMH burden in relation to underlying disease/population group) is referred to as spectrum bias (Schmidt et al., 2013). Eighteen studies did not clearly report clinical features and disease characteristics of individuals included in terms of WMH severity. Therefore, it was difficult to judge whether or not a wider and balanced spectrum of WMH burden was present in the sample and, consequently, if the methods were biased towards data with higher, medium or small burdens of WMH in a certain population group.

Data inclusion and exclusion criteria were not explained in 22/37 studies. Of the studies that reported demographic information, five recruited healthy controls (Griffanti et al., 2016; Sundaresan et al., 2019; Damangir et al., 2017; Rincón et al., 2017; Ding et al., 2020). One study stated that the data selection and manipulation were blinded to clinical information (i.e.,

## White Matter Hyperintensities Segmentation Review

avoided clinical review bias) (Dadar et al., 2017a). One study reported having selected the cases randomly (Atlason et al., 2019).

The magnet strength of the scanner used to acquire the data processed was reported in 35/37 studies (see Table 2). Twelve studies used data only acquired at 1.5T, and twelve used data only acquired at 3T. 11/37 studies used data acquired at both 1.5T and 3T (see Table 2).

We observed differential verification bias in 17 studies. These studies used different reference standards to verify segmentation methods' performances; i.e., more than one reviewer was involved in manual WMH delineation of different datasets, or each dataset was delineated by a different person using different strategies, without stating the degree of inter-observer reliability or whether or not the final reference segmentation was agreed between the observers involved. Only 8/37 studies made the code publicly available (Griffanti et al., 2016; Hong et al., 2020; Jiang et al., 2018; Li et al., 2018; Park et al., 2018; Rachmadi et al., 2018, 2020; Valverde et al., 2017). One study (Ling et al., 2018) evaluated different configurations of the method described by Griffanti et al. (2016) making recommendations of its use. We present risk of bias assessment of the 37 included studies using QUADAS 2 tool in Table 3. Out of the 37 studies, only 7 were judged as having low risk of bias overall.

Table 3: Risk of Bias Assessment of the studies reviewed using QUADAS 2 Tool

(<https://www.bristol.ac.uk/media-library/sites/quadas/migrated/documents/quadas2.pdf>).

Studies appear in alphabetical order of the first author's surname by year of publication.

| STUDY | RISK OF BIAS                         |                       |                            |                              |                           |                   | APPLICABILITY                   |                    |                        |
|-------|--------------------------------------|-----------------------|----------------------------|------------------------------|---------------------------|-------------------|---------------------------------|--------------------|------------------------|
|       | PATIENT SELECTION                    | INDEX TEST            | REFERENCE TEST             |                              | FLOW &TIMING              |                   | PATIENT SELECTION               | INDEX TEST         | REFERENCE TEST         |
|       | Could the selection of patients have | Could the method have | Is the ref. std. likely to | Is the ref. std. manipulated | Did all data had the same | Were all patients | The included patients match the | Are there concerns | Are there concerns re. |

## White Matter Hyperintensities Segmentation Review

|                                     | introdu-<br>ced bias? | introdu-<br>ced bias? | be<br>correct? | blind to<br>the<br>index<br>test? | ref.<br>std.? | included<br>? | review<br>question? | re. appli-<br>cability? | reprodu-<br>cibility? |
|-------------------------------------|-----------------------|-----------------------|----------------|-----------------------------------|---------------|---------------|---------------------|-------------------------|-----------------------|
| Roy et al.<br>(2015)                |                       |                       |                |                                   |               |               |                     |                         |                       |
| Sudre et al.<br>(2015)              |                       |                       |                |                                   |               |               |                     |                         |                       |
| Van<br>Opbroek et<br>al.<br>(2015a) |                       |                       |                |                                   |               |               |                     |                         |                       |
| Van<br>Opbroek et<br>al. (2015b)    |                       |                       |                |                                   |               |               |                     |                         |                       |
| Wang et al.<br>(2015)               |                       |                       |                |                                   |               |               |                     |                         |                       |
| Zhan et al.<br>(2015)               |                       |                       |                |                                   |               |               |                     |                         |                       |
| Damangir et<br>al.<br>(2016)        |                       |                       |                |                                   |               |               |                     |                         |                       |
| Griffanti et<br>al. (2016)          |                       |                       |                |                                   |               |               |                     |                         |                       |
| Stone et al.<br>(2016)              |                       |                       |                |                                   |               |               |                     |                         |                       |
| Bowles et al.<br>(2017)             |                       |                       |                |                                   |               |               |                     |                         |                       |
| Dadar et al.<br>(2017a)             |                       |                       |                |                                   |               |               |                     |                         |                       |
| Ghafoorian<br>et al. (2017)         |                       |                       |                |                                   |               |               |                     |                         |                       |
| Rincón et al.<br>(2017)             |                       |                       |                |                                   |               |               |                     |                         |                       |
| Sudre et al.<br>(2017)              |                       |                       |                |                                   |               |               |                     |                         |                       |
| Valverde et<br>al. (2017)           |                       |                       |                |                                   |               |               |                     |                         |                       |

## White Matter Hyperintensities Segmentation Review

|                                 |  |  |  |  |  |  |  |  |  |
|---------------------------------|--|--|--|--|--|--|--|--|--|
| Zhan et al. (2017)              |  |  |  |  |  |  |  |  |  |
| Bandeira<br>Diniz et al. (2018) |  |  |  |  |  |  |  |  |  |
| Guerrero et al. (2018)          |  |  |  |  |  |  |  |  |  |
| Jiang et al. (2018)             |  |  |  |  |  |  |  |  |  |
| Knight et al. (2018)            |  |  |  |  |  |  |  |  |  |
| Li et al. (2018)                |  |  |  |  |  |  |  |  |  |
| Ling et al. (2018)              |  |  |  |  |  |  |  |  |  |
| Manjón et al. (2018)            |  |  |  |  |  |  |  |  |  |
| Moeskops et al. (2016)          |  |  |  |  |  |  |  |  |  |
| Park et al. (2018)              |  |  |  |  |  |  |  |  |  |
| Qin et al. (2018)               |  |  |  |  |  |  |  |  |  |
| Rachmadi et al. (2018)          |  |  |  |  |  |  |  |  |  |
| Atlason et al. (2019)           |  |  |  |  |  |  |  |  |  |
| Schirmer et al. (2019)          |  |  |  |  |  |  |  |  |  |
| Sundaresan et al (2019)         |  |  |  |  |  |  |  |  |  |
| Wu et al. (2019)                |  |  |  |  |  |  |  |  |  |
| Wu, Zhang, Wang & Tang (2019)   |  |  |  |  |  |  |  |  |  |
| Ding et al. (2020)              |  |  |  |  |  |  |  |  |  |



|                        |                                                                                   |                                                                                   |                                                                                   |                                                                                   |                                                                                   |                                                                                     |                                                                                     |                                                                                     |                                                                                     |
|------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| Fiford et al. (2020)   |  |  |  |  |  |  |  |  |  |
| Hong et al. (2020)     |  |  |  |  |  |  |  |  |  |
| Liu et al. (2020)      |  |  |  |  |  |  |  |  |  |
| Rachmadi et al. (2020) |  |  |  |  |  |  |  |  |  |

 - High  - Unclear  - Low

3.4. Pre-Processing Methods

All studies which reported ground truth generation details, validated the WMH segmentation method with ground truth binary masks, generated using the FLAIR MRI sequence. However, only three studies reported having used only the FLAIR sequence in their segmentation framework (Bandeira Diniz et al., 2018; Knight et al., 2018; Schirmer et al., 2019). Of the rest (i.e., 34/37) which described using data from different sequences, 28 used a combination of more than one sequence (i.e., also known as “multispectral approach”), generally T1-weighted and FLAIR, to generate the final outcome. In general, after MRI acquisition, various pre-processing steps were conducted. These were often registration, brain extraction, intensity inhomogeneity correction, noise reduction and intensity normalisation. Table 4 summarises the publicly available tools used in the studies’ pipelines and Table 5 summarises the pre-processing steps used by each study. Only one study reported having conducted all the above-mentioned pre-processing steps prior to the segmentation method (Manjón et al., 2018), and one did not provide any information about pre-processing steps performed before segmentation (Liu et al., 2020). The latter selected MRI slices from already brain-extracted images downloaded from an image data repository, without specifying how the slice selection was performed (i.e., by visual inspection or automatically). Slice selection excluded 81 slices with haemorrhagic stroke and those at the top and bottom of the brain, which are more prone to have confounding artefacts.

Table 4. Publicly available software resources and tools used in the selected studies

## White Matter Hyperintensities Segmentation Review

| Software library / Repository                                                                                                                                                                                                                                                                         | Tool                                                                                                                                              | Use                                                                                           | References / Documentation                                                                                                                                                 |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| FMRIB software library (FSL)<br><a href="https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/">https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/</a>                                                                                                                                                                         | Brain Extraction Tool (BET)                                                                                                                       | Removal of non-brain tissues                                                                  | Smith (2002); Jenkinson, Pechaud and Smith (2005)                                                                                                                          |
|                                                                                                                                                                                                                                                                                                       | FMRIB's Linear Image Registration Tool (FLIRT)                                                                                                    | Linear (affine) intra- and inter-modal brain image registration                               | Jenkinson and Smith (2001)<br>Jenkinson, Banner, Brady and Smith (2002)                                                                                                    |
|                                                                                                                                                                                                                                                                                                       | FMRIB's Automated Segmentation Tool (FAST)                                                                                                        | Tissue segmentation and correction for bias field inhomogeneities                             | Zhang, Brady & Smith (2001)                                                                                                                                                |
| Statistic Parametric Mapping (SPM)<br><a href="https://www.fil.ion.ucl.ac.uk/spm/software/">https://www.fil.ion.ucl.ac.uk/spm/software/</a>                                                                                                                                                           | Set of MATLAB scripts and functions integrated in a graphic interface environment that can be used independently                                  |                                                                                               | ("SPM - Documentation", 2020)                                                                                                                                              |
| Advanced Normalization Tools (ANTs)<br><a href="https://sourceforge.net/projects/advants/v">https://sourceforge.net/projects/advants/v</a> , <a href="http://stnava.github.io/ANTs/">http://stnava.github.io/ANTs/</a> ,<br><a href="https://github.com/ANTsX/ANTs">https://github.com/ANTsX/ANTs</a> | antsRegistration                                                                                                                                  | Part of the ANTs suite of image registration tools                                            | ("Advanced Normalization Tools ( ANTs ) - SourceForge.net", 2020)<br>( <a href="http://stnava.github.io/ANTs/">http://stnava.github.io/ANTs/</a> )<br>("ANTsX/ANTs", 2020) |
|                                                                                                                                                                                                                                                                                                       | N3, N4                                                                                                                                            | Correction for bias field inhomogeneities                                                     | Tustison et al. (2010)                                                                                                                                                     |
| bric1936<br><a href="https://sourceforge.net/projects/bric1936/">https://sourceforge.net/projects/bric1936/</a>                                                                                                                                                                                       | bricBET (MATLAB wrappers to FSL-BET)                                                                                                              | Generation of the intracranial volume binary mask from combinations of multiple MRI sequences | <a href="https://sourceforge.net/projects/bric1936/files/Documentation/">https://sourceforge.net/projects/bric1936/files/Documentation/</a>                                |
| 3DSlicer<br><a href="https://www.slicer.org/">https://www.slicer.org/</a>                                                                                                                                                                                                                             | Modules for Image registration<br><a href="https://www.slicer.org/wiki/Slicer3:Registration">https://www.slicer.org/wiki/Slicer3:Registration</a> |                                                                                               | ("Documentation/4.10/Training - Slicer Wiki", 2020)                                                                                                                        |
| Medical Image Registration Tool (MIRTK)<br><a href="https://biomedica.doc.ic.ac.uk/software/mirtk/">https://biomedica.doc.ic.ac.uk/software/mirtk/</a><br><a href="https://github.com/BioMedia/MIRTK">https://github.com/BioMedia/MIRTK</a>                                                           |                                                                                                                                                   | Image registration                                                                            | ("MIRTK – BioMedia", 2020)                                                                                                                                                 |
| OptiBET<br><a href="https://montilab.psych.ucla.edu/fmri-wiki/optibet/">https://montilab.psych.ucla.edu/fmri-wiki/optibet/</a>                                                                                                                                                                        |                                                                                                                                                   | Brain extraction (uses FSL-BET)                                                               | Lutkenhoff et al. (2014)                                                                                                                                                   |
| Elastix<br><a href="https://en.freedownloadmanager.org/Windows-PC/Elastix-FREE.html">https://en.freedownloadmanager.org/Windows-PC/Elastix-FREE.html</a><br><a href="https://elastix.lumc.nl/download.php">https://elastix.lumc.nl/download.php</a>                                                   |                                                                                                                                                   | Image registration                                                                            | Klein et al. (2010); Shamonin et al. (2014)                                                                                                                                |
| Niftyreg<br><a href="https://sourceforge.net/projects/niftyreg/">https://sourceforge.net/projects/niftyreg/</a><br><a href="https://www.nitrc.org/projects/niftyreg/">https://www.nitrc.org/projects/niftyreg/</a>                                                                                    |                                                                                                                                                   | Image registration                                                                            | Modat et al. (2014)                                                                                                                                                        |

## White Matter Hyperintensities Segmentation Review

|                                                                                                                                                                                                                          |                                                                            |                                                                                                                                                  |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| MRICro<br><a href="https://www.mccauslandcenter.sc.edu/crnl/mricro">https://www.mccauslandcenter.sc.edu/crnl/mricro</a><br><a href="https://www.nitrc.org/projects/mricron/">https://www.nitrc.org/projects/mricron/</a> | Brain extraction and delineation of regions of interest                    | ("MRICro   CRNL", 2020)<br>("NITRC: MRICron: Document Manager: Display Document", 2020)                                                          |
| Freesurfer<br><a href="https://surfer.nmr.mgh.harvard.edu/">https://surfer.nmr.mgh.harvard.edu/</a>                                                                                                                      | Image segmentation (Anatomical segmentation of regions of interest)        | ("FreeSurferWiki - Free Surfer Wiki", 2020)                                                                                                      |
| Multi-cONtrast brain STRipping (MONSTR)<br><a href="https://www.nitrc.org/projects/monstr">https://www.nitrc.org/projects/monstr</a>                                                                                     | Brain extraction                                                           | Roy et al. (2017)                                                                                                                                |
| Analysis of Functional NeuroImages (AFNI)<br><a href="https://afni.nimh.nih.gov/">https://afni.nimh.nih.gov/</a>                                                                                                         | Space transformation, visualisation, and statistical analyses of fMRI data | <a href="https://afni.nimh.nih.gov/pub/dist/doc/htmldoc/index.html">https://afni.nimh.nih.gov/pub/dist/doc/htmldoc/index.html</a><br>(Cox, 1996) |

Table 5. Pre-processing in the studies involved. Studies appear in alphabetical order of the first author's surname by year of publication (as per previous tables).

| STUDY                      | REGISTRATION (details reported)                                                                              | BRAIN EXTRACTION                                                                      | INTENSITY INHOMOGENEITIES CORRECTION | NOISE REDUCTION          | INTENSITY NORMALISATION                                                                                                                                      |
|----------------------------|--------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|--------------------------------------|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Roy et al. (2015)          | Rigid-body – uses SPM 8                                                                                      | FSL-BET                                                                               | N3                                   | Information not provided | Histogram matching (Nyul et al., 2000)                                                                                                                       |
| Sudre et al. (2015)        | Intra-subject inter-modality co-registration, and statistical atlases warped to observed data using niftyreg | Performed using STEPS (Cardoso et al. 2013) followed by non-brain tissue mask filling | Information not provided             | Information not provided | Intensity rescaling from 0 to 1.                                                                                                                             |
| Van Opbroek et al. (2015a) | Information not provided                                                                                     | Information not provided                                                              | N4                                   | Information not provided | Three normalisation algorithms were evaluated:<br><br>1) Range-matching (maps the 4th and the 96th percentage of intensity within the brain mask to 0 and 1. |

## White Matter Hyperintensities Segmentation Review

|                            |                                                                                                                 |                                 |                          |                          |                                                                                                                                                                                                       |
|----------------------------|-----------------------------------------------------------------------------------------------------------------|---------------------------------|--------------------------|--------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                            |                                                                                                                 |                                 |                          |                          | <p>2) Linear intensity adjustment to the range [0,1].</p> <p>3) Method 1 followed by mapping of every tenth percentile within 0 and 1 to the mean intensity over all (training and target) images</p> |
| Van Opbroek et al. (2015b) | Information not provided                                                                                        | FSL-BET                         | N4                       | Information not provided | Range-matching procedure that scaled the voxels within a mask such that the voxels between the 4th and 96th percentage in intensity are mapped between 0 and 1                                        |
| Wang et al. (2015)         | Information not provided                                                                                        | Brain Extraction Tool in MRICro | N3                       | Information not provided | Image intensity rescaling from 0 to 255                                                                                                                                                               |
| Zhan et al. (2015)         | A mutual information-based registration method (Viola and Wells, 1997)                                          | FSL-BET                         | N3                       | Information not provided | Information not provided                                                                                                                                                                              |
| Damangir et al. (2016)     | Intra-subject, rigid, 3-D with mutual information - uses FSL FLIRT                                              | FSL-BET                         | N3                       | Information not provided | Information not provided                                                                                                                                                                              |
| Griffanti et al. (2016)    | Linear, with trilinear interpolation (in multisequence evaluation using T1-weighted and FLAIR) - uses FSL-FLIRT | Information not provided        | Information not provided | Information not provided | Used variance scaling                                                                                                                                                                                 |
| Stone et al. (2016)        | Rigid-body, linear intra-subject, of FLAIR, T1- and T2-weighted sequences – uses ANTs                           | ANTs                            | N4 bias correction       | Information not provided | Normalized to the intensity range 0–1.                                                                                                                                                                |

## White Matter Hyperintensities Segmentation Review

|                          |                                                                                                                                                                    |                                                                                         |                                         |                                                                                                   |                                                                                                                                                                         |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|-----------------------------------------|---------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Bowles et al. (2017)     | Rigid-body, linear, of T1-weighted to FLAIR followed by free-form deformation between T1-weighted image in FLAIR image space and an MNI template –uses MIRTk suite | pincram algorithm (uses label propagation and group agreement) (Heckemann et al., 2015) | N4 algorithm                            | Information not provided                                                                          | Linear scaling using the mean intensity of the healthy (i.e., gray and white matter) tissue in the cohort (estimated having the value of 1000). (Huppertz et al., 2011) |
| Dadar et al. (2017a)     | Intra-subject, linear rigid body, of T2-weighted, PD and FLAIR to T1-weighted, then non-linear based on intensity correlation coefficient to ADNI 150 template     | Information not provided                                                                | N3                                      | Automatic multi-threaded denoising method based on non-local means filtering (Manjón et al. 2010) | Linear intensity scaling using histogram matching to a template from 150 subjects (50 normal controls, 50 MCI and 50 dementia) from ADNI database (i.e., ADNI150)       |
| Ghafoorian et al. (2017) | Rigid-body, linear, of T1-weighted to FLAIR with trilinear interpolation and mutual information optimization – uses FSL-FLIRT                                      | FSL-BET on T1-weighted images                                                           | FSL-FAST                                | Information not provided                                                                          | Normalized per patient to be within the range of [0, 1]                                                                                                                 |
| Rincón et al. (2017)     | Linear, intra-subject, of T1-weighted to FLAIR – uses BRAINSFit from 3D-Slicer                                                                                     | White matter tissue and ventricles segmented using Freesurfer                           | SPM 8                                   | Information not provided                                                                          | Normal white matter modelled for Gaussian fit to generate parameters for FLAIR intensity normalisation using the EzyFit Toolbox ("EzyFit 2.44", 2020)                   |
| Sudre et al. (2017)      | Affine, linear, intra-subject, of FLAIR to T1-weighted. Non-linear (niftyreg) used for generating synthetic data.                                                  | Performed, but information not provided                                                 | Performed, but information not provided | Information not provided                                                                          | Information not provided                                                                                                                                                |
| Valverde et al. (2017)   | Affine followed by non-linear registration of the MNI-ICBM152 brain template to the native                                                                         | Information not provided                                                                | Information not provided                | Information not provided                                                                          | Information not provided                                                                                                                                                |

## White Matter Hyperintensities Segmentation Review

|                              |                                                                                                                                                                        |                                                                      |                          |                          |                                                                                                           |
|------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|--------------------------|--------------------------|-----------------------------------------------------------------------------------------------------------|
|                              | T1-weighted space - used niftyreg followed by the registration tool in SPM12.                                                                                          |                                                                      |                          |                          |                                                                                                           |
| Zhan et al. (2017)           | Information not provided                                                                                                                                               | FSL-BET on T1-weighted images                                        | N3 bias field correction | Information not provided | Linear contrast adjustment to match the intensities of the training and testing datasets                  |
| Bandeira Diniz et al. (2018) | Not applicable (uses only FLAIR images)                                                                                                                                | Algorithm by Bauer et. al. (2011)                                    | Information not provided | Information not provided | histogram matching algorithm                                                                              |
| Guerrero et al. (2018)       | T1-weighted registered to FLAIR - uses FSL-FLIRT                                                                                                                       | Information not provided                                             | Information not provided | Information not provided | Tissue intensities normalised to have zero mean and standard deviation of one (excluding the background). |
| Jiang et al. (2018)          | Rigid body, linear, of FLAIR to T1-weighted and from T1-weighted space to the Diffeomorphic Anatomical Registration Through Exponentiated (DARTEL) space - uses SPM 12 | SPM 12 – used brain mask in DARTEL space to remove non-brain tissues | FSL-FAST                 | Information not provided | Information not provided                                                                                  |
| Knight et al. (2018)         | Training images are resampled (trilinear) to 1.5mm isotropic voxel resolution in the MNI-ICBM152 space – uses the Segment tool in SPM 12                               | Segment tool in SPM 12                                               | Segment tool in SPM 12   | Information not provided | Information not provided                                                                                  |
| Li et al. (2018)             | (Retrospective data, intra-subject MRI sequences previously co-registered)                                                                                             | (data was already brain extracted)                                   | SPM12                    | Information not provided | Gaussian intensity normalisation                                                                          |
| Ling et al. (2018)           | Linear, of FLAIR to T1-weighted space and from T1-weighted to                                                                                                          | FSL-BET                                                              | Information not provided | Information not provided | Variance scaling within FSL- BIANCA                                                                       |

## White Matter Hyperintensities Segmentation Review

|                        |                                                                                                                        |                                                                            |                                                         |                                              |                                                                                                                                                         |
|------------------------|------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|---------------------------------------------------------|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|
|                        | MNI-ICBM152 standard space                                                                                             |                                                                            |                                                         |                                              |                                                                                                                                                         |
| Manjón et al. (2018)   | Linear registration to the MNI-ICBM152 space – uses ANTs                                                               | SPM12                                                                      | SPM12                                                   | Spatially Adaptive 3D Non-local Means Filter | All brain voxels intensities were divided by the median intensity within the brain region. Resulting intensities were squared to enhance image contrast |
| Moeskops et al. (2018) | Intra-subject, of T1-weighted (with and without inversion recovery pulse) to FLAIR – uses elastix (Klein et al., 2010) | SPM 12                                                                     | SPM 12                                                  | Information not provided                     | Information not provided                                                                                                                                |
| Park et al. (2018)     | Rigid-body – uses FSL-FLIRT                                                                                            | FSL-BET                                                                    | FSL-FAST                                                | Information not provided                     | Information not provided                                                                                                                                |
| Qin et al. (2018)      | Rigid-body, linear, of T1-weighted and FLAIR - uses FSL-FLIRT                                                          | Information not provided                                                   | As per the unified segmentation algorithm in SPM5       | Information not provided                     | As per the unified segmentation algorithm in SPM5                                                                                                       |
| Rachmadi et al. (2018) | Rigid-body, linear - uses FSL-FLIRT                                                                                    | OptiBET                                                                    | Information not provided                                | Information not provided                     | Histogram matching (Nyúl et al., 2000)                                                                                                                  |
| Atlason et al. (2019)  | Rigid registration to the MNI-ICBM152 template                                                                         | skull removal using MONSTR                                                 | inhomogeneity correction using N4 integrated into SegAE | Information not provided                     | Information not provided                                                                                                                                |
| Schirmer et al. (2019) | Affine – uses ANTs                                                                                                     | (Neuron-BE) that used a 2D U-Net convolutional neural network architecture | Information not provided                                | Information not provided                     | Mean-shift algorithm and corresponding full width half maximum (FWHM) of the peak in the intensity histogram of the total brain volume                  |



## White Matter Hyperintensities Segmentation Review

|                               |                                                                                                                       |                                                                                            |                                             |                          |                                                                 |
|-------------------------------|-----------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|---------------------------------------------|--------------------------|-----------------------------------------------------------------|
| Sundaresan et al. (2019)      | Rigid-body, linear – uses FSL FLIRT                                                                                   | FSL-BET                                                                                    | FSL-FAST                                    | Information not provided | Information not provided                                        |
| Wu et al. (2019)              | FLAIR to T1-weighted previously mapped in MNI-ICBM152 coordinates - uses SPM (version not specified)                  | Information not provided                                                                   | N4 bias correction                          | Information not provided | Histogram matching                                              |
| Wu, Zhang, Wang & Tang (2019) | Information not provided (Retrospective data, intra-subject MRI sequences previously co-registered)                   | FSL-BET                                                                                    | Information not provided                    | Information not provided | Gaussian intensity normalization (z-scores)                     |
| Ding et al. (2020)            | Linear within-subject co-registration of FLAIR and T1W sequences using FSL-FLIRT through fsr (Muschelli et al., 2018) | FSL-BET followed by erosion of the brain mask through fsr (Muschelli et al., 2018)         | N4                                          | Information not provided | z-scores calculated over the entire brain-only intensity signal |
| Fiford et al 2020             | Information not provided                                                                                              | skull-stripped and atlases obtained from label fusion GIF framework (Cardoso et al. 2015). | Method proposed by Van Leemput et al (1999) | Information not provided | Information not provided                                        |
| Hong et al. (2020)            | Rigid-body linear within-subject co-registration of T1W and FLAIR sequences                                           | Information not provided                                                                   | Information not provided                    | Information not provided | Information not provided                                        |
| Liu et al. (2020)             | Information not provided                                                                                              | Information not provided                                                                   | Information not provided                    | Information not provided | Information not provided                                        |
| Rachmadi et al. (2020)        | Rigid-body linear within-subject co-registration of T1W and FLAIR sequences using FSL-FLIRT                           | OptiBET and bricBET (applied to separate datasets)                                         | Information not provided                    | Information not provided | Information not provided                                        |

Legend: T1W: T1-weighted structural magnetic resonance (MRI) sequence, FLAIR: fluid-attenuated inversion recovery structural MRI sequence, MNI-ICBM152 template: Montreal Neurological Institute –

## White Matter Hyperintensities Segmentation Review

International Consortia for Brain Mapping brain template from 152 healthy young adults that includes both a set of coordinates and the associated anatomical labels,

Note: for list of software tools, please, refer to Table 4.

Most studies included in the review used linear transformation models to co-register the different image sequences from each subject data (Figure 4 (left hand side), Table 5). From the 27/37 studies that provided information on the tool used for image registration, ten used FSL FLIRT (Qin et al., 2018; Griffanti et al., 2016; Guerrero et al., 2017; Sundaresan et al., 2019; Damangir et al., 2017; Rachmadi et al., 2018, 2020; Ghafoorian et al., 2017; Hong et al., 2020; Ding et al., 2020), three used ANTs (Schirmer et al., 2019; Manjón et al., 2018; Stone et al., 2016), five used SPM (Wu et al., 2019; Roy et al., 2015; Jiang et al., 2018; Knight et al., 2018; Valverde et al., 2017), one study used elastix (Moskops et al., 2018), one study used 3DSlicer (Rincon et al., 2017), and one study used MIRTk (Bowles et al., 2017). The skull stripping method for removal of non-brain tissues was reported in 25/37 studies: FSL-BET, ANTs, OptiBET, Neuron BE, SPM, Freesurfer, MONSTR and MRlcro. Intensity inhomogeneity correction was reported in 17/37 studies, and it was always performed using a well-known tool: N3 (or N4), SPM, FSL-FAST or the Nu estimate. N3 and its newer version N4, were the tools most commonly used for intensity inhomogeneity correction (Stone et al., 2016; Wu et al., 2019; Bowles, et al., 2017; Dadar et al., 2017a; Van Opbroek, Ikram, Vernooij & de Bruijne, 2015a, 2015b; Damangir et al., 2017; Roy et al., 2015; Wang et al., 2015; Zhan et al., 2015; Zhan et al., 2017; Atlason et al., 2019; Ding et al., 2020). Non-local means was the only filtering technique used by the two studies that reported having included noise removal within their pre-processing steps (Manjón et al., 2018; Dadar et al., 2017a). Neither of these two studies selectively applied the filtering after analysing the signal. The 23/37 studies that provided information on intensity normalisation, reported the use of either variance / linear scaling or histogram matching, with variations in their implementation.

3.5. Segmentation Methods

From the studies included, 73% (i.e., 27/37) proposed a supervised segmentation method. Of them, 37% (i.e. 10/27) used deep learning. From the 10 studies that proposed an unsupervised segmentation method (i.e., 27% of the total number of studies included), one used deep learning (Atlason et al., 2019). In total, eleven studies used Convolutional Neural Networks (Rachmadi, et al., 2018; Li et al., 2018; Guerrero et al., 2017; Moeskops et al., 2018; Ghafoorian et al., 2016; Hong et al., 2020; Manjón et al., 2018; Wu et al., 2019; Liu et al., 2020; Diniz et al., 2018; Schirmer et al., 2019), four studies proposed a method based on k-nearest neighbours (k-NN) (Sundaresan et al., 2019; Jiang et al., 2018; Ling et al., 2018; Griffanti et al., 2016), four studies proposed regression models (Knight et al., 2018; Dadar et al., 2017a; Zhan et al., 2017; Ding et al., 2020), and three studies used Random forest (RF) in their proposed algorithms (Stone et al., 2016; Park et al., 2018; Roy et al., 2015). Two studies proposed a method based on Fuzzy C mean algorithm (Zhan et al., 2015; Valverde et al., 2017) and three proposed improvements to a Gaussian Mixture Model framework (Sudre et al., 2015, 2017; Fiford et al., 2020), both unsupervised methods. We summarize the segmentation methods included in the reviewed studies in Figure 4 (right hand side).




Figure 4. Co-registration procedures involved in the WMH segmentation frameworks (left) and types of WMH segmentation methods covered (right) by the articles reviewed.

### 3.5.1. Supervised WMH segmentation methods

#### 3.5.1.1. *k*-Nearest Neighbours (*k*-NN)

*k*-NN is a well-established pattern recognition method that, for WMH segmentation, compares each voxel's spatial (i.e., location) and intensity features with those extracted from a training set, and assigns a probability of being (or not) WMH based on the result. This algorithm was first proposed for this task in 2000 (Warfield, Kaus, Jolesz & Kikinis, 2000), further evaluated in 2004 (Anbeek et al., 2004) and improved by additionally using spatial tissue type priors in further works (De Boer et al., 2007; Steenwijk et al., 2013). Three of the four papers included in this review that use this method (Ling et al., 2018; Griffanti et al., 2016; Sundaresan et al., 2019), use the implementation Brain Intensity Abnormality Classification Algorithm (BIANCA) of the FMRIB Software Library (FSL). BIANCA (Griffanti et al., 2016) is a versatile, easy to use, freely available implementation, which offers different options for input modalities (i.e., only FLAIR or multi-sequence), weighting the spatial information, local spatial intensity averaging, and for the choice of the number and location of the training points. Ling et al. (2018) evaluated BIANCA using: 1) input modalities FLAIR alone vs FLAIR and T1-weighted; and 2) applying different thresholds to BIANCA's probabilistic output, and highlighted the high number of false positives observed when using the FLAIR sequence alone compared to those obtained when the multispectral approach is used. Sundaresan et al. (2019) improved BIANCA to accommodate variability of sources and automatically optimise the thresholding of the lesion probability map by adaptively determining local thresholds, instead of adopting a global threshold. For this purpose (i.e., calculating and generating the local thresholds), the study presents the Locally Adaptive Threshold Estimation (LOCATE) algorithm.

Jiang et al. (2018) incorporate WMH cluster size as a third feature in the *k*-NN algorithm, and integrate it in a pipeline called UBO detector, freely available from <https://cheba.unsw.edu.au/research-groups/neuroimaging/pipeline>. UBO detector merges registration and normal tissue segmentation functions available in two different software libraries (i.e., SPM the FMRIB Software Library) for pre-processing and uses T1-weighted and

FLAIR images as input. Although UBO uses a supervised algorithm for WMH segmentation, it can prescind from manual generated labels for training by taking candidate clusters from the priors generated in the pre-processing stage. As the authors recognise, the accuracy in segmenting WMH depends on the accuracy of the segmentation of candidate WMH clusters obtained from FSL-FAST.

#### 3.5.1.2. *Large Margin Classifiers*

Large margin algorithms maximise the margin around the decision boundary of a classifier to reduce the uncertainty in the classification, handling well, high-dimensional data (Wu et al., 2013). Qin et al. (2018) developed a supervised large margin algorithm (SLM) followed by a semi-supervised large margin algorithm (SSLM) in a framework that modifies a self-guided labelling procedure, namely unsupervised one-class learning (UOCL) (Liu et al., 2014), which discovers potential “outliers” in the data, being the WMH. Qin et al. (2018) introduced a new term in the objective function of the UOCL that maximises the average margin between the hyperintensities (i.e. considered outliers) and the decision boundary. The general SLM classifier minimises the objective function using a conjugate gradient method to learn from the training set and provides a rough WMH segmentation map. The SSLM, then, refines the given labels on the target data.

#### 3.5.1.3. *Multi-Atlas Segmentation*

Wu et al. (2019) presented a framework that simultaneously segments the brain and detects WMH. The proposed multi atlas-based detection and localization (MADL) framework uses a multi-atlas likelihood fusion approach to segment the brain tissues and structures, and identify WMH. It uses a multi-atlas library generated from 15 FLAIR images with minimal WMH load and atrophy ranging from minimal to moderate. The Bayes maximum a posteriori estimation generates a maximum posterior probability value for each voxel, of belonging to a certain (atlas) label. The WMH are identified as voxels with maximum posterior probability values below certain threshold empirically determined.

#### 3.5.1.4. *Feature Filters*

Rincón et al. (2017) present an object-based segmentation framework, namely amorphous object segmentation in 2D (AMOS -2D). This method uses a multi-level information approach consisting of a hierarchical multi-threshold WMH segmentation followed by an object-based filter that reduces the number of false-positives. After pre-processing T1-weighted and FLAIR images, AMOS-2D applies white-matter Gaussian modelling to determine the intensity distribution of the WMH. An initial WMH mask is generated using multi-threshold segmentation, which combines single grey-scale thresholding with a seed-based thresholding. In the latter, the higher threshold (i.e. seed) acts as WMH detector and the lower threshold (i.e. region) refines the contours. The optimum thresholds are determined ad-hoc from the training dataset. The filter that refines the “initial” WMH mask is an object-based classifier that uses support vector machine (SVM). The feature vector for this classifier initially consisted of 178 features, including normalised intensity, others derived from applying connected-component analysis, distance to white matter contour, distance to white matter skeleton, distance to ventricles, among others not specified. The dimensionality of the initial feature vector was reduced using correlation-based feature selection.

Roy et al. (2015) present two filtering approaches: one for generating probabilistic regions of interest (i.e. weighted candidate voxels) for the segmentation algorithm to operate, and another to post-process the classifier’s output. The first are contrast-based global probabilistic maps generated from a feature set containing enhanced intensity, anatomical and spatial information, and the second is an edge potential function based Markov Random Field model, which is used to remove false positives and obtain the final output.

#### 3.5.1.5. *Regression Models*

Dadar et al. (2017a) proposed a multispectral linear regression classifier that uses the least-squares parameters estimation to segment WMH. It combines intensity and location features from FLAIR, T1-, T2- and PD-weighted MRI and manually labelled training data, to provide a

continuous subject-specific WMH map displaying different levels of tissue damage along with a binary segmentation.

Knight et al. (2018) developed a supervised logistic regression framework exclusively for FLAIR sequences, called Voxel-Wise Logistic Regression. This method modifies the open source Lesion Segmentation Tool (LST) LPA by estimating the voxel-wise logistic regression parameters simultaneously across the image space for facilitating convergence during the parameters' estimation, instead of randomly sampling the image space. The logistic model, trained using the standardised FLAIR intensity levels of a training set, generates a set of parameters that are subsequently smoothed for their use in the lesion prediction for new images.

Zhan et al. (2017) developed a supervised method that integrated the multi-sequence and spatial information in a Bayesian framework for WM lesion detection from multi sequence MR images. The proposed method is based on a three-step approach: 1) multinomial logistic regression is employed to learn the conditional probability distributions of WMH and brain tissues from training data; 2) spatial information from Markov random field priors is merged with multi sequence information in the Bayesian framework to improve the accuracy of WMH segmentation; and 3) pathology background information is used to reduce false positives.

Ding et al. (2019) present a supervised segmentation method called OASIS-AD. This approach is derived from a previous scheme (i.e., OASIS, Sweeney et al., 2013) developed for MS lesion segmentation, which uses a logistic regression model involving several imaging modalities to determine the probability of a voxel being WMH or not. This model uses as input brain-extracted and normalised image data. The enhanced version OASIS-AD additionally erodes the brain-extracted binary mask generated in the pre-processing step and refines the probability map obtained from the regression model by applying a nearest neighbour feature construction approach that uses FSL-FAST (Zhang, Brady & Smith, 2001), followed by a Gaussian filter.

#### 3.5.1.6. *Random Forest (RF)*

## White Matter Hyperintensities Segmentation Review

Park et al. (2018) present a machine learning based pipeline called DEWS (DEep White-matter hyperintensity Segmentation framework). The authors segment the normal appearing white matter using FSL-FAST and use a combination of morphological operations and multi-level thresholding and inter-sequence registration to generate a normal white matter space that contained only deep WMH clusters in the FLAIR space. Then, a RF classifier uses size, texture and multi-parametric intensity statistical parameters from deep WMH (from a training set) as features for detecting small, superficially located deep WMH.

Stone et al. (2016) propose a multispectral framework that concatenates two RF classifiers, which the authors refer as a “two-stages” scheme. The first stage uses image intensity, symmetry, tissue segmentation voxel-wise probabilities, distance maps and neighbourhood statistics from the training data as features. These are used to produce the voxel-wise ‘voting maps’ (i.e. the classification count of each decision tree for each tissue label) of the first RF classifier for their use as tissue priors in a second multispectral 6-tissue segmentation that additionally uses a Markov Random Field as spatial prior. The second stage uses all Stage 1 features plus the Stage 1 voting maps and the resulting posterior probability images as features for the second RF classifier. The whole framework is constructed on Advanced Normalization Tools (ANTs) and ANTsR toolkits. Stone et al. (2016) suggested that proposed supervised method is suitable for large dataset. However, this method is tested in a small sample size.

Roy et al. (2015) use a set of nine features as input to the RF classifier. The first eight features contain multi-sequence (i.e. from T1-weighted and FLAIR) intensity, anatomical and spatial information per voxel. These are generated from probability maps of cerebrospinal fluid, grey and white matter, and normalised (x,y,z) coordinates in the MNI 152 space. The last feature is the global reference points-based contrast resulted from the filtering technique referred previously.



### 3.5.1.7. Support Vector Machine (SVM)

Van Opbroek et al. (2015a,b) evaluate different transfer-learning approaches in linear and non-linear SVM classifiers, all consisting of different strategies for weighting the feature vector. Both studies use data from different datasets acquired under different scanning protocols and conclude that their transfer learning strategy (i.e. weighting the feature vector) outperforms the conventional SVM using non-weighted features. In Van Opbroek et al. (2015a) authors evaluate two feature sets: one of size 6 and other of size 33. The former uses the intensity and x,y,z voxel coordinates of cerebrospinal fluid, white and grey matter probabilistic segmentations in FLAIR and the latter uses the same features but also for T1- and T2/PD-weighted, using Gaussian kernels of  $\sigma = 0.5, 1$  and  $2 \text{ mm}^3$ . In Van Opbroek et al. (2015b), the authors add the gradient magnitude and the Laplacian of the normalized intensities after convolution with the Gaussian kernel at different scales and recommend using always a feature vector higher than 10 in size.

Van Opbroek et al. (2015a) assign weights to each feature of the feature vector in a way that the sum of all weights equals the total number of training samples, and combine training data with the same intensity distribution with data with different distribution in three weighting schemes: 1) Weighted SVM; 2) Re-weighted SVM; and 3) TransAdaBoost. In (1) lower weights are assigned to misclassified training data with different distribution. In (2) the misclassified lower weights (i.e. from (1)) are iteratively reduced. TransAdaBoost increases the weights of misclassified same-distribution data and reduces those from misclassified different-distribution data, but this scheme was the worst performer. In the same study authors also evaluate the namely “Adaptive SVM” that uses a weighted vector from same-distribution data for training and is tested from different-distribution data.

Van Opbroek et al.(2015b) rather evaluates three different point distribution functions (PDFs) dissimilarity measures to generate the optimal weights for the Weighted SVM classifier – the winner scheme from those evaluated in (2015a)-, which in this case uses a Gaussian kernel. The weights are chosen in an unsupervised manner, by minimizing the difference between the PDFs

of the weighted training images and the PDF of the target image. The three PDF dissimilarity measures evaluated are: 1) the Kullback–Leibler divergence; 2) the Bhattacharyya distance; and 3) the squared Euclidean distance. The optimal weights are determined by minimizing these three dissimilarity criteria while constraining them to the range  $[0;1]$  and that the norm of all the weights should be 1, using the interior-reflective Newton method (Coleman and Li, 1996).

#### 3.5.1.8. *Neural Networks*

Moeskops et al. (2018) evaluated the 3-pathway multi-scale (i.e. patch-wise) convolutional neural network (CNN) scheme developed by the same group in 2016 for segmenting normal tissues in neonatal and young adults (Moeskops et al., 2016) to segment WMH in addition to normal tissues in MRI scans for older individuals / patients. In this occasion, the scheme uses the T1-weighted, T2-weighted, FLAIR and T1-weighted inversion recovery (IR) images as input. Along with WMH, the scheme segmented normal-appearing white matter, cortical grey matter, basal ganglia, thalamus, cerebellum, brain stem, lateral ventricular cerebrospinal fluid, and peripheral cerebrospinal fluid.

Bandeira Diniz et al. (2018) use Simple Linear Iterative Clustering (SLIC) to group pixels based in their location and intensities and generate candidates to lesion / non-lesion regions in each FLAIR axial slice. Authors design a single-pathway CNN for extracting implicit features from the “superpixels” of the FLAIR axial slices presented as input and classify them in lesion regions or non-lesion regions. The CNN seems to have a linear deep architecture, developed ad-hoc for this purpose. This approach resulted efficient in heterogeneously sourced data, reporting a negligible number of false positives.

Rachmadi et al. (2018) proposed an adaptation of a dual-pathway CNN scheme developed for segmenting brain lesions with considerable mass effect (Kamnitsas et al., 2017) to segment WMH. The authors introduced a way to integrate spatial information to the CNN scheme for WMH segmentation called global spatial information (GSI), and evaluate the performance of two configurations (i.e. with 8 and 5 convolutional layers) using only FLAIR vs. using a

## White Matter Hyperintensities Segmentation Review

combination of T1-weighted and FLAIR, and repeated the experiments using a single-pathway CNN architecture with and without GSI. Authors recommend the use of GSI in a multispectral (i.e. using more than one MRI sequence) dual-pathway scheme of the 2D CNN architecture evaluated.

Manjón et al. (2018) present an ensemble of patch-wise neural network classifiers for segmenting WMH on FLAIR images. After a lesion candidate ROI selection, a feature vector containing 58 features (voxel intensities from 3x3x3 and 5x5x5 patches, 3 spatial coordinates and one *a priori* lesion probability) is used by an ensemble of two one-hidden layer feedforward multilayer perceptron which performs the classification. The study evaluates two ways of configuring this ensemble: bagging (Bootstrap aggregating) and boosting. The first approach averages the outputs of the two neural network classifiers, independently trained on different randomly selected datasets. The second approach uses the output from one classifier to improve the next one by either iteratively giving more weight in the next classifier, to the samples wrongly classified in the first one, or non-randomly selecting (i.e. on the training dataset) with higher probability samples wrongly classified previously.

Guerrero et al. (2018) use the UResNet CNN architecture to segment WMH and distinguish them from stroke lesions. This method comprised an analysis path that gradually learned low- and high-level features, followed by a synthesis path, that gradually combined and up samples the low and high-level features into a class likelihood semantic segmentation. The authors confirmed that the CNN architecture performed well compared to other state of the art algorithms.

Li et al. (2018) propose a method using a 19-layer deep fully CNN scheme. In this method, WMH detected based on convolution-deconvolution architecture with long-range connections which simultaneously classified each pixel and locates objects of an input image. The scheme used ensemble models with random parameter initializations and shuffled data for voting the pixel labels in the final evaluation, all which conferred good adaptability on multi-scanners and

protocols and helped reduce overfitting. The authors pointed out that FLAIR and T1 sequences provide complementary information to detect WMH.

Schirmer et al. (2019) incorporate a deep learning CNN previously proposed by Dalca et al. in 2014, in a pipeline consisting of: 1) brain extraction using only clinical FLAIR images; 2) intensity normalisation to accommodate for multi-site heterogeneity; and 3) automatic atlas-based segmentation of WMH.

Ghafoorian et al. (2016) implement several deep CNN architectures which considered multi-scale patches or explicit location features while training, to integrate the anatomical location information into the network. The authors point out that the CNNs which incorporated location information significantly outperformed a conventional segmentation method with hand-crafted features and CNNs that did not integrate location information.

Wu, Zhang, Wang & Tang (2019) modified the U-Net CNN architecture by skipping connections between the down- and up-sampling convolutional branches of the original model, and named their model Skip Connection U-Net (i.e., SC U-Net). SC U-Net additionally connects the outputs of the 4th, 7th, 10th and 13th layers in the down-sampling convolutional branch of the original model to the outputs of the 15th, 18th, 21th and 24th layers in the up-sampling convolutional branch, and feeds them (i.e., the outputs of the 4th, 7th, 10th and 13th layers) to the 16th, 19th, 22th and 25th layers. Hence the resultant model consists of a shrinking part which aims to capture context, a symmetric expansive part that gradually combines features to enable a precise localization, and a skip connection part that alleviates the vanishing gradient problem and improves the speed of the optimization convergence facilitating the training.

Liu et al. (2020) present a multi-scale feature-based CNN model, called M2DCNN not only to segment WMH, but also to distinguish them from ischemic stroke lesions. M2DCNN contains two symmetric U-shaped subnets that produce multi-scale features through the inclusion of dense and dilated blocks. The former helps reducing the number of training parameters and alleviate the gradient vanishing problem. The latter helps enlarging the receptive fields of the

convolution blocks without reducing the feature map size. M2DCNN uses a loss function based on the Dice coefficient.

Hong et al. (2020) present a deep-learning architecture that concatenates two U-Net CNN models that use 3x3 kernels in their convolutional layers. The first U-Net consists of four down-sampling and four up-sampling convolutional layers, and generates WMH priors from brain-extracted co-registered T1W and FLAIR images. These WMH candidates, together with the brain-extracted co-registered T1W and FLAIR images, are input to the second U-Net, consisting of two down-sampling and two up-sampling convolutional layers, which reduces the false-positives.

### 3.5.2. *Unsupervised WMH Segmentation Methods*

Damangir et al. (2017) developed an unsupervised method that statistically defined WMH based on the one-tailed Kolmogorov-Smirnov test (Gail et al., 1976).

Zhan et al. (2015) present an unsupervised WMH segmentation method for T1 and FLAIR data. The T1 image is, first, segmented into different normal tissues, among which regions of white matter and grey matter are combined to provide a region of interest that is subsequently mapped to the FLAIR image. Secondly, the authors calculated the z-score of the intensities in the ROI and defined a threshold to find the abnormalities in normal tissues. They then employ a level set method to improve the preliminary thresholding-based segmentation results and extracted the WMH. The authors pointed out that LGDF energy aided to obtain precise segmentation results compared to other level set methods that used global intensity information.

Bowles et al. (2017) propose a method built upon previous work by the same authors, which can detect abnormally hyperintense regions on FLAIR, disregarding the underlying pathology or location by combining image synthesis, Gaussian mixture models and one-class support vector machines trained only on healthy tissue.

## White Matter Hyperintensities Segmentation Review

Valverde et al. (2017) integrate a partial-volume tissue segmentation with WM outlier rejection and filling, combining intensity, probabilistic and morphological prior maps in a pipeline consisting of five steps. These are: 1) Register three statistical a-priori tissue atlases (CSF, GM and WM) and a brain structure atlas to the patient space; 2) Perform atlas-based 5-tissue segmentation on the T1-weighted image; 3) Detect and refill WM outliers as normal-appearing WM based on the registered a-priori and hyper-intense FLAIR maps if available (using the segmentation from step 2); 4) Re-estimate (again) the 5-tissue classes; and 5) Reassign intermediate volume maps into CSF, GM and WM using both neighbour and spatial prior information.

Wang et al. (2015) model the WMH in FLAIR images as having either Gumbel or Fréchet histogram distributions (see Table 1) and compare the results of their algorithm with those from applying a trimmed likelihood estimator. Although results were not accurate for all degrees of lesion loads authors recommend the principle, especially using the Fréchet distribution, due to its simplicity, for studies of ageing and vascular dementia, likely to include subjects with moderate-to-high lesion load.

Atlason et al. (2019) present an autoencoder Segmentation Auto-Encoder (SegAE) consisting on a CNN with architecture similar to that of U-Net with an additional linear layer and parameter constraints to perform linear unmixing. In this model, down- and up-sampling are performed with strided convolutions of 3D kernels of size 2x2x2 and skip connections are added between activations of the same spatial resolution from the down-sampling to the up-sampling paths. The pure tissue, WMH and cerebrospinal fluid masks obtained during the segmentation were used as priors for the N4 algorithm. Thus, the b1 inhomogeneities are corrected during the training phase and segmentation takes place in presence of inhomogeneity artifacts.

Rachmadi et al. (2020) present an unsupervised segmentation method called Limited One-Time Sampling Irregularity Map (LOTS-IM). This method generates an irregularity map (IM) that represents all voxels as irregularity values ranging from 0 to 1 with respect to the ones considered "normal" based on the original FLAIR texture information. The scheme hierarchically

samples a limited number of target squared patches (i.e., 2D patches of 1x1, 2x2, 3x3 and 4x4) from a non-overlapping grid of source patches of the same size on each brain-extracted FLAIR image slice, assigning an irregularity value to each source patch. The final irregularity map is generated by blending the hierarchically generated irregularity maps, penalizing the result using the original FLAIR intensities, and normalizing the final values between 0 and 1. The WMH are obtained by thresholding the irregularity map.

Fiford et al. (2020) examined an unsupervised segmentation method Bayesian Model Selection (BaMoS) (Sudre et al., 2015), which models the data as a multivariate mixture of Gaussians, further optimized using the expectation-maximization algorithm. It uses an initial outlier map derived after convergence of the initial Gaussian mixture model to enhance sensitivity, as proposed by Sudre et al. (2017). Newly, this study incorporates a two-threshold selection of the candidate regions and selects the WMH clusters after applying connected component analyses twice, considering first 18 neighbourhoods, and then 6 neighbourhoods, to avoid discarding regions where artefacts and true WMH are present.

### 3.6. Additional Publications Evaluating WMH Segmentation Methods

Dadar et al. (2017b) compare the performance of 10 different linear and non-linear supervised classification methods segmenting WMH in brain scans from 201 subjects from four different datasets. The methods evaluated are: Naïve Bayes, Logistic Regression, Linear and Quadratic Discriminant Analyses, k-NN, decision trees, RF, AdaBoost, SVM and Bagging. Out of these methods, RF was the best performer.

Kuijff et al. (2019) comparatively evaluate 20 methods presented at the 20th International Conference on Medical Image Computing and Computer Assisted Intervention (MICCAI) WMH segmentation challenge in 2019. All algorithms are trained with 60 image datasets acquired in 3 different MR scanners, and evaluated on 110 image datasets from 5 MR scanners. All image data are composed of T1-weighted and FLAIR brain-extracted, bias-corrected and co-registered

images from patients with various degrees of age-related neurodegeneration and presenting different vascular pathologies. However, the WMH volume distribution across the dataset is skewed towards low-to-medium WMH burden. From the 20 methods evaluated 14 are neural network approaches, four involve RF, one uses logistic regression, and one a three-level Gaussian mixture model. The evaluation combines the results from five similarity metrics: DSC, a modified Hausdorff distance (95th percentile), absolute percentage volume difference, sensitivity (recall), and F1-score for individual lesions. The top-ranked methods use ensembles of neural networks

Rachmadi et al. (2017) compare the performance of two conventional machine learning classifiers (i.e. support vector machine (SVM) and RF) with the performance of three deep learning algorithms, namely the deep Boltzmann machine, convolutional encoder network and a CNN dual-pathway architecture developed specifically for brain lesion detection, for segmenting WMH on brains displaying only mild or no vascular pathology. The results from these five supervised machine-learning methods are also compared with the results from the unsupervised lesion growth algorithm (LGA) of the Lesion Segmentation Tool (LST) publicly available. The evaluation uses FLAIR and T1-weighted images from 20 subjects randomly selected from the ADNI database (<http://adni.loni.usc.edu/>), acquired in three consecutive years, and for which ground truth WMH segmentations from two different analysts were available. Authors adapted (and/or implemented) configurations that were reported to give the highest WMH segmentation accuracy in previous works. For SVM and RF this study evaluates several combinations of feature vectors with lengths ranging from 44 to 4000, all reported previously having generated results from similar quality. The optimum threshold that defines the boundaries of the probabilistic WMH segmentations differed across methods. Differences in methods' performance depending on the WMH burden prompted authors to conclude that deep-learning methods, in general, performed better than the two conventional machine learning classifiers (i.e. SVM and RF), being the patch-based CNN configuration the best approach only for scans with low burden of WMH.



### 3.7. Segmentation Descriptive Quality

We analyzed the descriptive segmentation method qualities using the scale developed by Byrne et al. (2016). The segmentation descriptive quality (SDQ) is rated on a three point scale: 1 - indicates description of the segmentation method; 2 - indicates explanation of the segmentation method, but no description of how each step is applied; and 3 – indicates full explanation of how the segmentation method proposed is applied. 23/37 studies scored 3.

### 3.8. Processing Time of Segmentation Methods

Processing time of machine learning based segmentation methods refers to: 1) time taken to load the image; and 2) time taken for application of automatic segmentation algorithm (Cruz et al., 2017). Only 9/37 studies reported the processing time of segmentation method (Ling et al., 2018; Rachmadi et al., 2018, 2020; Manjón et al., 2018; Dadar et al., 2017a; Jiang et al., 2018; Qin et al., 2018; Griffanti et al., 2016; Atlason et al., 2019). Out of these nine studies, four reported the time consumed by the segmentation per image. It ranged from 0.03 seconds to 9 seconds per MRI. The method proposed by Qin et al. (2018) consumed considerably less time (i.e., 0.03 seconds per image) compared to the rest.

### 3.9. Methods Evaluation

The proposed method was cross validated with leave-one-subject-out evaluation in six studies (Stone et al., 2016; Moskops et al., 2018; Jiang et al., 2018; Knight et al., 2018; Li et al., 2018; Wu et al., 2019). Fourteen studies described a method for false positive removal (Sudre et al., 2017; Manjón et al., 2018; Stone et al., 2016; Moskops et al., 2018; Rincón et al., 2017; Bowles et al., 2017; Li et al., 2018; Ghafoorian et al., 2017; Roy et al., 2015; Zhan et al., 2017; Hong et al., 2020; Fiford et al., 2020; Rachmadi et al., 2020; Ding et al., 2020). Fazekas visual rating scale was used in the validation of the results in nine studies (Rachmadi et al., 2018; Qin et al., 2018; Guerrero et al., 2017; Griffanti et al., 2016; Ling et al., 2018; Moskops et al., 2018; Jiang et al., 2018; Bowles et al., 2017; Rachmadi et al., 2020).

Of the 37 studies, 29 studies evaluated the performance of their WMH segmentation method using the Dice Similarity Coefficient (DSC) among other metrics that measure spatial concordance between the results of the method proposed and reference segmentations. Average DSC values ranged from 0.538 to 0.91 (Table 2). The unsupervised segmentation method proposed by Damangir et al. (2016) reported the highest average DSC value for WMH segmentation (DSC ranging from 0.85 up to 0.91), followed by the also unsupervised scheme proposed by Wang et al. (2015) (DSC ranging from 0.81 to 0.84), and the k-NN scheme proposed by Jiang et al. (2018) (UBO detector, DSC 0.85). The Bland Altman plot (Bland & Altman, 1986) was used in five studies to analyse the volumetric agreement between the method's result and manual segmentation (Qin et al., 2018; Guerrero et al., 2017; Ling et al., 2018; Sudre et al., 2015; Fiford et al., 2020). Only four studies validated their method in longitudinal samples (Sudre et al., 2017; Jiang et al., 2018; Rachmadi et al., 2018, 2020), and eight performed an additional validation (i.e., to the traditional comparison against reference standard measurements) using clinical parameters (Guerrero et al., 2018, Jiang et al., 2018; Qin et al., 2018; Rachmadi et al., 2018, 2020; Schirmer et al., 2019; Wu et al., 2019; Fiford et al., 2020). Comparison with other methods' performance was done in 29/37 studies. The reference algorithms for excellence were the Lesion Growth Algorithm (LGA) and the Lesion Prediction Algorithm (LPA), both unsupervised methods from the Lesion Segmentation Tool (LST) for SPM (<https://www.applied-statistics.de/lst.html>).

#### 4. DISCUSSION

In the five-year period evaluated, 37 studies proposed new, or adapted and re-purposed existing approaches, for segmenting WMH of presumed vascular origin from brain MRI. Of these, only 10 were unsupervised. Within the last two years, considerable efforts have been put into developing deep learning WMH segmentation methods particularly based on CNN architectures that have demonstrated success in similar tasks. From the supervised algorithms, 37% used state-of-the-art CNN and the rest used either conventional machine-learning

algorithms, the k-NN algorithm or logistic regression models. Despite the high accuracy usually reported by CNN algorithms, those reviewed do not outperform, in terms of spatial agreement with reference segmentations, the more traditional clustering (i.e. k-NN) and logistic regression supervised methods or the unsupervised methods published in this period. Probably the simplicity and strong priors of the k-NN and logistic regression methods make them easier to train with less data, and are less susceptible to overfitting when training data is limited, compared to the deep-learning schemes. The fact that most of these methods give probabilistic outputs, may be helpful in quantifying marginally pathological tissues like dirty-appearing white matter, and help in the characterisation of ill-defined WMH boundaries. However, it also conspires against their evaluation since these probabilistic results need to be binarised for comparison with manually-derived segmentation binary masks. Quality of reporting has a considerable effect on studies' value. Poor reporting of the pre-processing and segmentation methods' steps and lack of availability of the code significantly affects the applicability of various studies included in this review.

We evaluated the validity and accuracy of the segmentation methods reviewed. We refer to validity as the extent to which these algorithms measure what they intended to beyond the data used to develop (i.e., train) and validate them, thus including the applicability to other data. Most studies ignore the issues pertaining to validity and focus only on accuracy of their algorithms. The validity of the proposed segmentation methods was not always clear, mainly due to the different sources of bias in the reference used to evaluate the algorithms (i.e. observer bias in manually-delineated ground truth), the sample selection, and the data source (i.e. mainly from a single protocol and / or acquired from scanners with the same field strength). Many studies exhibited observer bias, either in training or in evaluating their algorithms, as manual outlines of WMH are always affected by the observer's perception in recognising a true lesion from an artefact and are influenced by the observer's experience and ability in delineating the lesion boundaries on MR images. Moreover, reference segmentations are generally obtained by manually refining a semi-automatic segmentation result, obtained generally by thresholding followed by a region-growing algorithm. Selecting the optimum

threshold to segment WMH from FLAIR MRI can also be a source of bias (Valdés Hernández et al., 2010). Additionally, not all the studies included were absent of having data selection bias, which can facilitate overfitting if this is not properly addressed. Data augmentation helps reducing overfitting and increasing the number of the training data. However, effects of bias cannot be balanced-out by increasing the sample size or by repetition (Schmidt et al., 2013).

It is important to describe the target population, which informs the individuals for whom the results of the study are intended to apply. It can be inferred from the data used in the method development. Studies that validated their methods on a dataset different from the one used to develop it, in terms of clinical and image acquisition characteristics, obtained lower spatial agreement in this validation dataset (Roy et al., 2015; Wang et al., 2015; Atlason et al., 2019) (Table 2). Many of the studies included analysed the WMH load in the sample, only expressing that it “was representative of the whole load of WMH burden”. However, representativeness does not mean “balanced”: unbalanced data biases the results in favour of the dominant data subgroup – generally patients with medium-to-large WMH burden. Also, many studies did not explain the rationale followed for data selection. For instance, patients with mild cognitive impairment, Alzheimer’s disease, and normal cognition were included in the same study without explaining the selection criteria and relevance for the main objective of the study, i.e. segmenting WMH. For sample sizes like the ones observed in the majority of studies included (e.g.  $n < 100$ ), cognitive status is not a proxy for WMH load (Damangir et al., 2017). It is, therefore, difficult to decide for which level of severity of a particular condition or for which neurological condition the segmentation method had performed well and, therefore, would be recommended.

Many of the included studies used the open access datasets or the datasets provided for the different Lesion Segmentation Challenges. Mendelson et al. (2017) pointed out that using an open access dataset to evaluate the performance of a segmentation method introduces selection bias (Mendelson et al., 2017). It indeed is practical, cost effective and allows comparability between methods, but only within the context of the dataset used, especially in the case of supervised methods. Hence, segmentation studies can suffer from limited high-

quality data, which is required for training, and poorly labelled region of interests (Challen et al., 2019). The full value of a large dataset depends on the accuracy and completeness of the data collection, which is expensive and time consuming. The use of a limited dataset in cross-validation can falsely show high performance. To evaluate the performance of a segmentation method, large collections of image data are required. Data augmentation and high quality synthetic data can help addressing this need.

Segmenting a medical image is a laborious task. In general, it requires two main steps: 1) image pre-processing; and 2) segmentation (Jude Hemanth & Anitha, 2012). Pre-processing steps generally involve registration, brain extraction, intensity inhomogeneity correction, noise reduction and intensity normalisation (García-Lorenzo et al., 2013). If task-unrelated pathologies (e.g., stroke lesions, SVD neuroradiological features) or imaging artefacts would affect the segmentation algorithm, their identification should be part of the segmentation framework. Main objectives of pre-processing are removal of noise and confounding features, and improving image quality. Not reporting all these steps can be interpreted as they not being necessary or part of the segmentation framework, affecting its reproducibility. Good reporting quality is extremely important, to ensure that accurate and trustworthy information is obtained from the published studies (Samuel et al., 2016). Quality of reporting research studies needs to be improved by following the guidelines outlined by various organisations ("Reporting guidelines | The EQUATOR Network", 2020).

Accuracy of the segmentation methods evaluated in this review refers to their ability to distinguish WMH from normal appearing white matter or other pathological features of similar appearance, as well as a reference or “ground truth” segmentation manually generated by experts. Accuracy was estimated with Bland Altman plots, Jaccard Index, intra class correlation coefficient, true and false positives and negatives and DSC. All these measures have advantages and drawbacks when applied to this context, as none of them alone gives the necessary information about the precision and further applicability of the results in a clinical context. For example, the Bland-Altman plot *per-se* only allows volumetric comparison between the target and reference methods. The Jaccard Index and DSC are equivalent and they reflect the spatial

agreement between the two masks, but do not express how well the algorithm identified the true WMH and/or excluded the non-WMH voxels, as the true positives and negatives are given by other measurements (e.g., true positive fraction, true negative fraction, positive predicted values, false negative/positive fractions). Finally, the correlation coefficient between quantitative WMH volumes and clinical visual ratings (e.g. Fazekas scores), although of clinical use, only gives a gross estimate of how close to the neuroradiological assessment the segmentation is. Most of the papers included did not analyse the results of these metrics combined. It reinforces the claim by Pellegrini et al. (2018) that the relevant literature of computational segmentation algorithms is still insufficiently intertwined with the clinical world. We believe this depends, at least in part, on a misalignment of targets and methods. The computer scientists' community still aims primarily for algorithm novelty and reaching high levels of precision, experimenting with methods largely inspired by recent developments in the field of computer vision. The clinical research community, on the other hand, aims to verify associations (e.g., biomarkers for outcome, effect of drugs vs. placebo) with clinically relevant features that reflect in an improvement in patient outcomes using statistical models. A combination of methods and aims is, therefore, of great importance. The fact that 22% of the studies analysed incorporated a clinical validation to their scheme is encouraging.

Only 26% of the studies included in the review reported the processing time of the proposed segmentation algorithm. Reporting the processing time of the segmentation method could also aid in its further translation to clinical practice, by highlighting its speed or the need of optimising its current implementation. Despite its importance, translating research into clinical practice is challenging. Aside from simply demonstrating superior efficacy, new technologies entering the medical field must also integrate with current practices and be effective in an individual case basis. Kristensen et al. (2015) have reported that it takes more than a decade to implement research results in clinical practice. The research required for this "personalised" medicine would only be possible through summarising and integrating enormous quantities of medical information. This review has shown that this is still unachieved.

This review systematically extracted, synthesised, critically appraised and presents information about a highly active research field. Its main strengths are: 1) careful selection of relevant studies amongst a vast number of initial candidates resultant from the search; 2) identification of the possible sources of bias of the studies; and 3) synthesis of the contributions of the included papers.

Limitations are that we only included the articles published in English language for which we have full access: we may have missed articles published in languages other than English or other articles for which we could not access the full text. Also, there might be other relevant papers missing as a result of incongruences between the search terms and the article keywords or indexing in the databases. By excluding articles published in conference proceedings it is possible that promising WMH segmentation methods could have been excluded.

## 5. CONCLUSION AND FUTURE WORKS

Despite the increasing popularity and high accuracy of CNN schemes applied to WMH segmentation, we found no evidence to favour their application in clinical research over the k-NN algorithm, linear regression or unsupervised methods. High-quality large-sized data availability continues to limit computational developments of segmentation methods, biasing the studies. Future works should carefully consider ways to reduce or compensate the effect of observer, spectrum and selection biases, and improve transparent reporting. Future studies should also analyse the combined effect of several metrics in evaluating the results of their algorithms, to inform on the applicability of the method in clinical research and practice. The lack of code availability of some algorithms presented, and information about the pre- and post-processing steps, and processing time of segmentation *per se* limited the analyses presented and the further reproducibility of the results: issues that we hope future studies overcome.

## 6. ACKNOWLEDGEMENTS

MVH is funded by Row Fogo Charitable Trust (Grant no. BROD.FID3668413). This work is supported by: the Row Fogo Centre for Research into Ageing and the Brain, and the UK Dementia Research Institute which receives its funding from DRI Ltd, funded by the UK MRC, Alzheimer's Society and Alzheimer's Research UK. Images were acquired at the Research MR scanners at the Edinburgh Imaging facilities, supported by the Scottish Funding Council through the Scottish Imaging Network, A Platform for Scientific Excellence (SINAPSE) Collaboration; the 3T scanner is funded by the Wellcome Trust (104916/Z/14/Z), Dunhill Trust (R380R/1114), Edinburgh and Lothians Health Foundation (2012/17), Muir Maxwell Research Fund and the University of Edinburgh.

## 7. REFERENCES

Advanced Normalization Tools (ANTs) - SourceForge.net. (2020). Retrieved 18 June 2020, from <https://sourceforge.net/projects/advants/files/>

Anbeek, P., Vincken, K. L., van Osch, M. J., Bisschops, R. H., & van der Grond, J. (2004). Probabilistic segmentation of white matter lesions in MR imaging. *NeuroImage*, 21(3), 1037–1044. doi: 10.1016/j.neuroimage.2003.10.012

Akudjedu, T. N., Nabulsi, L., Makelyte, M., Scanlon, C., Hehir, S., Casey, H., Ambati, S., Kenney, J., O'Donoghue, S., McDermott, E., Kilmartin, L., Dockery, P., McDonald, C., Hallahan, B., & Cannon, D. M. (2018). A comparative study of segmentation techniques for the quantification of brain subcortical volume. *Brain imaging and behavior*, 12(6), 1678–1695. doi: 10.1007/s11682-018-9835-y

ANTsX/ANTs. (2020). Retrieved 18 June 2020, from <https://github.com/ANTsX/ANTs>



## White Matter Hyperintensities Segmentation Review

Atlason, H. E., Love, A., Sigurdsson, S., Gudnason, V., & Ellingsen, L. M. (2019). SegAE: Unsupervised white matter lesion segmentation from brain MRIs using a CNN autoencoder. *NeuroImage: Clinical*, 24, 102085. doi: 10.1016/j.nicl.2019.102085

Aubert-Broche, B., Fonov, V., García-Lorenzo, D., Mouiha, A., Guizard, N., & Coupé, P. et al. (2013). A new method for structural volume analysis of longitudinal brain MRI data and its application in studying the growth trajectories of anatomical brain structures in childhood. *Neuroimage*, 82, 393-402. doi: 10.1016/j.neuroimage.2013.05.065

Ayrignac, X., Menjot de Champfleur, N., Menjot de Champfleur, S., Carra-Dallière, C., Deverdun, J., Corlobe, A., & Labauge, P. (2016). Brain magnetic resonance imaging helps to differentiate atypical multiple sclerosis with cavitory lesions and vanishing white matter disease. *European Journal Of Neurology*, 23(6), 995-1000. doi: 10.1111/ene.12931

Bauer, S., Nolte, L. P., & Reyes, M. (2011). Skull-stripping for tumour-bearing brain images. Annual Meeting of Swiss Society for Biomedical Engineering (Bern), page 2, Bern, April 2011. SSBE

Blair, G., Valdés Hernández, M., Thrippleton, M., Doubal, F., & Wardlaw, J. (2017). Advanced Neuroimaging of Cerebral Small Vessel Disease. *Current Treatment Options In Cardiovascular Medicine*, 19(7). doi: 10.1007/s11936-017-0555-1

Bowles, C., Qin, C., Guerrero, R., Gunn, R., Hammers, A., & Dickie, D. et al. (2017). Brain lesion segmentation through image synthesis and outlier detection. *Neuroimage: Clinical*, 16, 643-658. doi: 10.1016/j.nicl.2017.09.003

Byrne, N., Velasco Forte, M., Tandon, A., Valverde, I., & Hussain, T. (2016). A systematic review of image segmentation methodology, used in the additive manufacture of patient-specific 3D printed models of the cardiovascular system. *JRSM cardiovascular disease*, 5, 2048004016645467. doi: 10.1177/2048004016645467

## White Matter Hyperintensities Segmentation Review

Challen, R., Denny, J., Pitt, M., Gompels, L., Edwards, T., & Tsaneva-Atanasova, K. (2019). Artificial intelligence, bias and clinical safety. *BMJ Quality & Safety*, 28(3), 231-237. doi: 10.1136/bmjqs-2018-008370

Caligiuri, M. E., Perrotta, P., Augimeri, A., Rocca, F., Quattrone, A., & Cherubini, A. (2015). Automatic Detection of White Matter Hyperintensities in Healthy Aging and Pathology Using Magnetic Resonance Imaging: A Review. *Neuroinformatics*, 13(3), 261–276. doi: 10.1007/s12021-015-9260-y

Cardoso, M. J., Leung, K., Modat, M., Keihaninejad, S., Cash, D., Barnes, J., Fox, N. C., Ourselin, S., & Alzheimer's Disease Neuroimaging Initiative (2013). STEPS: Similarity and Truth Estimation for Propagated Segmentations and its application to hippocampal segmentation and brain parcellation. *Medical Image Analysis*, 17(6), 671–684. doi: 10.1016/j.media.2013.02.006

Cardoso, M. J., Modat, M., Wolz, R., Melbourne, A., Cash, D., Rueckert, D., & Ourselin, S. (2015). Geodesic information flows: Spatiallyvariant graphs and their application to segmentation and fusion. *IEEE Transactions on Medical Imaging*, 34(9), 1976–1988. doi: 10.1109/TMI.2015.2418298

Coleman, T., & Li, Y. (1996). An Interior Trust Region Approach for Nonlinear Minimization Subject to Bounds. *SIAM Journal On Optimization*, 6(2), 418-445. doi: 10.1137/0806023

Coupe, P., Yger, P., Prima, S., Hellier, P., Kervrann, C., Barillot, C. (2008). An optimized blockwise nonlocal means denoising filter for 3-D magnetic resonance images. *IEEE Transactions on Medical Imaging*, 27(4), 425-441. doi:10.1109/TMI.2007.906087

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*, 29(3), 162–173. doi: 10.1006/cbmr.1996.0014

## White Matter Hyperintensities Segmentation Review

Cruz, H., Eckert, M., Meneses, J. M., & Martínez, J. F. (2017). Fast evaluation of segmentation quality with parallel computing. *Scientific Programming*, 1-9. doi: 10.1155/2017/5767521

Dadar, M., Pascoal, T. A., Manitsirikul, S., Misquitta, K., Fonov, V. S., Tartaglia, M. C., Breitner, J., Rosa-Neto, P., Carmichael, O. T., Decarli, C., & Collins, D. L. (2017a). Validation of a Regression Technique for Segmentation of White Matter Hyperintensities in Alzheimer's Disease. *IEEE Transactions on Medical Imaging*, 36(8), 1758–1768. doi: 10.1109/TMI.2017.2693978

Dadar, M., Maranzano, J., Misquitta, K., Anor, C. J., Fonov, V. S., Tartaglia, M. C., Carmichael, O. T., De Carli, C., Collins, D. L., & Alzheimer's Disease Neuroimaging Initiative (2017b). Performance comparison of 10 different classification techniques in segmenting white matter hyperintensities in aging. *NeuroImage*, 157, 233–249. doi: [10.1016/j.neuroimage.2017.06.009](https://doi.org/10.1016/j.neuroimage.2017.06.009)

Damangir, S., Westman, E., Simmons, A., Vrenken, H., Wahlund, L. O., & Spulber, G. (2017). Reproducible segmentation of white matter hyperintensities using a new statistical definition. *Magma (New York, N.Y.)*, 30(3), 227–237. doi: 10.1007/s10334-016-0599-3

Debette, S., & Markus, H. (2010). The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: systematic review and meta-analysis. *BMJ*, 341(jul26 1), c3666-c3666. doi: 10.1136/bmj.c3666

De Boer, R., van der Lijn, F., Vrooman, H., Vernooij, M., Ikram, M., Breteler, M., & Niessen, W. (2007). Automatic Segmentation of Brain Tissue and White Matter Lesions in MRI. *Proceedings of the 2007 IEEE International Symposium on Biomedical Imaging: From Nano to Macro, Washington, DC, USA, April 12-16, 2007*. doi: 10.1109/ISBI.2007.356936

Despotović, I., Goossens, B., & Philips, W. (2015). MRI Segmentation of the Human Brain: Challenges, Methods, and Applications. *Computational and Mathematical Methods In Medicine*, 2015, 1-23. doi: 10.1155/2015/450341

## White Matter Hyperintensities Segmentation Review

Ding, T., Cohen, A. D., O'Connor, E. E., Karim, H. T., Crainiceanu, A., Muschelli, J., Lopez, O., Klunk, W. E., Aizenstein, H. J., Krafty, R., Crainiceanu, C. M., & Tudorascu, D. L. (2020). An improved algorithm of white matter hyperintensity detection in elderly adults. *NeuroImage. Clinical*, 25, 102151. doi: 10.1016/j.nicl.2019.102151

Diniz, P. H. B., Valente, T. L. A., Diniz, J. O. B., Silva, A. C., Gattass, M., Ventura, N., ... Gasparetto, E. L. (2018). Detection of white matter lesion regions in MRI using SLIC0 and convolutional neural network. *Computer Methods and Programs in Biomedicine*, 167, 49-63. doi: 10.1016/j.cmpb.2018.04.011

Documentation/4.10/Training - Slicer Wiki. (2020). Retrieved 18 June 2020, from <https://www.slicer.org/wiki/Documentation/UserTraining>

EzyFit 2.44. (2020). Retrieved 18 June 2020, from <https://www.mathworks.com/matlabcentral/fileexchange/10176-ezyfit-2-44>

Fazekas, F., & Wardlaw, J. (2013). The Origin of White Matter Lesions. *Stroke*, 44(4), 951-952. doi: 10.1161/STROKEAHA.111.000849

Fiford, C. M., Sudre, C. H., Pemberton, H., Walsh, P., Manning, E., Malone, I. B., Nicholas, J., Bouvy, W. H., Carmichael, O. T., Biessels, G. J., Cardoso, M. J., & Barnes, J. (2020). Automated white matter Hyperintensity segmentation using Bayesian model selection: Assessment and correlations with cognitive change. *Neuroinformatics*, 18(3), 429-449. doi: 10.1007/s12021-019-09439-6

FreeSurferWiki - Free Surfer Wiki. (2020). Retrieved 18 June 2020, from <https://surfer.nmr.mgh.harvard.edu/fswiki>

Gail, M., & Green, S. (1976). Critical Values for the One-Sided Two-Sample Kolmogorov-Smirnov Statistic. *Journal Of The American Statistical Association*, 71(355), 757-760. doi: 10.1080/01621459.1976.10481562

## White Matter Hyperintensities Segmentation Review

García-Lorenzo, D., Francis, S., Narayanan, S., Arnold, D. L., & Collins, D. L. (2013). Review of automatic segmentation methods of multiple sclerosis white matter lesions on conventional magnetic resonance imaging. *Medical Image Analysis*, 17(1), 1–18. doi:

10.1016/j.media.2012.09.004

Ghafoorian, M., Karssemeijer, N., Heskes, T., van Uden, I., Sanchez, C. I., Litjens, G., de Leeuw, F. E., van Ginneken, B., Marchiori, E., & Platel, B. (2017). Location Sensitive Deep Convolutional Neural Networks for Segmentation of White Matter Hyperintensities. *Scientific Reports*, 7(1), 5110. doi: 10.1038/s41598-017-05300-5

Griffanti, L., Zamboni, G., Khan, A., Li, L., Bonifacio, G., Sundaresan, V., Schulz, U. G., Kuker, W., Battaglini, M., Rothwell, P. M., & Jenkinson, M. (2016). BIANCA (Brain Intensity AbNormality Classification Algorithm): A new tool for automated segmentation of white matter hyperintensities. *NeuroImage*, 141, 191–205. doi: 10.1016/j.neuroimage.2016.07.018

Guerrero, R., Qin, C., Oktay, O., Bowles, C., Chen, L., Joules, R., Wolz, R., Valdés-Hernández, M. C., Dickie, D. A., Wardlaw, J., & Rueckert, D. (2017). White matter hyperintensity and stroke lesion segmentation and differentiation using convolutional neural networks. *NeuroImage. Clinical*, 17, 918–934. doi: 10.1016/j.nicl.2017.12.022

Hachinski, V.C., Potter, P., & Merskey, H. (1987) Leuko-araiosis. *Archives of Neurology*, 44(1), 21-3. doi: 10.1001/archneur.1987.00520130013009

Haralick, R., Shapiro, L. (1991). Glossary of computer vision terms. *Pattern Recognition*, 24(1), 69-93. doi: 10.1016/0031-3203(91)90117-n

Hasan, T. F., Barrett, K. M., Brott, T. G., Badi, M. K., Lesser, E. R., Hodge, D. O., & Meschia, J. F. (2019). Severity of White Matter Hyperintensities and Effects on All-Cause Mortality in the Mayo Clinic Florida Familial Cerebrovascular Diseases Registry. *Mayo Clinic Proceedings*, 94(3), 408–416. doi: 10.1016/j.mayocp.2018.10.024

## White Matter Hyperintensities Segmentation Review

Heckemann, R. A., Ledig, C., Gray, K. R., Aljabar, P., Rueckert, D., Hajnal, J. V., & Hammers, A. (2015). Brain Extraction Using Label Propagation and Group Agreement: Pincram. *Plos One*, 10(7). doi: 10.1371/journal.pone.0129211

Hong, J., Park, B., Lee, M. J., Chung, C., Cha, J., & Park, H. (2020). Two-step deep neural network for segmentation of deep white matter hyperintensities in migraineurs. *Computer Methods and Programs in Biomedicine*, 183, 105065. doi: 10.1016/j.cmpb.2019.105065

Huppertz H.-J., Wagner J., Weber B., House P., Urbach H. (2011). Automated quantitative FLAIR analysis in hippocampal sclerosis. *Epilepsy Research*, 97, 146-156. doi: 10.1016/j.epilepsyres.2011.08.001

Imaging in COVID-19 complications - ESR Connect. (2020). Retrieved 14 June 2020, from <https://connect.myesr.org/course/imaging-in-covid-19-complications/>

Inzitari, D. (2003). Leukoaraiosis. *Stroke*, 34(8), 2067-2071. doi: 10.1161/01.STR.0000080934.68280.82

Jenkinson, M., & Smith, S. (2001). A global optimisation method for robust affine registration of brain images. *Medical Image Analysis*, 5(2), 143–156. doi: 10.1016/s1361-8415(01)00036-6

Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the robust and accurate linear registration and motion correction of brain images. *NeuroImage*, 17(2), 825–841. doi: 10.1016/s1053-8119(02)91132-8

Jenkinson, M., Pechaud, M., Smith, S., M. (2004). BET2: MR-based estimation of brain, skull and scalp surfaces. *Eleventh Annual Meeting of the Organization for Human Brain Mapping, Toronto, Ontario*, p. 716

Jiang, J., Liu, T., Zhu, W., Koncz, R., Liu, H., Lee, T., Sachdev, P. S., & Wen, W. (2018). UBO Detector - A cluster-based, fully automated pipeline for extracting white matter hyperintensities. *NeuroImage*, 174, 539–549. doi: 10.1016/j.neuroimage.2018.03.050

Jude Hemanth, D., & Anitha, J. (2012). Image Pre-processing and Feature Extraction Techniques for Magnetic Resonance Brain Image Analysis. *Communications in Computer and Information Science*, 349-356. doi: 10.1007/978-3-642-35594-3\_47

Kamnitsas, K., Ledig, C., Newcombe, V., Simpson, J. P., Kane, A. D., Menon, D. K., Rueckert, D., & Glocker, B. (2017). Efficient multi-scale 3D CNN with fully connected CRF for accurate brain lesion segmentation. *Medical Image Analysis*, 36, 61–78. doi: 10.1016/j.media.2016.10.004

Keller, S. S., & Roberts, N. (2009). Measurement of brain volume using MRI: software, techniques, choices and prerequisites. *Journal of anthropological sciences = Rivista di antropologia : JASS*, 87, 127–151.

Kim, K. W., MacFall, J. R., & Payne, M. E. (2008). Classification of white matter lesions on magnetic resonance imaging in elderly persons. *Biological psychiatry*, 64(4), 273–280. doi: 10.1016/j.biopsych.2008.03.024

Klein, S., Staring, M., Murphy, K., Viergever, M. A., & Pluim, J. P. (2010). elastix: a toolbox for intensity-based medical image registration. *IEEE Transactions on Medical Imaging*, 29(1), 196–205. doi: 10.1109/TMI.2009.2035616

Knight, J., Taylor, G., & Khademi, A. (2018). Voxel-Wise Logistic Regression and Leave-One-Source-Out Cross Validation for white matter hyperintensity segmentation. *Magnetic Resonance Imaging*, 54, 119-136. doi: 10.1016/j.mri.2018.06.009

Kristensen, N., Nymann, C., & Konradsen, H. (2015). Implementing research results in clinical practice- the experiences of healthcare professionals. *BMC Health Services Research*, 16(1). doi: 10.1186/s12913-016-1292-y

Kuijff, H. J., Biesbroek, J. M., De Bresser, J., Heinen, R., Andermatt, S., Bento, M., Berseth, M., et al. (2019). Standardized Assessment of Automatic Segmentation of White Matter

## White Matter Hyperintensities Segmentation Review

Hyperintensities and Results of the WMH Segmentation Challenge. *IEEE Transactions on Medical Imaging*, 38(11), 2556–2568. doi: 10.1109/TMI.2019.2905770

Kwan, R. K. S., Evans, A. C., & Pike, B. (1999). MRI simulation-based evaluation of image-processing and classification methods. *IEEE Transactions on Medical Imaging*, 18 (11), 1085–1097. doi: 10.1109/42.816072

Labauge, P., Horzinski, L., Ayrignac, X., Blanc, P., Vukusic, S., Rodriguez, D., Mauguier, F., *et al.* (2009). Natural history of adult-onset eIF2B-related disorders: a multi-centric survey of 16 cases. *Brain : a journal of neurology*, 132(Pt 8), 2161–2169. doi: 10.1093/brain/awp171

Li, H., Jiang, G., Zhang, J., Wang, R., Wang, Z., Zheng, W. S., & Menze, B. (2018). Fully convolutional network ensembles for white matter hyperintensities segmentation in MR images. *NeuroImage*, 183, 650–665. doi: 10.1016/j.neuroimage.2018.07.005

Liu, W., Hua, G., & Smith, J. (2014). Unsupervised One-Class Learning for Automatic Outlier Removal. *2014 IEEE Conference On Computer Vision And Pattern Recognition*. doi: 10.1109/cvpr.2014.483

Ling, Y., Jouvent, E., Cousyn, L., Chabriat, H., & De Guio, F. (2018). Validation and Optimization of BIANCA for the Segmentation of Extensive White Matter Hyperintensities. *Neuroinformatics*, 16(2), 269–281. doi: 10.1007/s12021-018-9372-2

Liu, L., Chen, S., Zhu, X., Zhao, X., Wu, F., & Wang, J. (2020). Deep convolutional neural network for accurate segmentation and quantification of white matter hyperintensities. *Neurocomputing*, 384, 231–242. doi: 10.1016/j.neucom.2019.12.050

Lladó, X., Ganiler, O., Oliver, A., Martí, R., Freixenet, J., & Valls, L. *et al.* (2011). Automated detection of multiple sclerosis lesions in serial brain MRI. *Neuroradiology*, 54(8), 787–807. doi: 10.1007/s00234-011-0992-6



## White Matter Hyperintensities Segmentation Review

Llado, X., Oliver, A., Cabezas, M., Freixenet, J., Vilanova, J. C., Quiles, A., Valls, L., Ramió-Torrentà, Ll., Rovira, A. (2012). Segmentation of multiple sclerosis lesions in brain MRI: A review of automated approaches. *Information Sciences*, 186, 164-185. doi: 10.1016/j.ins.2011.10.011

Lutkenhoff, E., Rosenberg, M., Chiang, J., Zhang, K., Pickard, J., Owen, A., & Monti, M. (2014). Optimized Brain Extraction for Pathological Brains (optiBET). *Plos ONE*, 9(12), e115551. doi: 10.1371/journal.pone.0115551

Maino, G., & Foresti, G. (2011). *Image analysis and processing--ICIAP 2011*. Heidelberg: Springer. doi: 10.1007/978-3-642-24085-0

Maltais, M., de Souto Barreto, P., Moon, S. Y., Rolland, Y., Vellas, B., & MAPT/DSA Study Group (2019). Prospective association of white matter hyperintensity volume and frailty in older adults. *Experimental gerontology*, 118, 51–54. doi: 10.1016/j.exger.2019.01.007

Maniega, S. M., Valdés Hernández, M. C., Clayden, J. D., Royle, N. A., Murray, C., Morris, Z., Aribisala, B. S., Gow, A. J., Starr, J. M., Bastin, M. E., Deary, I. J., & Wardlaw, J. M. (2015). White matter hyperintensities and normal-appearing white matter integrity in the aging brain. *Neurobiology of aging*, 36(2), 909–918. doi: 10.1016/j.neurobiolaging.2014.07.048

Manjón, J. V., Coupé, P., Martí-Bonmatí, L., Collins, D. L., & Robles, M. (2010). Adaptive non-local means denoising of MR images with spatially varying noise levels. *Journal of magnetic resonance imaging : JMRI*, 31(1), 192–203. doi: 10.1002/jmri.22003

Manjón, J. V., Coupé, P., Raniga, P., Xia, Y., Desmond, P., Fripp, J., & Salvado, O. (2018). MRI white matter lesion segmentation using an ensemble of neural networks and overcomplete patch-based voting. *Computerized Medical Imaging and Graphics : the official journal of the Computerized Medical Imaging Society*, 69, 43–51. doi: 10.1016/j.compmedimag.2018.05.001

Martin Bland, J., & Altman, D. (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *The Lancet*, 1 (8476), 307-310.

## White Matter Hyperintensities Segmentation Review

Mendelson, A., Zuluaga, M., Lorenzi, M., Hutton, B., & Ourselin, S. (2017). Selection bias in the reported performances of AD classification pipelines. *Neuroimage: Clinical*, 14, 400-416. doi: 10.1016/j.nicl.2016.12.018

Miller, D. H., Grossman, R. I., Reingold, S. C., & McFarland, H. F. (1998). The role of magnetic resonance techniques in understanding and managing multiple sclerosis. *Brain : a journal of neurology*, 121 ( Pt 1), 3–24. doi: 10.1093/brain/121.1.3

MIRTK – BioMedIA. (2020). Retrieved 18 June 2020, from <https://biomedica.doc.ic.ac.uk/software/mirtk/>

Modat, M., Cash, D. M., Daga, P., Winston, G. P., Duncan, J. S., & Ourselin, S. (2014). Global image registration using a symmetric block-matching approach. *Journal of medical imaging (Bellingham, Wash.)*, 1(2), 024003. doi: 10.1117/1.JMI.1.2.024003

Mortazavi, D., Kouzani, A. Z., & Soltanian-Zadeh, H. (2012). Segmentation of multiple sclerosis lesions in MR images: a review. *Neuroradiology*, 54(4), 299–320. doi: 10.1007/s00234-011-0886-7

Moeskops, P., Viergever, M., Mendrik, A., de Vries, L., Benders, M., & Išgum, I. (2016). Automatic Segmentation of MR Brain Images With a Convolutional Neural Network. *IEEE Transactions On Medical Imaging*, 35(5), 1252-1261.

Moeskops, P., Bresser, J. D., Kuijf, H. J., Mendrik, A. M., Biessels, G. J., Pluim, J. P., & Išgum, I. (2018). Evaluation of a deep learning approach for the segmentation of brain tissues and white matter hyperintensities of presumed vascular origin in MRI. *NeuroImage: Clinical*, 17, 251–262. doi: 10.1109/TMI.2016.2548501

MRI Shows Brain Abnormalities in Some COVID-19 Patients. (2020). Retrieved 14 June 2020, from <https://www.diagnosticimaging.com/covid-19/mri-shows-brain-abnormalities-some-covid-19-patients>

## White Matter Hyperintensities Segmentation Review

Muschelli, J., Gherman, A., Fortin, J. P., Avants, B., Whitcher, B., Clayden, J. D., Caffo, B. S., & Crainiceanu, C. M. (2019). Neuroconductor: an R platform for medical imaging analysis. *Biostatistics* (Oxford, England), 20(2), 218–239. doi: 10.1093/biostatistics/kxx068

NITRC: MRlcron: Document Manager: Display Document. (2020). Retrieved 18 June 2020, from [https://www.nitrc.org/docman/?group\\_id=152](https://www.nitrc.org/docman/?group_id=152)

Nyúl, L. G., & Udupa, J. K. (1999). On standardizing the MR image intensity scale. *Magnetic Resonance in Medicine*, 42(6), 1072–1081. doi: 10.1002/(sici)1522-2594(199912)42:6<1072::aid-mrm11>3.0.co;2-m

Park, B. Y., Lee, M. J., Lee, S. H., Cha, J., Chung, C. S., Kim, S. T., & Park, H. (2018). DEWS (DEep White matter hyperintensity Segmentation framework): A fully automated pipeline for detecting small deep white matter hyperintensities in migraineurs. *NeuroImage. Clinical*, 18, 638–647. doi: 10.1016/j.nicl.2018.02.033

Pellegrini, E., Ballerini, L., Hernandez, M., Chappell, F. M., González-Castro, V., Anblagan, D., Danso, S., Muñoz-Maniega, S., Job, D., Pernet, C., Mair, G., MacGillivray, T. J., Trucco, E., & Wardlaw, J. M. (2018). Machine learning of neuroimaging for assisted diagnosis of cognitive impairment and dementia: A systematic review. *Alzheimer's & dementia (Amsterdam, Netherlands)*, 10, 519–535. doi: 10.1016/j.dadm.2018.07.004

Qin, C., Guerrero, R., Bowles, C., Chen, L., Dickie, D., & Valdes-Hernandez, M. et al. (2018). A large margin algorithm for automated segmentation of white matter hyperintensity. *Pattern Recognition*, 77, 150-159. doi: 10.1016/j.patcog.2017.12.016

Rachmadi, M. F., Valdés-Hernández, M. D., Li, H., Guerrero, R., Meijboom, R., Wiseman, S., Waldman, A., Zhang, J., Rueckert, D., Wardlaw, J., & Komura, T. (2020). Limited one-time sampling irregularity map (LOTS-IM) for automatic unsupervised assessment of white matter Hyperintensities and multiple sclerosis lesions in structural brain magnetic resonance

## White Matter Hyperintensities Segmentation Review

images. *Computerized Medical Imaging and Graphics*, 79, 101685. doi: 10.1016/j.compmedimag.2019.101685

Rachmadi, M. F., Valdés-Hernández, M., Agan, M., Di Perri, C., Komura, T., & Alzheimer's Disease Neuroimaging Initiative (2018). Segmentation of white matter hyperintensities using convolutional neural networks with global spatial information in routine clinical brain MRI with none or mild vascular pathology. *Computerized medical imaging and graphics: the official journal of the Computerized Medical Imaging Society*, 66, 28–43. doi: 10.1016/j.compmedimag.2018.02.002

Rachmadi, M., Valdés-Hernández, M., Agan, M., & Komura, T. (2017). Deep Learning vs. Conventional Machine Learning: Pilot Study of WMH Segmentation in Brain MRI with Absence or Mild Vascular Pathology. *Journal Of Imaging*, 3(4), 66. doi: 10.3390/jimaging3040066

Reporting guidelines | The EQUATOR Network. (2020). Retrieved 5 July 2020, from <https://www.equator-network.org/reporting-guidelines/>

Rincón, M., Díaz-López, E., Selnes, P., Vegge, K., Altmann, M., Fladby, T., & Bjørnerud, A. (2017). Improved Automatic Segmentation of White Matter Hyperintensities in MRI Based on Multilevel Lesion Features. *Neuroinformatics*, 15(3), 231–245. doi: 10.1007/s12021-017-9328-y

Roy, P., Bhuiyan, A., Janke, A., Desmond, P., Wong, T-Y., Abhayaratna, W., & Ramamohanarao, K. (2015). Automatic white matter lesion segmentation using contrast enhanced FLAIR intensity and Markov Random Field. *Computerized Medical Imaging and Graphics*. 45. doi: 10.1016/j.compmedimag.2015.08.005.

Roy, S., Butman, J. A., Pham, D. L., & Alzheimers Disease Neuroimaging Initiative (2017). Robust skull stripping using multiple MR image contrasts insensitive to pathology. *NeuroImage*, 146, 132–147. doi: 10.1016/j.neuroimage.2016.11.017

## White Matter Hyperintensities Segmentation Review

Samuel, G., Hoffmann, S., Wright, R., Lalu, M., Patlewicz, G., & Becker, R. et al. (2016). Guidance on assessing the methodological and reporting quality of toxicologically relevant studies: A scoping review. *Environment International*, 92-93, 630-646. doi: 10.1016/j.envint.2016.03.010

Schirmer, M. D., Dalca, A. V., Sridharan, R., Giese, A. K., Donahue, K. L., Nardin, M. J., Mocking, S., McIntosh, E. C., Frid, P., Wasselius, J., Cole, J. W., Holmegaard, L., Jern, C., Jimenez-Conde, J., Lemmens, R., Lindgren, A. G., Meschia, J. F., Roquer, J., Rundek, T., Sacco, R. L., ..., MRI-GENIE Investigators (2019). White matter hyperintensity quantification in large-scale clinical acute ischemic stroke cohorts - The MRI-GENIE study. *NeuroImage. Clinical*, 23, 101884. doi: 10.1016/j.nicl.2019.101884

Schmidt, R. L., & Factor, R. E. (2013). Understanding sources of bias in diagnostic accuracy studies. *Archives of pathology & laboratory medicine*, 137(4), 558–565. doi: 10.5858/arpa.2012-0198-RA

Shamonin, D. P., Bron, E. E., Lelieveldt, B. P., Smits, M., Klein, S., Staring, M., & Alzheimer's Disease Neuroimaging Initiative (2014). Fast parallel image registration on CPU and GPU for diagnostic classification of Alzheimer's disease. *Frontiers in neuroinformatics*, 7, 50. doi: 10.3389/fninf.2013.00050

Shinohara, R. T., Goldsmith, J., Mateen, F. J., Crainiceanu, C., & Reich, D. S. (2012). Predicting breakdown of the blood-brain barrier in multiple sclerosis without contrast agents. *AJNR. American journal of neuroradiology*, 33(8), 1586–1590. doi: 10.3174/ajnr.A2997

Sled, J. G., Zijdenbos, A. P., & Evans, A. C. (1998). A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE transactions on medical imaging*, 17(1), 87–97. doi: 10.1109/42.668698

Smith, S. (2002). Fast robust automated brain extraction. *Human Brain Mapping*, 17(3), 143-155. doi: 10.1002/hbm.10062

## White Matter Hyperintensities Segmentation Review

SPM - Documentation. (2020). Retrieved 18 June 2020, from

<https://www.fil.ion.ucl.ac.uk/spm/doc/>

Steenwijk, M. D., Pouwels, P. J., Daams, M., van Dalen, J. W., Caan, M. W., Richard, E., Barkhof, F., & Vrenken, H. (2013). Accurate white matter lesion segmentation by k nearest neighbor classification with tissue type priors (kNN-TTPs). *NeuroImage. Clinical*, 3, 462–469. doi: 10.1016/j.nicl.2013.10.003

Stone, J. R., Wilde, E. A., Taylor, B. A., Tate, D. F., Levin, H., Bigler, E. D., Scheibel, R. S., Newsome, M. R., Mayer, A. R., Abildskov, T., Black, G. M., Lennon, M. J., York, G. E., Agarwal, R., DeVillasante, J., Ritter, J. L., Walker, P. B., Ahlers, S. T., & Tustison, N. J. (2016). Supervised learning technique for the automated identification of white matter hyperintensities in traumatic brain injury. *Brain injury*, 30(12), 1458–1468. doi: 10.1080/02699052.2016.1222080

Styner, M., Lee, J., Chin, B., Chin, M., Commowick, O., Tran, H-H., Jewells, V., & Warfield, S. (2007). 3D segmentation in the clinic: A grand challenge II: MS lesion segmentation. MIDAS Journal. Accessed 20 June 2020 from <http://hdl.handle.net/10380/1449>

Sudre, C. H., Cardoso, M. J., Bouvy, W. H., Biessels, G. J., Barnes, J., & Ourselin, S. (2015). Bayesian model selection for pathological neuroimaging data applied to white matter lesion segmentation. *IEEE Transactions on Medical Imaging*, 34(10), 2079–2102. doi: 10.1109/TMI.2015.2419072

Sudre, C. H., Cardoso, M. J., Ourselin, S., & Alzheimer's Disease Neuroimaging Initiative (2017). Longitudinal segmentation of age-related white matter hyperintensities. *Medical Image Analysis*, 38, 50–64. doi: 10.1016/j.media.2017.02.007

Sundaresan, V., Zamboni, G., Le Heron, C., Rothwell, P., Husain, M., & Battaglini, M. et al. (2019). Automated lesion segmentation with BIANCA: Impact of population-level features, classification algorithm and locally adaptive thresholding. *Neuroimage*, 202, 116056. doi: 10.1016/j.neuroimage.2019.116056

## White Matter Hyperintensities Segmentation Review

Sweeney, E. M., Shinohara, R. T., Shiee, N., Mateen, F. J., Chudgar, A. A., Cuzzocreo, J. L., Calabresi, P. A., Pham, D. L., Reich, D. S., & Crainiceanu, C. M. (2013). OASIS is automated statistical inference for segmentation, with applications to multiple sclerosis lesion segmentation in MRI. *NeuroImage: Clinical*, 2, 402-413. doi: 10.1016/j.nicl.2013.03.002

Valdés Hernández, M.C., Ferguson, K. J., Chappell, F. M., Wardlaw, J. M. (2010). New multispectral MRI data fusion technique for white matter lesion segmentation: method and comparison with thresholding in FLAIR images. *European Radiology*, 20(7), 1684-1691. doi: 10.1007/s00330-010-1718-6

Valverde, S., Oliver, A., Roura, E., González-Villà, S., Pareto, D., Vilanova, J. C., Ramió-Torrentà, L., Rovira, À., & Lladó, X. (2017). Automated tissue segmentation of MR brain images in the presence of white matter lesions. *Medical image analysis*, 35, 446–457. doi: 10.1016/j.media.2016.08.014

Van Leemput, K., Maes, F., Vandermeulen, D., & Suetens, P. (1999). Automated model-based tissue classification of MR images of the brain. *IEEE Transactions on Medical Imaging*, 18(10), 897–908. doi: 10.1109/42.811270

Van Opbroek, A., Ikram, M., Vernooij, M., & de Bruijne, M. (2015a). Transfer Learning Improves Supervised Image Segmentation Across Imaging Protocols. *IEEE Transactions On Medical Imaging*, 34(5), 1018-1030. doi: 10.1109/TMI.2014.2366792

Van Opbroek, A. V., Vernooij, M. W., Ikram, M. A., & Bruijne, M. (2015b). Weighting training images by maximizing distribution similarity for supervised segmentation across scanners. *Medical image analysis*, 24(1), 245–254. doi: 10.1016/j.media.2015.06.010

“VisibleHuman.png.” *MRICro* / *CRNL*, [www.mccauslandcenter.sc.edu/crnl/mricro](http://www.mccauslandcenter.sc.edu/crnl/mricro).

Viola, P., Wells III, W. M. (1997) Alignment by maximization of mutual information. *IntJ Comput Vis*, 24(2), 137-154. doi: 10.1109/ICCV.1995.466930

## White Matter Hyperintensities Segmentation Review

Wang, R., Li, C., Wang, J., Wei, X., Li, Y., Zhu, Y., & Zhang, S. (2015). Automatic segmentation and volumetric quantification of white matter hyperintensities on fluid-attenuated inversion recovery images using the extreme value distribution. *Neuroradiology*, 57(3), 307–320. doi: 10.1007/s00234-014-1466-4

Warfield, S., Kaus, M., Jolesz, F., & Kikinis, R. (2000). Adaptive, template moderated, spatially varying statistical classification. *Medical Image Analysis*, 4(1), 43-55. doi: 10.1016/s1361-8415(00)00003-7

Wardlaw, J. M., Smith, E. E., Biessels, G. J., Cordonnier, C., Fazekas, F., Frayne, R., Lindley, R. I., O'Brien, J. T., Barkhof, F., Benavente, O. R., Black, S. E., Brayne, C., Breteler, M., Chabriat, H., Decarli, C., de Leeuw, F. E., Doubal, F., Duering, M., Fox, N. C., Greenberg, S., ... STandards for Reporting Vascular changes on nEuroimaging (STRIVE v1) (2013). Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. *The Lancet. Neurology*, 12(8), 822–838. doi: 10.1016/S1474-4422(13)70124-8

Wardlaw, J., Valdés Hernández, M., & Muñoz-Maniega, S. (2015). What are White Matter Hyperintensities Made of?. *Journal Of The American Heart Association*, 4(6). doi: 10.1161/JAHA.114.001140

Wu, Y., & Liu, Y. (2013). Adaptively Weighted Large Margin Classifiers. *Journal Of Computational And Graphical Statistics*, 22(2), 416-432. doi: 10.1080/10618600.2012.680866

Wu, J., Zhang, Y., Wang, K., & Tang, X. (2019). Skip connection U-net for white matter Hyperintensities segmentation from MRI. *IEEE Access*, 7, 155194-155202. doi: 10.1109/access.2019.2948476

Wu, D., Albert, M., Soldan, A., Pettigrew, C., Oishi, K., Tomogane, Y., Ye, C., Ma, T., Miller, M. I., & Mori, S. (2019). Multi-atlas based detection and localization (MADL) for location-dependent quantification of white matter hyperintensities. *NeuroImage. Clinical*, 22, 101772. doi: 10.1016/j.nicl.2019.101772



## White Matter Hyperintensities Segmentation Review

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE transactions on medical imaging*, 20(1), 45–57. doi: 10.1109/42.906424

Zhan, T., Liu, Z., Xiao, L., Zhan, Y., & Wei, Z. (2015). Automatic method for white matter lesion segmentation based on T1-fluid-attenuated inversion recovery images. *IET Computer Vision*, 9(4), 447–455. doi: 10.1049/iet-cvi.2014.0121

Zhan, T., Yu, R., Zheng, Y., Zhan, Y., Xiao, L. and Wei, Z., 2017. Multimodal spatial-based segmentation framework for white matter lesions in multi-sequence magnetic resonance images. *Biomedical Signal Processing and Control*, 31, pp.52-62. doi: 10.1016/j.bspc.2016.06.016

Zheng, J., Delbaere, K., Close, J., Sachdev, P., & Lord, S. (2011). Impact of White Matter Lesions on Physical Functioning and Fall Risk in Older People. *Stroke*, 42(7), 2086-2090. doi: 10.1161/strokeaha.110.610360