

## Supplementary material

### **Molecular characteristics of extraintestinal pathogenic *E. coli* (ExPEC), uropathogenic *E. coli* (UPEC), and multidrug resistant *E. coli* isolated from healthy dogs in Spain. Whole genome sequencing of canine ST372 isolates and comparison with human isolates causing extraintestinal infections.**

Saskia-Camille Flament-Simon<sup>1,2</sup>, María de Toro<sup>3</sup>, Vanesa García<sup>1,2</sup>, Jesús E. Blanco<sup>1,2</sup>, Miguel Blanco<sup>1,2</sup>, María Pilar Alonso<sup>2,4</sup>, Ana Goicoa<sup>5,6</sup>, Juan Díaz-González<sup>1</sup>, Marie-Hélène Nicolas-Chanoine<sup>7</sup>, and Jorge Blanco<sup>1,2\*</sup>

- <sup>1</sup> Laboratorio de Referencia de *E. coli* (LREC), Departamento de Microbiología e Parasitología, Facultad de Veterinaria, Universidad de Santiago de Compostela (USC), Lugo, Spain.
- <sup>2</sup> Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Spain.
- <sup>3</sup> Plataforma de Genómica y Bioinformática, Centro de Investigación Biomédica de La Rioja (CIBIR), Logroño, Spain.
- <sup>4</sup> Unidade de Microbiología, Hospital Universitario Lucus Augusti (HULA), Lugo, Spain.
- <sup>5</sup> Servicio de Medicina Interna, Hospital Veterinario Universitario Rof Codina, USC, Lugo, Spain.
- <sup>6</sup> Departamento de Anatomía, Producción Animal e Ciencias Clínicas Veterinarias, USC, Lugo, Spain
- <sup>7</sup> Université de Paris, IAME, UMR1137, INSERM, Paris, France.

**Table S3.** Prevalence of the phylogenetic groups in the 197 canine *E. coli* isolates.

Phylogenetic group	Number of isolates (%)
A	32 (16.2)
B1	26 (13.2)
B2	84 (42.6)
C	10 (5.1)
D	6 (3.0)
E	14 (7.1)
F	18 (9.1)
Clade V	1 (0.5)
Not typeable	6 (3.0)

**Table S4.** Comparison of the distribution of the phylogenetic groups among the 197 canine isolates according to the strain ExPEC and UPEC status.

Phylogenetic group	Number of isolates (%)		P-value <sup>1</sup>
	ExPEC and/or UPEC (n= 91)	non-ExPEC and non-UPEC (n=106)	
A	2 (2.2)	30 (28.3)	<0.00001
B1	2 (2.2)	24 (22.6)	0.00001
B2	78 (85.7)	6 (5.7)	<0.00001
C	0	10 (9.4)	0.00199
D	1 (1.1)	5 (4.7)	
E	1 (1.1)	13 (12.3)	0.00371
F	7 (7.7)	11 (10.4)	
Clade V	0	1 (0.9)	
Not typeable	0	6 (5.7)	0.03155

<sup>1</sup>Two-tailed *P* values by Fisher's exact probability test are shown where *P* < 0.05.

**Table S5.** Comparison of the distribution of the phylogenetic groups among the 197 canine isolates according to the strain multidrug resistant (MDR) status.

Phylogenetic group	Number of isolates (%)		P-value <sup>1</sup>
	MDR (n= 28)	non-MDR (n=169)	
A	4 (14.3)	28 (16.6)	
B1	8 (28.6)	18 (10.7)	0.01597
B2	2 (7.1)	82 (48.5)	0.00004
C	1 (3.6)	9 (5.3)	
D	5 (17.9)	1 (0.6)	0.00022
E	5 (17.9)	9 (5.3)	0.03238
F	3 (10.7)	15 (8.9)	
Clade V	0	1 (0.6)	
Not typeable	0	6 (3.6)	

<sup>1</sup>Two-tailed *P* values by Fisher's exact probability test are shown where *P* < 0.05.

**Table S6.** Comparison of the strain ExPEC and UPEC status among canine multidrug resistant (MDR) and non-MDR isolates.

Status	Number of isolates (%)		P-value <sup>1</sup>
	MDR (n= 28)	non-MDR (n=169)	
ExPEC	8 (28.6)	66 (39.1)	
UPEC	4 (14.3)	78 (46.2)	0.00156
ExPEC and/or UPEC	8 (28.6)	83 (49.1)	

<sup>1</sup>Two-tailed *P* values by Fisher's exact probability test are shown where *P* < 0.05.

Table S7. New sequence types observed in 18 canine *E. coli* isolates.

ST New	Allele numbers						
	<i>adk</i>	<i>fumC</i>	<i>gyrB</i>	<i>icd</i>	<i>mdh</i>	<i>purA</i>	<i>recA</i>
1	10	11	4	666	560	8	2
2	13	13	9	13	16	<sup>2</sup> 126-like	9
3	13	13	9	13	439	10	9
4	88	103	19	18	23	44	26
5	88	103	<sup>1</sup> 19-like	36	23	44	26
6	6	11	4	8	235	8	2
7	13	40	9	13	16	126	9
8	13	363	302	97	23	94	93
9	13	24	9	912	17	11	25
10	36	24	10	912	17	10	25
11	9	23	33	628	11	8	6
12	13	40	459	912	23	28	109
13	13	40	13	912	23	25	66
14	21	35	27	6	286	5	4
15	6	4	4	16	474	8	14
16	6	4	5	18	11	319	6
17	6	29	33	402	11	7	2
18	63	30	3	628	43	350	6


<sup>1</sup>Nucleotide sequence of *gyrB* 19-like, closest match: *Escherichia coli* Achtman MLST *gyrB*: 19 (1 difference found: <sup>49</sup>C→<sup>49</sup>T)

```
GGTCTGCACGGCGTTGGTGTTCGGTAGTAAACGCCCTGTCGCAAAAATTGGAGCTGGTTATCCAGCGGAGGGTAAAAATCACCGTCAGATCTACGAACA
CGGTGTACCGCAGGCCCGCTGGCGTTACCGCGGAGACTGAAAAAACCGGCACTATGGTGGCGTTCTGGCCAAGCCTTGAAACCTTCACCAATGTGACCG
AGTTCGAATATGACATCTGGCGAAACGCTCGCTGAGTTGTCGTTCCTCAACTCCGGCGTTCCATTCGTCGCGGACAAGCGCGACGGCAAAGAAGAC
CACTTCCACTATGAAGCGGCATCAAGGCGTTCGTTGAATATCTGAACAAGAACAACGCGGATCCACCCGAATATCTTCTACTTCTCCACCGAAAAAGA
CGGTATTGGCGTCGAAGTGGCGTTGCAGTGGACGATGGCTTCCAGGAAAAATCT
```

<sup>2</sup>Nucleotide sequence of *purA* 126-like, closest match: *Escherichia coli* Achtman MLST *purA*: 126 (1 difference found: <sup>118</sup>A→<sup>118</sup>G)

```
ATAACGCGCGTGAGAAAGCGCGTGGCGGAAAGCGATCGGCACCACCGGTCGAGGTATCGGGCCTGCTTATGAAGATAAAGTGGCACGTCGCGGCTGCGGT
GTTGGCGACCTTTTCGGCAAAGAAACCTTCGCTGAAAAACTGAAAGAAGTGAATATCACAACTCCAGTTGGTTAACTACTACAAAGCTGAAGCGGT
TGATTACGAAAAGTTCTGGATGATACGATGGCTGTTGCCGACATCCTGACTTCTATGGTTGTTGACGTTTCTGATCTGCTCGACCAAGCGCGTCAGCGTG
GCGATTTTCGTCATGTTTGAAGTGGCGAGGTACGCTGCTGGATATCGACCACGGCACTTATCCGTACGTAACCTTCTCCAACACTACTGCTGGTGGCGTG
GCGACCGGTTCCGCGCTGGGCCCGGTTATGTTGATTACGTTTTGGGTATCCTCAAAGCTTACTCCACTCGTGT
```

Figure S1. Comparison of contigs harbouring integrative conjugative elements (ICEs) from 22 ST372 *E. coli* genomes.



<sup>1</sup>The shadow parallelograms denote genetic regions that exhibit sequence homology among different segments. Light shadow denotes regions with a lower level of sequence identity (99%) by BLAST. The image was generated using EasyFig [59] (default parameters).