Article

Callus Culture of *Ocimum basilicum* L. cv 'Thai Basil' is an Effective Biological System for the Production of Antioxidants Compared to Leaves

Saher Nazir¹, Hasnain Jan¹, Duangjai Tungmunthum^{2,3}, Samantha Drouet², Muhammad Zia¹, Christophe Hano^{3*}, Bilal Haider Abbasi^{1*}

¹ Department of Biotechnology, Quaid-i-Azam University, Islamabad, 45320, Pakistan; saher_nazir158@yahoo.com (S.N.); hasnain849@gmail.com (H.J.); ziachaudhary@gmail.com (M.Z.); bhabbasi@qau.edu.pk (B.H.A.)

² Laboratoire de Biologie des Ligneux et des Grandes Cultures, INRAE USC1328, University of Orleans, 45067 Orléans CEDEX 2, France; samantha.drouet@univ-orleans.fr (S.D.); hano@univ-orleans.fr (C.H.)

³ Department of Pharmaceutical Botany, Faculty of Pharmacy, Mahidol University, Bangkok 10400, Thailand; duangjai.tun@mahidol.ac.th (D.T.)

* Correspondence: bhabbasi@qau.edu.pk (B.H.A.); hano@univ-orleans.fr (C.H.); Tel.: +33-77-698-41-48 (B.H.A.); +33-237-309-753 (C.H.)

Abstract: Thai basil is a renowned medicinal plant and a rich source of bioactive antioxidant compounds having several health benefits, with actions to prevent of cancer, diabetes and cardiovascular disease. Plant cell and tissue culture technologies can be routinely established as an important, sustainable and low-cost biomass source for the production of high-value phytochemicals. The current study aimed at developing an effective protocol for the production of Thai basil leaf derived callus cultures with sustainable and high production of biomass and antioxidants as an alternative of leaves production. MS basal medium with various concentrations of plant growth regulators (PGRs) compatible with nutraceutical applications (i.e., gibberellic acid (GA3) and 6-benzylaminopurine (BAP) either alone or in combination with naphthalene acetic acid (NAA)) were evaluated. Among all tested PGRs, the combination BAP:NAA (5 mg/L:1 mg/L) yield maximum biomass accumulation (fresh weight (FW): 190 g/L and dry weight (DW): 13.05 g/L) as well as enhanced phenolic (346.08 mg/L) production. HPLC quantification analysis indicated high productions of chicoric acid (35.77 mg/g DW) and rosmarinic acid (7.35 mg/g DW) under optimized callus culture conditions. Antioxidant potential was assessed using both *in vitro* cell free and *in vivo* cellular antioxidant assays. Maximum *in vitro* antioxidant activity DPPH (93.2 % of radical scavenging activity) and ABTS (1322 μ M Trolox equivalent antioxidant capacity) was also observed for the extracts from callus cultures grown on optimal conditions. *In vivo* cellular antioxidant activity assay confirmed the effective protection against oxidative stress of the corresponding extract by the maximum inhibition of ROS and RNS production. Compared to commercial leaves, callus extracts showed higher production of chicoric acid and rosmarinic acid associated with higher antioxidant capacity. In addition, this biological system also has a large capacity for continuous biomass production, thus demonstrating its high potential for possible nutraceutical applications.

Keywords: Antioxidants; Biological System; Caffeic Acid; Callus culture; Chicoric Acid; Phenolics; Plant *in vitro* Cultures; Rosmarinic Acid; Thai Basil

1. Introduction

Genus *Ocimum* (Family *Lamiaceae*) includes almost 150 species of herbs and shrubs that habitats the semi tropical and tropical regions of America (Central and South), Asia and Africa [1,2]. Since prehistoric times, basil was used as flavoring agent and for fragrance in food and cosmetics [3].

Ocimum species, but also cultivars within the same species, varied—considerably in their phytochemical compositions [4]. *Ocimum basilicum* L. cv 'Thai Basil' is used as herbal medicinal products [5]. Indeed, Thai basil is a reservoir of a pharmaceutically important metabolite, such as rosmarinic acid and chicoric acid [5,6]. These plant specialized metabolites play a pivotal role in plant-environment interactions, provide biotic and abiotic stress tolerance and are used as pharmaceutical agents because they have a variety of antioxidant, anticancer, antimicrobial or antidiabetic activities [7–10].

Plant specialized metabolites of pharmacological significance may be difficult to extract in high, sustainable and continuous quantities from field-grown plant material, but *in vitro* cultures have been identified as a possible biological system for the production of these metabolites [11]. *In vitro* culture techniques provide consistent production schemes for metabolites that are free from environmental limitations and ensure increased biomass and a high concentration of compounds compared to natural production systems [12]. In addition, in the case of basil varieties or cultivars which can be differentiated on the basis of color, aromatic composition and growth habitat, sometimes harvested from nature, thus complicating its botanical identity, the *in vitro* system presents a major advantage for the certification of plant authentication [4,13]. Various reports are available in the literature on the *in vitro* production of bioactive metabolites from the callus culture of medicinally important plants. For example, Smith [14] reported on the production of shikonin by *Lithospermum erythrorhizon* Siebold & Zucc., a natural dye used in food and/or cosmetic applications, as well as of the anticancer drug paclitaxel by *Taxus* L. species or the antimicrobial berberine by *Coptis japonica* Wall. cultures.

Previously, different varieties or cultivars of *O. basilicum* L. have been used to develop different types of *in vitro* cultures [10,15,16]. Following our works on *O. basilicum* L. cv Purple basil [10,17,18], the current study aimed at establishing a proficient protocol for Thai basil *in vitro* cultures that are effective in producing continuously high biomass and accumulating high levels of antioxidants under the treatment of various PGRs, either alone or in combination. Culture extracts were further analyzed for total phenolic production, HPLC quantification of chicoric acid, rosmarinic acid, and caffeic acid, as well as antioxidant capacity using both *in vitro* cell-free (DPPH, ABTS and FRAP assays) and cellular assays. These values were compared with those obtained from commercial leaves for extract. In the light of these results, the nutraceutical potential of the established Thai basil *in vitro* cultures is discussed.

2. Materials and Methods

2.1. Chemicals and Reagents

In the current research, solvents of analytical grade were used, supplied by Thermo Scientific (Illkirch, France). All reagents and standards were bought from Merck (Saint-Quentin Fallavier, France).

2.2. Plant Materials and Callus Culture Establishment

Seeds of Thai basil were obtained from National Agriculture Research Center (NARC, Islamabad, Pakistan). Procedure of Abbasi *et al.* [19] was adopted for seed surface sterilization. Seeds were immersed for 30 sec in mercuric chloride (0.1%) and afterward in ethanol and washed thrice with distilled water. MS basal medium [20] with agar (8 g/L) and sucrose (30 g/L) were utilized for seed germination. For callus culture establishment, 1-1.5 cm long sections of leaf explant from 28 days old *in vitro* plantlets were inoculated on MS basal medium with different concentrations of BAP, GA₃ alone or along with 1mg/L NAA (Table S1). Each concentration was replicated three times and the entire experiment was performed twice. Data of callogenic frequency was noted on weekly basis and callus was subcultured with the same PGRs concentration after every 4 weeks. Following this, callus cultures were harvested after the first subculture for the determination of fresh weight (FW) and dry weight (DW) respectively, as described previously [10].

In addition, 4 different samples of commercial leaves of Thai basil were purchased from organic markets (NaturéO, Chartres, France) for comparison purpose.

2.3. Sample Extraction

The extracts from fine powder of lyophilized dried sample were obtained using the validated green extraction method designed for the extraction of rosmarinic acid derivatives from *Lamiaceae* [21]. Shortly, the extraction consisted in S/M ratio of 25:1 ml/g DW mixed with 20 ml of pure ethanol (100%) as the extraction solvent extracted during 45 min in an ultrasound bath (USC1200TH, Prolabo, Sion, Switzerland) with operating at an ultrasound frequency of 30 kHz with extraction temperature set at 45 °C. After extraction, centrifugation was carried out for 15 min at 8,000 × g, and the resulting supernatant filtered (0.45 µm, Merck, Saint-Quentin Fallavier, France) prior further analysis.

2.4. Determination of Total Phenolic (TPC)

Determination of TPC was performed by adopting the procedure of *Singleton & Rossi* [22]. Extract (20 µl) was mixed with Folin-Ciocalteu (FC) reagent (Merck, Saint-Quentin Fallavier, France) (90 µl) and kept for 5 min before adding sodium carbonate (90 µl). Gallic acid as positive and DMSO as negative control were used. Absorbance was recorded at 630 nm with microplate reader. By using the following formula, the total phenolic production was estimated: $TPP \text{ (mg/L)} = TPC \text{ (mg/g)} \times DW \text{ (g/L)}$.

2.5. HPLC Analysis

Chicoric acid, rosmarinic acid and caffeic acid were estimated by HPLC using a method validated according to international standards of the association of analytical communities (AOAC) [10]. A reversed phase HPLC fitted with autosampler, Varian (Les Ulis, France) Prostar 230 pump and a Varian Prostar 335 photodiode array detector was utilized by following previously reported protocol [23]. A RP-18 Purospher (Merck) column (250 mm × 4.0 mm, internal diameter: 5 µm) was used at 35° C for separation. The mixture of acetic acid (0.2%), water and methanol were used as mobile phase. During 60 min run the composition of mobile phase was varied linearly from 5:95(v/v) to 100:0(v/v) at a uniform flow rate of 0.6 mL/min. Compounds were detected at 320 and 520 nm. Compounds identification was achieved by comparing with their commercial standards (Merck, Saint-Quentin Fallavier, France). Quantification was carried out using 5-point calibration curves 1 with at least 0.998 correlation coefficient [10]. Triplicate assays were conducted, and results were demonstrated as mg/g DW of the sample.

2.6. Antioxidant Activity

2.6.1. DPPH Assay

Free radical DPPH (2,2-diphenyl-1-picrylhydrazyl) scavenging potential was evaluated by following protocol of *Shah et al.* [24]. Shortly sample extract (20 µl) was blended with 180µl of DPPH solution and kept for 1h in dark. Using the microplate reader, readings were recorded at 517 nm. Free radical scavenging potential was estimated by following equation: % scavenging = $100 \times (1 - AE/AD)$. With, AE = dilution absorbance at 517 nm and AD = pure DPPH absorbance as a standard

2.6.2. ABTS Assay

Protocol of *Velioglu et al.* [25] was adapted for ABTS (2,2-Azinobis (3-ethylbenzthiazoline-6-sulphonic acid) assay. Briefly by combining equal amount of ABTS salt (7 mM) and potassium per sulphate (2.45 mM) solution was prepared. Then placed for 16 h in dark after adding callus extract. Absorbance was measured at 734 nm, after keeping the mixture for 15 min in dark at 25° C. Triplicate assays were performed and antioxidant power was demonstrated as TAEC.

2.6.3. Ferric Reducing Antioxidant Power (FRAP) Assay

To find out FRAP, protocol of *Benzie and Strain* [26] was adapted. Sample extract (10 µl) was blended with of FRAP solution (190 µl). Absorbance was recorded at 630 nm after incubating the mixture at 25° C for 15 min. Triplicate assays were performed and antioxidant power was demonstrated as TAEC.

2.6.4. Cellular Antioxidant Assay

Cellular antioxidant assay was performed with the detection of reactive nitrogen (RNS) and oxygen (ROS) species levels were determined by using a fluorescent dye Dihydrorhodamine-123 (DHR-123) as described previously [27]. Overnight incubation of yeast cells with Thai basil extracts was done. Then washed two times with PBS and kept at 30° C in the dark for 10 min. BioRad (Marnes-la-Coquette, France) Fluorimeter (λ_{em} = 535 nm, λ_{ex} = 505 nm) was used for detection of fluorescent signal after repeatedly washing with PBS.

2.7. Statistical Analysis

Means and standard deviations of three to three separate replicates were used to present the data. Statistical analysis was performed with XLSTAT 2019 (Addinsoft, Paris, France) following the manufacturer's instructions using Kruskal-Wallis test. Significant threshold at $p < 0.05$ was used for all statistical tests and was represented by different letters in the Tables and Figures. Correlation analysis was performed with Past 3.0 (Øyvind Hammer, Natural History Museum, University of Oslo, Oslo, Norway) using the Pearson parametric correlation test and visualized using Heatmapper [28].

3. Results and Discussion

3.1. Callus Induction and Biomass Production

Callogenesis from 28 days old leaf explants was evaluated for optimum biomass production in response to various PGRs (BAP, GA₃, NAA) treatments either alone or in combination (Table S1). The morphological aspects of the resulting Thai basil callus on different culture media are shown in Figure 1B. Morphological differences in the color and texture of callus were noted. Here, callogenesis originated at the cut end of explants after 7 to 12 days in response to all tested PGRs. However, the callogenic frequency was varied in response to all PGRs tested. A high callogenic frequency of 95% was observed for BAP:NAA (5 mg/L: 1 mg/L) followed by 92 % for BAP:NAA (2 mg/L: 1 mg/L). Callogenesis resulted around 20 % to 70 % for BAP treatments but the value improved to 95 % when NAA was used in combination with BAP (Table S1). No callus induction was observed in the MS basal medium without hormones, whereas the initiation of callogenesis was observed with NAA alone, but the callus rapidly turn brown, without noticeable growth and eventually die.

Maximum FW (190 g/L) and DW (13.05 g/L) biomasses were produced in presence of BAP:NAA (5 mg/L: 1 mg/L) (Figure 1B). While lowest FW (31.5 g/L) and DW (3.7 g/L) biomasses were noted for GA₃ (10.0 mg/L) (Figure 1B). Moreover, the maximal production of FW and DW biomasses were detected for all BAP concentrations (0.25-5.0 mg/L) combined with NAA (Figure 1B).

Earlier reports have shown that exogenous auxins in major angiosperm species are essential growth regulators for callus induction [29]. Cytokinin along with auxin is generally required for maximum callus induction by most plant species. In particular, the need for exogenous cytokinin to maintain the interdependent balance between cytokinin and auxin that promotes cell division has been reported as a significant strategy for the production of callus [30,31]. Likewise, in numerous other medicinal plant species, the combination of NAA with other PGRs resulted in maximum callogenic frequency [19,32]. However, higher concentrations of applied PGRs may limit the callogenesis, possibly due to the repression of some endogenous hormones [33]. The combination of BAP and NAA have been shown to produce optimum biomass for callus cultures of many medicinal plants such as the endangered species *Aquilaria malaccensis* Lam [34]. This may result from the synergistic interaction between BAP and NAA as observed for the production of *Brassica oleracea* L. callus culture biomass [35]. In our previous studies, on *O. basilicum* L. cv Purple basil *in vitro* cultures optimal production of biomass was achieved using the PGR thidiazuron (TDZ) [10]. However, TDZ

is no longer considered safe in Europe where its use is banned in particular for nutraceutical applications, which is the intended application.

Figure 1. (A) Effects of different PGRs on callogenesis from Thai basil leaves explants. (a1) 1 mg/L BAP, (a2) 5 mg/L BAP, (a3) BAP:NAA (0.5 mg/L: 1 mg/L), (a4) BAP:NAA (5 mg/L: 1 mg/L), (a5) 2 mg/L GA₃, (a6) 5 mg/L GA₃, (a7) GA₃:NAA (0.25 mg/L:1 mg/L), (a8) GA₃:NAA (5 mg/L: 1 mg/L); (B) Biomass accumulation in response to different PGRs. Fresh weight (FW) and dry weight (DW) from leaf derived callus grown on BAP (0.25, 0.5, 1, 2, 5, 10 mg/L) and GA₃ (0.25, 0.5, 1, 2, 5, 10 mg/L) alone or in combination with NAA (1 mg/L). Values are means $\pm$ standard deviations (SD) of at least 3 independent experiments. Different letters represent significant differences between the various extraction conditions ($p < 0.05$). Note that no differences were observed for the statistical groups for the FW and DW values.

3.2. Production of phenolic compounds

3.2.1. Total Phenolic Content (TPC)

In response to BAP combined with NAA (5 mg/L: 1 mg/L) callus yield maximum total phenolic content (TPC) of 132.6 mg/g DW (Table 2). On the contrary, BAP alone, whatever the concentration applied, resulted in a lower TPC ranging from 42.5 to 70.8 mg/g DW. We noted that for each BAP concentration used, the addition of NAA increased TPC. Minimum TPC values ranging from 21.8 to 37.0 mg/g DW have been detected for GA₃ treatments (0.25-10.0 mg/L). In addition to GA₃, NAA also stimulated the accumulation of total phenolics.

Interestingly, higher TPC was obtained in extract from *in vitro* callus cultures grown under optimal PGRs conditions as compared to the extract from commercial leaves of Thai basil (Table 1).

Table 1. Quantification of the main phenolic compounds accumulated in extracts from Thai basil callus cultures in response to different PGRs treatment and leaves.

PGR Treatments	TPC ¹ (mg/g DW)	Chicoric acid (mg/gDW)	Rosmarinic acid (mg/g DW)	Caffeic acid (mg/g DW)
0.25GA3	27.0±0.9 ^k	20.8±0.9 ^f	2.2±0.1 ^f	0.06±0.03 ^b
0.5GA3	31.8±1.5 ^k	23.8±0.7 ^e	3.1±0.2 ^e	0.06±0.02 ^b
1GA3	31.8±3.4 ^k	28.1±0.7 ^{cd}	3.0±0.2 ^e	0.06±0.02 ^b
2GA3	37.0±2.8 ^j	28.5±0.6 ^c	3.7±0.2 ^d	0.07±0.02 ^b
5GA3	32.7±3.2 ^k	27.0±0.5 ^d	3.9±0.2 ^d	0.06±0.03 ^b
10GA3	21.8±1.8 ^l	21.5±0.4 ^f	3.1±0.2 ^e	0.05±0.03 ^b
0.25GA3+NAA	52.7±3.6 ⁱ	22.7±0.4 ^{ef}	4.3±0.3 ^d	0.06±0.02 ^b
0.5GA3+NAA	71.7±4.3 ^{fg}	23.7±0.4 ^e	4.5±0.3 ^{cd}	0.06±0.02 ^b
1GA3+NAA	76.3±0.6 ^f	26.3±0.5 ^d	5.1±0.3 ^c	0.07±0.03 ^b
2GA3+NAA	79.9±2.1 ^e	21.6±0.4 ^f	5.4±0.3 ^{bc}	0.06±0.02 ^b
5GA3+NAA	85.4±4.7 ^{de}	16.7±0.3 ^g	5.5±0.3 ^{bc}	0.07±0.03 ^b
10GA3+NAA	89.9±1.3 ^d	15.4±0.3 ^h	4.9±0.3 ^c	0.08±0.03 ^b
0.25BAP	42.5±0.8 ^j	13.6±0.3 ⁱ	2.1±0.2 ^f	0.04±0.01 ^b
0.5BAP	61.5±2.7 ^h	15.3±0.3 ^h	2.3±0.2 ^f	0.04±0.02 ^b
1BAP	66.3±5.3 ^{gh}	17.6±0.3 ^g	2.8±0.2 ^{ef}	0.04±0.02 ^b
2BAP	70.8±3.9 ^g	14.6±0.3 ^h	2.8±0.2 ^{ef}	0.05±0.02 ^b
5BAP	45.4±4.6 ^{ij}	13.0±0.2 ⁱ	2.5±0.2 ^f	0.04±0.02 ^b
10BAP	43.5±3.1 ^j	14.8±0.2 ^h	2.4±0.2 ^f	0.04±0.02 ^b
0.25BAP+NAA	41.8±4.2 ^j	20.5±0.3 ^f	4.2±0.3 ^d	0.06±0.03 ^b
0.5BAP+NAA	63.6±3.6 ^{gh}	26.4±0.5 ^d	4.7±0.3 ^{cd}	0.07±0.03 ^b
1BAP+NAA	92.6±2.7 ^{cd}	31.3±0.8 ^b	5.9±0.3 ^b	0.07±0.02 ^b
2BAP+NAA	105.4±4.4 ^b	36.8±1.1 ^a	7.1±0.4 ^a	0.07±0.02 ^b
5BAP+NAA	132.6±3.1 ^a	35.8±1.2 ^a	7.4±0.7 ^a	0.07±0.02 ^b
10BAP+NAA	96.3±2.3 ^c	32.5±0.9 ^b	7.0±0.4 ^a	0.06±0.02 ^b
LEAVES	87.1±4.1 ^e	24.2±1.8 ^e	7.2±0.7 ^a	0.6±0.1 ^a

¹ Total phenolics content; Values are means ± SD of three independent replicates. Different letters represent significant differences between the various extraction conditions (p < 0.05).

Interestingly, there was a significant correlation between biomass production and total phenolic production (Figure 2, Table S2). In line with this observation, the biomass-dependent production of total phenolics has already been reported in different plant systems [36].

Figure 2. Pearson correlation analysis (PCC) of the relation between the main phytochemicals (TPC: total phenolics content; chicoric acid, rosmarinic acid and caffeic acid) in extracts from Thai basil callus cultures and the different antioxidant assays (*in vitro* cell-free: DPPH, ABTS, FRAP and *in vivo*: CAA (cellular antioxidant assay)). *** significant $p < 0.001$; ** significant $p < 0.01$; * significant $p < 0.05$; actual PCC values are indicated in Supplementary Materials Table S1.

Szopa & Ekiert [37] noted that PGRs have a direct impact on the biosynthesis of phenolics in medicinal plants. This result is supported by a number of studies which report that the inclusion of a higher BAP concentration and a lower NAA concentration resulted in the maximum TPC [38–42]. This observation could be related to stimulation of essential enzymes for the biosynthesis of various phenylpropanoids, such as phenylalanine/tyrosine ammonia-lyase (PAL/TAL), in response to several PGR concentrations [40]. In contrast, in accordance with our results, GA₃ has previously been shown to suppress the production of some phenylpropanoids, such as lignans in flax, in a dose-dependent manner [43].

3.2.2. HPLC-Based Evaluation of Simple Phenolics

The HPLC analysis confirmed treatment with different PGRs of leaf-derived callus culture as a tool for improving the synthesis of the main biologically active phenolic acids of Thai basil (Table 1). All BAP and GA₃ treatments alone and with NAA exhibit a wide range of metabolite production. Maximum accumulation of chicoric acid (36.8 and 35.8 mg/g DW) obtained for the BAP treatments at concentrations of 2 and 5 mg/L, respectively, in combination with NAA at 1 mg/L (Table 1). By contrast, treatments with BAP alone at varying concentrations resulted in the lowest accumulation of chicoric acid ranging from 13.0 mg/g DW to 17.6 mg/g DW (Table 1). Similarly, an optimum production of rosmarinic acid (7.4 mg/g DW) was observed for combined BAP and NAA (5 mg/L: 1 mg/L) treatment, while a minimum production was found for all tested BAP concentrations alone (Table 1). Compared to extract from commercial Thai basil leaves, chicoric acid accumulation was 2.6-fold higher in extract from *in vitro* callus produced under optimized conditions, while rosmarinic acid content was similar in both types of extracts (Table 1). In comparison, the accumulation levels in callus extracts of their biosynthetic precursor, caffeic acid, was much lower than that of chicoric acid and rosmarinic acid (Table 1). The concentration of caffeic acid in extract from *in vitro* callus cultures was 7.5-to 10-fold lower than in extract from commercial Thai basil leaves (Table 1).

Chicoric, rosmarinic and caffeic acids have previously been reported in different basil species and varieties [44], in particular in Thai basil [6]. It is generally accepted that the accumulation of different phenolic acids in plant cultures may be modulated by the exogenous

application of PGRs [37,43,45,46]. In line with this observation, our results shown that phenolic acids, chicoric acid and rosmarinic acid can be modulated by PGRs *in vitro* callus cultures of Thai basil. Similarly, [47] also revealed an increase in the concentration of chicoric acid in their callus cultures of *Echinacea purpurea* grown on NAA supplemented medium. Similarly, increased production of rosmarinic acid was observed in *O. basilicum* *in vitro* cultures grown on media supplemented with NAA in combination with other PGRs [48], and in *Ocimum americanum* callus culture grown on BAP in combination with IAA [49]. In the present study, chicoric acid and rosmarinic acid were both quantified and their contents were increased in response to PGR treatment. The lower levels of caffeic acid may be associated with its efficient biotransformation into its derivatives rosmarinic acid and chicoric acid [50,51].

3.3. Antioxidant Activities

3.3.1. In Vitro Cell-Free Antioxidant Assays

Three distinct *in vitro* cell-free antioxidant assays, namely DPPH, ABTS and FRAP were used to assess the antioxidant capacity of the extracts from Thai basil callus cultures. FRAP represents electron transfer (ET) antioxidant mechanism, while ABTS follows hydrogen atom transfer (HAT) antioxidant mechanism. Here, the results of both assays were calculated as TEAC (μM of Trolox C equivalent antioxidant capacity). DPPH assay can reveal both HAT-and ET-based antioxidants, and was calculated as a percentage of DPPH radical scavenging activity (RSA percentage). Extracts from callus grown on BAP:NAA (5 mg/L: 1 mg/L) presented the highest levels for DPPH (93.2 % RSA) and ABTS (1322.0 μM TEAC) antioxidant assays (Table 2), while lowest levels for DPPH (61.0 % RSA) and ABTS (713.1 μM TEAC) were measured for extract from callus grown on 10 mg/L GA₃ medium (Table 2). Maximum FRAP (523.3 μM TEAC) was measured for extract from callus grown on 2 mg/L BAP medium, and lowest value (335.3 μM TEAC) for extract resulting from callus grown on 10 mg/L GA₃ medium (Table 2).

Comparatively, extract from commercial Thai basil leaves presented a 71.4 % RSA for DPPH assay, as well as 914.2 and 387.7 μM TEAC for ABTS and FRAP assays, respectively. Based on these *in vitro* cell-free assays, the measured antioxidant capacity of the extract from commercial Thai basil leaves was within the range of those determined for callus extracts (Table 2). However, under optimized conditions, extracts from callus cultures showed significantly higher *in vitro* antioxidant capacity than extract from commercial leaves (Table 2).

Significant correlations between *in vitro* antioxidant activity and the presence of chicoric acid, rosmarinic acid and caffeic acid were observed for each assay (Figure 2, Table S2). TPC was also significantly correlated with ABTS antioxidant activity (Figure 2, Table S2).

Similarly, a significant relationship between phenolic acid accumulation and antioxidant capacity has been revealed for several medicinal plant species [52,53]. From a strictly chemical point of view, these *in vitro* cell-free antioxidant assays are interesting since they can provide information on the mechanism reaction involved in the antioxidant potential of an extract [54]. Here, the results obtained could reflect a preferential ET-type mechanism. However, these *in vitro* assays do not necessarily reflect the situation of the *in vivo* systems and, therefore, their validity should be considered to be strictly limited to interpretation within the meaning of the chemical reactivity with respect to the *in vitro* produced radicals, and thus should be confirmed by *in vivo* assays.

Table 2: *In vitro* cell-free antioxidant potential of extracts from Thai basil callus cultures in response to different PGRs treatment and leaves.

PGR Treatments	DPPH (% RSA)	ABTS (TEAC μM)	FRAP (TEAC μM)
0.25GA3	85.9±1.7 c	891.7±5.4 g	438.2±5.9 de
0.5GA3	74.2±1.9 e	907.5±5.9 g	446.5±5.5 d
1GA3	95.1±1.8 a	826.7±6.0 h	432.6±5.4 d
2GA3	86.9±1.8 b	759.4±4.9 i	420.7±5.5 e
5GA3	91.6±1.7 ab	814.3±4.9 h	385.5±5.3 f
10GA3	61.0±1.8 g	713.1±5.2 k	335.3±5.8 h
0.25GA3+NAA	69.8±1.7 ef	1034.1±7.9 d	403.0±5.1 f
0.5GA3+NAA	72.0±1.8 e	1100.2±7.6 c	445.7±5.3 d
1GA3+NAA+NAA	64.4±1.6 f	1170.4±6.7 b	450.4±4.9 d
2GA3+NAA	88.1±1.8 b	1030.5±6.9 d	499.7±5.2 b
5GA3+NAA	89.6±1.7 b	933.3±6.9 f	451.4±5.3 d
10GA3+NAA	79.2±1.5 d	852.8±7.6 h	349.3±4.1 h
0.25BAP	63.6±1.6 f	788.7±7.7 i	483.2±4.9 c
0.5BAP	67.4±1.6 f	784.2±7.5 i	505.5±5.2 ab
1BAP	73.5±1.5 e	1101.1±7.7 c	496.4±5.4 bc
2BAP	85.1±2.1 c	878.5±6.8 g	523.3±6.1 a
5BAP	77.8±2.2 d	897.4±7.1 g	517.3±5.7 a
10BAP	72.5±1.9 e	862.0±6.8 h	484.4±4.9 c
0.25BAP+NAA	74.7±2.0 e	849.7±6.4 h	502.9±6.1 b
0.5BAP+NAA	82.6±2.1 cd	991.2±6.9 e	454.6±4.8 d
1BAP+NAA	79.1±1.9 d	997.3±6.9 e	447.6±4.4 d
2BAP+NAA	82.6±2.1 cd	1194.3±7.9 b	415.2±4.3 e
5BAP+NAA	93.2±2.2 a	1322.0±7.7 a	384.7±4.1 f
10BAP+NAA	87.2±2.1 b	1080.1±6.9 c	376.7±4.3 g
LEAVES	71.4±3.2 e	914.2±7.3 f	387.7±8.2 f

Values are means ± SD of three independent replicates. Different letters represent significant differences between the various extraction conditions (p < 0.05).

3.3.2. Cellular Antioxidant Activity

Antioxidant activity was also evaluated *in vivo* through the ability of the extract to inhibit the formation of reactive nitrogen and oxygen species (RNS and ROS) using a cellular antioxidant assay (Figure 3). The results confirmed the protective action of Thai basil callus extracts against oxidative stress. Maximum inhibition of the ROS/RNS production (56.4 %) was measured with extract from callus grown on medium with BAP and NAA (5 mg/L: 1 mg/L), while minimum production inhibition ranging from 17.5 % to 27.4 % were recorded for extracts from callus grown on media supplemented with BAP alone (Figure 3).

Figure 3: *In vivo* antioxidant potential of extracts from Thai basil callus cultures in response to different PGRs treatment and leaves measured by cellular antioxidant assay (CAA) estimated as inhibition % of ROS/RNS production. Values are means $\pm$ SD of three independent replicates. Different letters represent significant differences between the various extraction conditions ($p < 0.05$).

Extract from commercial Thai basil leaves presented an inhibition of 35.6 % of the ROS/RNS production. As a result, extracts from callus cultures grown under optimized conditions, also showed significantly higher *in vivo* antioxidant capacity than extracts from commercial leaves (Table 2), thus demonstrating the interest of this culture system in potential nutraceutical applications.

The cellular antioxidant assay confirmed the interest of natural antioxidant from plant extracts [17,55,56]. In the past decade, natural antioxidants have generated increasing interest due to their possible use as an alternative to potentially hazardous synthetic antioxidants such as butylated hydroxyanisole (BHA) or butylated hydroxytoluene (BHT) in various food preparations and their possible use as nutraceuticals [57]. Some natural antioxidants have already been shown to be as effective as these synthetic antioxidants [58,59]. In the meantime, plant *in vitro* systems have been proposed as an efficient alternative for the production of natural antioxidants [60]. The present results indicate the potential use of an *in vitro* biological system as a tool for continuous and high production of natural antioxidants from Thai basil.

4. Conclusions

A protocol was proposed for an effective production of Thai basil leaf-derived callus cultures with sustainable and high production of biomass and antioxidants. Here, we showed that plant cell and tissue culture technologies can be routinely used for Thai basil as an important, sustainable and low-cost biomass source for the production of high-value antioxidant compounds, particularly when the optimal combination of PGRs is used. Indeed, under optimal conditions, compared to commercial leaves, the extracts of callus showed higher production of chicoric acid, and similar levels of rosmarinic acid, associated with higher antioxidant capacity. Furthermore, given that the large capacity of this biological system for continuous biomass production can be further expanded in the future with the use of cell suspension cultures in bioreactors, we anticipate that it could have a great potential for possible nutraceutical applications.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, **Table S1:** Callogenesis, morphology and induction frequency of *O. basilicum* callus culture under various PGRs. **Table S2:** Actual values for PCC (Pearson Correlation Coefficient) showing the relation between the different

phytochemicals (TPC: total phenolics content; chicoric acid, rosmarinic acid and caffeic acid) in extracts from Thai basil callus cultures and the different antioxidant assays (*in vitro* cell-free: DPPH, ABTS, FRAP and *in vivo*: CAA (cellular antioxidant assay)).

Author Contributions: Conceptualization, B.H.A. and C.H.; methodology, S.N., D.T., S.D. and C.H.; software, S.D. and C.H.; validation, B.H.A., D.T. and C.H.; formal analysis, B.H.A., D.T. and C.H.; investigation, S.N., H.J., D.T. and S.D.; resources, B.H.A. and C.H.; data curation, B.H.A., D.T. and C.H.; writing—original draft preparation, S.N., H.J. and C.H.; writing—review and editing, B.H.A., D.T., S.D., M.Z. and C.H.; visualization, S.N., H.J., D.T., M.Z., C.H. and B.H.A.; supervision, B.H.A. and C.H.; project administration, B.H.A. and C.H.; funding acquisition, B.H.A. and C.H.; All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by Cosmetosciences, a global training and research program dedicated to the cosmetic industry. Located in the heart of the Cosmetic Valley, this program led by University of Orléans is funded by the Région Centre-Val de Loire (VALBIOCOSM 17019UNI).

Acknowledgments: B.H.A. acknowledges research fellowship of Le Studium-Institute for Advanced Studies, Loire Valley, Orléans, France. S.D. acknowledges research fellowship of Région Centre-Val de Loire, France. D.T. and C.H. gratefully acknowledges the support of the French Government via the French Embassy in Thailand in the form of Junior Research Fellowship Program 2018. C.H. and D.T. also gratefully acknowledges the support of Campus France through the PHC SIAM (PNPIA, Project 44926WK). S.D. acknowledges post-doctoral research fellowship of Loire Valley Region.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Sahoo, Y.; Pattnaik, S. K.; Chand, P. K. *In vitro* clonal propagation of an aromatic medicinal herb *Ocimum basilicum* L. (sweet basil) by axillary shoot proliferation. *In Vitro Cell. Dev. Biol.* **1997**, *33*, 293–296.
2. Paton, A.; Harley, R. M.; Harley, M. M. *Ocimum*: an overview of classification and relationships. In *Basil*; CRC Press, 1999; pp. 11–46.
3. Vani, S. R.; Cheng, S. F.; Chuah, C. H. Comparative study of volatile compounds from genus *Ocimum*. *Am. J. Appl. Sci.* **2009**, *6*, 523.
4. Paton, A.; Putievsky, E. Taxonomic problems and cytotaxonomic relationships between and within varieties of *Ocimum basilicum* and related species (Labiatae). *Kew Bull.* **1996**, 509–524.
5. Selvi, M. T.; Thirugnanasampandan, R.; Sundarammal, S. Antioxidant and cytotoxic activities of essential oil of *Ocimum canum* Sims. from India. *J. Saudi Chem. Soc.* **2012**, *9*, 97–100.
6. Lee, J.; Scagel, C. F. Chicoric acid found in basil (*Ocimum basilicum* L.) leaves. *Food Chem.* **2009**, *115*, 650–656.
7. Nyarko, A. K.; Asare-Anane, H.; Ofosuhen, M.; Addy, M. E. Extract of *Ocimum canum* lowers blood glucose and facilitates insulin release by isolated pancreatic β -islet cells. *Phytomedicine* **2002**, *9*, 346–351.
8. Kaya, I.; Yigit, N.; Benli, M. Antimicrobial activity of various extracts of *Ocimum basilicum* L. and observation of the inhibition effect on bacterial cells by use of scanning electron microscopy. *African J. Tradit. Complement. Altern. Med.* **2008**, *5*, 363–369.
9. Behera, S.; Babu, S. M.; Ramani, Y. R.; Choudhury, P. K.; Panigrahi, R. Phytochemical investigation and study on antioxidant properties of *Ocimum canum* hydro-alcoholic leaf extracts. *J. Drug Deliv. Ther.* **2012**, *2*.
10. Nazir, M.; Tungmunthum, D.; Bose, S.; Drouet, S.; Garros, L.; Giglioli-Guivarc'h, N.; Abbasi, B. H.;

- 381 Hano, C. Differential Production of Phenylpropanoid Metabolites in Callus Cultures of *Ocimum*
 382 *basilicum* L. With Distinct in Vitro Antioxidant Activities and in Vivo Protective Effects against UV stress.
 383 *J. Agric. Food Chem.* **2019**, *67*, 1847–1859, doi:10.1021/acs.jafc.8b05647.
- 384 11. Sasheva, P.; Letkarska, G.; Ionkova, I. Biotechnological production of podophyllotoxin and
 385 podophyllotoxin-related lignans in cultures of *Linum thracicum* Degen. *Comptes Rendus l'Academie Bulg.*
 386 *des Sci.* **2013**, *66*, 1445–1450.
- 387 12. Khurshid, R.; Khan, T.; Zaeem, A.; Garros, L.; Hano, C.; Abbasi, B. H. Biosynthesis of precious
 388 metabolites in callus cultures of *Eclipta alba*. *Plant Cell, Tissue Organ Cult.* **2018**, *135*, 287–298.
- 389 13. Drouet, S.; Garros, L.; Hano, C.; Lainé, É. A Critical View of Different Botanical , Molecular , and
 390 Chemical Techniques Used in Authentication of Plant Materials for Cosmetic Applications. *Cosmetics*.
 391 **2018**, doi:10.3390/cosmetics5020030.
- 392 14. Smith, M. A. L. Valuable secondary products from in vitro culture. Chapter 24. *Plant Dev. Biotechnol.*
 393 *CRC Press LLC, Boca Rat.* **2005**.
- 394 15. Bais, H. P.; Walker, T. S.; Schweizer, H. P.; Vivanco, J. M. Root specific elicitation and antimicrobial
 395 activity of rosmarinic acid in hairy root cultures of *Ocimum basilicum*. *Plant Physiol. Biochem.* **2002**, *40*,
 396 983–995.
- 397 16. Gopi, C.; Ponmurugan, P. Somatic embryogenesis and plant regeneration from leaf callus of *Ocimum*
 398 *basilicum* L. *J. Biotechnol.* **2006**, *126*, 260–264.
- 399 17. Nazir, M.; Asad Ullah, M.; Mumtaz, S.; Siddiquah, A.; Shah, M.; Drouet, S.; Hano, C.; Abbasi, B. H.
 400 Interactive Effect of Melatonin and UV-C on Phenylpropanoid Metabolite Production and Antioxidant
 401 Potential in Callus Cultures of Purple Basil (*Ocimum basilicum* L. var. *s. purpurascens*). *Molecules* **2020**, *25*,
 402 1072.
- 403 18. Nazir, M.; Ullah, M. A.; Younas, M.; Siddiquah, A.; Shah, M.; Giglioli-Guivarc'h, N.; Hano, C.; Abbasi,
 404 B. H. Light-mediated biosynthesis of phenylpropanoid metabolites and antioxidant potential in callus
 405 cultures of purple basil (*Ocimum basilicum* L. var *purpurascens*). *Plant Cell, Tissue Organ Cult.* **2020**.
- 406 19. Abbasi, B. H.; Khan, M. A.; Mahmood, T.; Ahmad, M.; Chaudhary, M. F.; Khan, M. A. Shoot regeneration
 407 and free-radical scavenging activity in *Silybum marianum* L. *Plant Cell, Tissue Organ Cult.* **2010**, *101*, 371–
 408 376.
- 409 20. Murashige, T.; Skoog, F. A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue
 410 Cultures. *Physiol. Plant.* **1962**, *15*, 473–497, doi:10.1111/j.1399-3054.1962.tb08052.x.
- 411 21. Tungmunthum, D.; Garros, L.; Drouet, S.; Renouard, S.; Lainé, E.; Hano, C. Green Ultrasound Assisted
 412 Extraction of trans Rosmarinic Acid from *Plectranthus scutellarioides* (L.) R.Br. Leaves. *Plants* **2019**, *8*, 50,
 413 doi:10.3390/plants8030050.
- 414 22. Singleton, V. L.; Rossi, J. A. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid
 415 reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144–158.
- 416 23. Coricovac, D. O.; Soica, C. O.; Muntean, D. A.; Popovici, R. A.; Dehelean, C. A.; Hoge, E. L. Assessment
 417 of the effects induced by two triterpenoids on liver mitochondria respiratory function isolated from aged
 418 rats. *Rev. Chim* **2015**, *66*, 1707–1710.
- 419 24. Shah, M.; Ullah, M. A.; Drouet, S.; Younas, M.; Tungmunthum, D.; Giglioli-Guivarc'h, N.; Hano, C.;
 420 Abbasi, B. H. Interactive effects of light and melatonin on biosynthesis of silymarin and anti-
 421 inflammatory potential in callus cultures of *Silybum marianum* (L.) Gaertn. *Molecules* **2019**,
 422 doi:10.3390/molecules24071207.

- 423 25. Velioglu, Y. S.; Mazza, G.; Gao, L.; Oomah, B. D. Antioxidant Activity and Total Phenolics in Selected
424 Fruits, Vegetables, and Grain Products. *J. Agric. Food Chem.* **1998**, *46*, 4113–4117, doi:10.1021/jf9801973.
- 425 26. Benzie, I.; Strain, J. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”:
426 the FRAP assay. *Anal. Biochem.* **1996**, *239*, 70–76, doi:10.1006/abio.1996.0292.
- 427 27. Tungmunthum, D.; Abid, M.; Elamrani, A.; Drouet, S.; Addi, M.; Hano, C. Almond Skin Extracts and
428 Chlorogenic Acid Delay Chronological Aging and Enhanced Oxidative Stress Response in Yeast. *Life*
429 **2020**, *10*, 80.
- 430 28. Babicki, S.; Arndt, D.; Marcu, A.; Liang, Y.; Grant, J. R.; Maciejewski, A.; Wishart, D. S. Heatmapper:
431 web-enabled heat mapping for all. *Nucleic Acids Res.* **2016**, *44*, W147–W153, doi:10.1093/nar/gkw419.
- 432 29. Chang, C.; Chang, W. C. Plant regeneration from callus culture of *Cymbidium ensifolium* var. *misericors*.
433 *Plant Cell Rep.* **1998**, *17*, 251–255.
- 434 30. Ishii, Y.; Takamura, T.; Goi, M.; Tanaka, M. Callus induction and somatic embryogenesis of *Phalaenopsis*.
435 *Plant Cell Rep.* **1998**, *17*, 446–450.
- 436 31. Johri, M. M.; Mitra, D. Action of plant hormones. *Curr. Sci.* **2001**, *80*, 199–205.
- 437 32. Liu, S.; Gao, P.; Zhu, Q.; Luan, F.; Davis, A. R.; Wang, X. Development of cleaved amplified polymorphic
438 sequence markers and a CAPS-based genetic linkage map in watermelon (*Citrullus lanatus* [Thunb.]
439 Matsum. and Nakai) constructed using whole-genome re-sequencing data. *Breed. Sci.* **2016**, *66*, 244–259,
440 doi:10.1270/jsbbs.66.244.
- 441 33. Anjum, S.; Abbasi, B. H.; Hano, C. Trends in accumulation of pharmacologically important antioxidant-
442 secondary metabolites in callus cultures of *Linum usitatissimum* L. *Plant Cell. Tissue Organ Cult.* **2017**, *129*,
443 doi:10.1007/s11240-016-1158-3.
- 444 34. Jayaraman, S.; Daud, N. H.; Halis, R.; Mohamed, R. Effects of plant growth regulators, carbon sources
445 and pH values on callus induction in *Aquilaria malaccensis* leaf explants and characteristics of the
446 resultant calli. *J. For. Res.* **2014**, *25*, 535–540.
- 447 35. Ahmad, S.; Spoor, W. Effect of NAA and BAP on callus culture and plant regeneration in Curly kale
448 (*Brassica oleracea* L.). *Pakistan J. Biol. Sci.* **1999**.
- 449 36. Ali, M.; Abbasi, B. H. Light-induced fluctuations in biomass accumulation, secondary metabolites
450 production and antioxidant activity in cell suspension cultures of *Artemisia absinthium* L. *J. Photochem.*
451 *Photobiol. B Biol.* **2014**, *140*, 223–227.
- 452 37. Szopa, A.; Ekiert, H. Production of biologically active phenolic acids in *Aronia melanocarpa* (Michx.)
453 Elliott in vitro cultures cultivated on different variants of the Murashige and Skoog medium. *Plant*
454 *Growth Regul.* **2014**, *72*, 51–58.
- 455 38. Taveira, M.; Pereira, D. M.; Sousa, C.; Ferreres, F.; Andrade, P. B.; Martins, A.; Pereira, J. A.; Valentao, P.
456 In vitro cultures of *Brassica oleracea* L. var. *costata* DC: potential plant bioreactor for antioxidant phenolic
457 compounds. *J. Agric. Food Chem.* **2009**, *57*, 1247–1252.
- 458 39. Kartnig, T.; Kögl, G.; Heydel, B. Production of flavonoids in cell cultures of *Crataegus monogyna*. *Planta*
459 *Med.* **1993**, *59*, 537–538.
- 460 40. Maharik, N.; Elgengaihi, S.; Taha, H. Anthocyanin production in callus cultures of *Crataegus sinaica* boiss.
461 *Int J Acad Res* **2009**, *1*, 30–34.
- 462 41. Chaâbani, G.; Tabart, J.; Kevers, C.; Dommès, J.; Khan, M. I.; Zaoui, S.; Chebchoub, L.; Lachaâl, M.;

- 463 Karray-Bouraoui, N. Effects of 2, 4-dichlorophenoxyacetic acid combined to 6-Benzylaminopurine on
464 callus induction, total phenolic and ascorbic acid production, and antioxidant activities in leaf tissue
465 cultures of *Crataegus azarolus* L. var. *aronia*. *Acta Physiol. Plant.* **2015**, 37, 16.
- 466 42. El-Baz, F. K.; Mohamed, A. A.; Ali, S. I. Callus formation, phenolics content and related antioxidant
467 activities in tissue culture of a medicinal plant colocynth (*Citrullus colocynthis*). *Nov. Biotechnol.* **2010**, 10,
468 79–94.
- 469 43. Corbin, C.; Renouard, S.; Lopez, T.; Lamblin, F.; Lainé, E.; Hano, C. Identification and characterization
470 of cis-acting elements involved in the regulation of ABA- and/or GA-mediated LuPLR1 gene expression
471 and lignan biosynthesis in flax (*Linum usitatissimum* L.) cell cultures. *J. Plant Physiol.* **2013**, 170, 516–522,
472 doi:10.1016/j.jplph.2012.11.003.
- 473 44. Srivastava, S.; Adholeya, A.; Conlan, X. A.; Cahill, D. M. Acidic potassium permanganate
474 chemiluminescence for the determination of antioxidant potential in three cultivars of *Ocimum basilicum*.
475 *Plant foods Hum. Nutr.* **2016**, 71, 72–80.
- 476 45. Coste, A.; Vlase, L.; Halmagyi, A.; Deliu, C.; Coldea, G. Effects of plant growth regulators and elicitors
477 on production of secondary metabolites in shoot cultures of *Hypericum hirsutum* and *Hypericum*
478 *maculatum*. *Plant Cell, Tissue Organ Cult.* **2011**, 106, 279–288.
- 479 46. Ali, M.; Abbasi, B. H. Production of commercially important secondary metabolites and antioxidant
480 activity in cell suspension cultures of *Artemisia absinthium* L. *Ind. Crops Prod.* **2013**, 49, 400–406.
- 481 47. Ramezannezhad, R.; Aghdasi, M.; Fatemi, M. Enhanced production of cichoric acid in cell suspension
482 culture of *Echinacea purpurea* by silver nanoparticle elicitation. *Plant Cell, Tissue Organ Cult.* **2019**, 139,
483 261–273.
- 484 48. Kintzios, S.; Makri, O.; Panagiotopoulos, E.; Scapeti, M. In vitro rosmarinic acid accumulation in sweet
485 basil (*Ocimum basilicum* L.). *Biotechnol. Lett.* **2003**, 25, 405–408.
- 486 49. Rady, M. R.; Nazif, N. M. Rosmarinic acid content and RAPD analysis of in vitro regenerated basil
487 (*Ocimum americanum*) plants. *Fitoterapia* **2005**, 76, 525–533.
- 488 50. Petersen, M.; Simmonds, M. S. J. Rosmarinic acid. *Phytochemistry* **2003**, 62, 121–125.
- 489 51. Petersen, M.; Abdullah, Y.; Benner, J.; Eberle, D.; Gehlen, K.; Hücherig, S.; Janiak, V.; Kim, K. H.; Sander,
490 M.; Weitzel, C. Evolution of rosmarinic acid biosynthesis. *Phytochemistry* **2009**, 70, 1663–1679.
- 491 52. Kim, D.-O.; Chun, O. K.; Kim, Y. J.; Moon, H.-Y.; Lee, C. Y. Quantification of polyphenolics and their
492 antioxidant capacity in fresh plums. *J. Agric. Food Chem.* **2003**, 51, 6509–6515.
- 493 53. Djeridane, A.; Yousfi, M.; Nadjemi, B.; Boutassouna, D.; Stocker, P.; Vidal, N. Antioxidant activity of
494 some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem.* **2006**, 97, 654–660.
- 495 54. Hano, C.; Tungmunthum, D. Plant Polyphenols, More than Just Simple Natural Antioxidants:
496 Oxidative Stress, Aging and Age-Related Diseases. *Medecines*. **2020**, 7, (5):26
- 497 55. Garros, L.; Drouet, S.; Corbin, C.; Decourtil, C.; Fidel, T.; De Lacour, J. L.; Leclerc, E. A.; Renouard, S.;
498 Tungmunthum, D.; Doussot, J.; Abassi, B. H.; Maunit, B.; Lainé, É.; Fliniaux, O.; Mesnard, F.; Hano, C.
499 Insight into the influence of cultivar type, cultivation year, and site on the lignans and related phenolic
500 profiles, and the health-promoting antioxidant potential of flax (*Linum usitatissimum* L.) seeds. *Molecules*
501 **2018**, 23, doi:10.3390/molecules23102636.
- 502 56. Tungmunthum, D.; Drouet, S.; Kabra, A.; Hano, C. Enrichment in Antioxidant Flavonoids of Stamen
503 Extracts from *Nymphaea lotus* L. Using Ultrasonic-Assisted Extraction and Macroporous Resin

- 504 Adsorption. *Antioxidants* **2020**, *9*.
- 505 57. Williams, G. M.; Iatropoulos, M. J.; Whysner, J. Safety Assessment of Butylated Hydroxyanisole and
506 Butylated Hydroxytoluene as Antioxidant Food Additives. *Food chem toxicol.* 1999; Vol. 37 (9-10):1027-
507 1038.
- 508 58. Hano, C.; Corbin, C.; Drouet, S.; Quéro, A.; Rombaut, N.; Savoie, R.; Molinié, R.; Thomasset, B.; Mesnard,
509 F.; Lainé, E. The lignan (+)-secoisolariciresinol extracted from flax hulls is an effective protectant of
510 linseed oil and its emulsion against oxidative damage. *Eur. J. Lipid Sci. Technol.* **2017**, *119*,
511 doi:10.1002/ejlt.201600219.
- 512 59. Drouet, S.; Doussot, J.; Garros, L.; Mathiron, D.; Bassard, S.; Favre-Réguillon, A.; Molinié, R.; Lainé, É.;
513 Hano, C. Selective Synthesis of 3-O-Palmitoyl-Silybin, a New-to-Nature Flavonolignan with Increased
514 Protective Action against Oxidative Damages in Lipophilic Media. *Molecules* **2018**, *23*, 2594,
515 doi:10.3390/molecules23102594.
- 516 60. Matkowski, A. Plant in vitro culture for the production of antioxidants—a review. *Biotechnol. Adv.* **2008**,
517 *26*, 548–560.
- 518