

Comparative physiological and biochemical mechanisms in Di, Tri, and Tetraploid Watermelon (*Citrullus lanatus* L.) grafted by Branches

Kaseb M O^{1,2}, Muhammad Jawad Umer¹, Hailelassie Gebremeskel¹, Eftekhar Mahmud¹, Weinan Diao¹, Pingli Yuan¹, Hongju Zhu¹, Shengjie Zhao¹, Xuqiang Lu¹, Nan He¹, Chengsheng Gong¹, Muhammad Anees¹, El-Eslamboly A.A.S.A², and Wenge Liu¹, *

1. Zhengzhou Fruit Research Institute, Chinese Academy of Agricultural Sciences, Henan Joint International Research Laboratory of Fruits and Cucurbits Biological Science in South Asia Zhengzhou 450009

2. Horticulture Research Institute, Agriculture Research Center, Giza, 12119, Egypt

10

Abstract

Polyploid seeds production is laborious, complicated, and costly work. Tetraploid and triploid plants produce a fewer number of seeds/fruit, and triploid embryos are fairly weak, covered with a more hardened seed coat as compared to diploid seeds. Here we investigated the interactive effect of new grafting technique of polyploid watermelon scion onto rootstock on plants' survival rate, biochemical, and hormones contents. In this study, three different branches, apical meristem (AM), branch with 1 node (1N), and 2 nodes (2N) from di, Tri, and tetraploid watermelon plants, were used as scion and grafted onto squash rootstock. The results showed highly significant differences between polyploid watermelon when 1N using as a scion, tetraploid showed maximum survival rates, higher contents of hormones, and antioxidants (AOX) activities, compared to diploid, these may be the possible reasons for high compatibility in tetraploid and degrading the grafting zone in diploid. RTq-PCR results confirm that the expression of genes linked to compatibility is consistent with the hormonal and AOX activities. This study provides an alternative and economical approach to produce more tetraploid and triploid plants for breeding or seeds production by using branches as scions.

Keywords: watermelon, polyploidy, grafting, branches

31 **1. Introduction**

32 Watermelon is the fifth most-consumed flesh fruit worldwide and occupies 7% of the
33 cultivated land for fruits and vegetables (FAOSTAT 2018). Seedless watermelons are
34 the most desirable watermelon cultivars presently available to consumers and command
35 a high price and more excellent quality than seeded watermelon [1]. Seedless
36 watermelons are produced by crossing a tetraploid seed parent with a diploid pollen
37 parent [2]. Tetraploid induction can be done by different methods like applying aqueous
38 chemicals (colchicine, Oryzalin, and Dinitroaniline) solution to the growing apex of
39 diploid seedlings or by soaking diploid seeds before germination [3]; these chemicals
40 are toxic to the plant material treated, so the percentage of success was low [4].
41 However, seedless watermelon production has been hampered by high seed cost and
42 poor seed germination. High seed cost has generally been because of difficulties in
43 obtaining a sufficient number of tetraploid individuals and the low number of seeds
44 only 5-20 seeds/fruit as compared to diploid fruits (600 seeds/fruit). It is important to
45 find a more economical way to increase the number and quality of triploid seeds [5-13]
46 Grafting has a very long history, and evidence of its use has been found in ancient
47 civilizations, e.g., in 1560 BC in China, as discussed by Aristotle (384-322 BC) [14].
48 Recently, the practice of grafting watermelons has become popular in the world
49 because it can improve disease resistance, tolerance to abiotic stress, fruit quality, and
50 plant size [15-18]. In recent years, splice grafting is one of the most commonly used
51 grafting methods, has achieved about 90% of grafting success, because it is the fastest,
52 most efficient [19-23].
53 Graft compatibility can be defined as a complex biochemical and structural process that
54 begins with an initial wound response, followed by a callus formation, the creation of a
55 continuous cambium, and a functional vascular system between scion and rootstock
56 [24]. All of these steps can determine the future of a grafted plant [25]. However, still
57 now no general rules, on how these treatments affect incompatibility responses [26,27].
58 In fruit trees, graft incompatibility occurs due to the accumulation of phenolic
59 compounds [28]. [29] found that reduced auxin and lignification was associated with
60 high phenols in incompatible grafts and may suggest weak graft unions, or reactive
61 oxygen species [30-33], or hormone imbalance [26,34,35]. IAA, and cytokinin, play an
62 important role in regulating stock–scion interactions [36], and required for vascular
63 bundle regeneration in the graft union [37,38]. Plant hormones are important for

vascular formation; auxin is important for the differentiation of vascular tissues and wound healing [15,18,32,33,39-45]. Reactive Oxygen Species (ROS) accumulation leads to cell death, and oxidative damage depends on the balance between the production of antioxidant AOX enzymes and ROS [46,47]. Increase the activity of defense enzymes such as Peroxidase POD, and catalase CAT can scavenge ROS in plants, and improving the resistance of plants to stress [48,49]. POX and CAT convert H_2O_2 to H_2O [50]. So the difference relation between AOX and ROS during the healing process could be used as a rapid mechanism to verify incompatibility [51]. Lignin is abundant in woody plants and primarily contributes strength to the cell wall; so, decreased lignin would be antagonistic to a strong graft union [52].

Up to date, the cause and underlying mechanism for grafting incompatibility remain elusive, and there is no report on genes associated with grafting incompatibility [53,54]. More research work needs to be done to fully understand the mechanisms of graft compatibility and incompatibility [41,52].

The purpose of this research was to study the effect of genome duplication on graft compatibility, by comparing the factors and parameters which lead to graft compatibility or incompatibility. This study provides information at the molecular and physiological levels for the mechanism compatibility in polyploid watermelons, which should explain the mechanism of compatibility in polyploid watermelon, also to try a new technique for vegetable propagation by using branches as scions, three different branches were taken from mother plants and graft on squash rootstock to study the success of this method. This method will add an option for breeders and seed producers to save time and money using an asexual method to increase plant numbers.

2. MATERIALS AND METHODS

2. 1. Plant materials

Seedless watermelons are triploids ($3n=33$) produced by crossing a tetraploid seed parent with a diploid ($2n=22$) pollen parent. Tetraploid induction was done by applying colchicine to the growing apex of diploid seedlings. Polyploid seeds for one variety (*mimei*), which is homozygous and genetically stable, and passed the achievement appraisal of the Chinese Department of Agriculture in 1990 and won the second prize of science technology progress of the Department of Agriculture in 1991 [55], were used as a mother plant and squash interspecific hybrid (*xijiaqiangsheng*) which widely used in China as a rootstock, obtained from the polyploid watermelon group - Zhengzhou Fruit research institute (CAAS) China. Tetraploid, triploid, and diploid watermelon and rootstock seeds were sown in seedling cell trays with 32 cells at the intelligent greenhouse of Zhengzhou, Henan province, China. After 50-60 days from the date of transplanting, good and healthy mother plants free of pests and diseases, especially virus-free, were selected for obtaining different types of scions. From the healthy mother plants, suitable branches were chosen for scions. Three types of branches (Figure 1) were taken from mother plants, (a) apical meristem (AM), (b) the branch having one node and one leaf (1N), and (c) the branch has two nodes and one leaf (2N) as mentioned by [10]. The grafting process was performed after 15 and 20 days after rootstock seeds sowing. The splice grafting method was used [19] (Figure S1). Rootstock seedlings were subjected to adaptation before and after grafting to increase the survival rate, as mentioned by [56,57].




Figure 1. Three types of branches were taken from mother plants to use as scions in di, Tri, and tetraploid watermelon, (A) apical meristem (AM), (B) branch has one node, and one leaf (1N), and (C) branch has two nodes (2N)

As the survival rate by using one node was higher and had significant differences between Di, Tri and Tetra watermelon, the samples (grafting union) from branch with one node (1N) Were collected at three different stages, (0, 3, and 15 days after grafting) for the determination of hormones, AOX and RNA extraction. Plants were cultivated in the growth chamber and the growth condition was set to 25-30 °C temperature and 60-85% humidity. The sensor in the center of each experimental plot was used to record data for temperature and humidity (THtool-V151_En; Campbell Scientific Ltd., China).

2.2. Data collection

2.2.1. Survival rates

Survive rates were investigated after 15 days using the formula [49].

$$\text{Survival rate} = \frac{\text{Number of survived grafted plants}}{\text{Total number of grafted plants}} \times 100$$

2.2.2. Detection of Hormones Content by ELISA

Samples were collected from the graft union at 0, 3, and 15 days after grafting (DAG) with three biological replicates. The contents of endogenous Indole-3-acetic acid (IAA) and Zeatin Riboside (ZR) were measured with the Enzyme-Linked Immunosorbent Assay (ELISA) method (College of Agriculture and Biotechnology, China Agricultural University) with three biological replicates for each set of treatments. The determination of hormonal contents was performed as outlined by [36].

2.2.3. Assay for antioxidants enzymes activity and H₂O₂ contents

SOD assay kit/ YX-C-A500 was used for the measurement of superoxide dismutase activity at 560 nm wavelength. CAT assay kit/BC0200 for Catalase assay using 240 nm wavelength, POD assay kit/ YX-C-A502 for peroxidase activity using 570 nm wavelength, and H₂O₂ assay kit/ YX-C-A400 for Hydrogen peroxidase content using 415 nm wavelength, (Sino best biological technology co, Ltd, Beijing, China) according to the manufacturer's instructions. Three biological replications from grafting union at 0, 3, and 15 (DAG) from three different plants for each replicate were collected for analysis. The activities of the antioxidant enzymes were expressed as unit U/g FW sample.

2.2.4. Quantification of biochemical at different stages

All experiments were carried out using starch assay kit/ YX-C-C400 for starch content estimation at 620 nm wavelength, lignin assay kit/ YX-C-B636 for lignin content using 280 nm wavelength, and phenol assay kit/ YX-C-A507 for total phenols content using

760 nm wavelength. (Sino best biological technology co, Ltd, Beijing, China) according to the manufacturer's instructions. Three biological replications from grafting union at 0, 3, and 15 DAG from three different plants for each replicate were collected for analysis.

2.2.5. Identification of genes involved in the compatibility mechanisms.

The protein sequences of genes involved in compatibility mechanisms were downloaded from the watermelon database <http://cucurbitgenomics.org/organism/2>. A total of 26 genes involved in compatibility mechanisms were identified based on the similarity index between protein sequences. The protein sequences for compatibility mechanisms in watermelon were downloaded from the 97,103 Watermelon Genome Database. MEGA 6 software was used to draw phylogenetic trees of selected genes. The clustalW tool was first used for the alignment. Afterward, the neighbor-joining method, with 1000 bootstrap replicates, was used [58,59]. The phylogenetic trees were constructed using protein sequences of the compatibility mechanisms.

2.2.6. Characteristics and structural analysis of genes responsible for compatibility mechanisms in watermelon

Gene names, accession number of genes, genomic lengths, coding sequences (CDS) lengths, protein sizes, and isoelectric points (pIs) and Mw (Da) were downloaded from <http://cucurbitgenomics.org/organism/1> and ExPASy <http://web.expasy.org/computeipi/> databases. Gene Structure Display Server (GSDS), a web-based bioinformatics tool, was used to give a structural representation to selected watermelon genes responsible for sugars and organic acids metabolism and transporter genes, including upstream/downstream regions, coding sequences (CDS), and intron numbers were constructed [60].

2.2.7. Isolation of RNA and first-strand cDNA synthesis

Using a Total RNA Kit (Tiangene, China), we extracted the total RNA from the grafting union according to the manufacturer's instructions. Samples from grafting unions were taken at three stages (0, 3, and 15 days after grafting). We detected RNA degradation and contamination on a 1% agarose gel. Using a NanoDrop ND-1000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA) and an Agilent 2100 Bioanalyzer (Agilent Technologies, CA, USA), the quantity quality of RNA was checked according to manufacturer's instructions. To carry out quantitative reverse-

transcription PCR (RTq-PCR), cDNA was synthesized from RNA with M-MLV Reverse Transcriptase (Promega, USA) and diluted 20 ng/ul.

2.2.8. RTq-PCR expression analysis of genes involved in compatibility mechanisms

Genes linked to the compatibility mechanisms were selected from the watermelon genome database (<http://cucurbitgenomics.org/organism/21>). To check the expression patterns of selected genes in grafted watermelon at 0, 3, and 15 DAG, quantitative reverse transcription PCR (RTq-PCR) was performed. The entire data were analyzed using the $2^{-\Delta\Delta C_t}$ method [61]. Three independent biological replicates were used for gene expression analysis [62]. Actin “*cla016178*” was used as a reference gene [58].

2.2.9. Experimental design and statistical analysis

Data were statistically analyzed using analyses of variance (ANOVA), with the Stat soft statistical package MSTATC software program (Michigan State University, East Lansing, MI, USA). Mean values of all data for all parameters were subjected to analysis of variance (ANOVA) by using SAS (version 9) computer software programs [SAS inst., 2002]. A least significant difference (LSD) was used to compare significant means at 5% probability level.

3. RESULTS

3.1. Survival rates of Di, Tri, and tetraploid watermelon

Plant survival rates were recorded for all the grafting combinations among diploid (Di), triploid (Tri), and tetraploid (Tetra), grafted by different parts from mother plants on interspecific rootstock during two consecutive seasons, March and August 2018. The statistical analysis showed highly significant differences between Di, Tri, and Tetra when 1N was used as a scion, and the combined analysis for two seasons gave 45.45, 55.45, and 78.03% in Di, Tri, and Tetra watermelon respectively. Grafting by AM gave the highest survival rate 93.3, 96.67, and 96.67 % in the first season and 96.67, 100, and 100 % in the second season in Di, Tri, and Tetra watermelon, respectively, but no significant differences were observed between polyploids. Also, branch with 2 nodes (2N) did not show any significant differences between polyploids in terms of survival rate 86.67, 83.33 and, 83.33 % in the first season and 90, 93.33, and 96.67 % in the second season in Di, Tri, and Tetra watermelon respectively, (Figure: 2 and 3) (Figure S2, and S3). Our results also showed the suitability of vegetative propagation using AM, 2N, and 1N branches as scions in tetraploid watermelon, it is a new method for seedless watermelon seed producers.




Figure 2. The effect of Scion/rootstocks combinations on the survival rate of polyploid watermelon grafting union at 15 days after grafting observed a significant difference between AM and 1N. **AM:** Apical Meristem, **1N:** Branch with 1Node, **2N:** Branch with 2Nodes.



Figure 3. Effect of Scion/rootstocks combinations on the survival of polyploid watermelon grafting union at 15 days after grafting. No significant differences between polyploids when AM and 2N were used as scions, but 1N showed highly significant between di and tetraploid. New branches can be observed in cycles. (A): Apical Meristem (AM), (B) Branch with 2Node (2N), (C): Branch with 1Nodes (1N), DAG: Days after Grafting.

3.2. Measurement of IAA and ZR in the grafting union among Di, Tri, and tetraploid watermelon at different days after grafting

Ploidy level (Di, Tri, and Tetra) exerted a significant effect on IAA content in the graft union at all stages 0, 3, and 15 days after grafting (DAG) (Figure 4). At 0 DAG, the contents of IAA in Tetra were 2.26 and 2.35-fold higher than Di in the first and second season, respectively, while in Tri, IAA was 1.4 and 1.38-fold higher than Di in the first and second season, respectively. Also, the increment rate of IAA in the graft union at 3 DAG in all polyploids; were 23.88, 37.93, and 23.68% in the first season and 26.35, 35.82, and 25.34% in the second season in Di, Tri, and Tetra watermelon, respectively. The content of IAA was higher in Tetra, followed by Tri and Di watermelon at 0, 3, and 15 DAG, respectively.

Whereas ZR showed no significant differences between Di and Tetra at 0 DAG, while at 3 and 15 DAG, there were highly significant differences in both seasons (Figure 4). The results at 3 and 15 DAG showed that the content of the ZR starts to increase significantly in Tetra than Di and Tri. At 3 DAG, the content of ZR was increased by 59.43 and 42.9% during the first, and second season respectively, and by 18.69 and 37% at 15 DAG during both seasons.




Figure 4. Effect of Scion/rootstocks combinations on IAA and ZR hormones on polyploids watermelon grafting union at 0, 3, and 15 days during March 2018 and August 2018 cropping seasons. The values are means of fifteen independent plants (n = 15). Different letters (a, b, c, d, and e) in rows indicate significant differences at $P < 0.01$ using Duncan's multiple range test. **IAA:** Indole acetic acid, **ZR:** Zeatin Riboside, **Di:** Diploid, **Tri:** triploid, **Tetra:** Tetraploid, **DAG:** Days after grafting.

3.3 Measurement of Antioxidants (POD, SOD, and CAT) and H_2O_2 in the grafting union among Di, Tri, and tetraploid watermelon at different days after grafting

The increase of antioxidants (AOX) activity such as peroxidase (POD) and catalase (CAT) can scavenge reactive oxygen species (ROS) in plants, thus improving the resistance of plants to stress. Antioxidant enzyme activities were substantially affected by the grafting process between polyploid watermelon [48,49]. After grafting, the results of POD showed highly significant differences between different ploidy watermelons during both seasons (Figure 5). The highest POD activities 2.87 and 2.05-fold was recorded in Tetra at 0 DAG in comparison to Di, and 1.48 and 1.19-fold as compared to Tri in the first and second season, respectively. Higher POD activities in the graft union were observed at 3 DAG in all polyploids, with increment rates of 124.84, 98.88, and 89.83% in the first season and 22.34, 16.25, and 63.21% during the second season in Di, Tri, and Tetra watermelon respectively. While at 15 DAG, the increment rates were (46.86, 47.04, and 47.09%) in the first season and 80.53, 18.64, and 17.74% during the second season in Di, Tri, and Tetra watermelon, respectively.




Figure 5. Effect of Scion/rootstocks combinations on Peroxidase (POD), Superoxide dismutase (SOD), and Catalase (CAT) activities at polyploid watermelons at 0, 3, and 15 days during crop season March 2018 and August 2018 cropping seasons. The values are means of fifteen independent plants (n = 15). Different letters (a, b, c, d, e, and f) in rows indicate significant differences at P < 0.01 using Duncan's multiple range test. **Di:** Diploid, **Tri:** triploid, **Tetra:** Tetraploid, **DAG:** Days after grafting.

SOD activity was observed higher in Tetra and Tri watermelon as compared to Di watermelon (Figure 5). At 0 DAG, SOD activities were 3.76 and 3.19-fold higher in Tri than diploid in both seasons, respectively. At 3 DAG, SOD activities were 2.9 and 2.97-fold higher than Di in the first and second season, respectively. In comparison, at 15 DAG, the SOD activities in Tri were 4.32 and 2.7-fold higher than Di in both seasons. In Tetra SOD activities at 0 DAG were 3.64 and 4.49-fold higher than Di in the first and second season, respectively, while at 3 DAG, it was 2.53 and 3-fold higher than Di in the first and second season, respectively. Moreover, at 15 DAG, the contents of SOD in Tri were 4.47 and 4.19-fold higher than Di in the first and second season, respectively. Generally, there are significant differences between Tetra and Di at 0, 3, and 15 DAG.

Significantly higher CAT content was observed in Tetra than Di and Tri (Figure 5). At 0 DAG it was 1.85 and 2.35-fold higher in Tetra than Di in the first and second season respectively, and 1.31 and 1.7-fold higher than Tri in the first and second season,

respectively. At 0 DAG, CAT activity was 1.4 and 1.38-fold higher than Di in the first and second season, respectively. CAT activities in the graft union start to increase at 3 DAG in all polyploids, but the increase at Tri was more than Di and Tetra, with increment rates of 9.19, 122.29, and 33.47% in the first season and 26.35, 35.52, and 25.34% in the second season in Di, Tri, and Tetra watermelon respectively.

Reactive Oxygen Species (ROS) like hydrogen peroxide (H₂O₂) accumulation leads to cell death, and oxidative damage depends on the balance between the production of AOX and ROS [46,47]. H₂O₂ contents were higher in Triploid than Di and Tetra at 0 DAG stage (Figure 6). It was 1.49 and 1.65-fold higher than Di in the first and second season, respectively, and 1.14 and 1.2-fold higher than Tetra in the first and second season, respectively. But the increment rates of H₂O₂ activities in the graft union at Tetra starts to increase at 3 DAG less than Di, and, Tri; the increment rates were 51.51, 5.43, and 4.08% in the first season and 42.53, 8.62, and 8.79% in Di, Tri, and Tetra plants respectively. These may be because of the high activities of AOX in Tetra than Di and Tri during the grafting process.




Figure 6. Effect of Scion/rootstocks combinations on Hydrogen peroxide (H₂O₂) contents at polyploid watermelon grafting union at 0, 3, and 15 days during crop season March and August 2018. The values are means of fifteen independent plants (n = 15). Different letters (a, b, c, and d) in rows indicate significant differences at P < 0.01 using Duncan's multiple range test. **Di:** Diploid, **Tri:** triploid, **Tetra:** Tetraploid, **DAG:** Days after grafting.

3.4. Measurement of Lignin, phenols, and starch contents in the grafting union among Di, Tri, and tetraploid watermelon at different days after grafting

Lignin content was significantly different between Di and Tetra watermelons in both seasons (Figure 7). Lignin contents were higher in Tri, and Tetra than Di at 0 DAG, where it at Tetra were 2.7 and 2.41-fold higher than Di in the first and second season respectively, and at Tri were 2.67 and 2.41-fold higher than Di in the first and second season, respectively. But because of callus induction (branchime cells) in the graft union at 3 DAG stage [18], the lignin contents start to decrease; nevertheless, the decrement rate in Tetra was less than Di. The increment rates of lignin contents in the graft union at 3 DAG were 51.68, 21.5, and 6.33% in the first season, and 51.68, 16.98, and, 12.32% in the second season at Di, Tri, and, Tetra watermelon respectively. At 15 DAG, the lignin contents start to increase by 164.32, 55, 08, and, 59.18% in the first season, and 251.37, 264, and, 188.11% in the second season at Di, Tri, and, Tetra watermelon respectively.

The results of phenols showed significant differences between Di and Tetra watermelons in both seasons at 3 DAG (Figure 7). The contents of phenols in Tetra were 1.4, 1.7, and 1.1-fold higher than Di at 0, 3, and 15 DAG, respectively, in the first season, and 1.1, 1.3, and 1.2-fold higher than Di at 0, 3, and 15 DAG respectively, in the second season.

The results of starch showed highly significant differences between ploidy watermelons in the first season during the grafting process, while in the second season, no significant differences between ploidy levels (Figure 7). The results showed that, the starch contents in Tri and Tetra highly significant than di, where it at 0 DAG were (1.57 and 1.48-fold in Tri and Tetra higher than Di in the first season, while in the second season, no significant differences between polyploidy levels. Also, the contents of starch start to increase at 15 DAG in all polyploids, but the increment rates at Tri and Tetra were more than Di 9.68, 14.09, and 16.58% at the first season in Di, Tri, and Tetra watermelon, respectively.




Figure 7. Effect of Scion/rootstocks combinations on lignin, Phenols, and Starch contents at polyploid watermelon grafting union at 0, 3, and 15 days during crop season March and August 2018. The values are means of fifteen independent plants (n = 15). Different letters (a, b, c, d, e, and f) in rows indicate significant differences at P < 0.01 using Duncan's multiple range test. **Di:** Diploid, **Tri:** triploid, **Tetra:** Tetraploid, **DAG:** Days after grafting, **ns:** not significant.

3.5 Phylogenetic analysis of selected genes linked to compatibility mechanisms in the grafting union among Di, Tri, and tetraploid watermelon

BlastP searches in the watermelon Genome Database, using *Arabidopsis lyrata* subsp. *Lyrata*, *Arabidopsis thaliana*, *Citrullus colocynthis*, *Citrus sinensis*, *Coffea canephora*, *Cucumis melo*, *Cucumis sativus*, *Cucurbita maxima*, *Cucurbita moschata*, *Cucurbita pepo*, *Dimocarpus longan*, *Eutrema salsugineum*, *Glycyrrhiza uralensis*, *Gossypium arboreum*, *Hevea brasiliensis*, *Kandelia candel*, *Morus notabilis*, *Oryza punctate*, *Oryza sativa*, *Pinus Sylvestris*, *Pisum sativum*, *Populus tremula x Populus tremuloides* (Hybrid aspen), *Populus trichocarpa*, *Prunus persica*, *Ricinus communis*, *Salvia splendens*, *Solanum tuberosum*, *Theobroma cacao*, *Vitis vinifera*, *Zantedeschia aethiopica*, *Zea mays*, *Ziziphus jujube* and *Zostera marina* protein sequences for genes linked to H₂O₂, starch, POD, SOD, lignin, phenols, ZR, IAA, and CAT were used as a

query, permitted the identification of candidate genes in watermelon. Watermelon genes having high homology are summarized in the (Figure 8) (Table S1).




Figure 8. Maximum likelihood phylogeny of watermelon genes encoding key enzymes and transporters involved graft compatibility with those from *Arabidopsis lyrata* subsp. *Lyrata*, *Arabidopsis thaliana*, *Citrullus colocynthis*, *Citrus sinensis*, *Coffea canephora*, *Cucumis melo*, *Cucumis sativus*, *Cucurbita maxima*, *Cucurbita moschata*, *Cucurbita pepo*, *Dimocarpus longan*, *Eutrema salsugineum*, *Glycyrrhiza uralensis*, *Gossypium arboreum*, *Hevea brasiliensis*, *Kandelia candel*, *Morus notabilis*, *Oryza punctata*, *Oryza sativa*, *Pinus Sylvestris*, *Pisum sativum*, *Populus tremula* x *Populus tremuloides* (Hybrid aspen), *Populus trichocarpa*, *Prunus persica*, *Ricinus communis*, *Salvia splendens*, *Solanum tuberosum*, *Theobroma cacao*, *Vitis vinifera*, *Zantedeschia aethiopica*, *Zea mays*, *Ziziphus jujube*, and *Zostera marina*. The phylogenetic tree was constructed from protein sequences using the neighbor-joining method.

3.6 RT-qPCR analysis of genes regulating hormones, biochemical, and antioxidants in Di, Tri, and Tetraploid watermelon at different days after grafting

Expressions of genes regulating *CAT*, *POD*, *SOD*, *H₂O₂*, *IAA*, *Cytokinin (ZR)*, *phenols*, *lignin*, and *starch* were checked at three different stages (0, 3, 15 DAG). Primer 3 was used to design primers for RTq-PCR analysis (Table S2). The range of all PCR products was from 80 to 200bp. The specificity of PCR amplification was observed by monitoring the dissociation curves during qPCR using a Roche LightCycler 480 II [58,63]. Transcript levels of these genes were linked to the compatibility or incompatibility as they are involved in regulating important biochemical and

antioxidants. *WM-IAA-1* (*ClCG05G010400*), *WM-ZR-1* (*ClCG11G005450*), had higher expression at all the time points in tetra watermelon followed by triploid and diploid watermelon in both seasons. *WM-POD-1* (*ClCG07G008740*), *WM-CAT-1* (*ClCG11G018720*), *WM-SOD-1* (*ClCG04G005370*) had a higher expression in Tetra as compared to Tri and Di watermelon at all days after grafting (Figure 9). *WM-H₂O₂-1* (*ClCG10G013100*) had a lower expression level in Tetra watermelon than Tri and Di in both the seasons. Genes regulating starch and lignin contents in watermelon, including *WM-Starch-1* (*ClCG01G018090*), *WM-Lignin-1* (*ClCG11G016110*), had high expressions in Tetra watermelon as compared to Tri and Di watermelon confirming their active role in starch and lignin accumulation at all the time points of sampling. Among the genes controlling phenolic contents, *WM-Phenol-1* (*ClCG09G008770*) had higher expressions in Tri and Di watermelon than Tetra watermelon at all the sampling stages.




Figure 9: Heatmap of selected genes in Di, Tri, and Tetraploid watermelon at different days after grafting in two consecutive seasons.

4. Discussion:

Triploid watermelon seeds are produced by cross-pollination between Di and Tetra, but Tetra seeds production is a very hard process [11]. So it is important to find a more economical way to increase the number and quality of triploid seeds [13]. In this study, we tested three different parts from mother plants as a scion, apical meristem (AM), branch with 1, and 2 nodes (1N and 2N) to increase the number of plants using vegetative propagation from mother plants with grafting technique. Our results showed a higher survival rate (Figure 2) with three kinds of branches used in Tetra plants, whereas in case of triploid AM and 2N give a high survival rate while 1N gave a low survival rate, but in Di just AM gave high survival rate. Suggesting that Tetra grafted with branch with 1 and 2 nodes and apical meristem, triploid with grafted with AM and 2N and Di grafted with AM are the best possible way of vegetative propagation in watermelon.

Plant hormones play important roles in plant growth, development, and response to biotic and abiotic cues and vascular formation in the graft junction [15,32,33,40-45]. Auxin, and cytokinin, play an important role in regulating stock–scion interactions [18,36]. Cell divisions happened within 2-3 DAG in the graft junction and have the highest hormone levels during this period [18,39,40,64]. During the grafting process, the peak of IAA is observed to occur within 3 DAG in the scion, as reported by [65]. IAA and Zeatin Riboside (ZR) are required for vascular bundle regeneration in the graft union [37,38]. The main cause of incompatibility is the occurrence of hormonal imbalance [34]. A low indole-3-acetic acid (IAA) content in incompatible combinations may then affect the differentiation of xylem and phloem, as well as lignification [17,28,66]. We compared the IAA and ZR contents in the graft union between Di, Tri, and Tetra watermelon (Figure 4). The IAA concentration in the Tetra combination was significantly higher than that in the Di, and Tri combination ($p < 0.05$). Our results showed high compatibility in Tetra which has high content and high increment rates of hormones than Di especially at 3 DAG (critical period) and 15 DAG, this can explain high survival rates in Tetra, this results were agreement with [67-69]

On the other side, incompatibility results from the stress-induced during the healing response, as reported earlier by [41]. In this study in (Figure 5), POD, SOD, and CAT activities showed a high increase during the healing process in Tetra and Tri. Also, the content of H_2O_2 (Figure 6) didn't increase during the healing process in Tri, because of

the high activities of antioxidants, which leads to scavenging oxygen radicals. These results were in accordance with [70-74], who found that genome duplication gave high resistance for salt stress because of the high contents of hormones and highly activities of antioxidants more than Di. Also, ploidy results in chromosome doubling leading to gene doubling, which results in higher expression, thus causing an increase in protein contents [75]. Polyploidy is more tolerant of abiotic stress than Di it was reported in watermelon [76], rice [73,77], citrus [72], black locust [78], honeysuckle [79], kinnow mandarin [80], cotton [81], and rangpur lime [82]. Peroxidase and catalase activities were increased in the grafted plants [83].

Generally, antioxidant enzymes were usually studied in the responses to abiotic and biotic stresses, but not often for graft stress; in this study, we suggest grafting as stress because of, cut or wound stress, complete dark, and high humidity stress, especially in the first three days after grafting (healing response) [12,41]. The most important critical period in the grafting healing process was at 2 and 3 days after grafting [18]. The results in both seasons showed that the activities of SOD, POD, and CAT were significantly different at the grafting healing process between polyploid watermelons (Figure 6). It has been reported already by [49,83] that high activities of POD and CAT, during the healing process have a high ability to scavenge reactive oxygen species (ROS) and hydrogen peroxide (H_2O_2) in plants. H_2O_2 product when the plant is wounded or stressed and caused cell death [84].

Higher enzymatic activities and accumulation of phenolic compounds are typically associated with plant stress resistance [85]. Phenols contribute to the elimination of reactive oxygen species (ROS) [86]. In our results, the content of phenols was higher in Tetra than Di and Tri in all stages, but because of the accumulation of phenols leads to incompatibility [31,32], a higher decrease in phenol contents were observed at later stages of grafting (15 DAG) (Figure 7). The decrement rates between 3 and 15 DAG in Tetra were (38.24, and 30.77%) in the first and second season, respectively, while in Di, the content of phenols increase at 15 DAG in the first season and decrease (27%) in the second season. With Tri, the decrement rates were (15, and 18.7%) at 15 DAG in the first and second season, respectively. These results can explain the high compatibility of Tetra and low compatibility in a Di. Also, our results are similar to [85], who found that the phenolic content and lignin content were higher in the compatible combination.

451 **5. Conclusion**

452 Seedless watermelons seeds production faces many problems. The current study
453 concludes that vegetative propagation by branches grafting could be used in Tri and
454 Tetra watermelon seed production. The high contents of hormones and high activities
455 of antioxidants in Tetra more than Di because of the genome duplication, which gave
456 high compatibility and high survival rates of grafting. Grafting using branches can be a
457 good alternative to promote vegetative propagation in Tetra watermelon. These
458 findings add significant information to our existing knowledge of compatibility in
459 watermelon by ultimately helping in vegetative propagation, breeding watermelon
460 varieties of desirable values.

461

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgments

The authors thank all the members of the Polyploid Watermelon Research Group, Zhengzhou Fruit Research Institute; Chinese Academy of Agricultural Sciences.

Kaseb M O^{1,2}, Muhammad Jawad Umer¹, Haileslassie Gebremeskel¹, Eftekhar Mahmud¹, Weinan Diao ¹, Pingli Yuan¹, Hongju Zhu ¹, Shengjie Zhao¹, Xuqiang Lu ¹, Nan He ¹, Chengsheng Gong¹, Muhammad Anees¹, El-Eslamboly A.A.S.A², and Wenge Liu^{1, *}

1. Zhengzhou Fruit Research Institute, Chinese Academy of Agricultural Sciences, Henan Joint International Research Laboratory of Fruits and Cucurbits Biological Science in South Asia Zhengzhou 450009
2. Horticulture Research Institute, Agriculture Research Center, Giza, 12119, Egypt

Supplementary Materials:

- Figure S1:** Splice grafting method steps in polyploid watermelon plants
- Figure S2:** Effect of Scion/rootstocks combinations on the survival of polyploid watermelon grafting using (AM) as scion. (A): at 0 DAG, (B) 15 DAG, and (C): 25 DAG, DAG: Days after Grafting.
- Figure S3:** Effect of Scion/rootstocks combinations on the survival of polyploid watermelon grafting using (2N) as scion. (A): at 15 DAG, and (B) 25 DAG.
- Figure S4:** Structure of genes involved in graft compatibility during watermelon grafting process. Yellow boxes and black lines indicating exons and introns respectively, while blue boxes at the both end of each gene indicates untranslated regions (UTRs).
- Table S1:** Genes linked to graft compatibility having high homology with other genes selected from different plant species.
- Table S2:** Primers used for qRT-PCR in this study.

Author Contributions

KMO and WL conceived and designed the experiments. KMO performed the experiments. KMO and MJU analyzed the data and wrote the manuscript. HG, EM, WD, PY, HZ, SZ, XL, NH, CG, MA, and EL participated in the experimentations and

493 analysis. MJU, MA, HZ, and WL edited the manuscript. All authors read and approved
494 the last version.

495

FUNDING

This research was supported by the Agricultural Science and Technology Innovation Program (CAAS-ASTIP-2016-ZFRI), the National Key R&D Program of China (2018YFD0100704), the China Agriculture Research System (CARS-25-03), the Special Fund for Basic Scientific Research of Chinese Academy of Agricultural Sciences(1610192020103) and National Natural Science Foundation of China (No.31471893; 31672178).

Data availability

The authors declare that the data supporting the study findings are presented in the article, and Supplementary Information files are available from the corresponding author upon request.

508

6. References

509 1. Marr, C.W.; Gast, K.L.B. Charles W. Marr and Karen L.B. Gast 2 1. **1991**, 105-106.
510 2. Jaskani, M.J.; Kwon, S.W.; Kim, D.H. Comparative study on vegetative, reproductive
511 and qualitative traits of seven diploid and tetraploid watermelon lines. *Euphytica*
512 **2005**, *145*, 259-268.
513 3. Omran, S.; Guerra-Sanz, J.; Cardenas, J.G. Methodology of tetraploid induction and
514 expression of microsatellite alleles in triploid watermelon. In Proceedings of
515 Proceedings IXth Eucarpia meeting on genetics and breeding of Cucurbitaceae; pp.
516 381-384.
517 4. Nasr, M.; Habib, H.; Ibrahim, I.; Kapiel, T. In vitro induction of autotetraploid
518 watermelons using colchicine and four dinitroaniline compounds. In Proceedings of
519 Proceedings of International Conference of Genetic Engineering and Its Applications,
520 Sharm Elsheik, Egypt; pp. 8-11.
521 5. Data, R.U.S.A. Patent Application Publication (10) Pub . No .: US 2005 / 0034197 A1.
522 **2005**, *1*.
523 6. Pradesh, A. Studies to Improve Seed Germination in Tetraploids Watermelon (
524 *Citrullus lanatus* Thunb .). **2014**, *4*, 1-5.
525 7. NERSON, H.; PARIS, H.; KARCHI, Z.; SACHS, M. Seed treatments for improved
526 germination of tetraploid watermelon. *HortScience* **1985**, *20*, 897-899.
527 8. Cobb, B.G. Excess Moisture and Seedcoat Nicking Influence Germination of Triploid
528 Watermelon. **2000**, 10.21273/HORTSCI.35.7.1355, 1355-1357,
529 doi:10.21273/HORTSCI.35.7.1355.
530 9. Grange, S.; Leskovar, D.I.; Pike, L.M.; Cobb, B.G. Seedcoat Structure and Oxygen-
531 enhanced Environments Affect Germination of Triploid Watermelon. **2014**,
532 10.21273/JASHS.128.2.0253, doi:10.21273/JASHS.128.2.0253.
533 10. Khereba, A.H.A.; Bekhit, R.S.; Kamooh, A.A.; El-Eslamboly, A.A.S. New grafting method
534 for seedless watermelon plants propagation. *J. Agric. Sci. Mansoura Univ.*, **2008**,
535 *33(11)*, 8071-8090.
536 11. Ishtiaq Ahmad; Tanveer Hussain; Muhammad Nafees; Maryam; Moazam Jamil; Irfan
537 Ashraf; Muhammad Fakhar-u-Zaman Akhtar; Muhammad Iqbal; Muhammad Rafay;
538 Ali, T.R.a.L. Morphological Dissimilarity Between Tetraploid and Diploid Watermelon
539 (*Citrullus lanatus* Thunb.). *World Applied Sciences Journal* **2013**, *21 (6)*: 858-861,
540 doi:10.5829/idosi.wasj.2013.21.6.169.
541 12. Schwarz, D.; Rouphael, Y.; Colla, G.; Venema, J.H. Grafting as a tool to improve
542 tolerance of vegetables to abiotic stresses: Thermal stress, water stress and organic
543 pollutants. *Scientia Horticulturae* **2010**, *127*, 162-171,
544 doi:10.1016/j.scienta.2010.09.016.
545 13. Solmaz, İ.; Sari, N.; Kombo, M.D.; ŞİMŞEK, İ.; Hussein, S.; Namli, M. Rootstock capacity
546 in improving production and quality of triploid watermelon seeds. *Turkish Journal of*
547 *Agriculture and Forestry* **2018**, *42*, 298-308, doi:10.3906/tar-1801-59.
548 14. Tsutsui, H.; Notaguchi, M. The Use of Grafting to Study Systemic Signaling in Plants.
549 *Plant Cell Physiol* **2017**, *58*, 1291-1301, doi:10.1093/pcp/pcx098.
550 15. Goldschmidt, E.E. Plant grafting: new mechanisms, evolutionary implications.
551 *Frontiers in Plant Science* **2014**, *5*, doi:10.3389/fpls.2014.00727.
552 16. Lee, J.-M.; Kubota, C.; Tsao, S.J.; Bie, Z.; Echevarria, P.H.; Morra, L.; Oda, M. Current
553 status of vegetable grafting: Diffusion, grafting techniques, automation. *Scientia*
554 *Horticulturae* **2010**, *127*, 93-105, doi:10.1016/j.scienta.2010.08.003.
555 17. Koepke, T.; Dhingra, A. Rootstock scion somatogenetic interactions in perennial
556 composite plants. *Plant Cell Rep* **2013**, *32*, 1321-1337, doi:10.1007/s00299-013-1471-
557 9.

- 558 18. Melnyk, Charles W.; Schuster, C.; Leyser, O.; Meyerowitz, Elliot M. A Developmental
559 Framework for Graft Formation and Vascular Reconnection in *Arabidopsis thaliana*.
560 *Current Biology* **2015**, *25*, 1306-1318, doi:10.1016/j.cub.2015.03.032.
- 561 19. Dabirian, S.; Miles, C.A. Increasing survival of splice-grafted watermelon seedlings
562 using a sucrose application. *HortScience* **2017**, *52*, 579-583,
563 doi:10.21273/HORTSCI11667-16.
- 564 20. Khankahdani, H.H.; Zakeri, E.; Saeedi, G.; Shakerdargah, G. Evaluation of different
565 rootstocks and grafting techniques on graft union percent, yield and yield components
566 of Watermelon CV. 'Crimson sweet'. *World Applied Sciences Journal* **2012**, *18*, 645-
567 651, doi:10.5829/idosi.wasj.2012.18.05.1716.
- 568 21. Petropoulos, S.A. Evaluation of rootstocks for watermelon grafting with reference to
569 plant development , yield and fruit quality. **2014**.
- 570 22. Mohamed, F.H.; El-hamed, K.E.A.; Elwan, M.W.M.; Hussien, M.N.E. Scientia
571 Horticulturae Evaluation of different grafting methods and rootstocks in watermelon
572 grown in Egypt. *Scientia Horticulturae* **2014**, *168*, 145-150,
573 doi:10.1016/j.scienta.2014.01.029.
- 574 23. Monteiro, G.C.; Goto, R.; Minatel, I.O.; de Sousa da Silva, E.; da Silva, E.G.; Vianello, F.;
575 Lima, G.P.P. Grafting, Agrochemicals, and Oxidative Enzymes as Factor for Plant Biotic
576 Resistance. In *Plant Health Under Biotic Stress: Volume 1: Organic Strategies*, Ansari,
577 R.A., Mahmood, I., Eds. Springer Singapore: Singapore, 2019; 10.1007/978-981-13-
578 6043-5_2pp. 37-57.
- 579 24. Assunção, M.; Canas, S.; Cruz, S.; Brazão, J.; Zanol, G.C.; Eiras-Dias, J.E. Graft
580 compatibility of *Vitis* spp.: the role of phenolic acids and flavanols. *Scientia*
581 *Horticulturae* **2016**, *207*, 140-145, doi:10.1016/j.scienta.2016.05.020.
- 582 25. Pina, A.; Cookson, S.J.; Calatayud, A.; Trinchera, A.; Errea, P. Physiological and
583 molecular mechanisms underlying graft compatibility. *Vegetable Grafting: Principles*
584 *and Practices* **2017**, 10.1079/9781780648972.0132, 132-154,
585 doi:10.1079/9781780648972.0132.
- 586 26. Fery, R.L.; Hasegawa, P.K. American Society for Horticultural Science Editorial Board ,
587 Volume 23 Leo Gene Albrigo. **1999**, 23.
- 588 27. Tamilselvi, N.A.; Pugalandhi, L. Graft Compatibility and Anatomical Studies of Bitter
589 Gourd (*Momordica charantia* L.) Scions with Cucurbitaceous Rootstocks. *International*
590 *Journal of Current Microbiology and Applied Sciences* **2017**, *6*, 1801-1810,
591 doi:10.20546/ijcmas.2017.602.202.
- 592 28. Pina, A.; Errea, P. Influence of graft incompatibility on gene expression and enzymatic
593 activity of UDP-glucose pyrophosphorylase. *Plant Science* **2008**, *174*, 502-509,
594 doi:10.1016/j.plantsci.2008.01.015.
- 595 29. Pina, A.; Errea, P.; Martens, H.J. Graft union formation and cell-to-cell communication
596 via plasmodesmata in compatible and incompatible stem unions of *Prunus* spp.
597 *Scientia Horticulturae* **2012**, *143*, 144-150, doi:10.1016/j.scienta.2012.06.017.
- 598 30. Ren, Y.; Guo, S.r.; Shu, S.; Xu, Y.; Sun, J. Isolation and expression pattern analysis of
599 CmRNF5 and CmNPH3L potentially involved in graft compatibility in
600 cucumber/pumpkin graft combinations. *Scientia Horticulturae* **2018**, *227*, 92-101,
601 doi:10.1016/j.scienta.2017.09.022.
- 602 31. Ren, Y.; Xu, Q.; Wang, L.; Guo, S.; Shu, S.; Lu, N.; Sun, J. Plant Physiology and
603 Biochemistry Involvement of metabolic , physiological and hormonal responses in the
604 graft-compatible process of cucumber / pumpkin combinations was revealed through
605 the integrative analysis of mRNA and miRNA expression. **2018**, *129*, 368-380,
606 doi:10.1016/j.plaphy.2018.06.021.

- 607 32. Chen, Z.; Zhao, J.; Hu, F.; Qin, Y.; Wang, X.; Hu, G. Transcriptome changes between
608 compatible and incompatible graft combination of Litchi chinensis by digital gene
609 expression profile. *Sci Rep* **2017**, *7*, 3954, doi:10.1038/s41598-017-04328-x.
- 610 33. Melnyk, C.W.; Gabel, A.; Hardcastle, T.J.; Robinson, S.; Miyashima, S.; Grosse, I.;
611 Meyerowitz, E.M. Transcriptome dynamics at Arabidopsis graft junctions reveal an
612 intertissue recognition mechanism that activates vascular regeneration. *Proc Natl*
613 *Acad Sci U S A* **2018**, *115*, E2447-E2456, doi:10.1073/pnas.1718263115.
- 614 34. Aloni, B.; Karni, L.; Deventurero, G.; Levin, Z.; Cohen, R.; Katzir, N.; Lotan-Pompan, M.;
615 Edelstein, M.; Aktas, H.; Turhan, E., et al. Physiological and biochemical changes at the
616 rootstock-scion interface in graft combinations between Cucurbita rootstocks and a
617 melon scion. *Journal of Horticultural Science and Biotechnology* **2008**, *83*, 777-783,
618 doi:10.1080/14620316.2008.11512460.
- 619 35. Kumpers, B.M.; Bishopp, A. Plant grafting: making the right connections. *Curr Biol*
620 **2015**, *25*, R411-413, doi:10.1016/j.cub.2015.03.055.
- 621 36. Mo, Z.; Feng, G.; Su, W.; Liu, Z.; Peng, F. Transcriptomic Analysis Provides Insights into
622 Grafting Union Development in Pecan (Carya illinoensis). *Genes (Basel)* **2018**, *9*,
623 doi:10.3390/genes9020071.
- 624 37. Lu, S.; Song, Y. Relation between phytohormone level and vascular bridge
625 differentiation in graft union of explanted internode autografting. *Chinese science*
626 *bulletin* **1999**, *44*, 1874-1878.
- 627 38. Davis, A.R.; Perkins-Veazie, P.; Sakata, Y.; López-Galarza, S.; Maroto, J.V.; Lee, S.-G.;
628 Huh, Y.-C.; Sun, Z.; Miguel, A.; King, S.R., et al. Cucurbit Grafting. *Critical Reviews in*
629 *Plant Sciences* **2008**, *27*, 50-74, doi:10.1080/07352680802053940.
- 630 39. Melnyk, C.W.; Meyerowitz, E.M. Plant grafting. *Curr Biol* **2015**, *25*, R183-188,
631 doi:10.1016/j.cub.2015.01.029.
- 632 40. Yin, H.; Yan, B.; Sun, J.; Jia, P.; Zhang, Z.; Yan, X.; Chai, J.; Ren, Z.; Zheng, G.; Liu, H.
633 Graft-union development: A delicate process that involves cell-cell communication
634 between scion and stock for local auxin accumulation. *Journal of Experimental Botany*
635 **2012**, *63*, 4219-4232, doi:10.1093/jxb/ers109.
- 636 41. Gainza, F.; Opazo, I.; Muñoz, C. Graft incompatibility in plants: Metabolic changes
637 during formation and establishment of the rootstock/scion union with emphasis on
638 Prunus species. *Chilean journal of agricultural research* **2015**, *75*, 28-34,
639 doi:10.4067/s0718-58392015000300004.
- 640 42. Nanda, A.K.; Melnyk, C.W. The role of plant hormones during grafting. *Journal of Plant*
641 *Research* **2018**, *131*, 49-58, doi:10.1007/s10265-017-0994-5.
- 642 43. Donner, T.J.; Sherr, I.; Scarpella, E. Regulation of preprocambial cell state acquisition
643 by auxin signaling in Arabidopsis leaves. **2014**, 10.1242/dev.037028,
644 doi:10.1242/dev.037028.
- 645 44. Wang, J.; Jiang, L.; Wu, R. Plant grafting: how genetic exchange promotes vascular
646 reconnection. *New Phytologist* **2017**, *214*, 56-65, doi:10.1111/nph.14383.
- 647 45. Mazur, E.; Benková, E.; Friml, J. Vascular cambium regeneration and vessel formation
648 in wounded inflorescence stems of Arabidopsis. *Nature Publishing Group* **2016**,
649 10.1038/srep33754, 1-15, doi:10.1038/srep33754.
- 650 46. Zafar, S.A.; Patil, S.B.; Uzair, M.; Fang, J.; Zhao, J.; Guo, T.; Yuan, S.; Uzair, M.; Luo, Q.;
651 Shi, J., et al. DEGENERATED PANICLE AND PARTIAL STERILITY 1 (DPS1) encodes a
652 cystathionine β -synthase domain containing protein required for anther cuticle and
653 panicle development in rice. *New Phytologist* **2019**, *1*, doi:10.1111/nph.16133.
- 654 47. Hussain, H.A.; Men, S.; Hussain, S.; Chen, Y.; Ali, S.; Zhang, S.; Zhang, K.; Li, Y.; Xu, Q.;
655 Liao, C., et al. Interactive effects of drought and heat stresses on morpho-physiological
656 attributes, yield, nutrient uptake and oxidative status in maize hybrids. *Sci Rep* **2019**,
657 *9*, 3890, doi:10.1038/s41598-019-40362-7.

- 658 48. Noor, R.S.; Wang, Z.; Umair, M.; Yaseen, M.; Ameen, M.; Rehman, S.-U.; Khan, M.U.;
659 Imran, M.; Ahmed, W.; Sun, Y. Interactive Effects of Grafting Techniques and Scion-
660 Rootstocks Combinations on Vegetative Growth, Yield and Quality of Cucumber
661 (*Cucumis sativus* L.). *Agronomy* **2019**, *9*, doi:10.3390/agronomy9060288.
- 662 49. Meng, J.; Wu, S.; Wang, X.; Yu, X.; Jiang, R. Effects of Different Rootstocks on Plant
663 Growth and Fruit Quality of Watermelon. *Agricultural Biotechnology* **2019**, *8* (1) : 64 -
664 68, 77 **2019**.
- 665 50. Ahmad, P.; Wani, M.R. Physiological mechanisms and adaptation strategies in plants
666 under changing environment: Volume 1. *Physiological Mechanisms and Adaptation*
667 *Strategies in Plants Under Changing Environment: Volume 1* **2014**, 10.1007/978-1-
668 4614-8591-9, 1-376, doi:10.1007/978-1-4614-8591-9.
- 669 51. Kohatsu, D.S.; Zucareli, V.; Brambilla, W.P.; Ono, E.O.; Doutores, P.; Cca, U.E.M.;
670 Agronomia, D.C.A.; Pr, U. Peroxidase and polyphenol oxidase activity on the yield of
671 grafted and ungrafted cucumber plants. *African Journal of Agricultural Research* **2013**,
672 *8*, 279-283, doi:10.5897/AJAR12.1560.
- 673 52. Adams, S.W. The Effects of Rootstock, Scion, Grafting Method and Plant Growth
674 Regulators on Flexural Strength and Hydraulic Resistance of Apple. *Thesis (master of*
675 *science)* **2016**, UTAH STATE UNIVERSITY Logan, Utah.
- 676 53. Errea, P.; Felipe, A.; Herrero, M. Graft establishment between compatible and
677 incompatible *Prunus* spp . **1994**, *45*, 393-401.
- 678 54. Dogra, K.; Kour, K.; Kumar, R.; Bakshi, P.; Kumar, V. Graft-Incompatibility in
679 Horticultural Crops Graft-Incompatibility in Horticultural Crops. **2018**,
680 10.20546/ijcmas.2018.702.218, doi:10.20546/ijcmas.2018.702.218.
- 681 55. Suying, T.; Jiwei, L.; Xiuqiang, H.; Sijun, L. Selection of New Watermelon Variety ‘
682 Mimei Tetraploid’ . *果树科学* **1993**, *5*.
- 683 56. Oda, I. New Grafting Methods for Fruit-Bearing Vegetables in Japan. **1995**, *194*, 187-
684 194.
- 685 57. Oda, M. Vegetable seedling grafting in japan. *Acta Horticulturae* **2007**, *759*, 175-180,
686 doi:10.17660/ActaHortic.2007.759.13.
- 687 58. Jawad, U.M.; Gao, L.; Gebremeskel, H.; Safdar, L.B.; Yuan, P.; Zhao, S.; Xuqiang, L.;
688 Nan, H.; Hongju, Z.; Liu, W. Expression pattern of sugars and organic acids regulatory
689 genes during watermelon fruit development. *Scientia Horticulturae* **2020**, *265*,
690 109102.
- 691 59. Zhu, Q.; Gao, P.; Liu, S.; Zhu, Z.; Amanullah, S.; Davis, A.R.; Luan, F. Comparative
692 transcriptome analysis of two contrasting watermelon genotypes during fruit
693 development and ripening. *BMC genomics* **2017**, *18*, 3.
- 694 60. Hu, B.; Jin, J.; Guo, A.-Y.; Zhang, H.; Luo, J.; Gao, G. GSDS 2.0: an upgraded gene
695 feature visualization server. *Bioinformatics* **2015**, *31*, 1296-1297.
- 696 61. Chi, Q.; Guo, L.; Ma, M.; Zhang, L.; Mao, H.; Wu, B.; Liu, X.; Ramirez-Gonzalez, R.H.;
697 Uauy, C.; Appels, R. Global transcriptome analysis uncovers the gene co-expression
698 regulation network and key genes involved in grain development of wheat (*Triticum*
699 *aestivum* L.). *Functional & Integrative Genomics* **2019**, *19*, 853-866.
- 700 62. Kariyanna, B.; Prabhuraj, A.; Asokan, R.; Babu, P.; Jalali, S.K.; Venkatesan, T.; Gracy,
701 R.G.; Mohan, M. Identification of suitable reference genes for normalization of RT-
702 qPCR data in eggplant fruit and shoot borer (*Leucinodes orbonalis* Guenée). *Biologia*
703 **2020**, *75*, 289-297.
- 704 63. Gao, L.; Zhao, S.; Lu, X.; He, N.; Zhu, H.; Dou, J.; Liu, W. Comparative transcriptome
705 analysis reveals key genes potentially related to soluble sugar and organic acid
706 accumulation in watermelon. *PloS one* **2018**, *13*, e0190096.
- 707 64. Melnyk, C.W. Plant grafting: insights into tissue regeneration. *Regeneration* **2017**, *4*, 3-
708 14, doi:10.1002/reg2.71.

709 65. BING SONG ZHENG; HUAI LIANG CHU; SONG HENG JIN; YOU JUN HUANG; ZHENG JIA
710 WANG; CHEN, M.; HUANG, J.Q. cDNA-AFLP analysis of gene expression in hickory
711 (*Carya cathayensis*) during graft process. *Tree Physiology* **30**, 297–303 **2009**, 30, 297–
712 303, doi:10.1093/treephys/tpp102.

713 66. Aloni, B.; Cohen, R.; Karni, L.; Aktas, H.; Edelstein, M. Hormonal signaling in rootstock–
714 scion interactions. *Scientia Horticulturae* **2010**, 127, 119-126,
715 doi:10.1016/j.scienta.2010.09.003.

716 67. Melnyk, C.W.; Schuster, C.; Leyser, O.; Meyerowitz, E.M. A Developmental Framework
717 for Graft Formation and Vascular Reconnection in *Arabidopsis thaliana*. *Curr Biol* **2015**,
718 25, 1306-1318, doi:10.1016/j.cub.2015.03.032.

719 68. Cookson, S.J.; Clemente Moreno, M.J.; Hevin, C.; Nyamba Mendome, L.Z.; Delrot, S.;
720 Trossat-Magnin, C.; Ollat, N. Graft union formation in grapevine induces
721 transcriptional changes related to cell wall modification, wounding, hormone
722 signalling, and secondary metabolism. *J Exp Bot* **2013**, 64, 2997-3008,
723 doi:10.1093/jxb/ert144.

724 69. Schaller, G.E.; Bishopp, A.; Kieber, J.J. The yin-yang of hormones: cytokinin and auxin
725 interactions in plant development. *Plant Cell* **2015**, 27, 44-63,
726 doi:10.1105/tpc.114.133595.

727 70. Tu, Y.; Jiang, A.; Gan, L.; Hossain, M.; Zhang, J.; Peng, B.; Xiong, Y.; Song, Z. Genome
728 duplication improves rice root resistance to salt stress. **2014**, 10.1186/s12284-014-
729 0015-4, 1-13, doi:10.1186/s12284-014-0015-4.

730 71. Ganie, S.A.; Molla, K.A.; Henry, R.J.; Bhat, K.V.; Mondal, T.K. Advances in
731 understanding salt tolerance in rice. *Theoretical and Applied Genetics* **2019**, 132, 851-
732 870, doi:10.1007/s00122-019-03301-8.

733 72. Ruiz, M.; Quiñones, A.; Martínez-Alcántara, B.; Aleza, P.; Morillon, R.; Navarro, L.;
734 Primo-Millo, E.; Martínez-Cuenca, M.-R. Effects of salinity on diploid (2x) and doubled
735 diploid (4x) *Citrus macrophylla* genotypes. *Scientia Horticulturae* **2016**, 207, 33-40,
736 doi:10.1016/j.scienta.2016.05.007.

737 73. Jiang, A.; Gan, L.; Tu, Y.; Ma, H.; Zhang, J.; Song, Z.; He, Y.; Cai, D.; Xue, X. The effect of
738 genome duplication on seed germination and seedling growth of rice under salt stress.
739 *Australian Journal of Crop Science* **2013**, 7, 1814.

740 74. Tu, Y.; Jiang, A.; Gan, L.; Hossain, M.; Zhang, J.; Peng, B.; Xiong, Y.; Song, Z.; Cai, D.; Xu,
741 W. Genome duplication improves rice root resistance to salt stress. *Rice* **2014**, 7, 1-13.

742 75. Doyle, J.J.; Coate, J.E. Polyploidy, the nucleotype, and novelty: The impact of genome
743 doubling on the biology of the cell. *International Journal of Plant Sciences* **2019**, 180,
744 1-52, doi:10.1086/700636.

745 76. Zhu, H.; Zhao, S.; Lu, X.; He, N.; Gao, L.; Dou, J.; Bie, Z.; Liu, W. Genome duplication
746 improves the resistance of watermelon root to salt stress. *Plant Physiol Biochem* **2018**,
747 133, 11-21, doi:10.1016/j.plaphy.2018.10.019.

748 77. Yi Tu; Aiming Jiang; Lu Gan; Mokter Hossain; Jinming Zhang; Bo Peng; Yuguo Xiong;
749 Zhaojian Song; Detian Cai; Weifeng Xu, et al. Genome duplication improves rice root
750 resistance to salt stress. *Rice* **2014**, 7:15.

751 78. Meng, F.; Luo, Q.; Wang, Q.; Zhang, X.; Qi, Z.; Xu, F.; Lei, X.; Cao, Y.; Chow, W.S.; Sun,
752 G. Physiological and proteomic responses to salt stress in chloroplasts of diploid and
753 tetraploid black locust (*Robinia pseudoacacia* L.). *Scientific reports* **2016**, 6, 23098.

754 79. Yan, K.; Wu, C.; Zhang, L.; Chen, X. Contrasting photosynthesis and photoinhibition in
755 tetraploid and its autodiploid honeysuckle (*Lonicera japonica* Thunb.) under salt
756 stress. *Frontiers in Plant Science* **2015**, 6, 227.

757 80. Balal, R.M.; Shahid, M.A.; Vincent, C.; Zotarelli, L.; Liu, G.; Mattson, N.S.;
758 Rathinasabapathi, B.; Martínez-Nicolas, J.J.; Garcia-Sanchez, F. Kinnow mandarin
759 plants grafted on tetraploid rootstocks are more tolerant to Cr-toxicity than those

- 760 grafted on its diploids one. *Environmental and Experimental Botany* **2017**, 140, 8-18,
 761 doi:10.1016/j.envexpbot.2017.05.011.
- 762 81. Said, J.I.; Lin, Z.; Zhang, X.; Song, M.; Zhang, J. A comprehensive meta QTL analysis for
 763 fiber quality, yield, yield related and morphological traits, drought tolerance, and
 764 disease resistance in tetraploid cotton. *BMC Genomics* **2013**, 14, 776,
 765 doi:10.1186/1471-2164-14-776.
- 766 82. Allario, T.; Brumos, J.; Colmenero-Flores, J.M.; Iglesias, D.J.; Pina, J.A.; Navarro, L.;
 767 Talon, M.; Ollitrault, P.; Morillon, R. Tetraploid Rangpur lime rootstock increases
 768 drought tolerance via enhanced constitutive root abscisic acid production. *Plant Cell*
 769 *Environ* **2013**, 36, 856-868, doi:10.1111/pce.12021.
- 770 83. Fernandez-Garcia, N.; Carvajal, M.; Olmos, E. Graft union formation in tomato plants:
 771 peroxidase and catalase involvement. *Ann Bot* **2004**, 93, 53-60,
 772 doi:10.1093/aob/mch014.
- 773 84. North, J.J. Effects of antioxidants, plant growth regulators and wounding on phenolic
 774 compound excretion during micropropagation of *Strelitzia reginae*. *International*
 775 *Journal of the Physical Sciences* **2012**, 7, 638-646, doi:10.5897/ijps11.786.
- 776 85. Xu, Q.; Guo, S.-R.; Li, H.; Sun, J.; Du, N.-S.; Shu, S. Physiological Aspects of
 777 Compatibility and Incompatibility in Grafted Cucumber Seedlings. *J.*
 778 *AMER.SOC.HORT.SCI.* 140(4):299–307 **2015**.
- 779 86. Silva, E.S.; Menezes, D.V.; Silva, E.G.; Goto, R.; Lima, G.P.P. Different methods of
 780 grafting and activity of antioxidant enzymes in tomato. *Revista Brasileira de Ciências*
 781 *Agrárias - Brazilian Journal of Agricultural Sciences* **2016**, 11, 267-271,
 782 doi:10.5039/agraria.v11i4a5392.

783