

## Climatic trend in different bioclimatic zones in Chitwan Annapurna Landscape, Nepal

**Dol Raj Luitel<sup>1,2</sup>, Pramod K. Jha<sup>1</sup>, Mohan Siwakoti<sup>1</sup>, Madan Lall Shrestha<sup>3</sup>, and Rangaswamy Munniappan<sup>4</sup>**

<sup>1</sup>Central Department of Botany, Tribhuvan University, Kritipur, Kathmandu, Nepal

<sup>2</sup>Departments of Environment, Ministry of Forests and Environment, Kathmandu, Nepal

<sup>3</sup>Nepal Academy of Science and Technology, Lalitpur, Nepal

<sup>4</sup>IPM-IL Virginia Tech, Blacksburg, USA

Corresponding email: [luiteldr2@gmail.com](mailto:luiteldr2@gmail.com); [pkjhaprof@gmail.com](mailto:pkjhaprof@gmail.com)

### Abstract

Depending upon altitudinal gradient in the Himalayas, the rate of climate change varies from lowland to upland. The Chitwan Annapurna Landscape (CHAL) is the central part of the Himalayas and covers all bioclimatic zones. Analysis of time series data (1970-2019) of temperature and precipitation was carried out in seven bioclimatic zones extending from lowland Terai to higher Himalayas. The non-parametric Mann-Kendall test was applied to determine the trend, which was quantified by Sen's slope. Annual and decade interval average temperature, precipitation trends, and lapse rate were analyzed in each bioclimatic zone. Out of seven bioclimatic zones, four zones showed a decreasing precipitation trend (lower tropical, upper tropical, upper subtropical, and alpine bioclimatic zones) at the rate of 1.8, 1.98, 2.06, and 1.80 mm/year, and in lower sub-tropical, temperate, and lower subalpine bioclimatic zones, increasing at the rate of 0.45, 1.81 and 1.28mm/year, respectively. Precipitation did not show any particular trend at decade intervals. The average annual temperature at different bioclimatic zones clearly indicates that temperature at higher elevations is significantly increasing more than at lower elevations. In lower tropical bioclimatic zone (LTBZ), upper tropical bioclimatic zone(UTBZ), lower subtropical bioclimatic zone (LSBZ), upper subtropical bioclimatic zone(USBZ), and temperate bioclimatic zone(TBZ), the average temperature increased by 0.022, 0.030, 0.036, 0.042 and 0.051°C/year, respectively. The decade level temperature scenario revealed that the hottest decade was from 1999-2009. The average temperature was found as 24.1, 21.8, 19.7, 17.5, and 13.3°C in LTBZ, UTBZ, LSBZ, USBZ, and TBZ, respectively, and the average annual precipitation in LTBZ, UTBZ, LSBZ, USBZ, TBZ, LBZ, and ABZ was 2002.1, 2613.1, 2223.9, 3146.9, 1447.2, 952.1, and 361.7mm/year, respectively, in CHAL. With the impact of climate

change site and region-specific, this information highlights the need to mitigate climate change in different bioclimatic zones.

**Key words:** bioclimatic zones, climate change, precipitation, temperature, trend

## 1. Introduction

Elevation-dependent warming is higher in the Himalayas than any other part of the world[1,2].Globally, the average yearly temperature increases by  $0.02^{\circ}\text{C}$  [3], but the average maximum temperature of Nepal increases yearly by  $0.056^{\circ}\text{C}$  [2],varying in different areas and bioclimatic zones[4]. This indicates that the impact of climate change is severe and more rapid in both the upland and lowland area of Nepal [5], which could lead to several unpredictable problems for livelihoods [6], ecosystem processes, biodiversity, and natural resources in general.

The Chitwan-Annapurna Landscape (CHAL) area in Central Nepal extends from 164m to 8091m altitudes within a short north-south distance of about 185km. It covers  $32,090\text{ km}^2$  land area, encompassing 19 political districts that cover 22% of land in Nepal. It is known that altitude and aspect along the latitude create a wide range of climatic conditions and that CHAL also encounters almost all types of bioclimates – from tropical to alpine – which makes it an appropriate landscape for climatic study. The climate change analysis and trend information would be useful for biodiversity analysis, biological invasion, natural disaster, and vulnerability and impact studies.

Since 1993, numerous analyses of climatic trends at national and regional levels[7,8,9,10, 11,12, 13, 14,15, 2, 16,17,18, 9,20] have been conducted in Nepal. Most of these studies have conducted thorough analyses of selected meteorological stations of Nepal and adjoining regions. However, systematic climatic trends from a complete set of meteorological stations at different bioclimatic zones of a landscape have not been studied. Due to topographic variation and orographic effects, the average temperature and precipitation assessment is a challenge in a region like CHAL. Simply averaging the data from stations at different altitudes cannot distinguish variations in trends among bioclimatic zones.

This study fulfils a gap in knowledge of the rate of warming in different elevation levels within CHAL by i) analyzing the trends of temperature and precipitation at different bioclimatic zones,


ii) examining decade interval climate change patterns in temperature and precipitation at different bioclimatic zones, and iii) determining the average temperature, precipitation, and lapse rate of temperature and precipitation in different bioclimatic zones of the area.

## 2. Materials and Methods

### 2.1 Study area

The study area is Chitwan Annapurna Landscape (CHAL), located in central Nepal between  $27^{\circ}35'$  and  $29^{\circ}33'$  N latitude and  $82^{\circ}88'$  and  $85^{\circ} 80'$  E longitude [21]. The CHAL includes all or part of 19 districts (*Mustang, Manang, Gorkha, Rasuwa, Nuwakot, Dhading, Lamjung, Tanahu, Syangja, Kaski, Palpa, Parbat, Baglung, Myagdi, Gulmi, Arghakhachi, Makwanpur, Chitwan and Nawalparasi*) (Figure-1). Based on altitude, topography, and climate, Nepal is generally divided into Terai (59-200 m), Siwaliks (200-1500 m), middle mountains (1000-2500 m), high mountains (2200-4000m), and high Himalayas ( $>4000$  ml). In this study, CHAL is categorized into seven bioclimatic zones [22]and[23]with slight modification to evaluate the climatic trends as lower tropical bioclimatic zone ( $<500$ m) (LTBZ), upper tropical bioclimatic zone (500-1000m) (UTBZ), lower subtropical bioclimatic zone (1000-1500m) (LSBZ), upper subtropical bioclimatic zone (1500-2000m) (USBZ), temperate bioclimatic zone (TBZ), lower sub-alpine bioclimatic zone (3000-3500m) (LABZ), and upper sub-alpine bioclimatic zone ( $>3500$ m asl) (ABZ).

A wide range of climatic conditions exist in CHAL, including the tropical humid climate in lowland of Chitwan and Nawalparasi districts, the alpine in the high mountains, and the cold dry climate in the trans-Himalaya in parts of Mustang and Manang districts. The mean temperature of Terai/Siwalik is more than  $25^{\circ}\text{C}$ , about  $20^{\circ}\text{C}$  in the middle hills, and in between  $10^{\circ}\text{C}$  to  $20^{\circ}\text{C}$  in the high mountains [24]. The average annual rainfall ranges from 165mm at Lomanthang (Mustang) in the northern part to 5,244mm at Lumle (highest rainfall in the country) in mid-hills [2]. Orographic effects cause high spatial variation in precipitation in different zones of the landscape. Nearly 78%of the total annual precipitation occurs during the monsoon season between June and September [16]. Occasional winter rainfall is common including short rainfall in the Siwalik and mid-hills and snowfall is common in high altitude regions.



**Figure 1** Map of Nepal with Chitwan Annapurna Landscape (CHAL) area


## 2.2 Climate analysis

Both precipitation and temperature data for all stations in CHAL were procured from the Department of Hydrology and Meteorology (DHM), Government of Nepal. There are 81 and 32 weather stations to measure rainfall and temperature within CHAL, respectively (Table-1). However, time series data (1970-2019) of daily temperature and rainfall of 26 and 52 stations, respectively, were only used in this analysis due to presence of complete data set of different bioclimatic zones in CHAL (Table-1). The distribution scenarios of functional stations used to depict inferences along different bioclimatic zones are presented in Figs2 and 3. The station details, including the data available date, are presented in Annex 1 and 2.

The rainfall and temperature trends were calculated in spatial (seven bioclimatic zones based upon altitude, climate, and vegetation) and temporal (annual and decade) scales.

**Table 1** Total number of precipitation and temperature stations in each bioclimatic zone in CHAL. Numbers in parentheses indicate stations having complete data sets.

Bioclimatic zones	Precipitation stations	Temperature stations
LTBZ(<500m)	14(11)	6(5)
UTBZ (500-1000m)	16(11)	10(8)
LSBZ(1000-1500m)	16(10)	4(4)
USBZ (1500-2000m)	14(8)	5(4)
TBZ (2000-3000m)	10(7)	7(5)
LABZ (3000-3500m)	3(2)	-
ABZ (>3500m)	8(3)	-
<b>Total</b>	<b>81(52)</b>	<b>32(26)</b>



**Figure 2** Stations recording precipitation at different bioclimatic zones in CHAL


The average temperature of individual stations averaged and average sum of precipitation from all stations in each bioclimatic zone was calculated.

The temperature was analyzed only from LTBZ to TBZ in this study. Although two climatological stations, Jomsom and Thakmarpha (station Id 601 and 604), lie in between 2500

and 3000m altitudes in Mustang district of CHAL, the data of these stations cannot represent temperature of all bioclimatic zones in upper temperate regions. Mustang lies within the Trans-Himalayan region i.e. rain shadow area. Therefore, stations of lower temperate and upper temperate bioclimatic zones were merged and averaged to calculate the trend.

### 2.3 Data management

The stations level data were manually checked to determine the odd and unusual patterns. The station data that showed outliers and unusual values were corrected for individual stations by replacing them with the average of the value from the previous day and the next day [2]. If there is an outlier or missing data for about a month or more, that particular year is omitted from the analysis.



**Figure 3** Temperature stations at different bioclimatic zones in CHAL

## 2.4 Trend Analysis

Mann-Kendall test and Sen's slope methods were used to analyze the climatic trend, magnitude, significance of temperature, and precipitation data in bioclimatic zones. The Mann-Kendall test is a non-parametric tool and one of the best methods to analyze the presence and significance of monotonic positive or negative trend in time series climatic data [25, 26, 2]. Existence of positive or negative trend among the considered climatic variables was determined by using Mann-Kendall trend test, its quantification was done by using Sen's slope method, and significance by Mann-Kendall Methods in R package version 3.4.4 [27]. The Mann-Kendall method of significance test uses the hypothesis testing approach. In the testing mechanism null hypothesis ( $H_0$ ) there is no monotonic trend in climatic data, and with alternative hypothesis ( $H_1$ ), there is a monotonic trend in climatic data at significant level. Significant tests at 0.05 confidence levels were used.

## 2.5 Lapse rate of temperature and precipitation

The lapse rate of temperature and precipitation in different bioclimatic zones along altitudinal gradient in CHAL were calculated following [28] and [29] with slight modification. The lapse rate of precipitation was derived by analyzing annual average precipitation sums of the stations lying in each bioclimatic zone by using the following equation:

$$PPtLR = \frac{P_1 - P_2}{Z_1 - Z_2}$$

Where PPt LR is precipitation lapse rate,  $P_1$  and  $P_2$  are the precipitation sums of the highest and lowest bioclimatic zones (in mm) and  $Z_1$  and  $Z_2$  are their respective elevations of higher and lower bioclimatic zones. The value of  $Z_1-Z_2$  considered 1 because our calculation is converted into 1 kilometer along elevation change.

All temperature data were aggregated to yearly values for the period of 50 years. Temperature lapse rates were calculated through temperature difference between two successive bioclimatic zones i.e. temperature-elevation space [28]. Average temperature is normally assumed to decrease linearly with increasing elevation in CHAL. So, Temp LR was calculated as:

$$TempLR = \frac{T_1 - T_2}{Z_1 - Z_2}$$


Where Temp LR is Temperature lapse rate,  $T_1$  and  $T_2$  are the yearly average temperatures of the highest and lowest temperature in each bioclimatic zone ( $^{\circ}\text{C}$ ), and  $Z_1$  and  $Z_2$  are their average elevations (m). We calculated the lapse rates based on the strong relationship between ambient air temperature and elevation [30, 28, 29].

### 3. Result

#### 3.1 Annual trend of precipitation

A mixed pattern of precipitation in different bioclimatic zones was noted in CHAL. Out of seven bioclimatic zones, four zones showed decreasing precipitation and three zones showed increasing trend (Table-2). Except UTBZ and LSBZ, the trend was statistically significant in these bioclimatic zones. At LTBZ ( $<500\text{m}$ ), the average precipitation significantly decreased at the rate of 1.8 mm/year. In UTBZ, the precipitation decreased at the rate of 1.98mm/year. Precipitation in USBZ had increasing yearly of 0.45mm, but at USBZ a decreasing rate of 2.06mm/year (Table-3 and Figure-4).

Above from 2000m elevation, the precipitation showed increasing trend up to 3500m and then again decreasing. At TBZ (2000-3000m), the precipitation increased at the rate of 1.81mm/year and at LABZ (3000-3500m) at the rate of 1.28mm/year. At ABZ ( $>3500\text{m}$ ), precipitation had a decreasing trend at the rate of 1.80mm/year (Table-3 and Figure-4).



**Figure 4**Average annual rainfall trends in different bioclimatic zones in CHAL

### 3.2 Precipitation trend in different decade


Average decade rainfall in bioclimatic zones in CHAL was analyzed. In general, precipitation in all bioclimatic zones increased until 2000AD, but decreased after 2000AD. At UTBZ, average rainfall increased from 1970-1979 to 1980-1990 and again showed decreasing and increasing pattern. In the last decade (2010-2019), the average rainfall was found decreasing from preceding decades. From Sen's slope value, the rainfall was found decreasing at the rate of 19.8mm per decade.

USBZ received maximum average rainfall in CHAL (Figure-5); however, overall trend shows that average precipitation was decreasing at the rate of 20.6mm/decade (Tabel-2).

In TBZ, from the first (1970-1979) to the second decade (1980-1989), the rainfall trend was found increasing and decreasing thereafter. However, overall decade level trend analysis showed increasing rainfall at the rate of 18.1mm/decade. Similarly, particular trend was not seen in LABZ, but average decade rainfall was found increasing at the rate of 12.8mm/decade.

**Table 2** Decade wise average precipitation (mm) trend in different bioclimatic zones in CHAL

Bioclimatic zones	1970-1979	1980-1989	1990-1999	2000-2009	2010-2019	trend between 1970-2019		
						Sen's slope per year	P value (mann'sKendall test)	Significance
LTBZ	1964.4	2030.5	2022.3	2169.3	1823.6	-1.8	<b>0.029</b>	No
UTBZ	2620.8	2604.6	2665.7	2677.8	2496.3	-1.98	0.47	No
LSBZ	2145.8	2216.0	2362.1	2242.8	2152.8	0.45	0.89	No
USBZ	3222.5	3422.9	3066.9	3105.5	2913.79	-2.06	0.03	Yes
TBZ	1230.2	1558.1	1519.65	1498.75	1429.25	1.81	0.002	Yes
LABZ	886.2	854.8	992.1	971.1	1056.2	1.28	0.0001	Yes
ABZ	366.8	444.3	351.1	298.4	348.0	-1.80	0.05	Yes

**Figure 5** Decade-wise precipitation trend along different bioclimatic zones in CHAL

### 3.3. Annual trend of temperature


**Table 3** Average temperature trend (1970-2019) in different bioclimatic zones in CHAL

Bioclimatic zones	Sen's slope	P value (Mann's Kendel test)	Significance
LTBZ	0.022	<0.001	Yes
UTBZ	0.030	<0.00001	Yes
LSBZ	0.036	<0.0001	Yes
USBZ	0.042	<0.00001	Yes
TBZ	0.051	<0.0001	Yes

The average temperature increased by 0.022°C/year during 1970 to 2019 in LTBZ. However, in UTBZ, LSBZ, USBZ, and TBZ, the average annual temperature increased by 0.030, 0.036, 0.042, and 0.051°C/year, respectively. This result clearly indicated that temperature at higher elevation has been significantly increasing more than at lower elevations (Table-3 and Figure-6). Although average temperature increased in all bioclimatic zones, the increasing rate of average annual temperature at higher elevation was high.

### 3.4. Temperature in different decades

Average temperature in every bioclimatic zone of CHAL has been increasing over the past five decades (Table-4). On average, at a rate of 0.2°C/decade, temperature significantly increased at LTBZ of CHAL. At LTBZ, from 1989-1999, the highest average temperature increase was 0.27°C/decade. From 1979-1989, the average temperature rose to 0.26°C/decade, but the last two decades experienced slightly low temperature gain (Table-4). Similarly, 1999-2009 was a comparatively hotter decade in higher altitudes above 1000m (Table-4).



**Figure 6** Average temperature trends in different bioclimatic zones of CHAL (1970-2019)

**Table 4** Increased average temperature (°C) per decade in different bioclimatic zones of CHAL since 1979

Bioclimatic zones	1979-1989	1989-1999	1999-2009	2009-2019	1979-2019	
					Sen's slope	Significance
<b>LTBZ</b>	0.26	0.27	0.16	0.19	0.022	Yes
<b>UTBZ</b>	0.28	0.28	0.33	0.32	0.030	Yes
<b>LSBZ</b>	0.17	0.39	0.44	0.42	0.036	Yes
<b>USBZ</b>	0.36	0.39	0.44	0.47	0.042	Yes
<b>TBZ</b>	0.48	0.51	0.56	0.52	0.051	Yes

### 3.5 Average annual temperature and precipitation

The average temperature and precipitation values of all stations in each bioclimatic zone have been analyzed for 1970-2019 (Table-5). The results show 24.1, 21.8, 19.7, 17.5, and 13.3°C average annual temperature at LTBZ, UTBZ, LSBZ, USBZ, and TBZ, respectively, in CHAL.

Similarly, average annual precipitation in LTBZ, UTBZ, LSBZ, USBZ, TBZ, LBZ, and ABZ was 2002.1, 2613.1, 2223.9, 3146.9, 1447.2, 952.1, and 361.7mm/year, respectively, in CHAL. This result indicates that with increasing altitude from LTBZ to UTBZ, annual average precipitation increases, but in other BZ the precipitation successively decreases (Table-5).

**Table 5** Lapse rate and average temperature and precipitation along different bioclimatic zones in CHAL

Bioclimatic zones	AAT (°C)	TLR (C/500m)	TLR (°C/km)	ANR (mm)	PLR (mm/500m)	PLR (mm/km)
Lower tropical bioclimatic zone(<500m)	24.1		-	2002.1		-
Upper tropical bioclimatic zone(500-1000m)	21.8	-2.3	-4.6	2613.1	611	1222
Lower subtropical bioclimatic zone(1000-1500m)	19.7	-2.1	-4.2	2223.9	-389.2	-778.4
Upper subtropical bioclimatic zone(1500-2000m)	17.5	-2.2	-4.4	3146.4	922.5	1845
Temperate bio-climatic zone(2000-3000m)	13.3	-2.2	-4.4	1447.2	-849.6	-1699.2
Lower subalpine bioclimatic zone(3000-3500m)				952.1	-495.1	-990.2
Alpine bioclimatic zone(>3500m)				361.7	-590.4	-1180.8

**Note:** AAT= Average annual temperature, TLR= Temperature lapse rate, PLR= precipitation lapse rate

### 3.6 Lapse rate of temperature and precipitation lapse rate

The temperature lapse rate showed 2.1 to 2.3°C decreases with every increase of 500m altitude along different bioclimatic zones in CHAL (Table-5). It implies that every one-kilometer that altitude rises, the temperature decreases by 4.2 to 4.6°C/km in CHAL.

#### 4. Discussion

A mixed pattern of precipitation in different bioclimatic zones is in congruence with the precipitation pattern for the whole of Nepal [31]. Previous work in different parts of CHAL were localized to districts or whole regions at small scale analysis based on some meteorological stations of Lamjung[32], Syangja [33], and other districts [2]. According to Poudel and Shaw[32], annual precipitation was increasing in Khudi (855m) and Kunchha (823m) stations of Lamjung district at the rate of 0.063mm/year and 4.42mm/year, respectively. However, in Gharedhunga (1120m) station, in the same district over the course of 32 years (1980-2012), there was a decreasing trend at the rate of 3.48mm/year within CHAL. Our findings in general conclude that for the whole CHAL, annual average precipitation varies because of precipitation amount varying in different years.

According to DHM [2], normal annual precipitation at district level in CHAL showed that Mustang, a trans-Himalaya region, receives the lowest rainfall (<4000mm), whereas Kaski, Parbat, Tanahu, Lamjung, and Nuwakot receive more than 2000mm of annual rain. Similarly, High Himalayas receives (>4000m) the least amount of rain and the remaining regions receive 1500-2000mm, which is consistent with present results. The annual precipitation trend of Nepal along different physiographic regions showed inconsistency. According to [2]in Terai, annual average rainfall was found increasing at the rate of 0.49mm/year, but Siwalik, Mid-mountain, High mountains, and High Himalayas showed decreasing at the rate of 1.48mm, 1.58mm, 3.17mm, and 1.46mm/year, respectively. For the whole of Nepal, total annual precipitation since the 1960s has been decreasing at the rate of 3.7 mm per year [34]. District wise, annual precipitation trend in CHAL showed a different pattern. Among 19 districts, annual average rainfall significantly decreased in 10 districts and decreased in nine districts [2], which is consistent with the present study. There is no previous comparison at the bioclimatic zone level in CHAL.

In Nepal, average annual rainfall is 1858.6 mm [35]; however, the spatial pattern of annual rainfall in the country depends on topography. Altitude further affects rainfall pattern; total annual rainfall increases with altitude up to approximately 3,000m above sea level and then diminishes at higher elevations [36]. However, in CHAL, average annual rainfall was found inconsistently increasing and decreasing from lower to higher altitudes because of mountains

blocking the monsoon wind. Seasonal analysis of station-based rainfall pattern in Gandaki river basin (major part of CHAL) revealed significantly decreasing winter rainfall but increasing monsoon rainfall [16]. The precipitation was found significantly increasing in the high mountains of CHAL.

Precipitation in each bioclimatic zone in the mountains depends upon many factors such as topography, strength of moisture-bearing wind, and the orientation of the mountain range with respect to the prevailing wind direction. Thus, precipitation processes and distribution in a region is influenced by aforementioned factors, including steep altitudinal contrast [37]. Variation in elevation within a very short distance of about 185km creates dissimilarity in precipitation within particular regions in Nepal [31, 38]. In this study, we found that the rainfall in LTBZ and UTBZ is decreasing annually but in LSBZ it is increasing, and again significantly decreasing in USBZ. This unusual pattern of precipitation along different bioclimatic zones of CHAL is therefore due to orography and the spatial arrangement of topographic gradients, which control the precipitation patterns [37]. These spatial arrangements of topographic gradients, wind direction, aspects, and slopes of mountain may alter consistency in precipitation along different ecological zones within a short distance in CHAL.

In the higher elevation above 2000m, annual precipitation was found to be much lower than precipitation in lower elevations. Increases in annual precipitation from LTBZ to higher elevation up to 2000m at USBZ, and decreases in precipitation above TBZ, was well noted. The highest annual rainfall was found in USBZ in CHAL. The Lumle station in CHAL received the highest amount of annual precipitation in Nepal of about 5500mm [39, 35] as it lies on the windward side of the Annapurna-Machhapuchhre mountain range. The lowest precipitation site was recorded in Upper Mustang Dhiee, Lomanthang area of Mustang district within CHAL, with a mean annual precipitation of less than 200mm. Both of these highest and lowest precipitation sites of the country are in the Annapurna area of present study site.

According to [15], there are two high monsoonal rainfall zones in Nepal, one around 600m and another around 2100m, with different rain patterns. This record of gradual increasing precipitation from LTBZ to USBZ, with highest at around 2000m altitude, may be due to presence of high monsoonal rainfall zones. From the comprehensive precipitation observations in the Annapurna range, [40] showed that the annual precipitation gradually increases from lowland and had a strong peak at about 3000m altitude, then decreases as elevation increases. However,

present analysis showed that strong peak of precipitation at 2000m then decreases as elevation increases in CHAL, which is inconsistent with the result of [40].

According to the precipitation trend analyzed from data of 80 stations in Nepal, most of the Terai area and Western Nepal observe a negative trend [41]. The hills and mountains of Western Nepal and the northern part of Eastern Nepal have a positive trend. Based on data from 1947 to 1993, [42] found that the precipitation trend in the Koshi Basin (Eastern Nepal) showed an increasing trend. The overall average trend for Nepal indicates that the annual average precipitation is decreasing at the rate of 9.8mm/decade [5]. Our findings on decade level precipitation trend in different bioclimatic zones of CHAL showed consistency with previous findings from Nepal.

According to [43], global mean surface temperature has increased on average by 0.8°C in the 20<sup>th</sup> century and by 0.6°C from 1975 to 2005. [9] reported the maximum temperature increased in mountains of Nepal during 1971-1994 by 0.06 to 0.12°C/year, while the southern plains increased by less than 0.03°C/year, and during 1970-2015 by the maximum temperature increased by 0.056°C/year in Nepal[2]. In the CHAL, rise in average temperature was found considerably higher than the global average.

At the district level, the highest significant positive trend was observed in Manang district (0.12°C/yr) [2]. Average annual temperature increased in Terai (<200m), Siwalik (200-1500m), Middle mountain (1500-2500m), High mountain (2200-4000m), and High Himalayas at the rate of 0.020, 0.023, 0.031, 0.032, and 0.035°C/year, respectively, which is a similar pattern with the present study. The temperature increase was noted higher in our study than DHM (2017). This speedy warming trend in high elevation zones compared to lower altitudes is due to melting of snow and ice [44] and cold air pooling along local heating by combination of topography and synoptic conditions [45].

In Nepal, average temperature and precipitation analysis for different bioclimatic zones in particular areas has not been assessed before. However, average temperature and precipitation analysis along different physiographic regions had been done. The range of temperature in Terai and Siwalik, mid-hills, and mountain physiographic regions is 20-25, 15-20, 10-20, <3-10°C, respectively [24, 36]. [20] analyzed the variation in temperature by increasing every two hundred meters altitudes and found that temperature at <200m was 24.8°C. The temperature at <3800m was 7.8°C without any consistency pattern of increasing temperature in between lowest to higher elevation.

Thus, these reasonable elevation-dependent temperature and precipitation trends vary due to elevation-based differential changes in climate drivers, such as snow/ice cover, clouds, water vapor, aerosols, and soil moisture, or differential sensitivities of surface warming due to changes in these drivers at different elevations. However, mountain systems are inherently difficult to understand as a result of their complex topography, which leads to a high level of spatial and temporal variability in their climatic responses [4].

According to [29], the annual average TLR on the southern slope of the central Himalayas is  $-5.2^{\circ}\text{C}/\text{km}$ , which is  $-0.4$  to  $0.8^{\circ}\text{C}/\text{km}$  higher than lapse rate at different bioclimatic zones of CHAL. The lapse rate of temperature depends on surface air temperature on elevation and varies due to different seasonality, interplay of radiation, slope and aspect of the mountains, and several local factors [46].

However, the precipitation lapse rate shows irregular patterns in different bioclimatic zones in CHAL. When altitudes rise from LTBZ to UTBZ, PLR shows positive trends of 611mm/500m elevation i.e. 1222mm/km. However, moving upwards from UTBZ to LSBZ, the precipitation lapse rate (PLR) shows negative rate of 778.4mm/km, and then again shows the rate of 1845mm/km up to 3000m. Then, PLR continuously decreases as altitude increases. The decreasing rate is higher at uppermost elevation (Table-5). This unusual pattern of increasing and decreasing PLR is consistent with [47] in Langtang valley within CHAL. The difference in PLR between different bioclimatic zones may be due to the local effects of topography, slope, and aspects of mountain as well as the direction of wind and many other local factors.

## 5. Conclusion

The result of the present study contributes to understanding temperature and precipitation trends in different bioclimatic zones in Nepal. There was considerable variation in precipitation and a mixed pattern from lower tropical to alpine bioclimatic zones. The average precipitation is significantly decreasing at the rate of  $-2.06$  mm/year in USBZ, but increasing at the rate of 1.81mm/yr in TBZ. A similar pattern of precipitation was observed in the last five decades, except LABZ and ABZ in all bioclimatic zones. The temperature at different bioclimatic zones of CHAL at higher elevation was significantly increasing, more so with elevational rise i.e. 0.022 to  $0.051^{\circ}\text{C}$ .

The average temperature was 24.1, 21.8, 19.7, 17.5, and  $13.6^{\circ}\text{C}$  at LTBZ, UTBZ, LSBZ, USBZ and TBZ, respectively, in CHAL. It indicates  $2.1$ - $2.3^{\circ}\text{C}$  temperature lapse rate is decreasing with

increasing altitude by 500m in CHAL. Similarly, average annual precipitation in LTBZ, UTBZ, LSBZ, USBZ, TBZ, LBZ, and ABZ was 2002.1, 2613.1, 2223.9, 3146.9, 1447.2, 952.1, and 361.7mm/year, respectively. These findings would be useful for the planning of mitigation and adaptation strategies for climate change impacts.

### **Author Contributions:**

DRL generated idea and designed study, data collection, management, analysis and final manuscript write up, PKJ, MS and MLS supervised and corrected the manuscript and RM arranged the financial details and final english editing. Finally all authors read and approve the final version of the manuscript.

**Funding:** This research was fully funded by the USAID Bureau of food security under the cooperative agreement No AID-OAA-L-15-00001.

### **Acknowledgements:**

This work was funded in whole by the United States Agency for International Development (USAID) Bureau of Food Security under the Cooperative Agreement No. AID-OAA-L-15-00001 as a part of Feed the Future Innovation Lab for Integrated Pest Management. Thank to Prof. Dr. Ram Kailash Prasad Yadav, Head, Central Department of Botany, TU for encouragement. We especially thank to Mr. Hari Prasad Pandey, Undersecretary, Ministry of Forests and Environment for guiding Mann Kendall test and Sen's slope determination in R package. Thanks to Department of Hydrology and Meteorology, Government of Nepal, for providing climatic data. Thanks to Ms Sara Hendery for language editing.

### **Conflicts of Interest:**

The authors declare no conflict of interest.

### **References**

1. Palazzi, E.;Filippi, L.; Hardenberg, J.V. Insights into elevation dependent warming in the Tibetan plateau-Himalayas from CMIP5 model simulations. *ClimDyn.* 2016, DOI 10.1007/s00382-016-3316-z.
2. DHM. *Observed Climate Trend Analysis in the Districts and Physiographic Regions of Nepal (1971-2014)*. Department of Hydrology and Meteorology, Kathmandu, Nepal, 2017.

3. IPCC. *Summary for Policymakers*. In: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)], Cambridge University Press: Cambridge, UK, 2018.
4. Rangwala, I.; Miller, J. R. Climate change in mountains - a review of elevation dependent warming and its possible causes. *Climatic Change* 2012, 114, 527–547. DOI 10.1007/s10584-012-0419-3.
5. MoFE. *Climate change scenarios for Nepal for National Adaptation Plan (NAP)*. Ministry of Forests and Environment, Kathmandu, 2019.
6. Parmesan, C.; Yohe, G. A. globally coherent fingerprint of climate change impacts across natural systems. *Nature*. 2003, 42(1), 37-42.
7. Shrestha, M.L. Development of climate change scenarios with reference to Nepal. *Proceedings of the workshop on climate change in Nepal*. Department of Hydrology and Meteorology, Kathmandu. 1997, 16-32.
8. DHM. *Proceedings of the workshop on climate change in Nepal*. Department of Hydrology and Meteorology, Kathmandu. 1997.
9. Shrestha, A. B.; Wake, C. P.; Mayewski, P. A.; Dibb, J. E. Maximum temperature trends in the Himalaya and its vicinity: An analysis based on temperature records from Nepal for the period 1971-94. *Journal of Climate*. 1999, 12(9), 2775-2786.
10. Baidya, S.K.; Regmi, R.K.; Shrestha, M.L. Climate profile and observed climate change and climate variability in Nepal. APN CAPABLE Report. 2007.
11. Karmacharya, J.; Shrestha, A.; Shrestha, M. L. *Climate change scenarios for South Asia and central Himalayan region based on GCM Ensemble*. APN CAPABLE Report. 2007.
12. Baidya, S.K.; Shrestha, M.L.; Shiekh, M.M. Trends in daily extremes of temperature and precipitation in Nepal. *Journal of Hydrology and Meteorology*. 2008, 5(1), 38-51.
13. NCVST. *Vulnerability through the eyes of vulnerable: climate change induced uncertainties' and Nepal's development predicaments*. Institute for Social and

Environmental Transition-Nepal (ISET-N, Kathmandu) and Institute for Social and Environmental Transition (ISET, Boulder, Colorado) for Nepal Climate Vulnerability Study Team, Kathmandu, Nepal.2009.

14. Practical Action. *Understanding disaster management in practice with reference to Nepal*. Practical Action. Kathmandu, Nepal. 2009.
15. Shrestha, U. B.; Gautam, S.; Bawa, K. S. Widespread climate change in the Himalayas and associated changes in local ecosystems. *PLOS ONE*. 2012, 7(5), e36741. (doi:10.1371/journal.pone.0036741).
16. Panthi, J.; Dahal, P.; Shrestha, M.; Aryal, S.; Krakauer, N.; Pradhanang, S.; Lakhankar, T.; Jha, A.; Sharma, M.; Karki, R. Spatial and temporal variability of rainfall in the Gandaki River Basin of Nepal Himalaya. *Climate*. 2015, 3, 210–226. (DOI:10.3390/cli3010210).
17. Nepal, S. Impacts of climate change on the hydrological regime of the Koshi river basin in the Himalayan region. *J. Hydro-Environment Res.* 2016, 10, 76–89. (DOI:10.1016/j.jher.2015.12.001).
18. Khatiwada, K.; Panthi, J.; Shrestha, M.; Nepal, S. Hydro-climatic variability in the Karnali river basin of Nepal Himalaya. *Climate*. 2016, 4, 17. (DOI:10.3390/cli4020017).
19. Agrawal, A.; Babel, M.S.; Maskey, S.; Shrestha, S.; Kawasaki, A.; Tripathi, N.K. Analysis of temperature projections in the Koshi River Basin, Nepal. *International Journal of Climatology*. 2016, 36: 266–279.
20. Manadal, R.A. Weather station and annual temperature dynamics in the elevation gradient (spatial and temporal analysis of Chitwan-Annapurna, Nepal). *J His Arch & Anthropol Sci*. 2020, 5(1):37–44.
21. MoFSC. *Strategy and Action Plan 2016-2025, Chitwan Annapurna Landscape, Nepal*. Ministry of Forests and Soil Conservation, Kathmandu Nepal. 2015.
22. Dobremez, J.F. *Ecology and Biogeography of Nepal*. National scientific research centre Paris, France. (In French). 1976.
23. TISC. *Forest and Vegetation Types of Nepal*, Tree Improvement and Silviculture Component (TISC), TISC documentation series No 105. Department of Forest, HMG/NARMSAP, Kathmandu. 2002.

24. MoE. National Adaptation Programme of Action (NAPA) to Climate Change. Ministry of Environment, Kathmandu, Nepal: Government of Nepal. 2011.
25. Karpouzos, D.K.; Kavalieratou, S.; Babajimopoulos, C. Trend analysis of precipitation data in Pieria Region (Greece). *Eur. Water.* 2010, 30: 31–40.
26. Partal, T.; Kahya, E. Trend analysis in Turkish precipitation data. *Hydrol. Processes.* 2006, 20(9), 2011–2026.
27. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. 2013 (URL <http://www.R-project.org/>).
28. Petersen, L.; Pellicciotti, F. Spatial and temporal variability of air temperature on a melting glacier: Atmospheric controls, extrapolation methods and their effect on melt modeling, Juncal Norte Glacier, Chile, *J. Geophys. Res.* 2011, 116, D23109, doi:10.1029/2011JD015842.
29. Kattel, D. B.; Yao, T.; Yang, K.; Tian, L.; Yang, G.; Joswiak, D. Temperature lapse rate in complex mountain terrain on the southern slope of the central Himalayas, *Theor. Appl. Climatol.* 2013, 113(3–4), 671–682, doi:10.1007/s00704-012-0816-6.
30. Tang, Z., Fang, J. Temperature variation along the northern and southern slopes of Mt. Taibai, China. *Agricultural and Forest Meteorology.* 2006, 139, 200–207. doi:10.1016/j.agrformet.2006.07.001.
31. Shrestha, A.B.; Wake, C.P.; Dibb, J.E.; Mayewski, P.A. Precipitation fluctuations in the Nepal Himalaya and its vicinity and relationship with some large scale climatological parameters. *Int J Climatol.* 2000, 20: 317–327.
32. Poudel, S.; Shaw, R. The relationships between climate variability and crop yield in a mountainous environment: a case study in Lamjung district, Nepal. *Climate.* 2016, 4, 13; doi:10.3390/cli4010013.
33. Luitel, D.R.; Siwakoti, M.; Jha, P.K. Relationship between Climatic Variables and Finger Millet Yield in Syangja, Central Nepal. *Journal of Forest and Livelihood.* 2018, 16 (2), 1–13.
34. McSweeney, C.; New, M.; Lizcano, G. Nepal. UNDP climate change country profiles series. 2012. Retrieved from <http://www.geog.ox.ac.uk/research/climate/projects/undp-cp/index.html?country=Nepal&d1=Reports>.

35. DHM. Study of climate and climatic variation over Nepal. A final report submitted to Ministry of science, technology and environment, Department of Hydrology and Meteorology, Kathmandu, Nepal. 2015.

36. MoSTE. *Nepal Second National Communication to United Nations Framework Convention on Climate Change*. Ministry of Science, Technology and Environment.2014.<http://unfccc.in/t/resource/docs/natc/nplnc2.pdf>.

37. Shrestha, M. L. Inter-annual variation of summer monsoon rainfall over Nepal and its relation to Southern Oscillation Index. *Met. Atmos. Phys.* 2000, 75, 21-28.

38. Dawadi, B. Climatic records and linkage along an altitudinal gradient in the southern slope of Nepal Himalaya. *Journal of Nepal Geological Society*.2017, 53, 47-56.

39. Kansakar, S.R.; Hannah, D.M.; Gerrard, J.; Rees, G. Spatial pattern in the precipitation regime in Nepal. *Int. J. Climatol.* 2004, 24, 1645–1659.

40. Putkonen, J. K. Continuous Snow and Rain Data at 500 to 4400 m altitude near Annapurna, Nepal, 1999–2001, *Arctic, Antarctic, and Alpine Research*. 2004,36:2, 244-248.

41. WESC. Water resources of Nepal in the Context of Climate Change .Water and energy commission secretariat, Singhadurbar, Kathmandu Nepal, 2011.

42. Sharma, K.P.; Vorosmarty, C.J.; Moore III, B. Sensitivity of the Himalayan Hydrology to Land-use and Climatic Changes. In: *Climatic Change* 47, Kluwer Academic Publishers, Dordrecht, the Netherlands. 2000,117-139.

43. Hansen, J.; Sato, M.; Ruedy, R.; Lo, K.; Lea, D.W.;Elizade, M. *Global temperature change*. Proc Natl Acad Sci USA. 2006, 103: 14288–14293. doi:10.1073/pnas.0606291103.

44. Chen, B.; Chao, W.C.; Liu, X. Enhanced climatic warming in the Tibetan plateau due to doubling CO<sub>2</sub>: A model study. *Climate Dynamics*.2003, 20,401–413.

45. ICIMOD. *Snow Cover Statistics- Nepal*. E-bulletin: International Centre for Integrated Mountain Development (ICIMOD), Nepal. 2013.

46. Heynen, M.; Miles, E.; Ragettli, S.; Buri, P.; Immerzeel, W.W.;Pellicciotti, F. Air temperature variability in a high-elevation Himalayan catchment. *Annals of Glaciology*. 2016, 57(71) doi: 10.3189/2016AoG71A076.

47. Immerzeel, W. W.; Petersen, L.; Ragettli, S.; Pellicciotti, F. The importance of observed gradients of air temperature and precipitation for modeling runoff from a glacierized watershed in the Nepalese Himalayas, *Water Resour. Res.* 2014, 50, 2212–2226, doi:10.1002/2013WR014506.

### Annex-1

Stations used to analyze the temperature trends in CHAL, 1970-2019

Station Name	Station ID	Station type	District	Latitude	Longitude	Altitude (m)	Data available
<b>Lower tropical bio climatic zone(&lt;500m)</b>							
Dumkauli	706	Agro-meteorology	Nawalparasi	28.1	84.22	154	1976
Simari	728	Climatology	Nawalparasi	28.12	83.75	154	1981
Chapakot	810	Climatology	Syangja	27.88	83.82	460	1979
Rampur	902	Agro-meteorology	Chitwan	28.02	84.42	256	1970
Hetauda NFI	906	Climatology	Makawanpur	27.68	85.05	474	1971
<b>Upper tropical bio climatic zone(500-1000m)</b>							
Baglung	605	Climatology	Baglung	27.93	83.6	984	1978
Beni bazar	609	Climatology	Myagdi	27.53	83.57	835	1989
Kusma	614	Climatology	Parbat	28.28	83.7	891	1989

Khudi bazar	802	Climatology	Lamjung	28.3	84.37	823	1970
Pokhara airport	804	aeronautical	Kaski	28.03	84.0	827	1970
Syangja	805	Climatology	Syangja	27.62	83.88	868	1976
Malepatan	811	Agro-meteorology	Kaski	27.92	84.12	856	1970
Khairenitar	815	Climatology	Tanahu	28.1	84.1	500	1970
<b>Lower subtropical bio climatic zone(1000-1500)</b>							
Tansen	702	Climatology	Palpa	28.22	83.53	1067	1972
Gorkha	809	Agro-meteorology	Gorkha	28	84.62	1097	1970
Nuwakot	1004	Climatology	Nuwakot	27.93	85.02	1003	1970
Pensayakhola	1057	Climatology	Nuwakot	28.28	85.12	1240	1975
<b>Upper subtropical bio climatic zone(1500-2000)</b>							
Khachikot	715	Climatology	Argakhanchi	28	83.15	1760	1977
Tamghas	725	Climatology	Gulmi	27.88	83.25	1530	1981
Lumle	814	Agro-meteorology	Kaski	27.8	83.8	1740	1970
Dhunche	1055	Climatology	Rasuwa	28.1	85.3	1982	1989
<b>Temperate bio climatic zone(2000-2500m)</b>							
Jomsom	601	Climatology	Mustang	27.87	83.72	2744	1970
Thakmarpha	604	Agro-meteorology	Mustang	27.68	83.7	2566	1970
Daman	905	Climatology	Makawanpur	27.87	85.08	2314	1972
Lete	607	Climatology	Mustang	28.07	83.6	2384	1998
Kakani	1007	Agro-meteorology	Nuwakot	28.07	85.25	2064	1972

Source: Department of Hydrology and Meteorology

## Annex-2

### Stations used to analyzed the precipitation trends in CHAL

Station Name	Station ID	Station type	District	latitude	Longitude	Altitude
<b>Lower tropical bioclimatic zone(&lt;500m)</b>						
Ridi bazar	701	Precipitation	Gulmi	27.95	83.43	442
Beluwa (girwari)	704	Precipitation	Nawalparasi	27.68	84.05	150
Dumkauli	706	Agrometeorology	Nawalparasi	27.68	84.22	154
Dumkibas	710	Precipitation	Nawalparasi	27.58	83.87	164
Chapkot	810	Climatology	Syangja	27.88	83.82	460
Damauli	817	Climatology	Tanahun	27.97	84.28	358
Rampur	902	Agrometeorology	Chitawan	27.62	84.42	256
Jhawani	903	Precipitation	Chitawan	27.58	84.53	270
Hetaundan.f.i.	906	Climatology	Makwanpur	27.42	85.05	474
Beluwa(manahari)	920	Precipitation	Makwanpur	27.55	84.82	274
Bharatpur	927	Climatology	Chitawan	27.67	84.43	205
<b>Upper tropical bioclimatic zone(500-1000m)</b>						
Baglung	605	Climatology	Baglung	28.27	83.60	984
Beni bazar	609	Climatology	Myagdi	28.35	83.57	835
Kushma	614	Climatology	Parbat	28.22	83.70	891
Garakot	726	Precipitation	Palpa	27.87	83.80	500
Khudi bazar	802	Climatology	Lamjung	28.28	84.37	823
Pokhara airport	804	Aeronatical	Kaski	28.22	84.00	827
Syangja	805	Climatology	Syangja	28.10	83.88	868
Kunchha	807	Precipitation	Lamiung	28.13	84.35	855
Bandipur	808	Climatology	Tanahun	27.93	84.42	965
Malepatan	811	Agrometeorology	Kaski	28.12	84.12	856
Khairinitar	815	Agrometeorology	Tanahun	28.03	84.10	500
Arughatbazar	1002	Precipitation	Dhading	28.05	84.82	518
<b>Lower subtropical bioclimatic zone (1000-1500)</b>						
Tatopani	606	Precipitation	Myagdi	28.48	83.65	1243
Tansen	702	Climatology	Palpa	27.87	83.53	1067
Musikot	722	Precipitation	Gulmi	28.17	83.27	1280
Jagat (setibas)	801	Precipitation	Gorkha	28.37	84.90	1334
Gorkha	809	Agrometeorology	Gorkha	28.00	84.62	1097
Lamachaur	818	Precipitation	Kaski	28.27	83.97	1070
PAMDUR	830	Precipitation	Kaski	28.27	84.78	1160
Makwanpurgadhi	919	Precipitation	Makwanpur	27.42	85.17	1030
Nuwakot	1004	Climatology	Nuwakot	27.92	85.17	1003
Dhading	1005	Precipitation	Dhading	27.87	84.93	1420
Pansayakhola	1057	Precipitation	Nuwakot	28.02	85.12	1240
<b>Upper subtropical bioclimatic zone (1500-2000)</b>						
Karki neta	613	Precipitation	Parbat	28.18	83.75	1720
Khanchikot	715	Climatology	Arghakhanchi	27.93	83.15	1760
Tamghas	725	Climatology	Gulmi	28.07	83.25	1530
Lumle	814	Agrometeorology	Kaski	28.30	83.80	1740

Ghandruk	821	Precipitation	Kaski	28.38	83.80	1960
Chisapanigadhi	904	Precipitation	Makwanpur	27.55	85.13	1706
Markhugaun	915	Precipitation	Makwanpur	27.62	85.15	1530
Timure	1001	Climatology	Rasuwa	28.28	85.38	1900
Thamachit	1054	Precipitation	Rasuwa	28.17	85.32	1847
Dhunche	1055	Climatology	Rasuwa	28.10	85.30	1982
<b>Lower temperate bioclimatic zone (2000-2500m)</b>						
Lete	607	Climatology	Mustang	28.63	83.60	2384
Bobang	615	Precipitation	Baglung	28.40	83.10	2273
Daman	905	Climatology	Makwanpur	27.60	85.08	2314
Kakani	1007	Agrometeorology	Nuwakot	27.80	85.25	2064
<b>Upper temperate bioclimatic zone (2500-3000)</b>						
Jomsom	601	Climatology	Mustang	28.78	83.72	2744
Thakmarpha	604	Agrometeorology	Mustang	28.75	83.70	2566
Gurjakhani	616	Climatology	Myagdi	28.60	83.22	2530
Chame	816	Climatology	Manang	28.55	84.23	2680
<b>Lower subalpine bioclimatic zone (3000-3500m)</b>						
Ghami (mustang)	610	Precipitation	Mustang	29.05	83.88	3465
Manangbhot	820	Climatology	Manang	28.67	84.02	3420
<b>Alpine bioclimatic zone (3500-4000m)</b>						
Ranipauwa (m.nath)	608	Precipitation	Mustang	28.82	83.88	3609
Mustang(lomangthang)	612	Climatology	Mustang	29.18	83.97	3705
Larkesamdo	806	Precipitation	Gorkha	28.67	84.62	3650

Source: Department of Hydrology and Meteorology,GoN