

Photoimmunotherapy using cationic and anionic photosensitizer-antibody conjugates against HIV Env-expressing cells

Mohammad Sadraei¹, Calise Bahou², Edgar Ferreira da Cruz³, Luiz Mário Ramos Janini³,
Ricardo Sobhie Diaz^{*3}, Ross W. Boyle^{*4}, Vijay Chudasama^{*2}, Francisco E. G. Guimarães^{*1}

¹ Instituto de Física de São Carlos, Universidade de São Paulo, Caixa Postal 369, São Carlos,
SP, CEP 13560-970, Brazil

² Department of Chemistry, University College London, London WC1H 0AJ, UK

³ Laboratório de Retrovirologia, Universidade Federal de São Paulo, São Paulo, Brazil

⁴ Department of Chemistry, University of Hull, Cottingham Road, Hull, HU6 7RX, UK

*** Corresponding authors:**

Francisco Eduardo Gontijo Guimarães, email: guimaraes@ifsc.usp.br, +55(16)33739792

Vijay Chudasama, email: v.chudasama@ucl.ac.uk

Ross W. Boyle, email: r.w.boyle@hull.ac.uk

Ricardo Sobhie Diaz, email: rsdiaz@catg.com.br

*** Current addresses:**

Instituto de Física de São Carlos, Universidade de São Paulo, Caixa Postal 369, São Carlos,
SP, CEP 13560-970, Brazil

Word count: Abstract 184; Main text 2187;

ABSTRACT

Different therapeutic strategies have been investigated to target and eliminate HIV-1-infected cells by using armed antibodies specific to viral proteins, with varying degrees of success. Herein, we propose a new strategy by combining photodynamic therapy (PDT) with HIV Env-targeted immunotherapy, and refer to it as HIV photoimmunotherapy (PIT). A human anti-gp41 antibody (7B2) was conjugated to two photosensitizers with different charges through different linking strategies; “Click” conjugation by using an azide-bearing porphyrin attached *via* a disulfide bridge linker with a drug-to-antibody ratio (DAR) of exactly 4, and “Lysine” conjugation by using phthalocyanine IRDye 700DX dye with average DARs of 2.1, 3.0 and 4.4. These photo-immunoconjugates (PICs) were compared *via* biochemical and immunological characterizations regarding the dosimetry, solubility, and cell targeting. Photo-induced cytotoxicity of the PICs were compared using assays for apoptosis, reactive oxygen species (ROS), photo-cytotoxicity, and confocal microscopy. Targeted phototoxicity seems to be primarily dependent on the binding of PS-antibody to the HIV antigen on the cell membrane, whilst being independent of the PS type. This is the first report of the application of PIT for HIV immunotherapy by killing HIV Env-expressing cells.

Keywords: HIV immunotherapy, photoimmunotherapy, photodynamic Therapy, porphyrin, phthalocyanine, HIV-infected cell, monoclonal antibody

1. Introduction

Antibody-based therapies have become important clinical tools in treating chronic diseases. Different immunotherapeutic strategies, with limited success, have been investigated to target and eliminate viral-infected cells by using armed antibodies specific to viral proteins (1). This

can be achieved by linking a drug to monoclonal antibodies (MAbs), to form immunotoxins (IT) (2), radio-labeled antibodies (3,4) or other cell-binding conjugates (5,6) capable of specifically killing infected cells, such as HIV infected cells (1,7). Photoimmunotherapy (PIT) is an emerging targeted therapy that uses photosensitizer-conjugated MAbs. PIT has certain advantages over immunotoxins or RIT to eradicate infected cells (8). In immunotherapy based on immunotoxins, a MAbs is conjugated to an immunogenic toxin such as ricin (9), pulchellin (10,11) or shiga toxin (12), which can elicit an immunogenicity exemplified by the anti-toxin response (3,13). PIT is a minimally invasive treatment, which is safer and cheaper than immunotoxins or RIT (14). While HIV RIT is drawing the attention of researchers (4), there are no studies on the possibility of HIV treatment by PIT.

In this paper, we propose HIV immunotherapy *via* arming HIV monoclonal antibodies with photosensitizers (PSs). Anti-HIV immunoconjugates must be targeted to the HIV envelope spike (Env) that consists of gp160 (9,15), gp120 (16), and gp41 glycoproteins (11,16). PSs can be activated by visible light of a specific wavelength to initially form the excited singlet state, followed by transition to the longer-lived excited triplet state (17). This triplet state can undergo photochemical reactions in the presence of oxygen *via* Type I photochemical mechanisms (electron transfer to produce superoxide, hydrogen peroxide, and hydroxyl radicals), typical of phenothiazinium dyes (18), or Type II mechanisms (energy transfer process), typical of porphyrins (19,20), and Rose Bengal (21,22). Reactive oxygen species (ROS) can oxidize lipids, proteins, and nucleic acids, leading to cell damage and death through necrosis where cells swell and lyse to release their intracellular content causing inflammation, or alternatively cells can undergo apoptosis (23,24). The conventional treatment using only PS and light is called photodynamic therapy (PDT), which can destroy cells nonspecifically when exposed to light (23). Related to this, PIT is the targeted form of conventional PDT, achieved through the conjugation of PS with MAbs targeting specific cell surface receptors (25,26). PIT can also

reduce other issues, such as dark toxicity and PS aggregation in aqueous media which reduce PDT efficacy (14). The conjugation of a photosensitizer to antibodies can be achieved through lysine modification (27) or by functionalization of thiols generated *via* reduction of interchain disulfide bonds without functional rebridging; however, neither of these conjugation strategies is ideal (28,29). Lysine modification generates heterogeneous products, while conventional methods of cysteine modification following interchain disulfide reduction avoid this issue, they result in the permanent loss of structural disulfide bonds, which may impact Fc function (30) and negatively affect the stability of the antibody *in vivo* (31,32).

Recently, we described a novel technology of disulfide modification based on the insertion of pyridazinediones (PDs) bearing orthogonal 'clickable' handles into the native inter-strand disulfide bonds of trastuzumab full antibody (33–36). This was then combined with mild bio-orthogonal conjugation of porphyrin photosensitizers and the conjugate targeted Her2 positive breast cancer cells (37,38).

In this study, we utilized lysine and disulfide modifications to identify an appropriate photo-immunoconjugate (PIC) for our suggested PIT. Briefly, a human anti-gp41 antibody (7B2) (39) was conjugated with two different photosensitizers, cationic porphyrin with a net charge of 3+ and anionic IR700 with a net charge of 4-. We employed two different strategies for conjugation to the antibody; lysine conjugation by using phthalocyanine IRDye700DX dye (27), and "Click" conjugation by using an azide-bearing porphyrin with a strained alkyne attached *via* a disulfide bridge linker (28). The comparison between PICs is of interest with regard to the dosimetry and solubility of PICs and mechanism of *in vitro* cytotoxicity (Figure 1A).




Figure1. Structure and function of photo-immunoconjugates (PICs). **(A)** Anti-gp41 monoclonal antibody (7B2 MAb) conjugated with 4 porphyrins can specifically bind to the HIV infected cells expressing HIV Env (gp41 and gp120). The presence of CD4 may increase the binding ability of 7B2-porphyrin to the gp41 on the cell membrane. Porphyrins can be activated by visible light of a specific wavelength (herein 405 nm) to initially generate singlet oxygen or ROS reaction, and consequently kill the target cell. Furthermore, the emission of red fluorescent by porphyrin can be observed during irradiation. **(B)** Schematic picture demonstrates two generations of PICs in the study; Lysine modification leading to heterogeneous IR700-PICs with “average distribution” of DARs 2, 3 and 4 with undefined physical and pharmacokinetic properties. In a click approach, disulfide re-bridging by a pyridazinedione construct yields a homogeneous porphyrin-PIC with a constant DAR4.

2. Results and Discussion

2.1. Production and Characterization of Photoimmunoconjugates PICs

Two generations of PICs were produced; Porphyrin-7B2 PIC was produced through the use of the strain-promoted azide–alkyne cycloaddition (SPAAC) methodology in two steps: firstly, the

insertion of a strained alkyne harboring pyridazinediones (PD) into native interchain disulfide bonds in humanized IgG1 MAb (7B2), then, the antibody-alkyne reacted with azide-bearing water-soluble porphyrin.

As a side by side comparison, a well-known lysine-reactive and relatively hydrophilic dye, IRDye 700DX, in different molar concentrations, was covalently conjugated with primary amine groups on the structure of 7B2 MAb using a *N*-hydroxysuccinimide (NHS) ester of the photosensitizer (Figure 1B).

Chemical characterization by UV-Vis spectrophotometry showed a PD loading of 4.3 for the rebridged antibody and a DAR of 4.1 for the final porphyrin-PIC species. IR700-PICs showed DARs of 2.1, 3.0 and 4.4, which were equal to molar ratios of 3.5, 7, and 14, before conjugation, respectively (Figure 2A). The DLS results showed that the attachment of small molecule drugs to the 7B2 MAb slightly increases the molecule's average size (hydrodynamic radius, R_h) and polydispersity (%Pd), indicating the presence of some high molecular weight species in this region (Figures 2A and 2D). 7B2-porphyrin showed the highest %Pd and R_h in the main peak. The zeta potential of 7B2 antibody changed very slightly from +0.05 mV towards +0.01 and - 0.04 for cationic porphyrin and anionic IR-700 DAR4, respectively (Figure 2A). The binding of the 7B2 based-PICs to gp41 loop peptide antigen was examined within the conjugation. ELISA results showed that an increase of IR700-antibody-ratio decreases the binding ability, as 7B2-IR700 DAR4 showed the lowest binding to the gp41. In contrast, the conjugation of porphyrin with 7B2 MAb did not significantly affect the immunologic specificity of the antibody (Figure 2B). Results of microcapillary electrophoresis of non-reduced antibody (H and L chains) and PICs demonstrated the molecular weights (MWs) with 151.4, 166.1 and 166.9 kDa for naked antibody, rebridged antibody and clicked porphyrin-antibody. The increase in the MWs of IR700-PICs were in agreement of determined DARs, with 154.1, 154.7 and 156.8 kDa for DARs 2, 3 and 4, respectively (Figures 2A and 2C). In the presence of reducing agent (TCEP), the light

and heavy chains of either 7B2 or 7B2-IR700 were observed, due to the reduction of all available interchain disulfide bridges. The reduced 7B2-porphyrin did not show the separation of chains, confirming the accuracy of click chemistry with DAR4 (Figure 2C).




Figure2. Characterization of photo-immunoconjugates (PICs). **(A)** The table shows products of 7B2 MAb conjugated with different molar concentrations of IR700, rebridged antibody with pyridazinedione and porphyrin. Chemical characterization by UV-Vis spectrophotometry shows the antibody to PS ratio (DAR). In DLS studies, the zeta potential, ζ , was estimated according to

the Huckel equation. The increase in the molecular weights (MWs) of conjugations, measured by microcapillary electrophoresis, are in confirmation of determined DARs. **(B)** ELISA plates coated with gp41 antigen, as a peptide representing 7B2's epitope. Results are representative of means of triplicate values with three individual experiments. 924 MAb was used as an isotype control. **(C)** Microcapillary electrophoresis of reduced and non-reduced 7B2 MAb and PICs, before and after irradiation. Unlike 7B2 and 7B2-IR700, the reduced 7B2-porphyrin did not show the separation of light and heavy chains, confirming the accuracy of click chemistry with DAR4. Size standards are indicated on the side of each "gel". **(D)** DLS Histograms of hydrodynamic radius (R_h) for 7B2 MAb and PICs, monitoring how the R_h preserved during the conjugation process. When IR700 was conjugated with an antibody molecule, ligand dissociation from IR700 affected the shape and solubility of the antibody to which it is attached as well as antibody-antigen complexes.

In total, the characterization of PICs showed that the shape, solubility and binding ability of IR700-PICs was affected by increasing the drug-to-antibody ratio (DAR), while they were preserved for water-soluble homogeneous porphyrin-7B2 with a constant DAR of 4, due to the retention of rigid structural bridges on the antibody.

2.2. Flow Cytometry analysis

By using Cary Eclipse UV-Vis-NIR spectroscopy, we analyzed the excitation and emission spectra before flow cytometry study without using a secondary antibody. We showed the excitation of porphyrin by blue laser at 488 nm would not interfere with the emission intensity. The excitation by violet laser at 405 nm caused red emission at 725 nm with 1.163 (a. u.) intensity, while the maximum red fluorescent intensity (3.779 a. u.) can be observed by the excitation of Soret band of porphyrin at 432 nm. The results were in agreement with the theoretical analysis of UV-Vis spectra (Supplementary Figure S1).

Specific cell binding to native Env was examined by flow cytometry. Without using a secondary antibody, the red fluorescent emission from IR700-PICs and porphyrin-PIC were directly detected on the Env-transfected cells, by filter Qdot 705 and filter Qdot 655, respectively (Figure 3A). By using FITC-secondary antibody, PICs showed equal binding ability into Env-transfected cells, while 7B2-IR700 DAR4 showed somewhat unspecific binding into the control 293T cells (Figure 3B). Not surprisingly, the presence of soluble CD4 (sCD4) showed a synergic effect on the binding ability of porphyrin-7B2, in comparison to the absence of sCD4 (Figure 3C). This result is in agreement with our previous studies on immunotoxins (11,16), as we showed sCD4 enhances epitope exposure.




Figure 3. The flow cytometry histogram of PICs by using 293T control cells and Env-transfected 293T cells incubated in PBS + 1% BSA + sodium azide (0.2%). **(A)** PIC binding to 293T cells

transfected with 92UG037.8 gp160 (293T/92UG) cells was directly detected through red fluorescent emission from IR700-PICs and porphyrin-PIC, by filter Qdot 705 and filter Qdot 655, respectively. **(B)** By using FITC-secondary immunofluorescence, PICs showed equal binding ability into Env-transfected cells, while 7B2-IR700 DAR4 showed somewhat unspecific binding into the control 293T cells. Results are representative of at least three independent experiments. Isotype control (chimeric RAC18) is shown as blue shaded histogram.

Interestingly, IR700-PICs with different DARs (Figure 2A) and different binding abilities (Figure 2B) showed equal fluorescence intensity, not only with the direct red emission by filter Qdot 705 (Figure 3A), also with indirect emission by FITC-secondary antibody (Figure 3B). This can be due to the quenching of chromophores, as some that are too close to each other will prevent the chromophores from being excited by laser light and then bleaching (40). There is no linear relationship between the number of chromophores per antibody and fluorescence intensity (41), while there is a correlation between the trends in fluorescence quenching and efficacy of heterogeneous IR700-PICs (42). We assume this quenching would not happen in homogeneous porphyrin-PIC with disulfide modification.

2.3. ROS generation

In cellular studies, to have an equal irradiation for both porphyrin-PIC and IR700-PIC, for a side by side comparison, a custom-made LED device with a broad spectrum of light (380–780 nm) was utilized, as the blue spectra can excite the Soret band of porphyrin at 432 nm, meanwhile, red spectra excites IR700-7B2 at 689 nm (Figure 4C).

To detect the ROS generation, the incubated cells by PICs and H2 DCF-DA fluorescent probe were irradiated with 50 J/cm². H2DCF-DA fluorescent probe was applied to detect ROS production in PIT-treated cells by using flow cytometry. The Env-transfected cells incubated with PICs exhibited a significant increase in ROS levels, whereas control 293T cells did not show

ROS production (Figure 4A). In parallel, in the positive control group, we observed ROS production in 293T cells which were treated with 80 μM H_2O_2 for 24 hr in the dark (Figure 4B). The irradiated cells, with either no incubation or incubated with 7B2 antibody, showed an increase in the fluorescence intensity in comparison to the unstained cells in darkness.




Figure 4. ROS generation and apoptosis assay by PICs. **(A)** Samples were kept in the dark (represented by grey) or were irradiated with 50 J/cm^2 . **(B)** As a positive control, 293T cells were treated with 80 μM H_2O_2 for 24 h in dark. Control group (without H_2O_2) was incubated under the same conditions. **(C)** The diagram demonstrates a broad spectrum of light (380–780 nm) by a homemade LED device (30 mW/cm^2) for irradiation of both IR700 and porphyrin PICs, at the same time. Blue spectra can excite the Soret band of porphyrin at 432 nm, meanwhile red spectra can excite 7B2-IR700 at 689 nm. **(D)** In apoptosis assay by Annexin V - FITC/PI, the Env-transfected cells were incubated with PICs for 4 hr, then irradiated. PICs showed cell death due to late apoptosis after 4 hr incubation. Control 7B2 antibody showed neither cell death nor apoptosis sign. ($n = 3$).

2.4. Apoptosis Assay

For apoptosis assays, the Env-transfected cells were incubated by PICs for 4 hr, irradiated, then Annexin V - FITC/PI dye added before flow cytometry analysis. No features related to early stages of apoptosis, such as extracellular exposure of phosphatidylserine, was observed in response to PICs. While, both porphyrin-PIC and IR700-PIC showed cell death due to late apoptosis (FITC emission) after 4 hr incubation, and signs of necrosis (PI emission). Control 7B2 antibody showed neither cell death nor signs of apoptosis (Figure 4D).

2.5. Photo-cytotoxicity Assay

Env-transfected 293 T cells (293 T/92UG) and control 293T cells were incubated with a serial dilution of PICs starting at 500 nM in the presence of sCD4 (anti-gp120) for 4 hr then were assayed by photoimmunotherapy (PIT) with 20 J/cm². IR700-PIC with a DAR of 3 appeared more cytotoxic than porphyrin-PIC with a constant DAR of 4. PICs showed neither dark cytotoxicity on transfected cells nor nonspecific killing on 293T control cells. The cell growth inhibition was observed due to the irradiation on the cells without PIC treatment (Green stars in Figure 5A). Optical microscopic observations before and after PIT demonstrated the bleb formation as signs of necrotic cell death. Cells PIT-treated with IR700-7B2 showed not only rapid bleb formation but also cell debris (Supplementary Figure S2).

To study how the combination of both PICs may affect the cytotoxicity, an equal molarity of porphyrin-PIC was mixed with IR700-PIC. Interestingly, this combination showed a significant increase in the cytotoxicity in compare to the average cytotoxicity of both PICs, due to the synergic effect of two PICs with two different mechanism (Figure 5B).

Furthermore, the photo-cytotoxicity of porphyrin-7B2 increased twice by increasing the time of incubation from 1 hr to 4 hr, indicating the internalization-dependency of porphyrin-mAb. In

contrast, the photo-cytotoxicity of IR700-7B2 did not change significantly on increasing the time of incubation at 37 °C (Figure 5C).




Figure 5. Comparing the photo-cytotoxicity and targeting of PICs by direct cytotoxicity assay.

(A) Env-transfected 293 T cells (293 T/92UG) and control 293T cells were incubated with 7B2-IR700 (DAR 3), 7B2-porphyrin and an equal molarity mixture of both PICs in the presence of sCD4 (anti-gp120) for 4 h then were assayed by PIT. Data are \pm means S.E.M. ($n = 3$) with three individual experiments. Where no error bars are visible, they are obscured by the symbol.

(B) An equal molarity mixture of porphyrin-PIC with IR700-PIC may increase the cytotoxicity in compare to porphyrin-PIC or the average cytotoxicity of both PICs. **(C)** Incubation of transfected cells in two different times (1 and 4 h) before PIT showed the internalization-dependency of

Mab-porphyrin to have more photo-cytotoxicity, while the membrane binding of 7B2-IR700 was sufficient to induce cell death. Data are \pm means S.E.M. ($n = 3$).

2.6. Confocal Microscopy Study

The cells were treated with IR700-7B2 or 7B2-porphyrin for 1 hour, then washed and irradiated by 2-photons, 800 nm, 2.2 mW/cm. Red fluorescent emission from IR700-7B2 was directly detected (Figure 6A and Supplementary Video 1). As 7B2-porphyrin did not show strong red fluorescent emission, FITC anti-human IgG secondary antibody was applied to detect PIC indirectly (Figure 6B and Supplementary Video 2). After 10 minutes of adding porphyrin-7B2, the cellular swelling and bleb formation were observed as signs of necrotic cell death. The same necrotic signs were observed by adding IR700-7B2, but in 5 minutes of irradiation. After 30 minutes irradiation, Orange Acrylamide and Ethidium Bromide were added as Live-dead dyes. Interestingly, the irradiated and non-irradiated regions were clearly distinguishable in red and green colors, respectively, confirming no dark toxicity by PICs (Figure 6C).




Figure 6. Target-specific cell death in response to PIT in HIV Env-expressing 293T cells. The cells were treated with IR700-7B2 (**A**) or 7B2-porphyrin (**B**) in the presence of sCD4 for 1 hour before laser irradiation by microscope. (**A**) The emitted red fluorescent from IR700-7B2 was directly detected. (**B**) 7B2-porphyrin was detected indirectly by using FITC anti-human IgG secondary antibody. After 30 minutes irradiation, the cell viability was studied by using Orange Acrylamide and Ethidium Bromide (Live/dead dye). White arrows indicate the bleb formation caused the necrotic cell death. The white bar indicates 10 μ m. (**C**) The red square shows the

irradiated region that the cells are dead (red color). The non-irradiated region (green color) signifies live cells, confirming no dark toxicity by PIC.

In conclusion, we compared the mechanism of two photosensitizers in the formation of PICs regarding the induction of cell death in HIV-transfected cells. We also addressed some of the main issues associated with conventional PDT including low selectivity, controlled drug dosimetry, dark toxicity and low water solubility of photosensitizer. Finally, we studied the possibility of PIT application to directly killing HIV infected-cell. Targeted phototoxicity seems to be primarily dependent on the binding of PS-antibody to the antigen on the cell membrane or HIV Envelope, whilst being independent of the PS type. The ability to optimize the excitation of PSs in deep tissue may give us a highly flexible theranostic platform for HIV researches. To the best of our knowledge, this is the first report to demonstrate the utility of photoimmunotherapy for HIV Env-expressing cells. The results of this strategy can potentially translate to viral photoimmunotherapy against other enveloped viruses.

3. Materials and Methods

3.1. Consumable materials.

3.1.1. Chemical reagents.

All reagents are from ThermoFisher Scientific (Waltham, MA, USA), unless otherwise stated.

3.1.2. Cell lines.

293T cell lines stably express clade A clinical isolate 92UG037.8 gp160 as native gp120/gp41 trimers (293T/92UG) (43). 293T cells were used as uninfected control cells. The transfected and non-transfected 293T cells were maintained at 37 °C in 5% CO₂ in DMEM

medium with 10% fetal bovine serum (Gibco Invitrogen, Grand Island NY). In this paper, Env-transfected cells refer to the 293T cells stably transfected with 92UG037.8 gp160.

3.1.3. Antibodies.

MAb 7B2 (Genbank accession numbers JX188438 and JX188439) is a human IgG1 which binds HIV gp41 at AA 598–604 (CSGKLIC) in the helix-loop-helix region (39,44). HY (Genbank accession numbers JX188440 and JX188441), an affinity matured version of the anti-CD4 binding site Ab b12 (45). MAbs were purified from hybridoma supernatant on Protein A agarose beads (Invitrogen) and eluted with 0.1 M glycine, pH 2.5, immediately neutralized, and dialyzed vs. PBS. Two types of soluble CD4 were used to observe CD4-mediated effects (44); CD4-183 contains the first two domains of CD4 including the region in domain 1 that binds the HIV coat protein gp120. CD4-IgG2 is a tetrameric fusion protein comprising human IgG2 in which the Fv portions of both heavy and light chains have been replaced by the V1 and V2 domains of human CD4. Goat anti-human IgG (Invitrogen) was conjugated to either alkaline phosphatase (AP) or fluorescein isothiocyanate (FITC).

3.2. Porphyrin-Antibody conjugation by click chemistry.

Conjugation between azide porphyrin and mAb 7B2 was carried out in two steps: antibody functionalization, then click chemistry conjugation. The process is a modification of our protocol described previously (36,46,47) and described in detail in the supplementary methods.

3.3. Conjugation and optimization of IR700-antibody by lysine modification.

IRDye 700DX (LI-COR Biosciences; IR700) was covalently linked to purified MAb 7B2 through an *N*-hydroxysuccinimide reactive group, according to the manufacturer's instructions (27). The process of conjugation and optimization are described in detail in the supplementary methods.

3.4. UV-Vis spectroscopy

UV-Vis spectroscopy was used to determine protein concentrations and photosensitizer to antibody ratios, firstly by using Nanodrop 1000 UV-Visible spectrophotometer (ThermoFisher Scientific) and then a Varian Cary 100 Bio UV-Visible spectrophotometer operating at 21 °C. Sample buffer was used as blank for baseline correction with extinction coefficients; $\epsilon_{280} = 250,440 \text{ M}^{-1} \text{ cm}^{-1}$ for antibodies, $\epsilon_{335} = 9,100 \text{ M}^{-1} \text{ cm}^{-1}$ for pyridazinedione scaffolds, $\epsilon_{422} = 165,175 \text{ M}^{-1} \text{ cm}^{-1}$ for porphyrin and $\epsilon_{689} = 165,000 \text{ M}^{-1} \text{ cm}^{-1}$ for IRDye 700DX. A Correction Factor at 280 nm of 0.25 (at A_{335}) was employed for pyridazinedione scaffolds, as described elsewhere (37).

3.5. Electrophoresis

Molecular size, purity and accuracy of the conjugation of products were determined using Non/reducing glycine-SDS-PAGE and then confirmed by microcapillary electrophoresis in the presence or absence of TCEP·HCl (6.0 μL , 20 mM in d.d water, 12 eq.), as a reductive agent (Agilent Bioanalyzer, GE Healthcare), following standard lab procedures.

3.6. ELISA

ELISAs were performed for Ag-binding specificity analysis and titration of purified MAbs and photo-immunoconjugates (PICs) in wells coated with gp41 antigen (1 $\mu\text{g/mL}$), as described elsewhere (16). The gp41 antigen was a linear peptide HIV-1 consensus clade B sequence [LGIWGCSGKLICTT] representing the epitope of 7B2. Binding of antibody to the antigen was detected with AP-conjugated secondary antibody (goat anti-human IgG) (Zymed Laboratories, South San Francisco, CA). Data are reported as optical density at 405 nm and represent means of triplicate values with three independent experiments.

3.7. Dynamic Light Scattering (DLS) and Zeta Potential

Hydrodynamic radii, electrophoretic mobility, zeta potential, and polydispersity of naked antibody and PICs were measured. Samples with 70 μ L volume at 1 mg/mL in UV-transparent 96-well plates were measured by using a DLS Wyatt Möbius (Wyatt Technologies, Dernbach, Germany) with incident light at 532 nm, at an angle of 163.5°. Samples were equilibrated at 25 ± 0.1 °C for 600 s prior to the measurements, and this temperature was held constant throughout the experiments. All samples were measured in triplicate with 10 acquisitions and a 5 s acquisition time. The change in cumulant fitted hydrodynamic radius in nanometers was monitored during the storage period. Results were calculated using the Dynamics 7.1.7 software (Wyatt Technologies). By using an Agilent Technologies Cary Series UV-VIS-NIR spectrofluorimeter (Cary 5000), we showed the incident laser beam at 532 nm is not within the fluorescent sample's (IR700-7B2 and porphyrin-7B2) band of excitation; as the analysis would not be disrupted or tainted.

3.8. Direct and indirect immunofluorescence using Flow cytometry

We used indirect immunofluorescence and flow cytometry to analyze the binding of photo-immunoconjugates (PICs) to Env-transfected 293T cells. 8×10^4 cells were incubated with PBS/BSA/0.01% sodium azide (PBA) for 30 min in RT, and then PICs added to reach the final concentrations of 10 μ g/mL in the presence of 5 μ g/mL soluble CD4. Cells were incubated with PICs for 1 hr at room temperature. For indirect assay, half of the samples were washed, and then stained with FITC conjugated goat anti-human IgG secondary antibody (2 μ g/mL) for 1 hr. All the samples, for direct and indirect assays, were washed twice and fixed in 100 μ L of 2% paraformaldehyde. After a minimum of 4 hr, 150 μ L of PBS was added. Cells (10,000) were studied on LSR Fortessa optics (Becton-Dickson), analyzed by Flow-Jo software (Treestar Inc).

Mean fluorescence of the gated cell population labelled with immunoconjugate was calculated in relation to the mean fluorescence of cells labelled with MAb 7B2.

Direct immunofluorescence without adding FITC conjugated secondary antibody was applied to analyze the binding of PICs to Env-transfected 293T cells. For IR700-PICs, the cells were excited by either violet laser at 405 nm or red laser at 640 nm, and the red fluorescence emission was detected by 710±50 band-pass filter (Qdot 705). For porphyrin-PICs, the cells were excited by violet laser at 405 nm, and the red fluorescence emission was detected by 660±20 band-pass filter (Qdot 655).

3.9. ROS detection

ROS generation in the cell lines was detected with the fluorescent probe H2DCF-DA (2',7'-Dichlorodihydrofluorescein diacetate). 2×10^5 cells were incubated with 1 μ M of native antibody, porphyrin-PIC, or IR700-PIC in 96 well plates in phenol-free DMEM for 45 minutes. The cells were washed and incubated with 1 μ M H2DCF-DA solution prepared in PBS, and washed again and incubated with phenol-free DMEM before irradiation with 50 J/cm². A mirror dark-plate, submitted to the same procedure, except for light exposure, contained the dark control groups. The cells incubated with 40 mM H₂O₂, without irradiation, were used as positive control. Fluorescence intensity (Ex/Em = 485/535 nm) was measured with the flow cytometry, with the procedure as mentioned above.

3.10. Apoptosis / cell death assay

Annexin V-FITC/PI cell staining was performed following 6 hr of standard photoimmunotherapy. Briefly, cells were incubated with Annexin V binding buffer (10 mM HEPES, pH 7.4; 140 mM NaCl; 2.5 mM CaCl₂) and stained with Annexin V-FITC (ThermoFisher Scientific) for 15 min. After three washes, cells were incubated with 2 μ g/mL of propidium iodide (PI) (Life Technologies, Carlsbad, CA, USA) for 15 min. The mean fluorescence intensity was measured

for 10,000 events. The data were collected using linear amplification for FSC and SSC, and logarithmic amplification for related filters, and analyzed by associated software as mentioned above.

3.11. Cell proliferation assay after photoimmunotherapy.

Naked 7B2 antibody, IR700-7B2, porphyrin-7B2 and a mixture of both PICs with equal molarity were diluted in the phenol-free DMEM medium, but without FCS, to give a range of five final concentrations of 500 nM, 250 nM, 125 nM, 62.5 nM and 31.2 nM into 5 wells of a 96-well plate. HIV Env-transfected cells and 293T cells were adjusted to a concentration of 1×10^6 cells/mL. Soluble CD4 was added to each well in a final concentration of 500 ng/mL. The plate incubated for 4 h at 37 °C and 5% CO₂. Identical plates were kept protected from light for the same period to obtain the proper dark control groups. After incubation, the cells were washed with a 4X excess of serum-free phenol-free DMEM to eliminate any unbound conjugate. Each plate was irradiated with 20 J/ cm² of light (380–780 nm) by a homemade LED array (30 mW/cm²) in two equal doses separated by 10 min, as described elsewhere (48,49). After irradiation, 10 µL of FCS was added to each well and the plates were returned to the incubator for 3 days. The growth of the cells in each group was also photographed under a microscope (IX81, Olympus America) at each time point, before and after irradiation. MTS/PMS substrate (Promega, Madison, WI) was added to each well and plates read hourly at 490 nm. Results represent the mean and SEM of triplicate samples (representative of three independent experiments), and are plotted as A₄₉₀ with the no cell background subtracted. The results are expressed as % of cell viability versus PIC concentration.

3.12. Live imaging by two-photon confocal microscopy.

Live cell imaging was performed using a Zeiss LSM 780 confocal inverted microscope with a Coherent Chameleon laser (Ti:sapphire) as a source for two-photons (2P) excitation at 800 nm.

The images were obtained by the average of 2 scans and no appreciate variation was observed. The spatial resolution was approximately 350 nm (considering the numerical aperture and the wavelength of excitation), as described previously (50). Detailed conditions and settings are described in the supplementary methods.

3.13. Statistical analyses.

Statistical analyses were performed using GraphPad Prism, v5.0. Data are shown as mean and SEM of the indicated number of replicate values. If no error bar appears present, the error bars are smaller than, and obscured by, the symbol. The method for statistical comparison is by unpaired two-tailed Student's t-test, unless specifically indicated otherwise.

Data Availability: All data generated or analyzed during this study are included in this published article and its Supplementary Information file. The detailed method of live microscopy with extra videos can be found at

https://docs.google.com/forms/d/e/1FAIpQLSdOC3q1uWNlpVj_iPcz5G8rLrQRjuffNowE2U-DcUrKI3C_Aw/viewform?usp=sf_link

Password: USP4030

Author Contributions: F.E.G.G. and M.S. wrote the main manuscript text and prepared figures. F.E.G.G., R.W.B., V. C. and M.S. designed and performed the biochemical, immunological and microscopy experiments. R.S.D., E.F.C., M.S. and L.M.R.J. designed and performed the viral assays. C. B. performed the small molecule synthesis and performed the bioconjugation experiments. F.E.G.G. supervised the study. All authors reviewed the manuscript.

Funding: The authors acknowledge the support provided by: FAPESP (Sao Paulo Research Foundation) –grant numbers: 2017/10910-5 (M. Sadraeian - Pós-Doutorado - Fluxo Contínuo), 2013/07276-1(CEPOF – CEPID Program), 2009/54035-4 (EMU). The reagents related to HIV were supported by NIH AIDS Research and Reference Reagent Program.

Acknowledgements: The authors acknowledge the support provided by: FAPESP (Sao Paulo Research Foundation) –grant numbers: 2017/10910-5 (M. Sadraeian - Pós-Doutorado - Fluxo Contínuo), 2013/07276-1(CEPOF – CEPID Program), 2009/54035-4 (EMU). We also thank NIH AIDS Research and Reference Reagent Program for supporting the reagent related to HIV. We thank Seth Pincus, M.D. for his supports in the Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT. We thank Juliana Galinskas (Unifesp) and Bruno Andrade Ono (IFSC) for their valuable expertise in quantification assay and fluorescence microscopy, respectively.

Conflicts of interest: V.C. is a Director of ThioLogics. Other authors declare no competing interests.

Abbreviations

DAR: Drug-Antibody-Ratio

MAB: Monoclonal Antibody

PDT: Photodynamic Therapy

PIT: Photoimmunotherapy

PIC: Photo-immunoconjugate

Additional Information: Supplementary information accompanies this paper at
<http://www.mdpi.com/>

References:

1. Caskey M, Klein F, Nussenzweig MC. Broadly neutralizing anti-HIV-1 monoclonal antibodies in the clinic. *Nat Med* [Internet]. 2019;25(4):547–53.
2. Polito L, Bortolotti M, Battelli MG, Calafato G, Bolognesi A. Ricin: An ancient story for a timeless plant toxin. *Toxins (Basel)*. 2019;11(6):1–16.
3. Larson SM, Carrasquillo JA, Cheung NK V., Press OW. Radioimmunotherapy of human tumours. *Nat Rev Cancer* [Internet]. 2015;15(6):347–60.
4. Helal M, Dadachova E. Radioimmunotherapy as a Novel Approach in HIV, Bacterial, and Fungal Infectious Diseases. *Cancer Biother Radiopharm*. 2018;33(8):330–5.
5. Ventola CL. Cancer Immunotherapy: Challenges and Future Trends. *P&T*. 2017;42(8):514–21.
6. Nsole Biteghe FA, Toung Chalomie NE, Mungra N, Vignaux G, Gao N, Vergeade A, et al. Antibody-based immunotherapy: Alternative approaches for the treatment of metastatic melanoma. *Biomedicines*. 2020;8(9):327.
7. Ferrari G, Haynes BF, Koenig S, Nordstrom JL, Margolis DM, Tomaras GD. Envelope-

- specific antibodies and antibody-derived molecules for treating and curing HIV infection. Nat Rev Drug Discov [Internet]. 2016; 15(12):823-34.
8. Tsukrov D, Dadachova E. The potential of radioimmunotherapy as a new hope for HIV patients. Expert Rev Clin Immunol [Internet]. 2014;10(5):553–5.
 9. Pincus SH, Song K, Maresh GA, Hamer DH, Dimitrov DS, Chen W, et al. Identification of Human Anti-HIV gp160 Monoclonal Antibodies That Make Effective Immunotoxins. J Virol [Internet]. 2017;91(3):JVI.01955-16.
 10. Sadraeian M. Production and Characterization of Pulchellin A chain conjugated to HIV mAbs, and study its selective cytotoxicity against cells expressing HIV envelope [Internet]. Vol. 01, São Carlos: , Instituto de Física de São Carlos. [São Carlos]: Doctoral dissertation, Universidade de São Paulo; 2017.
 11. Sadraeian, Mohammad. Francisco E. G. Guimarães, Ana P. U. Araújo, David K. Worthylake, Louis LeCour Jr SHP. Selective cytotoxicity of a novel immunotoxin based on pulchellin A chain for cells expressing expressing HIV envelope. Sci Rep. 2017;7(1):1-12.
 12. Sadraeian M, Khoshnood Mansoorkhani MJ, Mohkam M, Rasoul-Amini S, Hesarak M, Ghasemi Y. Prevention and Inhibition of TC-1 Cell Growth in Tumor Bearing Mice by HPV16 E7 Protein in Fusion with Shiga Toxin B-Subunit from shigella dysenteriae. Cell J [Internet]. 2013;15(2):176–81.
 13. Med AN, Hanaoka YHH. Applying near-infrared photoimmunotherapy to B-cell lymphoma : comparative evaluation with radioimmunotherapy in tumor xenografts. Ann Nucl Med. 2017; 31(9):669-77.
 14. Sandland J, Boyle RW. Photosensitizer Antibody-Drug Conjugates: Past, Present, and

- Future. *Bioconjug Chem.* 2019;30(4):975–93.
15. Pincus SH, Song K, Maresh GA, Frank A, Worthylake D, Chung H, et al. Design and In Vivo Characterization of Immunoconjugates Targeting HIV gp160. *J Virol.* 2017;91(3):1–15.
 16. Craig RB, Summa CM, Corti M, Pincus SH. Anti-HIV Double Variable Domain Immunoglobulins Binding Both gp41 and gp120 for Targeted Delivery of Immunoconjugates. *PLoS One.* 2012;7(10):1–13.
 17. Buzzá HH, Silva AP Da, Vollet Filho JD, Ramirez DP, Trujillo JR, Inada NM, et al. Photodynamic therapy: Progress toward a scientific and clinical network in Latin America. *Photodiagnosis Photodyn Ther [Internet].* 2016;13:261–6.
 18. Wainwright M, O’Kane C, Rawthore S. Phenothiazinium photosensitisers XI. Improved toluidine blue photoantimicrobials. *J Photochem Photobiol B Biol [Internet].* 2016;160:68–71.
 19. Auwärter W, Écija D, Klappenberger F, Barth J V. Porphyrins at interfaces. *Nat Chem [Internet].* 2015;7(2):105–20.
 20. Fujita AKL, Rodrigues PGS, Requena MB, Escobar A, da Rocha RW, Nardi AB De, et al. Fluorescence evaluations for porphyrin formation during topical PDT using ALA and methyl-ALA mixtures in pig skin models. *Photodiagnosis Photodyn Ther [Internet].* 2016;15:236–44.
 21. Valkov A, Nakonechny F, Nisnevitch M. Polymer-immobilized photosensitizers for continuous eradication of bacteria. *Int J Mol Sci.* 2014;15(9):14984–96.
 22. Nakonechny F, Barel M, David A, Koretz S, Litvak B, Ragozin E, et al. Dark antibacterial

- activity of rose Bengal. *Int J Mol Sci.* 2019;20(13):3196.
23. Dougherty TJ, Gomer CJ, Henderson BW, Jori G, Kessel D. Photodynamic Therapy. *J Natl Cancer Inst.* 1998;17(9):889–905.
 24. Bacellar IOL, Tsubone TM, Pavani C, Baptista MS. Photodynamic efficiency: From molecular photochemistry to cell death. *Int J Mol Sci.* 2015;16(9):20523–59.
 25. Crous A, Abrahamse H. Effective gold nanoparticle-antibody-mediated drug delivery for photodynamic therapy of lung cancer stem cells. *Int J Mol Sci.* 2020;21(11):3742.
 26. Sandland J, Boyle RW. Photosensitizer Antibody-Drug Conjugates: Past, Present, and Future. *Bioconjug Chem.* 2019;30(4):975–93.
 27. Caudle AS, Yang WT, Mittendorf E a, Kuerer HM. Cancer Cell-Selective In Vivo Near Infrared Photoimmunotherapy Targeting Specific Membrane Molecules. *Nat Med.* 2016;150(2):137–43.
 28. Bryden F, Maruani A, Savoie H, Chudasama V, Smith MEB, Caddick S, et al. Regioselective and stoichiometrically controlled conjugation of photodynamic sensitizers to a HER2 targeting antibody fragment. *Bioconjug Chem.* 2014;25(3):611–7.
 29. Maruani A, Smith MEB, Miranda E, Chester K a, Chudasama V, Caddick S. A plug-and-play approach to antibody-based therapeutics via a chemoselective dual click strategy. *Nat Commun [Internet].* 2015;31;6(1):1-9.
 30. Bahou C, Love EA, Leonard S, Spears RJ, Maruani A, Armour K, et al. Disulfide Modified IgG1: An Investigation of Biophysical Profile and Clinically Relevant Fc Interactions. *Bioconjug Chem.* 2019;30(4):1048–54.
 31. Chester KA, Baker R, Caddick S. Next generation maleimides enable the controlled

- assembly. *R Soc Chem*. 2014;10(5):1–52.
32. Jackson DY. Processes for Constructing Homogeneous Antibody Drug Conjugates. *Org Process Res Dev*. 2016;20(5):852–66.
33. Bryden F, Boyle RW. A mild, facile, one-pot synthesis of zinc azido porphyrins as substrates for use in click chemistry. *Synlett*. 2013;24(15):1978–82.
34. Cheng MHY, Maruani A, Savoie H, Chudasama V, Boyle RW. Synthesis of a novel HER2 targeted aza-BODIPY-antibody conjugate: Synthesis, photophysical characterisation and: In vitro evaluation. *Org Biomol Chem*. 2018;16(7):1144–9.
35. Lee MTW, Maruani A, Richards D a., Baker JR, Caddick S, Chudasama V, et al. Enabling the controlled assembly of antibody conjugates with a loading of two modules without antibody engineering. *Chem Sci [Internet]*. 2017;8(3):2056-60.
36. Castañeda L, Wright ZVF, Marculescu C, Tran TM, Chudasama V, Maruani A, et al. A mild synthesis of N-functionalised bromomaleimides, thiomaleimides and bromopyridazinediones. *Tetrahedron Lett*. 2013;54(27):3493–5.
37. Robinson E, Nunes JPM, Vassileva V, Maruani A, Nogueira JCF, Smith MEB, et al. Pyridazinediones deliver potent, stable, targeted and efficacious antibody-drug conjugates (ADCs) with a controlled loading of 4 drugs per antibody. *RSC Adv*. 2017;7(15):9073–7.
38. Bryden F, Maruani A, Rodrigues JMM, Cheng MHY, Savoie H, Beeby A, et al. Assembly of High-Potency Photosensitizer-Antibody Conjugates through Application of Dendron Multiplier Technology. *Bioconjug Chem*. 2018;29(1):176–81.
39. Pincus SH, Fang H, Wilkinson R a, Marcotte TK, Robinson JE, Olson WC. In vivo efficacy

- of anti-glycoprotein 41, but not anti-glycoprotein 120, immunotoxins in a mouse model of HIV infection. *J Immunol.* 2003;170(4):2236–41.
40. Kobayashi H, Choyke PL. Target-Cancer-Cell-Specific Activatable Fluorescence Imaging Probes: Rational Design and in Vivo Applications. *Acc Chem Res* [Internet]. 2011 Feb 15;44(2):83–90.
 41. Kratochvil HT, Ha DG, Zanni MT. Counting tagged molecules one by one: Quantitative photoactivation and bleaching of photoactivatable fluorophores. *J Chem Phys.* 2015;143(10):104201.
 42. Fujimura D, Inagaki F, Okada R, Rosenberg A, Furusawa A, Choyke PL, et al. Conjugation Ratio, Light Dose, and pH Affect the Stability of Panitumumab–IR700 for Near-Infrared Photoimmunotherapy. *ACS Med Chem Lett* [Internet]. 2020;11(8):1598-604.
 43. Kovacs JM, Noeldeke E, Ha HJ, Peng H, Rits-Volloch S, Harrison SC, et al. Stable, uncleaved HIV-1 envelope glycoprotein gp140 forms a tightly folded trimer with a native-like structure. *Proc Natl Acad Sci U S A* [Internet]. 2014;111(52):18542–7.
 44. Pincus SH, McClure J. Soluble CD4 enhances the efficacy of immunotoxins directed against gp41 of the human immunodeficiency virus. *Proc Natl Acad Sci U S A.* 1993;90(1):332–6.
 45. Mchugh L, Hu S, Lee BK, Santora K, Kennedy PE, Berger EA, et al. Increased Affinity and Stability of an Anti-HIV-1 Envelope Immunotoxin by Structure-based Mutagenesis *. *J Biol Chem.* 2002;277(37):34383–90.
 46. Giuntini F, Bryden F, Daly R, Scanlan EM, Boyle RW. Huisgen-based conjugation of water-soluble porphyrins to deprotected sugars: towards mild strategies for the labelling

- of glycans. *Org Biomol Chem* [Internet]. 2014;12(8):1203–6.
47. Bahou C, Richards DA, Maruani A, Love EA, Javaid F, Caddick S, et al. Highly homogeneous antibody modification through optimisation of the synthesis and conjugation of functionalised dibromopyridazinediones. *Org Biomol Chem* [Internet]. 2018;16(8):1359–66.
48. Inada NM, da Costa MM, Guimarães OCC, da Silva Ribeiro E, Kurachi C, Quintana SM, et al. Photodiagnosis and treatment of condyloma acuminatum using 5-aminolevulinic acid and homemade devices. *Photodiagnosis Photodyn Ther* [Internet]. 2012;9(1):60–8.
49. Romano R a, Silva AP, Kurachi C, Guimara FEG. Light-driven photosensitizer uptake increases *Candida albicans* photodynamic inactivation. 2017;9:1–10.
50. Mello BL, Alessi AM, Riaño-Pachón DM, DeAzevedo ER, Guimarães FEG, Espírito Santo MC, et al. Targeted metatranscriptomics of compost-derived consortia reveals a GH11 exerting an unusual exo-1,4- β -xylanase activity. *Biotechnol Biofuels*. 2017;10(1):1–17.