

An Introduction to the *Callithrix* Genus and Overview of Recent Advances in Marmoset Research

AUTHORS

Joanna Malukiewicz is a Marie Skłodowska-Curie Post-Doctoral Fellow at Primate Genetics Laboratory, German Primate Research Center, Leibniz Institute for Primate Research, Goettingen, Germany and an associate post-doctoral research at the Faculty of Medicine, University of São Paulo, São Paulo, Brazil.

Vanner Boere is an associate professor at the Institute of Humanities, Arts, and Sciences at the Federal University of Southern Bahia, Itabuna, Bahia, Brazil.

Maria Adelia Borstelmann de Oliveira is a professor in the Department of Animal Morphology and Physiology at the Federal Rural University of Pernambuco, Recife, Pernambuco, Brazil.

Mirela D’Arc is a postdoctoral researcher at the Viral Diversity and Diseases Laboratory, Department of Genetics, Health Science Center, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil.

Jessica Viviane Amorim Ferreira is a researcher at that Centro de Conservação e Manejo de Fauna da Caatinga, Universidade Federal do Vale do São Francisco, Petrolina, Pernambuco, Brazil.

Jeffery French is a professor in the Department of Psychology at the University of Nebraska Omaha, Omaha, Nebraska, USA.

Genevieve Housman is a postdoctoral fellow in Genetic Medicine at the University of Chicago, Chicago, Illinois, USA.

Claudia Almeida Igayara de Souza is a medical veterinarian at the Guarulhos Municipal Zoo, Guarulhos, São Paulo, São Paulo, Brazil.

Leandro Jerusalinsky is the director the Center for Research and Conservation of Brazilian Primates under the Chico Mendes Institute for Biodiversity Conservation, João Pessoa, Paraíba, Brazil.

Fabiano Rodrigues de Melo is an associate professor at Department of Forest Engineering at the Federal University of Viçosa, Viçosa, Minas Gerais, Brazil.

Mônica Mafra Valença-Montenegro is an environmental analyst at the Center for Research and Conservation of Brazilian Primates under the Chico Mendes Institute for Biodiversity Conservation, João Pessoa, Paraíba, Brazil.

Silvia Bahadian Moreira is a medical veterinarian at the Centro de Primatologia do Rio de Janeiro, Guapimirim, Rio de Janeiro, Rio de Janeiro, Brazil.

Ita de Oliveira e Silva is an associate professor at the Institute of Humanities, Arts, and Sciences at the Federal University of Southern Bahia, Itabuna, Bahia, Brazil.

Felipe Santos Pacheco is a doctoral student in the Postgraduate Program in Animal Biology at the Federal University of Viçosa, Viçosa, Minas Gerais, Brazil.

Jeff Rogers is an associate professor of molecular and human genetics at Baylor College of Medicine, Houston, Texas, USA.

Alcides Pissinatti is the director of the Centro de Primatologia do Rio de Janeiro, Guapimirim, Rio de Janeiro, Rio de Janeiro, Brazil.

Ricardo del Rosario is a computational biologist at the Broad Institute, Cambridge, Massachusetts, USA.

Corinna Ross is associate professor of biology at Texas A&M San Antonio and population health at the Texas Biomedical Research Institute, San Antonio, Texas, USA.

Carlos R. Ruiz-Miranda is an associate professor at the Laboratory of Environmental Sciences, Center for Biosciences and Biotechnology at Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, Rio de Janeiro, Brazil.

Luiz C.M. Pereira is the director of Centro de Conservação e Manejo de Fauna da Caatinga, Universidade Federal do Vale do São Francisco, Petrolina, Pernambuco, Brazil.

Nicola Schiel is an associate professor in the Post-Graduate Programs in Ecology as well as Ethnobiology and Nature Conservation at the Federal Rural University of Pernambuco, Recife, Pernambuco, Brazil.

Fernanda de Fátima Rodrigues da Silva is an affiliate researcher at the Centro de Conservação dos Saguís-da-Serra at Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil.

Antonio Souto is a full professor at the Laboratory of Ethology, Department of Zoology, Federal University of Pernambuco, Recife, Brazil.

Vedrana Šlipogor is a postdoctoral research fellow at the Department of Behavioral and Cognitive Biology, University of Vienna, Austria.

Suzette Tardif is professor emeritus at the Southwest National Primate Research Center, Texas Biomedical Research Institute, San Antonio, Texas, USA.

*Address correspondence to Joanna Malukiewicz jmalukie@gmail.com

Primate Genetics Laboratory, German Primate Research Center, Leibniz Institute for Primate Research, 37077 Goettingen, Germany

ABSTRACT

We provide here a current overview of marmoset (*Callithrix*) evolution, hybridization, species biology, basic/biomedical research, and conservation initiatives. Composed of two subgroups, the *aurita* group (*C. aurita* and *C. flaviceps*) and the *jacchus* group (*C. geoffroyi*, *C. jacchus*, *C. kuhlii* and *C. penicillata*), this relatively young primate radiation is endemic to the Brazilian Cerrado, Caatinga, and Atlantic Forest biomes. Significant impacts on *Callithrix* within these biomes resulting from anthropogenic activity include: (1) population declines, particularly for the *aurita* group; (2) widespread geographic displacement, biological invasions, and range expansions of *C. jacchus* and *C. penicillata*; (3) anthropogenic hybridization; and (4) epizootic Yellow Fever and Zika viral outbreaks. A number of Brazilian legal and conservation initiatives are now in place to protect the threatened *aurita* group and increase research into them. Due to their small size and fast life history, marmosets are prized biomedical models. As a result, there are increasingly sophisticated genomic *Callithrix* resources available and burgeoning marmoset functional, immuno-, and epi- genomic research. In both the laboratory and the wild, marmosets have given us insight into cognition, social group dynamics, human disease, and pregnancy. *Callithrix jacchus* and *C. penicillata* are emerging Neotropical primate models for arbovirus disease, including Dengue and Zika. Wild marmoset populations are helping us understand sylvatic transmission and human spillover of Zika and Yellow Fever viruses. All of these factors are positioning marmosets as preeminent models to facilitate understanding of facets of evolution, hybridization, conservation, human disease, and emerging infectious diseases.

Keywords: marmoset, Neotropical, Brazil, biomedical, arbovirus, conservation, hybridization, biological invasion

BACKGROUND

The small, charismatic Brazilian monkeys that compose the *Callithrix* genus have special relevance in biomedicine, evolutionary/hybridization studies, and burgeoning conservation work. In this chapter, we first give a general overview of the six *Callithrix* species by discussing their biogeography, evolutionary relationships, and demographic history. Then we focus on unique behavioral, ecological, demographic, and/or conservation aspects of individual species. The final section of this chapter summarizes a number of recent, novel, and neglected aspects of captive and wild *Callithrix* research. Specifically, we discuss: (1) unique aspects of *Callithrix* biology relative to biomedical research; (2) recent comparative, functional, and population *Callithrix* genomics work; (3) biomedical relevance of *Callithrix* in viral studies, *Callithrix* susceptibility to viral disease, and the *Callithrix* virosphere; (4) natural and anthropogenic *Callithrix* hybridization; (5) legislative and conservation aspects of *Callithrix* population management; and (6) anthropogenic impacts on *Callithrix* populations. We conclude this chapter by identifying new avenues for research with wild and captive *Callithrix*.

AN OVERVIEW of the *CALLITHRIX* GENUS

CALLITHRIX BIOGEOGRAPHY and EVOLUTIONARY HISTORY

The *Callithrix* genus is composed of six naturally allo- and parapatric species (Figure 1) endemic to Brazil [1]. Phylogenetic analysis of mitogenomes (Figure 2) shows that this genus diverged from the Amazonian *Cebuella/Mico* clade approximately 6.83 million years ago (MYA) (Node E, Figure 2) [2]. Then the genus separated into two subgroups, the *aurita* group (*C. aurita* and *C. flaviceps*) and *jacchus* group (*C. geoffroyi*, *C. jacchus*, *C. kuhlii*, *C. penicillata*) approximately 3.54 MYA (Node D, Figure 2). Within the *jacchus* group, *C. geoffroyi* represents the most basal lineage, having arisen




Figure 1. Maps of natural Brazilian *Callithrix* ranges following [1], and natural hybrid zones [21].

Each range is color-coded by species and respective hybrid zones are shown as points. Abbreviations are Brazilian states- AL (Alagoas); BA (Bahia); ES (Espírito Santo); CE (Ceará); GO (Goiás); MA (Maranhão); MG (Minas Gerais); PB (Paraíba); PI (Piauí); PE (Pernambuco); PR (Paraná); RJ (Rio de Janeiro); RN (Rio Grande do Norte); SE (Sergipe); SP (São Paulo); TO (Tocantins). Hybrid zone abbreviations: JP: *C. jacchus* x *C. penicillata*; AF: *C. aurita* x *C. flaviceps*; FG: *C. flaviceps* x *C. geoffroyi*; PG: *C. geoffroyi* x *C. penicillata*; PK: *C. penicillata* x *C. kuhlii*.




Figure 2. BEAST Bayesian phylogeny showing divergence patterns and temporal divergence estimates between *Callithrix* mitogenomic lineages, modified from [2]. Boxed letters represent major divergence nodes among *Callithrix* clades. ‘Ma’ refers to millions of years.

approximately 1.18 MYA (Node C, Figure 2). Mitochondrial (mtDNA) phylogenies show *C. kuhlii* and *C. penicillata* clades as polyphyletic, likely due to incomplete lineage sorting [2]. Finally, the *C. jacchus* clade arose approximately 0.5 MYA as the sister clade to one of the polyphyletic *C. penicillata* clades (Node A, Figure 2).

Callithrix marmosets likely originated within the southeastern Atlantic Forest, after a vicariant event split them from the Amazonian marmosets, and then spread northward along the coast and into the Brazilian interior [3]. *Callithrix aurita*, *C. flaviceps*, *C. geoffroyi*, and *C. kuhlii* are endemic to the Atlantic Forest biome [1, 3]. However, *C. geoffroyi* is also found in the semi-arid regions of northeastern Minas Gerais and southern Bahia [1]. *Callithrix jacchus* can be found in the Atlantic Forest and the semi-arid Caatinga biome of northeastern Brazil [1]. *Callithrix penicillata* mostly occurs in the semi-arid Cerrado of central and central-western Brazil [1], but also is found in the Caatinga [4, pers obs. JM, LCP].

CALLITHRIX NATURAL GEOGRAPHIC RANGES

Callithrix aurita, or the buffy-tufted-ear marmoset, inhabits submontane and montane forests of the Atlantic Forest biome at altitudes above 500 m [5]. Environmental factors such as altitude, climate and phytophysiology may play key roles in determining *C. aurita*-suitable habitats [6,7]. The northwestern limits of the *C. aurita* range reach transitional regions between the Atlantic Forest and Cerrado in São Paulo and Minas Gerais states [1,5]. Hershkovitz [8] placed the northeastern limit of the species' distribution in Minas Gerais state at the right bank of the Muriaé River. However, [9] noted that the species also inhabits the region north of the Rio Doce State Park, MG, but does not exceed the northern limits of the Doce river. The southern limit of *C. aurita*'s range is undefined, but may occur at the intersection of the Tietê and Piracicaba rivers in São Paulo state [10].

With a geographical distribution of 30,815 km², *Callithrix flaviceps*, or the buffy-headed marmoset, possesses the smallest *Callithrix* geographic distribution [11]. This species occurs in

mountainous regions of Espírito Santo and Minas Gerais at altitudes up to at least 1,200 m [11]. Its range begins south of the Doce River [5] and continues into the Serra da Mantiqueira in the south of Espírito Santo where this state borders Rio de Janeiro state [1]. However, the *C. flaviceps* range may reach to the far north of Rio de Janeiro state [1,12]. An unconfirmed *Callithrix flaviceps* occurrence was reported in the municipality of Varre-Sai, northern Rio de Janeiro state [13].

The northern distribution limit of *Callithrix kuhlii*, or Wied's marmoset, occurs at the Contas river in Bahia state and the southern limit is the Jequitinhonha River in Minas Gerais state [1]. There are recent reports of *C. kuhlii* to the north beyond the Rio de Contas, in the region of Valença, Bahia [12], and within the vicinity of Todos os Santos Bay, Bahia [14, pers. obs. VB and IOS]. The eastern limit of *C. kuhlii*'s distribution reaches the Atlantic Ocean, and the Cerrado likely terminates the species' natural distribution to the west [15].

Callithrix geoffroyi, or Geoffroy's tufted-ear marmoset, generally occurs in the Atlantic Forest at altitudes below 500-700 m [1]. The southern bank of the Jequitinhonha River and eastern bank of the Araçuaí river represent upper limits of the *C. geoffroyi* range [1]. *Callithrix geoffroyi* occurrences have also been recorded in the Espinhaço Mountains, Minas Gerais (see Figure 1), which are located at the western Atlantic Forest limit [16]. Additionally, the species has been observed in the region between the Jequitinhonha and Doce rivers, in the north and northeast of Minas Gerais state, but it is not clear whether *C. geoffroyi* is introduced there [15,17]. While *C. geoffroyi* is considered a lowland species, it has been observed in the Serra do Cipó National Park, Minas Gerais state at 1274 m above sea level [1,12]. The species has also been observed in a semi-arid region of northeastern Minas Gerais state on the western bank of the Araçuaí River in Berilo municipality [pers. obs., JM].

Callithrix penicillata, or the black-tufted marmoset, has the widest *Callithrix* geographic distribution, with an area of about 1,300,000 km² [15,16,18]. The northern *C. penicillata* range seems limited by the Grande and São Francisco rivers [15,16]. The eastern limits of the *C. penicillata* range seem to be bounded by the Araguaia River's course through Goiás, Mato Grosso and Tocantins states. This species can also be found in the north of São Paulo, north of the Tietê and




Figure 3. Light blue and darker blue regions respectively represent natural *C. jacchus* and *C. penicillata* ranges [1]. Maroon and black dots respectively represent unpublished *C. jacchus* and *C. penicillata* occurrences observed by LCMP and JM. State abbreviations follow Figure 1.

Piracicaba rivers [8], as well as eastern Minas Gerais and southern Bahia, where *C. penicillata* has contact zones with *C. geoffroyi*, *C. aurita*, *C. flaviceps* and probably *C. kuhlii* [1,15,16].

For the currently recognized *C. penicillata* distribution, several unfilled gaps still exist in the occurrence of this species, particularly within the Caatinga biome. Although the southern portion of the Caatinga is recognized by [1] as part of the natural distribution of *C. penicillata*, we have observed occurrences of *C. penicillata* in northern parts of the Caatinga (Figure 3) (pers. obs. LCMP and JM). The dataset of [19] lists several occurrences of *C. penicillata* as ‘alien’ in the northern Caatinga where no *Callithrix* are thought to occur naturally. We, however, would like to note that the Caatinga *sensu stricto* is one of the youngest Brazilian vegetation formations, originating about 15-10 years ago MYA [20]. Further the *C. penicillata* Caatinga clade and the *C. jacchus* clades represent the two youngest clades within the phylogeny shown in Figure 2 [2]. Thus, these occurrences of *C. penicillata* in the Caatinga may be part of a natural expansion from the Cerrado and into the Caatinga.

Callithrix jacchus, or the white-tufted marmoset, occurs in northeastern Brazil [1]. The southern portion of its geographic distribution extends to the northern bank of the São Francisco River and the west bank of the Rio Grande [1]. Whereas the northern and eastern portion of the *C. jacchus* natural distribution is limited by the Brazilian Atlantic coast, the species’ western limits are less clear [1]. Currently recognized limits of the natural *C. jacchus* distribution are at the boundary between Tocantins and Maranhão states, but the species may also occur in the northeast of Tocantins [1].

NATURAL *CALLITHRIX* HYBRIDIZATION

Callithrix species are a relatively recent primate radiation, and secondary contact between *Callithrix* species at the borders of their natural geographical distributions can result in natural hybridization (Figure 1). One natural hybrid zone exists in the mountains of Espírito Santo state where hybrids of *C. flaviceps* and *C. geoffroyi* occur in an area of overlap between altitude limits for each parental species [5]. A number of natural hybrid zones occur at major river boundaries between *C. penicillata*’s natural distribution and that of other congeners (Figure 1) [21]. *Callithrix aurita* and *C.*

flaviceps hybridize naturally in a region that roughly overlaps the borders of Minas Gerais, Rio de Janeiro, and Espírito Santo states. Climatic factors that shape *C. flaviceps*' natural distribution and past climatic fluctuations may determine locations of natural *C. aurita* and *C. flaviceps* hybrid zones [22]. We have observed natural *C. jacchus* and *C. penicillata* hybrids along the banks of the São Francisco River between Petrolina, PE and Juazeiro, BA [4,21] as well as in Xique-Xique, BA (pers. obs., LCMP). Natural hybrids between the two species have also been observed at the northern tip of the Espinhaço mountain range where there is a mixture of Cerrado, Caatinga, and semi-deciduous forests (pers. obs. LCMP).

CHARACTERISTICS and DIFFERENCES between *CALLITHRIX* SPECIES

CALLITHRIX AURITA

The total population size adult *C. aurita* (Figure 4) is estimated to be 10,000 individuals, and the species is currently listed as endangered [10]. *Callithrix aurita* social groups usually possess a single male-female breeding pair, although groups with multiple breeding females have been noted [23-24]. Group size varies from 5-11 individuals [7, 25]. Population density for *C. aurita* also varies from 2.8 individuals/km² in the Serra do Brigadeiro State Park, MG [26] to 14.76 individuals/km² in the Pouso Alegre Municipal Natural Park, MG [27]. The home range size for a *C. aurita* group is approximately 11 hectares [28]. The suitability of a given habitat for *C. aurita*, considering topography and forest composition, may ultimately determine population density within a given region [6]. While conducting linear transect surveys, [6] estimated a general population of 1892 individuals in an 250.7 km² area of Atlantic Forest in southern São Paulo state.

Several morphological differences distinguish *C. aurita* from other *Callithrix* species. *Callithrix aurita* is perhaps the largest *Callithrix* species considering body weight and overall body (Figure 5 and Table 1). *Callithrix aurita* shares morphological similarities with *C. flaviceps* [e.g., 29], likely due to the close evolutionary relationship between them [5]. For example, [29] noted a larger

brain case size in the *aurita* group, relative to the *jacchus* group, as well as retrusions of the upper and lower jaws. Differences in cranial morphology between *C. aurita* and *C. flaviceps* include greater lateral expansion of the parietal bone and elongation of the occipital bone in *C. aurita* but more lateral nasal compression and larger incisors in *C. flaviceps* [29].

Morphological differences between the *aurita* and *jacchus* groups are likely related to stronger morphological specialization for gummivory in the latter group [28, 29]. In a 17 ha forest fragment in Minas Gerais, *C. aurita* consumed 50.5% gums, 11% fruits, and 38.5% prey [30]. Gums are generally accessed by *C. aurita* from sources that do not require gouging [10]. The species also consumes fungi fruiting bodies from bamboo of the Poaceae family [10].

CALLITHRIX FLAVICEPS

Callithrix flaviceps (Figure 6) is one of the least studied *Callithrix* taxa. The species is listed as critically endangered [11], as a total of approximately 2000 adult *C. flaviceps* individuals likely remain in the wild [11]. In secondary forests, *C. flaviceps* occupies areas sized between 15 ha to 35.5 ha [31-33], but in denser forests, 138.5 ha home ranges have been observed [34]. These home range sizes are among the largest for any *Callithrix* species [32,33]. On average, *C. flaviceps* travels 1222.5 m per day, but food scarcity during the dry season likely causes the species to have longer daily travel than in the rainy season [7]. The size *C. flaviceps* social groups is between 3-15 members [31]. Although social groups tend to have one reproductive female, there occasionally are two or more reproductive females and several adult helpers that care for offspring [35].



Figure 4. Image of *C. aurita*. Photo credit: Orlando Vital.




Figure 5. Boxplots of body weight (A), tail length (B), body length (C), and intercranial length for select *Callithrix* species and hybrid types. Taxa abbreviations are A=*C. aurita*, CC= unidentified hybrid, J=*C. jacchus*, PG= *C. penicillata* x *C. geoffroyi* hybrid, AH=*C. aurita* hybrid, G= *C. geoffroyi*, P=*C. penicillata*, PJ=*C. penicillata* x *C. jacchus* hybrid. Modified from Malukiewicz et al. (unpublished data).

Table 1. Means of thirteen morphometric measures. Numbers in parentheses represent the number of samples for a given measure. Measure abbreviations are as follows: IC=intercranial distance, FO=cranial frontal-occipital distance, Zyg=distance between zygomatic arches, jaw= intermandibular distance. Taxa abbreviations follow Figure 4. Modified from Malukiewicz et al. (unpublished data). Taxon abbreviations follow Figure 5.

Taxa	Weight (g)	Tail (cm)	Body (cm)	IC (mm)	FO (mm)	Zyg (mm)	Jaw (mm)	Hand (mm)	Humerus (cm)	Forearm (cm)	Femur (cm)	Tibia (cm)	Foot (mm)
A	440.6 (25)	32.3 (26)	21.9 (27)	33.1 (27)	42.6 (27)	31.4 (23)	23.7 (23)	30.5 (21)	5.4 (26)	5.2 (27)	6.5 (27)	7.2 (27)	61.1 (24)
AH	408.1(9)	29.7 (9)	21.4 (9)	32.6 (9)	38.8 (9)	30.3 (9)	26.3 (9)	39.0 (9)	5.2 (9)	5.2 (9)	6.4 (9)	7.2 (9)	57.9 (9)
G	386.2 (14)	30.7 (14)	22.2 (14)	30.2 (14)	40.2 (14)	30.1 (14)	25.7 (14)	36.3 (13)	5.3 (14)	4.8 (14)	6.3 (14)	7.1 (14)	55.4 (14)
J	322.6 (30)	27.4 (30)	19.9 (29)	27.4 (29)	39.7 (24)	28.7 (29)	22.2 (29)	35.4 (28)	4.7 (29)	4.5 (29)	5.5 (29)	6.6 (29)	54.7 (27)
P	308.4 (54)	27.7 (54)	20.9 (52)	28.4 (54)	39.8 (50)	28.6 (51)	22.9 (52)	35.2 (48)	4.6 (54)	4.6 (54)	5.8 (54)	6.5 (54)	54. (54)2
PG	355.8 (18)	30.7 (18)	21.4 (18)	29.8 (18)	37.7 (18)	30.3 (18)	22.7 (18)	36.8 (9)	5.1 (18)	4.8 (18)	6.6 (18)	7.1 (18)	53.3 (16)
PJ	317.9 (53)	28.7 (53)	21.3 (54)	28.9 (54)	39.2 (49)	29.1 (53)	23.7 (53)	34.4 (50)	4.7 (54)	4.6 (54)	5.8 (54)	6.7 (54)	54.4 (54)

The *C. flaviceps* dentition, like that of *C. aurita*, is less specialized for gummivory than that of the *jacchus* group [36], and the species' consumption of fruits, insects and fungi varies seasonally [34]. *Callithrix flaviceps* uses a “scan and pounce” approach to foraging for small prey [37]. The

consumption of fungi is especially high in this species, and can compose up 65% of the total items consumed [34, 38]. Gum consumption by *C. flaviceps* is mostly opportunistic, as this species exploits gums from tree holes made by wood-digging insects [7].

CALLITHRIX GEOFFROYI

Callithrix geoffroyi is characterized by a white face and black voluminous ear tufts. The species appears to be the largest *jacchus* group for average body weight and size (Figure 5 and Table 1). One *C. geoffroyi* social group in an Atlantic Forest patch was composed of a single female, two males, and two juveniles [39]. The same group had a total home range size of 23.3 ha, but that varied between 4.9 ha in the rainy season and 7.2 ha in the dry season [39], which is likely related to distribution of nutritional resources. The group's daily ranging distances varied from 480 to 1980 meters [39]. In a 110 ha forest fragment in Espírito Santo, major food categories for one *C. geoffroyi* group were, on average, gums (68.6%) fruits (15%), and prey (invertebrates 14.6% and vertebrates 0.8%), although utilization of these items varied seasonal [40]. This same group preyed upon insects, arachnids, gastropods, lizards, and frogs [41]. Population densities for *C. geoffroyi* in northern Espírito Santo ranged from 0.7 groups/km² (Putiri) to 7.5 groups/km², generally being higher in larger fragments (approximately 20,000 ha) than in smaller fragments (approximately 200 ha) [41].

In general, *C. geoffroyi* seems highly flexible in the types of habitats that it can occupy. Forested environments occupied by *C. geoffroyi* include dense and semi-deciduous forests of the Atlantic Forest as well as deciduous Caatinga forests [15]. The species is tolerant of secondary habitats, for example being able to utilize forest fragments surrounded by eucalyptus monoculture



Figure 6. Image of *C. flaviceps*. Photo credit: Sarisha Trindade.

CALLITHRIX JACCHUS

Callithrix jacchus (Figure 7) is one of the smallest *Callithrix* species (Table 1, Figure 5), and several of its biological traits seem associate with a high level of specialization for exudivory [28]. Exudivory adaptations of *C. jacchus* include chewing muscles for production of relatively large tree gouging gapes [42] and an enlarged caecum as part of a digestive system that optimizes digestion of tree gums [43]. Further, [29] attributed the compressed brain case and protruding zygomatic arches of the *jacchus* group as tree gouging adaptations. *Callithrix jacchus* and its sister species, *C. penicillata*, also share derived dentition patterns [36], that also likely represent exudivory adaptations.

Few *Callithrix* species have home ranges as small as those of *Callithrix*. In comparing Atlantic Forest living marmosets and Caatinga living marmosets, the former had a mean group composition of 8.3 individuals while Caatinga mean group size was 5.9 [44]. When comparing the animals' home range between these two environments, Atlantic forest living animals had a slightly smaller mean home range (4.1 ha) than Caatinga living animals (mean 5.3 ha) [44]. Castro and Araújo [45] found that *C. jacchus* feed on fruit, gums, and invertebrates, but fruit and gum consumption were negatively related and influenced by seasonality. In contrast, [46] did not observe a seasonal difference in gum consumption in Caatinga living individuals. However, insect consumption was significantly higher in the rainy season.

Common marmosets are highly adaptive [44,46,47] and their inhabitation of the Caatinga represents an ecological challenge related to heat stress and limited resources like water [48]. The species' adaptability warrants the question as to what makes their ecological success possible. The species' funnel-shaped teeth and ability to cling vertically on tree trunks that facilitates exudate extraction [49]. Having a cooperative breeding system and production of two litters per year [50] coupled with complex cognitive abilities [51] certainly contribute to *C. jacchus* ecological success. However, behavioural adjustments might be the key factor for navigating the challenges of the




Figure 7. Image of *C. jacchus*. Photo Credit: Antonio Souto.

Caatinga [46, 47]. Behavioral modification for dealing with heat stress can be observed by Caatinga common marmosets who rest twice as much as Atlantic Forest marmosets [47], live in smaller groups and appear to have a higher ratio of surface area to body mass [49]. Added to that Caatinga marmosets eat cactus, which requires cognitive skills to access the resource while avoiding injury [46].

CALLITHRIX KUHLII

The home range of *C. kuhlii* varies from 10 to 58.3 ha [52], and daily ranging distance is estimated to be as far as 1498 m, considered the longest daily among *Callithrix* species [53]. In this species, group size averages 4.3 individuals, with 1 breeding female and 1–2 adult males [52]. The diet of Wied's marmoset is heavily fruit- and nectar-based, with invertebrates and small vertebrates complementing the diet [53]. In many regions, *C. kuhlii* co-occurs with golden-headed lion tamarins (*Leontopithecus chrysomelas*) although it is not clear whether there is a niche overlap of the two species [52].

Callithrix kuhlii utilizes various habitats that include restinga forest, riverside forests, secondary forests, mangroves, fruit groves, coconut and palm oil palm trees [52]. In the region of Itabuna and Ilhéus, Bahia state, the species also occupies old growth forests where cocoa is cultivated [52]. The species has a preference for occupying degraded areas, close to wood edges [52]. The species is common in urban areas of Southern Bahia state, where social groups can be seen in backyards, orchards and public squares [54]. Urban *C. kuhlii* groups travel along poles, electrical wires, and houses [54]. At dusk the animals retreat to clumps of bromeliads and vines which is believed to be an anti-predatory and thermoregulatory strategy [52].

CALLITHRIX PENICILLATA

Like its sister species, *C. penicillata* is among the smallest *Callithrix* species (Table 1, Figure 5). The home range of *C. penicillata* varies between 2.0 ha [55] to 18.5 ha [56], and social groups are composed of 2-19 individuals [28, 57]. *Callithrix penicillata* exploits many sources of food such as fruit, buds, flowers, leaves, young stems, small prey, and bird eggs [28,56]. Additionally, this species also heavily exploits tree gums, which seems to give a great advantage in adapting to degraded environments and secondary forests [28]. Given the close evolutionary relationship between *C. jacchus* and *C. penicillata* [2], the latter likely possesses similar adaptations for gummivory as the former.

Cosmopolitan *C. penicillata* populations have significantly expanded their ranges beyond their natural distribution [2], likely in part due to illegal pet trafficking of the species between northeastern and southeastern Brazil [58]. The species has been introduced to portions of Bahia, Espírito Santo, Minas Gerais, Rio de Janeiro, Paraná, Santa Catarina, São Paulo, and Rio Grande do Sul states [58, 59, pers obs. JM, VB, IOS]. Mathematical models predict that the expansion of *C. penicillata* may be greater than expected, and that this species may even replace the species of marmosets native to areas of the Atlantic Forest [59-61].

While deforestation threatens other marmoset species, *C. penicillata* may have instead benefited through the opening of new niches favoring a population expansion of the species [28]. Additionally, *C. penicillata* shows great flexibility in occupying altered areas and secondary vegetation [62]. Other anthropogenic factors that may favor the range expansion of *C. penicillata*, especially within urban areas, include planting of exotic trees and reductions of predators such as constrictor snakes, birds of prey and wild cats.

A SUMMARY of RECENT *CALLITHRIX* RESEARCH

RELEVANCE of *CALLITHRIX* BIOLOGICAL TRAITS in BIOMEDICAL RESEARCH

REPRODUCTION and GROWTH PATTERNS in CALLITHRIX

Small body size and a fast life history, characteristics that make marmosets prized primate biomedical models (e.g., pediatric obesity [63]; Zika infection during pregnancy [64], are related to the evolution of *Callithrix* reproductive function. Four marmoset traits related to size and life history are likely inextricably linked: reproductive suppression, miniaturization, production of litters, and chimerism. However, it is important to note that, as biomedical models, marmosets have a delayed period of embryogenesis relative to humans [65]. Understanding this difference is extremely important to ensuring appropriate use of the species to model pregnancy and development [64]).

Evidence for positively selected growth factor-insulin like growth factor axis genes that may have driven miniaturization in *C. jacchus* has been found at *GHSR*, *IGF2*, *IGF1R*, *IGFBP2*, and *IGFBP7* [66,67]. Further, *Callithrix* growth patterns are likely tied to selection in marmosets for routine ovulation of multiple ova that leads to dizygotic twinning [68]. Altered placental and fetal growth patterns in marmosets may provide protection against discordant growth and placental pathologies of gestating litters in a simplex uterus [67]. The four positively selected genes with a likely role in regulating ova number and multiple gestations in *C. jacchus* are *GDF9*, *BMP15*, *BMP4*, and *WFIKKN1* [67].

Several factors associated with small body size, twin-rearing, and the fact that marmosets do not display anovulation have led to a marmoset social structure in which generally one female reproduces regardless of the number of other adult females within the group. This reproductive suppression has been the subject of extensive study over the past 30 years, but the mechanisms behind it are still somewhat unclear [68]. Impaired ovulatory function, which can be maintained for up to several years, is associated with suppressed pituitary secretion of luteinizing hormone and chorionic gonadotropin. Suppressed pituitary function is in turn likely associated with enhanced negative-feedback sensitivity to low levels of estrogen or blunted responsiveness to increasing estrogen. Some

marmoset daughters escape suppression while in the presence of their mothers and ovulate, though they often display impaired luteal function, which raises questions about the fertility of these females. However, even if daughters ovulate, as long as a social group remains intact, with only the daughter's father and brothers and potential breeding partners, these females rarely become pregnant.

CALLITHRIX CHIMERISM

Perhaps the most poorly understood *Callithrix* derived trait is chimerism, which occurs when a single organism possesses individual cells with >1 distinct genotype. Marmoset litters usually consist of twins or triplets that share a bidiscoid placenta with a fused chorion and vascular anastomoses [69]. Fusion of the chorionic membranes during embryonic development may allow blastocyst stem and precocial cells to cross the anastomoses, rendering littermates chimeric [70]. Marmosets are therefore hematopoietic chimeras with blood derived tissues containing cells of the individual as well as their littermates [71]. However the selection pressures that may have led to this trait remain obscure.

Benirschke et al. [71] noted that approximately 90% of *C. jacchus* pregnancies result in chimerism, but the degree of marmoset chimerism within a tissue and the extent of the tissues involved are not resolved. An evaluation of microsatellites from *C. kuhlii* pregnancies with known parentage and samples from multiple tissue lineages estimated that all tissues could be chimeric, but blood-derived tissues were significantly more likely to be chimeric than other tissues [70]. Accordingly, [70] showed that blood-derived tissues from deceased and living *C. kuhlii* being between approximately 50%-100% chimeric. Malukiewicz et al. [4] found similar results for *C. jacchus* when comparing blood and skin tissues via microsatellite genotyping. In another chimerism tissue analysis based on the SRY gene concluded that germ line cells were not chimeric and only hematopoietic lineages were [72]. Therefore, the contribution of embryonic cells to the overall phenotypic impact of chimerism on individual marmosets remains unknown.

An interesting question is how chimerism in marmosets impacts different levels of gene regulation. With respect to the epigenome, it is known that both underlying genetic signatures and environmental factors influence epigenetic changes [73,74]. Thus, in marmosets, the complication of how chimerism levels vary between tissue types, may result in different genetic and epigenetic

signatures across tissues. Further, the external signals that chimeric cells provide to one another within particular tissue environments may promote additional epigenetic changes. This idea has led to the hypothesis that chimerism may be a mechanism by which founder populations can rapidly increase their genetic diversity and gene expression variance, making them more adaptable to a range of novel environments [75]. However, while chimerism may affect such phenotypic plasticity through variation in genetic and epigenetic regulation, this relationship has not explicitly been tested in marmosets.

In most cases of biomedical modeling, it remains unclear whether chimerism may play a role in the variability found between individuals. However, chimerism in marmosets may be particularly well suited to questions designed to take advantage of the close relatedness of multizygotic litters using techniques such as adoptive transfer or paired design using litter mates as controls versus treatment subjects in immunology studies. Additionally, marmosets may be a natural model in which to study the impact of genomic conflict for questions of stem cell transfer, maternal-fetal microchimerism, or transplants.

CALLITHRIX COGNITION

Cognition studies is one of the growing areas of research for common marmosets (reviewed in [51]). Although laboratory studies dominate *C. jacchus* research from the last five years, field studies should gain more attention because of their complementary value in relation to laboratory studies [51]. In this regard, field studies can have particular importance in areas that need integration between cognition and ecology. For instance, a recent study showed that common marmosets use spatial cognitive abilities to effectively obtain food in nature [76]. Studies like this, in which wild animals face the constraints and limitations imposed by the environment, can help us better understand the selective pressures that ultimately shaped their cognitive abilities.

Common marmosets are a particularly promising model for studying evolutionary and functional questions in primate personality due to their explorative and highly social character, and adaptability to different natural environments [44, 51]. Marmosets display personality traits when

assessed with tools such as personality tests [e.g., 77] and behavioural observations [e.g., 78]. In particular, marmosets exhibit short-term and long-term consistency of non-social and social personality traits [79]. Personality structure under captive and wild conditions is rather similar [79]. Inoue-Murayama et al. [78] found that *C. jacchus* sociability was positively linked with increased subjective well-being and cortisol levels, whereas neuroticism and dominance were associated with certain genetic polymorphisms. Interestingly, group level similarity in marmoset personality has been found [77], possibly as a product of shared social environment, genetics or social facilitation. Future studies are needed to further explore links between inter-individual behavioural and cognitive variation in this species.

CALLITHRIX GENOMICS and GENETICS

CALLITHRIX GENOMIC ASSEMBLIES

A high-quality genome is required for animal models, and in 2014 *C. jacchus* became the first New World monkey whose genome was sequenced and assembled. Sanger sequences from a female marmoset were generated by an international consortium and assembled into extended continuous arrays of nucleotides (contigs). The ultimate published sequence (designated CalJac3 in the NCBI genomics database) consisted of 2.26 billion base pairs covering all *Callithrix* chromosomes [66]. However, the large number of continuous sequences were broken by gaps so that no chromosome sequence was complete end-to-end. Nonetheless, this genome allowed analysis of marmoset twinning genes [66, 67], and detection of primate-specific constrained elements [80].

Tables 2 and 3 list several genomic assemblies now available for *C. jacchus*, some which

Table 2. Marmoset genome assemblies from short-read next-generation sequencing.

Genbank Accession	Submitter	Date	Sequencing Technology	Contig N50 (Kb)	Scaffold N50 (Mb)	Number contigs	Number Scaffolds

GCA_002754865.1 (ASM275486v1)	Broad Institute	2017/11/06	Illumina HiSeq	155.3	129.2	88,439	39,944
GCA_001269965.1 (CIEA01)	Keio University	2015/08/07	Illumina GAIIx, Illumina HiSeq	61.0	NA	109,198	NA
GCA_000832365.1 (ASM83236v1)	UN-NHPGC	2015/01/29	Ion Torrent	0.859	NA	1,992,178	NA
GCA_000004665.1 (calJac3)	Washington University (WashU)	2010/01/22	ABI 3730	29.3	5.2	201,371	16,399

Table 3. Marmoset genome assemblies from long-read sequencing.

Genbank Accession	Submitter	Date	Sequencing Technology	Contig N50 (Kb)	Scaffold N50 (Mb)	Number contigs	Num Scaffolds
GCA_013373975.1 (CJA1912RKC)	Keio University / RIKEN	2020/06/03	PacBio RSII	24,828.9	132.3	3,719	1,872
GCA_009663435.2 (cj1700_1.1)	McDonnell Genome Institute at Washington University	2020/05/22	PacBio, Illumina	25,227.1	98.2	1,336	994
GCA_011078405.1 (mCalJa1c.mat , maternal alternate haplotype)	Vertebrate Genomes Project	2020/03/10	PacBio Sequel I CLR; Illumina NovaSeq; Arima Genomics Hi-C; Bionano Genomics DLS	8,609.0	146.9	1,284	216
GCA_011100535.1 (mCalJac1.pat ; paternal alternate haplotype)	Vertebrate Genomes Project	2020/03/10	PacBio Sequel I CLR; Illumina NovaSeq; Arima Genomics Hi-C; Bionano Genomics DLS	14,650.8	137.0	1,124	336
GCA_011100555.1 (mCalJac1.pat.X , principal haplotype of the diploid genome)	Vertebrate Genomes Project	2020/03/10	PacBio Sequel I CLR; Illumina NovaSeq; Arima Genomics Hi-C; Bionano Genomics DLS	13,255.6	137.9	1,312	353
GCA_009811775.1 (CJ2019)	Keio University	2019/12/13	PacBio RSII	6,375.4	143.9	1,812	65

have been published [e.g., 81]. One unpublished *C. jacchus* genome assembly by the Vertebrate Genomes Project is based on a novel technique called trio-binning in which the reads from parental and maternal haplotypes are assembled separately, resulting in a diploid assembly [82]. The most recent assembly is designated cj1700_1.1, which has now replaced calJac3 as the *C. jacchus* reference genome. Cj1700_1.1 has a contig N50 (defined as the length of an individual contig for which half of all the available sequence occurs in contigs larger than that figure) of 25.2 megabases, an approximately 860-fold improvement over calJac3 in terms of contiguity.

More accurate and (largely) complete reference genome sequences for marmosets are opening tremendous opportunities for novel research. For example, CRISPR/Cas9 gene knock-in [83] generated transgenic marmosets to model human disease. Further, the newer *Callithrix* genomic assemblies contain less gaps and better assembly of duplicated regions, which will provide better genomic annotation, leading to improved analysis of assays that require alignment of short reads to a reference genome (RNA-seq, ATAC-seq, ChIP-seq, etc). The diploid assemblies will facilitate the study of haplotype structure and structural variation in marmosets [84].

CALLITHRIX FUNCTIONAL GENOMICS and EPIGENOMICS

Marmoset epigenetics studies have examined DNA methylation patterns in a wide sampling of tissues like bone [85] and placental tissue [86]. DNA methylation variation can be associated with variation in transient traits like body weight, and when such changes in gene regulation occur in developmentally important tissues like the placenta, they can potentially impact phenotypic development in offspring [86]. Interestingly, [85] found that DNA methylation variation in marmosets is not strongly associated with variation in static traits like bone morphology. This study also noted higher levels of DNA methylation heterogeneity as compared to other nonhuman primate species, which led these researchers to speculate whether such differences might be due to chimerism. Such epigenetic results suggest that it is particularly relevant to examine to what degree chimerism affects epigenetic variation within specific tissues or cell types, as levels of chimerism are known to vary across different marmoset tissues [4,70,72].

Brain development, evolution, and plasticity is one area of marmoset functional genomics that is being explored. Current work has focused on characterizing candidate gene expression patterns in brain tissues [87,88], but some research has also assayed candidate gene expression responses following severe psychological perturbations such as parental separation [89]. Interestingly, while most brain regions show conserved levels of candidate gene expression between marmosets and mice, some areas of the brain do show marmoset-specific expression patterns, such as the early visual cortical area and afferent areas of the hippocampus [87].

Now is a particularly relevant time to pursue functional genomics research, especially in primates [90], as technologies can characterize whole transcriptomes and epigenomes at both bulk and single-cell resolutions. Already, some comparative cross-species studies that include marmosets have begun utilizing these single-cell methods [91,92]. Further, these technologies can be used on primary tissue samples, as well as *in vitro* cell culture systems. Marmoset cell lines that are readily used include embryonic stem cells (ESCs) [e.g., 93] and induced pluripotent stem cells (iPSCs) [e.g., 94,95]. ESCs and iPSCs are ideal for functional genomics research, especially in the context of disease, as they are immortalized and self-renewing, have the potential to differentiate into cells from any germ layer, and can be readily used in controlled experiments [93]. Another benefit is the relative ease with which these cell lines can be genetically modified [83].

Finally, consolidating functional genomics data for better access and comparison is essential in these efforts. Larger consortia that compile large -omics datasets for marmosets are being built [96,97]. For instance, functional genomics data from marmoset brain tissues have been collected and incorporated into the NIH Brain initiative [98]. This is of particular relevance as the common marmoset has been identified as an important model organism for neuroscience research that can bridge the gap between mice and humans [96,99]. Such foundational databases have enabled subsequent explorations such as single-cell RNA-seq of *C. jacchus* and other mammals have identified primate-specific interneuron subtypes [100] and common and divergent features of preimplantation development [101].

CALLITHRIX IMMUNOGENETICS

Although New World primate immunogenetic studies lags behind that of Old World primates, complex immunogene families like Major Histocompatibility Complex (MHC) Class I and the Natural Killer Complex (NKC) have been sequenced and annotated in *C. jacchus*. The MHC, an evolutionary hallmark of enormous genetic variability and genomic diversity, determines immune responsiveness and is associated with many diseases. MHC class I proteins bind pathogen-derived peptides for presentation to specialized cells for immune response initiation [102]. MHC class I proteins also interact with natural killer (NK) cell receptors to prevent NK cell destruction of healthy cells [103].

The *C. jacchus* MHC class I region is composed of three segments- *Caja-B/C* segment, the *Caja-E* segment, and the *Caja-G/F* segment [104,105]. In humans, six MHC class I loci are known as HLA (human leukocyte antigen)-A/B/C/D/E/F/G, and *Caja-B/C* corresponds to HLA-B/C [104] and *Caja-G/F* corresponds to the HLA-A/G/F region [104]. Shiina [105] described *Caja-B/C* as a 1,079 kb fragment containing nonclassical (low polymorphism) MHC class I genes and [105] characterized a 854 kb tract of the *Caja-G/F* segment, which harbors classical (polymorphic) MHC class I genes. *Caja-B/C* contains 54 genes among which there are nine functional MHC genes and four MHC pseudogenes [105]. *Caja-G/F* encompasses six functional MHC genes as well some non-MHC genes and MHC pseudogenes [105]. These MHC I regions likely evolved by segmental duplication [104,105].

Interestingly, the human HLA-G locus shows limited polymorphism and grants immuno-tolerance between mother and fetus during pregnancy [106]. The homologous *Caja-G* locus instead shows high allelic polymorphism levels, and expression of *Caja-G/F* alleles in various tissues suggests that this locus has taken on the role of classical MHC I function in *C. jacchus* [106]. Evidence for genetic conversion in the role of *Caja-G/F* is seen further when comparing exon diversity between *Caja-G/F* and HLA-G gene associated with MHC protein binding of pathogen peptides. *Caja-G/F* MHC genes show a high level of polymorphism that sharply contrasts that of HLA-G MHC genes [106].

Two structurally unrelated classes of NK cell receptors exist in primates- the killer cell immunoglobulin-like receptors (KIR) encoded by the leukocyte receptor complex (LRC) and killer

cell lectin-like receptors encoded by the natural killer complex (NKC) [107]. The KIR lineage is present in all simian primates [103], but in *C. jacchus* there are apparently only two KIR loci [66]. In marmosets, NKC genes cluster together and are orthologous to humans, but the marmoset NKC segment is 1.5x smaller [107]. Marmoset NKC genes also show moderate polymorphism [106]. Averdam et al. [106] found that the *C. jacchus* NKC encodes a single inhibitory heterodimer (CD94/NKG2A) and two activating NK cell receptors, (CD94/NKG2CE and NKG2D). In humans, CD94/NKG2A and CD94/NKG2C are ligands for HLA-E, a nonclassical MHC class I molecule that plays an important function in cell recognition by NK cells [107], but the role of *Caja*-E remains to be determined.

CALLITHRIX VIRUSES and the VIROSPHERE

CALLITHRIX as *MODELS* of *VIRAL DISEASES* and *VACCINE DEVELOPMENT*

Callithrix jacchus has served as a model for several aspects of biomedical research related to viruses. In humans, Epstein-Barr virus infection may enhance risk of developing multiple sclerosis, and natural infection of *C. jacchus* with the related γ 1-herpesvirus CalHV3 has been harnessed to model human autoimmune disease [108]. Additionally, *Callithrix jacchus* has been used to model hepatitis A infection [109]. The species was also used for *in vitro* transformation of two permanent and virus producing lymphoblastoid cell lines with a nasopharyngeal carcinoma derived Epstein-Barr viral strain [110]. Other viral pathogens that have been explored in *Callithrix* include parainfluenza virus type 1 [111], Flaviviridae-like viruses [112], Oropouche virus [113], and Simian foamy virus [114].

More recently, *Callithrix* is emerging as an infection model of several arthropod borne viruses (arboviruses) of single stranded-RNA viruses from the genus *Flavivirus*. These viruses include dengue virus (DENV), now considered the most prevalent and quickly spreading human arboviral disease [115]. DENV infection causes disease in humans that ranges from high fever and joint pain to shock and death (<https://www.cdc.gov/dengue/index.html>). Both *C. jacchus* and *C. penicillata* are candidate systems to model DENV pathogenesis and for discovery of antiviral drugs and vaccines [116,117]. In

C. penicillata, DENV infection produced marked changes in microglial cells of the central nervous system [118]. Both *Callithrix* species show elevated proinflammatory cytokines upon DENV infection, which may model aspects of human DENV pathogenesis. Another *Flavivirus*, Zika virus (ZIKV) was originally discovered in Uganda [119], but is now the newest *Flavivirus* arrival in the New World. Pregnant women infected with ZIKV are at risk for miscarriage and fetus microcephaly, and adults can develop neurological condition like Guillain-Barré syndrome [120]. A number of recent studies have found that *C. jacchus* replicates many of these human clinical and gestational symptoms [64,120].

EXPOSURE of WILD CALLITHRIX to PATHOGENIC ARBOVIRUSES

Yellow fever virus (YFV), another flavivirus, was anthropogenically introduced to the Americas in the 15th and 16th centuries from Africa, and exists in Brazil as part of an established sylvatic cycle [121]. Serological surveys of YFV antibodies in wild primates have shown some positive results in *Callithrix* [122,123]. Retrospective data on surveillance of primate epizootics shows that *Callithrix* represented 8.7% of the positive non-human primates (NHPs) for YFV in Brazil between 1996 and 2016, which increased to 17.1% between 2016 to 2019 [123]. Since 2016, free-living *Callithrix* infection has been demonstrated in southeastern Brazil during a recent major YFV outbreak, suggesting increasing importance of *Callithrix* for YFV epidemiology in Brazil. Between 2017 and 2018, [124] analyzed 2,099 NHP samples from southeastern and northeastern Brazil suspected of YFV. *Callithrix* were among the most frequently infected YFV- infect NHPs, with 140 (9.3%) *Callithrix* being positive among 1505 tested marmosets. These percentages may be even higher for epizootic diseases, as classification of NHPs in epidemiological data usually doesn't go beyond the genus level [125]. Cunha et al. [126] observed that *Callithrix* found positive for YFV through molecular methods did not necessarily demonstrate any associated histopathological changes or immunohistochemical YFV positivity.

The immunization of NHPs against the yellow fever virus is still the subject of intense debates in the scientific community. These debates are likely due to the possibility of interfering in the use of these animals as sentinels for the transmission of the virus in the natural environment. Although there are no extensive studies on the safety and effectiveness of the vaccine in NHPs, promising initiatives to test the safety and effectiveness of the vaccine are under development in a partnership between the Institute of Technology in Immunobiologicals (Biomanguinhos / Fiocruz) with the Center of Primatology of Rio de Janeiro.

Callithrix susceptibility has been shown for other arboviruses like DENV, ZIKV, Mayaro (MAYV), and Chikungunya (CHIKV), all which can potentially establish sylvatic cycles in Brazil and other parts of South America [127]. Currently, only a limited number of studies have looked at arbovirus exposure in wild Brazilian primates, and [128] found no exposure evidence to the above arboviruses in urban and peri-urban *jacchus* group marmosets sampled in northeastern Brazil between 2012 and 2017, except one *C. jacchus* that showed serological CHIKV exposure. Also, [129] found no evidence of ZKV infection in *Callithrix* sampled in southeastern Brazil between 2015 and 2018. On the other hand, evidence for ZKV infection was reported in free-ranging peri-domestic *C. jacchus* sampled between 2015 and 2016 in Ceara state [130]. Additionally, [131] found evidence of widespread ZIKV exposure in over 50 *Callithrix* sampled Minas Gerais and São Paulo states.

In order for a translocated arbovirus to enzootically establish itself within the Neotropics, wildlife exposure, zoonotic infection, and persistent enzootic transmission of the translocated zoonosis need to occur [132]. *Callithrix* characteristics that can facilitate arbovirus transmission include a great adaptability to human environments, frequent interactions with human beings in urban and peri-urban environments also occupied by anthropophilic arbovirus vectors such as *Aedes aegypti* and *A. albopictus* [124]. The presence of short-lived, fast reproducing primates also facilitates the persistence arbovirus transmission [132], which are certainly characteristics that define many *Callithrix* species and hybrids.

OTHER PATHOGENS of CALLITHRIX

Human herpesvirus 1 (HV1) is transmissible from humans to *Callithrix* marmosets, which has resulted in a number of fatal epizootics events [133,134]. Transmission occurs through the close contact that these animals have developed with humans, facilitated by the habit of offering food, since the virus can be transmitted through contaminated saliva, aerosols and fomites. Transmission can also occur spontaneously between marmosets within the same group [133,135]. Affected animals are usually found dead or with severe neurological symptoms, resulting from virus-induced encephalitis [134]. The virus also causes death from meningitis [134], erosions and ulcers in the oral cavity, in addition to hemorrhages and focal necrosis in several organs [135].

Rabies is a zoonotic disease caused by lyssaviruses, transmitted through saliva, and infection causes fatal acute encephalomyelitis. The disease is transmissible from animals to humans by bites. *Callithrix jacchus*, has played an important role in the transmission of rabies in northeastern Brazil, where it maintains a distinct transmission cycle [136]. Infected marmosets often show no clinical signs, but rather sudden death preceded by sadness or weakness. A recent case of rabies in *Callithrix* sp. occurred in urban Niterói, Rio de Janeiro, Brazil [137]. The proximity of marmosets to urban environments and the practice of capturing and maintaining these animals as pets are two factors that greatly increase the risk of marmoset rabies transmission [138].

CALLITHRIX VIROME RESEARCH

Although the virosphere incorporates the most abundant and diverse group of organisms on Earth, it remains largely neglected, with most attention given to viruses of high biomedical relevance. It is estimated that there are over 500,000 undiscovered animal viruses that are transferable to people [139]. Although *Callithrix* virome work is still at an infancy, one pioneering study helped to characterize two novel *Papillomavirus* genomes (CpenPV-1 and CpenPV-2) in captive *C. penicillata* [140]. Other viral families also have been sequenced in this work, but remain to be published. An important future role for *Callithrix* virome studies will be to help understand the capacity of *Callithrix*

to be YFV disease reservoirs and further understand why infected *Callithrix* show relatively low YFV viral loads. Also, virome studies can help to understand if this genus could actually represent a potential risk for a severe urban YFV outbreak [124,126] and even unknown zoonotic virus spillover. Overall, virome studies can greatly enhance our ability to detect known and novel viral sequences and can also help to understand the entire dynamic and interactions of the *Callithrix* virosphere.

CALLITHRIX SOCIAL DYNAMICS

While long-term studies are still necessary to understand wild *Callithrix* social dynamics, some insight is available from *C. jacchus* groups at the Tapacurá Field Station, São Lourenço da Mata, Pernambuco (08° 07' S, 34 55' W, Figure 8). A typical Tapacurá *C. jacchus* social group may contain eight individuals- three reproductive adults, two juveniles, two infant co-twins, and an adult or subadult that may or may not be related to other group members [141]. The three breeding adults may include a dominant female and two males that take on variable roles in offspring care and group defense. One breeding male will be the principal infant carrier, while the second breeding male aids the breeding female in choosing group feeding, resting, and sleeping sites, and in territorial defense [141]. A breeding female depends on her group's support to maintain dominant status, but the loss of the reproductive position may not result in her departure. There are instances at Tapacurá where breeding females were displaced by one or two daughters, but remained in the group to suckle grandchildren along with the new dominant female(s) [141].




Figure 8. Study area of the Tapacurá Field Station, São Lourenço da Mata, Pernambuco, Brazil, with a 50m \times 50m grid of trails. Shown are the home ranges of five focal *C. jacchus* study groups (EC, EM, EP, ET, EV) represented by continuous lines, peripheral groups (orange dotted lines) and temporary groups (pale blue dotted lines) between May 1994 and May 1996.

In regards to sexual behavior studied at Tapacurá, between 1994-1996, 52 copulations were observed, of which 16 were intra-group, and only two involved non-reproductive females. Of 33 extra-group copulations, 17 were between neighboring focal study groups, and 16 were between members of a focal group and peripheral groups. To illustrate extra-group copulations, the reproductive female of one focal group was seen copulating seven times with two dominant males (2 to 3 times each) in neighboring groups. In general, sexually mature females take all opportunities to copulate, independent of their phase in the estrus cycle. Females also disperse more than males (Figure 9), leaving their natal groups to increase their chances of breeding [142].

Interestingly, the strongest social interactions within *C. jacchus* groups are between the male care-giver (one of the probable fathers in the group) and infants. The findings help to explain, at least in part, why males disperse less than females, as nutritional provisioning of infants by the male caregiver demands dedication and skill. On one occasion, a male carer was seen to dexterously secure the floral pedicel of a cashew fruit with both hands and, while balancing with extended legs, lifting it above his head after biting the hypocarpium several times to allow the infants to lick the juice (Figure 10). During the infant weaning phase, the carer male also teaches infants skills needed to travel, forage, confront predators, and how to recognize and catch and manipulate food items [141]. The male carer is the last to abandon a dead infant, which illustrates a deep social bond between a marmoset male and infants [143].

ANTHROPOGENIC EFFECTS on *CALLITHRIX*: PET TRADE, SPECIES INVASIONS, and HYBRIDIZATION

PET TRADE

While it is illegal in Brazil to have native primates as pets [144,145] and in São Paulo state there is a moratorium on captive breeding of marmosets by authorized commercial breeding




Figure 9. Changes in group composition (males in the top and females below) of five *C. jacchus* Tapacurá Field Station focal study groups between May 1994 and May 1996. Each square panel represents a single social group, with male members on top and female members on the bottom. Bars represent individual marmosets and arrows represent their movement between social groups.



Figure 10. Illustration of a *C. jacchus* breeding male caregiver hydrating marmoset infants with cashew hypocarpium juice (*Anacardium occidentale*), an act which shows dexterity and skill.

facilities [146], there still exists an underground Brazilian, illegal pet market [147]. Marmosets, most often *C. jacchus* and *C. penicillata*, are taken from the wild by several methods including trapping, killing of adults to take the infants, rescue of infants after habitat loss, and then either sold locally or transported to large urban areas [147-149]. Confiscated marmosets from animal traffickers are often stressed, malnourished and dehydrated - conditions that favor the development and transmission of diseases. Transport stress and inadequate hygiene conditions also contribute to the development of infections.

Captured marmosets appear to be trafficked using the same illegal wildlife trade routes as other wildlife. In the case of *C. penicillata*, construction of roads and increased vehicular traffic between the Cerrado and southeastern Brazil, has greatly facilitated illegal trafficking of this species [58]. There are few studies that quantify the pet trade, but *Callithrix* individuals, especially *C. jacchus* and *C. penicillata*, have been among the top five species of mammals confiscated and sent to government triage centers. Between 1999 and 2006, out of 4631 primates received by Brazilian triage centers, 54.2% were from the *Callithrix* genus [148]. However, the actual numbers of marmosets taken from the wild for commercial purposes are likely underestimated, as many confiscated individuals do not go to the triage centers, instead, they are released in the nearest forests by authorities, rural landowners, or dealers/owners. Important consequences of the illegal primate trade include the release of marmosets outside of their natural areas of distribution, anthropogenic hybridization, saturation of captive facilities with hybrids and non-hybrid marmosets, and biological invasions that threaten endangered, native *Callithrix* species [2,150,151].

INVASIVE CALLITHRIX SPECIES in the SOUTHEASTERN BRAZILIAN ATLANTIC FOREST

Allochthonous *Callithrix* species began appearing in portions of the southeastern Brazilian Atlantic Forest within approximately the last 20-30 years [2,152,153]. Now, three *Callithrix* species, *C. geoffroyi*, *C. jacchus*, and *C. penicillata*, have high invasive potential [61,62], and are spreading

throughout the southeastern Atlantic Forest due to the legal and illegal pet trades [21,58,154]. As a result, *Callithrix jacchus* and *C. penicillata* have established several allochthonous populations in southeastern Brazil [2,21,58,150-155]. Mitogenomic data shows that source populations of allochthonous *Callithrix* species actually come from across broad geographic origins outside of the southeastern Brazilian Atlantic Forest [2]. As discussed below, anthropogenic hybridization between allochthonous marmosets and either other allochthonous or autochthonous congeners occurs widely within the southeastern Brazilian Atlantic Forest [2,4,21,153-155].

Contemporary allochthonous *Callithrix* species are usually found in urban or peri-urban areas of the southeastern Brazilian Atlantic Forest [21, 58,151,152,155-157]. Within these contexts, allochthonous *Callithrix* have frequent human contact and exposure, and receive anthropogenic food supplementation [54, 58,158]. These allochthonous marmosets can also be found within or around natural reserves [13,58,151,152,155, 157], and in some areas, such marmoset populations are larger than those of native, endangered callitrichids [e.g., 151]. Such introductions could alter the ecological relationships among taxa [13, 159], as the main threats posed by allochthonous marmosets are competition for food resources, increased predation of native fauna, the introduction and maintenance of disease, and hybridization [160-162]. The threats can become exacerbated when introduced marmosets become abundant in a landscape of small fragments.

ANTHROPOGENIC CALLITHRIX HYBRIDIZATION

Due to the legal and illegal pet trades, *C. jacchus* and *C. penicillata* have established several allochthonous populations in southeastern Brazil that hybridize with allochthonous and autochthonous congeners (Figure 11; [19,21,163]). It is highly likely that *C. aurita* faces competition, conservation, and genetic threats from allochthonous *jacchus* group species and anthropogenic hybrids [2,10,154]. *Callithrix flaviceps* also likely faces similar pressures. Some cases exist of native *C. aurita* and *C. flaviceps* meeting up and interbreeding with hybrid and allochthonous *Callithrix* at urban fringes (reviewed in [21]). Such interactions likely facilitate gene flow from allochthonous *jacchus* group

species into native marmoset populations in southeastern Brazil, with consequences that may include outbreeding depression, admixture, hybrid swamping, or introgressive replacement [164-166]. Indeed, [2] showed for the first time, evidence of introgression of mitochondrial DNA from allochthonous *C. jacchus* into the genetic background of native *C. aurita* from the São Paulo metropolitan area. These data also showed the first genetic evidence for cryptic hybridization within the *aurita* group marmosets. A few anthropogenic *Callithrix* hybrid zones have also been studied using mitochondrial, nuclear, and Y-chromosome markers [2,4,21,155, <https://tede.ufrj.br/jspui/handle/jspui/3009>].

Anthropogenic hybridization with *C. jacchus* and/or *C. penicillata* also represents a potential risk for genetic extinction of the other two *jacchus* group species, *C. geoffroyi* and *C. kuhlii*, and [58] recently showed that *C. penicillata* is encroaching on the native range of *C. geoffroyi*. Anthropogenic hybridization of *jacchus* group species generally results in the formation of hybrid swarms, admixed populations that lost parental phenotypes and genotypes [4,21,153]. Should large numbers of exotic *C. jacchus* or *C. penicillata* ever invade native ranges of *C. kuhlii* or *C. geoffroyi*, the latter two species may be threatened with genetic swamping by the former two species, a process through which parental lineages are replaced by hybrids that have admixed genetic ancestry [169]. Further, biological invasions by other marmosets present potential conservation risks for *C. kuhlii*, which is already considered vulnerable.

CALLITHRIX POPULATION DECLINES and CONSERVATION

CALLITHRIX POPULATION DECLINES

All *Callithrix* species face population declines across their native ranges as a result of habitat loss and land conversion, principally for urbanization, agriculture, and livestock production [11, 10, 12,168,169]. The estimates of remaining original forest area are 63% for the Cerrado, 47% for the Caatinga, and 11-28% for the Atlantic forest [170,171]. Furthermore, between 2006 and 2015, the accumulated area of lost natural vegetation of Brazilian biomes was 300,000 km² [172].




Figure 11. Anthropogenic *Callithrix* hybrid zones. Each natural species range is color-coded as in Figure 1. Abbreviations of Brazilian states follow Figure 1. Hybrid zone abbreviations: JP: *C. jacchus* x *C. penicillata*; PG: *C. geoffroyi* x *C. penicillata*; AH: *C. aurita* x *Callithrix* sp.; AP: *C. aurita* x *C. penicillata*; JA: *C. jacchus* x *C. aurita*; JH: *C. jacchus* x *Callithrix* sp; PH: *C. penicillata* x *Callithrix* sp. Hybrid zone locations adapted from [19,21,168]

For arboreal primates, marmosets included, continued deforestation in Brazil is the most serious threat to the persistence of populations and species [12,173].

The three most threatened *Callithrix* species are *C. flaviceps*, *C. aurita* and *C. kuhlii*; with the first considered “critically endangered” [11], the second listed as “endangered” [10], and the third considered “vulnerable” [174]. *Callithrix aurita* has also been listed among the 25 most threatened primates on the planet [175]. Projections based on forest loss and other anthropic factors predict reductions of over 50% of *C. aurita* and *C. flaviceps* populations in the next 18 years [10,11]. For instance, forests within the geographical distribution of *C. aurita* were reduced by 43% between 1990 to 2008 [176]. For *C. flaviceps*, agricultural activities have profoundly modified the environment where the species naturally occurs, with cattle breeding and coffee plantation occupy formerly forested areas [58]. The remaining *C. flaviceps* sub-populations, estimated at ~4,440 individuals, are restricted to the margins of riparian forests, and steep hillsides unsuitable for raising cattle, or planting coffee or eucalyptus [11]. Between 2015-2017, the wide-spread Brazilian YFV epidemic decimated > 90% the *C. flaviceps* population at RPPN Feliciano Miguel Abdala, a private reserve in Caratinga, Minas Gerais [177]. The virus may have caused other such reductions across the *C. flaviceps* range [11]. It is interesting to note that the global climate changes that are happening could lead to reductions in the distribution area as high as 95% for *C. flaviceps* and 27% for *C. aurita* [60]. For *C. kuhlii*, whose total population is estimated to be >10,000 individuals, a population decline of 30% is expected by 2031, because much of its native habitat is being converted to cattle ranches and agriculture [174].

The other three *Callithrix* species are of “least concern”, nonetheless their populations are declining throughout their native geographic ranges [168,169,178]. These changes are likely driven by the illegal wildlife trade and agricultural/ranching activities. The urban expansion that accompanies agriculture expansion is characterized by activities that limit dispersal or increase mortality: building of roads, topographical changes caused by erosion, and the presence of domestic animals such as dogs that enter the forests for hunting [180]. For the three “least concern” *jacchus* group species, the human activities that devastated marmoset habitats also drive social groups into urbanized areas, where they are more vulnerable to disease, accidents and poaching- perhaps making urban *Callithrix* populations less viable.

Nonetheless, marmoset populations manage to persist in urbanized areas [181, pers. obs. IOS, JM, LCMP, VB]. The use of urban environments demands plasticity and perhaps adaptations in behavior, diet, metabolism, toxicants, immune defense and relationships with pathogens. Marmosets are indeed capable of showing adaptive plasticity in social behavior and sleeping site selection in urban, disturbed environments [182,183].

CALLITHRIX CONSERVATION LEGISLATION, SPECIES ACTION PLANS, and CURRENT INITIATIVES

The Brazilian constitution grants protection for all wildlife, and Brazil has devised a number of legal instruments related to the protection and conservation of *Callithrix* species (Table 4). The Brazilian National Biodiversity Policy [184] was formulated to comply with the Convention on Biological Diversity of 2002. Three instruments of this policy deal directly with species conservation [185]: the Official National Lists of Endangered Species; the Red Books of Endangered Brazilian Species; and the Conservation of Endangered Species National Action Plans (NAPs). NAPs define both *in situ* and *ex situ* actions for the conservation and recovery of endangered species. Initially, conservation of *Callithrix* species came under the Conservation of Mammals in the Central Atlantic Forest NAP [186]. In 2018, *Callithrix* species became part of the NAP for the Conservation of The Primates of the Atlantic Forest and the Collared Sloth [187]. The Saguis-da-serra Conservation Program (PCSS) is an initiative to put into practice the actions foreseen by such NAPs, along with the help of several researchers and institutions.

Recently, a studbook was created to track genealogical relationships between individuals of the Brazilian captive *C. aurita* population [154]. The *C. aurita* Studbook Keeper, C. Igayara, estimates that there are approximately 33 individuals that make up the current captive population. An important objective in breeding and maintaining a captive *C. aurita* population is to maximize genetic diversity individuals [154,175]. Currently, no *C. flaviceps* are maintained in captivity (pers obs., FR de Melo).

Another important *Callithrix* conservation initiative has been the creation of the Center for the Conservation of Saguís-da-serra at the Federal University of Viçosa, Brazil (CCSS-UFV) - the first primatology center in the world devoted exclusively to develop both *in situ* and *ex situ* conservation activities (pers obs., FR de Melo). Among objectives of the CCSS-UFV are to develop directives for the management of *in situ* populations of *C. aurita* and *C. flaviceps*, and to help guide researchers in choosing the best management strategies for particular marmoset populations. Additionally, protocols are being developed that aim to standardize data collection for mountain marmosets, and prioritizing locations for further *in situ* research (pers obs., FR de Melo).

In the last decade, regulations for the management of fauna have emerged in Brazil [e.g., 188,189] including, more recently, the management for the control of invasive alien and allochthonous species [146,190,191]. Currently, many confiscated and apprehended animals end up in Brazilian animal triage centers [192], and these captive facilities do not have further capacity to receive a higher influx of marmosets, nor to keep them for long. In addition, most management institutions have no interest in maintaining “common” or non-threatened species. From an ecological perspective, allochthonous marmosets can be considered introduced and potential or *de facto* invasive species. If allochthonous marmoset species are considered exotic, then the proper course of action would be to eradicate them during the early invasion stages or control established populations. Introduced and hybrid marmosets could be also removed from areas in which they pose a direct conservation problem, but the major obstacle for this course of action is resolving the destination of these animals. Ultimately, although these kinds of actions are supported by national Brazilian legislation and NAPs, effective action implementation runs into problems concerning funding and reaching a common consensus for the best plan of action.

CONCLUSION

Marmosets have been the quintessential laboratory primate model for a wide variety of health studies. Nowadays, with the increased adoption of the concept of OneHealth, and the opportunities afforded by established field sites for several species under a variety of environmental conditions, marmosets can become a model group to study the evolution and ecology of infectious diseases, the relationships among environment, health, ageing and genetics, adaptive responses to emerging diseases, immune system genetics, physiology, epidemics and other public health topics. These studies can contribute both to the understanding of the dynamics of diseases of human concern and to the conservation of several endangered species of marmosets and other primates with which they are sympatric. This scenario offers an unprecedented opportunity for synergy between evolutionary biology, human health and species conservation concerns. The stage is prime to establish long term scientific exchange and collaboration among scientists from various disciplines and different countries to address issues of world-wide importance.

Table 4. List and description of the main legal instruments related to the protection and

conservation of *Callithrix* species.

Topics	Legal Instrument	Description	Observations
Protection and Conservation of Wildlife	Law N° 5.197/1967	It provides for the protection of fauna and provides other measures	“Art. 1º Os animais de qualquer espécie, em qualquer fase do seu desenvolvimento e que vivem naturalmente fora do cativeiro constituindo a fauna silvestre, bem como seus ninhos, abrigos e criadouros naturais, são propriedade do Estado, sendo proibida a sua utilização, perseguição, destruição, caça ou apanha”
	Law N° 7.653/1988 - Wildlife Protection Act	Altera a redação dos arts. 18, 27, 33 e 34 da Lei nº 5.197, de 3 de janeiro de 1967, que dispõe sobre a proteção à fauna, e dá outras providências .	renames, modifies and reenacts LAW No. 5,197/1967.

	FEDERAL CONSTITUTION of 1988	Art. 225. All have the right to the ecologically balanced environment , well of common use of the people and essential to the healthy quality of life, imposing on the public power and the collectivity the duty to defend it and preserve it for the present and future generations.	Paragraph 1, item VII, points out the burden given to the Public Power to safeguard fauna and flora, being prohibited, in the form of law, practices that endanger their ecological function, that promote the extinction of species or subject animals to cruelty.
	LAW No. 9,605/1998 - Environmental Crimes Act	Dispõe sobre as sanções penais e administrativas derivadas de condutas e atividades lesivas ao meio ambiente, e dá outras providências .	

	<p>DECREE N° 4.339/2002</p>	<p>Establishes principles and guidelines for the implementation of the National Biodiversity Policy</p>	<p>Created to be implemented with the participation of federal, district, state and municipal governments, and civil society, considering:</p> <ol style="list-style-type: none"> 1. The commitments made by Brazil to sign the Convention on Biological Diversity (CBD) during the United Nations Conference on Environment and Development - UNCED in 1992; 2. The provisions of Article 225 of the Constitution, in Law No. 6,938, of August 31, 1981, which provides for the National Environment Policy, the Rio Declaration and Agenda 21, both signed by Brazil in 1992, during UNCED, and the other current rules on biodiversity; And 3. That the development of national biodiversity strategies, policies, plans and programmes is one of the main commitments made by the member countries of the Convention on Biological Diversity.
--	-----------------------------	---	---

	<p>JOINT ORDINANCE MMA AND ICMBIO Nº 316/2009</p>	<p>Apply the following instruments for the implementation of the National Biodiversity Policy aimed at the conservation and recovery of endangered species: I - Official National Lists of Endangered Species; II - Red Books of Endangered Brazilian Species; II - National Action Plans for the Conservation of Endangered Species</p>	
--	---	--	--

	<p>COMPLEMENTARY LAW No. 140/2011</p>	<p>It lays down rules, pursuant to item III, VI and VII of the caput and the sole paragraph of Art. 23 of the Federal Constitution, for cooperation between the Union, the States, the Federal District and the Municipalities in administrative actions arising from the exercise of common competence relating to the protection of notable natural landscapes, the protection of the environment , the protection of pollution in any of its forms and the preservation of forests , fauna and flora.</p>	
--	---------------------------------------	--	--

	Ordinance MMA Nº 43/2014	Establish the National Program for the Conservation of Endangered Species - Pro-Species, with the objective of adopting prevention, conservation , management and management actions, with a view to minimizing threats and the risk of extinction of species.	The instruments of the Pro-Species are: I - Official National Lists of Endangered Species; II - National Action Plans for Conservation of Endangered Species- PAN; III - Databases and information systems aimed at supporting extinction risk assessments, as well as the process of planning actions for conservation.
--	-----------------------------	--	---

	Law Nº 13.052/2014.	It amends Article 25 of Law No. 9,605 of February 12, 1998, which provides for criminal and administrative sanctions arising from conduct and activities harmful to the environment and provides other measures to determine that seized animals are released primarily into their habitat and establish conditions necessary for the welfare of these animals.	"§ 1º Animals shall be released primarily into their habitat or, if such a measure is not feasible or not recommended for sanitary reasons, delivered to zoos, foundations or similar entities, for custody and care under the responsibility of technicians habilitados."
	Ordinance MMA Nº444/2014	Recognize as endangered brazilian fauna species those constant "Official National List of Endangered	<i>Callithrix aurita</i> and <i>C. flaviceps</i> are considered endangered (EN) in the List.

		Fauna Species".	
Conducting Research and Management Activities	JOINT NORMATIVE INSTRUCTION ICMBio/IBAMA 01/2014	It establishes procedures between ICMBio and IBAMA for the management and conservation of Brazilian wildlife species.	“Art.1º Establish, within the scope of the Brazilian Institute of the Environment and Renewable Natural Resources - IBAMA and the Chico Mendes Institute for Biodiversity Conservation - ICMBio, the procedures for the shared use of information and for the complementarity of actions with regard to the management and conservation of wildlife. Single paragraph. The shared use of information will cover its use for environmental licensing and control over the resources exercised by IBAMA and for the actions of authorization, monitoring and conservation of biodiversity promoted by ICMBio.”

	<p>NORMATIVE INSTRUCTION IBAMA 23/2014</p>	<p>It defines the guidelines and procedures for the disposal of wild animals seized, rescued by a competent authority or voluntarily delivered by the population, as well as for the operation of the Wild Animal Triage Centers of IBAMA - CETAS.</p>	<p>Os espécimes híbridos e exóticos apenas poderão ser destinados para instituições de manejo em cativeiro (Artigos 21, 22 e 23).</p>
--	--	--	---

	<p>NORMATIVE INSTRUCTION ICMBio Nº 03/2014</p>	<p>It sets standards for the use of the Authorization and Information System on Biodiversity - SISBio, in the form of the guidelines and conditions provided for in this Normative Instruction, and regulates the availability, access and use of data and information received by the Chico Mendes Institute for Biodiversity Conservation through SISBio.</p>	<p>“Art. 3. Fix a standard on the performance of the following activities, with the purpose of scientific or didactic, in the national territory, on the continental shelf, in the territorial sea and in the exclusive economic zone:</p> <p>I - collection of biological material;</p> <p>II - capture or marking of wild animals in situ;</p> <p>III - temporary maintenance of specimens of wild fauna in captivity;</p> <p>IV - transport of biological material; And</p> <p>V - conducting research in federal conservation unit or in underground natural cavity.</p> <p>§ 1 - The activities with didactic purposes provided for in the caput are restricted to those performed in the scope of higher education.</p> <p>§ 2 - This Normative Instruction does not apply to the collection and transportation of material biological species:</p> <p>I - domesticated or cultivated, except when related to research carried out in federal conservation units in the public domain; And</p> <p>II - exotic wild in ex situ condition.”</p> <p>When the research activities are carried out within state or municipal Conservation Units, in addition to SISBio authorization, it is also necessary to request authorization from the respective competent environmental agencies:</p> <p>- São Paulo: submit a project to the Technical-Scientific Committee of the Forestry Institute (COTEC), in addition to the board of management.</p> <p>- Minas Gerais: follow the standards of the State Forest Institute (IEF), according to Ordinances 130/2017 and 17/2019.</p> <p>- Rio de Janeiro: request the INEA (State Institute of the Environment) based on Ordinance IEF/RJ/PR No. 227/2007.</p>
--	--	---	---

			<p>- Espírito Santo: request the IEMA (State Institute of Environment and Water Resources), based on Decree 4.225-N/1998.</p>
--	--	--	---

Management of hybrids and exotics	RESOLUTION No. 7/2018 OF THE NATIONAL BIODIVERSITY COMMISSION - CONABIO/MMA	Provides on the National Strategy for Invasive Alien Species	Objective of the Strategy: To guide the implementation of measures to prevent the introduction and dispersion and significantly reduce the impact of invasive alien species on Brazilian biodiversity and ecosystem services, control or eradicate invasive alien species. Exotic species - species, subspecies or upper hierarchy is occurring outside its past or present natural distribution area; includes any part, such as gametes, seeds, eggs or propagules that can survive and subsequently reproduce.
	INSTRUÇÃO NORMATIVA ICMBio Nº 6/2019	It provides for the prevention of introductions and the control or eradication of exotic or invasive species in federal Conservation Units and their damping zones.	“Art. 2º The Guidance guide for the management of invasive alien species in Federal Conservation Units is established, available in http://www.icmbio.gov.br/cbc/publicacoes , which includes methods already approved by ICMBio and is considered the guiding document for project analysis.”

	RESOLUTION SMA No. 164/2018 (São Paulo)	It establishes procedures for reproduction of specimens of the <i>genus</i> <i>Callithrix</i> kept in wildlife enterprises in captivity in the State of São Paulo.	It prohibits the production of hybrid individuals of the <i>genus Callithrix</i> and the reproduction of <i>individuals belonging to the species</i> <i>Callithrix jacchus</i> and <i>Callithrix</i> <i>penicillata</i> (exception given only to scientific and commercial breeding sites using these species as matrices, but marketed animals should be sterilized). To <i>C. penicillata</i> reproduction may be authorised only for individuals of known origin and within the programme of conservação, by Zoos and scientific breeding sites for conservation purposes.
Conservation Research and Management (more targeted at endangered <i>Callithrix</i> species)	ORDINANCE ICMBio N° 134/2010	Approve the National Action Plan for the Conservation of Mammals in the Central Atlantic Forest	It comprises objectives and actions for the conservation of 23 mammalian dollar sons, <i>including Callithrix aurita and</i> <i>Callithrix flaviceps</i> , for a period of five years from its date of publication.

	ORDINANCE ICMBio N° 702/2018	Approves the National Action Plan for the Conservation of Atlantic Forest primates and collared sloth - PAN PPMA, contemplating fourteen taxons threatened with extinction, establishing their general objective, specific objectives, lead time, scope and forms of implementation and supervision.	It covers and establishes priority conservation strategies for 13 (thirteen) primate species and for collared sloth, considered endangered, listed in the National List (MMA Ordinance No. 444/2014), including <i>Callithrix aurita</i> and <i>Callithrix flaviceps</i> , for a period of five years, from its date of publication.
--	------------------------------------	---	---

	<p>COOPERATION AGREEMENT ON PROCESS</p> <p>No. 02070.003869/2018-</p> <p>BIODIVERSITY CONSERVATION signed between ICMBio and AZAB</p>	<p>This Cooperation Agreement is based on cooperation in the preparation, implementation, maintenance and coordination of Ex situ Management Programs for Endangered Species in Brazilian Zoos and Aquariums, as detailed in the Work Plan.</p>	<p>Ex Situ Conservation Program for <i>Callithrix aurita</i></p>
--	---	---	--

	<p>RED BOOK OF ENDANGERED BRAZILIAN FAUNA (ICMBio/MMA, 2018)</p>	<p>This edition of the Red Book of Endangered Brazilian Fauna brings the result of the process of evaluation of species coordinated by the Chico Mendes Institute, made official by the Ministry of the Environment through Ordinances No. 444 and 445 of 2014.</p>	<p>Presents factsheets with information on the species officially categorized as endangered, such as <i>Callithrix aurita</i> and <i>Callithrix flaviceps</i>, both considered Endangered (EN).</p>
--	--	---	---

	THE IUCN RED LIST OF THREATENED SPECIES	<p>“The International Union for Conservation of Nature’s Red List of Threatened Species has evolved to become the world’s most comprehensive information source on the global conservation status of animal, fungi and plant species.”</p>	<p>It presents factsheets with information on the species according to the overall assessment. <i>Callithrix aurita</i> e <i>and Callithrix flaviceps</i> were considered Endangered and Critically Endangered (CR), respectively.</p> <p>- Melo, F., Bicca-Marques, J., Ferraz, D. da S., Jerusalinsky, L., Mittermeier, R.A., Oliveira, L.C., Port-Carvalho, M., Ruiz-Miranda, C.R., Valença Montenegro, M., da Cunha, R. & do Valle, R.R. 2020. <i>Callithrix aurita</i> (amended version of 2019 assessment). The IUCN Red List of Threatened Species 2020: e.T3570A166617776. https://dx.doi.org/10.2305/IUCN.UK.2020-1.RLTS.T3570A166617776.en.</p> <p>- <i>Callithrix flaviceps</i>: Ferrari, S., Mendes, S.L. & Rylands, A.B. 2020. <i>Callithrix flaviceps</i>. The IUCN Red List of Threatened Species 2020: e.T3571A17936805. https://dx.doi.org/10.2305/IUCN.UK.2020-2.RLTS.T3571A17936805.en.</p>
--	---	--	---

	<p>CITES - Convention on International Trade in Endangered Species of Wild Fauna and Flora</p>	<p>“CITES (the Convention on International Trade in Endangered Species of Wild Fauna and Flora) is an international agreement between governments . Its aim is to ensure that international trade in specimens of wild animals and plants does not threaten their survival.”</p>	<p><i>Callithrix aurita</i> and <i>Callithrix flaviceps</i> are listed in Appendix I.</p>
--	--	--	---

	<p>THE WORLD’S 25 MOST ENDANGERED PRIMATES: 2018–2020 (2019)</p>	<p>“The tenth iteration of the biennial listing of a consensus of the 25 primate species considered to be among the most endangered worldwide and the most in need of conservation measures.”</p>	<p><i>Callithrix aurita</i> is among the 25.</p> <p>Rodrigo S. Carvalho, Sally J. Fransen, Mônica M. Valença-Montenegro, Nicholas J. Dunn, Cláudia A. Igayara-Souza, Márcio Port-Carvalho, Dominic Wormell, Fabiano R. Melo, Alessandro Silva, Wagner R. Lacerda & Leandro Jerusalinsky. 2019. Buffy-tufted-ear marmoset. <i>Callithrix aurita</i> É. Geoffroy Saint-Hilaire, 1812 Brazil (2018). In: C. Schwitzer, R.A. Mittermeier, A.B. Rylands, F. Chiozza, E.A. Williamson, D. Byler, S. Wich, T. Humle, C. Johnson, H. Mynott and G. McCabe (eds.), <i>Primates in Peril: The World’s 25 Most Endangered Primates 2018–2020</i>, pp. 24-27. IUCN SSC Primate Specialist Group, International Primatological Society, Global Wildlife Conservation, and Bristol Zoological Society, Washington, DC.</p>
--	--	---	---

ACKNOWLEDGEMENTS

We would like to thank Dr. Anthony Rylands for helping translate portions of this text from Portuguese into English.

FUNDING

Portions of this work have been supported by a Brazilian CNPq Jovens Talentos Postdoctoral Fellowship, an American Society of Primatologists Conservation Small Grant, an International Primatological Society Research Grant, a Brazilian CNPq DCR grant (300264/2018-6), a Goldberg Research Grant, and a Marie Skłodowska-Curie Post-Doctoral Fellowship to JM.

REFERENCES

1. Rylands AB, Coimbra-Filho AF, Mittermeier RA. The systematics and distributions of the marmosets (*Callithrix*, *Callibella*, *Cebuella*, and *Mico*) and *Callimico* (*Callimico*) (Callitrichidae, Primates). In: Ford SM, Porter LM, Davis LC, editors. The smallest anthropoids: the marmoset/Callimico radiation. New York: Springer; 2009. p. 25-61.
2. Malukiewicz J, Cartwright RA, Curi NHA et al. Mitogenomic Phylogeny of *Callithrix* with Special Focus on Human Transferred Taxa. bioRxiv 2020.08.12.247692.
3. Buckner JC, Lynch-Alfaro JW, Rylands AB et al. Biogeography of the marmosets and tamarins (Callitrichidae). *Mol Phylogenet Evol.* 2015; 82B:413–425.
4. Malukiewicz J, Boere V, Fuzessy LF et al. Natural and anthropogenic hybridization in two species of eastern Brazilian marmosets (*Callithrix jacchus* and *C. penicillata*). *PLoS ONE.* 2015; 10(6):e0127268.
5. Mendes SL. Padrões biogeográficos e vocais em *Callithrix* do grupo *jacchus* (Primates, Callitrichidae). Doctoral Dissertation. Instituto de Biologia, Universidade Estadual de Campinas (UNICAMP). 1997.
1. Norris D, Rocha-Mendes F, Marques R et al. Density and spatial distribution of buffy-tufted-ear marmosets (*Callithrix aurita*) in a continuous Atlantic Forest. *International Journal of Primatology*, 2011; 32(4):811-829.
2. Ferrari SF, Corre HKM, Coutinho PEG. Ecology of the “southern” marmosets (*Callithrix aurita* and *Callithrix flaviceps*): How different, how similar? In: Norconk MA, Rosenberger

- AL, Garber PA, editors. Adaptive Radiations of Neotropical Primates. New York:Plenum Press; 1996. p. 157–171.
3. Hershkovitz P. Living New World monkeys (Platyrrhini) with an introduction to primates: Vol 1. Chicago Chicago University Press. 1977.
 4. Mittermeier RA, Coimbra-Filho AF, Constable ID et al. Conservation of primates in the Atlantic forest region of eastern Brazil. *Int Zoo Yearb.* 1982; 22:2–17.
 5. Melo F, Bicca-Marques J, Ferraz DS et al. (2020) *Callithrix aurita* (amended version of 2019 assessment). The IUCN Red List of Threatened Species 2020: e.T3570A166617776. <https://dx.doi.org/10.2305/IUCN.UK.2020-1.RLTS.T3570A166617776.en>. Downloaded on 27 July 2020.
 6. Ferrari S, Mendes SL, Rylands AB. (2020) *Callithrix flaviceps*. The IUCN Red List of Threatened Species 2020: e.T3571A17936805. <https://dx.doi.org/10.2305/IUCN.UK.2020-2.RLTS.T3571A17936805.en>. Download on 27 July 2020.
 7. Rylands AB, Fonseca GAB, Leite YL et al. Primates of the Atlantic Forest: Origen, distribution, endemism, and communities. In: Norconk MA, Rosenberger AL, Garber PA, editores. Adaptive Radiations of Neotropical Primates. New York:Plenum Press; 1996. p. 21-51.
 8. Oliveira LC, Grelle CEV. Introduced primate species of an Atlantic Forest region in Brazil: present and future implications for the native fauna. *Tropical Conservation Science.* 2012; 5(1):112-120.
 9. Oliver W LR, Santos IB. Threatened endemic mammals of the Atlantic forest region of south-east Brazil. In: Wildl. Preserv. Trust, Special Scientific Report 4. 1991. 126pp.
 10. Rylands AB, Spironelo W, Tornisielo VL et al. Primates of the Rio Jequitinhonha valley, Minas Gerais, Brazil. *Primate Conserv.* 1988; 9: 100–109.
 11. Vivo M. Taxonomia de *Callithrix Erxleben, 1777* (Callitrichidae. Primates). Belo Horizonte:Fundação Biodiversitas .1991.
 12. Silva FFR, Malukiewicz J, Silva LC et al. A survey of wild and introduced marmosets (*Callithrix*: Callitrichidae) in the southern and eastern portions of the state of Minas Gerais, Brazil. *Primate Conservation.* 2018;32:1–18.
 13. Vale C, Neto LM, Prezoto F. Distribution and invasive potential of the black-tufted marmoset *Callithrix penicillata* in the Brazilian territory. *Scientia Plena.* 2020;16:1-19.
 14. da Rosa CA, Ribeiro BR, Bejarano V. Neotropical alien mammals: a data set of occurrence and abundance of alien mammals in the Neotropics. *Ecology.* 2020. doi:10.1002/ecy.3115
 15. Ferreira FM, Queiroz LP.. Vegetação e flora da Caatinga. *Cienc Cult.* 2018; 70(4): 51-56.
 16. Malukiewicz J. A Review of experimental, natural, and anthropogenic hybridization in *Callithrix* marmosets. *Int J Primatol.* 2019; 40:72–98.
 17. Grelle CEV, Cerqueira R. Determinantes da distribuição geográfica de *Callithrix flaviceps* (Thomas) (Primates, Callitrichidae). *Rev. Bras. Zool.* 2006;23:414-420.
 18. Ferrari SF, Diego VH. Long-term changes in a wild marmoset group. *Folia Primatologica.* 1992; 58(4): 215-218.
 19. Coutinho PE, Corrêa KM. Polygyny in a free-ranging group of buffy-tufted-ear marmosets, *Callithrix aurita*. *Folia primatologica.* 1995; 65(1): 25-29.
 20. Muskin A. Field notes and geographic distribution of *Callithrix aurita* in Eastern Brazil. *Am J Primatol.* 1984;7: 377-380.

21. Cosenza BAP, Melo FR. Primates of the Serra do Brigadeiro State Park, Minas Gerais, Brazil. *Neotropical Primates*. 1998;86(1):18-20.
22. Costa MD, Fernandes FAB, Hilário RR et al. Densidade, tamanho populacional e conservação de primatas em fragmento de Mata Atlântica no sul do estado de Minas Gerais, Brasil. *Iheringia, Série Zoologia, Porto Alegre*. 2012.102(1):5-10.
23. Rylands AB, Faria DS. Habitats, feeding ecology, and home range size in the genus *Callithrix*. In: Rylands AB, editor, Marmosets and tamarins: Systematics, behaviour, and ecology. Oxford: Oxford University Press. 1993. p.262–272
24. Souza VB. Variação do crânio e da mandíbula em *Callithrix* Erxleben, 1777 (Platyrrhini, Callitrichidae): resultados de uma abordagem através de morfometria geométrica. Master Thesis. Universidade Federal de Viçosa. 2016.
25. Martins MM, Setz EZF. Diet of buffy tufted-eared Marmosets (*Callithrix aurita*) in a Forest Fragment in Southeastern Brazil. *International Journal of Primatology*. 2000; 21: 467–476
26. Alves MC. Observações sobre o *Callithrix flaviceps* (Thonas, 1903) na Estação Ecológica de Caratinga - EBC/FBCN, Minas Gerais (Callitrichidae, Primates). In: Mello MT, editor. *A primatologia no Brasil - 2*. Brasília, Sociedade Brasileira de Primatologia. 1986. p. 205-206
27. Ferrari SF. Preliminary report on the field study of *Callithrix flaviceps*, In: Rylands AB, and Bernardes AT, editors. *A Primatologia no Brasil - 3*.1991. Belo Horizonte; Fundação Biodiversitas e Sociedade Brasileira de Primatologia. 1991. p.159-171.
28. Guimarães A. Ecologia, comportamento reprodutivo e marcação de cheiro em um grupo de *Callithrix flaviceps* (Callitrichidae, Primates), na Estacao Biologia de Caratinga, Minas Gerais. Master Thesis. Universidade Federal de Minas Gerais.1998.
29. Hilário RR, Ferrari SF. Feeding ecology of a group of buffy-headed marmosets (*Callithrix flaviceps*): fungi as a preferred resource. *Am J Primatol*. 2010; 72:515-521.
30. Hilario RR, Ferrari SF. Four breeding females in a free ranging group of buffy-headed marmosets (*Callithrix flaviceps*). *Folia Primatol*. 2010; 81:31–40
31. Natori M. Interspecific relationships of *Callithrix* based on the dental characters. *Primates*. 1986; 27:321–336.
32. Ferrari SF. The ecology and behaviour of the buffy-headed marmoset, *Callithrix flaviceps* (O. Thomas, 1903). Doctoral Dissertation. University of London. 1988.
33. Hilário RR, Ferrari SF. Why feed on fungi? The nutritional content of sporocarps consumed by buffy-headed marmosets, *Callithrix flaviceps* (Primates: Callitrichidae), in Southeastern Brazil. *J Chem Ecol*. 2011; 37:145–149.
34. Passamani M, Rylands AB. Home range of a Geoffroy's marmoset group, *Callithrix geoffroyi* (Primates, Callitrichidae) in South-eastern Brazil. *Rev. Brasil. Biol*. 2000; 60:275-281.
35. Passamani M, Rylands AB. Feeding behavior of Geoffroy's marmoset (*Callithrix geoffroyi*) in an Atlantic Forest fragment of South-eastern Brazil. *Primates*. 2000; 41:27-38.
36. Chiarello AG. Mammalian community and forest structure of an Atlantic forest fragment of southeastern Brazil. Doctoral thesis. University of Cambridge. 1997.
37. Taylor A, Vinyard C. Comparative analysis of masseter fiber architecture in tree-gouging (*Callithrix jacchus*) and nongouging (*Saguinus oedipus*) callitrichids. *J. Morphol*. 2004;261:276-285.
38. Canton JM, Hill DM, Hume JD, Crook GA. The digestive strategy of the common marmoset, *Callithrix jacchus*. *Comparative Biochemical Physiology*. 1996; 144:1-8.

39. Garber PA, Caselli CB, McKenney AC, et al. Trait variation and trait stability in common marmosets (*Callithrix jacchus*) inhabiting ecologically distinct habitats in northeastern Brazil. *Am J Primatol*. 2019; 81:e23018.
40. Castro CSS, Araujo A. Diet and feeding behavior of marmoset, *Callithrix jacchus*. *Braz. J. Ecol*. 2007;7:14–19.
41. Abreu F, De la Fuente MFC, Schiel N et al. Feeding ecology and behavioral adjustments: Flexibility of a small neotropical primate (*Callithrix jacchus*) to survive in a semiarid environment. *Mammal Research*; 2016;61:221–229.
42. De la Fuente MFC, Souto A, Sampaio MB et al. Behavioral adjustments by a small Neotropical primate (*Callithrix jacchus*) in a semiarid Caatinga environment. *The Scientific World Journal*. 2014;326524:1–8.
43. Streilein KE. Ecology of small mammals in the semiarid brazilian Caatinga. II. *Water relations*. *Ann Carnegie Mus*.1982; 51(6):109–126
44. Garber PA. Vertical clinging, small body size, and the evolution of feeding adaptations in the Callitrichidae. *Am J Phys Anthropol*.1992;88:469–482
45. Tardif SD, Smucny DA, Abbott DH, Mansfield K, Schultz Darken N, Yamamoto ME. Reproduction in captive common marmosets (*Callithrix jacchus*). *Compar Med*. 2003; 53:364–368.
46. Schiel N, Souto A. The common marmoset: An overview of its natural history, ecology and behavior. *Developmental Neurobiology*. 2017;77(3):244–262.
47. Raboy BE, Canale GR, Dietz JM. Ecology of *Callithrix kuhlii* and a review of eastern Brazilian Marmosets. *Int J Primatol* .2008;29:449–467.
48. Rylands AB. Sympatric Brazilian callitrichids: the black-tufted-ear marmoset, *Callithrix kuhlii*, and the golden-headed lion tamarin, *Leontopithecus chrysomelas*. *Journal of Human Evolution*. 1989;18(7): 679-695.
49. Rodrigues NN, Martinez RA. Wildlife in our backyard: interactions between Wied's marmoset *Callithrix kuhlii* (Primates: Callitrichidae) and residents of Ilhéus, Bahia, Brazil. *Wildlife Biology*. 2014;20:91-96.
50. Fonseca GAB, Lacher TE Jr, Alves C Jr, Magalhães-Castro B. Some ecological aspects of free-living black tufted-ear marmosets (*Callithrix jacchus penicillata*). *Antropologia Contemporâneo*.1980;3:197.
51. Miranda GHB, Faria DS. Ecological aspects of black-pincelled Marmoset (*Callithrix penicillata*) in the Cerradão and dense Cerrado of the Brazilian Central Plateau. *Brazilian Journal Biology*. 2001;61(3):397-404.
52. Silva IO. Socialidade e acesso a recursos alimentares por fêmeas de saguis (*Callithrix penicillata*) em grupos em ambiente natural. Doctoral Dissertation. Universidade de Brasília. 2008.
53. Silva FFR, Malukiewicz J, Silva LC, Carvalho RS, Ruiz-Miranda CR, et al. A survey of wild and introduced marmosets (*Callithrix*: Callitrichidae) in the southern and eastern portions of the state of Minas Gerais, Brazil. *Primate Conservation*. 2018;32: 1–18.
54. Bicca-Marques J, Jerusalinsky L, Mittermeier RA et al. 2018. *Callithrix penicillata*. The IUCN Red List of Threatened Species 2018: e.T41519A17935797. <https://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T41519A17935797.en>. Downloaded on 11 October 2020.
55. Braz AG, Lorini ML Vale, MM. Climate change is likely to affect the distribution but not parapatry of the Brazilian marmoset monkeys (*Callithrix* spp.). *Divers Distrib*. 2019; 25: 536–550.

56. Moraes AM, Vancine MH, Moraes AM et al. Predicting the potential hybridization zones between native and invasive marmosets within Neotropical biodiversity hotspots. *Global Ecol Conserv.* 2019; 20:e00706.
57. Vale CA, Neto LM, Prezoto F. Distribution and invasive potential of the black-tufted marmoset *Callithrix penicillata* in the Brazilian territory. *Scientia Plena.* 2020:052401.
58. Power ML, Ross, CN, Schulkin J, Tardif SD. The development of obesity begins at an early age in captive common marmosets (*Callithrix jacchus*). *Am J Primatol.* 2012;74:261–269.
59. Seferovic M, Sánchez-San Martín C, Tardif SD. et al. Experimental Zika virus infection in the pregnant common marmoset induces Spontaneous fetal loss and neurodevelopmental abnormalities. *Sci Rep.* 2018; 8:6851.
60. Rutherford JN, Tardi SD. Placental efficiency and intrauterine resource allocation strategies in the common marmoset pregnancy. *Am J Phys Anthropol.* 2008;137:60-68.
61. Worley K, Warren W, Rogers J et al. The common marmoset genome provides insight into primate biology and evolution. *Nat Genet.* 2014;46:850–857.
62. Harris RA, Tardif SD, Vinar T et al. Evolutionary genetics and implications of small size and twinning in callitrichine primates. *Proc Natl Acad Sci U S A.* 2014;111(4):1467-72.
63. Saltzman W, Digby LJ, Abbott DH. Reproductive skew in female common marmosets: what can proximate mechanisms tell us about ultimate causes? *Proceedings of the Royal Society B.* 2009;276:389-399.
64. Wislocki GB. Observations on twinning in marmosets. *Am J Anat.* 1939;64: 445-483.
65. Ross CN, French JA, Ortí G. Germ-line chimerism and paternal care in marmosets (*Callithrix kuhlii*). *Proceedings of the National Academy of Sciences.* 2007; 104:6278-6282.
66. Benirschke K, Anderson JM, Brownhill LE. Marrow chimerism in marmosets. *Science.* 1962;138(3539):513-5.
67. Sweeney CG, Curran E, Westmoreland SV, Mansfield KG, Vallender EJ. Quantitative molecular assessment of chimerism across tissues in marmosets and tamarins. *BMC Genomics.* 2012;13:98.
68. Bell JT, Pai AA, Pickrell JK, Gaffney DJ et al. DNA methylation patterns associate with genetic and gene expression variation in HapMap cell lines. *Genome Biol.* 2011;12(1):R10.
69. Feil R, Fraga MF. Epigenetics and the environment: emerging patterns and implications. *Nat Rev Genet.* 2012 Jan 4;13(2):97-109.
70. Ben-Shlomo R. Invasiveness, chimerism and genetic diversity. *Molecular Ecology.* 2017;26(23):6502–6509.
71. Abreu F, Souto A, Schiel N. Wild common marmosets (*Callithrix jacchus*) employ spatial cognitive abilities to improve their food search and consumption: an experimental approach in small-scale space. *Primates.* 2020. doi: 10.1007/s10329-020-00826-1. Epub ahead of print. PMID: 32383127.
72. Šlipogor V, Gunhold-de Oliveira T, Tadić Z et al. Consistent inter-individual differences in common marmosets (*Callithrix jacchus*) in Boldness-Shyness, Stress-Activity, and Exploration-Avoidance. *Am J Primatol.* 2016;78(9):961-73.
73. Inoue-Murayama M, Yokoyama C, Yamanashi Y et al.. Common marmoset (*Callithrix jacchus*) personality, subjective well-being, hair cortisol level and AVPR1a, OPRM1, and DAT genotypes. *Sci Rep.* 2018;8(1):10255.

74. Šlipogor V, Massen JJM, Schiel N et al. (accepted). Temporal consistency and ecological validity of personality structure in common marmosets (*Callithrix jacchus*): a unifying field and laboratory approach. *American Journal of Primatology*.
75. del Rosario RC, Rayan NA, Prabhakar S. Noncoding origins of anthropoid traits and a new null model of transposon functionalization. *Genome Res.* 2014;24(9):1469-84.
76. Jayakumar V, Ishii H, Seki M. et al. An improved de novo genome assembly of the common marmoset genome yields improved contiguity and increased mapping rates of sequence data. *BMC Genomics.* 2020 Apr 2;21(Suppl 3):243.
77. Koren S, Rhie A, Walenz BP et al.. De novo assembly of haplotype-resolved genomes with trio binning. *Nat Biotechnol.* 2018;10.1038/nbt.4277.
78. Yoshimatsu S, Sato T, Yamamoto M et al. Generation of a male common marmoset embryonic stem cell line DSY127-BV8VT1 carrying double reporters specific for the germ cell lineage using the CRISPR-Cas9 and PiggyBac transposase systems. *Stem Cell Res.* 2020;44:101740.
79. Chin CS, Peluso P, Sedlazeck FJ et al. Phased diploid genome assembly with single-molecule real-time sequencing. *Nat Methods.* 2016;13(12):1050-1054.
80. Housman G, Quillen EE, Stone AC. Intraspecific and interspecific investigations of skeletal DNA methylation and femur morphology in primates. *Am J Phys Anthropol.* 2020;173(1):34-49.
81. Narapareddy L, Wildman DE, Armstrong DL et al. Maternal weight affects placental DNA methylation of genes involved in metabolic pathways in the common marmoset monkey (*Callithrix jacchus*). *Am J Primatol.* 2020;82(3):e23101.
82. Mashiko H, Yoshida AC, Kikuchi SS et al. Comparative anatomy of marmoset and mouse cortex from genomic expression. *J Neurosci.* 2012;32(15):5039-53.
83. Matsunaga E, Nambu S, Oka M, Iriki A. Comparative analysis of developmentally regulated expressions of Gadd45a, Gadd45b, and Gadd45g in the mouse and marmoset cerebral cortex. *Neuroscience.* 2015 Jan 22;284:566-580.
84. Law AJ, Pei Q, Feldon J et al. Gene expression in the anterior cingulate cortex and amygdala of adolescent marmoset monkeys following parental separations in infancy. *Int J Neuropsychopharmacol.* 2009 Jul;12(6):761-72.
85. Housman G, Gilad Y. Prime time for primate functional genomics. *Current opinion in genetics and development.* 2020; 62:1-7.
86. Bakken TE, Jorstad NL, Hu Q et al. Evolution of cellular diversity in primary motor cortex of human, marmoset monkey, and mouse. *BioRxiv*, 2020.03.31.016972.
87. Geirsdottir L, David E, Keren-Shaul H et al. Cross-Species Single-Cell Analysis Reveals Divergence of the Primate Microglia Program. *Cell.* 2019. 2;179(7):1609-1622.e16.
88. Aravalli RN, Steer CJ. Utility of common marmoset (*Callithrix jacchus*) embryonic stem cells in liver disease modeling, tissue engineering and drug metabolism. *Genes (Basel).* 2020;11(7):729.
89. Debowski K, Warthemann R, Lentjes J et al. Non-viral generation of marmoset monkey iPS cells by a six-factor-in-one-vector approach. *PLoS One.* 2015;10(3):e0118424.
90. Vermilyea SC, Guthrie S, Meyer M et al. Induced pluripotent stem cell-derived dopaminergic neurons from adult common marmoset fibroblasts. *Stem Cells Dev.* 2017;26(17):1225-1235.
91. Okano H, Sasaki E, Yamamori T et al. Brain/MINDS: A Japanese national brain project for marmoset neuroscience. *Neuron.* 2016;92(3):582-590.

92. Shimogori T, Abe A, Go Y et al. Digital gene atlas of neonate common marmoset brain. *Neurosci Res.* 2018;128:1-13.
93. Koroshetz W, Gordon J, Adams A et al. The State of the NIH BRAIN Initiative. *J Neurosci.* 2018;38(29):6427-6438.
94. Kishi N, Sato K, Sasaki E, Okano H. Common marmoset as a new model animal for neuroscience research and genome editing technology. *Dev Growth Differ.* 2014 Jan;56(1):53-62.
95. Krienen FM, Goldman M, Zhang Q et al. Innovations in Primate Interneuron Repertoire. bioRxiv 709501; doi: <https://doi.org/10.1101/709501>
96. Boroviak T, Stirparo GG, Dietmann S et al. Single cell transcriptome analysis of human, marmoset and mouse embryos reveals common and divergent features of preimplantation development. *Development.* 2018;145(21):dev167833.
97. Klein J, Figueroa F. Evolution of the major histocompatibility complex. *Crit Rev Immunol.* 1986;6(4):295-386.
98. Walter L. Major histocompatibility complex class-I-interacting natural killer cell receptors of nonhuman primates. *J Innate Immun.* 2011;3(3):236-41.
99. Shiina T, Kono A, Westphal N et al. Comparative genome analysis of the major histocompatibility complex (MHC) class I B/C segments in primates elucidated by genomic sequencing in common marmoset (*Callithrix jacchus*). *Immunogenetics.* 2011;63(8):485-99.
100. Kono A, Brameier M, Roos C et al. Genomic sequence analysis of the MHC class I G/F segment in common marmoset (*Callithrix jacchus*). *J Immunol.* 2014;192(7):3239-46.
101. van der Wiel MK, Otting N, de Groot NG et al. The repertoire of MHC class I genes in the common marmoset: evidence for functional plasticity. *Immunogenetics.* 2013;65(12):841-9.
102. Averdam A, Petersen B, Rosner C, Neff J et al. A novel system of polymorphic and diverse NK cell receptors in primates. *PLoS Genet.* 2009;5(10):e1000688.
103. Hart BA, Kap YS. An essential role of virus-infected B cells in the marmoset experimental autoimmune encephalomyelitis model. *Mult Scler J Exp Transl Clin.* 2017;3(1):2055217317690184.
104. Pinto MA, Marchevsky RS, Baptista ML et al. Experimental hepatitis A virus (HAV) infection in *Callithrix jacchus*: early detection of HAV antigen and viral fate. *Exp Toxicol Pathol.* 2002;53(6):413-20.
105. Falk L, Deinhardt F, Wolfe L et al. Epstein-Barr virus: experimental infection of *Callithrix jacchus* marmosets. *Int J Cancer.* 1976;17(6):785-8.
106. Sutherland SD, Almeida JD, Gardner PS et al.. Rapid diagnosis and management of parainfluenza I virus infection in common marmosets (*Callithrix jacchus*). *Lab Anim.* 1986;20(2):121-6.
107. Bukh J, Apgar CL. Five new or recently discovered (GBV-A) virus species are indigenous to New World monkeys and may constitute a separate genus of the Flaviviridae. *Virology.* 1997;229(2):429-36.
108. Tilston-Lunel NL, Hughes J, Acrani GO et al. Genetic analysis of members of the species Oropouche virus and identification of a novel M segment sequence. *J Gen Virol.* 2015;96(Pt 7):1636-50.
109. Muniz CP, Zheng H, Jia H et al. A non-invasive specimen collection method and a novel simian foamy virus (SFV) DNA quantification assay in New World primates reveal

aspects of tissue tropism and improved SFV detection. *PLoS One*. 2017 Sep 1;12(9):e0184251.

110. Guzman MG, Gubler DJ, Izquierdo A et al. Dengue infection. *Nat Rev Dis Primers*. 2016;2:16055.
111. Omatsu T, Moi ML, Takasaki T et al. Changes in hematological and serum biochemical parameters in common marmosets (*Callithrix jacchus*) after inoculation with dengue virus. *J Med Primatol*. 2012;41(5):289-96.
112. Ferreira MS, de Castro PH, Silva GA et al. *Callithrix penicillata*: a feasible experimental model for dengue virus infection. *Immunol Lett*. 2014;158(1-2):126-33.
113. Vasconcelos BC, Vieira JA, Silva GO, et al. Antibody-enhanced dengue disease generates a marked CNS inflammatory response in the black-tufted marmoset *Callithrix penicillata*. *Neuropathology*. 2016 Feb;36(1):3-16.
114. Dick GWA. Trans Roy Soc Trop Med Hyg. 1952;46: 521–534.
115. Chiu CY, Sánchez-San Martín C, Bouquet J et al. Experimental Zika Virus Inoculation in a New World Monkey Model Reproduces Key Features of the Human Infection. *Sci Rep*. 2017;7(1):17126.
116. Chippaux JP, Chippaux A. Yellow fever in Africa and the Americas: a historical and epidemiological perspective. *J Venom Anim Toxins Incl Trop Dis*. 2018;24:20.
117. Freitas ACP, Oviedo-Pastrana ME, da Rocha DA et al. Diagnóstico de animais ilegais recebidos no centro de triagem de animais silvestres de Belo Horizonte, Estado de Minas Gerais, no ano de 2011. *Ciência Rural*; 2015;45(1):163-170.
118. Silva NIO, Sacchetto L, de Rezende IM et al. Recent sylvatic yellow fever virus transmission in Brazil: the news from an old disease. *Viol J*. 2020;17(1):9.
119. Mares-Guia MAMM, Horta MA, Romano A et al. Yellow fever epizootics in non-human primates, Southeast and Northeast Brazil (2017 and 2018). *Parasit Vectors*. 2020;13(1):90.
120. Cavalcante KRLJ, Tauil PL. Epidemiological characteristics of yellow fever in Brazil, 2000-2012. *Epidemiol. Serv. Saúde, Brasília*. 2016;25(1):11-20.
121. Cunha MS, da Costa AC, de Azevedo Fernandes NCC et al. Epizootics due to Yellow Fever Virus in São Paulo State, Brazil: viral dissemination to new areas (2016-2017). *Sci Rep*. 2019;9(1):5474.
122. Figueiredo LTM. Emergent arboviruses in Brazil. *Rev. Soc. Bras. Med. Trop*. 2007;40(2): 224-229.
123. Moreira-Soto A, Torres MC, Lima de Mendonça MC et al. Evidence for multiple sylvatic transmission cycles during the 2016-2017 yellow fever virus outbreak, Brazil. *Clin Microbiol Infect*. 2018;1019.e1-1019.e4.
124. Abreu FVS, Ferreira-de-Brito A, Azevedo AS et al. Survey on Non-Human Primates and Mosquitoes Does not Provide Evidences of Spillover/Spillback between the Urban and Sylvatic Cycles of Yellow Fever and Zika Viruses Following Severe Outbreaks in Southeast Brazil. *Viruses*. 2020;12(4):364.
125. Favoretto SR, Araujo DB, Duarte NFH et al. Zika Virus in Peridomestic Neotropical Primates, Northeast Brazil. *Ecohealth*. 2019 Mar;16(1):61-69.
126. Terzian ACB, Zini N, Sacchetto L, Rocha RF et al. Evidence of natural Zika virus infection in neotropical non-human primates in Brazil. *Sci Rep*. 2018;8(1):16034.

127. Guth S, Hanley KA, Althouse BM et al.. Ecological processes underlying the emergence of novel enzootic cycles: Arboviruses in the neotropics as a case study. *PLOS Neglected Tropical Diseases*.2020;14(8):e0008338.
128. Costa EA, Luppi MM, Malta MC et al.Outbreak of human herpesvirus type 1 infection in nonhuman primates (*Callithrix penicillata*). *J Wildl Dis*. 2011;47(3):690-3.
129. Longa CS, Bruno SF, Pires AR et al. Human herpesvirus 1 in wild marmosets, Brazil, 2008. *Emerg Infect Dis*. 201;17(7):1308-10.
130. Mätz-Rensing K, Jentsch KD, Rensing S et al.. Fatal Herpes simplex infection in a group of common marmosets (*Callithrix jacchus*). *Vet Pathol*. 2003;40(4):405-11.
131. Kotait I, Oliveira RN, Carrieri ML et al. Non-human primates as a reservoir for rabies virus in Brazil. *Zoonoses Public Health*.2019;66(1):47-59.
132. Moutinho FFB, Correa DMS, Marcanth N, Nunes VMA, Boges FVB, Neto FF, Bruno SF. Surveillance of epizootic diseases in non-human primates in Niterói, RJ, Brazil. *Revista Brasileira de Geografia Médica e da Saúde Hygeia*.2020;16:37-48.
133. Aguiar TDF, Costa EC, Rolim BN, Romijn PC, Moraes NB, Teixeira MFS. Risks of transmitting rabies virus from captive domiciliary common marmoset (*Callithrix jacchus*) to human beings, in the metropolitan region of Fortaleza, State of Ceará, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical*. 2011; 44(3):356-363.
134. Carroll D, Daszak P, Wolfe ND et al. The Global Virome Project. *Science*. 2018.359(6378):872-874.
135. D'arc M, Moreira FRR, Dias CA et al.. The characterization of two novel neotropical primate papillomaviruses supports the ancient within-species diversity model. *Virus Evol*. 2020;6(1):veaa036.
136. Monteiro da Cruz MAO. Dinâmica reprodutiva de uma população de saguis-do-Nordeste (*Callithrix jacchus*) na Estação Ecológica do Tapacurá, PE. São Paulo University Thesis. São Paulo, 1998.
137. Faulkes CG, Arruda MF, da Cruz MAOM Genetic Structure Within and Among Populations of the Common Marmoset, *Callithrix jacchus*: Implications for Cooperative Breeding. In: Ford SM; Porter LM, Davis LC, editors. The Smallest Anthropoids: The Marmoset/Callimico Radiation. 2009. p. 103-117.
138. Thompson CL, Hrit R, Melo LCO et al. Callitrichid responses to dead and dying infants: the effects of paternal bonding and cause of death. *Primates*. 2020 Sep;61(5):707-716. doi: 10.1007/s10329-020-00824-3. Epub 2020 May 14. PMID: 32409994.
139. Brasil. 1967. Lei nº 5.197, de 3 de janeiro de 1967. Dispõe sobre a proteção à fauna e dá outras providências. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 3 jan. 1967. Disponível em: http://www.planalto.gov.br/ccivil_03/leis/L5197compilado.htm
140. Bastos LF, Luz VLF, dos Reis I J et al. Apreensão de espécimes da fauna silvestre em Goiás–situação e destinação. *Revista De Biologia Neotropical*. 2008; 5(2):51-63.
141. São Paulo (estado). Secretaria de Estado do Meio Ambiente. Resolução. SMA Nº 164, DE 27 DE NOVEMBRO DE 2018. Estabelece procedimentos para reprodução de espécimes do gênero *Callithrix* mantidos em empreendimentos de fauna silvestre em cativeiro no Estado de São Paulo. Disponível em: <file:///C:/Users/Usuario/Downloads/Resolu%C3%A7%C3%A3o%20SMA%20164%20-%202018-processo-2029-2016-estabelece-procedimentos-para-reproducao-de-callithrix.pdf> Acesso 08 de setembro de 2020.

142. RENTAS. 2001. 1º Relatório Nacional Sobre o Tráfico de Fauna Silvestre. RENTAS, Brasília. Website: http://www.rentas.org.br/wp-content/uploads/2014/02/REL_RENTAS_pt_final.pdf Accessed 08 September 2020.
143. Levacov D, Jerusalinsky L, Fialho MDS. 2011. Levantamento dos primatas recebidos em Centros de Triagem e sua relação com o tráfico de animais silvestres no Brasil. In: Melo FR, Mourthé I, editors. A Primatologia no Brasil – 11, Belo Horizonte: Sociedade Brasileira de Primatologia, 2011. pp.281–305.
144. Silva NM. Espécimes recebidos no centro de triagem de animais silvestres de Salvador, BA durante os anos de 2012 a 2014. Monograph. Federal University of Bahia. 2015.
145. de Moraes Jr MM, Ruiz-Miranda CR, Grativol AD et al. Os sagüis, *Callithrix jacchus penicillata*, como espécies invasoras na região de ocorrência do mico-leão dourado. Conservação do mico-leão-dourado. 2008;28820:86.
146. Ruiz Miranda CR, De Moraes Jr MM, de Paula VR et al. O problema dos saguis introduzidos no Rio de Janeiro: vítimas ou vilões? *Ciência Hoje*, 2011;48:44-49.
147. Ruiz-Miranda, CR, Affonso AG, Martins A et al. Distribuição do sagui (*Callithrix jacchus*) nas áreas de ocorrência do mico-leão-dourado (*Leontopithecus rosalia*) no estado do Rio de Janeiro. *Neotropical Primates*, 2000;8: 98–100.
148. Fuzessy LF, Silva IO, Malukiewicz, J et al (2014). Morphological variation in wild marmosets (*Callithrix penicillata* and *C. geoffroyi*) and their hybrids. *Evolutionary Biology*, 2014;41:480–493.
149. Carvalho RS, Bergallo HG, Cronemberger C et al. *Callithrix aurita*: A tiny primate on the edge of extinction in the Brazilian Atlantic Forest. *Neotropical Primates*. 2018; 24:1–8.
150. Carvalho RS. Conservação do saguis-da-serra-escuro (*Callithrix aurita* (Primates)) – Análise molecular e colorimétrica de populações do gênero *Callithrix* e seus híbridos. PhD dissertation, Universidade do Estado do Rio de Janeiro. 2015
151. Secco H, Grilo C, Bager A. Habitat selection by the black-tufted marmoset *Callithrix penicillata* in human-disturbed landscapes. *Journal of Tropical Ecology*. 2018;34(02):135-144.
152. Detogne N, Ferregueti AC, Mello JFF. et al Spatial distribution of buffy-tufted-ear (*Callithrix aurita*) and invasive marmosets (*Callithrix* spp.) in a tropical rainforest reserve in southwestern Brazil. *American Journal of Primatology*. 2017;79:e22718.
153. Paula HMG, Távora RS, Almeida MV et al. Estudos preliminares da presença de sagüis no município de Bauru, São Paulo, Brasil. *Neotrop. Primates*. 2005; 13: 6–1.
154. Crispo E, Moore J-S, Lee-Yaw JA et al. Broken barriers: human-induced changes to gene flow and introgression in animals: an examination of the ways in which humans increase genetic exchange among populations and species and the consequences for biodiversity. *BioEssays*. 2011; 33:508–518. doi:10.1002/bies.201000154
155. Lyra-Neves RM, Oliveira MAB, Telino-Júnior WR et al. Comportamentos interespecíficos entre *Callithrix jacchus* (Linnaeus) (Primates, Callitrichidae) e algumas aves de Mata Atlântica, Pernambuco, Brasil. *Revista Brasileira de Zoologia*. 2007; 24(3):709-716.
156. Silva IdO, Alvarenga ABB, Boere V. Occasional Field Observations of the Predation on Mice, Dove and Ants by Black-Tufted-Ear Marmosets (*Callithrix penicillata*). *Neotropical Primates*. 2008;15(2):59-62.
157. Sales IdS, Ruiz-Miranda CR, Santos CdP. Helminths found in marmosets (*Callithrix penicillata* and *Callithrix jacchus*) introduced to the region of occurrence of golden lion tamarins (*Leontopithecus rosalia*) in Brazil. *Veterinary Parasitology*. 2010;171(1-2):123-129.

158. Silva FFR. Distribuição do gênero *Callithrix* no estado de Minas Gerais: introdução de espécies e hibridação. 2014. Master Thesis. Federal University of Viçosa.
159. Rhymer JM, Simberloff D. Extinction by hybridization and introgression. *Annual Review of Ecology and Systematics*. 1996;27; 83-109
160. McFarlane SE, Pemberton JM. Detecting the true extent of introgression during anthropogenic hybridization. *Trends Ecol Evol*. 2019;34(4):315-326.
161. Allendorf F, Leary R, Spruell P et al. The problems with hybrids: setting conservation guidelines. *TREE*.. 2001;16: 613–622.
162. Todesco M, Pascual MA, Owens GL, et al. Hybridization and extinction. *Evol Appl*. 2016; 9: 892-908. doi:10.1111/eva.12367
163. Bezerra B, Bicca-Marques J, Miranda J. et al. (2018) *Callithrix jacchus*. The IUCN Red List of Threatened Species 2018: e.T41518A17936001. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T41518A17936001.en>
164. Bicca-Marques J, Jerusalinsky L, Mittermeier RA et al. (2018) *Callithrix penicillata*. The IUCN Red List of Threatened Species 2018: e.T41519A17935797. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T41519A17935797.en>
165. Beuchle R, Grecchi RC, Shimabukuro Y et al. Land cover changes in the Brazilian Cerrado and Caatinga biomes from 1990 to 2010 based on a systematic remote sensing sampling approach. *Applied Geography*. 2015;58,2015:116-127,
166. Rezende C, Scarano LFR, Assad ED et al. From hotspot to hopespot: An opportunity for the Brazilian Atlantic Forest. *Perspectives in Ecology and Conservation*. 2018;16;208-214
167. Crouzeilles R, Ferreira MS, Chazdon RL et al. Ecological restoration success is higher for natural regeneration than for active restoration in tropical forests. *Sci Adv*. 2017;3(11):e1701345.
168. Jerusalinsky L, Melo FR. Conservação de primatas no Brasil: perspectivas e desafios. In: Urbani B, Kowalewski M, Cunha RGT et al, editors.. *La primatología en Latinoamérica 2 – A primatologia na America Latina 2*. Tomo I Argentina-Colombia. Ediciones IVIC. Caracas: Instituto Venezolano de Investigaciones Científicas (IVIC). 2018. p. 161-186.
169. Nieves L, Bicca-Marques J, Jerusalinsky L et al. (2019) *Callithrix kuhlii*. The IUCN Red List of Threatened Species 2019: e.T3575A17936243. <http://dx.doi.org/10.2305/IUCN.UK.2019-3.RLTS.T3575A17936243.en>.
170. Carvalho RS, Franssen SJ, Valença-Montenegro MM et al. Buffy-tufted-ear marmoset *Callithrix aurita* É. Geoffroy Saint-Hilaire, 1812, Brazil. In: Schwitzer C, Mittermeier RA, Rylands AB, et al, editors. *Primates in Peril: The world's 25 most endangered primates 2018-2020*. 2019. p.79-81.
171. Bechara IM. Abordagens metodológicas em Biogeografia da Conservação para avaliar risco de extinção de espécies: um estudo de caso com *Callithrix aurita* (Primates: Callitrichidae). Master Thesis. Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ. 20123.
172. Possamai CB, Mendes SL, Strier, KB. Decline of a primate community following a yellow fever outbreak in the Brazilian Atlantic Forest. In: Lutz C, Fragaszy D, Wood EK, editors *Program of the Forty-Second Meeting of The American Society of Primatologists*. Wiley. 2019; p. 36-37.
173. Rylands AB, Mendes SL. (2018) *Callithrix geoffroyi*. The IUCN Red List of Threatened Species 2018: e.T3572A17936610. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T3572A17936610.en>

174. Estrada A, Garber PA, Rylands AB et al. Impending extinction crisis of the world's primates: Why primates matter. *Sci Adv.* 2017 Jan 18;3(1):e1600946. doi: 10.1126/sciadv.1600946. PMID: 28116351; PMCID: PMC5242557.
175. Estrada A, Garber PA, Mittermeier RA et al. Primates in peril: the significance of Brazil, Madagascar, Indonesia and the Democratic Republic of the Congo for global primate conservation. *PeerJ.* 2018 Jun 15;6:e4869. doi: 10.7717/peerj.4869.
176. Teixeira B, Hirsch A, Goulart VDLR et al. Good neighbours: distribution of black-tufted marmoset (*Callithrix penicillata*) in an urban environment. *Wildlife Research.* 2015; 42:579-589.
177. Duarte MHL, Young RJ: Sleeping Site Selection by Urban Marmosets (*Callithrix penicillata*) under conditions of exceptionally high predator density. *International Journal of Primatology.* 2011; 32:29–334.
178. Goulart VDLR, Teixeira CP, Young RJ: Analysis of callouts made in relation to wild urban marmosets (*Callithrix penicillata*) and their implications for urban species management. *European Journal of Wildlife Research.* 2010;56(4):641-649.
179. http://www.planalto.gov.br/ccivil_03/decreto/2002/d4339.htm
180. Brasil. 2009. Portaria Conjunta MMA/ICMBIO nº 316, de 09 de setembro de 2009. Dispõe sobre os instrumentos de implementação da Política Nacional da Biodiversidade voltados para a conservação e recuperação de espécies ameaçadas de extinção. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 10 set. 2009. Disponível em: https://www.mma.gov.br/estruturas/179/_arquivos/portaria_316_mma_icmbio_179.pdf
181. Brasil. 2010. Portaria ICMBio nº 134 de 23 de dezembro de 2010. Aprova o Plano de Ação Nacional para a Conservação dos Mamíferos da Mata Atlântica Central. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 24 dez. 2010. Disponível em: <https://www.icmbio.gov.br/portal/images/stories/docs-pan/pan-mamac/1-ciclo/pan-mamac-portaria-aprovacao.pdf>
182. Brasil. 2018. Portaria ICMBio nº 702 de 21 de março de 2018. Aprova o Plano de Ação Nacional para a Conservação dos Primatas da Mata Atlântica e da Preguiça-de-coleira. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 22 mar. 2018. Disponível em: https://www.in.gov.br/web/guest/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/7526450/do1-2018-03-22-portaria-n-702-de-21-de-marco-de-2018-7526446
183. Brasil. 2014. Instrução Normativa ICMBio/IBAMA nº01, de 08 de dezembro de 2014. Estabelece procedimentos entre o ICMBio e o IBAMA para o manejo e a conservação de espécies da fauna silvestre brasileira. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 19 dez. 2014. Disponível em: https://www.icmbio.gov.br/cepsul/images/stories/legislacao/Instrucao_normativa/2014/in_icmbio_ibama_01_2014_procedimentos_entre_icmbio_ibama_manejo_conserva%C3%A7%C3%A3o_fauna_br.pdf
184. Brasil. 2014. Instrução Normativa IBAMA nº23, de 31 de dezembro de 2014. Define as diretrizes e os procedimentos para a destinação de animais silvestres apreendidos, resgatados por autoridade competente ou entregues voluntariamente pela população, bem como para o funcionamento dos Centros de Triagem de Animais Silvestres do IBAMA - CETAS. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 2 jan. 2015. Disponível em: https://www.in.gov.br/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/32094529/do1-2015-01-02-instrucao-normativa-n-23-de-31-de-dezembro-de-2014-32094512

1. Brasil. 2018. Resolução da Comissão Nacional de Biodiversidade – CONABIO/MMA nº 7, de 29 de maio de 2018. Aprova a Estratégia Nacional para Espécies Exóticas Invasoras. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 13 jun. 2018. Disponível em:<https://www.mma.gov.br/images/arquivo/80049/Conabio/Resolucoes/RESOLUCAO%20N%207%20DE%2029%20DE%20MAIO%20DE%202018%20-%20Diario%20Oficial%20da%20Uniao%20-%20Imprensa%20Nacional.pdf>
2. Brasil. 2019. Instrução Normativa ICMBio nº 6, de 25 de julho de 2019. Regulamenta o inciso XX do artigo 2º do Anexo I do Decreto 8.974, de 24 de janeiro de 2017, que dispõe sobre a prevenção de introduções e o controle ou erradicação de espécies exóticas ou invasoras em Unidades de Conservação federais e suas zonas de amortecimento. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 19 ago. 2019. Disponível em:<https://www.in.gov.br/en/web/dou/-/instrucao-normativa-n-6-de-25-de-julho-de-2019-211213911#:~:text=Regulamenta%20o%20inciso%20XX%20do,e%20suas%20zonas%20de%20amortecimento.>
3. Brasil. 2018. Resolução CONAMA nº 489, de 26 de outubro de 2018. Define as categorias de atividades ou empreendimentos e estabelece critérios gerais para a autorização de uso e manejo, em cativeiro, da fauna silvestre e da fauna exótica. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 29 out. 2018. Disponível em:https://www.in.gov.br/materia/-/asset_publisher/Kujrw0TZC2Mb/content/id/47542644/do1-2018-10-29-resolucao-n-489-de-26-de-outubro-de-2018-47542603