

Article

Influence of a major mountainous landscape barrier (Mount Cameroon) on the spread of metabolic (*GSTE2*) and target-site (*Rdl*) resistance alleles in the African malaria vector *Anopheles funestus*

Nathalie Amvongo-Adjia ^{1,2,3,6,*}, Jacob M. Riveron ³, Flobert Njiokou ^{1,3}, Samuel Wanji ^{2,4} and Charles S. Wondji ^{3,5,*}

¹ Department of Animal Biology and Physiology, Faculty of Science, University of Yaoundé 1, P.O. BOX 812 Yaoundé, Cameroon; amvongo.n@gmail.com, njiokou@yahoo.com

² Research Foundation for Tropical Diseases and Environment (REFOTDE), University of Buea, P.O. BOX 474 Buea, Cameroon; swanji@yahoo.fr

³ Centre for Research in Infectious Diseases (CRID), LSTM Research Unit, P.O. Box 13591 Yaoundé, Cameroon; jacob.riveron_miranda@syngenta.com

⁴ Department of Microbiology and Parasitology, University of Buea, P.O. BOX 63 Buea, Cameroon;

⁵ Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool L35QAUK, United Kingdom; charles.wondji@lstmed.ac.uk

⁶ Medical Research Centre, Institute of Medical Research and Medicinal Plants Studies (IMPM), P.O. BOX 13033 Yaoundé, Cameroon.

* Correspondence: amvongo.n@gmail.com (N.A.A); Charles.wondji@lstmed.ac.uk (C.S.W.)

Abstract: Increased levels of insecticide resistance in major malaria vectors such as *Anopheles funestus* threaten the effectiveness of insecticide-based control programmes. Understanding the landscape features impacting the spread of resistance makers is necessary to design suitable resistance management strategies. Here we examined the influence of the highest mountain in West Africa (Mount Cameroon; 4,100 meters elevation) on the spread of metabolic and target-site resistance alleles in *An. funestus* populations. Vector composition varied across the four localities surveyed along the altitudinal cline with major vectors exhibiting high parity rate (80.5%). *Plasmodium* infection rates ranged from 0.79% (*An. melas*) to 4.67% (*An. funestus*). High frequencies of *GSTE2^R* (67% - 81%) and *Rdl^R* (49% - 90%) resistance alleles were observed in *An. funestus* throughout the study area, with *GSTE2^R* frequency increasing with altitude whereas the opposite is observed for *Rdl^R*. Patterns of genetic diversity and population structure analyses revealed high levels of polymorphisms with 12 and 16 haplotypes respectively for *GSTE2* and *Rdl*. However, the reduced diversity patterns of resistance allele carriers revealed signatures of positive selection on the two genes across the study area irrespective of the altitude. Despite slight variations associated with the altitude, the spread of resistance alleles suggest that control strategies could be implemented against malaria vectors across mountainous landscapes.

Keywords: Malaria, *Anopheles funestus*, Insecticide resistance, Mount Cameroon.

1. Introduction

Vector control of mosquitoes is a critical part of the global strategy to manage mosquito-associated diseases, and insecticides are the most important component in this effort. Four insecticide classes namely organochlorines, organophosphates, carbamates and pyrethroids are mainly recommended for use against adult mosquitoes [1]. Pyrethroids are the most widely used insecticides for indoor spraying against mosquitoes worldwide, owing to their efficacy and safety [2]. They are the mainly chemicals approved to treat mosquito nets [3, 4], the main tool for human protection from malaria-carrying mosquitoes especially in Africa [1]. In the past, massive sprayings

of insecticides greatly limited mosquito-borne diseases and even eradicated malaria in a few areas [5, 6]. However, the widespread development of resistance in mosquitoes to the most commonly used insecticides negatively impacted the fight against mosquitoes in many areas [7]. This has resulted in a number of outbreaks of mosquito-borne diseases in recent years [8, 9].

In Cameroon, malaria is the leading cause of morbidity and mortality accounting for an estimated 6.2 million clinical cases and 12,500 deaths [10]. *Anopheles funestus sensu stricto* (s.s.) is one of the four major malaria vectors in the country [11, 12], being mostly prevalent in the Sudan savanna domain. However, this species had been found transmitting *Plasmodium falciparum* malaria parasite in Cameroon highlands such as the Mount Cameroon region [13] and the locality of Santchou in the western Cameroon [14]. The resistance profile of *An. funestus* s.s. has previously been explored for some populations, with multiple resistance to pyrethroids, dichlorodiphenyltrichloroethane (DDT) and carbamates reported in the localities of the soudano-sahelian [15-17] and forested [18, 19] zones of Cameroon. Two mutations, the *L119F* of the glutathione s-transferases epsilon 2 (GSTe2) gene conferring resistance to DDT and pyrethroids [20], and the *A296S* of the γ -aminobutyric acid (GABA) gene implicated in resistance to dieldrin (*Rdl*) [15], have been identified in *An. funestus* s.s. African populations. These markers have been respectively associated to separated resistance mechanisms: the increased metabolic detoxification of insecticides commonly known as metabolic resistance, and the decreased sensitivity of the target proteins on which an insecticide acts, so called target-site resistance [21]. Additionally, cytochromes *P450s* genes have been associated to pyrethroids resistance in the same vector [22, 23].

The spread of these resistance alleles are not uniformed across the African continent, with *A296S-Rdl* and *L119F-GSTe2* being present in West and Central Africa but absent in southern Africa [15, 20], whereas the *CYP6P9a/b* resistance alleles present an opposite distribution [24, 25]. Presence of barriers to gene flow has been suggested to explain the restriction in the spread of these alleles [26, 27]. Major landscape modifications such as Rift Valley have been suggested as the main continent-wide barrier to the spread of resistance alleles in *An. funestus* [26, 27]. It remains to know whether other major but local landscape modifications such as major mountain chains could also restrict the spread of these resistance alleles. The Mount Cameroon chain with its highest peak at 4,100 meters above the sea level (m a.s.l.), is the highest mountain in West Africa and thus constitutes a major landscape variation in the region with potential to impact patterns of gene flow between populations of species across this region as. This potential influence on the spread of resistance alleles across the region remains unknown.

The Mount Cameroon region is also an area of prime importance to the Cameroon agriculture sector with industrial banana, rubber and palm plantations, activities relying on the intensive use of agrochemicals likely to select resistance in local mosquito populations. This has already been shown in *An. coluzzii* and *An. gambiae* which exhibit high resistance level to pyrethroids in this region [28]. Thus, similar resistance profile is expected in *An. funestus* vectors.

Understanding the influence of the high altitude of Mount Cameroon in the spread of resistance alleles could help to improve the management of related resistance. Here we assess the impact of the Mount Cameroon on the spread of both metabolic and target-site resistance by assessing the genetic diversity of both GSTe2 and GABA receptor genes across an altitudinal transect. Based on analysis of GSTe2 and *Rdl* markers, genotype and allele frequency distributions were compared between *An. funestus* s.s. populations collected in four localities across the Mount Cameroon region. Furthermore, we assessed signatures of selection these genes and established pattern of genetic structure between the four *An. funestus* s.s. Beside this, the *Plasmodium* infection rate of *An. funestus* s.s. was also assessed.

2. Materials and Methods

2.1. Study site and adult mosquito sampling

The Mount Cameroon is the highest active volcano in western Africa, rising from the Atlantic Ocean at the Gulf of Guinea and having an elevation of 4095 meters above the sea level (m a.s.l.). The study was conducted in low, mid and high altitude areas within the lowland rain forest from sea level to 800 meters elevation across the Mount Cameroon area (precisely in the Fako division, South west region of Cameroon), having different landscape and environmental patterns [29]. Entomological surveys were restricted at this altitudinal range (from sea level to 800 m a.s.l.) due to previous surveys where the absence of *Anopheles* mosquitoes above 800 m elevation (precisely in the localities of Bokova and Bonakanda situated between 800 and 870 m a.s.l.) was reported during four consecutive seasons [30].

The localities surveyed during this study were: Tiko village (4°3' N, 9°22' E and elevation 9 m a.s.l.) and Likomba (4°5' N, 9°20' E and elevation 70 m a.s.l) located in the Tiko municipality and considered as lowlands (Tiko village and Likomba were further considered as a single collection site: Tiko), Mutengene (4°05'57" N, 9°18'29" E, altitude 220 m a.s.l.) and Meanja (4°16' N, 9°23' E, altitude 305 m a.s.l.) respectively located in the Tiko and Muyuka municipalities were considered as mid altitude areas, and Likoko (4°8'41" N, 9°13'38" E, elevation 800 m a.s.l.) in the Buea municipality, a highland area. Localities of Tiko, Likomba, Mutengene and Likoko follow an altitudinal transect on the southwest and west edge of the mountain whereas Meanja was selected as a sided-elevated area on the eastern edge of the mountain to assess if contrasting events occur across the mountain (Figure 1). Malaria transmission is perennial in low and mid altitude areas whereas in highland it is heterogeneous [31, 32]. The region has an equatorial climate modified by a double influence of the Atlantic Ocean and the mountain and it is characterized by a unique rainy season from March to October while the dry season goes from November to February [33]. *Anopheles gambiae*, *An. coluzzii* and *An. melas* sibling species of the *An. gambiae* s.l. complex, and *An. funestus* s.s. are the prevalent malaria vectors [13, 34, 35].




Figure 1. Map of the study area

Prior to the entomological field activities, authorisations were sought from village's chiefs, sub-chiefs and quarter's heads. The permission to collect mosquitoes from households was obtained from the household heads. Mosquito collectors were invited to sign a consent form before participating in night collections and could withdraw whenever wanted. In addition, presumptive malaria prophylaxes were given to them at the end of collections.

Female adult *Anopheles* mosquitoes were collected from the year 2010 to 2014 (2010: May; 2011: February-May-August-November; 2012: February-August-November; 2013: August-November; 2014: February) (Supplementary Table S1), indoor and outdoor households between 06:00 PM to 06:00 AM using the landing catch on volunteers' technique as previously reported [35].

2.2. Species identification and *Plamodium* infection rate

Female *Anopheles* caught were morphologically identified according to the key of Gillies and de Meillon [36], and ovaries were dissected for physiological age-grading [37]. For each specimen identified as belonging to either the *An. gambiae* complex or the *An. funestus* group, genomic DNA was extracted following the DNA extraction buffer protocol [38] on whole mosquitoes. Sibling species of the *An. gambiae* complex were distinguished using conventional PCR [39] and PCR-RFLP [40] of the ribosomal intergenic spacer IGS of the nuclear rDNA gene, whereas members of the *An. funestus* group were characterized by amplifications of the internal transcribed spacer ITS2 [41]. Mosquito infection status was assessed using TaqMan assay as described by Bass, *et al.* [42] and confirmed using multiplex single-round [43] and nested [44] PCR assays.

2.3. Genotyping of *GSTe2* and *Rdl* resistance markers

GSTe2 mutation associated to DDT and pyrethroids' resistance in *An. funestus* was genotyped using an allele-specific PCR (AS-PCR) assay previously described by Tchouakui, *et al.* [45]. The same was done for the *Rdl* mutation conferring resistance to dieldrin as described by Riveron, *et al.* [46]. The primers used for these assays can be found as Supplementary Table S2. Frequency distribution of *GSTe2* and *Rdl* genotypes and alleles were compared using the Chi-square (χ^2) test, with statistical significance set at $p < 0.05$.

2.4. Polymorphism analysis of resistance genes

2.4.1. Genetic variability of *An. funestus* s.s. across the Mount Cameroon region based on *GSTe2* and *Rdl* full gene sequencing

Full-length sequences (all exons and introns) of *GSTe2* and *Rdl* were individually amplified for forty samples (ten samples per locality each: Tiko, Mutengene, Meanja and Likoko) according to previous protocols [20, 46]. Details of the primers used are provided in Supplementary Table S2. Thereafter, successful amplicons were purified using Exo-SAP clean up protocol (ThermoFisher Scientific, Santa Clara, CA95051, USA) and directly sequenced on both strands. Other *An. funestus* DNA samples have previously been sequenced based on *GSTe2* mutation in Benin, Cameroon, Ghana, Malawi, Mozambique and Uganda in a separated study [20]. These sequences were used in the present study to assess the genetic variability patters of *GSTe2* in *An. funestus* populations across Africa. The polymorphic positions were identified through a manual analysis of sequence chromatograms using BioEdit 7.2.5 [47] based on sequence differences in multiple alignments using ClustalW [48]. Genetic diversity parameters (number of polymorphic sites S^* , number of haplotypes h , haplotype diversity Hd , Synonymous mutations Syn , Non-synonymous mutations $Nsyn$, nucleotide diversity π , mean number of nucleotide differences k ; Tajima D and Fu and Li F^* statistics) were computed using dnaSP 5.10 [49]. Haplotype networks were then built using the TCS program [50, 51] to assess the connection between haplotypes. The level of pairwise genetic differentiation were estimated using the K_{ST} statistics [52] as implemented in dnaSP 5.10. The significance of the K_{ST} estimates were assessed by permutation of subpopulation identities and re-calculating K_{ST} 10,000 times.

2.4.2. Phylogenetic trees of haplotypes

Prior to the construction of the Maximum-Likelihood (ML) phylogenetic trees for *GSTe2* and *Rdl* haplotypes, the best-fit substitution model for each dataset was assessed based on the Bayesian Information Criterion (BIC) in MEGA 10.1.6 [53]. These analyses indicated that the Tamura

3-parameter, Kimura 2-parameter and Hasegawa-Kishino-Yano models best described *GSTe2* haplotype dataset of Mount Cameroon, *GSTe2* haplotype dataset for Africa-wide analysis and *Rdl* haplotype dataset, respectively. These models were then used to build the respective ML tree using MEGA 10.1.6 with 500 bootstrap replications for the robustness of the trees. Neighbour-Joining trees were also constructed with pairwise *GST* genetic distances [54] between subpopulations still in MEGA 10.1.6.

3. Results

3.1. Mosquito species composition

A total of 4,911 female mosquitoes were collected across the study localities during the four seasons of collections (Supplementary Figure S1 and Table S1). Four mosquito genera were found present in the Mount Cameroon region, these are: 13.5% *Mansonia* spp., 19.9% *Culex* spp., 1.6% *Aedes* spp. and 65% *Anopheles* spp. The highest number of mosquitoes was collected from Mutengene ($n = 2,094$) and the least number from Likoko ($n = 319$). A significant difference based on Chi-square test was noted in mosquito distribution between all four sites ($p < 0.0001$).

Morphological identification of the 3,194 females *Anopheles* revealed that they mostly belong to two major malaria vector species, *An. gambiae* complex (86%) and *An. funestus* group (12.2%) and three secondary vector species (1.6% *An. hancocki*, 0.2% *An. nili* s.l. and 0.03% *An. ziemanni*) (Supplementary Figure S2), with a significant difference noted in the frequency distributions of *Anopheles* sp. between localities ($p < 0.0001$). PCR-species identification performed from 3,135 females morphologically identified as *An. gambiae* s.l. and *An. funestus* s.l. revealed that three *An. gambiae* siblings coexist in the study area: *An. coluzzii* (44.7%), *An. gambiae* (28.6%) and *An. melas* (12%), in addition to 23 hybrids *An. coluzzii* × *An. gambiae* (0.7%). Whereas, all *An. funestus* s.l. individuals were identified as *An. funestus* s.s.

3.2. Parity and *Plasmodium* infection rates

Of the 3,135 female *Anopheles* identified as either *An. funestus* s.s. or belonging to the *An. gambiae* complex, the tracheal filament of 2,490 (79.4%) were analysed for physiological age determination (parity rate). Of these, 2,004 (80.5%) dissected females were found parous (Supplementary Figure S3). The highest parity rate was observed in *An. melas* (87%; 301/346), whereas 82.5% (558/676) *An. gambiae*, 81% (243/300) *An. funestus* s.s., 77.2% (885/1147) *An. coluzzii* and 76.2% (16/21) *An. coluzzii* × *An. gambiae* hybrids were detected parous. Parity rates showed significant differences between dissected mosquito species ($p = 0.003$) and between the surveyed localities except for the lowland Tiko and Likoko highland ($p = 0.764$).

Similarly, a total of 2,746 *An. gambiae* s.l. specimens and 364 *An. funestus* s.s. were screened for *P. falciparum* (falcip+) and *P. ovale*/*P. vivax*/*P. malariae* (OVM+) using TaqMan assay on genomic DNA extracted from whole mosquitoes (Supplementary Figure S3). An overall infection rate (IR) of 2.35% (73/3110) was obtained, with *An. funestus* s.s. being more infected (4.67%, 17/364) while 0.79% *An. melas* (3/381), 1.61% *An. coluzzii* (23/1428), 3.17% *An. gambiae* (29/914) and 4.35% *An. coluzzii* × *An. gambiae* hybrids (1/23) were tested positive for *Plasmodium* spp. infection. Of the 73 *Anopheles* mosquitoes tested positive for *Plasmodium* spp. infection, 57 mosquitoes were falcip+ (78.1%), 14 OVM+ (19.2%) and 2 were Pf/OVM+ (2.7%). The multiplex PCR assays performed with infected mosquitoes confirmed TaqMan results and determined that the 16 OVM+ individuals specifically corresponds to 15 *P. malariae* and one *P. vivax*. Mosquitoes presenting mixed infection Pf/OVM+ were mix-infected with *P. falciparum* and *P. malariae*. A significant difference was noted between the infection rates of the different *Anopheles* species tested ($p = 0.04$). However, no significant differences were observed in mosquito's infection rates between Tiko/Meanja ($p = 0.203$) and Mutengene/Likoko ($p = 0.863$).

3.3. Detection of mutations associated to DDT and dieldrin resistance in *An. funestus* s.s. mosquitoes

The presence of *GSTe2* and *Rdl* mutations was investigated respectively in 339 and 218 *An. funestus* s.s. specimens across the study area. Genotyping results showed high frequencies of *GSTe2*^R (66.7% - 97.1%) (Error! Reference source not found.A) and *Rdl*^R (68.4% - 90%) (Error! Reference source not found.B) resistance cases in *An. funestus* s.s. populations, with an overall resistant cases estimated at 93.5% (317/339) for *GSTe2* gene and 74.3% (162/218) for *Rdl* gene. It was interesting to note that land elevation seems to reversely influence the occurrence of both mutations across the study area. In fact, the frequency of *GSTe2* resistant population increased with altitude whereas the occurrence of dieldrin resistant specimens decreased with altitude. Conversely, while the frequency of *GSTe2* susceptible specimen decreased with altitude, the proportion of dieldrin susceptible mosquitoes increased with climb in altitude. However, no significant differences based on Chi-square test were noted in the overall distribution of both markers in *An. funestus* s.s. populations across the study area (*GSTe2* $p = 0.446$ and *Rdl* $p = 0.543$).




Figure 2. Frequency distribution of *GSTe2* (A) and *Rdl* (B) mutations in *An. funestus* s.s. populations from the localities surveyed.

Specifically, *GSTe2* mutation had an overall frequency of 64.3% (218/339) for homozygote resistant mosquitoes (RR) and 29.2% (99/339) heterozygotes (RS) whereas 6.5% (22/339) were homozygotes susceptible (SS). A predominance of homozygote resistant genotypes was observed throughout the study area being 50% (3/6) in Tiko, 68.2% (180/264) in Mutengene, 45.7% (16/35) in Meanja and 55.9% (19/34) in Likoko (Error! Reference source not found.A). Interestingly, an increase in the frequencies of *GSTe2* heterozygotes was noted with the climb in altitude whereas the occurrence of susceptible mosquitoes decreases with land elevation. Meanwhile for *Rdl* mutation, although homozygotes resistant mosquitoes almost dominated across the study area (Tiko: 90% (9/10), Mutengene: 40.6% (25/128), Meanja: 64.3% (27/42), Likoko: 28.9% (11/38)), frequencies of heterozygotes and susceptible mosquitoes followed reverse tendencies than that of *GSTe2*. Here, the frequencies of heterozygotes seemed to decrease with altitude while the occurrence of susceptible mosquitoes increased with land elevation. For both makers, Chi-square test statistical analysis of genotype frequency distributions between localities showed significant differences except in Tiko/Meanja (*GSTe2*: $p = 0.273$, *Rdl*: $p = 0.244$), Mutengene/Likoko (*GSTe2*: $p = 0.17$, *Rdl*: $p = 0.423$) and Meanja/Likoko (only *GSTe2*: $p = 0.354$).




Figure 3. Genotype (A) and allele (B) frequency distributions of *GSTe2* and *Rdl* mutations in *An. funestus* s.s. populations for the different localities surveyed (altitude in m a.s.l.).

Analysis of the allele frequencies of *GSTe2* and *Rdl* mutations in *An. funestus* s.s. populations from the four localities revealed same pattern than those obtained with genotyping results (**Error! Reference source not found.B**). *GSTe2^R* and *Rdl^R* resistant alleles were predominantly represented in almost all the collection sites except in Likoko where the *Rdl^S* susceptible allele was predominant (with a frequency of 0.51). However, both markers showed contrary evolution in their alleles' frequency distributions when considering land elevation. For resistant alleles, while the frequency of *GSTe2^R* seemed to increase with altitude, *Rdl^R* frequency decreased with altitude. Rather for susceptible alleles, as the *GSTe2^S* frequency is decreasing with altitude, the *Rdl^S* increased with land elevation.

3.4. Analysis of the polymorphism of *GSTe2* and *Rdl* genes across Mount Cameroon *An. funestus* s.s. populations

3.4.1. Sequence analysis of full length *GSTe2* and *Rdl* genes

GSTe2 and *Rdl* full length fragments were successfully amplified in 36 and 34 (respectively for both makers) *An. funestus* s.s. genomic DNA samples with specific primers for both mutations (**Error! Reference source not found.2**). The alignment and comparison of sequences obtained with that referenced in Genbank confirmed the presence of glutathione S-transferase Epsilon 2 protein of *An. funestus* (AHC31021.1) and exon 7 encoding the M2 transmembrane domain region of *An. funestus* GABA-receptor gene (AZB49494.1) respectively for *GSTe2* and *Rdl* amplicons.

A point mutation (CTT to TTT) at position codon 119, inducing an amino acid change of leucine to phenylalanine (L119F) and which confers resistance to pyrethroids in *An. funestus* s.s. mosquitoes was observed in all *GSTe2* resistant samples of the Mount Cameroon (Figure 4A). Likewise, the GCA

(alanine) to TCA (serine) mutation at position codon 296 (A296S) was observed in *An. funestus* s.s. dieldrin resistant samples (Figure 4B).




Figure 4. Sequence chromatograms of a 729 bp fragment in *GSTe2* gene (A) and a 1006 bp fragment of the GABA-receptor gene (B); for susceptible (top), homozygote resistant (middle) and heterozygote (bottom) *An. funestus* s.s.

3.4.2. Haplotype distribution of the *GSTe2* gene

A total of 14 variable or polymorphic sites which led to the formation of 12 haplotypes were observed within the 729 bp fragment of *GSTe2* gene of *An. funestus* s.s. mosquitoes across the Mount Cameroon region (Figure 5A and Table 1). The nucleotide sequences of the haplotypes were submitted to Genbank (accession numbers: MN562756, MN562757, MN562760, MN562764 – MN562766, MN562768 – MN562771, MN562774 and MN562775).

Overall, the *GSTe2* polymorphism level was average (haplotype diversity: $Hd = 0.56$). The number of haplotypes (h) and its associated diversity indice (Hd) seemingly decreased (0.68 – 0.51) from the lowland Tiko (9 – 70 m a.s.l.) to Likoko situated at the highest altitude (800 m a.s.l.). Samples carrying L119 susceptible allele were found highly polymorphic ($h = 9$ and $Hd = 0.91$) as compared to 119F resistant allele carriers ($h = 5$ and $Hd = 0.36$).

Table 1. Genetic variability parameters of *GSTe2* full mutation.

Samples	N	S*	h (Hd)	Syn	NSyn	π (k)	D	F*
<i>Per allele</i>								
S	18	11	9 (0.91)	4	2 (D61E, L119F)	0.003 (2.69)	- 0.84 ^{ns}	- 0.70 ^{ns}
R	54	6	5 (0.36)	2	1 (L119F)	0.001 (0.60)	- 1.36 ^{ns}	- 0.27 ^{ns}
<i>Per locality</i> (altitude in m a.s.l.)								
Tiko (9 – 70)	12	8	6 (0.68)	4	1 (L119F)	0.003 (1.97)	- 1.38 ^{ns}	- 1.77 ^{ns}
Mutengene (220)	20	8	7 (0.58)	4	1 (L119F)	0.002 (1.38)	- 1.31 ^{ns}	- 2.13 ^{ns}
Meanja (305)	20	7	5 (0.56)	2	2 (D61E, L119F)	0.001 (1.04)	- 1.55 ^{ns}	- 2.54 ^{ns}
Likoko (800)	20	7	5 (0.51)	1	1 (L119F)	0.002 (1.32)	- 1.40 ^{ns}	- 0.76 ^{ns}
<i>All</i>	72	14	12 (0.56)	6	2	0.002 (1.36)	- 1.62 ^{ns}	- 1.02 ^{ns}

N = number of sequences (2n); S*, number of polymorphic sites; h, number of haplotypes (Hd = haplotype diversity); Syn, Synonymous mutations; Nsny, Non-synonymous mutations; π , nucleotide diversity (k = mean number of nucleotide differences); D and F* Tajima's and Fu and Li's statistics; ns, not significant; S = susceptible; R = resistant; m a.s.l. = meters above the sea level.

Haplotype networks (Figures Figure 5B and Figure 5C) were built using TCS software. In one hand, haplotype network representation with respect to collection sites (Figure 5B) showed the

presence of a unique major haplotype (H2: 47/72 sequences) distributed from lowland to highland across the study area. The ancestral haplotype (H1) appeared in lowland (Tiko: 9 – 70 m a.s.l.; one sequence) and midlands (Mutengene: 220 m a.s.l., 2 sequences and Meanja: 305 m a.s.l., 4 sequences) but not in highland (Likoko: 800 m a.s.l.). Three haplotypes (H4, H9 and H11) were half-shared between Tiko and Mutengene located at 9 to 220 m a.s.l.; another one haplotype (H3) was concomitantly found in both Tiko (9 – 70 m a.s.l.; one sequence), Meanja (305 m a.s.l.; one sequence) and Likoko (800 m a.s.l.; 2 sequences), and one other (H5) only found in Meanja (one sequence) and Likoko (2 sequences). Out of the twelve haplotypes identified, five occurred as singletons and were distributed as follows: two (H10 and H12) in Mutengene (220 m a.s.l.), one (H8) in Meanja (305 m a.s.l.) and two (H6 and H7) in Likoko (800 m a.s.l.) (Figure 5A).




Figure 5. Genetic diversity and polymorphism patterns of *GSTe2* DNA sequences across the Mount Cameroon. (A) Haplotype diversity patterns of the 729bp *GSTe2* fragment in Tiko (Tik), Mutengene (Mut), Meanja (Mea) and Likoko (Lik); H = haplotype; R = resistant; S = susceptible; polymorphic sites are in red. TCS haplotype networks showing haplotype's generation within *GSTe2* gene in *An. funestus* s.s with respect to localities (B) and allelic profiles (C): haplotypes are presented in circular shape scaled to reflect their respective frequencies; Ancestral haplotype in red. Maximum likelihood phylogenetic trees of *GSTe2* DNA sequences among localities (D) and allelic profiles (E) showing an apparent separation between susceptible samples and resistant ones. (F) Neighbour-joining tree of the genetic distances showing a genetic relatedness to the landscape along the Mountain altitudinal transect with Tiko and Mutengene (9 – 220 m a.s.l., both located on the southwest edge) clustering together than Likoko (800 m a.s.l., west edge) and Meanja (305 m a.s.l., eastern edge).

Comparatively, haplotype network representation with respect to the allele type (either susceptible or resistant) carried by the sequences analysed (Figure 5C) showed that out of the twelve haplotypes recorded only two (H1 and H2) were common to both L119 and 119F carriers whereas three haplotypes (H3, H6 and H9) were strictly found in 119F carriers and the remaining seven haplotypes (H4, H5, H7, H8, and H10 – H12) appeared only in L119 carriers. It was noted that common haplotypes to both L119 and 119F carriers were predominantly found in resistance sequences (H1: 4 sequences of 119F and 3 sequences of L119; H2: 41 sequences of 119F and 6 sequences of L119). Also, out of five haplotypes which appeared as singletons, four were strictly associated to L119 carriers (H7, H8, H10 and H12) whereas only one haplotype (H6) originated from a 119F carrier.

Evidence of selection acting on *GSTE2* gene could be noted as 119F-resistant allele carriers exhibited low diversity parameters ($h = 5$, $Hd = 0.36$ and $\pi = 0.001$), unless negative and non-significant values were obtained from Tajima D and Fu and Li F* neutrality tests (Table 1).

- *Gste2* gene Africa-wide analysis

The alignment of the 72 *GSTE2* sequences analysed in the present work with 90 other sequences [17] carrying the same mutation and previously reported in Africa (Malawi: 18 sequences, Mozambique: 10 sequences, Uganda: 10 sequences, Benin: 24 sequences, Ghana: 16 sequences and North Cameroon: 12 sequences) revealed the presence of a total of 58 polymorphic sites resulting to the formation of 53 haplotypes across Africa (Supplementary Figure S4 and Table S3). The most prevalent haplotype H18 (48/162 sequences) appeared in both L119-susceptible (8 sequences) and 119F-resistance (40 sequences) allele carriers. It was shared by Mount Cameroun (8 sequences of L119 and 39 sequences of 119F) and Ugandan (one 119F-sequence) populations (Supplementary Figure S5). Another major haplotype H3 (16/162 sequences) was found in both L119 (11 sequences) and 119F (5 sequences) allele carriers of Mount Cameroon (two L119 and five 119F sequences), Malawi (six L119-sequences), Mozambique (two L119-sequences) and Uganda (one L119-sequence); whereas two minor haplotypes still observed in L119 and 119F allele carriers were one shared by Mount Cameroon/Malawian samples (H4: one L119 sequence from Malawi and one 119F sequence from Mount Cameroon-Mutengene) and the other only noted in the Mount Cameroon area (H50: one L119-Mutengene sequence and one 119F-Tiko sequence). TCS haplotype networks also showed the presence of a major resistant haplotype (H26: 32/162 sequences) restricted to West and Central African populations (22 in Benin, 5 in Ghana, 4 from Mont Cameroon and one from North Cameroon). Overall, it was noted a high proportion of singletons (39/53 haplotypes) being mostly of L119-susceptible allele carriers (30/39 haplotypes); thus suggesting a high polymorphism with reduced numbers of mutational steps, in addition to a signature of selection within the *GSTE2* gene across some *An. funestus* African populations where a *GSTE2*-based resistance to pyrethroids/DDT has been reported (haplotype diversity, $Hd = 0.86$; nucleotide diversity $\pi = 0.005$; Tajima D = -2.03 with $p < 0.05$) (Supplementary Figures S4 and S5 and Table S3).

The construction of maximum-likelihood (ML) phylogenetic trees revealed that haplotypes cluster according to allelic profiles and irrespectively of their geographical origin. Rather, on a neighbour-joining (NJ) distance tree Mount Cameroon clustered alongside the North Cameroon *An. funestus* s.s. population, whereas the East-South (Uganda – Malawi and Mozambique) and West (Benin and Ghana) African populations formed separated clusters (Supplementary Figure S5).

3.4.3. Haplotype distribution of the GABA-receptor gene across Mount Cameroon populations of *An. funestus* s.s.

Analysis of 68 sequences for the 1006 bp fragment of the GABA-receptor gene (accession numbers: Genbank MN562780 – MN562795) showed the presence of 16 polymorphic sites and the formation of an equal number of haplotypes (Figure 6A and Table 2). Genetic variability parameters with respect to allelic profiles and among the four tested *An. funestus* s.s. populations showed that the high number of haplotypes occurred in A296-susceptible allele carriers ($h = 10$, with an equivalent high haplotype diversity $Hd = 0.90$) and for the localities of Mutengene (at 220 m a.s.l., $h =$

9 and $Hd = 0.85$) and Likoko (at 800 m a.s.l., $h = 7$ and $Hd = 0.83$) while the lowest number of haplotypes was found in Tiko ($h = 1$, $Hd = 0.00$) at the lowest base of the Mount Cameroon (9 – 70 m a.s.l.) with no polymorphic site detected.

Table 2. Genetic variability parameters of *Rdl* full mutation.

Samples	N	S*	h (Hd)	Syn	NSyn	π (k)	D	F*
<i>Per Allele</i>								
S	25	11	10 (0.90)	0	0	0.003 (2.53)	- 0.44 ^{ns}	- 0.53 ^{ns}
R	43	6	6 (0.30)	1	0	0.0005 (0.54)	- 1.61 ^{ns}	- 1.65 ^{ns}
<i>Per locality</i> (altitude in m a.s.l.)								
Tiko (9 – 70)	12	0	1 (0.00)	0	0	0 (0.00)	n.a.	n.a.
Mutengene (220)	20	9	9 (0.85)	1	1 (A296S)	0.002 (1.97)	- 0.76 ^{ns}	- 0.39 ^{ns}
Meanja (305)	20	10	6 (0.68)	0	1 (A296S)	0.002 (2.00)	- 1.02 ^{ns}	0.04 ^{ns}
Likoko (800)	16	8	7 (0.83)	0	1 (A296S)	0.002 (2.38)	- 0.05 ^{ns}	- 0.21 ^{ns}
<i>All</i>	68	16	16 (0.71)	1	1	0.002 (1.85)	- 1.32 ^{ns}	- 1.87 ^{ns}

N = number of sequences (2n); S*, number of polymorphic sites; h, number of haplotypes (Hd = haplotype diversity); Syn, Synonymous mutations; Nsyn, Non-synonymous mutations; π , nucleotide diversity (k = mean number of nucleotide differences); D and F* Tajima's and Fu and Li's statistics; ns, not significant; S = susceptible; R = resistant; m a.s.l. = meters above the sea level.

The most common haplotype (H2: 36/68 sequences) was found to be carrying the 296S-resistant allele and was distributed throughout the study area (Figure 6A, Figure 6B and Figure 6C), unlike the ancestral haplotype (H4: 6/36 sequences) which carried the A296-susceptible allele and distributed at low (Tiko: 9 – 70 m a.s.l.) and mid (Mutengene: 220 m a.s.l.; Meanja: 305 m a.s.l.) altitude but not in highland (Likoko: 800 m a.s.l.). Approximately one third of the haplotypes (6/16) appeared as singletons shared half by A296 (H6, H12 and H13) and 296S (H7, H8 and H15) allele carriers, and identified from mid (Mutengene: H12, H13 and H15; Meanja: H8) to highland (Likoko: H6 and H7).




Figure 6. Genetic diversity and polymorphism patterns of the GABA-receptor gene across the Mount Cameroon area. (A) Haplotype diversity patterns of a 1006bp fragment of the GABA-receptor gene in Tiko (Tik), Mutengene (Mut), Meanja (Mea) and Likoko (Lik); H = haplotype; R = resistant; S = susceptible; polymorphic sites are in red. TCS haplotype networks showing a high polymorphism of the GSTe2 gene in *An. funestus* s.s with reduced number of mutational steps between haplotypes with respect to localities (B) and allelic profiles (C); haplotypes are presented in circular shape scaled to reflect their respective frequencies; Ancestral haplotype in red. Maximum likelihood phylogenetic trees of *Rdl*-DNA sequences among localities (D) and allelic profiles (E) showing a marked separation between A296 and 296S allele carriers. (F) Neighbour-joining tree of the genetic distances

showing an apparent genetic relatedness associated with the altitude along the Mount Cameroon with Tiko being genetically differentiated from other localities.

Overall, there were negative and non-significant values of the selection test from Tajima D and Fu and Li F* (Table 2). However, the positive but not significant F* in Meanja (F*: 0.04), in addition to low values of haplotype and nucleotide diversity recorded in 296S-resistant allele carriers ($H_d = 0.30$ and $\pi = 0.0005$ respectively) could be indicators of an ongoing selection acting on the GABA receptor gene within the Mount Cameroon *An. funestus* s.s. populations.

3.5. Population structure at *GSTe2* and *Rdl* mutations in *An. funestus* s.s. across Mount Cameroon

Analysis of the genetic structure at the 729bp *GSTe2* fragment supported the genetic variability observed between low (Tiko), mid (Mutengene and Meanja) and highland (Likoko). Construction of the maximum likelihood (ML) phylogenetic tree with respect to localities (Figure 5D) showed the presence of major consensus cluster across the Mount Cameroon, in addition to three apparent separated clusters formed by haplotypes of low- to midland, midland and mid- to highland. Whereas the construction of ML tree with respect to allelic profile (Figure 5E) highlighted the reduced diversity of 119F resistant allele carriers across the study area. The patterns of clustering according to locations was further supported by low values of genetic differentiation estimates ($-0.0006 \leq K_{ST} \leq 0.010$; all not significant) obtained between the four localities tested and the Nm gene flow index which showed a marked genetic closeness between Tiko/Mutengene (low- and midland) and Meanja/Likoko (mid- and highland) as illustrated on the NJ tree of genetic distances (Figure 5F and Table 3).

Table 3. Patterns of genetic differentiation between *An. funestus* s.s. populations based on K_{ST} estimates from *GSTe2* mutation with (Nm).

	Tiko	Mutengene	Meanja
Mutengene	- 0.027 ^{ns} (0.00)		
Meanja	0.002 ^{ns} (57.05)	0.003 ^{ns} (43.40)	
Likoko	- 0.007 ^{ns} (0.00)	0.010 ^{ns} (12.19)	- 0.0006 ^{ns} (0.00)

PERMTEST calculates Hudson's K_{ST} statistic of genetic differentiation. K_{ST} is equal to $12KS/KT$, where KS is a weighted mean of K_1 and K_2 (mean number of differences between sequences in subpopulations 1 and 2, respectively) and KT represents the mean number of differences between two sequences regardless of their subpopulation; ns, not significant.

Similarly, analysis of the ML phylogenetic tree at the GABA –receptor for *Rdl* mutation based on localities highlighted the presence of a main consensus cluster across the Mount Cameroon and three other apparent clusters generated by midland, mid-/highland and highland haplotypes (Figure 6D). Whereas the ML tree with respect to allelic profiles highlighted the marked reduced diversity of 296S resistant allele carriers (Figure 6E). The construction of a NJ tree of genetic distance (Figure 6F) revealed that the lowland Tiko clusters separately from other localities aligned following the same altitudinal transect (Mutengene and Likoko) or situated eastward of the mountain (Meanja), all at higher elevation level (200 – 800 m a.s.l.). This latter observation was further supported by the consistency of high and significant values of genetic differentiation estimates observed between Tiko and the other three localities ($0.037 < K_{ST} < 0.158$) (Table 4).

Table 4. Patterns of genetic differentiation between *An. funestus* s.s. populations based on K_{ST} estimates from *Rdl* mutation with (Nm).

	Tiko	Mutengene	Meanja
Mutengene	0.125** (1.24)		
Meanja	0.037* (2.08)	0.0098 ^{ns} (15.82)	
Likoko	0.158** (1.16)	0.007 ^{ns} (25.82)	0.035* (14.54)

PERMTEST calculates Hudson's K_{ST} statistic of genetic differentiation. K_{ST} is equal to $12KS/KT$, where KS is a weighted mean of $K1$ and $K2$ (mean number of differences between sequences in subpopulations 1 and 2, respectively) and KT represents the mean number of differences between two sequences regardless of their subpopulations. *, $0.01 < P < 0.05$; **, $0.001 < P < 0.01$; ns, not significant.

4. Discussion

The implementation of effective insecticide resistance management strategies relies on the good understanding of the direction and speed of spread of resistance alleles among mosquito populations. This study assessed the influence of the Mount Cameroon on the spread of both GST-mediated metabolic resistance and *Rdl*-based target-site resistance among population of *An. funestus* s.s. malaria vector.

Results from this study correlates with observations from previous reports [35, 55] indicating the predominance of *Anopheles* mosquitoes within the overall mosquito fauna found in the Mount Cameroon region. *Anopheles* mosquitoes were collected throughout the study area (from lowland to highland), and a total of seven *Anopheles* species were identified; these included: sibling species (*An. coluzzii*, *An. gambiae* and *An. melas*) of the *An. gambiae* sensu lato (s.l.) complex, *An. funestus* s.s., *An. hancocki*, *An. nili* and *An. ziemanni*. The same species have already been found in other elevated areas such as those of the western [14] and north-western Cameroon [56]. *Anopheles funestus* s.s. appeared as the second dominant vector species after *An. gambiae* s.l., confirming previous reports in these sites [13, 57]. Overall, *An. funestus* s.s. abundance increased from the coastal lowland Tiko to Likoko highland, with relatively high abundances observed in mid-altitude areas (Mutengene and Meanja) probably due to its larval habitat preference. Breeding sites of *An. funestus* s.s. are limited to large permanent waters with aquatic vegetation [58-60], habitats which are quite abundant in Mutengene and Meanja areas [57].

In the Mount Cameroon area, *An. funestus* s.s. vector seemed to supplement malaria transmission more than *An. gambiae* s.l. vectors especially during the dry season months (November – February). This attested the role of *An. funestus* s.s. in bridging the gap of malaria transmission between rainy and dry seasons [58], and compensating the lack of malaria transmission induced by microclimatic conditions in highlands [14].

The genotyping of GSTe2-based DDT resistance maker showed the presence at high frequencies of *GSTe2^R* genotypes and alleles in all *An. funestus* s.s. populations tested throughout the Mount Cameroon altitudinal gradient (7 – 800 m a.s.l.). This presence of GST-based metabolic resistance was further confirmed after sequencing by the detection of a single substitution of nucleotide at position 119 inducing an amino acid change of leucine (CTT) to phenylalanine (TTT) [20]. GST-based metabolic resistance is common in a number of anopheline species including *An. gambiae* s.l. [61, 62] and *An. funestus* s.s. [16, 45, 63], reflecting the heavy use of DDT and pyrethroids for malaria control over several decades [7, 64]. In Cameroon, the intensive use of these insecticide classes to control agriculture pests especially in agroecosystems such as those found in the Mount Cameroon region (especially Tiko, Mutengene and Meanja in this study), may have contaminated mosquitoes breeding sites, thus exerting significant and constant selection pressure on *Anopheles* populations [55].

Similarly, *An. funestus* s.s. populations from the Mount Cameroon exhibited high levels of mutations in the *Rdl* gene which encoded for the GABA-receptor subunit. Dieldrin resistance have already been documented in insect species [65] including the malaria vectors *An. stephensi* [66], *An.*

gambiae s.l. [67] and *An. funestus* [15]. In Cameroon, high frequencies of *Rdl* mutation have also been reported in areas other than the Mount Cameroon region [16, 46, 68, 69] despite the fact that cyclodienes are no longer used for control programs. Moreover, observations made in this study raise concerns on the use of agrochemicals targeting the GABA-receptor in the agricultural environment. Thus, understanding the factors which could possibly explain the persistence of this resistance in *An. funestus* populations are greatly needed in order to get insights on resistance management.

Thirty-nine percent (132/339) of *An. funestus* s.s. tested specimens were found carrying both *GSTe2^R* and *Rdl^R* alleles simultaneously, confirming report of Menze, *et al.* [16] who had identified multiple resistance markers in *An. funestus* s.s. populations from Gounougou, in the Northern part of the country. The extensive use of agrochemicals for crop protection coupled to long lasting insecticidal nets (LLINs) massively distributed since 2011 for public health activities through the National Malaria Control Program of the Ministry of Public Health, might have also greatly influenced the selection of these mutations in vector populations. However, investigations are required on the frequency and distribution of metabolic and target-site resistant alleles as well as on the validation of pyrethroids resistance in *An. funestus* s.s. populations from a thorough fully representative sites across this agricultural domain which is highly favorable for human implantation. These will represent an added value in further understanding the linkage between resistance markers and the use of agrochemical and malaria public health activities.

Our results showed substantial variation in *GSTe2*-based and dieldrin resistance trends within the Mount Cameroon domain. Interestingly, *GSTe2^R* allelic frequencies increased with land elevation whereas *Rdl^R* frequencies decreased with climb in altitude. These observations suggest that altitude could positively favour the establishment of *An. funestus* *GSTe2*-resistant populations from mid-(Mutengene and Meanja) to highland (Likoko) areas, unlike to dieldrin-resistant populations which seemed to be more adapted to Tiko, the lowest elevated site of the Mount Cameroon region as investigated in this study. Land elevation had previously been reported as one important influential predictor of increase in pyrethroid resistance in the *An. gambiae* species complex in West Africa though not in East Africa [70]. This study highlights some uncertainties of the potential influence of altitude on the maintenance of insecticide resistances in malaria vector populations under specific environmental conditions similar to that of the study area. In such areas, field sampling to measure resistance is the only means of informing resistance management decisions alongside an assessment of the historical and contemporary role of pesticide usage and the role of public health insecticide use in the development of insecticide resistance in malaria vectors as previously reported [71].

It has been reported that landscape variations are associated with the risk of presence and insecticide resistance for malaria vectors [72-74]. Topography and land use influence vector densities, level of exposure to insecticide and resistance development in mosquitoes, especially in areas with combined exposure to insecticide from agricultural and /or vector control activities [75]. Distribution of chromosomal inversions such as those found in resistance to insecticides had already been extensively associated with latitudinal but less frequently with altitudinal changes [76-78], thus investigations are needed in order to further elucidate the association between altitudinal changes and the geographical distribution of insecticide resistance.

Patterns of genetic differentiation based on *GSTe2* mutation revealed that Tiko (9 – 70 m a.s.l.) and Mutengene (220 m a.s.l.) populations of *An. funestus* s.s. are genetically differentiated to that of Meanja (305 m a.s.l.) and Likoko (800 m a.s.l.) as they formed a unique cluster compared to others on the neighbour-joining tree of distance. Out of twelve haplotypes identified, *An. funestus* s.s. populations from Tiko and Mutengene appeared to share six haplotypes of which three are exclusively found in these localities. The causes of this clustering could be associated with the similar geographical position of both populations around the Mount Cameroon (southwestern edge) or the presence of the mountain itself which affects the population genetic structure and speed of spread of *GSTe2^R* allele between *An. funestus* s.s. populations. Patterns of *GSTe2* population genetic support the contrast in resistance patterns between *An. funestus* s.s. populations and further suggest the presence of barriers to gene flow between these populations. Similar geographical barriers to the

spread of resistance alleles has been mentioned for other resistance makers such as P450-based metabolic resistance in *An. funestus* [79] or knock-down resistance (*kdr*) mutations in *An. gambiae* [9].

Comparatively, genetic variability patterns within the GABA-receptor based *Rdl* mutation showed that *An. funestus* s.s. population from Tiko at the base of the Mount Cameroon is more genetically differentiated to mid- and highland vector populations as it separately clusters to other localities (Table 2). This strong differentiation observed on *Rdl* mutation in Tiko was confirmed by the high and significant values of K_{ST} statistics of genetic differentiation obtained for *An. funestus* s.s. population from Tiko (Table 4). Once more, it can be hypothesized that the presence of the Mount Cameroon (mountain) influences the contrast in *Rdl* resistance patterns between populations of *An. funestus* s.s. in the study area thus suggesting the presence of barriers of gene flow between *An. funestus* s.s. populations. This is further supported by the reduced genetic diversities parameters (h and Hd), positive value of F_u and F_l F^* index and significant K_{ST} statistics obtained in Meanja midland (eastern edge of the Mount Cameroon) compared to mid- (Mutengene) and highland (Likoko) localities of the Great West (Table 2 and Table 4). Nevertheless, investigations of more vector populations from both sides of the Mount Cameroon are needed to validate such hypothesis.

A strong selection process was observed on both 119F and 296S resistance allele carriers. This is seen by the reduced genetic diversity parameters (h , Hd , π and k) with limited number of mutational steps (polymorphic sites) between haplotypes in resistance allele carriers compared to susceptible allele carriers which maintained high diversity parameters (Table 1 and Table 2). Analysis of ML tree based on allelic profiles (Figure 5 and Figure 6) further illustrated the reduced diversity of resistance allele carriers in both cases. Furthermore, the most predominant haplotype is found in 85% (GSTe2-based mutation) to 100% (*Rdl* mutation) resistance allele carriers which is indicative of ongoing selection on 119F and 296S alleles contrasting with L119 and A296 susceptible alleles which maintained high number of singletons. Similar selection patterns have been observed in P450 [79] and GSTe2 [20] genes in *An. funestus* populations from other African regions. The selection process could be due to intensive use of insecticides through routine integrated control carried out by the National Malaria Control Program in the Mount Cameroon region, particularly in the localities surveyed. In addition, positive selection could also be associated with adaptation of mosquito larval stages to agricultural pesticides and other adverse conditions, such as temperature and landscape [80, 81].

Based on GSTe2 gene, comparing populations of *An. funestus* s.s. from the Mount Cameroon region with other *An. funestus* African populations revealed that Cameroonian (including Mount Cameroon and North Cameroon) populations of *An. funestus* seemingly present similar patterns of genetic diversity. Despite the presence of *GSTEeR* allele recorded in the Mount Cameroon (at high frequencies), high values of genetic diversity parameters were still obtained compared to that of Benin where a marked reduced diversity was noted (Supplementary Table S3). However, the high diversity observed in the Mount Cameroon besides high frequencies of *GSTE2R* allele, reflects a situation of moderate selection of the 119F allele which could progressively change in the future to become as Benin where the *GSTE2R* allele is nearly driven to fixation due to a greater selection of resistance [20]. The high diversity obtained in Cameroon is further reinforced by the high level of diversity in L119 susceptible allele carriers. Thereby, if selective pressure continues to act within *An. funestus* populations, resistance profiles are likely to changes as previously reported for P450-based metabolic resistance where CYP6P9a-R allele frequency increased from 7.5% in 2002 to 100% in 2017 due to scale-up of insecticidal bed-nets [82].

5. Conclusions

Broadly, this study highlights for the first time the presence of GSTe2-based metabolic resistance and the GABA-receptor target-site mutation associated to dieldrin resistance in the malaria vectors *An. funestus* s.s. across the Mount Cameroon domain. Both *GSTE2R* and *RdlR* alleles were found with high frequencies in almost all the localities surveyed; however, the speed of spread of these two molecular mechanisms appears to be influenced by the presence of a major mountainous barrier, the Mount Cameroon which contrasts the resistance and diversity patterns of

these two genes between populations of *An. funestus* s.s. in the study area. Furthermore, we provide evidence of positive selection occurring on *GSTE2^R* and *Rdl^R* throughout the Mount Cameroon region, which if not adequately monitor could drive to fixation in response to a greater selection of resistance in the future. This emphasizes the need of molecular studies of multiple collections sites throughout such mountainous landscapes to fully elucidate the role of environmental changes on the acquisition of insecticide resistance in *Anopheles* vector populations and to mitigate against further spread of resistance through the development of new vector management strategies.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, **Figure S1.** Mosquito genera identified in each surveyed locality; **Figure S2.** Female's anophelines species composition across the study area; **Figure S3.** Parity and Plasmodium spp. infection rates within *An. gambiae* species complex and of *An. funestus* s.s.; **Figure S4.** Nucleotide sequences of the 53 *GSTE2* haplotypes identified across Africa; **Figure S5.** Africa-wide diversity pattern of a 729 bp fragment of the *GSTE2* gene. **Table S1:** Detailed period (in years) of collections; **Table S2:** List of primers used in this study; **Table S3:** Genetic diversity parameters of *GSTE2* gene across Africa.

Author Contributions: Conceptualization, S.W., FN and C.S.W.; methodology, N.A.A; validation, S.W., F.N., JMR and C.S.W.; formal analysis, N.A.A. and C.S.W.; writing—original draft preparation, N.A.A., J.M.R. and C.S.W. All authors have read and agreed to the published version of the manuscript.

Acknowledgements: This work and the cost to publish in open access was supported by a Wellcome Trust Research Career Development Fellowship (083515/Z/07/Z) and a Wellcome Trust Senior Research Fellowship in Biomedical Sciences (101893/Z/13/Z) to CSW.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. WHO *Test procedure for insecticides resistance monitoring in malaria vector mosquitoes*. 2nd ed.; World Health Organisation: 2016; p 6-8.
2. Zaim, M.; Aitio, A.; Nakashima, N. Safety of pyrethroid-treated mosquito nets. *Med Vet Entomol* **2000**, *14* (1), 1-5, doi:10.1046/j.1365-2915.2000.00211.x.
3. WHO *Long-lasting insecticidal nets for malaria prevention - a manual for malaria programme managers*. Trial ed.; World Health Organisation: 2007; p 5-9.
4. WHO *WHO recommended long-lasting insecticidal mosquito nets*; World Health Organisation: Geneva, 2009.
5. Pluess, B.; Tanser, F.C.; Lengeler, C.; Sharp, B.L. Indoor residual spraying for preventing malaria. *Cochrane Database Sys Rev* **2010**, *2010* (4), doi:10.1002/14651858.CD006657.pub2.
6. Bhatt, S.; Weiss, D.; Cameron, E.; Bisanzio, D.; Mappin, B.; Dalrymple, U.; Battle, K.; Moyes, C.; Henry, A.; Eckhoff, P.; Wenger, E.; Briët, O.; Penny, M.; Smith, T.; Bennett, A.; Yukich, J.; Eisele, T.; Griffin, J.; Fergus, C.; Lynch, M.; Lindgren, F.; Cohen, J.; Murray, C.; Smith, D.; Hay, S.; Cibulskis, R.; Gething, P. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature* **2015**, *526* (7572), 207-211, doi:10.1038/nature15535.
7. Hemingway, J.; Ranson, H. Insecticide resistance in insect vectors of human disease. *Annu Rev Entomol* **2000**, *45*, 371-391, doi:10.1146/annurev.ento.45.1.371.
8. Hemingway, J.; Field, L.; Vontas, J. An overview of insecticide resistance. *Science* **2002**, *298*, 96-97, doi:10.1126/science.1078052.
9. Ranson, H.; N'Guessan, R.; Lines, J.; Moiroux, N.; Nkuni, Z.; Corbel, V. Pyrethroid resistance in African anopheline mosquitoes: what are the implications for malaria control? *Trends Parasitol* **2010**, *xx*, 1-8, doi:10.1016/j.pt.2010.08.004.
10. WHO *World Malaria Report 2019*; World Health Organisation: Geneva, 2019.
11. Sinka, M.; Bangs, M.; Manguin, S.; Coetzee, M.; Mbogo, C.; Hemingway, J.; Patil, A.; Temperley, W.; Gething, P.; Kabaria, C.; Okara, R.; Van-Boeckel, T.; Godfray, H.; Harbach, R.; Hay, S. The dominant *Anopheles* vectors of human malaria in Africa, Europe and the Middle East: occurrence data, distribution maps and bionomic précis. *Parasit Vectors* **2010**, *3* (1), 117, doi:10.1186/1756-3305-3-117.
12. Coetzee, M.; Hunt, R.; Wilkerson, R.; Della Torre, A.; Coulibaly, M.; Besansky, N. *Anopheles coluzzii* and *Anopheles amharicus*, new members of the *Anopheles gambiae* complex. *Zootaxa* **2013**, *3619*, 246-274, <http://dx.doi.org/10.11646/zootaxa.3619.3.2>.
13. Wanji, S.; Tanke, T.; Atanga, S.N.; Ajonina, C.; Tendongfor, N.; Fontenille, D. *Anopheles* species of the mount Cameroon region: biting habits, feeding behaviour and entomological inoculation rates. *Trop Med Int Health* **2003**, *8* (7), 643-649.

14. Tchuinkam, T.; Simard, F.; Lélé-Defo, E.; Téné-Fossog, B.; Tateng-Ngouateu, A.; Antonio-Nkondjio, C.; Mpoame, M.; Toto, J.; Njiné, T.; Fontenille, D.; Awono-Ambéné, P. Bionomics of Anopheline species and malaria transmission dynamics along an altitudinal transect in Western Cameroon. *BMC Infect Dis* **2010**, *10*, 119, doi:10.1186/1471-2334-10-119.
15. Wondji, C.; Dabire, R.; Tukur, Z.; Irving, H.; Djouaka, R.; Morgan, J. Identification and distribution of a GABA receptor mutation conferring dieldrin resistance in the malaria vector *Anopheles funestus* in Africa. *Insect Biochem Mol Biol* **2011**, *41*, 484-491, doi:10.1016/j.ibmb.2011.03.012.
16. Menze, B.; Riveron, J.; Ibrahim, S.; Irving, H.; Antonio-Nkondjio, C.; Awono-Ambene, P.; Wondji, C. Multiple insecticide resistance in the malaria vector *Anopheles funestus* from northern Cameroon is mediated by metabolic resistance alongside potential target site insensitivity. *PLoS ONE* **2016**, *11* (10), e0163261, doi:10.1371/journal.pone.0163261.
17. Irving, H.; Wondji, C. Investigating knockdown resistance (*kdr*) mechanism against pyrethroids/DDT in the malaria vector *Anopheles funestus* across Africa. *BMC Genetics* **2017**, *18*, 76, doi:10.1186/s12863-017-0539-x.
18. Ndo, C.; Kopya, E.; Donbou, M.A.; Njiokou, F.; Awono-Ambene, P.; Wondji, C. Elevated *Plasmodium* infection rates and high pyrethroid resistance in major malaria vectors in a forested area of Cameroon highlight challenges of malaria control. *Parasit Vectors* **2018**, *11* (1), 157, doi:10.1186/s13071-018-2759-y.
19. Ndo, C.; Kopya, E.; Irving, H.; Wondji, C. Exploring the impact of glutathione S-transferase (GST)-based metabolic resistance to insecticide on vector competence of *Anopheles funestus* for *Plasmodium falciparum* [version 2; peer review: 2 approved, 1 approved with reservations]. *Wellcome Open Research* **2019**, *4* (52), doi:10.12688/wellcomeopenres.15061.2.
20. Riveron, J.; Yunta, C.; Ibrahim, S.; Djouaka, R.; Irving, H.; Menze, B.; Ismail, H.; Hemingway, J.; Ranson, H.; Albert, A.; Wondji, C. A single mutation in the GSTe2 gene allows tracking of metabolically based insecticide resistance in a major malaria vector. *Genome Biol* **2014**, *12*, R27.
21. Lui, N. Insecticides resistance in mosquitoes: impact, mechanisms, and research directions. *Annu Rev Entomol* **2015**, *60*, 537-559, doi:10.1146/annurev-ento-010814-020828.
22. Riveron, J.; Irving, H.; Ndula, M.; Barnes, K.; Ibrahim, S.; Paine, M.; Wondji, C. Directionally selected cytochrome P450 alleles are driving the spread of pyrethroid resistance in the major malaria vector *Anopheles funestus*. *Proc Natl Acad Sci U S A* **2013**, *110* (1), 252-257, doi:10.1073/pnas.1216705110.
23. Riveron, J.; Ibrahim, S.; Chanda, E.; Mzilahowa, T.; Cuamba, N.; Irving, H.; Barnes, K.; Ndula, M.; Wondji, C. The highly polymorphic CYP6M7 cytochrome P450 gene partners with the directionally selected CYP6P9a and CYP6P9b genes to expand the pyrethroid resistance front in the malaria vector *Anopheles funestus* in Africa. *BMC Genomics* **2014**, *15* (1), 817, doi:10.1186/1471-2164-15-817.
24. Weedall, G.; Mugenzi, L.; Menze, B.; Tchouakui, M.; Ibrahim, S.; Amvongo-Adjia, N.; Irving, H.; Wondji, M.; Tchoupo, M.; Djouaka, R.; Riveron, J.; Wondji, C. A cytochrome P450 allele confers pyrethroid resistance on a major African malaria vector, reducing insecticide-treated bednet efficacy. *Science Translational Medicine* **2019**, *11*, eaat7386.
25. Mugenzi, L.M.J.; Menze, B.D.; Tchouakui, M.; Wondji, M.J.; Irving, H.; Tchoupo, M.; Hearn, J.; Weedall, G.D.; Riveron, J.M.; Wondji, C.S. Cis-regulatory CYP6P9b P450 variants associated with loss of insecticide-treated bed net efficacy against *Anopheles funestus*. *Nat Commun* **2019**, *10* (1), 4652, doi:10.1038/s41467-019-12686-5.
26. Barnes, K.G.; Irving, H.; Chiumia, M.; Mzilahowa, T.; Coleman, M.; Hemingway, J.; Wondji, C.S. Restriction to gene flow is associated with changes in the molecular basis of pyrethroid resistance in the malaria vector *Anopheles funestus*. *Proc Natl Acad Sci U S A* **2017**, *114* (2), 286-291, doi:10.1073/pnas.1615458114.
27. Weedall, G.D.; Riveron, J.M.; Hearn, J.; Irving, H.; Kamdem, C.; Fouet, C.; White, B.J.; Wondji, C.S. An Africa-wide genomic evolution of insecticide resistance in the malaria vector *Anopheles funestus* involves selective sweeps, copy number variations, gene conversion and transposons. *PLoS genetics* **2020**, *16* (6), e1008822-e1008822, doi:10.1371/journal.pgen.1008822.
28. Boussougou-Sambe, S.T.; Eyiap, W.E.; Tasse, G.C.T.; Mandeng, S.E.; Mbakop, L.R.; Enyong, P.; Etang, J.; Fokam, E.B.; Awono-Ambene, P.H. Insecticide susceptibility status of *Anopheles gambiae* (s.l.) in South-West Cameroon four years after long-lasting insecticidal net mass distribution. *Parasit vectors* **2018**, *11* (1), 391, doi:10.1186/s13071-018-2979-1.
29. Awono, A.; Akombi, A.; Owona, H.; Barreau, E. REDD+ around Mount Cameroon, southwest region of Cameroon. In *REDD+ on the ground: A case book of subnational initiatives across the globe*, Sills, E.; Stibniati S Atmadja, S.; de Sassi, C.; Duchelle, A.; Kweka, D.; Resosudarmo, I.; Sunderlin, W., Eds. CIFOR: Indonesia, 2014; pp 188-202.
30. Wanji, S.; Enyong, P.; Tendongfor, N.; Ndamukong, J.; Mafo, F.; Kengne-Ouaf, A. Biodiversity of *Anopheles* mosquito species across the Mount Cameroon region. **2006** (Unpublished work).

31. Wanji, S.; Kengne-Ouafo, A.J.; Eyong, E.E.J.; Kimbi, H.K.; Tendongfor, N.; Ndamukong-Nyanga, J.L.; Nana-Djeunga, H.C.; Bourguinat, C.; Sofeu-Feugaing, D.D.; Charvet, C.L. Genetic diversity of *Plasmodium falciparum* merozoite surface protein-1 block 2 in sites of contrasting altitudes and malaria endemicities in the Mount Cameroon region. *The American journal of tropical medicine and hygiene* **2012**, *86* (5), 764-774, doi:10.4269/ajtmh.2012.11-0433.

32. Anchang-Kimbi, J.K.; Nkweti, V.N.; Ntonifor, H.N.; Apinjoh, T.O.; Tata, R.B.; Chi, H.F.; Achidi, E.A. *Plasmodium falciparum* parasitaemia and malaria among pregnant women at first clinic visit in the mount Cameroon Area. *BMC Infect Dis* **2015**, *15*, 439-439, doi:10.1186/s12879-015-1211-6.

33. Suchel, J.-B. Les climats du Cameroun. Thèse de doctorat d'Etat, Université de Saint-Etienne, France, 1988.

34. Bigoga, J.; Manga, L.; Titanji, V.; Coetzee, M.; Leke, R. Malaria vectors and transmission dynamics in coastal south-western Cameroon. *Malar J* **2007**, *6* (5), doi:10.1186/1475-2875-6-5.

35. Amvongo-Adjia, N.; Wirsiy, E.L.; Riveron, J.M.; Chounna Ndongmo, W.P.; Enyong, P.A.; Njiokou, F.; Wondji, C.S.; Wanji, S. Bionomics and vectorial role of anophelines in wetlands along the volcanic chain of Cameroon. *Parasit Vectors* **2018**, *11* (1), 471-471, doi:10.1186/s13071-018-3041-z.

36. Gillies, M.; de Meillon, B. *The Anophelinae of Africa South of the Sahara (Ethiopian Zoogeographical Region)* South African Institute for Medical Research: Johannesburg, 1968; Vol. 2nd.

37. Detinova, T. Age grouping methods in Diptera of medical importance, with special reference to some vectors of malaria. *World Health Organization Monographs series*, **1962**, 47.

38. Cornel, A.; Collins, F. PCR of the ribosomal DNA intergenic spacer regions as a method for identifying mosquitoes in the *Anopheles gambiae* complex. *Methods Mol Biol* **1996**, *50*, 321-332, doi:10.1385/0-89603-323-6:321.

39. Scott, J.A.; Brogdon, W.G.; Collins, F.H. Identification of single specimens of the *Anopheles gambiae* complex by the polymerase chain reaction. *The American journal of tropical medicine and hygiene* **1993**, *49* (4), 520-529, doi:10.4269/ajtmh.1993.49.520.

40. Fanello, C.; Santolamazza, F.; della Torre, A. Simultaneous identification of species and molecular forms of *Anopheles gambiae* complex by PCR-RFLP. *Med Vet Entomol* **2002**, *16*, 461-464, doi:10.1046/j.1365-2915.2002.00393.x.

41. Koekemoer, L.; Kamau, L.; Hunt, R.; Coetzee, M. A cocktail polymerase chain reaction assay to identify members of the *Anopheles funestus* (Diptera: Culicidae) group. *The American journal of tropical medicine and hygiene* **2002**, *66*, 804-811, doi:10.4269/ajtmh.2002.66.804.

42. Bass, C.; Nikou, D.; Blagborough, M.; Vontas, J.; Sinden, E.; Williamson, S.; Field, M. PCR-based detection of *Plasmodium* in *Anopheles* mosquitoes: a comparison of a new high-throughput assay with existing methods. *Malar J* **2008**, *7*, 177, doi:10.1186/1475-2875-7-177.

43. Padley, D.; Moody, A.H.; Chiodini, P.L.; Saldanha, J. Use of a rapid, single-round, multiplex PCR to detect malarial parasites and identify the species present. *Annals of tropical medicine and parasitology* **2003**, *97*, 131-137, doi:10.1179/000349803125002977.

44. Snounou, G.; Pinheiro, L.; Goncalves, A.; Fonseca, L.; Dias, F.; Brown, K.N.; Dorosario, V.E. The importance of sensitive detection of malaria parasites in the human and insect hosts in epidemiological studies, as shown by the analysis of field samples from Guinea Bissau. *Trans R Soc Trop Med Hyg* **1993**, *87*, 646-653, doi:10.1016/0035-9203(93)90274-t.

45. Tchouakui, M.; Chiang, M.-C.; Ndo, C.; Kuicheu, C.; Amvongo-Adjia, N.; Wondji, M.; Tchoupo, M.; Kusimo, M.; Riveron, J.; Wondji, C. A marker of glutathione S-transferase-mediated resistance to insecticides is associated with higher *Plasmodium* infection in the African malaria vector *Anopheles funestus*. *Sci Rep* **2019**, *9*, 5772, doi:10.1038/s41598-019-42015-1.

46. Riveron, J.; Chiumia, M.; Menze, B.; Barnes, K.; Irving, H.; Ibrahim, S.; Weedall, G.; Mzilahowa, T.; Wondji, C. Rise of multiple insecticide resistance in *Anopheles funestus* in Malawi: a major concern for malaria vector control. *Malar J* **2015**, *14*, 344, doi:10.1186/s12936-015-0877-y.

47. Hall, T. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **1999**, *41*, 95-98.

48. Thompson, J.; Higgins, D.; Gibson, T. CLUSTALW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res* **1994**, *22* (22), 4673-4680, doi:10.1093/nar/22.22.4673.

49. Librado, P.; Rozas, J. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* **2009**, *25* (11), 1451-1452, doi:10.1093/bioinformatics/btp187.

50. Clement, M.; Posada, D.; Crandall, K. TCS: a computer program to estimate gene genealogies. *Mol Ecol* **2000**, *9* (10), 1657-1660, doi:10.1046/j.1365-294x.2000.01020.x.

51. Santos, A.; Cabezas, M.; Tavares, A.; Xavier, R.; Branco, M. tcsBU: a tool to extend TCS network layout and visualization. *Bioinformatics* **2015**, *32* (4), 627-628, doi:10.1093/bioinformatics/btv636.

52. Hudson, R.R.; Slatkin, M.; Maddison, W.P. Estimation of levels of gene flow from DNA sequence data. *Genetics* **1992**, *132* (2), 583-589, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1205159/>.

53. Kumar, S.; Stecher, G.; Li, M.; Knyaz, C.; Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **2018**, *35* (6), 1547-1549, doi:10.1093/molbev/msy096.

54. Nei, M. Analysis of gene diversity in subdivided populations. *Proc Natl Acad Sci U S A* **1973**, *70* (12), 3321-3323, doi:10.1073/pnas.70.12.3321.

55. Antonio-Nkondjio, C.; Ndo, C.; Njiokou, F.; Bigoga, J.; Awono-Ambene, H.; Etang, J.; Ekobo, A.S.; Wondji, C. Review of malaria situation in Cameroon: technical viewpoint on challenges and prospects for disease elimination. *Parasit Vectors* **2019**, *12* (1), 501, doi:10.1186/s13071-019-3753-8.

56. Tabue, R.; Nem, T.; Atangana, J.; Bigoga, J.; Patchoké, S.; Tchouine, F.; Fodjo, B.; Leke, R.; Fondjo, E. *Anopheles ziemanni* a locally important malaria vector in Ndop health district, north west region of Cameroon. *Parasit Vectors* **2014**, *7*, 262, doi:10.1186/1756-3305-7-262.

57. Wanji, S.; Mafo, F.; Tendongfor, N.; Tanga, M.; Tchuente, F.; Bilong Bilong, C.; Njiné, T. Spatial distribution, environmental and physicochemical characterization of *Anopheles* breeding sites in the Mount Cameroon region. *J Vector Borne Dis* **2009**, *46*, 75-80, <https://pubmed.ncbi.nlm.nih.gov/19326712/>.

58. Minakawa, N.; Sonye, G.; Mogi, M.; Githeko, A.; Yan, G. The effects of climatic factors on the distribution and abundance of malaria vectors in Kenya. *J Med Entomol* **2002**, *39*, doi:10.1603/0022-2585-39.6.833.

59. Minakawa, N.; Munga, S.; Atieli, F.; Mushinzimana, E.; Zhou, G.; Githeko, A.; Yan, G. Spatial distribution of anopheline larval habitats in Western Kenyan highlands: effects of land cover types and topography. *The American journal of tropical medicine and hygiene* **2005**, *73* (1), 157-165, <https://pubmed.ncbi.nlm.nih.gov/16014851/>.

60. Kweka, E.; Munga, S.; Himeidan, Y.; Githeko, A.; Yan, G. Assessment of mosquito larval productivity among different land use types for targeted malaria vector control in the western Kenya highlands. *Parasit Vectors* **2015**, *8*, 356, doi:10.1186/s13071-015-0968-1.

61. Prapanthadara, L.; Ketterman, A. Qualitative and quantitative changes in glutathione S-transferases in the mosquito *Anopheles gambiae* confer DDT-resistance. *Biochem Soc Trans* **1993**, *21* (Pt 3), 304S, doi:10.1042/bst021304s.

62. Ranson, H.; Rossiter, L.; Ortelli, F.; Jensen, B.; Wang, X.; Roth, C.; Collins, F.; Hemingway, J. Identification of a novel class of insect glutathione S-transferases involved in resistance to DDT in the malaria vector *Anopheles gambiae*. *Biochem J* **2001**, *359* (Part 2), 295-304, doi:10.1042/0264-6021:3590295.

63. Tchakounte, A.; Tchouakui, M.; Mu-Chun, C.; Tchapga, W.; Kopia, E.; Soh, P.; Njiokou, F.; Riveron, J.; Wondji, C. Exposure to the insecticide-treated bednet PermaNet 2.0 reduces the longevity of the wild African malaria vector *Anopheles funestus* but GSTe2-resistant mosquitoes live longer. *PloS ONE* **2019**, *14* (3), e0213949, doi:10.1371/journal.pone.0213949.

64. Hemingway, J.; Hawkes, N.; Carroll, L.; Ranson, H. The molecular basis of insecticide resistance in mosquitoes. *Insect Biochem Mol Biol* **2004**, *34* (7), 653-665, doi:10.1016/j.ibmb.2004.03.018.

65. Ffrench-Constant, R. The molecular and population genetics of cyclodiene insecticide resistance. *Insect Biochem Mol Biol* **1994**, *24* (4), 335-345, doi:10.1016/0965-1748(94)90026-4.

66. Andreasen, M.; Ffrench-Constant, R. In situ hybridization to the Rdl locus on polytene chromosome 3L of *Anopheles stephensi*. *Med Vet Entomol* **2002**, *16* (4), 452-455, doi:10.1046/j.1365-2915.2002.00390.x.

67. Du, W.; Awolola, T.S.; Howell, P.; Koekemoer, L.L.; Brooke, B.D.; Benedict, M.Q.; Coetzee, M.; Zheng, L. Independent mutations in the Rdl locus confer dieldrin resistance to *Anopheles gambiae* and *An. arabiensis*. *Insect Mol Biol* **2005**, *14* (2), 179-183, doi:10.1111/j.1365-2583.2005.00544.x.

68. Etang, J.; Manga, L.; Chandre, F.; Guillet, P.; Fondjo, E.; Mimpfoundi, R.; Toto, J.-c.; Fontenille, D. Insecticide Susceptibility Status of *Anopheles gambiae* s.l. (Diptera : Culicidae) in the Republic of Cameroon. *J Med Entomol* **2003**, *40* (4), 491-497, doi:10.1603/0022-2585-40.4.491.

69. Nwane, P.; Etang, J.; Chouaïbou, M.; Toto, J.C.; Koffi, A.; Mimpfoundi, R.; Simard, F. Multiple insecticide resistance mechanisms in *Anopheles gambiae* s.l. populations from Cameroon, Central Africa. *Parasit Vectors* **2013**, *6*, 41-41, doi:10.1186/1756-3305-6-41.

70. Hancock, P.; Hendriks, C.; Tangena, J.-A.; Gibson, H.; Hemingway, J.; Coleman, M.; Gething, P.; Cameron, E.; Bhatt, S.; Moyes, C. Mapping trends in insecticide resistance phenotypes in African malaria vectors. *PLOS Biology* **2020**, *18* (6), e3000633, doi:10.1371/journal.pbio.3000633.

71. Matiya, D.; Philbert, A.; Kidima, W.; Matowo, J. Dynamics and monitoring of insecticide resistance in malaria vectors across mainland Tanzania from 1997 to 2017: a systematic review. *Malaria Journal* **2019**, *18* (1), 102, doi:10.1186/s12936-019-2738-6.

72. Chang, X.; Zhong, D.; Lo, E.; Fang, Q.; Bonizzoni, M.; Wang, X.; Lee, M.-C.; Zhou, G.; Zhu, G.; Qin, Q.; Chen, X.; Cui, L.; Yan, G. Landscape genetic structure and evolutionary genetics of insecticide resistance gene mutations in *Anopheles sinensis*. *Parasit Vectors* **2016**, *9* (1), 228, doi:10.1186/s13071-016-1513-6.

73. Souris, M.; Marcombe, S.; Laforet, J.; Brey, P.T.; Corbel, V.; Overgaard, H.J. Modeling spatial variation in risk of presence and insecticide resistance for malaria vectors in Laos. *PLoS ONE* **2017**, *12* (5), e0177274, doi:10.1371/journal.pone.0177274.

74. Mandeng, S.E.; Awono-Ambene, H.P.; Bigoga, J.D.; Ekoko, W.E.; Binyang, J.; Piameu, M.; Mbakop, L.R.; Fesuh, B.N.; Mvondo, N.; Tabue, R.; Nwane, P.; Mimpfoundi, R.; Toto, J.C.; Kleinschmidt, I.; Knox, T.B.; Mnzava, A.P.; Donnelly, M.J.; Fondjo, E.; Etang, J. Spatial and temporal development of deltamethrin resistance in malaria vectors of the *Anopheles gambiae* complex from North Cameroon. *PLoS ONE* **2019**, *14* (2), e0212024, doi:10.1371/journal.pone.0212024.

75. Wanjala, C.L.; Kweka, E.J. Malaria vectors insecticides resistance in different agroecosystems in western Kenya. *Front Public Health* **2018**, *6*, 55-55, doi:10.3389/fpubh.2018.00055.

76. Touré, Y.T.; Petrarca, V.; Traoré, S.F.; Coulibaly, A.; Maïga, H.M.; Sankaré, O.; Sow, M.; Di Deco, M.A.; Coluzzi, M. Ecological genetic studies in the chromosomal form Mopti of *Anopheles gambiae* s.str. in Mali, West Africa. *Genetica* **1994**, *94* (2), 213-223, doi:10.1007/BF01443435.

77. Petrarca, V.; Nugud, A.D.; Ahmed, M.A.E.; Haridi, A.M.; Di Deco, M.A.; Coluzzi, M. Cytogenetics of the *Anopheles gambiae* complex in Sudan, with special reference to *An. arabiensis*: relationships with East and West African populations. *Med Vet Entomol* **2000**, *14* (2), 149-164, doi:10.1046/j.1365-2915.2000.00231.x.

78. Ayala, D.; Ullastres, A.; González, J. Adaptation through chromosomal inversions in *Anopheles*. *Front Genet* **2014**, *5* (129), doi:10.3389/fgene.2014.00129.

79. Barnes, K.; Weedall, G.; Ndula, M.; Irving, H.; Mzihalowa, T.; Hemingway, J.; Wondji, C. Genomic footprints of selective sweeps from metabolic resistance to pyrethroids in African malaria vectors are driven by scale up of insecticide-based vector control. *PLOS Genetics* **2017**, *13* (2), e1006539, doi:10.1371/journal.pgen.1006539.

80. Simard, F.; Lehmann, T.; Lemasson, Jean-Jacques; Diatta, M.; Fontenille, D. Persistence of *Anopheles arabiensis* during the severe dry season conditions in Senegal: an indirect approach using microsatellite loci. *Insect Mol Biol* **2000**, *9* (5), 467-479, doi:10.1046/j.1365-2583.2000.00210.x.

81. Yang, M.; Ma, Y.; Wu, J. Mitochondrial genetic differentiation across populations of the malaria vector *Anopheles lesteri* from China (Diptera: Culicidae). *Malar J* **2011**, *10* (1), 216, doi:10.1186/1475-2875-10-216.

82. Riveron, J.; Huijben, S.; Tchapga, W.; Tchouakui, M.; Wondji, M.; Tchoupo, M.; Irving, H.; Cuamba, N.; Maquina, M.; Paaijmans, K.; Wondji, C. Escalation of pyrethroid resistance in the malaria vector *Anopheles funestus* induces a loss of efficacy of piperonyl butoxide-based insecticide-treated nets in Mozambique. *The Journal of infectious diseases* **2019**, *220* (3), 467-475, doi:10.1093/infdis/jiz139.