

The beneficial potential of genetically modified stem cells in the treatment of stroke: a review

Mohammad Saied Salehi¹, Anahid Safari^{2*}, Sareh Pandamooz², Benjamin Jurek³, Etrat Hooshmandi¹, Maryam Owjifard¹, Mahnaz Bayat¹, Seyedeh Shaghayegh Zafarmand¹, Jaleel A Miyan⁴, Afshin Borhani-Haghighi¹

1. Clinical Neurology Research Center, Shiraz University of Medical Sciences, Shiraz Iran
2. Stem cell Technology Research Center, Shiraz University of Medical Sciences, Shiraz Iran
3. Institute of Molecular and Cellular Anatomy, University of Regensburg, Regensburg, Germany
4. Faculty of Biology, Medicine & Health, Division of Neuroscience & Experimental Psychology, The University of Manchester, Manchester, UK

Correspondence:

Anahid Safari, Stem cell Technology Research Center, Shiraz University of Medical Sciences, Shiraz Iran. Tel/Fax: +98-7136281547, Email: a_safari@sums.ac.ir

Abstract

The last two decades have witnessed a surge in investigations proposing stem cells as a promising strategy to treat stroke. Since growth factor release is considered as one of the most important aspects of cell-based therapy, stem cells over-expressing growth factors are hypothesized to yield higher levels of therapeutic efficiency. In pre-clinical studies of the last 15 years that were investigating the efficiency of stem cell therapy for stroke, a variety of stem cell types were genetically modified to over-express various factors. In this review we summarize the current knowledge on the therapeutic efficiency of stem cell-derived growth factors, encompassing techniques employed and time points to evaluate. In addition, we discuss several types of stem cells, including the recently developed model of epidermal neural crest stem cells, and genetically modified stem cells over-expressing specific factors, which could elevate the restorative potential of naive stem cells. The restorative potential is based on enhanced survival/differentiation potential of transplanted cells, apoptosis inhibition, infarct volume reduction, neovascularization or functional improvement. Since the majority of studies have focused on the short-term curative effects of genetically engineered stem cells, we emphasize the need to address their long-term impact.

Key words: Cell therapy; MCAO; Stroke; Genetic engineering, Growth factors, Stem cells

1. Introduction

Stroke is considered as one of the main causes of disability and death all around the world [1, 2]. Reperfusion modalities, such as mechanical thrombectomy and thrombolysis, have been applied for treating ischemic stroke as the most common type of stroke. However, some issues reduce their clinical application, such as development of hemorrhagic transformation, a narrow therapeutic window, and restricted availability [3]. Therefore, seeking new alternatives with fewer limitations and higher efficacy, especially for therapy resistant patients is of paramount necessity.

Reduced oxygen levels in the infarct region following stroke eventually leads to activation of a wide variety of growth factor-related signaling cascades, which exert neuroprotective, anti-inflammatory and anti-apoptotic effects [4, 5]. Nevertheless, endogenous growth factors are insufficient to induce cell survival, neurogenesis and angiogenesis. Hence, it has been suggested that the increment of growth factors by protein or gene therapy could be a potential strategy for stroke. Although, the restorative effects of various growth factors have been investigated for stroke treatment [4], their short half-life has limited their direct application.

The last two decades have witnessed a surge in investigations proposing stem cells as a promising strategy to treat stroke (Figure 1). Mesenchymal stem cells [6-12] from different sources including bone marrow [13, 14] adipose tissue [15] and umbilical cord [16] as well as neural stem cells [17-23] are the most commonly used types of stem cells that have been employed for this purpose. Furthermore, their effectiveness and mechanisms of action have been thoroughly reviewed during the last few years. Moreover, other types of stem cells, such as bone marrow mononuclear cells [24, 25], induced pluripotent stem cells [26-29], dental pulp stem cells [30, 31], menstrual blood-derived stem cells [32, 33] and epidermal neural crest stem cells [34] have been evaluated.




Figure 1: Number of published manuscripts per year listed in PubMed from 1999 to 2019 based on “Stem cell” and “Stroke” MeSH terms.

Releasing growth factors by stem cells is considered as one of the most important therapeutic aspects of cell-based therapy [35]. Therefore, the hypothesis has been raised that stem cells over-expressing growth factors have higher levels of restorative efficiency. During the last 15 years, different types of stem cells were genetically engineered to over-express various kinds of genes in pre-clinical investigations in the context of stroke (Figure 2), which are discussed in the current review. Furthermore, since obtained results are highly time and method dependent, here we discussed evaluated experiments in details with a focus on the employed techniques and time points.




Figure 2: Genetically modified stem cells over-expressing specific factors could elevate the restorative potential of naive stem cells in the treatment of stroke.

2. Over-expression of neurotrophins

2.1. Brain-derived neurotrophic factor (BDNF)




BDNF, one of the most well-studied members of the neurotrophin family, is considered as the neurotrophin with the highest expression level in the adult central nervous system. BDNF plays fundamental roles in synaptic remodeling and brain plasticity, neurite extension, neuronal development and cell survival. Following stroke, it has been reported that BDNF expression in the infarct region is permanently reduced [36], and single allele deficiency for BDNF (+/-) resulted in larger infarct area [37]. Furthermore, it has been shown that BDNF treatment led to a reduction in lesion size [38, 39] and improved functional motor performance [39-41]; however, due to the short half-life of BDNF, this therapeutic approach does not provide long-term effects. Therefore, in


order to provide long-lasting effects of BDNF post-ischemia, one possible strategy is to transplant stem cells that stably over-express this protein.

As the first published experiment using genetically modified stem cells to treat stroke, Kurozumi et al. [42] evaluated therapeutic effects of human bone marrow mesenchymal stem cells (BM-MSCs), genetically modified to over-express BDNF (MSCs^{BDNF}). The stem cells were transplanted into the striatum of male rats, one day after 90min middle cerebral artery occlusion (MCAO). Limb placement and treadmill tests showed functional improvements at days 8 and 15 after stroke in rats that received BM-MSCs^{BDNF} compared to the non-transplanted (control) group. Also, T2-weighted imaging (T2WI, see Box 1) on days 7 and 14 post-MCAO revealed a prominent infarct volume reduction in the MSCs^{BDNF} group, compared to controls. Wild-type MSCs neither improved functional recovery, nor reduced infarct size at the evaluated time points. MSC^{BDNF} also decreased the number of terminal deoxynucleotidyltransferase dUTP nick-end labeling (TUNEL)-positive apoptotic cells in the ischemic boundary zone one week after stroke compared to control. Again, wild-type MSCs did not show any therapeutic effect. At this time point, some transplanted stem cells expressed neuronal (NeuN) or glial (GFAP) markers and MSCs^{BDNF} showed features similar to MSCs. Hence, curative potential of MSCs^{BDNF} might be independent of their differentiation potential.

Moreover, Nomura, Honmou (43) evaluated the efficacy of systemic delivery of human MSCs transfected with the BDNF gene in a male rat model of permanent MCAO (pMCAO). In their study, stem cells were intravenously injected 6h after induction of unilateral pMCAO to evaluate if cellular delivery of BDNF by hMSCs could affect functional recovery and infarct volume. The T2WI showed wild-type hMSCs decreased the size of the lesion area on days 1, 3, and 7 after pMCAO, compared to control group; BDNF overexpression enhanced the positive outcome

compared to hMSCs at day seven. In addition, 2,3,5-triphenyltetrazolium chloride (TTC, see Box 1) staining showed similar results one-week post-MCAO. Behavioral performance, as assessed by treadmill stress test, revealed both MSCs and MSCs^{BDNF} elicited a functional improvement compared to controls; however, the effect was greater in MSCs^{BDNF} group. Lastly, a small number of implanted cells expressed FOX-3 (NeuN, ~8%), neurofilament (NF, ~7%) or glial fibrillary acidic protein (GFAP, 7.5%) proteins at day 7, revealing neuronal and glial developmental lines.

| Box 1: Most frequently employed techniques to visualize infarct area. | | |
|--|---|---|
| Technique | Features | Sample figure |
| TTC staining | The 2,3,5-triphenyltetrazolium chloride (TTC) staining is the most frequently used tool to macroscopically differentiate viable tissue from infarction. This technique is fast and cheap and the results can be obtained in less than an hour. In the viable tissue, the colorless TTC is enzymatically reduced to a red formazan product by dehydrogenases, which are most abundant in mitochondria. Hence, the injured area without viable mitochondria remains unstained. However, transient dehydrogenase activity impairment can lead to over-estimation of the infarct area [44, 45]. |  |
| Nissl staining | Cresyl violet staining is one of the commonly used techniques to visualize experimental brain infarctions. Cresyl violet stains Nissl substance in neurons. TTC and Nissl staining are showing a high degree of correlation in infarct areas; however Nissl staining is rather time consuming [46]. |  |
| HE staining | Hematoxylin / eosin (HE) staining has been used for more than a century to recognize different cell types, as well as morphological changes. Hematoxylin stains nucleic acids, while eosin stains proteins nonspecifically. Although HE stainings have been employed to detect infarct areas in some studies, its |  |

| | | |
|------------|--|---|
| | lack of specificity constitutes a disadvantage compared to Nissl or TTC stainings. | |
| MRI | Unlike all above mentioned techniques that are executed at the end of each set of experiments, magnetic resonance imaging (MRI) can be performed during the experiments to assess the infarct area. Diffusion-weighted imaging (DWI), a form of MR imaging, is more useful for lesion volume detection during the acute infarction phase (usually 3 to 6 hours after stroke). T2 weighted image (T2WI) is one of the basic pulse sequences in MRI, and is usually used to detect infarct areas, days to weeks after stroke [47]. |  |

Another approach by Lee et al. [48] made use of stable BDNF overexpressing human neural stem cell (hNSCs) line (HB1.F3), in a mouse model of intracerebral hemorrhage (ICH). ICH was induced by injection of collagenase type IV into the striatum and seven days later stem cells were administrated into the ipsilateral striatum. Stem cells improved motor function, which was assessed by rotarod and limb placement tests (See Box 2), up to eight weeks post-transplantation. NSCs^{BDNF} enhanced motor outcome compared to NSCs. Over-expression of BDNF increased survival of transplanted cells in the striatum by 1.6-fold after two weeks, and 3-fold after 8 weeks post administration compared to NSCs. At 2- and 8-weeks post-transplantation, large portion of NSCs^{BDNF} expressed the light and heavy chain of neurofilament, MAP2 and GFAP in the vicinity of the hemorrhagic core. Also, immunostaining for von Willebrandt factor (vWF) revealed NSCs^{BDNF} increased number of microvessels by 2-3 fold compared to NSCs and 6-8 fold compared to the control group. Transplanted stem cells reduced the number of apoptotic (TUNEL positive) cells in the hemorrhage core border areas, and the number of apoptotic cells was decreased in the group receiving BDNF over-expressing stem cells. Although ICH increased proapoptotic protein expression of caspase 3, p21 and p53 in the ipsilateral hemisphere, stem cells and more specifically

NSCs^{BDNF} prevented over-expression of those mentioned proapoptotic proteins, 2 and 8 weeks after transplantation. In line with that, the pro- survival signaling molecule Akt1 as well as the antiapoptotic Bcl-2 protein were elevated. Therefore, it seems that NSCs^{BDNF} could exert neoangiogenic and neuroprotective effects in ICH injury sites.

Zhu, Zhao (49) evaluated the effectiveness of rat NSCs transfected with the human BDNF gene. Stem cells were labelled with bromodeoxyuridine (BrdU) and intrastrially implanted three days after 120 min MCAO. The neurological function deficit was assessed using the neurological severity score (NSS) 2 to 12 weeks after transplantation, with no significant differences being observed between the experimental groups in the first 6 weeks. However, NSCs^{BDNF} improved functional recovery from week 8 on compared to non-transplanted control, and improved functional recovery compared to non-transfected NSCs from week 10 on. To track the transplanted stem cells, doublestaining against BrdU and BDNF was performed one week after transplantation and obtained results revealed that around 25% of implanted cells survived in the ischemic areas and expressed BDNF protein. Also, doublestaining against BrdU and neurofilament, 12 weeks after implantation, showed that 20-38% of transplanted NSCs^{BDNF} were able to differentiate into neurons. However, the authors did not provide any data regarding the differentiation ability of naive NSCs.

In another study, Tao, Ji (50) employed human amniotic membrane-derived mesenchymal stem cells (hAMSCs) that were transfected with the BDNF gene. Stem cells implanted into the dorsolateral striatum of female rats subjected to a transient 40min MCAO one day after stroke. On day 21 after hAMSCs therapy, some transplanted cells expressed nestin as neuronal progenitor marker and MAP2 as neuronal marker. At this time point, hAMSCs reduced infarct volume without significant differences between modified and non-modified hAMSCs, and decreased

caspase-3 and iNOS protein expression. Based on beam-walking and rotarod tests (see box 2 for more details) from days 6 to 21, stem cells improved the functional recovery compared to controls and hAMSCs^{BDNF} induced functional motor improvements at earlier time points and/or with greater effects.

| Box 2: Behavioral tests usually employed to evaluate neurological function | |
|---|--|
| Behavioral test | Description |
| Adhesive-removal test | In this test, adhesive paper dots are used as tactile stimuli on the wrist of each forelimb and the time necessary for animals to remove the tape is recorded. |
| Beam-walking test | In this test, the animal is placed on a beam, and beam walking is graded from easily traversed the beam, to unable to walk on the beam. |
| Bilateral grasp test | In this test, the animal's paws is placed on the edge of a box and strength of the hemiparetic paw is graded from grasp well, to unable to grasp with forepaw. |
| Cylinder rearing test | In this test, the animal is placed in a transparent cylinder and the weight-bearing forepaw(s) to contact the wall during a full rear is recorded. |
| Foot-fault test | In this test, the total number of steps (movement of each forelimb) that an animal uses to cross the grid and the total numbers of foot faults for each forelimb is recorded. |
| Hemiparesis score | In this test, the hind limbs of each animal are gently extended with a round-tipped forceps and the flexor response is scored. |
| Ipsilateral circling test | In this test, the extent of circling to the side of the infusion is graded from no circling to always circling. |
| Limb placement test | In this test, four limbs of animals are evaluated by using the edges and top of a counter top. |
| Modified neurological severity score (mNSS) test | NSS is a composite of balance, reflex, sensory, and motor tests and the higher scores represent increased severity in injury. |
| Morris Water Maze test | In this test, the animal is placed in a tank of water with a hidden platform. During training trials, the latency to find the platform location is recorded. During experimental trials, the platform is removed, and the percentage of time spent in the quadrant that normally contains the platform is compared to the time spent in other quadrants. |
| Posture score | In this test, the animal is suspended by the tail and forelimb flexion and body twisting is scored. |
| Roger's tests | In this test, the movement of an animal is scored from no deficit, to not responding to stimulation. |
| Rotarod test | In this test, the animal is placed on a rotating rod and the speed of the rotation is gradually increased. This test usually uses to assess sensorimotor coordination. |

| | |
|-------------------------|---|
| Stepping test | In this test, a wooden ramp with a length around 1 meter connect a platform to the animal's home cage. Initiation time, stepping time and step length is recorded when the animals move from the platform to their home cage. |
| Treadmill (stress) test | In this test, animals are placed on an accelerating treadmill and they should run and maintain their median position on the belt as its speed steadily increases. In the treadmill stress test, to avoid foot-shocks, animals have to move forward. |

Furthermore, Chang, Lee (51) evaluated the therapeutic and neurogenic ability of human NSC line (HB1.F3) over-expressing BDNF following transplantation into the contralateral striatum on day 7 after stroke in a male rat model of 90min MCAO. In this study, just effects of BDNF-overexpressing NSCs were compared to non-transplanted control. Up to eight weeks post implantation, behavioral improvements from week 3 in the rotarod- as well as stepping-tests were obtained. From week 4 on, improvements in the modified neurological severity score (mNSS) test were recorded. From day 5 after contralateral administration, transplanted stem cells were detected in the infarct area and eight weeks after transplantation, a high portion of NSCs^{BDNF} migrated to the ipsilateral damaged area. At this time point, some of the transplanted cells were nestin-, doublecortin (DCX, as migrating neuroblasts)- or MAP2-positive, which shows that some implanted stem cells preserved their stemness potential, while others differentiated toward the neuronal lineage. In addition, some transplanted stem cells co-stained with tyrosine hydroxylase or glutamic acid decarboxylase 65/67 (GAD65/67), indicating differentiation of the transplanted stem cells into dopaminergic and GABAergic neurons, respectively. Also, some migrated cells expressed dopamine- and cAMP-regulated neuronal phosphoprotein (DARPP-32) as striatal projection neurons marker. Furthermore, some NSCs^{BDNF} transplanted cells expressed CXCR4 as chemokine receptor 4 marker that can response to inflammatory signals, such as stromal derived factor-1, following cerebral ischemia.

Jeong and colleagues [52] investigated the effects of intra-striatal transplantation of BDNF modified human bone marrow (BM)-MSCs, three days after 90min MCAO. MSCs^{BDNF} improved the functional recovery from day 14 on up until day 28 after implantation, based on the adhesive-removal test. However, rotarod stress test revealed functional improvements in the MSCs^{BDNF} group just on day 28. Transplanted stem cells decreased the infarct area, which was assessed by TTC staining 14 days after stroke, and BDNF over-expression led to more reduction in lesion size. Immunostaining 7 and 28 days after MCAO showed that cell therapy increased the number of BrdU, BrdU/DCX and BrdU/NeuN positive cells in the sub-ventricular zone indicating enhanced endogenous neurogenesis. For all of these parameters, MSCs^{BDNF} had greater outcome. Stem cell transplantation decreased the number of TUNEL positive cells in the ischemic lesion, and the number of apoptotic cells was much lower in the BDNF over-expressing group. Then, MSCs^{BDNF} could enhance endogenous neurogenesis and protect neural cells from apoptotic death.

van Velthoven et al. [53] investigated the curative potential of mouse MSCs that were genetically modified to over-express BDNF, epidermal growth factor-like 7 (EGFL7), persephin, and sonic hedgehog in the perinatal mouse model of hypoxia-ischemia. Cerebral ischemia was induced by right common carotid artery occlusion, followed by 45min exposure to 10% oxygen and stem cells were administrated intranasally 10 days later. Based on the cylinder rearing test, MSCs^{BDNF} and MSCs^{EGFL7} improved motor function compared to control and MSCs groups at days 21 and 28 after ischemia. Persephin over-expression just improved functional recovery on day 21 compared to control. Mice treated with MSCs^{sonic hedgehog} showed even decreased functional improvements compared to MSCs. In this study, infarct volume was assessed at day 28 by immunostaining against microtubule-associated protein and myelin basic protein to visualize gray and white matters, respectively. Obtained data revealed that MSCs, MSCs^{BDNF} and MSCs^{persephin} decreased

infarct size compared to control, while MSCs^{EGFL7} had no effect. MSCs^{sonic hedgehog} even increased lesion size compared to MSCs. At this time point, although MSCs^{BDNF} elevated number of BrdU-positive cells in the hippocampus, cell therapy did not affect the number of Ki67 positive cells.

To evaluate the therapeutic effects of the BDNF or/and noggin-modified BM-MSCs, Lu, Liu (54) intravenously administered stem cells one day after 60min MCAO. One week after transplantation, transplanted stem cells improved mNSS, and genetically modified stem cells enhanced functional improvements. The recovery was even amplified in the group co-transfected with BDNF and noggin. Western blot and immunohistochemistry analysis in the ipsilateral cerebral cortex showed that modified MSCs increased the expression of VEGF, Bcl-2, p-GSK3 β , and p-Akt, but decreased Bax, TLR4, and MyD88 compared to non-transplanted control and MSCs groups. Also, an ELISA assay revealed transplanted stem cells reduced levels of matrix metalloproteinase 9 as well as reactive oxygen species and co-transfected MSCs^{BDNF-noggin} had the most powerful effect to decrease these factors.

2.2. Neurotrophin-3 (NT-3)

NT-3 is a member of the neurotrophin family that could promote survival of nearly all types of neurons and is involved in differentiation promoting pathways [55]. It was reported that NT-3 elevates the differentiation potential of stem cells toward neurons [56-58], and NT-3 knockout mice showed impaired neurogenesis [59]. In addition, NT-3 was able to stimulate neovascularization in a limb model of ischemia, which was induced by surgically excising the femoral artery [60].

Park and co-workers [61] evaluated the effects of mouse NSCs over-expressing NT3 in hypoxic–ischemic brain injury. Hypoxic–ischemic injury in 7-day old mice was induced by right common

carotid artery occlusion, followed by 2-3h exposure to 8% O₂ and 92% N₂. Stem cells were transplanted 3 days later in two sites at the ipsilateral infarcted hemisphere, which were 1–1.5 mm apart, or the contralateral lateral ventricle. Immunostainings of the brain slices 2-4 weeks after implantation showed NSCs^{NT3} differentiated into neurons at a higher rate in infarct cavity (20%) and penumbra (81%) than NSCs alone (5%). Small portion of NSCs^{NT3} expressed oligodendroglial (0.4%) and astroglial markers (1%). A subpopulation of NSCs^{NT3} derived neurons were either cholinergic (25–40%), glutamatergic (10–20%) or GABAergic (40–50%).

Furthermore, Zhang et al. [62] transplanted NSCs over-expressing human NT3 into the striatum ipsilateral to the injury of adult rats, 7 days after 2h MCAO. The rats that received stem cells exhibited enhanced functional recovery in neurological and behavioral tests, 7 and 14 days after transplantation. The neurological severity score of rats in both modified and non-modified stem cells groups were similar within seven days after implantation. However, animals that received NSCs^{hNT3} displayed a better functional performance than animals transplanted with wild type NSCs, fourteen days after transplantation.

3. Over-expression of glial-derived neurotrophic factor (GDNF)

GDNF belongs to the transforming growth factor beta family, which is able to promote survival of several types of central and peripheral neurons. Previous studies have shown that intracerebroventricular and intracortical [63] or intrastriatal [64] administration of GDNF resulted in protection of cerebral hemispheres from damage and neuroregenerative responses.

Kurozumi, Nakamura (65) compared therapeutic effects of human BM-MSCs over-expressing BDNF, GDNF, NT3 and ciliary neurotrophic factor (CNTF) in a rat model of cerebral ischemia, induced by 90min MCAO. Stem cells transplanted into ipsilateral striatum one day after stroke. At

day 8 and 15 after MCAO, rats that received MSCs^{GDNF} achieved higher limb placement scores compared to control. Similarly, MSCs^{GDNF} could reduce infarct area 7 and 14 days after ischemia. In addition, discosoma red fluorescent protein- positive MSCs were more numerous in the injection site of animals treated with MSCs^{GDNF} than MSCs group, 14 days after MCAO.

Horita et al. [66] investigated whether hMSCs overexpressing GDNF could contribute to functional recovery in a rat pMCAO model. The pMCAO was induced by intraluminal vascular occlusion and β -galactosidase (LacZ) transfected hMSCs^{GDNF} were intravenously administered three hours later. The β -galactosidase positive hMSCs^{GDNF} were detected throughout the injured hemisphere mainly in the penumbra lesion, one week after cell therapy. An estimate of infarct volume was obtained using *in vivo* diffusion-weighted (see Box 1) and T2-weighted magnetic resonance imaging. Based on the diffusion-weighted imaging, stem cells curative effects to reduce infarct size were detected only in the first week after transplantation. However, on T2W imaging, infarct size was decreased in stem cell transplanted groups in all evaluated time points from day 1 up to day 28 after stroke and hMSCs^{GDNF} had better outcome at all time points to reduce infarct area. Also, intravenous delivery of hMSCs-GDNF led to reduction in lesion size, as estimated from TTC staining 1 week after MCAO. By using the treadmill stress test, stem cell transplanted groups showed improvement in treadmill velocity up to 31 days. Again, hMSCs-GDNF had better outcome in all evaluated time points.

In addition, Chen and colleagues [67] transplanted rat NSCs transfected with GDNF into the ipsilateral lateral ventricle, three days after 120min MCAO. Stem cell transplantation led to decreased neurological severity scores from week 1 to 7 compared to non-transplanted control group. NSCs^{GDNF} only at weeks 2 and 3 showed behavioral improvements, compared to naive NSCs. Stem cells reduced infarct volume from week 1 up to week 7 compared to control;

NSCs^{GDNF} only at week 1 had better results compared to NSCs. Also, more GFP and BrdU positive stem cells were detected in the NSCs^{GDNF} group, compared to NSCs up to 7 weeks after stroke. Cell therapy and more specifically NSCs^{GDNF} enhanced synaptophysin and PSD-95 immunoreactivity whereas decreased number of caspase-3 and TUNEL positive cells in all evaluated time points. Elevated expression of synaptic proteins could suggest the reconstruction of neural circuitries and/or enhanced neuronal functioning following stem cell grafting.

Ou and co-workers [68] evaluated the therapeutic effects of human umbilical cord blood (HUCB) CD34+ cells over-expressing GDNF, in spontaneous hypertensive rats exposed to 120min MCAO. At 6h after stroke, stem cells were administered into the tail vein. Cell therapy reduced infarct size 7 days after transplantation and UCBCs^{GDNF} had greater outcome compared to UCBCs. GDNF over-expression increased survival of GFP positive stem cells compared to NSCs in the injured hemisphere. Double immunostaining against GFP and GFAP on day 28 showed more than 75% of the total cells were GFP-GFAP suggesting glial preferential differentiation of transplanted cell. Stem cells improved mNSS scores on days 7 and 28 and UCBCs^{GDNF} enhanced behavioral improvements compared to UCBCs.

Wang, Geng (69) evaluated the therapeutic effect of transplanted GDNF modified rat BM-MSCs on an experimental ischemic brain injury. The stem cells were injected three days after 120min MCAO via the caudal veins. The animals were evaluated at 3, 14, and 28 days after transplantation. At all time points, animals that underwent cell therapy showed behavioral improvements, which was obtained from ipsilateral circling, bilateral grasp and beam walking tests. At days 14 and 28, GDNF modified MSCs enhanced behavioral improvements compared to non-transfected MSCs. Also stem cells reduced TUNEL positive apoptotic cells as well as increased Bcl-2 positive cells

around the ischemic region in evaluated time points and MSCs-GDNF group showed enhanced outcome compared to MSCs.

4. Over-expression of angiogenic factors

4.1. Vascular endothelial growth factor (VEGF)

VEGF has been involved in all aspects of vascular formation including vasculogenesis, angiogenesis and arteriogenesis. In addition, VEGF has neuroprotective effects; therefore it is considered as one of the key elements in stroke [70]. It was reported that intracerebroventricular administration of VEGF peptide after cerebral ischemia in mice and rats led to angiogenesis stimulation, neurological performance improvement, infarct size reduction and enhanced newborn neurons survival in the subventricular zone and dentate gyrus [71, 72].

Considering evidence of functional recovery in stroke animal models following stem cell transplantation as well as VEGF treatment, Miki et al. [73] prepared gene-modified rat BM-MSCs that strongly expressed human VEGF. Stem cells were transplanted intracerebrally 24 hours after 120 min MCAO. Between 14 and 28 days after stroke, the rats treated with MSCs or MSCs-VEGF showed mNSS improvement compared to control. In addition, the MSCs^{VEGF} showed remarkable functional recovery compared to the non-transfected MSCs. On day 14 after ischemia, the infarct size assessed by TTC staining decreased in the stem cell transplanted groups and rats receiving MSCs^{VEGF} showed more reduction than native MSCs group. Brain water content was also measured 2 and 7 days after stroke and no significant difference was detected among the experimental groups at either analysis point.

In another study, Lee and co-workers [74] induced intracerebral hemorrhage by the administration of collagenase type IV into the striatum of mice (gender was not defined). Seven days later,

immortalized cell lines of human neural stem cells (HB1.F3), over-expressing VEGF were transplanted into the ipsilateral striatum. Stem cell transplantations led to a functional improvement in rotarod and modified limb placement tests from eight days post-implantation and the curative effects lasted for up to eight weeks. Enhanced behavioral recovery in NSCs^{VEGF} vs NSCs groups were observed 3 to 9 weeks after grafting. VEGF over-expression led to a two-fold increase in cell survival of implanted NSCs at two weeks post-administration and a three-fold increase at eight weeks after implantation. The majority of grafted NSCs^{VEGF} cells differentiated into either astrocytes (55-65%) or neurons (35-45%). NSCs^{VEGF} promoted proliferation of host endothelial cells/microvessels compared to the non-transfected NSCs at two- and eight-weeks post-grafting. The number of TUNEL-positive apoptotic cells in the hemorrhage core border areas, was lower in the stem cell transplanted groups eight weeks after implantation. Western blot analysis revealed increased expression of proapoptotic proteins Bax and caspase 3 in the control group; however, marked reduction of these proteins was detected in the stem cell transplanted groups. On the other hand, the levels of survival signal molecules Akt1, p110 and p58 as well as anti-apoptotic proteins Bcl-XL and Bcl-2 were higher in the stem cell grafted groups.

4.2. Angiopoietin

Angiopoietin belongs to the vascular growth factors family that plays a role in remodeling, stabilization and maturation of vessels. Angiopoietin-1 (Ang1) is the well-known member of this family and binds to Tie2, a receptor tyrosine kinase. Tie2 is expressed in the choroid plexus as well as on endothelial cells lining blood vessels which induces neovascularization in the brain [75].

In a study conducted by Onda, Honmou (76), human BM-MSCs had been transfected with the Ang-1 gene and were intravenously administered 6h after induction of unilateral permanent cerebral ischemia in rats. At 1, 3, and 7 days after pMCAO, MRI-estimated infarct size was less

in both MSC and MSCs-Ang groups without any significant differences between them. Similar results were obtained by TTC staining 7 days after pMCAO. Stem cell therapy increased capillary vascular volume ratio (ipsilateral/contralateral) at 7 and 28 days after pMCAO. The vascular volume ratio of MSCs^{Ang} treated group was higher than MSCs group in both time points. In addition, stem cell transplantation led to increased regional cerebral blood flow in some regions up to 7 days post-MCAO. From 1 to 7 days after transplantation, both MSCs and MSCs^{Ang} groups had greater maximum velocity on a motor-driven treadmill than control, but the MSCs^{Ang} group attained a higher velocity than the non-transfected MSCs.

Toyama and colleagues [77] investigated whether the combination of Ang1 and VEGF gene-modified human BM-MSCs contribute to functional recovery in a rat model of pMCAO. Stem cells were intravenously administered 6h after occlusion. To evaluate the relative efficacy of MSCs, MSCs^{Ang}, MSCs^{VEGF} and MSCs^{Ang-VEGF} transplantation, infarct volume was estimated using *in vivo* MRI analysis. At 1, 3, 7 and 14 days, infarct size reduced in the MSCs, MSCs^{Ang} and MSCs^{Ang-VEGF} groups; however, infarct area tended to increase in the MSCs^{VEGF} group. The reduction was not significant between MSCs and MSCs^{Ang}, but it was statistically significant between MSCs^{Ang-VEGF} and other groups. Similar results were obtained by TTC staining at day 7. Based on the three-dimensional capillary imaging obtained from systemically perfused fluorescein isothiocyanate –dextran 7 days after MCAO, stem cells led to higher capillary vascular volume ratio and MSCs^{Ang-VEGF} treated group showed higher ratios than others. Also, stem cells except MSCs-VEGF led to increased regional cerebral blood flow in some regions 7 days post transplantation. The treadmill stress test showed that the maximum speed at which the rats could run on a motor driven treadmill was faster in MSCs, MSCs^{Ang} and MSCs^{Ang-VEGF} treated rats from

1 up to 14 days post-MCAO. The MSCs^{VEGF} even reduced maximum velocity compared to the control at day 14.

4.3. Placental growth factor (PlGF)

The angiogenic factor, PlGF, belongs to the VEGFs gene family, and has a 53% homology to VEGF. PlGF has a crucial role in the placental chorion angiopoiesis and maintenance of the placenta development and normal growth. PlGF is also expressed in the CNS and plays a role in cerebral ischemic injury [70]. In this regard, the neuroprotective properties of PlGF have been reported in an *in vitro* model of ischemic condition [78], and PlGF-knockout mice showed a deficit in hypoxia-induced cerebral angiogenesis [79].

In the Liu, Honmou (80) study, PlGF modified human BM-MSCs were intravenously administered 3h after induction of unilateral pMCAO in a rat. An estimated infarct size obtained by *in vivo* MRI revealed that, although lesion volumes 3h after occlusion were the same among the experimental groups, infarct size was smaller 6h after MCAO in the MSCs^{PlGF} group. At 1, 3, 4 and 7 days, infarct size was smaller in both MSCs and MSCs^{PlGF} groups compared to control. The reduction in infarct volume was greater in MSCs^{PlGF} compared to MSCs at 3, 4, and 7 days post-occlusion. Similar results were also obtained by TTC staining at day 7. Three-dimensional analysis of capillary vessels in the lesion site 7 days after MCAO showed the capillary vascular volume ratio was higher in stem cells treated group and the angiogenesis was greater in the MSCs^{PlGF}. This trend was also observed in the number of apoptotic cells; fewer TUNEL-positive cells were found in rats receiving MSCs^{PlGF} than those in MSC and control groups. Based on the limb placement and treadmill stress behavioral tests, MSCs^{PlGF} and MSCs improved functional recovery from day 1 and day 3 respectively up to day 7 after transplantation.

4.4. Hypoxia-inducible factor 1 (HIF-1)

HIF-1 is a master regulator of hypoxia-responsive genes. The expression of a wide variety of genes that facilitate adaptation to low O₂ levels is regulated by HIF-1. It targets different molecules with vast range of functions, including energy metabolism, cell proliferation, erythropoiesis, angiogenesis and vasomotor control. All of these genes may potentially contribute to the recovery of neuronal cells following cerebral ischemia and reperfusion. Due to potential key roles of genes modulated by ischemia-activated HIF-1, it has been proposed that modulating induction and accumulation of HIF-1 is a curative strategy for cerebral ischemia. HIF-1 as a heterodimeric transcription factor contains two subunits of HIF-1 α and HIF-1 β . HIF-1 α protein synthesizes and degrades continuously, therefore it is almost absent in normoxic cells. On the other hand, HIF-1 β does not response to O₂ levels and is constitutively expressed in cells; however it is necessary for hypoxia-induced transcriptional changes mediated by HIF-1. Hence, activation of HIF-1 mainly depends on the HIF-1 α protein level [81, 82].

In this regard, Wu, Chen (83) investigated the efficacy of HIF-1 α modified NSCs in a rat model of stroke induced by a 90 min MCAO. BrdU-labelled stem cells were transplanted one day after MCAO into the lateral ventricle. From day 7 up to day 28 post ischemia, animals receiving NSCs^{HIF1 α} scored better in mNSS than other groups. At days 21 and 28, NSCs group also showed greater functional recovery compared to control. The numbers of BrdU-positive cells in the infarct area were greater in NSCs^{HIF1 α} than NSCs that could differentiate into both neuronal and glial cells. Also, NSCs^{HIF1 α} increased factor VIII-positive endothelial cells at the lesion sites at day 35 after MCAO.

Ye, Ye (84) injected HIF1 α over-expressing BM-MSCs into rat tail vein 3h after permanent MCAO. Rats treated with MSCs^{HIF1 α} had functional improvement on mNSS from day 7 post-

MCAO compared to control. MSCs showed neurological functional recovery compared to control from 14 days after ischemia. MSCs^{HIF1 α} led to better outcome on days 14 and 28. Based on the Morris Water Maze test (see Box 2), stem cells transplantations decreased latency periods and travel distances compared to control at both days 14 and 28 post MCAO without significant difference between transfected and non-transfected MSCs. Just MSCs^{HIF1 α} had a potential to reduce infarct volume at day 7. Tracking of GFP-labelled transplanted MSCs showed the presence of stem cells in the olfactory area at day 1, in the boundary area of the injured cortex, mainly including hypothalamus, thalamus, internal ventricle at day 7 and in the striatum and motor cortex fields adjacent to the infarct region at day 14. Although stem cell administration did not affect the number of apoptotic cells in the cortex, MSCs^{HIF1 α} reduced apoptosis in the hippocampus compared to control at day 7 after MCAO. Double immunofluorescent staining in the hippocampus region at day 7 and cortex at day 14 post MCAO revealed increased number of pax6/DCX cells in the MSCs^{HIF1 α} group, indicating enhanced neuronal proliferation.

Yang, Liu (85) injected BM-MSCs over-expressing HIF1 α into the rat tail vein 6h after 120 min MCAO. Only MSCs^{HIF1 α} improved the mNSS score from days 14 to 28 and reduced infarct size at days 14 and 28. Western blot analysis showed that MSCs^{HIF1 α} elevated VEGF protein expression in the injured hemisphere. Angiogenesis detected by CD105 staining in the MSCs group showed a small amount of scattered microvascularization at day 7 after stroke; the revascularization reached a maximum on day 14 and decreased gradually afterward. In the MSCs-HIF1 α group, density of neomicrovascular vessels was denser at day 7, reached its maximum on day 14 and kept at a relatively high level until day 28.

Ye, Chen (86) evaluated the therapeutic effects of NSCs over-expressing HIF1 α in a rat model of stroke induced by 90 min MCAO. Stem cells were injected into cortical peri-infarct regions of the

rat brain at 24h after MCAO. NSCs^{HIF1 α} treatment showed an improvement in NSS scores compared to control on days 7, 14, 21 and 28 post ischemia. NSCs treatment had a significant functional recovery compared to control on days 21 and 28; however, NSCs^{HIF1 α} had better outcome at these time points. Nissl staining (see Box 1) 5 weeks after ischemia showed that cortical and hemispheric atrophy of the lesion sites was reduced in the NSCs^{HIF1 α} group. Also, the number of BrdU labelled transplanted stem cells in the peri-ischemic area was higher in the genetically modified group, 35 days after MCAO. Double staining against BrdU/NF-200 and BrdU/GFAP showed that transplanted NSCs could differentiate into neuronal and glial cells at day 35. Number of VEGF and vWF positive cells were also increased in the peri-ischemic area in the NSCs-HIF1 α compared to control, 7 and 35 days after ischemia, respectively.

Lin, Zhou (87) aimed to evaluate therapeutic efficiency of rat BM-MSCs over-expressing HIF-1 α in hypoxic–ischemic brain damage. Hypoxic–ischemic injury in rats was induced by left common carotid artery occlusion followed by 2h exposure to 8% O₂ and 92% N₂. chloromethylbenzamido dialkylcarbocyanine (CM-DiI) labelled stem cells intravenously injected into the tail 24h after ischemia. Tracking of CM-DiI labelled stem cells showed the presence of positive cells in the hippocampus on day 7 after ischemia. From day 7 to day 21 after ischemia, the number of CM-DiI positive cells in the hippocampus elevated gradually in a time-dependent manner. Over-expression of HIF-1 α enhanced the migration potential of MSCs toward the injury sites. The spatial performance was assessed by Morris Water Maze on day 14 after ischemia. There was an increase of time in the target quadrant in stem cell transplanted groups compared to control, however, a higher increase in the amount of time have been found in MSCs^{HIF1 α} group. Furthermore, hematoxylin and eosin staining (see Box 1) of the hippocampus showed that stem cells were able

to ameliorate pathological changes that occurred after ischemia, whereas MSCs^{HIF1 α} enhanced therapeutic efficiency.

5. Over-expression of trophic factors

5.1. Fibroblast growth factor (FGF)

The fibroblast growth factors (FGFs) are a family of cell signaling proteins that participate in a broad range of processes, especially as pivotal factors for normal development. FGFs are also crucial for the adult brain maintenance. Hence, FGFs are one of the main elements involved in neuronal survival and synaptic plasticity during both development and adulthood [88]. Previous studies reported that exogenous FGF can reduce infarct size area in acute stroke (3-6h after stroke) and enhance neurological recovery in chronic stroke (72-168h after stroke) [89].

Ikeda, Nonoguchi (90) evaluated the effect of FGF-2 modified BM-MSC in a rat ischemic stroke model. Stem cells administered stereotactically in the striatum 24h after 2h MCAO. Seven days post MCAO, the rats treated with MSC^{FGF2} showed improvements in the mNSS score, compared to control. At 14 and 21 days after cerebral ischemia, functional improvements were also found in the MSC group compared to control; however, MSC^{FGF2} had a better outcome at both time points. Just MSC^{FGF2} decreased infarct volume, 14 days after MCAO.

Ghazavi, Hoseini (91) evaluated the effect of an acute intravenous injection of adipose-derived mesenchymal stem cells transfected with the *Fgf1* gene, 30 min after 30 min MCAO. The main outcomes of this study were assessed 24 h post MCAO. Stem cell transplanted groups showed a functional improvement, as assessed by rotarod and Rotor's tests and reduced apoptotic cells stained by TUNEL. For both parameters, MSC^{FGF1} had better outcome than MSC. Also, based on

TTC data, stem cells reduced lesion volume and there was no significant difference between stem cell transplanted groups.

Zhang, Zhu (92) evaluated the effects of intravenous administration of FGF2 modified NSCs (C17.2 cell line) in a rat model of stroke. Stem cells were transplanted 24h after 2h MCAO. To determine the proliferation potential of administered NSCs in the injured area, animals received BrdU after anesthesia via tail vein on the day of implantation and every day, for 28 days. Behavioral performance showed a remarkable advance in neurological severity scores from 7 days post-MCAO in NSCs^{FGF2} and 14 days post-MCAO in NSCs groups up to 28 days after transplantation. However, no significant differences were observed in the infarct size among experimental groups compared to control. Using pre-labeled cells with CM-DiI, red fluorescent dye, it has been found that FGF2 promoted the migration of NSCs into the injured brain and increased the survival of cells in the infarcted area at day 28. Also, at this time point, around 45% of the transplanted NSCs^{FGF2} and 30% of the NSCs overlapped with BrdU staining. Furthermore, NSCs^{FGF2} elevated the NeuN-positive cells co-localization with CM-DiI, in comparison to the NSCs. The number of GFAP immunoreactive cells co-localized with CM-DiI cells was not significantly different between the two groups. Nestin co-localization with CM-DiI cells was reduced in the NSCs^{FGF2} group, compared to the NSCs group. Hence, it seems that FGF2 accelerated differentiation of NSCs into mature neurons.

5.2. Hepatocyte growth factor (HGF)

HGF is a potent pleiotropic cytokine that is involved in angiogenesis, morphogenesis, mitogenesis, tissue regeneration, and anti-apoptosis in various cell types. Exogenous HGF has been reported to improve the neurological sequelae by decreasing the infarct area size after stroke [93].

Zhao, Nonoguchi (94) evaluated the effects of BM-MSCs overexpressing HGF, which were intrastriatally implanted in the superacute (2h) and acute (24h) phase of rat MCAO model. Superacute MSCs^{HGF} transplantation led to better mNSS from day 4 post-ischemia; however, MSCs improved the behavioral function from day 14 and these effects continued until day 35. Beneficial effects of stem cells that improved neurological deficits after acute transplantation were observed at day 7 and 14 in MSCs^{HGF} transplanted rats, and at day 14 in MSCs group. Three days after superacute transplantation of stem cells, the infarct volume was reduced just in the MSCs^{HGF} transplanted rats compared to control; acute administration did not affect infarct area size at this time point. Superacute transplantation of stem cells led to reduced infarct size on day 14; however, by acute administration, infarct volume was reduced in the MSCs^{HGF} compared to MSCs or control groups. On day 7, the ischemic boundary zone showed a significantly reduced percentage of apoptotic cells in the MSCs^{HGF} treated group compared to other groups. The result was the same when the MSC group was compared to non-transplanted control.

5.3. Pigment epithelium-derived factor (PEDF)

PEDF is a broadly expressed multifunctional member of the serine proteinase inhibitor (serpin) family. This broadly investigated protein plays key roles in a variety of pathophysiological and physiological processes such as inflammation, fibrogenesis, angiogenesis and neuroprotection.

Huang, Ding (95) investigated the protective effect of exosomes isolated from PEDF over-expressing adipose derived stem cells in the rat cerebral ischemia-reperfusion injury model. Daily administration of exosomes through the lateral cerebral ventricle was started 3 days before MCAO. Three days after reperfusion, PEDF-exosome reduced the infarct volume, decreased relative apoptosis rate and prevented MCAO-induced apoptotic pathway activation through reducing cleaved/total caspase-3 as well as cleaved/total caspase-9.

6. Over-expression of genes involved in cell survival and/or migration

6.1. Akt1

A serine/threonine kinase, Akt, plays a crucial role in the regulation of cell survival, growth and proliferation. Akt1 protein is known as a general mediator of cell survival signals in the NSCs. Lee and colleagues reported that grafted human NSCs had low survival rates in ischemia and ICH mice; with less than 50% of transplanted NSCs survived in ICH at 2-weeks post grafting and 30% at 8-weeks [74, 96]. Therefore, it was hypothesized that implanting NSCs that overexpress Akt1 into the damaged area could improve the viability of hNSCs.

Lee, Kim (97) induced ICH by intrastriatal administration of bacterial collagenase type IV and seven days later, human NSCs (HB1.F3), overexpressing Akt1 were transplanted into ipsilateral striatum of mice. Stem cell transplantation resulted in an improved behavioral performance based on the rotarod and limb placement test from 2 up to 8 weeks post transplantation without significant difference between NSCs and NSCs^{Akt1}. Eight weeks after cell therapy, around 35-54% transplanted NSCs^{Akt1} differentiated to NF-H positive neurons; however just around 4% of them were GFAP positive astrocytes. Immunostaining against human nuclear matrix antigen (hNuMA) showed that Akt1 overexpression resulted in a 40% increase in cell survival of transplanted NSCs at 2 weeks post-transplantation and 100% increase at 8 weeks post-transplantation in the hemorrhage core border areas. Also, NSCs^{Akt1} were able to migrate to the hippocampus at week 8. Transplanted NSCs or NSCs^{Akt1} were immuno-negative for the cell proliferation marker Ki-67, indicating that grafted stem cells did not continue to proliferate following transplantation. Furthermore, H&E-stained sections of mouse brains transplanted with NSCs or NSCs^{Akt1} six months post-transplantation showed no sign of tissue distortion or tumor formation.

6.2. B-cell lymphoma 2 (Bcl2)

Bcl-2 is the founding member of the Bcl-2 family of regulator proteins that regulate cell death by either inducing or inhibiting apoptosis. Wei, Cui (98) evaluated the therapeutic effects of mouse embryonic stem cells (ESCs) over-expressing Bcl2 in a rat stroke model induced by 120 min MCAO. BrdU-labelled stem cells were transplanted seven days after MCAO into 4 cerebral regions (Figure 3).




Figure 3: Wei et al. (2005) transplanted embryonic stem cells into 4 different coordinates.

TUNEL staining 3 days after transplantation revealed fewer cell death in ESCs-Bcl2 compared to ESCs group in the ischemic core region. Fourteen days after cell therapy, the number of neuron-like NeuN positive transplanted ESCs or ESCs-Bcl2 were $34 \pm 11\%$ and $58 \pm 7\%$, respectively. Also, some transplanted cells in the post-ischemic cortex and striatum stained positively for GFAP (astrocyte marker), neural/glial antigen-2 (NG-2, oligodendrocyte precursor marker) and adenomatous polyposis coli (APC, mature oligodendrocyte marker). Seven weeks after transplantation, the formation of neural structures was detected by NF staining and possible neovascularization was detected by Glut-1, a marker for differentiated endothelial cells. Stem cell transplantations led to better neurological severity scores from day 7 to 35. After 21 days of

transplantation, animals transplanted with ESCs^{Bcl2} showed an enhanced neurological recovery rate than animals transplanted with ESCs.

6.3. C-X-C chemokine receptor type 4 (CXCR-4)

SDF1 (stromal cell-derived factor 1), also known as CXCL12 (C-X-C motif chemokine 12), belongs to the CXC subfamily of chemokines. SDF1 is considered to preserve adult and embryonic NSCs and being involved in the recruitment of NSCs to damaged regions to enhance recovery [99]. Furthermore, it has been reported that SDF-1 is expressed in the ischemic brain and facilitates the migration of transplanted cells into the ischemic zone [100, 101]. However, due to the low levels of endogenous SDF-1 receptor (CXCR4), stem cells migrate slowly toward the injury. Therefore, it was hypothesized that overexpression of CXCR4 might accelerate the mobilization of stem cells toward the ischemic area.

In this regard, Yu, Chen (102) intravenously injected rat BM-MSCs over-expressing CXCR4, one day after 120min MCAO and the target parameters were evaluated 7 days post-ischemia. Analysis of eGFP-labelled stem cells showed that the percentage of migrating MSCs in the ischemic brain, especially within the ischemic boundary zone, was higher in the MSCs^{CXCR4} group. Total neurological scores and infarct volume decreased in stem cell transplanted groups with better outcome in the CXCR4 over-expressing group. Also, co-staining of eGFP/neuron specific enolase (NSE), eGFP/GFAP and eGFP/vWF revealed that transplanted MSCs^{CXCR4} could differentiate into neurons, astrocytes and vascular-endothelial cells. An increased capillary vascular volume ratio was observed in the stem cell groups with a higher ratio in MSCs^{CXCR4}.

In a very similar experiment, Bang, Jin (103) just used human BM-MSCs over-expressing CXCR4. At days 3 and 7 post-ischemia, stem cell treated groups showed better mNSS compared to control.

At a later time point (days 14), the degree of improvement was more pronounced in MSCs^{CXCR4} than MSCs. However, MRI estimations of the lesion size showed no significant difference between the groups at day 14. Immunohistochemical stainings for human nuclear and also polymerase chain reaction for human chromosome 7 alphoid repeats were performed to evaluate the migration of stem cells. The obtained results showed that the migration of stem cells to the brain and in the ischemic boundary zone was improved in the genetically modified MSCs compared to naive ones, 3 days after injection.

6.4. Copper/zinc-superoxide dismutase (SOD1)

Copper/zinc-SOD is an oxidoreductase enzyme responsible for the very rapid two-step dismutation of the toxic superoxide radical to molecular oxygen and hydrogen peroxide, through alternate reduction and oxidation of the active-site copper. Sakata, Niizuma (104), isolated NSCs from postnatal day 1 wild-type, heterozygous SOD1 over-expressing and SOD1 knockout mice. Stem cells were injected into the cortex at 3 coordinates, two days after ischemia. NSCs-SOD1 reduced number of TUNEL-positive cells by 42% compared to NSCs in the peri-infarct cortex, 2 days after transplantation. Twenty-eight days after MCAO, the number of surviving GFP positive implanted cells in the injured brains was higher in the NSCs^{SOD1} group compared to NSCs group. In contrast, transplanted cell survival was lower in the knockout group than in the NSCs. At this time point, the percentage of beta III tubulin⁺ neurons (8.9 ± 1.0 , 8.8 ± 1.1 , 9.3 ± 1.4) and GFAP⁺ astrocytes (40.1 ± 4.3 , 37.1 ± 5.9 , 38.3 ± 8.0) differentiating from the implanted NSCs, was similar among the wild-type, SOD over-expressing and SOD knockout stem cells. To assess blood vessel density, 14 days after ischemia, DyLight 594-labeled Lycopersicon esculentum lectin were injected into the jugular vein and 30 min later animals were killed. Higher blood vessel density was observed in the NSCs^{SOD1} group than in the non-transplanted and NSCs groups. However, enhanced angiogenesis

was not observed in the wild-type and knockout groups. Similar results were also obtained by ELISA analysis of VEGF in the cortex, 4 days after stroke. Hematoxylin and eosin staining 28 days after ischemia showed that NSCs^{SOD1} decreased the cortical infarct size by 19.8 and 13.1% compared to non-transplanted and NSCs groups. No changes were observed in striatal infarct size among the experimental groups. According to the Rotarod test, functional improvements were observed in NSCs^{SOD1} compared to the non-transplanted and NSCs groups from day 7 and 21, respectively. However, statistical significance was not reached by the modified neurological severity scores. Significant behavioral improvement was not observed in the wild-type and knockout NSCs.

6.5. Interleukin-10 (IL-10)

IL-10 is a pleiotropic anti-inflammatory cytokine that regulates inflammatory responses. IL-10 is mainly synthesized by type-2 helper T cells and inhibits pro-inflammatory cytokine release, T cell proliferation and macrophage activation [105]. It has been reported that intracerebroventricular injection of IL-10 [106] or using transgenic mice over-expressing murine IL-10 [107] ameliorated devastating conditions of stroke by upregulating anti-apoptotic proteins and attenuating pro-inflammatory signals.

In line with these studies, Nakajima, Nito (108) intravenously injected human BM-MSCs over-expressing IL-10, at 0 or 3h after 90min MCAO. Quantitative analysis of infarct volume by TTC staining showed that the transplantation of stem cells immediately after MCAO led to reduction of infarct size 3 and 7 days after stroke, with better results in MSCs^{IL10}. However, when stem cells were injected 3h after MCAO, MSCs^{IL10} reduced the infarct area, when compared to MSCs and control groups. Stem cell therapy immediately after MCAO led to improvements in neurological scores and motor function 7 days after stroke. However, when stem cells were injected 3h after

MCAO, MSCs^{IL10} improved posture score, hemiparesis score and rotarod performance (see Box 2). Immunostaining against ionized calcium binding adaptor molecule 1 (marker of microglial activation), TNF α (pro-inflammatory cytokine) and Fluoro-Jade C (marker of neurodegeneration) in the cortical ischemic boundary zone showed fewer numbers of positive cells in the stem cell-transplanted groups compared to control at 24 and 72h after ischemia; the reduction was markedly lower in MSCs^{IL10} group. Similar results were obtained by ELISA in the levels of IL-6, IL-1 β and TNF α in the ischemic hemisphere extracts at 72h post-ischemia. Finally, quantification of engrafted MSCs using real-time polymerase chain reaction with human-specific Alu sequences showed higher expression in MSCs^{IL10} than MSCs at 3 and 7 days after ischemia in the ipsilateral hemisphere.

6.6. Survivin (SVV)

SVV is one of the members of the apoptosis inhibitor family. The SVV protein functions to inhibit caspase activation, therefore causing negative regulation of programmed cell death or apoptosis. It has been reported that SVV-modified MSCs can further improve the cardiac performance of rats after myocardial infarction by enhancing survival of the transplanted cells [109]. Therefore, Liu, Zhang (110) evaluated therapeutic potential of rat BM-MSCs over-expressing SVV in the rat model of stroke induced by 120min MCAO. Stem cells were transplanted into tail vein 24h after ischemia. Number of the GFP-positive MSCs in the SVV group elevated by about 1.3-fold at 4 days after administration, and by 3.4-fold at 14 days after administration in MSC group. There were very few GFP-positive cells co-expressing NeuN in the stem cell transplantation groups. Stem cell administration resulted in higher protein expression levels of VEGF and FGF-2 in the injured cerebral tissue, 4 days after cell therapy; however, just MSCs^{SVV} could elevate the target protein levels at day 14. TTC staining, fourteen days after transplantation, showed reduced infarct

size in the stem cell treated groups compared to non-transplanted control with higher reduction in the MSCs^{SVV} group. Similar patterns were also observed in the behavioral improvements at this time point.

7. Over-expression of microRNAs

7.1. microRNA 133b (miR-133b)

miRNAs are 18–25 nucleotide, non-protein coding, evolutionarily conserved, transcripts that control gene expression via translational repression or mRNA degradation or both. miRNAs exert key roles in variety of regulatory mechanisms including host-pathogen interactions and developmental timing as well as tumorigenesis, apoptosis, proliferation and differentiation in different organisms [111].

In this regard, Xin, Li (112) used rat BM-MSCs, genetically modified for miR-133b in a rat model of cerebral ischemia. Stem cells were injected via the tail vein one day after 120min MCAO. Although, MSCs improved adhesive-removal and foot-fault functional scores (see Box 2) compared to non-transplanted control 14 days after stroke, functional improvements in the MSCs^{miR133b} group were seen from day 7 on with a better outcome at day 14. Intracortical axonal density was increased in the MSCs compared to control group at day 14 after MCAO. MSCs^{miR133} treatment further increased cortical axonal density at this time point compared to MSC. Compared to control, the positive areas of synaptophysin (marker for synapses), NF-200 (marker for apical dendrites of large cortical pyramidal neurons) and Bielshowsky silver (marker of neuronal fibers) staining increased at day 14 following ischemia along the ischemic boundary zone in the MSC treated group. MSCs^{miR133} enhanced the positive areas compared to MSC.

In another study, Huang, Jiang (113) intravenously transplanted rat BM-MSCs genetically modified for miR-133b, three days after 90min MCAO. Only MSCs^{miR133} could improve the modified neurological severity score, 14 days after treatment. Also, immunostaining against NeuN showed that exogenous stem cells survived and differentiated to neurons; however, the differentiation potential of MSCs is unaffected by miR-133b modification.

8. Epidermal neural crest stem cells as a promising candidate in stroke

Based on the above presented studies, BM-MSCs and neural stem cells are the most abundantly used cell-types that can be genetically modified to enhance their potential when transplanted in animal models of stroke. However, our investigations revealed that epidermal neural crest stem cells (EPI-NCSCs) can also benefit the inhospitable context of rat model of ischemic stroke [34] as well as *ex vivo* [114] and *in vivo* [115] models of spinal cord injury. EPI-NCSCs are located in the bulge area of the hair follicle, and retain the differentiation potential of their neural crest origin, so that they can differentiate into neural lineages [116]. Unlike BM-MSCs [117], that their proliferation and differentiation capacity dramatically decline with aging, recent reports revealed that EPI-NCSCs of aged donors can maintain their multipotency both *in vitro* and *in vivo* [118]. This advantage over BM-MSCs supports the application of these stem cells to elderly individuals, who have an increased incidence of stroke. According to accumulating evidence, EPI-NCSCs express several trophic factors, which their rate of expression can be manipulated through the use of different chemical agents [119-122] to achieve optimum efficiency after transplantation. In addition, administration of EPI-NCSCs via intra-arterial or intra venous routes following reperfusion, created a comparable outcome to intra-arterial grafted BM-MSCs, 7 days after cerebral ischemia [34]. It has been implicated that grafted EPI-NCSCs are mostly acting through the secretion of various trophic factors such as BDNF, GDNF, NT3, NGF and VEGF that can

impose different modulatory functions [123]. Interestingly, according to unpublished data from our group, the expression level of all aforementioned trophic factors, except NT3, is higher than cultured astrocytes. This striking data supports the beneficial function of these grafted stem cells in damaged tissues. Furthermore, EPI-NCSCs showed superior immunomodulation properties in animal models of nerve injury and stroke, as they upregulated the anti-inflammatory cytokines, whilst reducing the expression of pro-inflammatory cytokines [34, 124, 125]. Since EPI-NCSCs can easily be transduced to overexpress GFP [126], genetic manipulations to elevate neurotrophic factor expression seems feasible, and with this review we hope to prompt further studies investigating the beneficial characteristics of those cells. Taken together, owing to neuromodulatory properties of EPI-NCSCs and their ability to release trophic factors, these stem cells can be considered as a promising cell type to treat various conditions such as stroke or even the newly widespread COVID-19 infections [127].

9. Conclusion

Up until now, therapeutic benefits of a wide variety of stem cells have been demonstrated in the context of stroke. Furthermore, as reviewed here, genetically modified stem cells over-expressing specific proteins, could elevate the restorative potential of naive stem cells, by the enhanced survival/differentiation potential of transplanted cells, apoptosis inhibition, infarct volume reduction or neovascularization which eventually may lead to functional improvements (Table 1). Among the different cell types that were discussed here, epidermal neural crest stem cells are attractive candidate that can be considered for genetic modifications in the context of stroke, due to their expression profile of neurotrophic factors and neuromodulatory cytokines. In addition, since the majority of investigations have focused on the short-term curative effects of genetically engineered stem cells, further studies are required to clarify their long-term impacts.

Acknowledgments

This study was financially supported by Shiraz University of Medical Sciences (grant number: 99-01-106-23536). Partial support was also provided by International Brain Research Organization (IBRO) Research Fellowship award 2019 received by the first author.

Conflict of Interest

The authors have no conflicts of interest to declare that are relevant to the content of this article.

10. References

1. Johnson, C. O., Nguyen, M., Roth, G. A., Nichols, E., Alam, T., Abate, D., et al. (2019). Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology*, 18(5), 439-458.
2. Borhani-Haghighi, A., Safari, R., Heydari, S. T., Soleimani, F., Sharifian, M., Kashkuli, S. Y., et al. (2013). Hospital mortality associated with stroke in southern Iran. *Iranian Journal of Medical Sciences*, 38(4), 314.
3. Shahtaheri, R. A., Borhani Haghighi, A., Safari, A., Cruz-Flores, S. (2012). Recombinant tissue plasminogen activator (rtPA) and stroke unit for acute ischaemic stroke in developing countries, are they cost-effective? *International Journal of Stroke*, 7(7), E9-E9.
4. Rhim, T., Lee, M. (2016). Targeted delivery of growth factors in ischemic stroke animal models. *Expert Opinion on Drug Delivery*, 13(5), 709-723.
5. Safari, A., Safari, R., Borhani-Haghighi, A. (2016). Immunology of stroke. *Galen Medical Journal*, 5(Suppl 1), 10-17.
6. Dulamea, A. O. (2015). The potential use of mesenchymal stem cells in stroke therapy—from bench to bedside. *Journal of the Neurological Sciences*, 352(1-2), 1-11.
7. Lucia Maria Ferri, A., Bersano, A., Lisini, D., Boncoraglio, G., Frigerio, S., Parati, E. (2016). Mesenchymal stem cells for ischemic stroke: progress and possibilities. *Current Medicinal Chemistry*, 23(16), 1598-1608.
8. Hsuan, Y. C. Y., Lin, C. H., Chang, C. P., Lin, M. T. (2016). Mesenchymal stem cell-based treatments for stroke, neural trauma, and heat stroke. *Brain and Behavior*, 6(10), e00526.

9. Jiang, J., Wang, Y., Liu, B., Chen, X., Zhang, S. (2018). Challenges and research progress of the use of mesenchymal stem cells in the treatment of ischemic stroke. *Brain and Development*, 40(7), 612-626.
10. Toyoshima, A., Yasuhara, T., Date, I. (2017). Mesenchymal stem cell therapy for ischemic stroke. *Acta Medica Okayama*, 71(4), 263-268.
11. Turnbull, M. T., Zubair, A. C., Meschia, J. F., Freeman, W. D. (2019). Mesenchymal stem cells for hemorrhagic stroke: status of preclinical and clinical research. *NPJ Regenerative Medicine*, 4(1), 1-10.
12. Wang, F., Tang, H., Zhu, J., Zhang, J. H. (2018). Transplanting mesenchymal stem cells for treatment of ischemic stroke. *Cell Transplantation*, 27(12), 1825-1834.
13. Li, G., Yu, F., Lei, T., Gao, H., Li, P., Sun, Y., et al. (2016). Bone marrow mesenchymal stem cell therapy in ischemic stroke: mechanisms of action and treatment optimization strategies. *Neural Regeneration Research*, 11(6), 1015-1024.
14. Kuroda, S. (2016). Current opinion of bone marrow stromal cell transplantation for Ischemic stroke. *Neurologia Medico-Chirurgica*, 56(6), 293-301.
15. Gutiérrez-Fernández, M., Otero-Ortega, L., Ramos-Cejudo, J., Rodríguez-Frutos, B., Fuentes, B., Díez-Tejedor, E. (2015). Adipose tissue-derived mesenchymal stem cells as a strategy to improve recovery after stroke. *Expert Opinion on Biological Therapy*, 15(6), 873-881.
16. Li, Y., Hu, G., Cheng, Q. (2015). Implantation of human umbilical cord mesenchymal stem cells for ischemic stroke: perspectives and challenges. *Frontiers of Medicine*, 9(1), 20-29.
17. Baker, E. W., Kinder, H. A., West, F. D. (2019). Neural stem cell therapy for stroke: A multimechanistic approach to restoring neurological function. *Brain and Behavior*, 9(3), e01214.
18. Bernstock, J. D., Peruzzotti-Jametti, L., Ye, D., Gessler, F. A., Maric, D., Vicario, N., et al. (2017). Neural stem cell transplantation in ischemic stroke: a role for preconditioning and cellular engineering. *Journal of Cerebral Blood Flow & Metabolism*, 37(7), 2314-2319.
19. Boese, A. C., Le, Q.-S. E., Pham, D., Hamblin, M. H., Lee, J.-P. (2018). Neural stem cell therapy for subacute and chronic ischemic stroke. *Stem Cell Research & Therapy*, 9(1), 1-17.
20. Sinden, J. D., Hicks, C., Stroemer, P., Vishnubhatla, I., Corteling, R. (2017). Human neural stem cell therapy for chronic ischemic stroke: charting progress from laboratory to patients. *Stem Cells and Development*, 26(13), 933-947.

21. Zhang, G.-L., Zhu, Z.-H., Wang, Y.-Z. (2019). Neural stem cell transplantation therapy for brain ischemic stroke: Review and perspectives. *World Journal of Stem Cells*, 11(10), 817.
22. Chen, L., Zhang, G., Gu, Y., Guo, X. (2016). Meta-analysis and systematic review of neural stem cells therapy for experimental ischemia stroke in preclinical studies. *Scientific Reports*, 6, 32291
23. Zhang, L., Tao, W., Feng, H., Chen, Y. (2017). Transcriptional and genomic targets of neural stem cells for functional recovery after hemorrhagic stroke. *Stem Cells International*, 2017, 2412890.
24. Kumar, A., Prasad, M., Jali, V., Pandit, A., Misra, S., Kumar, P., et al. (2017). Bone marrow mononuclear cell therapy in ischaemic stroke: a systematic review. *Acta Neurologica Scandinavica*, 135(5), 496-506.
25. Vahidy, F. S., Rahbar, M. H., Zhu, H., Rowan, P. J., Bambhroliya, A. B., Savitz, S. I. (2016). Systematic review and meta-analysis of bone marrow–derived mononuclear cells in animal models of ischemic stroke. *Stroke*, 47(6), 1632-1639.
26. Zhu, Y., Wan, S., Zhan, R.-y. (2012). Inducible pluripotent stem cells for the treatment of ischemic stroke: current status and problems. *Reviews in the Neurosciences*, 23(4), 393-402.
27. Kokaia, Z., Tornerio, D., Lindvall, O. Transplantation of reprogrammed neurons for improved recovery after stroke. *Progress in Brain Research*. 231: Elsevier; 2017. p. 245-263.
28. Yu, F., Li, Y., M Morshead, C. (2013). Induced pluripotent stem cells for the treatment of stroke: the potential and the pitfalls. *Current Stem Cell Research & Therapy*, 8(5), 407-414.
29. Zents, K., Copray, S. (2016). The therapeutic potential of induced pluripotent stem cells after stroke: evidence from rodent models. *Current Stem Cell Research & Therapy*, 11(2), 166-174.
30. Gancheva, M. R., Kremer, K. L., Gronthos, S., Koblar, S. A. (2019). Using dental pulp stem cells for stroke therapy. *Frontiers in Neurology*, 10, 422.
31. Lan, X., Sun, Z., Chu, C., Boltze, J., Li, S. (2019). Dental pulp stem cells: an attractive alternative for cell therapy in ischemic stroke. *Frontiers in Neurology*, 10, 824.
32. Rodrigues, M. C. O., Lippert, T., Nguyen, H., Kaelber, S., Sanberg, P. R., Borlongan, C. V. Menstrual blood-derived stem cells: in vitro and in vivo characterization of functional effects. *Biobanking and Cryopreservation of Stem Cells*: Springer; 2016. p. 111-121.

33. Rodrigues, M. C. O., Dmitriev, D., Rodrigues jr, A., Glover, L. E., Sanberg, P. R., Allickson, J. G., et al. (2012). Menstrual blood transplantation for ischemic stroke: therapeutic mechanisms and practical issues. *Interventional Medicine and Applied Science*, 4(2), 59-68.
34. Salehi, M. S., Pandamooz, S., Safari, A., Jurek, B., Tamadon, A., Namavar, M. R., et al. (2020). Epidermal neural crest stem cell transplantation as a promising therapeutic strategy for ischemic stroke. *CNS Neuroscience & Therapeutics*, 26(7), 670-681.
35. Lin, L., Du, L. (2018). The role of secreted factors in stem cells-mediated immune regulation. *Cellular Immunology*, 326, 24-32.
36. Ferrer, I., Krupinski, J., Goutan, E., Marti, E., Ambrosio, S., Arenas, E. (2001). Brain-derived neurotrophic factor reduces cortical cell death by ischemia after middle cerebral artery occlusion in the rat. *Acta Neuropathologica*, 101(3), 229-238.
37. Endres, M., Fan, G., Hirt, L., Fujii, M., Matsushita, K., Liu, X., et al. (2000). Ischemic brain damage in mice after selectively modifying BDNF or NT4 gene expression. *Journal of Cerebral Blood Flow & Metabolism*, 20(1), 139-144.
38. Yamashita, K., Wiessner, C., Lindholm, D., Thoenen, H., Hossmann, K.-A. (1997). Post-occlusion treatment with BDNF reduces infarct size in a model of permanent occlusion of the middle cerebral artery in rat. *Metabolic Brain Disease*, 12(4), 271-280.
39. Zhang, Y., Pardridge, W. M. (2006). Blood-brain barrier targeting of BDNF improves motor function in rats with middle cerebral artery occlusion. *Brain Research*, 1111(1), 227-229.
40. Schabitz, W.-R., Berger, C., Kollmar, R., Seitz, M., Tanay, E., Kiessling, M., et al. (2004). Effect of brain-derived neurotrophic factor treatment and forced arm use on functional motor recovery after small cortical ischemia. *Stroke*, 35(4), 992-997.
41. Andsberg, G., Kokaia, Z., Klein, R. L., Muzyczka, N., Lindvall, O., Mandel, R. J. (2002). Neuropathological and behavioral consequences of adeno-associated viral vector-mediated continuous intrastriatal neurotrophin delivery in a focal ischemia model in rats. *Neurobiology of Disease*, 9(2), 187-204.
42. Kurozumi, K., Nakamura, K., Tamiya, T., Kawano, Y., Kobune, M., Hirai, S., et al. (2004). BDNF gene-modified mesenchymal stem cells promote functional recovery and reduce infarct size in the rat middle cerebral artery occlusion model. *Molecular Therapy*, 9(2), 189-197.

43. Nomura, T., Honmou, O., Harada, K., Houkin, K., Hamada, H., Kocsis, J. (2005). IV infusion of brain-derived neurotrophic factor gene-modified human mesenchymal stem cells protects against injury in a cerebral ischemia model in adult rat. *Neuroscience*, 136(1), 161-169.
44. Benedek, A., Moricz, K., Juranyi, Z., Giggler, G., Levay, G., Harsing, L. G., Jr., et al. (2006). Use of TTC staining for the evaluation of tissue injury in the early phases of reperfusion after focal cerebral ischemia in rats. *Brain Research*, 1116(1), 159-165.
45. Zille, M., Farr, T. D., Przesdzing, I., Muller, J., Sommer, C., Dirnagl, U., et al. (2012). Visualizing cell death in experimental focal cerebral ischemia: promises, problems, and perspectives. *Journal of Cerebral Blood Flow & Metabolism*, 32(2), 213-231.
46. Tureyen, K., Vemuganti, R., Sailor, K. A., Dempsey, R. J. (2004). Infarct volume quantification in mouse focal cerebral ischemia: a comparison of triphenyltetrazolium chloride and cresyl violet staining techniques. *Journal of Neuroscience Methods*, 139(2), 203-207.
47. Ding, G. L., Chopp, M., Li, L., Zhang, L., Zhang, Z. G., Li, Q. J., et al. (2014). Magnetic resonance imaging of stroke in the rat. *Bo Pu Xue Za Zhi*, 31(1), 116-132.
48. Lee, H. J., Lim, I. J., Lee, M. C., Kim, S. U. (2010). Human neural stem cells genetically modified to overexpress brain-derived neurotrophic factor promote functional recovery and neuroprotection in a mouse stroke model. *Journal of Neuroscience Research*, 88(15), 3282-3294.
49. Zhu, J., Zhao, Y., Chen, S., Zhang, W., Lou, L., Jin, X. (2011). Functional recovery after transplantation of neural stem cells modified by brain-derived neurotrophic factor in rats with cerebral ischaemia. *Journal of International Medical Research*, 39(2), 488-498.
50. Tao, J., Ji, F., Liu, B., Wang, F., Dong, F., Zhu, Y. (2012). Improvement of deficits by transplantation of lentiviral vector-modified human amniotic mesenchymal cells after cerebral ischemia in rats. *Brain Research*, 1448, 1-10.
51. Chang, D.-J., Lee, N., Choi, C., Jeon, I., Oh, S.-H., Shin, D. A., et al. (2013). Therapeutic effect of BDNF-overexpressing human neural stem cells (HB1. F3. BDNF) in a rodent model of middle cerebral artery occlusion. *Cell Transplantation*, 22(8), 1441-1452.
52. Jeong, C. H., Kim, S. M., Lim, J. Y., Ryu, C. H., Jun, J., Jeun, S.-S. (2014). Mesenchymal stem cells expressing brain-derived neurotrophic factor enhance endogenous neurogenesis in an ischemic stroke model. *BioMed Research International*, 2014.

53. Van Velthoven, C. T., Braccioli, L., Willemsen, H. L., Kavelaars, A., Heijnen, C. J. (2014). Therapeutic potential of genetically modified mesenchymal stem cells after neonatal hypoxic-ischemic brain damage. *Molecular Therapy*, 22(3), 645-654.
54. Lu, H., Liu, X., Zhang, N., Zhu, X., Liang, H., Sun, L., et al. (2016). Neuroprotective effects of brain-derived neurotrophic factor and noggin-modified bone mesenchymal stem cells in focal cerebral ischemia in rats. *Journal of Stroke and Cerebrovascular Diseases*, 25(2), 410-418.
55. Patapoutian, A., Reichardt, L. F. (2001). Trk receptors: mediators of neurotrophin action. *Current Opinion in Neurobiology*, 11(3), 272-280.
56. Li, X., Yang, Z., Zhang, A. (2009). The effect of neurotrophin-3/chitosan carriers on the proliferation and differentiation of neural stem cells. *Biomaterials*, 30(28), 4978-4985.
57. Lu, H.-x., Hao, Z.-m., Jiao, Q., Xie, W.-l., Zhang, J.-f., Lu, Y.-f., et al. (2011). Neurotrophin-3 gene transduction of mouse neural stem cells promotes proliferation and neuronal differentiation in organotypic hippocampal slice cultures. *Medical Science Monitor*, 17(11), BR305.
58. Ohtsuka, M., Fukumitsu, H., Furukawa, S. (2009). Neurotrophin-3 stimulates neurogenetic proliferation via the extracellular signal-regulated kinase pathway. *Journal of Neuroscience Research*, 87(2), 301-306.
59. Shimazu, K., Zhao, M., Sakata, K., Akbarian, S., Bates, B., Jaenisch, R., et al. (2006). NT-3 facilitates hippocampal plasticity and learning and memory by regulating neurogenesis. *Learning & Memory*, 13(3), 307-315.
60. Cristofaro, B., Stone, O. A., Caporali, A., Dawbarn, D., Ieronimakis, N., Reyes, M., et al. (2010). Neurotrophin-3 is a novel angiogenic factor capable of therapeutic neovascularization in a mouse model of limb ischemia. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 30(6), 1143-1150.
61. Park, K. I., Himes, B. T., Stieg, P. E., Tessler, A., Fischer, I., Snyder, E. Y. (2006). Neural stem cells may be uniquely suited for combined gene therapy and cell replacement: evidence from engraftment of neurotrophin-3-expressing stem cells in hypoxic-ischemic brain injury. *Experimental Neurology*, 199(1), 179-190.
62. Zhang, Z.-h., Wang, R.-z., Li, G.-l., Wei, J.-j., Li, Z.-j., Feng, M., et al. (2008). Transplantation of neural stem cells modified by human neurotrophin-3 promotes functional recovery after transient focal cerebral ischemia in rats. *Neuroscience Letters*, 444(3), 227-230.

63. Wang, Y., Lin, S.-Z., Chiou, A.-L., Williams, L. R., Hoffer, B. J. (1997). Glial cell line-derived neurotrophic factor protects against ischemia-induced injury in the cerebral cortex. *Journal of Neuroscience*, 17(11), 4341-4348.
64. Kobayashi, T., Ahlenius, H., Thored, P. r., Kobayashi, R., Kokaia, Z., Lindvall, O. (2006). Intracerebral infusion of glial cell line-derived neurotrophic factor promotes striatal neurogenesis after stroke in adult rats. *Stroke*, 37(9), 2361-2367.
65. Kurozumi, K., Nakamura, K., Tamiya, T., Kawano, Y., Ishii, K., Kobune, M., et al. (2005). Mesenchymal stem cells that produce neurotrophic factors reduce ischemic damage in the rat middle cerebral artery occlusion model. *Molecular Therapy*, 11(1), 96-104.
66. Horita, Y., Honmou, O., Harada, K., Houkin, K., Hamada, H., Kocsis, J. D. (2006). Intravenous administration of glial cell line-derived neurotrophic factor gene-modified human mesenchymal stem cells protects against injury in a cerebral ischemia model in the adult rat. *Journal of Neuroscience Research*, 84(7), 1495-1504.
67. Chen, B., Gao, X.-Q., Yang, C.-X., Tan, S.-K., Sun, Z.-L., Yan, N.-H., et al. (2009). Neuroprotective effect of grafting GDNF gene-modified neural stem cells on cerebral ischemia in rats. *Brain Research*, 1284, 1-11.
68. Ou, Y., Yu, S., Kaneko, Y., Tajiri, N., Bae, E. C., Chheda, S. H., et al. (2010). Intravenous infusion of GDNF gene-modified human umbilical cord blood CD34+ cells protects against cerebral ischemic injury in spontaneously hypertensive rats. *Brain Research*, 1366, 217-225.
69. Wang, Y., Geng, T., Ni, A., Yin, H., Han, B. (2011). Effects of transplanted GDNF gene modified marrow stromal cells on focal cerebral ischemia in rats. *Frontiers in Integrative Neuroscience*, 5, 89.
70. Greenberg, D. A., Jin, K. (2013). Vascular endothelial growth factors (VEGFs) and stroke. *Cellular and Molecular Life Sciences*, 70(10), 1753-1761.
71. Kaya, D., Gürsoy-Özdemir, Y., Yemisci, M., Tuncer, N., Aktan, S., Dalkara, T. (2005). VEGF protects brain against focal ischemia without increasing blood–brain permeability when administered intracerebroventricularly. *Journal of Cerebral Blood Flow & Metabolism*, 25(9), 1111-1118.
72. Sun, Y., Jin, K., Xie, L., Childs, J., Mao, X. O., Logvinova, A., et al. (2003). VEGF-induced neuroprotection, neurogenesis, and angiogenesis after focal cerebral ischemia. *The Journal of Clinical Investigation*, 111(12), 1843-1851.

73. Miki, Y., Nonoguchi, N., Ikeda, N., Coffin, R. S., Kuroiwa, T., Miyatake, S.-I. (2007). Vascular endothelial growth factor gene-transferred bone marrow stromal cells engineered with a herpes simplex virus type 1 vector can improve neurological deficits and reduce infarction volume in rat brain ischemia. *Neurosurgery*, 61(3), 586-595.
74. Lee, H. J., Kim, K. S., Park, I. H., Kim, S. U. (2007). Human neural stem cells over-expressing VEGF provide neuroprotection, angiogenesis and functional recovery in mouse stroke model. *PloS one*, 2(1), e156.
75. Ward, N. L., Lamanna, J. C. (2004). The neurovascular unit and its growth factors: coordinated response in the vascular and nervous systems. *Neurological Research*, 26(8), 870-883.
76. Onda, T., Honmou, O., Harada, K., Houkin, K., Hamada, H., Kocsis, J. D. (2008). Therapeutic benefits by human mesenchymal stem cells (hMSCs) and Ang-1 gene-modified hMSCs after cerebral ischemia. *Journal of Cerebral Blood Flow & Metabolism*, 28(2), 329-340.
77. Toyama, K., Honmou, O., Harada, K., Suzuki, J., Houkin, K., Hamada, H., et al. (2009). Therapeutic benefits of angiogenetic gene-modified human mesenchymal stem cells after cerebral ischemia. *Experimental Neurology*, 216(1), 47-55.
78. Du, H., Li, P., Pan, Y., Li, W., Hou, J., Chen, H., et al. (2010). Vascular endothelial growth factor signaling implicated in neuroprotective effects of placental growth factor in an in vitro ischemic model. *Brain Research*, 1357, 1-8.
79. Freitas-Andrade, M., Carmeliet, P., Charlebois, C., Stanimirovic, D. B., Moreno, M. J. (2012). PlGF knockout delays brain vessel growth and maturation upon systemic hypoxic challenge. *Journal of Cerebral Blood Flow & Metabolism*, 32(4), 663-675.
80. Liu, H., Honmou, O., Harada, K., Nakamura, K., Houkin, K., Hamada, H., et al. (2006). Neuroprotection by PlGF gene-modified human mesenchymal stem cells after cerebral ischaemia. *Brain*, 129(10), 2734-2745.
81. Shi, H. (2009). Hypoxia inducible factor 1 as a therapeutic target in ischemic stroke. *Current Medicinal Chemistry*, 16(34), 4593-4600.
82. Davis, C. K., Jain, S. A., Bae, O.-N., Majid, A., Rajanikant, G. (2019). Hypoxia mimetic agents for ischemic stroke. *Frontiers in Cell and Developmental Biology*, 6, 175.
83. Wu, W., Chen, X., Hu, C., Li, J., Yu, Z., Cai, W. (2010). Transplantation of neural stem cells expressing hypoxia-inducible factor-1 α (HIF-1 α) improves behavioral recovery in a rat stroke model. *Journal of Clinical Neuroscience*, 17(1), 92-95.

84. Ye, Z., Ye, W., Deng, Y., Wang, J., Zhou, G., Zhang, X. (2013). HIF-1-modified BMSCs improve migration and reduce neuronal apoptosis after stroke in rats. *Chinese Science Bulletin*, 58(28-29), 3519-3528.
85. Yang, C., Liu, H., Liu, D. (2014). Mutant hypoxia-inducible factor 1 α modified bone marrow mesenchymal stem cells ameliorate cerebral ischemia. *International Journal of Molecular Medicine*, 34(6), 1622-1628.
86. Ye, H., Chen, M.-r., Wu, W.-f. (2016). Transplantation of hypoxia-inducible factor-1 α gene modified neural stem cells increases cell survival and angiogenesis after cerebral ischemia. *Stem Cell and Translational Investigation*, 3.
87. Lin, D., Zhou, L., Wang, B., Liu, L., Cong, L., Hu, C., et al. (2017). Overexpression of HIF-1 α in mesenchymal stem cells contributes to repairing hypoxic-ischemic brain damage in rats. *Comptes Rendus Biologies*, 340(1), 18-24.
88. Guillemot, F., Zimmer, C. (2011). From cradle to grave: the multiple roles of fibroblast growth factors in neural development. *Neuron*, 71(4), 574-588.
89. Lin, D. A., Finklestein, S. P. (1997). Review: Basic fibroblast growth factor: a treatment for stroke? *The Neuroscientist*, 3(4), 247-250.
90. Ikeda, N., Nonoguchi, N., Zhao, M. Z., Watanabe, T., Kajimoto, Y., Furutama, D., et al. (2005). Bone marrow stromal cells that enhanced fibroblast growth factor-2 secretion by herpes simplex virus vector improve neurological outcome after transient focal cerebral ischemia in rats. *Stroke*, 36(12), 2725-2730.
91. Ghazavi, H., Hoseini, S. J., Ebrahimzadeh-Bideskan, A., Mashkani, B., Mehri, S., Ghorbani, A., et al. (2017). Fibroblast growth factor type 1 (FGF1)-overexpressed adipose-derived mesenchymal stem cells (AD-MSC FGF1) induce neuroprotection and functional recovery in a rat stroke model. *Stem Cell Reviews and Reports*, 13(5), 670-685.
92. Zhang, J.-J., Zhu, J.-J., Hu, Y.-B., Xiang, G.-H., Deng, L.-C., Wu, F.-Z., et al. (2017). Transplantation of bFGF-expressing neural stem cells promotes cell migration and functional recovery in rat brain after transient ischemic stroke. *Oncotarget*, 8(60), 102067.
93. Zeng, W., Ju, R., Mao, M. (2015). Therapeutic potential of hepatocyte growth factor against cerebral ischemia. *Experimental and Therapeutic Medicine*, 9(2), 283-288.
94. Zhao, M.-Z., Nonoguchi, N., Ikeda, N., Watanabe, T., Furutama, D., Miyazawa, D., et al. (2006). Novel therapeutic strategy for stroke in rats by bone marrow stromal cells and ex vivo

HGF gene transfer with HSV-1 vector. *Journal of Cerebral Blood Flow & Metabolism*, 26(9), 1176-1188.

95. Huang, X., Ding, J., Li, Y., Liu, W., Ji, J., Wang, H., et al. (2018). Exosomes derived from PEDF modified adipose-derived mesenchymal stem cells ameliorate cerebral ischemia-reperfusion injury by regulation of autophagy and apoptosis. *Experimental Cell Research*, 371(1), 269-277.

96. Lee, H. J., Kim, K. S., Kim, E. J., Choi, H. B., Lee, K. H., Park, I. H., et al. (2007). Brain transplantation of immortalized human neural stem cells promotes functional recovery in mouse intracerebral hemorrhage stroke model. *Stem Cells*, 25(5), 1204-1212.

97. Lee, H. J., Kim, M. K., Kim, H. J., Kim, S. U. (2009). Human neural stem cells genetically modified to overexpress Akt1 provide neuroprotection and functional improvement in mouse stroke model. *PloS one*, 4(5), e5586.

98. Wei, L., Cui, L., Snider, B. J., Rivkin, M., Steven, S. Y., Lee, C.-S., et al. (2005). Transplantation of embryonic stem cells overexpressing Bcl-2 promotes functional recovery after transient cerebral ischemia. *Neurobiology of Disease*, 19(1-2), 183-193.

99. Li, M., Hale, J. S., Rich, J. N., Ransohoff, R. M., Lathia, J. D. (2012). Chemokine CXCL12 in neurodegenerative diseases: an SOS signal for stem cell-based repair. *Trends in Neurosciences*, 35(10), 619-628.

100. Shichinohe, H., Kuroda, S., Yano, S., Hida, K., Iwasaki, Y. (2007). Role of SDF-1/CXCR4 system in survival and migration of bone marrow stromal cells after transplantation into mice cerebral infarct. *Brain Research*, 1183, 138-147.

101. Hill, W. D., Hess, D. C., Martin-Studdard, A., Carothers, J. J., Zheng, J., Hale, D., et al. (2004). SDF-1 (CXCL12) is upregulated in the ischemic penumbra following stroke: association with bone marrow cell homing to injury. *Journal of Neuropathology & Experimental Neurology*, 63(1), 84-96.

102. Yu, X., Chen, D., Zhang, Y., Wu, X., Huang, Z., Zhou, H., et al. (2012). Overexpression of CXCR4 in mesenchymal stem cells promotes migration, neuroprotection and angiogenesis in a rat model of stroke. *Journal of the Neurological Sciences*, 316(1-2), 141-149.

103. Bang, O. Y., Jin, K. S., Hwang, M. N., Kang, H. Y., Kim, B. J., Lee, S. J., et al. (2012). The effect of CXCR4 overexpression on mesenchymal stem cell transplantation in ischemic stroke. *Cell Medicine*, 4(2), 65-76.

104. Sakata, H., Niizuma, K., Wakai, T., Narasimhan, P., Maier, C. M., Chan, P. H. (2012). Neural stem cells genetically modified to overexpress cu/zn-superoxide dismutase enhance amelioration of ischemic stroke in mice. *Stroke*, 43(9), 2423-2429.
105. Kyurkchiev, D., Bochev, I., Ivanova-Todorova, E., Mourdjeva, M., Oreshkova, T., Belemezova, K., et al. (2014). Secretion of immunoregulatory cytokines by mesenchymal stem cells. *World Journal of Stem Cells*, 6(5), 552.
106. Liesz, A., Bauer, A., Hoheisel, J. D., Veltkamp, R. (2014). Intracerebral interleukin-10 injection modulates post-ischemic neuroinflammation: an experimental microarray study. *Neuroscience Letters*, 579, 18-23.
107. De Bilbao, F., Arsenijevic, D., Moll, T., Garcia-Gabay, I., Vallet, P., Langhans, W., et al. (2009). In vivo over-expression of interleukin-10 increases resistance to focal brain ischemia in mice. *Journal of Neurochemistry*, 110(1), 12-22.
108. Nakajima, M., Nito, C., Sowa, K., Suda, S., Nishiyama, Y., Nakamura-Takahashi, A., et al. (2017). Mesenchymal stem cells overexpressing interleukin-10 promote neuroprotection in experimental acute ischemic stroke. *Molecular Therapy-Methods & Clinical Development*, 6, 102-111.
109. Fan, L., Lin, C., Zhuo, S., Chen, L., Liu, N., Luo, Y., et al. (2009). Transplantation with survivin-engineered mesenchymal stem cells results in better prognosis in a rat model of myocardial infarction. *European Journal of Heart Failure*, 11(11), 1023-1030.
110. Liu, N., Zhang, Y., Fan, L., Yuan, M., Du, H., Cheng, R., et al. (2011). Effects of transplantation with bone marrow-derived mesenchymal stem cells modified by Survivin on experimental stroke in rats. *Journal of Translational Medicine*, 9(1), 105.
111. Fiore, R., Siegel, G., Schratt, G. (2008). MicroRNA function in neuronal development, plasticity and disease. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, 1779(8), 471-478.
112. Xin, H., Li, Y., Liu, Z., Wang, X., Shang, X., Cui, Y., et al. (2013). MiR-133b promotes neural plasticity and functional recovery after treatment of stroke with multipotent mesenchymal stromal cells in rats via transfer of exosome-enriched extracellular particles. *Stem Cells*, 31(12), 2737-2746.

113. Huang, B., Jiang, X.-C., Zhang, T.-Y., Hu, Y.-L., Tabata, Y., Chen, Z., et al. (2017). Peptide modified mesenchymal stem cells as targeting delivery system transfected with miR-133b for the treatment of cerebral ischemia. *International Journal of Pharmaceutics*, 531(1), 90-100.
114. Pandamooz, S., Salehi, M. S., Zibaii, M. I., Ahmadiani, A., Nabiuni, M., Dargahi, L. (2018). Epidermal neural crest stem cell-derived glia enhance neurotrophic elements in an ex vivo model of spinal cord injury. *Journal of Cellular Biochemistry*, 119(4), 3486-3496.
115. Hu, Y. F., Gourab, K., Wells, C., Clewes, O., Schmit, B. D., Sieber-Blum, M. (2010). Epidermal neural crest stem cell (EPI-NCSC)--mediated recovery of sensory function in a mouse model of spinal cord injury. *Stem Cell Reviews and Reports*, 6(2), 186-198.
116. Pandamooz, S., Naji, M., Alinezhad, F., Zarghami, A., Pourghasem, M. (2013). The influence of cerebrospinal fluid on epidermal neural crest stem cells may pave the path for cell-based therapy. *Stem Cell Research & Therapy*, 4(4), 84.
117. Ganguly, P., El-Jawhari, J. J., Giannoudis, P. V., Burska, A. N., Ponchel, F., Jones, E. A. (2017). Age-related changes in bone marrow mesenchymal stromal cells: a potential impact on osteoporosis and osteoarthritis development. *Cell Transplant*, 26(9), 1520-1529.
118. Moghadasi Boroujeni, S., Koontz, A., Tseropoulos, G., Kerosuo, L., Mehrotra, P., Bajpai, V. K., et al. (2019). Neural crest stem cells from human epidermis of aged donors maintain their multipotency in vitro and in vivo. *Scientific Reports*, 9(1), 9750.
119. Baharvand, Z., Nabiuni, M., Tahmaseb, M., Amini, E., Pandamooz, S. (2020). Investigating the synergic effects of valproic acid and crocin on BDNF and GDNF expression in epidermal neural crest stem cells. *Acta Neurobiologiae Experimentalis*, 80(1), 38-46.
120. Pandamooz, S., Salehi, M. S., Safari, A., Azarpira, N., Heravi, M., Ahmadiani, A., et al. (2019). Enhancing the expression of neurotrophic factors in epidermal neural crest stem cells by valproic acid: A potential candidate for combinatorial treatment. *Neuroscience Letters*, 704, 8-14.
121. Pandamooz, S., Jafari, A., Salehi, M. S., Jurek, B., Ahmadiani, A., Safari, A., et al. (2020). Substrate stiffness affects the morphology and gene expression of epidermal neural crest stem cells in a short term culture. *Biotechnology and Bioengineering*, 117(2), 305-317.
122. Salehi, M. S., Borhani-Haghighi, A., Pandamooz, S., Safari, A., Dargahi, L., Dianatpour, M., et al. (2019). Dimethyl fumarate up-regulates expression of major neurotrophic factors in the epidermal neural crest stem cells. *Tissue and Cell*, 56, 114-120.

123. Neirinckx, V., Coste, C., Rogister, B., Wislet-Gendebien, S. (2013). Concise review: adult mesenchymal stem cells, adult neural crest stem cells, and therapy of neurological pathologies: a state of play. *Stem Cells Translational Medicine*, 2(4), 284-296.
124. Li, Y., Yao, D., Zhang, J., Liu, B., Zhang, L., Feng, H., et al. (2017). The effects of epidermal neural crest stem cells on local inflammation microenvironment in the defected sciatic nerve of rats. *Frontiers in Molecular Neuroscience*, 10, 133.
125. Zhang, L., Li, B., Liu, B., Dong, Z. (2019). Co-transplantation of epidermal neural crest stem cells and olfactory ensheathing cells repairs sciatic nerve defects in rats. *Frontiers in Cellular Neuroscience*, 13, 253.
126. Pandamooz, S., Salehi, M. S., Nabiuni, M., Dargahi, L., Pourghasem, M. (2016). Evaluation of epidermal neural crest stem cells in organotypic spinal cord slice culture platform. *Folia Biologica (Praha)*, 62(6), 263-267.
127. Salehi, M. S., Pandamooz, S., Jurek, B. (2020). Epidermal neural crest stem cells as a perspective for COVID-19 treatment. *Stem Cell Reviews and Reports*, doi: 10.1007/s12015-020-10028-3.

Table 1: List of summarized manuscripts.

| References | Animal model | Type of ischemia | Type of stem cell | Root of administration | Time of transplantation after stroke | Number of transplanted cells | Modified gene | Evaluated parameters (employed method [time of evaluation after transplantation]) | Key findings |
|------------|---------------------------------|---|--------------------------------------|------------------------------|--------------------------------------|------------------------------|-------------------------|--|--|
| [42] | Adult male Wistar rats | MCAO (90 min) | Human BM-MSCs | Intra-striatum (ipsilateral) | 1 day | 5×10^5 | Over-expression of BDNF | 1. Functional recovery (limb placement and treadmill tests [7d, 14d]) 2. Infarct size (MRI [1d, 6d, 13d]) 3. Apoptosis (TUNEL [6d]) 4. Differentiation (NeuN and GFAP staining [6d]) | <ul style="list-style-type: none">Functional recovery: CTRL=MSCs<MSCs^B_{DNF}Infarct size: CTRL=MSCs>MSCs^B_{DNF}Apoptotic cells: CTRL=MSCs>MSCs^B_{DNF} |
| [43] | Adult male Sprague–Dawley rats | Permanent MCAO | Human BM-MSCs | IV | 6h | 1×10^7 | Over-expression of BDNF | 1. Functional recovery (treadmill stress test [24h, 72h, 7d]) 2. Infarct size (MRI [6h, 24h, 72h, 7d]; TTC staining [7d]) 3. Differentiation (NeuN, NF and GFAP staining [7d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^B_{DNF}Infarct size: CTRL>MSCs>MSCs^B_{DNF} |
| [48] | Adult mice (gender not defined) | Intracerebral hemorrhage (injection of collagenase type IV into striatum) | Human neural stem cell line (HB1.F3) | Intra-striatum (ipsilateral) | 7 days | 2×10^5 | Over-expression of BDNF | 1. Functional recovery (rotarod and limb placement tests [1d, 7d, 14d, 21d, 28d, 35d, 42d, 49d, 56d]) 2. Survival (HuNuA staining [14d, 56d]) 3. Differentiation (NF-L, NF-H, MAP2, GFAP staining [14d, 56d]) 4. Angiogenesis (vWF staining [14d, 56d]) 5. Apoptosis (TUNEL staining [56d]) 6. Gene expression (Western blot analyses of p53, p21, caspase-3, Bcl-2, Akt1 [14d, 56d]) | <ul style="list-style-type: none">Functional recovery: CTRL<NSCs<NSCs^{BD}_{NF}Survival: NSCs<NSCs^{BDNF}Angiogenesis: NSCs<NSCs^{BDNF}Apoptotic cells: CTRL>NSCs>NSCs^{BD}_{NF}Expression of pro-apoptotic proteins: CTRL>NSCs>NSCs^{BD}_{NF} |

| | | | | | | | | | |
|------|--------------------------------|----------------|---|--------------------------------|--------|-----------------|-------------------------|--|--|
| [49] | Adult male Sprague–Dawley rats | MCAO (120 min) | Rat neural stem cells | Intra-striatum (ipsilateral) | 3 days | 5×10^7 | Over-expression of BDNF | <ol style="list-style-type: none">1. Functional recovery (mNSS [14d, 28d, 42d, 56d, 70d, 84d])2. Survival (BrdU and BDNF staining [7d])3. Differentiation (neurofilament staining [84d]) | Functional recovery: CTRL<NSCs<NSCs ^{BDNF} |
| [50] | Female Sprague–Dawley rats | MCAO (40 min) | Human amniotic mesenchymal cells (hAMSCs) | Intra-striatum (ipsilateral) | 1 day | 8×10^5 | Over-expression of BDNF | <ol style="list-style-type: none">1. Functional recovery (neurologic, beam walking and rotarod tests [3d, 6d, 9d, 12d, 15d, 18d, 21d])2. Infarct size (TTC staining [21d])3. Differentiation (MAP2 and Nestin staining [21d])4. Gene expression (Western blot analysis of caspase-3 and iNOS [21d]) | <ul style="list-style-type: none">• Functional recovery: CTRL<hAMSCs<hAMSCs^{BDNF}• Infarct size: CTRL>hAMSCs=hAMSCs^{BDNF}• Expression of caspase-3 and iNOS: CTRL>hAMSCs |
| [51] | Adult male Sprague–Dawley rats | MCAO (90 min) | Human neural stem cell line (HB1.F3) | Intra-striatum (contralateral) | 7 days | 4×10^5 | Over-expression of BDNF | <ol style="list-style-type: none">1. Functional recovery (rotarod, stepping and mNSS tests [7d, 14d, 21d, 28d, 35d, 42d, 49d, 56d]; apomorphine-induced rotation test [14d, 28d, 42d, 56d])2. Survival and migration (MRI [5d, 18d, 32d]; HNu and hMito staining [56d])3. Differentiation (nestin, DCX, NeuN, TH, GAD65/67, DARPP-32, CXCR4, vWF, PCNA staining [56d]) | Functional outcome: CTRL<NSCs ^{BDNF} |
| [52] | Adult male Sprague–Dawley rats | MCAO (90 min) | Human BM-MSCs | Intra-striatum (ipsilateral) | 3 days | 5×10^5 | Over-expression of BDNF | <ol style="list-style-type: none">1. Functional recovery (rotarod and adhesive-removal tests [4d, 11d, 25d])2. Infarct size (staining [11d])3. Differentiation (DCX and NeuN staining [4d, 25d])4. Apoptosis (TUNEL staining [11d]) | <ul style="list-style-type: none">• Functional recovery: CTRL<MSCs<MSCs^{BDNF}• Infarct size: CTRL>MSCs>MSCs^{BDNF}• Endogenous neurogenesis: |

| | | | | | | | | | |
|------|--|---|-------------|------------|---------|-------------------|---|--|--|
| | | | | | | | | | <div>CTRL<MSCs<MSCs^B DNF</div> <ul style="list-style-type: none">Apoptotic cells: CTRL>MSCs>MSCs^B DNF |
| [53] | Postnatal day 9, C57Bl/6J male and female mice | Right common carotid artery occlusion followed by exposure to 10% oxygen in nitrogen (45 min) | Mouse MSCs | Intranasal | 10 days | 5×10 ⁵ | Over-expression of BDNF, EGFL7, persephin or sonic hedgehog | <div>1. Functional recovery (cylinder rearing test [11d, 18d])</div> <div>2. Infarct volume (MAP2 and myelin basic protein staining [18d])</div> <div>3. Proliferation (BrdU and Ki67 staining [18d])</div> | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^B DNF CTRL<MSCs<MSCs^E GFL7 CTRL<MSCs>MSCs^{per} sephin CTRL<MSCs>MSCs^{so} nic hedgehogInfarct size: CTRL>MSCs^{BDNF} CTRL=MSCs^{EGFL7} CTRL=MSCs^{persephin} |
| [7] | Sprague Dawley rats | MCAO (60 min) | Rat BM-MSCs | IV | 1 day | Not defined | Over-expression of BDNF or/and Noggin | <div>1. Functional recovery (mNSS [7d])</div> <div>2. Gene expression (Western blot and IHC analyses of VEGF, BAX, Bcl2, GSK3β, p-GSK3β, Akt, p-Akt, TLR4 and MyD88 [7d]; ELISA analyses of MMP-9 and ROS [7d]).</div> | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^B DNF MSCs- Noggin<MSCs^{BDNF}- NogginVEGF, Bcl-2, p-GSK3β, p-Akt expression: CTRL<MSCs<MSCs^B DNF/NogginBax, TLR4, MyD88 expression: CTRL>MSCs>MSCs^B DNF/Noggin |

| | | | | | | | | | |
|------|----------------------------------|---|---|--|--------|---|--|--|---|
| | | | | | | | | | <ul style="list-style-type: none">MMP-9 and ROS expression: CTRL>MSCs>MSCs^B DNF/Noggin->MSCs^{BDNF}-Noggin |
| [61] | Postnatal day 7, CD1 mice | Right common carotid artery occlusion followed by 2-3h exposure to 8% O ₂ and 92% N ₂ | Mouse neural stem cells (C17.2 cell line) | Two sites at ipsilateral infarcted hemisphere OR the contralateral lateral ventricle | 3 days | 3 ×10 ⁵ (ipsilateral hemisphere); 1 ×10 ⁵ (contralateral lateral ventricle) | Over-expression of NT-3 | 1. Functional recovery (treadmill stress test [1d, 3d, 5d, 7d, 11d, 15d, 19d, 23d, 27d, 31d]) 2. Differentiation (NeuN, MAP2, neurofilament, CNPase, O1, GFAP, glutamate, GABA and vesicular acetylcholine transporter staining [14-28d]) | Functional recovery: CTRL<NSCs<NSCs ^{NT3} |
| [62] | Adult male Sprague–Dawley rats | MCAO (120 min) | Rat neural stem cells | Intra-striatum (ipsilateral) at 2 different sites | 7 days | 1×10 ⁵ at each site | Over-expression of NT-3 | Functional recovery (NSS [1d, 7d, 14d]) | Functional recovery: CTRL<NSCs<NSCs ^{NT3} |
| [65] | Adult male Wistar rats | MCAO (90 min) | Human BM-MSCs | Intra-striatum (ipsilateral) | 1 day | 5×10 ⁵ | Over-expression of GDNF, BDNF, NT3 or CNTF | 1. Functional recovery (limb placement test [7d, 14d]) 2. Infarct size (MRI [1d, 6d, 13d]; TTC staining [13d]) 3. Survival (discosoma red fluorescent protein [13d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs^{BDNF}, MSCs^{GDNF}Infarct size: CTRL>MSCs^{BDNF}, MSCs^{GDNF}Survival: MSCs<MSCs^{BDNF}, MSCs^{GDNF} |
| [66] | Adult female Sprague-Dawley rats | Permanent MCAO | Human BM-MSCs | IV | 3h | 1×10 ⁷ | Over-expression of GDNF | 1. Functional recovery Infarct size (MRI [3h, 1d, 3d, 7d, 14d, 28d]; TTC staining [7d]) | <ul style="list-style-type: none">Functional outcome: CTRL<MSCs<MSCs^{GDNF}Infarct size: CTRL>MSCs>MSCs^{GDNF} |
| [67] | Adult male Wistar rats | MCAO (120 min) | Rat neural stem cells | Intra-lateral ventricle (ipsilateral) | 3 days | 5×10 ⁵ | Over-expression of GDNF | 1. Functional recovery (mNSS [4d, 11d, 18d, 32d, 46d]) | <ul style="list-style-type: none">Functional recovery: CTRL<NSCs<NSCs^{GDNF} |

| | | | | | | | | | |
|------|--|----------------|--|------------------------------|--------|-------------------|-------------------------|---|--|
| | | | | | | | | <div>2. Infarct size (H&E [4d, 11d, 18d, 32d, 46d])</div> <div>3. Survival (GFP and BrdU staining [4d, 11d, 18d, 32d, 46d])</div> <div>4. Apoptosis (TUNEL [4d, 11d, 18d, 32d, 46d])</div> <div>5. Gene expression (IHC analyses of synaptophysin PSD-95 and caspase-3 [4d, 11d, 18d, 32d, 46d]; Western blot analyses of BDNF and NT-3 [4d, 11d, 18d, 32d, 46d])</div> | <div>• Infarct size: CTRL>NSCs>NSCs^{GD}_{NF}</div> <div>• Survival: NSCs<NSCs^{GDNF}</div> <div>• Apoptotic cells: CTRL>NSCs>NSCs^{GD}_{NF}</div> <div>• Synaptophysin, PSD-95 and BDNF expression: NSCs<NSCs^{GDNF}</div> <div>• Caspase-3 expression: NSCs>NSCs^{GDNF}</div> |
| [68] | Adult male spontaneous hypertensive rats | MCAO (120 min) | Human umbilical cord blood CD34+ cells | IV | 6h | 1×10 ⁷ | Over-expression of GDNF | <div>1. Functional recovery (mNSS [1d, 7d, 14d, 21d, 28d])</div> <div>2. Survival (GFP staining [7d, 28d])</div> <div>3. Infarct size (TTC staining [7d])</div> <div>4. Differentiation (NeuN, GFAP staining [7d, 28d])</div> | <div>• Functional recovery: CTRL<UCBCs<UCBCs^{GDNF}</div> <div>• Survival: UCBCs<UCBCs^{GDNF}</div> <div>• Infarct size: CTRL>UCBCs>UCBCs^{GDNF}</div> |
| [69] | Adult Sprague–Dawley rats | MCAO (120 min) | Rat BM-MSCs | IV | 3 days | 5×10 ⁶ | Over-expression of GDNF | <div>1. Functional recovery (ipsilateral circling, bilateral grasp and beam walking tests [3d, 14d, 28d])</div> <div>2. Apoptosis (TUNEL staining [3d, 14d, 28d])</div> <div>3. Gene expression (Bcl2 staining [3d, 14d])</div> | <div>• Functional recovery: CTRL<MSCs<MSCs^G_{DNF}</div> <div>• Apoptotic cells: MSCs>MSCs^{GDNF}</div> <div>• Bcl2 expression: CTRL<MSCs<MSCs^G_{DNF}</div> |
| [73] | Adult male Wistar rats | MCAO (120 min) | Rat BM-MSCs | Intra-striatum (ipsilateral) | 1 day | 1×10 ⁶ | Over-expression of VEGF | <div>1. Functional recovery (mNSS [2d, 6d, 13d, 20d, 27d])</div> <div>2. Infarct volume (TTC staining [13d])</div> | <div>• Functional recovery: CTRL<MSCs<MSCs^V_{EGF}</div> |

| | | | | | | | | | |
|------|--------------------------------|--|--------------------------------------|------------------------------|--------|-------------------|-------------------------|---|---|
| | | | | | | | | 3. Brain water content ([1d, 6d]) | <ul style="list-style-type: none">• Infarct size: CTRL>MSCs>MSCs^{VEGF}• Brain water content: CTRL=MSCs=MSCs^{VEGF} |
| [74] | adult mice | Intracerebral hemorrhage (injection of collagenase type IV into striatum) | Human neural stem cell line (HB1.F3) | Intra-striatum (ipsilateral) | 7 days | 2×10 ⁵ | Over-expression of VEGF | <ol style="list-style-type: none">1. Functional recovery (rotarod and modified limb placement tests [1d, 7d, 14d, 21d, 28d, 35d, 42d, 49d, 56d])2. Survival (HuNuA staining [14d, 56d])3. Differentiation (NF-L, NF-H, MAP2 and GFAP staining [56d])4. Angiogenesis (vWF staining [14d, 56d])5. Apoptosis (TUNEL staining [56d])6. Gene expression (Western blot analysis of caspase 3, Bax, Bcl-2, Bcl-xL, Akt1, PI3 kinase p85 and p110) | <ul style="list-style-type: none">• Functional recovery: CTRL<NSCs<NSCs^{VEGF}• Survival: NSCs<NSCs^{VEGF}• Angiogenesis: CTRL<NSCs<NSCs^{VEGF}• Apoptotic cells: CTRL>NSCs• Pro-apoptotic expression: CTRL>NSCs, NSCs^{VEGF}• Anti-apoptotic expression: CTRL<NSCs, NSCs^{VEGF} |
| [76] | Adult male Sprague–Dawley rats | Permanent MCAO | Human BM-MSCs | IV | 6h | 1×10 ⁶ | Over-expression of Ang1 | <ol style="list-style-type: none">1. Functional recovery (treadmill stress test [1d, 3d, 7d])2. Infarct size (MRI [1d, 3d, 7d]; TTC staining [7d])3. Angiogenesis (three-dimensional image acquisition [7d, 28d])4. Regional cerebral blood flow (perfusion weighted imaging [6h, 3d, 7d]) | <ul style="list-style-type: none">• Functional recovery: CTRL<MSCs<MSCs^{An}_{g1}• Infarct size: CTRL>MSCs=MSCs^{An}_{g1}• Angiogenesis: CTRL<MSCs<MSCs^{An}_{g1} |

| | | | | | | | | | |
|------|--------------------------------|----------------|---------------|----|----|-------------------|--|--|---|
| | | | | | | | | | <ul style="list-style-type: none">• rCBF: CTRL<MSCs, MSCs^{Ang1} |
| [77] | Adult male Sprague–Dawley rats | Permanent MCAO | Human BM-MSCs | IV | 6h | 1×10 ⁶ | Over-expression of Ang1, VEGF or Ang1+VEGF | <ol style="list-style-type: none">1. Functional recovery (treadmill stress test [1d, 3d, 7d, 14d])2. Infarct size (MRI [1d, 3d, 7d, 14d]; TTC staining [7d]).3. Angiogenesis (three-dimensional image acquisition [7d])4. Differentiation (vWF staining [7d])5. Regional cerebral blood flow (perfusion weighted imaging [7d]) | <ul style="list-style-type: none">• Functional recovery: CTRL=MSCs^{VEGF}<MSCs, MSCs^{Ang}<MSCs^{Ang-VEGF}• Infarct size: CTRL=MSCs^{VEGF}>MSCs=MSCs^{Ang}>MSCs^{Ang-VEGF}• Angiogenesis: CTRL<MSCs=MSCs^V_{EGF}<MSCs^{Ang}<MSCs^{Ang-VEGF}• Differentiation: CTRL<MSCs=MSCs^V_{EGF}<MSCs^{Ang}<MSCs^{Ang-VEGF}• rCBF: CTRL, MSCs^{VEGF}<MSCs, MSCs^{Ang}<MSCs^{Ang-VEGF} |
| [80] | Adult male Sprague–Dawley rats | Permanent MCAO | Human BM-MSCs | IV | 3h | 1×10 ⁷ | Over-expression of PIGF | <ol style="list-style-type: none">1. Functional recovery (treadmill stress and limb placement tests [3h, 1d, 3d, 4d, 7d])2. Infarct size (MRI [3h, 6h, 1d, 3d, 4d, 7d]; TTC staining [7d])3. Apoptosis (TUNEL staining [7d])4. Angiogenesis (three-dimensional image acquisition [7d]) | <ul style="list-style-type: none">• Functional recovery: CTRL<MSCs<MSCs^{PI}_{GF}• Infarct size: CTRL>MSCs>MSCs^{PI}_{GF}• Apoptotic cells: CTRL>MSCs>MSCs^{PI}_{GF}• Angiogenesis: |

| | | | | | | | | | |
|------|--------------------------------|----------------|-----------------------|---------------------------------------|-------|-------------------|----------------------------------|--|--|
| | | | | | | | | | CTRL<MSCs<MSCs ^{PI} GF |
| [83] | Male Sprague-Dawley rats | MCAO (90 min) | Rat neural stem cells | Intra-lateral ventricle (ipsilateral) | 1 day | 1×10 ⁶ | Over-expression of HIF1 α | 1. Functional recovery (mNSS [6d, 13d, 20d, 27d]) 2. Survival (BrdU staining [34d]) 3. Differentiation (neurofilament and GFAP [34d]) 4. Angiogenesis (factor VIII [34d]) | <ul style="list-style-type: none">Functional recovery: CTRL<NSCs<NSCs^{HIF} 1αSurvival: CTRL<NSCs<NSCs^{HIF} 1αAngiogenesis: CTRL<NSCs<NSCs^{HIF} 1α |
| [84] | Adult male Sprague-Dawley rats | Permanent MCAO | Rat BM-MSCs | IV | 3h | 2×10 ⁶ | Over-expression of HIF1 α | 1. Functional recovery (mNSS [1d, 7d, 14d, 28d]; Morris water-maze test [14d, 28d]) 2. Infarct size (TTC staining [7d]) 3. Apoptosis (TUNEL staining in the hippocampus and cortex [7d]) 4. Gene expression/ differentiation (pax6/DCX staining in the hippocampus [7d] and cortex [14d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{HI} FlαInfarct size: CTRL>MSCs>MSCs^{HI} FlαApoptotic cells: CTRL>MSCs>MSCs^{HI} Flαpax6/DCX expression: CTRL<MSCs<MSCs^{HI} Flα |
| [85] | Adult male Sprague Dawley rats | MCAO (120 min) | Rat BM-MSCs | IV | 6h | 5×10 ⁶ | Over-expression of HIF1 α | 1. Functional recovery (mNSS [7d, 14d, 21d, 28d]) 2. Infarct size (TTC staining [1d, 14d, 28d]) 3. Gene expression (Western blot analysis of VEGF [7d, 14d, 28d]) 4. Microvessel density (CD105 staining [7d, 14d, 28d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{HI} FlαInfarct size: CTRL>MSCs>MSCs^{HI} FlαVEGF expression: CTRL<MSCs<MSCs^{HI} FlαMicrovessel density: |

| | | | | | | | | | |
|------|--------------------------------|--|------------------------------------|------------------------------|--------|-------------------|--------------------------|---|--|
| | | | | | | | | | CTRL<MSCs<MSCs ^{HIF1α} |
| [86] | Adult male Sprague-Dawley rats | MCAO (90 min) | Rat neural stem cells | Intra-cerebral (ipsilateral) | 1 day | 1×10 ⁶ | Over-expression of HIF1α | 1. Functional recovery (mNSS [6d, 13d, 20d, 27d]) 2. Brain atrophy (Nissl staining [34d]) 3. Survival (BrdU staining [34d]) 4. Differentiation (NF-200 and GFAP staining [34d]) 5. Gene expression (VEGF and vWF staining [6d]) | <ul style="list-style-type: none">Functional recovery: CTRL<NSCs<NSCs^{HIF1α}Brain atrophy: CTRL>NSCs>NSCs^{HIF1α}Survival: NSCs<NSCs^{HIF1α}VEGF and vWF expression: NSCs<NSCs^{HIF1α} |
| [87] | Male Sprague-Dawley rats | Left common carotid artery occlusion followed by exposure to 8% oxygen in nitrogen (120 min) | Rat BM-MSCs | IV | 1 day | 5×10 ⁵ | Over-expression of HIF1α | 1. Functional recovery (Morris water maze test [13d]) 2. Migration (Cm-Dil staining [6d, 13d, 20d]) 3. Histopathology (H&E [6d, 13d, 20d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{HIF1α}Migration: MSCs<MSCs^{HIF1α}Pathological changes: MSCs<MSCs^{HIF1α} |
| [90] | Adult male Wistar rats | MCAO (120 min) | Rat BM-MSCs | Intra-striatum (ipsilateral) | 1 day | 1×10 ⁶ | Over-expression of FGF2 | 1. Functional recovery (mNSS [2d, 6d, 13d, 20d]) Infarct size (TTC staining [2d, 13d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{FGF2}Infarct size: CTRL>MSCs>MSCs^{FGF2} |
| [91] | Adult male Wistar rats | MCAO (30 min) | Rat adipose mesenchymal stem cells | IV | 30 min | 2×10 ⁶ | Over-expression of FGF1 | 1. Functional recovery (rotarod and Roger's tests [1d]) 2. Infarct size (TTC staining [1d]) 3. Apoptosis (TUNEL staining [1d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{FGF1}Infarct size: CTRL>MSCs=MSCs^{FGF1}Apoptotic cells: |

| | | | | | | | | | |
|------|--------------------------------|---|---|---|--------------------------------|----------------------------------|--------------------------|---|--|
| | | | | | | | | | CTRL>MSCs>MSCs ^{FGF1} |
| [92] | Adult Sprague–Dawley male rats | MCAO (120 min) | Mouse neural stem cells (C17.2 cell line) | IV | 1 day | 5×10 ⁶ | Over-expression of FGF2 | 1. Functional recovery (mNSS [1d, 3d, 7d, 10d, 14d, 21d, 28d]) 2. Infarct size (TTC staining [7d]) 3. Survival (cell tracker CM-DiI dye [28d]) 4. Differentiation (NeuN, GFAP and nestin staining [28d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{FGF2}Infarct size: CTRL>MSCs=MSCs^{FGF2}Survival: MSCs<MSCs^{FGF2} |
| [94] | Adult male Wistar rats | MCAO (120 min) | Rat BM-MSCs | Intra-striatum (ipsilateral) | 2 or 24h | 1×10 ⁶ | Over-expression of HGF | 1. Functional recovery (mNSS [1d, 4d, 7d, 14d, 21d, 28d, 35d or 3d, 6d, 13d]) 2. Infarct size (TTC staining [3d, 14d]) 3. Apoptosis (TUNEL staining [7d]) | <ul style="list-style-type: none">Functional recovery: CTRL<MSCs<MSCs^{HGF}Infarct size: CTRL>MSCs^{HGF}Apoptotic cells: CTRL>MSCs>MSCs^{HGF} |
| [95] | Adult male Sprague–Dawley rats | MCAO (60 min) | Exosomes from PEDF over-expressing rat adipose stem cells | ICV | Daily for 3 days prior to MCAO | | Over-expression of PEDF | 1. Infarct size (TTC staining [3d after MCAO]) 2. Apoptosis (TUNEL staining [3d after MCAO]) 3. Gene expression (Western blot analyses of caspase-3, caspase-9 [3d after MCAO]) | <ul style="list-style-type: none">Infarct size: CTRL>PEDF-exosomeApoptotic cells: CTRL>PEDF-exosome |
| [97] | Mice | Intracerebral hemorrhage (injection of collagenase type IV into striatum) | Human neural stem cell line (HB1.F3) | Intra-cortical (ipsilateral) | 7 days | 2×10 ⁵ | Over-expression of Akt-1 | 1. Functional recovery (rotarod and modified limb placement tests [1d, 7d, 14d, 21d, 28d, 35d, 42d, 49d, 56d]) 2. Survival (hNuMA staining [14d, 56d]) 3. Differentiation (NF-L, NF-H, MAP2, GFAP staining [56d]) | <ul style="list-style-type: none">Functional recovery: CTRL<NSCs=NSCs^{Akt1}Survival: NSCs<NSCs^{Akt1} |
| [98] | Adult male Wistar rats | MCAO (120 min) | Mouse embryonic stem cell | Intra-cerebral at 4 different sites (ipsilateral) | 7 days | 2.5×10 ⁴ at each site | Over-expression of Bcl-2 | 1. Functional recovery (NSS [1d, 7d, 14d, 21d, 35d]) 2. Survival (M2 and M6 staining [7d]) | <ul style="list-style-type: none">Functional recovery: CTRL<ESCs=ESCs^{Bcl2}Survival: |

| | | | | | | | | | |
|-------|--------------------------------|----------------|--|---|--------|--------------------------------|-------------------------------------|--|--|
| | | | | | | | | <div>3. Differentiation (NeuN, GFAP, neural/glial antigen-2, adenomatous polyposis coli, neurofilament staining [14d])</div> <div>4. Vascularization (Glut-1 staining [42d])</div> <div>Apoptosis (TUNEL staining [3d])</div> | <div>ESCs<ESCs^{Bcl2}</div> <ul style="list-style-type: none">• Differentiation: ESCs<ESCs^{Bcl2}• Apoptotic cells: ESCs>ESCs^{Bcl2} |
| [102] | Adult male Sprague–Dawley rats | MCAO (120 min) | Rat BM-MSCs | IV | 1 day | 2×10 ⁶ | Over-expression of CXCR4 | <div>1. Functional recovery (postural reflex and forelimb placing tests [6d])</div> <div>2. Infarct size (TTC staining [6d])</div> <div>3. Differentiation (vWF, neuron specific enolase and GFAP staining [6d])</div> <div>4. Angiogenesis (three-dimensional image acquisition [6d])</div> <div>5. Migration (<i>In vivo</i> migration assay [6d])</div> | <ul style="list-style-type: none">• Functional recovery: CTRL<MSCs<MSCs^C_{XCR4}• Infarct size: CTRL>MSCs>MSCs^C_{XCR4}• Angiogenesis: CTRL<MSCs<MSCs^C_{XCR4}• Migration: MSCs<MSCs^{CXCR4} |
| [103] | Adult male Sprague–Dawley rats | MCAO (120 min) | Human BM-MSCs | IV | 1 day | 2×10 ⁶ | Over-expression of CXCR4 | <div>1. Functional recovery (mNSS [2d, 6d, 13d])</div> <div>2. Infarct size (MRI [13d])</div> <div>3. Survival (HuNuA staining [3d])</div> <div>4. Distribution (PCR [3d])</div> <div>5. Differentiation (NeuN, GFAP staining [3d])</div> | <ul style="list-style-type: none">• Functional recovery: CTRL<MSCs<MSCs^C_{XCR4}• Infarct size: CTRL=MSCs=MSCs^C_{XCR4}• Survival: MSCs<MSCs^{CXCR4} |
| [104] | Adult male C57BL/6 mice | MCAO (45 min) | Neural stem cells isolated from wild-type, SOD1 over-expressing and SOD1 knockout mice | Intra-cortical at 3 different sites (ipsilateral) | 2 days | 1×10 ⁵ at each site | Over-expression or knockout of SOD1 | <div>1. Functional recovery (mNSS and rotarod tests [5d, 12d, 19d, 26d])</div> <div>2. Survival (GFP staining [26d])</div> <div>3. Infarct size (H&E staining [26d])</div> <div>4. Apoptosis (TUNEL staining [2d])</div> <div>5. Differentiation (βIII-tubulin and GFAP staining [26d])</div> | <ul style="list-style-type: none">• Functional recovery, rotarod test: CTRL<NSCs<NSCs^{SO}_{D1}• Functional recovery, mNSS: CTRL=NSCs=NSCs^{SO}_{D1} |

| | | | | | | | | | |
|-------|--------------------------------|---------------|---------------|----|---------|-------------------|-------------------------|--|---|
| | | | | | | | | <div>6. Protein determination (ELISA analysis for VEGF level in the cortex [2d])</div> <div>7. Blood vessel density (DyLight 594-labeled lycopersicon esculentum lectin staining [12d])</div> | <div>• Survival: NSCs<NSCs^{SOD1}</div> <div>• Striatal infarct size: CTRL=NSCs=NSCs^{SO}_{D1}</div> <div>• Cortical infarct size: CTRL>NSCs>NSCs^{SO}_{D1}</div> <div>• Apoptotic cells: NSCs>NSCs^{SOD1}</div> <div>• Differentiation: CTRL=NSCs=NSCs^{SO}_{D1}</div> <div>• VEGF level: CTRL<NSCs<NSCs^{SO}_{D1}</div> <div>• Blood vessel density: CTRL<NSCs<NSCs^{SO}_{D1}</div> |
| [108] | Adult male Sprague-Dawley rats | MCAO (90 min) | Human BM-MSCs | IV | 0 or 3h | 1×10 ⁶ | Over-expression of IL10 | <div>1. Functional recovery (hemiparesis, abnormal posture and rotarod tests [7d])</div> <div>2. Infarct size (TTC staining [3d, 7d])</div> <div>3. Gene expression (ionized calcium binding adaptor molecule 1 and TNF-α staining [1d, 3d])</div> <div>4. Neuronal degeneration (fluoro-jade C staining [1d, 3d])</div> <div>5. Protein determination (ELISA analysis for TNF-α, IL-1β and IL-6 in the ischemic hemisphere [3d])</div> | <div>• Functional recovery: CTRL<MSCs<MSCs^{IL}₁₀</div> <div>• Infarct size: CTRL>MSCs>MSCs^{IL}₁₀</div> <div>• Pro-inflammatory cytokine expression: CTRL>MSCs>MSCs^{IL}₁₀</div> <div>• Neuronal degeneration: CTRL>MSCs>MSCs^{IL}₁₀</div> |

| | | | | | | | | | |
|-------|--------------------------------|----------------|-------------|----|--------|-------------------|-----------------------------|--|--|
| [110] | Adult male Sprague-Dawley rats | MCAO (120 min) | Rat BM-MSCs | IV | 24h | 3×10 ⁶ | Over-expression of SVV | <div>1. Functional recovery (mNSS test [1d, 14d])</div> <div>2. Infarct size (TTC staining [14d])</div> <div>3. Survival (GFP staining [4d, 14d])</div> <div>4. Differentiation (NeuN staining [4d, 14d])</div> <div>5. Gene expression (Western blot analysis of VEGF and bFGF in injured cerebral tissues [4d, 14d])</div> | <div>• Functional recovery: CTRL<MSCs<MSCs^{SV}_v</div> <div>• Infarct size: CTRL>MSCs>MSCs^{SV}_v</div> <div>• Survival: MSCs<MSCs^{SV}_v</div> <div>• VEGF and bFGF expression: CTRL<MSCs<MSCs^{SV}_v</div> |
| [112] | Adult male Wistar rats | MCAO (120 min) | Rat BM-MSCs | IV | 1 day | 3×10 ⁶ | Over-expression of MiR-133b | <div>1. Functional recovery (adhesive-removal and foot-fault tests [2d, 6d, 13d])</div> <div>2. Axonal plasticity (biotinylated dextran amine staining [13d])</div> <div>3. Neurite remodeling (Bielshowsky silver, NF-200, synaptophysin staining [13d])</div> | <div>• Functional recovery: CTRL<MSCs<MSCs^{mi}_{R133}</div> <div>• Axonal density: CTRL<MSCs<MSCs^{mi}_{R133}</div> <div>• Neurite remodeling: CTRL<MSCs<MSCs^{mi}_{R133}</div> |
| [113] | Adult male Sprague-Dawley | MCAO (90 min) | Rat BM-MSCs | IV | 3 days | 2×10 ⁶ | Over-expression of MiR-133b | <div>1. Functional recovery (mNSS [7d, 14d])</div> <div>2. Differentiation (NeuN staining [14d])</div> | <div>Functional recovery: CTRL<MSCs^{miR133b}</div> |

Ang1: Angiopoietin-1; **BDNF:** Brain-derived neurotrophic factor; **CNTF:** Ciliary neurotrophic factor; **CXCR4:** C-X-C chemokine receptor type 4; **DARPP-32:** Dopamine- and cAMP-regulated neuronal phosphoprotein; **DCX:** Doublecortin; **FGF:** Fibroblast growth factor; **GAD65/67:** Glutamic acid decarboxylase; **GDNF:** Glial cell-derived neurotrophic factor; **GFAP:** Glial fibrillary acidic protein; **GFP:** Green fluorescent protein; **H&E:** Hematoxylin and eosin; **HGF:** Hepatocyte growth factor; **HIF1α:** Hypoxia-inducible factor-1α; **hMito:** human mitochondria; **HuNuA:** Human nuclear matrix antigen; **IHC:** Immunohistochemistry; **IL:** Interleukin; **iNOS:** Inducible nitric oxide synthase; **MCAO:** Middle cerebral artery occlusion; **mNSS:** Modified neurological severity score; **MRI:** Magnetic resonance imaging; **NeuN:** Neuronal nuclei; **NF:** Neurofilament; **NF-H:** Neurofilament heavy chain; **NF-L:** Neurofilament light chain; **NT-3:** Neurotrophin-3;

PCNA: Proliferating cell nuclear antigen; **PCR**: Polymerase chain reaction; **PEDF**: Pigment epithelium-derived factor; **PIGF**: Placental growth factor; **rCBF**: Regional blood flow; **ROS**: Reactive oxygen species; **SOD**: Copper/zinc-superoxide dismutase; **SVV**: Survivin; **TH**: Tyrosine hydroxylase; **TNF- α** : Tumor necrosis factor alpha; **TTC**: Triphenyl tetrazolium chloride; **TUNEL**: Terminal deoxynucleotidyl transferase dUTP nick end labeling; **VEGF**: Vascular endothelial growth factor; **vWF**: von Willebrand factor.