

Review

# Multiparametric monitoring of hypnosis and nociception-antinociception balance during general anesthesia - A New Era in Patient Safety Standards and Healthcare Management

Alexandru Florin Rogobete <sup>1,2,3</sup>, Ovidiu Horea Bedreag <sup>1,2,3,#</sup>, Marius Papurica <sup>1,2,3,#</sup>, Sonia Elena Popovici <sup>1,2,3,\*</sup>, Lavinia Melania Bratu <sup>1,\*</sup>, Andreea Rata <sup>4</sup>, Claudiu Rafael Barsac <sup>1,2,3</sup>, Andra Maghiar <sup>1,2</sup>, Dragos Nicolae Garofil <sup>6</sup>, Mihai Negrea <sup>7</sup>, Laura Bostangiu Petcu <sup>8</sup>, Daiana Toma <sup>2,3</sup>, Corina Maria Dumbuleu <sup>2,3</sup>, Samir Rimawi <sup>3</sup> and Dorel Sandesc <sup>1,2,3,#</sup>

<sup>1</sup> Faculty of Medicine, "Victor Babes" University of Medicine and Pharmacy, Timisoara 300041, Romania; alexandru.rogobete@umft.ro (A.F.R.), bedreag.ovidiu@umft.ro (O.H.B.), marius.papurica@gmail.com (M.P.), popovici.sonia@yahoo.com (S.E.P.), claudiu\_barsac@yahoo.com (C.R.B.), andramaghiar@yahoo.com (A.M.), dsandesc@yahoo.com (D.S.), bratu.lavinia@umft.ro (L.M.B.); [dsandesc@yahoo.com](mailto:dsandesc@yahoo.com) (D.S.).

<sup>2</sup> Anaesthesia and Intensive Care Research Center, "Victor Babes" University of Medicine and Pharmacy, Timisoara 300041, Romania;

<sup>3</sup> Clinic of Anaesthesia and Intensive Care, Emergency County Hospital "Pius Brinzeu", Timisoara 300723, Romania; [daiana.toma@yahoo.com](mailto:daiana.toma@yahoo.com) (D.T.); [corina.maria.d@gmail.com](mailto:corina.maria.d@gmail.com) (C.M.D.); [rimawi.samir@gmail.com](mailto:rimawi.samir@gmail.com) (S.R.);

<sup>4</sup> Department of Vascular Surgery, "Victor Babes" University of Medicine and Pharmacy, Timisoara 300041, Romania; [rataandreealuciana@gmail.com](mailto:rataandreealuciana@gmail.com) (A.R.)

<sup>5</sup> Clinic of Vascular Surgery, Emergency County Hospital "Pius Brinzeu", Timisoara 300723, Romania;

<sup>6</sup> Faculty of Medicine, "Carol Davila" University of Medicine and Pharmacy, Bucharest 020021, Romania; [dragosgarofil@gmail.com](mailto:dragosgarofil@gmail.com) (D.N.G.)

<sup>7</sup> Faculty of Political, Administrative and Communication Sciences, "Babes Bolyai" University, Cluj Napoca, Romania; [negrea.mihai@gmail.com](mailto:negrea.mihai@gmail.com) (M.N.)

<sup>8</sup> Faculty of Management, The Bucharest University of Economic Studies, Bucharest 020021, Romania. [laurabostangiu@yahoo.com](mailto:laurabostangiu@yahoo.com) (L.B.P.)

\* Correspondence: [bratu.lavinia@umft.ro](mailto:bratu.lavinia@umft.ro) (L.M.B.) and [popovici.sonia@yahoo.com](mailto:popovici.sonia@yahoo.com) (S.E.P.); Tel.: +40 728 001 971

# O.H.B, M.P. and D.S. have equal contribution

**Abstract:** With the development of general anesthesia techniques and anesthetic substances, brought new horizons for the expansion and improvement of surgical techniques. Nevertheless, more complex surgical procedures brought a higher complexity and longer duration for general anesthesia that led to a series of adverse events such as hemodynamic instability, under- or overdosage of anesthetic drugs, as well as an increased number of post-anesthetic events. In order to adapt the anesthesia according to the particularities of each patient, the multimodal monitoring of these patients is highly recommended. Classically, general anesthesia monitoring consists of the analysis of vital functions and gas exchange. Multimodal monitoring refers to the concomitant monitoring of the degree of hypnosis and the nociceptive-antinociceptive balance. By titrating anesthetic drugs according to these parameters, clinical benefits can be obtained, such as hemodynamic stabilization, reduction of awakening times, and the reduction of post-operative complications. Another important aspect is the impact on the status of inflammation and the redox balance. By minimizing inflammatory and oxidative impact one can achieve a faster recovery that will lead to both increased patient satisfaction and an increase in patient safety. The purpose of this literature review is to present the most modern multimodal monitoring techniques, respectively to discuss the particularities of each technique.

**Keywords:** hypnosis; multimodal monitoring; entropy; qNOX; qCON; bispectral index; surgical plethysmographic index; general anaesthesia; patient safety

---

## 1. Introduction

The rapid developments in the field of anesthesia, including new drugs, new anesthetic techniques and new monitoring devices led to an increased trust in the anesthetic act from the general population and increased addressability towards the surgical services, also promoting the development of more complex surgical techniques. In order to keep up with the demand the multiparametric monitoring techniques in general anesthesia needed to be rapidly adapted leading to shorter waiting times, less postoperative adverse events as well as to an increase in patient safety [1–9].

The state of consciousness is represented by a series of variables that can be experienced and felt, such as perceptions, sensations, emotions and memories. These are a few of the reasons that make the quantitative analysis of these states impossible. One of the first theories was launched in 1949 when Hebb postulated that the physical transposition of a mental representation is given by the neuro-cellular assembly and by the neuronal interconnections [5]. The N-metil-D-aspartate (NMDA) synapses were discovered based on this first theory and after numerous studies, synapses that are predominantly found in the cortex [6–8]. Diverse interactions, ionic exchanges, the production of nitric oxide, and the electrical stimulation generated by the opening and closing of ion channels leads to the formation of inter-neuronal connections and to a complex neuronal activity. The loss of consciousness can have a number of causes, such as anesthesia, cerebral lesions or sleep. In the case of anesthesia the responses of the central nervous system are totally suppressed; the state is reversible and is an attribute of the development of modern medicine that made possible the development of modern surgery and invasive therapeutic and diagnostic techniques [10-25].

Multimodal monitoring techniques in general anesthesia refer to the utilization of all parameters that characterize this process. Therefore we talk about monitoring of the degree of hypnosis, of the nociception-antinociception balance, and of the functionality of the autonomic nervous system [23]. In the classical state of things general anesthesia monitoring includes the evaluation of vital functions such as heart rate, blood pressure, temperature, and other subjective clinical findings. In this situation there is always a risk of under- or overdosage of anesthetics leading to either awareness or an excessive degree of hypnosis with serious impact on the outcome and prognosis of these patients. Clinical signs such as hypertension, tachycardia, and tearing have long been used for guiding general anesthesia but it has already been proven that they are subjective and cannot guide general anesthesia in an individualized manner, in accordance with the real needs of each patient [2,24,25].

The electroencephalography (EEG) represents the recording of postsynaptic potentials in the pyramidal cells of the cerebral cortex. EEG is classified then based on the frequency. It can be recorded on the scalp and forehead using surface electrodes and reflects the metabolic activity of the brain. The metabolic activity of brain cells needs energy. Problems or changes in energy production (increased demand or decrease production) by brain cells can profoundly affect the EEG activity [10–12]. Monitoring of the level of consciousness during general anesthesia based on electroencephalography has become routine practice in the operating room. Both for the patient and for the anesthetist the main concern during general anesthesia is the state of unconsciousness, mainly avoiding the risk of awareness. EEG models are well known to change in pattern with the deepening of anesthesia; therefore the evaluation of the degree of hypnosis requires measurements of the electrical activity of the brain [13–15]. Anesthetics have the brain as a target effect site; therefore this organ needs to be monitored together with the spinal reflexes and the cardiovascular changes such as mean arterial pressure and heart rate. Measuring the depth of anesthesia is based on continuous EEG monitoring. Certain algorithms have been developed able to translate changes in the EEG signals into simple numerical indices that correspond to a certain level of anesthesia, from awake state to deep sleep [3,16,17]. Monitoring the state of consciousness is a complex endeavor and, although this clinical area has evolved rapidly, the benefits of EEG monitoring based anesthesia are still controversial. The

problem lies in the fact that our understanding of the human consciousness state is still incomplete and we still lack information about the exact effects of general anesthesia on the brain. The depth of anesthesia is neither stable, nor constant; it is more of a dynamic action that depends on the balance between the dosage of anesthetic medication and the pain caused by the surgical intervention [18–20].

Using EEG signal in order to monitor the depth of general anesthesia should reduce the incidence of intraoperative awareness and should lead to a reduction in anesthetic medication consumption, as well as reduce the incidence of adverse effects related to anesthesia and lead to shorter recovery times [21,22].

## 2. Multimodal monitoring techniques for the degree of hypnosis

In the current clinical practice achieving an individualized prediction of the response to sedation and hypnosis is not accurate without multiparametric monitoring because of complex factors and variables that interfere with the anesthetic act. Among these the most common are the concomitant administration of anesthetic agents, as well as the different pharmacokinetic response and the individual pharmacodynamic variability. Therefore, real-time monitoring of the effects induced by general anesthesia can bring an important contribution to the optimization of anesthetic dosage and hemodynamic control by the individualized titration of these medications. In order to limit perioperative adverse effects induced by the anesthetic drugs it is recommended to titrate the doses based on the individual clinical response [26–28]. Some of the most common techniques for the evaluation and quantification of the degree of hypnosis are represented by the bispectral index (BIS, Medtronic-Covidien, Dublin, Ireland), Entropy (GE Healthcare, Helsinki, Finland), composite auditory evoked potential index (cAAI, AEP Monitor / 2, Danmeter A/S, Odense, Denmark) and Narcotrend index (NCT, MonitorTechnik, Germany).

The bispectral analysis is a statistical technique that reveals non-linear phenomena such as the electroencephalogram (EEG). The conventional analysis of EEG signals using standard procedures can bring important information regarding the frequency, power and phase of EEG signals. The bispectral analysis of these signals represents a separate technique that analyses sinusoidal segments of the EEG spectrum, showing quantifiable variables in the form of an index (BIS) with clinical applicability. From a practical viewpoint BIS is represented by a numeric interval between 0 and 100, where 0 represents the complete electrical abolition translated through cortical suppression, and 100 is characterized by the conscious (awake) state on the EEG [29].

Another technique used for monitoring and individualizing the degree of hypnosis in patients undergoing general anesthesia is Narcotrend (Monitor Technik, Bad Bramstedt, Germany). It is based on analyzing EEG signal and it classifies the degree of hypnosis in different levels such as A (awake) and F (electrical silence), quantified by the Narcotrend Index, that ranges from 100 (awake) to 0 (electrical silence) [30]. Kreuer et al., in a study that compared the performance of BIS and Narcotrend Index have reported similar effects of the two techniques. This research group have obtained a prediction probability,  $P(K)$  for Narcotrend of  $0.88 \pm 0.03$ , while the  $P(K)$  for BIS was  $0.85 \pm 0.04$ . Furthermore, the mean drug effect,  $k(e0)$  was  $0.2 \pm 0.05 \text{ min}^{-1}$  for Narcotrend and  $0.16 \pm 0.07 \text{ min}^{-1}$  for BIS [31]. A similar study was carried out by Kreuer et al., who also reported similarities between the two techniques. Their study included 50 patients undergoing orthopedic surgery and has reported statistically significant correlations between the D and E segments of Narcotrend and the 64–40 range of BIS [32]. Another study on the impact of hypnosis monitoring by Narcotrend Index in the pediatric patient population has reported strong correlations between the Narcotrend Index and the minimum alveolar concentration (MAC) in patients over 4 months of age [33].

The Auditory evoked potentials (AEPs) represent another technique used for monitoring the degree of hypnosis in patients under general anesthesia [34]. Mantzaridis et al., have studied the AEPs index in patients undergoing orthopedic surgery. The mean value for the index at the beginning of surgery was  $72.5 \pm 11.2$ , followed by a decrease to  $39.6 \pm 6.9$  that correlated with loss of consciousness. After recovery from anesthesia the mean value for AEPs index was  $66.9 \pm 12.5$ , leading to the conclusion that this index is suitable for being used in the current medical practice [35].

On the other hand, the concept of Entropy derives from thermodynamics and is successfully used in the current clinical practice, being applied in the EEG signal analysis. Regarding the mechanism of analysis, the EEG signal is first of all analyzed based on the "Fast Fourier" [28-31] concept for the identification of the sinusoidal compounds. After identifying the spectrum the Shannon function is applied in order to identify the specific values for each identified frequency. The sum of these values results in the numerical values called Spectral Entropy. The first algorithm ever to be used in the clinical practice has been defined and applied in the M-Entropy modules S/5 (GE Healthcare, Helsinki, Finland) [10,29-45]. The EEG data are collected through an adhesive sensor made of three electrodes applied on the fronto-temporal region. Applying this concept for general anesthesia led to the idea that when the brain is in the "awake status" the EEG signals are complex and present with a high degree of irregularity. When the patient is asleep/under general anesthesia the neuronal activity progressively decreases and the EEG complexes become more regular. Applying the principle in the case of Entropy, in patients under general anesthesia a significant difference has been observed regarding the wave spectrum generated, this wave spectrum being directly proportional with the neuronal activity. Because the EEG signals are measured from electrodes placed on the frontal region, a high number of signals are represented by the activity of the muscles from the forehead region and are translated through an electromyography signal (EMG). Therefore, the EEG signals are defined by frequencies up to 32 Hz, while the EMG activity includes signals above 32 Hz. The M-Entropy module (GE Healthcare, Helsinki, Finland) distinguished these two frequencies and generates two different parameters, both having important clinical significance – "State Entropy – SE" and "Response Entropy - RE". SE (0.8 – 32 Hz) reflects the cortical status of the patient, while RE (0.8-47 Hz) includes both the EEG and the EMG activity [12,34-36]. The values of SE are between 0 (suppressed EEG) and 91 ("awake status"), while RE is characterized by values between 0-100. In clinical practice it is recommended to maintain RE/SE between 40 and 60 in order to achieve an adequate degree of hypnosis. Spectral Entropy is based on the analysis of frontal EEG and EMG variations and is a safe and reliable method for monitoring the depth of anesthesia. The Entropy module transforms the irregular content of the EEG signal in an index that reflects the depth of anesthesia. Normally, the signal is acquired from the skin on the forehead and temporal area; hence, it encompasses both an EEG and an EMG component [37]. The index is then calculated as follows: high levels of entropy during anesthesia demonstrate awareness, while very low entropy levels are correlated with a profound state of unconsciousness. Using this parameter will lead to a more rapid awakening of the patient at the end of surgery, lower dosage of anesthetic drugs, and will at the same time prevent intra-anesthetic awareness episodes [32,38-40].

Changes in neuronal activity can be analyzed indirectly through computed tomography with integrated positron emission (PET-CT). This analysis is based especially on the changes in certain variables such as neuronal activity, cerebral blood flow, and cellular metabolism [41]. Thus, specific changes in the glucose metabolism rate and cerebral blood flow can be quantified by using [<sup>18</sup>F] – fluorodeoxyglucose and [<sup>15</sup>O] H<sub>2</sub>O. General anesthetic agents such as sevoflurane and propofol reduce the cerebral blood flow, this effect being more important in the case of propofol. Maksimow et al., have carried out a study regarding the changes in neuronal activity under general anesthesia and have mapped the cerebral areas that better correlated with the EEG signals. The analysis of the regional cerebral blood flow has been studied at different degrees of hypnosis measured by Minimum Alveolar Concentration (MAC): MAC:1, MAC:1.5, and MAC:2 for sevoflurane, and different half maximal effective concentration for propofol (EC<sub>50</sub>) at 30 minutes intervals. For patients in the sevoflurane group they analyzed the End-Tidal Sevoflurane (Et-Sevo): 0% Et-Sevo (patient awake), 2 % Et-Sevo (1 MAC), 3% Et-Sevo (1.5 MAC) and 4 % Et-Sevo (2 MAC), while for the propofol group they analyzed: 0 microg/mL (patient awake), 6 microg/mL (1 EC<sub>50</sub>), 9 microg/mL (1.5 EC<sub>50</sub>) and 12 microg/mL (2 EC<sub>50</sub>). In both groups the Entropy was reduced inversely proportional with the sevoflurane and propofol concentrations, from 73.5 ± 6.5 to 12.2 ± 9.4, and from 70.4 ± 7.1 to 0.6 ± 1 respectively, in the frontal region. In the temporo-occipital region the Entropy analysis was similar following the same dose-dependent trend. Regarding the correlation between EEG/SE analysis and computed tomography, the researchers have found statistically significant correlations for both

drugs, at similar concentrations (1.5 MAC,  $r = 0.81$  și 1.5 EC50,  $r = 0.83$ ). Following this study, Maksimow et al. have validated the fact that spectral Entropy can be used for both sevoflurane and propofol, showing the same regional neuronal activity confirmed through non-invasive PET-CT analysis. The usage of monitoring techniques for the degree of hypnosis in the case of pediatric patients is limited and is not validated. Numerous studies analyze the statistical correlations between BIS and Entropy for different age groups but have not identified strong statistical correlations between BIS/Entropy values and anesthetic drugs concentrations in infants vs. pediatric patients (aged over 1 y.o.) [42]. Davidson et al., have carried out a study regarding the performance of BIS and Entropy for different age groups in pediatric patients. They have analyzed four age groups: 0 – 1 years old (n=8), 1-2 years old (n=10), 2-4 years old (n=18), and 4-12 years old (n=14). Regarding the comparison between Entropy and BIS, above the initial status (awake) they have identified statistically significant differences in the 0-1 years old group, as follows: RE/BIS:45 vs. 84,  $P=0.003$ , SE/BIS:36 vs. 78 ( $P=0.02$ ). Following this study no statistically significant differences have been proven for BIS or for Entropy especially in the 0-1 age group. Interestingly there were no performance differences between BIS and Entropy but applying these techniques in the case of infants should be done with caution. In Table 1 are summarized a series of implications for different monitoring techniques for the degree of hypnosis on the clinical prognostic of patients undergoing general anesthesia [43].

**Table 1.** The impact of monitoring the degree of hypnosis on anesthetic drugs consumption and on time recovery

Author	Parameter/Monitoring technique	Type of general anesthesia	Observations	Reference
Kim et al.	State Entropy (SE)	78 children (age: 3-12) Sevoflurane	↓ sevoflurane consumption ↓ postoperative recovery time ↓ sevoflurane consumption	[44]
Wu et al.	State Entropy (SE)	64 patients Sevoflurane	↓ consumption of antihypertensive drugs ↑ hemodynamic stability	[45]
Vakkuri et al.	State Entropy (SE)	368 patients propofol-alfentanil-N <sub>2</sub> O	↓ propofol consumption ↓ postoperative recovery time	[46]
Talawar et al.	Entropy (SE/RE)	50 patients isofluran-N <sub>2</sub> O	↓ postoperative recovery time	[47]
Elgebaly et al.	Entropy (SE/RE)	propofol	↓ propofol consumption ↑ hemodynamic stability	[48]
Gan et al.	Bispectral index (BIS)	302 patients propofol-alfentanil-N <sub>2</sub> O	↓ propofol consumption ↓ postoperative recovery time ↓ consumption of anesthetic drugs	
Liu et al.	Bispectral index (BIS)	1383 patients Day surgery	↓ incidence of adverse effects (nausea, vomiting) ↓ postoperative recovery time	
Bhardwaj et al.	Bispectral index (BIS)	50 pediatric pts propofol	No effects have been observed regarding the consumption of anesthetic drugs.	[49]




Aime et al.	Bispectral index (BIS) and Entropy (RE/SE)	115 patients Sevoflurane;	No effects on the postoperative recovery time BIS & Entropy: ↓ sevoflurane consumption	[50]
Liao et al.	Bispectral index (BIS) and A-line autoregressive index (AAI)	116 patients; Sevoflurane;	BIS & AAI: ↓ sevoflurane consumption ↓ postoperative recovery time;	[51]
Weber et al.	Composite auditory evoked potential index (cAAI)	20 pediatric patients; TIVA propofol and remifentanil;	↓ propofol consumption; ↑ hemodynamic stability	[52]
Lai et al.	Narcotrend	40 patients; propofol and fentanyl;	↓ propofol consumption ↓ postoperative recovery time; No effects on PONV; No effects on propofol/remifentanil consumption;	[53]
Rundshagen et al.	Narcotrend	48 patients propofol and remifentanil	No effects on postoperative recovery time	[54]

One other widely discussed risk is the incidence of intraoperative awareness that can lead to long term posttraumatic stress disorder. Sebel et al., have carried out a study on the incidence of intraanesthetic awareness analyzing 19575 patients. They have identified 25 patients that presented with awareness, resulting in an incidence of 0.13%. This research group has not found any statistically significant differences regarding the incidence based on sex or age, but increased incidence was associated with higher ASA scores (odds ratio, 2.41; 95%CI, 1.04-5.60 ASA III-V vs. ASA I-II) [55]. Sebel et al., have estimated a rough number of 26000 cases of intraanesthetic awareness throughout a year in the United States, this number being reported to approximately 20 million general anesthesia procedures [55]. Bruhn et al., reported in a similar study an incidence of 0.11% out of 10811 patients [34]. Ekman et al., reported a 0.18% incidence of awareness in a retrospective study that included 7826 patients [56]. For all listed studies the incidence of awareness was lower in the groups of patients where techniques for monitoring the degree of hypnosis were used [34,55-56].

### 3. Monitoring techniques for the nociception-antinociception balance

Another important aspect in the clinical practice is represented by the continuous monitoring of the nociception-antinociception balance. The aim of these parameters is to come as an aid to the clinician in deciding the adequate analgesia dosage for each patient. While monitoring the degree of hypnosis is achieved through the direct evaluation of the EEG signals, the nociception-antinociception balance can be monitored indirectly [9,12], by evaluating certain variables such as the vasomotor reflex, pupillary size, the H reflex, and the hemodynamic response [57-58] (Figure 1).



**Figure 1.** Technologies/parameters used for monitoring the nociception-antinociception balance [59–66]. ANI – analgesia nociception index; qNOX index - index of nociception; qCON index - index of consciousness; SPI – Surgical Plethysmographic Index; NOL index - Nociception Level Index

One of the most widely studied technologies is the analysis of hemodynamic changes and includes the evaluation of the normalized heart beat intervals (HBIs) and of the amplitude of the plethysmographic waves, both correlating with sympathetic and parasympathetic tones. A higher sympathetic tone correlates with the intensity of the surgical stimuli and results in a suppressed plethysmographic amplitude (PPGA). For the correct calculation of the surgical plethysmographic index (SPI), after normalizing these parameters by transforming the histogram, the SPI formula will be used where  $SPI = 100 - (0.67 \times PPGA_{norm} + 0.33 \times HBI_{norm})$ . The SPI value can be influenced by certain factors, such as cardiac pacemakers, cardiac arrhythmias, antiarrhythmic medication, beta-1 adrenergic antagonists, or alpha2-adrenergic agonists. Bonhomme et al., have evaluated the Surgical Pleth Index (SPI, GE Healthcare, Helsinki, Finland) trend and have made correlations with variability in mean arterial pressure and heart rate. Following this study they have shown that there is a strong correlation between all these variables and therefore SPI values depend on the doses of opioid medication administered during the anesthesia [58]. Bergmann et al, have carried out a randomized study that included 170 patients receiving general anesthesia with propofol and remifentanyl. The patients were divided in two study groups, one study group received opioids based on SPI values, while the other group received the doses of opioids based on standard monitoring parameters, both clinical and hemodynamic monitoring. Statistically significant differences were shown in both propofol ( $p < 0.05$ ,  $6.0 \pm 2.1$  vs.  $7.5 \pm 2.2$  mg/kg/h) remifentanyl ( $p < 0.05$ ,  $0.06 \pm 0.04$  vs.  $0.08 \pm 0.05$   $\mu\text{g}/\text{kg}/\text{min}$ ) consumption. The impact on post-anesthesia recovery time was evaluated by the time needed to open the eyes and time to extubation. The results presented statistical significance for both the evaluated features, extubation time ( $p < 0.05$ ,  $1.2 \pm 4.4$  min vs.  $4.4 \pm 4.5$  min) and eye-opening time ( $p < 0.05$ ,  $-0.08 \pm 4.4$  min vs.  $3.5 \pm 4.3$  min). The conclusion was that dose reduction and shorter recovery times can be achieved by adapting general anaesthesia based on the SPI [10]. Huiku et al confirmed in a similar study that SPI monitoring has a beneficial impact on anesthetic drugs used doses, increasing patient safety and the quality of the medical services [67].

Another parameter used for the evaluation of the nociception-antinociception balance is the Analgesia Nociception Index (ANI) [68]. The technology is based on the assessment in heart rate variability. To be used in the clinical setting ANI values lie between 0 and 100. In this case 0 represents a very low degree of parasympathetic modulation, and 100 represent a very high degree of parasympathetic activity. From a clinical point of view ANI=0 represents a high level of stress, while ANI=100 represents low stress levels.

Dostalova et al., have carried out a study in which they compared the impact the two monitoring techniques have on general anesthesia. They had three study groups: the group where doses of opioids were titrated based on ANI values, the SPI group and the control group. They have shown statistically significant differences regarding the decrease in opioid consumption and shorter recovery times after anesthesia [68]. Table 2 summarizes a series of studies regarding the impact of monitoring techniques on the nociception-antinociception balance and on the clinical outcome of patients.

**Table 2.** The impact of nociception-antinociception monitoring techniques on anesthetic drugs consumption and on recovery time

Author	Technique/parameter	Type of anesthesia Type of intervention	Observations	Reference
Funcke et al.	SPI & Pupillary Pain Index (PPI) & Nociception Level (NOL)	48 patients radical retropubic prostatectomy	SPI: ↓ hormonal response to stress PPI: ↓ sufentanil consumption, ↑ hormonal response to stress; No effect on: postoperative recovery time. ↓ propofol consumption; ↓ remifentanil consumption;	[69]
Bergmann et al.	Surgical Pleth Index (SPI)	170 patients orthopedic surgery	↓ postoperative recovery time. ↓ PONV; ↓ postoperative pain;	[10]
Jain et al.	Surgical Pleth Index (SPI)	140 patients Laparoscopic cholecystectomy;	↑ fentanyl consumption; No impact on hemodynamic stability ↓ oxycodone consumption;	[70]
Won et al.	Surgical Pleth Index (SPI)	45 patient; Elective thyroidectomy	↓ postoperative recovery time; ↓ extubation time; ↓ remifentanil consumption;	[71]
Chen et al.	Surgical Stress Index (SSI) – former Surgical Pleth Index (SPI)	80 patients Elective ENT surgical interventions	↓ postoperative adverse effects ↑ hemodynamic stability	[72]
Theerth et al.	Analgesia Nociception Index (ANI)	60 patients Oncological surgery	↓ fentanyl consumption; No impact on postoperative pain; ↓ opioid consumption;	[73]
Soral et al.	Analgesia Nociception Index (ANI)	102 patients Procedural sedation	No impact of propofol and ketamine consumption	[74]



Gall et al.	Analgesia Nociception Index (ANI)	60 patients Bariatric surgery	↓ sufentanyl consumption; No impact on PONV and postoperative pain;	[75]
-------------	---	----------------------------------	--	------

Numerous studies have shown that opioid overdose during anesthesia is responsible for a series of adverse effects, such as increased recovery times and opioid induced hyperalgesia, and they also lead to hypotension, having a major impact on perioperative hemodynamic stability [66-71]. Won et al., have reported that using SPI monitoring during general anesthesia reduces opioid consumption, improves hemodynamic stability, and reduced postoperative recovery times [71]. A similar study was carried out by Jain et al., and has reported a statistically significant decrease in the number of hemodynamic adverse events when SPI was used for the titration of opioid medication ( $p < 0.05$ ) [70].

Another system used for monitoring the nociception-antinociception balance is the index of nociception (qNOX) (qCON 2000 Monitor, Quantum Medical, Fresenius Kabi, Mataro, Spain). This parameter is based on the evaluation of EEG and EMG patterns, with values between 0 and 99. Jensen et al., carried out a study on 60 patients undergoing general anesthesia with propofol and remifentanyl and have shown a series of statistically significant correlations concluding that qNOX can detect fine changes in the nociception-antinociception balance [76]. The Nociception Level Index (NOL index, Medasense, Ramat Gau, Israel) is another widely used technology for titrating analgesic drugs during general anesthesia. It analyses the photoplethysmographic wave, temperature, skin galvanic conductance response, and accelerometry [63].

#### 4. The impact of multimodal monitoring on the hemodynamic status

During general anesthesia maintaining adequate tissue perfusion represents one of the most important goals in the perioperative management of the patient. Hypotension frequently occurs especially after the induction of anesthesia or, better said, during the time between the moment of induction and the start of surgery. Reich et al., have reported a decrease in mean arterial pressure (MAP) of over 40% ( $\text{MAP} < 70 \text{ mmHg}$  or  $\text{MAP} < 60 \text{ mmHg}$ ) in the first 10 minutes after induction ( $p < 0.001$ ) [77]. Moreover, this study ( $n = 2406$  patients) reported an increase in the time spent in the recovery room (13.3%,  $p < 0.05$ ) and in postoperative mortality rates (8.6%,  $p < 0.02$ ) in patients that presented perioperative hypotension. Another interesting phenomenon presented by the group was that post-induction hypotension was more frequent in the 5-10 minutes interval, in comparison to the 0-5 minutes interval after induction of general anesthesia [77]. A similar study carried out by Hug et al., reported that over 15% of the patients that present a decrease in systolic blood pressure (SBP) under 90 mmHg after induction with propofol in the first 10 minutes after administration [78]. Studies have shown that induction with sevoflurane maintains hemodynamic stability and decreases the risk of hypotension in comparison to induction with propofol, by this technique is not as well tolerated by the patients. Thwaites et al., made a study on the satisfaction of patients regarding the induction technique used: sevoflurane (inhalational induction, 8%) vs. propofol (i.v. induction). Over 14% of the patients consider inhalational induction unpleasant in comparison to 10% in the case of propofol. Furthermore, over 24% of the patients would not choose sevoflurane induction the second time [79].

Cerebral ischemia is one of the main causes for cognitive impairment, with a very high global degree of mortality, while motor and cognitive dysfunctions seriously affect the quality of life of these patients. Cerebral reperfusion after an ischemic episode can induce organ damage such as neurovascular injury, neuronal death, cerebral edema, and neuro-hemorrhagic changes. The most common cellular mechanisms involved are represented by apoptosis, inflammation, and excessive production of free radicals [80].

The impact of hypotension during general anesthesia on the postoperative outcome and on the development of postoperative adverse events has been widely studied. Intraoperative hypotension (IHO) is a common effect of general anesthesia and has been associated with an increased incidence of one year mortality after surgery [81,82].

The most important predictors for perioperative morbidity and mortality are the associated comorbidities, the determinants of the surgical procedure, and the specific aspects of perioperative

management and of general anesthesia. Apart from monitoring the hemodynamic parameters, quantification of the degree of hypnosis "depth of anesthesia" represents one of the most important parameters in modern general anesthesia. At the time monitoring the degree of hypnosis is possible by using techniques based on the analysis of electroencephalography signals (EEG) [83].

Monk et al., studied the one year prognosis of patients that have undergone non-cardiac surgery under general anesthesia. The research group has carried out complex statistical analysis in order to determine if death at one year after can be associated with significant clinical features of the patient or with the management of general anesthesia. In order to control the degree of hypnosis they used the Bispectral Index<sup>®</sup> (BIS<sup>®</sup>), with the same type of electrodes for all patients included in the study (A1050BIS Monitor, BIS sensors, Aspect Medical Systems, Newton, MA).


Global mortality at a year was 5.5% (n=1604) and 10.3% for patients aged over 65 (n=243). Regarding variables that correlated with mortality Monk et al., have reported 3 statistically significant segments: 1. patient comorbidities (relative risk 6.116, p<0.05), 2. general anesthesia overdose/ anesthesia to deep, BIS<45 (relative risk 1.244/h, p<0.05), 3. systolic hypotension during surgery (relative risk 1.036/min, p<0.05) [83]. They concluded that prolonged intraoperative hypotension can be associated with an increased incidence in mortality at one year [83]. Although there are numerous studies that focus on perioperative hypotension, at the time there is no clear definition for IHO [84]. Most of the studies address statistical associations and correlations between different numerical intervals, and correlations with the clinical changes. Sun et al., have carried out a study on the impact of IHO on acute kidney injury (AKI). Furthermore, the research group investigated the implications IHO time have on the incidence of AKI. They have correlated the AKI incidence with different IHO intervals as follows: MAP < 55 mmHg, MAP < 60 mmHg and MAP < 65 mmHg [85]. This was a retrospective study that included 5127 patients between 2009 and 2012. The results have shown an AKI incidence of 6.3% (324 patients) for MAP<60 mmHg and an IHO time between 11-20 minutes, and MAP<55 for an IHO time>10 minutes. Sun et al., reported a strong statistical correlation between sustained episodes of IHO with a MAP<50 mmHg and MAP<60 mmHg and AKI incidence. For the evaluation of AKI they have considered a 50% increase in creatinine levels or 0.3 mg/dl in the first 2 days after surgery. A similar study was developed by Walsh et al., regarding the implications of IHO on the incidence of AKI and myocardial injury. They have evaluated 33330 patients that had undergone non-cardiac surgery making statistical correlations between the incidence of AKI and myocardial injury in patients that had presented with IHO with a MAP<55mmHg and MAP<75 mmHg. Following statistical analysis they have identified 2478 patients that had developed AKI (7.4%) and 770 (2.3%) with myocardial injuries. For both groups MAP was under 55 mmHg. Interestingly, the risk for developing renal and myocardial lesions is increased even for short times of IHO [86]. In a similar context, a meta-analysis carried out by Wesselink et al., reported ischemic organ damage when MAP<80mmHg for longer than 10 minutes. This research group showed an increase in risk with any decrease in blood pressure [84].

## 5. The impact of general anaesthesia multimodal monitoring on inflammation/redox

Another important aspect that also has an impact on the clinical outcome of surgical patients is represented by the inflammatory status and by the oxidoreduction response (REDOX) [87–91]. The excessive production of free nitrogen and oxygen radicals has a direct involvement in the augmentation of the pro-inflammatory status. Under physiological conditions the balance between the production of free radicals and that of endogenous antioxidant substances maintains the oxidoreduction equilibrium and the body does not suffer. Under surgical stress, in the case of ischemia-reperfusion syndrome or hypotension an excessive amount of free radicals will be produced, as well as pro-inflammatory mediators. All these factors will also decrease the production capacity for antioxidant molecules [92].

Particularly in the case of patients under GA or in mechanically ventilated patients, oxygen plays an essential role in the therapeutic management. In the case of general anesthesia increased oxygen inspiratory fractions (FiO<sub>2</sub>) are administered before endotracheal intubation and after extubation in order to maintain an adequate oxygen plasma concentration. Under physiological conditions P<sub>a</sub>O<sub>2</sub>=

80-100 mmHg. When  $P_{aO_2}$  exceeds 100 mmHg the patient is characterized by hyperoxia, the most important systemic effect being the increased and accelerated production of reactive oxygen species (ROS) and the development of oxidative stress (OS) [93–99]. The most important mechanisms through which OS is augmented in the case of general anesthesia are represented by the increase in molecular oxygen offerings at the mitochondria, the interaction with reactive nitrogen species (RNS), and lipid peroxidation with destruction of cellular membranes [25,100–103]. (Figure 2)



**Figure 2.** Schematic representation of the oxidative response in patients under general anesthesia.

Nunes et al. have studied the implications of general anesthesia (GA) on the redox profile of surgical patients that underwent intravenous GA, as well as the implications of multimodal monitoring based on Entropy on the oxidoreduction activity. 20 patients were included in the study, divided into two study groups: the first in which Entropy values were maintained in the 45-59 interval and the second in which Entropy was maintained in the 30-44 interval in order to evaluate the impact of anesthetic overdosage on the redox balance. The patients were evaluated at different moments in time: M1 – right after the administration of anesthetic drugs, M2 – after endotracheal intubation, M3 – 5 minutes after endotracheal intubation, M4 – immediately after surgical pneumoperitoneum, M5 – 1 minute after pneumoperitoneum, M6 – 1 hour after the end of surgery. The researchers determined the plasma concentrations for Glutathione and TBARS (tiobarbituric acid reactive species). Following the analysis they have identified significant increases in the Glutathione and TBARS concentrations at M5 in both groups. There were statistically significant differences between the two study groups with higher values of both Glutathione and TBARS in the group where Entropy was maintained between 30 and 44 ( $p < 0.05$ ). In regard to the anesthetic management, recovery times were significantly shorter for the group where Entropy levels were kept between 45 and 59 ( $7.70 \pm 1.24$  min vs.  $10.20 \pm 0.90$  min,  $P < 0.05$ ). The increase in redox imbalance markers for the patients that received a deeper hypnosis (Entropy 20-44) reveals an increase in anaerobic metabolism, possibly because of an accentuated suppression of the autonomic nervous system [92].

Ferrari et al., carried out a study regarding the genotoxicity of sevoflurane on the DNA structure in isolated lymphocytes in 20 patients undergoing orthopedic surgery under GA. They have shown important changes in DNA structure and in redox activity that correlated statistically with the sevoflurane concentration [104]. Compared to the exposure to propofol, the group that was exposed

to sevoflurane presented a marked increase in the expression of tumor necrosis factor alpha (TNF-alpha) and a decrease for interleukin 10 (IL-10) [104-105].

## 6. The impact on drug consumption and recovery

Gan et al. led an important study regarding the implications of monitoring the degree of hypnosis. They included 302 patients divided in two groups. In the study group GA was guided based on BIS monitoring, while in the control group anesthesia was guided with basic monitoring. BIS values were measured in both groups [106]. In the study group the dosage of anesthetic agents were optimized in order to achieve a mean BIS value between 40 and 60 based on current guidelines and recommendations. Interestingly enough the BIS values in the control group were under 40, indicating a tendency to overdosing the anesthetic agents. The total propofol consumption was lower in the study group compared to the control. Another important variable is the time to extubation that was 7.27 minutes shorter (95% CI 6.23-8.28 min) in the study group compared to 11.22 min in the control group (95% CI, 8.51 – 13.60 min). Song et al. have designed a similar study that also showed a decrease in extubation times in patients that received general anesthesia modulated based on BIS, with a reduction from  $6.5 \pm 4.3$  min to  $3.6 \pm 1.5$  min (> 40%) for Desflurane, and from  $7.7 \pm 3.5$  min to  $5.5 \pm 2.2$  min for sevoflurane [107].

Vakkuri et al., have carried out a multicenter study on the impact the monitoring of degree of hypnosis through Entropy (GE Healthcare, Helsinki, Finland) has on anesthetic drug consumption and on postoperative recovery time. In the final analysis of the study they included 308 patients, divided homogeneously in two groups: the control group and the study group where GA was modulated based on Entropy. For propofol consumption there were statistically significant differences between the two study groups, the median for the control group being 0.11 (0.03, 0.21) mg/kg/min vs. 0.10 (0.04, 0.23) mg/kg/min for the group where Entropy was used.

The analysis of the implications multimodal monitoring has on the postoperative recovery showed a decrease in the time to spontaneous breathing in the study group 4.74 (0.00, 18.0) vs the median in the control group 7.07 (1.00-28.5). The time to extubation was also decrease by using Entropy from 9.16 (1.67, 32.3) minutes to 5.80 (3.00, 27.3) with  $p < 0.05$ . The patients in the target group opened their eyes to verbal command faster than the control group 6.08 (0.15, 37.5) minutes vs 10.8 (2.23, 43.2) minutes ( $p < 0.05$ ) and they were transferred in the Post-Anesthesia Care Unit (PACU) faster, at 10.3 (1.17, 48.7,  $p < 0.05$ ) minutes vs. 13.0 (5.0, 49.8). Mean State Entropy (SE) during general anesthesia was 50 (34-78), while the mean Response Entropy (RE) was 52 (35-84). [46]. A similar study was developed by El Hor et al., reporting an increase in sevoflurane consumption in the case of patients that could not benefit from advanced monitoring of the degree of hypnosis vs. patients for which Entropy monitoring was applied ( $5.2 \pm 1.4$  mL/h vs.  $3.8 \pm 1.5$  mL/h,  $P < 0.05$ ) [108]. Regarding hemodynamic stability the researchers found statistically significant differences between the groups: 10 hypertension episodes were reported in the control group vs. 7 hypertension episode in the target group, while for hypotension the ratio was 3 (control group) vs. 0 in the target group ( $p < 0.05$ ). Tachycardia episodes were reported as 5 (control group) vs. 8 (target group), while bradycardia 1 (control group) vs. 0 (study group)

Wu et al have analyzed the impact of multiparametric monitoring based on Entropy (GE Datex-Ohmeda S/5) on the recovery time and anesthetic drugs consumption in patients undergoing orthopedic surgery. This research group has included 68 patients in their analysis, divided into 2 groups: target group with Entropy monitoring and the control group with classical anesthesia monitoring. Sevoflurane consumption was significantly lower in the target group ( $P < 0.05$ ),  $27.79 \pm 7.4$  mL / patient vs.  $31.42 \pm 6.9$  mL / patient. Statistically significant differences have been reported also for hemodynamic stability, as the target group presented less hypertensive episodes compared to the control,  $0.94 \pm 1.15$  vs  $1.48 \pm 1.41$ ,  $P < 0.05$ . Following this study the research group concluded that using Entropy-based multimodal monitoring significantly reduces both sevoflurane consumption and the consumption of antihypertensive agents [45].

The impact of multiparametric monitoring on the anesthetic drugs consumption was proven in another study by Tewari et al., in patients undergoing gynecological and obstetrical surgery. They



have analyzed 120 patients that were divided into two study groups based on monitoring technique: Entropy group vs. classical monitoring of general anesthesia. They have shown that Entropy monitoring leads to a reduction of propofol doses (6.7% reduction,  $p=0.01$ ), but also that the Fentanyl doses were 10.9% larger in this group ( $p=0.07$ ). They did not find any statistically significant differences for recovery time and discharge time from PACU [109]. Refaat et al. in their study on the impact of Entropy on sevoflurane consumption in major hepatic surgery showed a marked decrease in the doses [110].

## 7. Conclusions

General anesthesia techniques are much more advanced nowadays compared to latter decades, in accordance with the surgical needs and with the needs of the general population. Medical services tend to become more and more complex, managing to solve a wide range of pathologies in all surgical fields. In order to increase both patient safety and medical act quality, as well as to decrease waiting times and to be able to answer the needs of an increasing number of patients, the endowment with modern multiparametric monitoring techniques for general anesthesia is necessary. In conclusion, we can state that by using monitoring techniques for the degree of hypnosis, for the nociception-antinociception balance, and for the hemodynamic status, patient safety is markedly increased. Furthermore, by reducing postoperative recovery times and by reducing anesthetic drugs doses, one can highlight the positive impact, both short- and long-term, that multiparametric monitoring has from an economic viewpoint.

**Author Contributions:** Conceptualization, A.F.R. and D.S.; methodology, D.N.G., M.N., C.M.D. and L.B.P. software, O.H.B; validation, D.S. and M.P.; formal analysis, A.M.; investigation, A.F.R.; resources, C.R.B., A.M.; data curation, L.M.B.; writing—original draft preparation, A.F.R. and D.T.; writing—review and editing, A.R., L.M.B. and C.M.D.; visualization, D.S.; supervision, D.S.; project administration, O.H.B. and M.P.; funding acquisition, D.S. and A.F.R.. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Dabouz, R.; Davidson, J.; Deloste, J.; Rahali, N.; Zadam, A.; Boselli, E.; Bouvet, L.; Be, G. Prediction of immediate postoperative pain using the analgesia / nociception index : a prospective observational study. *2017*, *112*, 715–721, doi:10.1093/bja/aet407.
2. Guerrero, J.L.; Matute, E.; Alsina, E.; Del Blanco, B.; Gilsanz, F. Response entropy changes after noxious stimulus. *J. Clin. Monit. Comput.* **2012**, *26*, 171–175, doi:10.1007/s10877-012-9351-3.
3. Lehmann, A.; Schmidt, M.; Zeitler, C.; Kiessling, A.-H.; Isgro, F.; Boldt, J. Bispectral index and electroencephalographic entropy in patients undergoing aortocoronary bypass grafting. *Eur. J. Anaesthesiol.* **2007**, *24*, 751–760, doi:10.1017/S0265021506002249.
4. Health Quality Ontario Bispectral index monitor: an evidence-based analysis. *Ont. Health Technol. Assess. Ser.* **2004**, *4*, 1–70.
5. Hebb, M.O.; McArthur, D.L.; Alger, J.; Etchepare, M.; Glenn, T.C.; Bergsneider, M.; Martin, N.; Vespa, P.M. Impaired percent alpha variability on continuous electroencephalography is associated with thalamic injury and predicts poor long-term outcome after human traumatic brain injury. *J Neurotrauma* **2007**, *24*.
6. Petras, M.; Tatarikova, Z.; Kovalska, M.; Mokra, D.; Dobrota, D.; Lehotsky, J.; Drgova, A. Hyperhomocysteinemia as a risk factor for the neuronal system disorders. *J. Physiol. Pharmacol.* **2014**, *65*, 15–23, doi:10.1056/NEJMra1313875.
7. Sleight, J.; Harvey, M.; Voss, L.; Denny, B. Ketamine - more mechanisms of action than just NMDA blockade. *Trends Anaesth. Crit. Care* **2014**, *4*, 76–81, doi:10.1016/j.tacc.2014.03.002.
8. Abushik, P.A.; Niittykoski, M.; Giniatullina, R.; Shakirzyanova, A.; Bart, G.; Fayuk, D.; Sibarov, D.A.; Antonov, S.M.; Giniatullin, R. The role of NMDA and mGluR5 receptors in calcium mobilization and neurotoxicity of homocysteine in trigeminal and cortical neurons and glial cells. *J. Neurochem.* **2014**, *129*, 264–274, doi:10.1111/jnc.12615.



9. Dinu, A.R.; Rogobete, A.F.; Popovici, S.E.; Bedreag, O.H.; Papurica, M.; Dumbuleu, C.M.; Velovan, R.R.; Toma, D.; Georgescu, C.M.; Trache, L.I.; et al. Impact of general anesthesia guided by state entropy (SE) and response entropy (RE) on perioperative stability in elective laparoscopic cholecystectomy patients - A prospective observational randomized monocentric study. *Entropy* **2020**, *22*, doi:10.3390/e22030356.
10. Bergmann, I.; Göhner, A.; Crozier, T.A.; Hesjedal, B.; Wiese, C.H.; Popov, A.F.; Bauer, M.; Hinz, J.M. Surgical pleth index-guided remifentanil administration reduces remifentanil and propofol consumption and shortens recovery times in outpatient anaesthesia. *Br. J. Anaesth.* **2013**, *110*, 622–628, doi:10.1093/bja/aes426.
11. Aho, A.J.; Kamata, K.; Jäntti, V.; Kulkas, A.; Hagihira, S.; Huhtala, H.; Yli-Hankala, A. Comparison of Bispectral Index and Entropy values with electroencephalogram during surgical anaesthesia with sevoflurane. *Br. J. Anaesth.* **2015**, *115*, 258–266, doi:10.1093/bja/aev206.
12. Freye, E.; Levy, J. V. Cerebral monitoring in the operating room and the intensive care unit: An introductory for the clinician and a guide for the novice wanting to open a window to the brain; 2005; Vol. 19; ISBN 1087700507.
13. Klockars, J.G.M.; Hiller, A.; Münte, S.; Van Gils, M.J.; Taivainen, T. Spectral entropy as a measure of hypnosis and hypnotic drug effect of total intravenous anesthesia in children during slow induction and maintenance. *Anesthesiology* **2012**, *116*, 340–351, doi:10.1097/ALN.0b013e3182410b5e.
14. Bibian, S.; Dumont, G.A.; Zikov, T. Dynamic behavior of BIS, M-entropy and neuroSENSE brain function monitors. *J. Clin. Monit. Comput.* **2011**, *25*, 81–87, doi:10.1007/s10877-010-9266-9.
15. Shortal, B.P.; Hickman, L.B.; Mak-McCully, R.A.; Wang, W.; Brennan, C.; Ung, H.; Litt, B.; Tarnal, V.; Janke, E.; Picton, P.; et al. Duration of EEG suppression does not predict recovery time or degree of cognitive impairment after general anaesthesia in human volunteers. *Br. J. Anaesth.* **2019**, *123*, 206–218, doi:10.1016/j.bja.2019.03.046.
16. Schultz, A.; Siedenberg, M.; Grouven, U.; Kneif, T.; Schultz, B. Comparison of narcotrend index, bispectral index, spectral and entropy parameters during induction of propofol-remifentanil anaesthesia. *J. Clin. Monit. Comput.* **2008**, *22*, 103–111, doi:10.1007/s10877-008-9111-6.
17. Singh, S.; Bansal, S.; Kumar, G.; Gupta, I.; Thakur, J.R. Entropy as an indicator to measure depth of anaesthesia for Laryngeal Mask Airway (LMA) insertion during sevoflurane and propofol anaesthesia. *J. Clin. Diagnostic Res.* **2017**, *11*, UC01–UC03, doi:10.7860/JCDR/2017/27316.10177.
18. Iannuzzi, M.; Iannuzzi, E.; Rossi, F.; Berrino, L.; Chiefari, M. Relationship between Bispectral Index , electroencephalographic state entropy and effect-site EC 50 for propofol at different clinical endpoints. **2005**, *94*, 492–495, doi:10.1093/bja/aei075.
19. Wildemeersch, D.; Peeters, N.; Saldien, V.; Vercauteren, M.; Hans, G. Pain assessment by pupil dilation reflex in response to noxious stimulation in anaesthetized adults. *Acta Anaesthesiol. Scand.* **2018**, *62*, 1050–1056, doi:10.1111/aas.13129.
20. Kumar, A.; Anand, S. A depth of anaesthesia index from linear regression of EEG parameters. *J. Clin. Monit. Comput.* **2006**, *20*, 67–73, doi:10.1007/s10877-005-9004-x.
21. Jagadeesan, N.; Wolfson, M.; Chen, Y.; Willingham, M.; Avidan, M.S. Brain monitoring during general anesthesia. *Trends Anaesth. Crit. Care* **2013**, *3*, 13–18, doi:10.1016/j.tacc.2012.12.003.
22. Johansen, J.W.; Sebel, P.S. Development and clinical application electroencephalographic bispectrum monitoring. *Anesthesiology* **2000**, *93*, 1336–1344, doi:10.1097/0000542-200011000-00029.
23. Rogobete, A.F.; Bedreag, O.H.; Sandesc, D. Entropy-Guided Depth of Anesthesia in Critically Ill Polytrauma Patients. *J. Interdiscip. Med.* **2017**, *2*, 7–8, doi:10.1515/jim-2017-0027.
24. Klockars, J.G.M.; Hiller, A.; Ranta, S.; Talja, P.; Van Gils, M.J.; Taivainen, T. Spectral entropy as a measure of hypnosis in children. *Anesthesiology* **2006**, *104*, 708–717, doi:10.1097/0000542-200604000-00015.
25. Cota, A.; Grințescu, I.M. Entropy — The Need of an Ally for Depth of Anesthesia Monitoring in Emergency Surgery. **2019**, 13–15, doi:10.35995/ceacr1010001.
26. Lonjaret, L.; Lairez, O.; Minville, V.; Geeraerts, T. Optimal perioperative management of arterial blood pressure. *Integr. Blood Press. Control* **2014**, *7*, 49–59, doi:10.2147/IBPC.S45292.
27. Südfeld, S.; Brechnitz, S.; Wagner, J.Y.; Reese, P.C.; Pinnschmidt, H.O.; Reuter, D.A.; Saugel, B. Post-induction hypotension and early intraoperative hypotension associated with general anaesthesia. *Br. J. Anaesth.* **2017**, *119*, 57–64, doi:10.1093/bja/aex127.
28. Hoppe, P.; Kouz, K.; Saugel, B. Perioperative hypotension: clinical impact, diagnosis, and therapeutic approaches. *J. Emerg. Crit. Care Med.* **2020**, *4*, 8–8, doi:10.21037/jeccm.2019.10.12.

29. Barnard, J.P.; Bennett, C.; Voss, L.J.; Sleight, J.W.; Hospital, W.; Zealand, N. NEUROSCIENCES AND NEUROANAESTHESIA Can anaesthetists be taught to interpret the effects of general anaesthesia on the electroencephalogram? Comparison of performance with the BIS and spectral entropy. *2007*, *99*, 532–537, doi:10.1093/bja/aem198.
30. Chen, S.J.; Peng, C.J.; Chen, Y.C.; Hwang, Y.R.; Lai, Y.S.; Fan, S.Z.; Jen, K.K. Comparison of FFT and marginal spectra of EEG using empirical mode decomposition to monitor anesthesia. *Comput. Methods Programs Biomed.* **2016**, *137*, 77–85, doi:10.1016/j.cmpb.2016.08.024.
31. Kreuer, S.; Wilhelm, W. The Narcotrend monitor. *Best Pract. Res. Clin. Anaesthesiol.* **2006**, *20*, 111–119, doi:10.1016/j.bpa.2005.08.010.
32. Kreuer, S.; Bruhn, J.; Larsen, R.; Bialas, P.; Wilhelm, W. Comparability of Narcotrend TM index and bispectral index during propofol anaesthesia. *2004*, *93*, 235–240, doi:10.1093/bja/ae182.
33. Kreuer, S.; Biedler, A.; Larsen, R.; Schoth, S.; Altmann, S.; Wilhelm, W. The Narcotrend--a new EEG monitor designed to measure the depth of anaesthesia. A comparison with bispectral index monitoring during propofol-remifentanil-anaesthesia. *Anaesthesist* **2001**, *50*, 921–925, doi:10.1007/s00101-001-0242-0.
34. Dennhardt, N.; Arndt, S.; Beck, C.; Boethig, D.; Heiderich, S.; Schultz, B.; Weber, F.; Sümpelmann, R. Effect of age on Narcotrend Index monitoring during sevoflurane anesthesia in children below 2 years of age. *Paediatr. Anaesth.* **2018**, *28*, 112–119, doi:10.1111/pan.13306.
35. Dong, X.; Suo, P.; Yuan, X.; Yao, X. Use of auditory evoked potentials for intra-operative awareness in anesthesia: a consciousness-based conceptual model. *Cell Biochem. Biophys.* **2015**, *71*, 441–447, doi:10.1007/s12013-014-0221-0.
36. Mantzaridis, H.; Kenny, G.N. Auditory evoked potential index: a quantitative measure of changes in auditory evoked potentials during general anaesthesia. *Anaesthesia* **1997**, *52*, 1030–1036, doi:10.1111/j.1365-2044.1997.185-az0327.x.
37. Khan, J.; Mariappan, R.; Venkatraghavan, L. Entropy as an indicator of cerebral perfusion in patients with increased intracranial pressure. *J. Anaesthesiol. Clin. Pharmacol.* **2014**, *30*, 409, doi:10.4103/0970-9185.137280.
38. Illman, H.; Antila, H.; Olkkola, K.T. Reversal of neuromuscular blockade by sugammadex does not affect EEG derived indices of depth of anesthesia. *J. Clin. Monit. Comput.* **2010**, *24*, 371–376, doi:10.1007/s10877-010-9257-x.
39. Schartner, M.M.; Carhart-Harris, R.L.; Barrett, A.B.; Seth, A.K.; Muthukumaraswamy, S.D. Increased spontaneous MEG signal diversity for psychoactive doses of ketamine, LSD and psilocybin. *Sci. Rep.* **2017**, *7*, 1–12, doi:10.1038/srep46421.
40. Abdelmageed, W.M.; Al Taher, W.M. Preoperative paracetamol infusion reduces sevoflurane consumption during thyroidectomy under general anesthesia with spectral entropy monitoring. *Egypt. J. Anaesth.* **2014**, *30*, 115–122, doi:10.1016/j.egja.2013.12.003.
41. Tiefenthaler, W.; Colvin, J.; Steger, B.; Pfeiffer, K.P.; Moser, P.L.; Walde, J.; Lorenz, I.H.; Kolbitsch, C. How bispectral index compares to spectral entropy of the EEG and A-line ARX index in the same patient. *Open Med.* **2018**, *13*, 583–596, doi:10.1515/med-2018-0087.
42. Bergmann, I.; Göhner, A.; Crozier, T.A.; Hesjedal, B.; Wiese, C.H.; Popov, A.F.; Bauer, M.; Hinz, J.M. Surgical pleth index-guided remifentanil administration reduces remifentanil and propofol consumption and shortens recovery times in outpatient anaesthesia. *Br. J. Anaesth.* **2013**, *110*, 622–628, doi:10.1093/bja/aes426.
43. Maksimow, A.; Särkelä, M.; Långsjö, J.W.; Salmi, E.; Kaisti, K.K.; Yli-Hankala, A.; Hinkka-Yli-Salomäki, S.; Scheinin, H.; Jääskeläinen, S.K. Increase in high frequency EEG activity explains the poor performance of EEG spectral entropy monitor during S-ketamine anesthesia. *Clin. Neurophysiol.* **2006**, *117*, 1660–1668, doi:10.1016/j.clinph.2006.05.011.
44. Davidson, A.J.; Huang, G.H.; Rebmann, C.S.; Ellery, C. Performance of entropy and Bispectral Index as measures of anaesthesia effect in children of different ages. *Br. J. Anaesth.* **2005**, *95*, 674–679, doi:10.1093/bja/aei247.
45. Kim, Y.H.; Choi, W. Effect of preoperative anxiety on spectral entropy during induction with propofol. **2013**, *65*, 108–113.
46. Wu, S.; Wang, P.; Liao, W.; Shih, T.; Chang, K.; Lin, K.; Chou, A. Use of Spectral Entropy Monitoring in Reducing the Quantity of Sevoflurane as Sole Inhalational Anesthetic and in Decreasing the Need for

- Antihypertensive Drugs in Total Knee Replacement Surgery. *Acta Anaesthesiol. Taiwanica* **2008**, *46*, 106–111, doi:10.1016/S1875-4597(08)60003-X.
47. Vakkuri, A.; Ph, D.; Yli-hankala, A.; Ph, D.; Sandin, R.; Ph, D.; Mustola, S.; Ph, D. Spectral Entropy Monitoring Is Associated with Reduced Propofol Use and Faster Emergence in Propofol – Nitrous Oxide – Alfentanil Anesthesia. **2005**, *0*, 274–279.
  48. Talawar, P.; Chhabra, A.; Trikha, A.; Arora, M.K.; Chandralekha Entropy monitoring decreases isoflurane concentration and recovery time in pediatric day care surgery—a randomized controlled trial. *Paediatr. Anaesth.* **2010**, *20*, 1105–1110, doi:10.1111/j.1460-9592.2010.03441.x.
  49. Elgebaly, A.S.; El Mourad, M.B.; Fathy, S.M. The role of entropy monitoring in reducing propofol requirements during open heart surgeries. A prospective randomized study. *Ann. Card. Anaesth.* **2020**, *23*, 272–276, doi:10.4103/aca.ACA\_184\_18.
  50. Bhardwaj, N.; Yaddanapudi, S. A randomized trial of propofol consumption and recovery profile with BIS-guided anesthesia compared to standard practice in children. *Paediatr. Anaesth.* **2010**, *20*, 160–167, doi:10.1111/j.1460-9592.2009.03240.x.
  51. Aime, I.; Taylor, G. Does Monitoring Bispectral Index or Spectral Entropy Reduce Sevoflurane Use ? **2006**, *103*, doi:10.1213/01.ane.0000246838.93153.23.
  52. Liao, W.-W.; Wang, J.-J.; Wu, G.-J.; Kuo, C.-D. The effect of cerebral monitoring on recovery after sevoflurane anesthesia in ambulatory setting in children: a comparison among bispectral index, A-line autoregressive index, and standard practice. *J. Chin. Med. Assoc.* **2011**, *74*, 28–36, doi:10.1016/j.jcma.2011.01.004.
  53. Weber, F.; Geerts, N.J.E.; Roeleveld, H.G.; Warmenhoven, A.T.; Liebrand, C.A. The predictive value of the heart rate variability-derived Analgesia Nociception Index in children anaesthetized with sevoflurane: An observational pilot study. *Eur. J. Pain (United Kingdom)* **2018**, *22*, 1597–1605, doi:10.1002/ejp.1242.
  54. Lai, R.-C.; Lu, Y.-L.; Huang, W.; Xu, M.-X.; Lai, J.-L.; Xie, J.-D.; Wang, X.-D. [Application of a narcotrend-assisted anesthesia in-depth monitor in the microwave coagulation for liver cancer during total intravenous anesthesia with propofol and fentanyl]. *Chin. J. Cancer* **2010**, *29*, 117–120, doi:10.5732/cjc.009.10244.
  55. Rundshagen, I.; Hardt, T.; Cortina, K.; Pragst, F.; Fritzsche, T.; Spies, C. Narcotrend-assisted propofol/remifentanyl anaesthesia vs clinical practice: Does it make a difference? *Br. J. Anaesth.* **2007**, *99*, 686–693, doi:10.1093/bja/aem231.
  56. Sebel, P.S.; Bowdle, T.A.; Ghoneim, M.M.; Rampil, I.J.; Padilla, R.E.; Gan, T.J.; Domino, K.B. The incidence of awareness during anesthesia: A multicenter United States study. *Anesth. Analg.* **2004**, *99*, 833–839, doi:10.1213/01.ANE.0000130261.90896.6C.
  57. Ekman, A.; Lindholm, M.-L.; Lennmarken, C.; Sandin, R. Reduction in the incidence of awareness using BIS monitoring. *Acta Anaesthesiol. Scand.* **2004**, *48*, 20–26, doi:10.1111/j.1399-6576.2004.00260.x.
  58. Fiedler, M.O.; Schätzle, E.; Contzen, M.; Gernoth, C.; Weiß, C.; Walter, T.; Viergutz, T.; Kalenka, A. Evaluation of Different Positive End-Expiratory Pressures Using Supreme™ Airway Laryngeal Mask during Minor Surgical Procedures in Children. *Medicina (Kaunas)*. **2020**, *56*, 1–9, doi:10.3390/medicina56100551.
  59. Bonhomme, V.; Uutela, K.; Hans, G.; Maquoi, I.; Born, J.D.; Brichant, J.F.; Lamy, M.; Hans, P. Comparison of the Surgical Pleth Index™ with haemodynamic variables to assess nociception-anti-nociception balance during general anaesthesia. *Br. J. Anaesth.* **2011**, *106*, 101–111, doi:10.1093/bja/aeq291.
  60. Boselli, E.; Bouvet, L.; Bégou, G.; Dabouz, R.; Davidson, J.; Deloste, J.Y.; Rahali, N.; Zadam, A.; Allaouchiche, B. Prediction of immediate postoperative pain using the analgesia/nociception index: A prospective observational study. *Br. J. Anaesth.* **2014**, *112*, 715–721, doi:10.1093/bja/aet407.
  61. Ledowski, T.; Ang, B.; Schmarbeck, T.; Rhodes, J. Monitoring of sympathetic tone to assess postoperative pain: Skin conductance vs surgical stress index. *Anaesthesia* **2009**, *64*, 727–731, doi:10.1111/j.1365-2044.2008.05834.x.
  62. Nie, F.; Liu, T.; Zhong, L.; Yang, X.; Liu, Y.; Xia, H.; Liu, X.; Wang, X.; Liu, Z.; Zhou, L.I.; et al. MicroRNA-148b enhances proliferation and apoptosis in human renal cancer cells via directly targeting MAP3K9. **2016**, 83–90, doi:10.3892/mmr.2015.4555.
  63. Jess, G.; Pogatzki-Zahn, E.M.; Zahn, P.K.; Meyer-Frießem, C.H. Monitoring heart rate variability to assess experimentally induced pain using the analgesia nociception index. *Eur. J. Anaesthesiol.* **2016**, *33*, 118–125, doi:10.1097/EJA.0000000000000304.

64. Edry, R.; Recea, V.; Dikust, Y.; Sessler, D.I. Preliminary Intraoperative Validation of the Nociception Level Index: A Noninvasive Nociception Monitor. *Anesthesiology* **2016**, *125*, 193–203, doi:10.1097/ALN.0000000000001130.
65. Gruenewald, M.; Dempfle, A. Analgesia/nociception monitoring for opioid guidance: Meta-Analysis of randomized clinical trials. *Minerva Anesthesiol.* **2017**, *83*, 200–213, doi:10.23736/S0375-9393.16.11602-5.
66. Wang, Y.L.; Kong, X.Q.; Ji, F.H. Effect of dexmedetomidine on intraoperative Surgical Pleth Index in patients undergoing video-assisted thoracoscopic lung lobectomy. *J. Cardiothorac. Surg.* **2020**, *15*, 1–7, doi:10.1186/s13019-020-01346-1.
67. Kim, J.H.; Jwa, E.K.; Choung, Y.; Yeon, H.J.; Kim, S.Y.; Kim, E. Comparison of Pupillometry With Surgical Pleth Index Monitoring on Perioperative Opioid Consumption and Nociception During Propofol-Remifentanyl Anesthesia: A Prospective Randomized Controlled Trial. *Anesth. Analg.* **2020**, *131*.
68. Huiku, M.; Uutela, K.; van Gils, M.; Korhonen, I.; Kymalainen, M.; Merilainen, P.; Paloheimo, M.; Rantanen, M.; Takala, P.; Viertio-Oja, H.; et al. Assessment of surgical stress during general anaesthesia. *Br. J. Anaesth.* **2007**, *98*, 447–455, doi:10.1093/bja/aem004.
69. Dostalova, V.; Schreiberova, J.; Bartos, M.; Kukralova, L.; Dostal, P. Surgical pleth index and analgesia nociception index for intraoperative analgesia in patients undergoing neurosurgical spinal procedures: A comparative randomized study. *Minerva Anesthesiol.* **2019**, *85*, 1265–1272, doi:10.23736/S0375-9393.19.13765-0.
70. Funcke, S.; Saugel, B.; Koch, C.; Schulte, D.; Zajonz, T.; Sander, M.; Gratarola, A.; Ball, L.; Pelosi, P.; Spadaro, S.; et al. Individualized, perioperative, hemodynamic goal-directed therapy in major abdominal surgery (iPEGASUS trial): study protocol for a randomized controlled trial. *Trials* **2018**, *19*, 273, doi:10.1186/s13063-018-2620-9.
71. Jain, N.; Gera, A.; Sharma, B.; Sood, J.; Chugh, P. Comparison of Surgical Pleth Index-guided analgesia using fentanyl versus conventional analgesia technique in laparoscopic cholecystectomy. *Minerva Anesthesiol.* **2019**, *85*, 358–365, doi:10.23736/S0375-9393.18.12954-3.
72. Won, Y.J.; Lim, B.G.; Lee, S.H.; Park, S.; Kim, H.; Lee, I.O.; Kong, M.H. Comparison of relative oxycodone consumption in surgical pleth index-guided analgesia versus conventional analgesia during sevoflurane anesthesia. *Medicine (Baltimore)*. **2016**, *95*, e4743, doi:10.1097/MD.0000000000004743.
73. Chen, X.; Thee, C.; Gruenewald, M.; Ilies, C.; Höcker, J.; Hanss, R.; Steinfath, M.; Bein, B. Correlation of Surgical Pleth Index with Stress Hormones during Propofol-Remifentanyl Anaesthesia. *Sci. World J.* **2012**, *2012*, 1–8, doi:10.1100/2012/879158.
74. Theerth, K.A.; Sriganesh, K.; Reddy, K.M.; Chakrabarti, D.; Umamaheswara Rao, G.S. Analgesia Nociception Index-guided intraoperative fentanyl consumption and postoperative analgesia in patients receiving scalp block versus incision-site infiltration for craniotomy. *Minerva Anesthesiol.* **2018**, *84*, 1361–1368, doi:10.23736/S0375-9393.18.12837-9.
75. Soral, M.; Altun, G.T.; Dinçer, P.Ç.; Arslantaş, M.K.; Aykaç, Z. Effectiveness of the analgesia nociception index monitoring in patients who undergo colonoscopy with sedo-analgesia. *Turkish J. Anaesthesiol. Reanim.* **2020**, *48*, 50–57, doi:10.5152/TJAR.2019.45077.
76. Gall, O.; Champigneulle, B.; Schweitzer, B.; Deram, T.; Maupain, O.; Montmayeur Verchere, J.; Orliaguet, G. Postoperative pain assessment in children: A pilot study of the usefulness of the analgesia nociception index. *Br. J. Anaesth.* **2015**, *115*, 890–895, doi:10.1093/bja/aev361.
77. JENSEN, E.W.; VALENCIA, J.F.; LÓPEZ, A.; ANGLADA, T.; AGUSTÍ, M.; RAMOS, Y.; SERRA, R.; JOSPIN, M.; PINEDA, P.; GAMBUS, P. Monitoring hypnotic effect and nociception with two EEG-derived indices, qCON and qNOX, during general anaesthesia. *Acta Anaesthesiol. Scand.* **2014**, *58*, 933–941, doi:https://doi.org/10.1111/aas.12359.
78. Reich, D.L.; Hossain, S.; Krol, M.; Baez, B.; Patel, P.; Bernstein, A.; Bodian, C.A. Predictors of hypotension after induction of general anesthesia. *Anesth. Analg.* **2005**, *101*, 622–8, table of contents, doi:10.1213/01.ANE.0000175214.38450.91.
79. Hug, C.C.J.; McLeskey, C.H.; Nahrwold, M.L.; Roizen, M.F.; Stanley, T.H.; Thisted, R.A.; Walawander, C.A.; White, P.F.; Apfelbaum, J.L.; Grasela, T.H. Hemodynamic effects of propofol: data from over 25,000 patients. *Anesth. Analg.* **1993**, *77*, S21-9.
80. Thwaites, A.; Edmonds, S.; Smith, I. Inhalation induction with sevoflurane: a double-blind comparison with propofol. *Br. J. Anaesth.* **1997**, *78*, 356–361, doi:10.1093/bja/78.4.356.



81. Liu, T.J.; Zhang, J.C.; Gao, X.Z.; Tan, Z. Bin; Wang, J.J.; Zhang, P.P.; Cheng, A. Bin; Zhang, S.B. Effect of sevoflurane on the ATPase activity of hippocampal neurons in a rat model of cerebral ischemia-reperfusion injury via the cAMP-PKA signaling pathway. *Kaohsiung J. Med. Sci.* **2018**, *34*, 22–33, doi:10.1016/j.kjms.2017.09.004.
82. Bijker, J.B.; Van Klei, W.A.; Kappen, T.H.; Van Wolfswinkel, L.; Moons, K.G.M.; Kalkman, C.J. Incidence of intraoperative hypotension as a function of the chosen definition: Literature definitions applied to a retrospective cohort using automated data collection. *Anesthesiology* **2007**, *107*, 213–220, doi:10.1097/01.anes.0000270724.40897.8e.
83. Chang, H.S.; Hongo, K.; Nakagawa, H. Adverse effects of limited hypotensive anesthesia on the outcome of patients with subarachnoid hemorrhage. *J. Neurosurg.* **2000**, *92*, 971–975, doi:10.3171/jns.2000.92.6.0971.
84. Monk, T.G. Processed EEG and patient outcome. *Best Pract. Res. Clin. Anaesthesiol.* **2006**, *20*, 221–228, doi:10.1016/j.bpa.2005.10.004.
85. Wesselink, E.M.; Kappen, T.H.; Torn, H.M.; Slooter, A.J.C.; van Klei, W.A. Intraoperative hypotension and the risk of postoperative adverse outcomes: a systematic review. *Br. J. Anaesth.* **2018**, *121*, 706–721, doi:10.1016/j.bja.2018.04.036.
86. Ristovic, V.; de Roock, S.; Mesana, T.; van Diepen, S.; Sun, L. The Impact of Preoperative Risk on the Association between Hypotension and Mortality after Cardiac Surgery: An Observational Study. *J. Clin. Med.* **2020**, *9*, 2057, doi:10.3390/jcm9072057.
87. Walsh, M.; Kurz, A.; Turan, A.; Rodseth, R.N.; Cywinski, J.; Thabane, L.; Ph, D.; Sessler, D.I. Relationship between Intraoperative Mean. *Anesthesiology* **2013**, *119*, 507–515.
88. Bedreag, O.H.; Sandesc, D.; Chiriac, S.D.; Rogobete, A.F.; Cradigati, A.C.; Sarandan, M.; Dumache, R.; Nartita, R.; Papurica, M. The Use of Circulating miRNAs as Biomarkers for Oxidative Stress in Critically Ill Polytrauma Patients. *Clin. Lab.* **2016**, *62*, 263–274, doi:10.7754/Clin.Lab.2015.150740.
89. Bratu, L.M.; Rogobete, A.F.; Sandesc, D.; Bedreag, O.H.; Tanasescu, S.; Nitu, R.; Popovici, S.E.; Crainiceanu, Z.P. The Use of Redox Expression and Associated Molecular Damage to Evaluate the Inflammatory Response in Critically Ill Patient with Severe Burn. *Biochem. Genet.* **2016**, *54*, doi:10.1007/s10528-016-9763-8.
90. Bedreag, O.H.; Rogobete, A.F.; Sandesc, D.; Cradigati, C.A.; Sarandan, M.; Popovici, S.E.; Dumache, R.; Horhat, F.G.; Vernic, C.; Sima, L. V; et al. Modulation of the Redox Expression and Inflammation Response in the Critically Ill Polytrauma Patient with Thoracic Injury. Statistical Correlations between Antioxidant Therapy. *Clin. Lab.* **2016**, *62*, 1747–1759, doi:10.7754/Clin.Lab.2016.160206.
91. Papurica, M.; Rogobete, A.F.; Sandesc, D.; Dumache, R.; Nartita, R.; Sarandan, M.; Cradigati, A.C.; Luca, L.; Vernic, C.; Bedreag, O.H. Redox Changes Induced by General Anesthesia in Critically Ill Patients with Multiple Traumas. *Mol. Biol. Int.* **2015**, *2015*, 238586, doi:10.1155/2015/238586.
92. Dinu, A.R.; Rogobete, A.F.; Bratu, T.; Popovici, S.E.; Bedreag, O.H.; Papurica, M.; Bratu, L.M.; Sandesc, D. Cannabis Sativa Revisited – Crosstalk between Stress , and Endocannabinoid Response System in. 1–23.
93. Nunes, R.R.; Nora, F.S.; Maia, D.; Dumaresq, H.; Maria, R.; Cavalcante, A.; Costa, A.A.; Moreira, L.; Carneiro, M.; Cesar, J.; et al. Influence of Total Intravenous Anesthesia , Entropy and Laparoscopy on Oxidative Stress. *Brazilian J. Anesthesiol.* **2012**, *62*, 484–501, doi:10.1016/S0034-7094(12)70147-2.
94. Gruber, J.; Fong, S.; Chen, C.; Yoong, S.; Pastorin, G.; Schaffer, S.; Cheah, I.; Halliwell, B. Mitochondria-targeted antioxidants and metabolic modulators as pharmacological interventions to slow ageing. *Biotechnol. Adv.* **2013**, *31*, 563–592, doi:10.1016/j.biotechadv.2012.09.005.
95. Lipina, C.; Hundal, H.S. Modulation of cellular redox homeostasis by the endocannabinoid system. **2016**.
96. Heyland, D.K.; Dhaliwal, R.; Suchner, U.; Berger, M.M. Antioxidant nutrients: a systematic review of trace elements and vitamins in the critically ill patient. *Intensive Care Med.* **2005**, *31*, 327–337, doi:10.1007/s00134-004-2522-z.
97. Sailaja Rao, P.; Kalva, S.; Yerramilli, A.; Mamidi, S. Free Radicals and Tissue Damage: Role of Antioxidants. *Free Radicals Antioxidants* **2011**, *1*, 2–7, doi:10.5530/ax.2011.4.2.
98. Hagar, H.H. An insight into the possible protective effect of pyrrolidine dithiocarbamate against lipopolysaccharide-induced oxidative stress and acute hepatic injury in rats. *Saudi Pharm. J.* **2009**, *17*, 259–267, doi:10.1016/j.jsps.2009.10.001.
99. Sorato, E.; Menazza, S.; Zulian, A.; Sabatelli, P.; Gualandi, F.; Merlini, L.; Bonaldo, P.; Canton, M.; Bernardi, P.; Di Lisa, F. Monoamine oxidase inhibition prevents mitochondrial dysfunction and apoptosis in myoblasts from patients with collagen VI myopathies. *Free Radic. Biol. Med.* **2014**, *75*, 40–47, doi:10.1016/j.freeradbiomed.2014.07.006.



100. Horhat, F.G.; Rogobete, A.F.; Papurica, M.; Sandesc, D.; Tanasescu, S.; Dumitrascu, V.; Licker, M.; Nitu, R.; Cradigati, C.A.; Sarandan, M.; et al. The Use of Lipid Peroxidation Expression as a Biomarker for the Molecular Damage in the Critically Ill Polytrauma Patient. **2016**, 1–7, doi:10.7754/Clin.Lab.2016.160306.
101. Moise, A.; Balescu-arion, C. Central European Annals of Clinical Research The Vitamin D and the Immune System . When ? Why ? How ? **2020**, 2, 1–9, doi:10.35995/ceacr2010001.
102. Georgescu, D.; Reisz, D.; Petre, I.; Ionita, I. Ischemic Stroke Secondary to Cerebral Venous Thrombosis : A Case Report. **2019**, 1–7, doi:10.35995/ceacr1010007.
103. Timar, C.; Negrău, M.; Pantiș, C.; Daina, C.; Stanciu, S.-D.; Hodoșan, V.; Cotrău, P. Septic Shock with Chlamydia Pneumoniae Secondary to Prostatic Abscess: A Rare Case Report. *Cent. Eur. Ann. Clin. Res.* **2019**, 1, 1, doi:10.35995/ceacr1010004.
104. Papurica, M.; Rogobete, A.F.; Sandesc, D.; Dumache, R.; Cradigati, C.A.; Sarandan, M.; Nartita, R.; Popovici, S.E.; Bedreag, O.H. Advances in biomarkers in critical ill polytrauma patients. *Clin. Lab.* **2016**, 62, 977–986, doi:10.7754/Clin.Lab.2015.151103.
105. Ferrari, R.S.; Andrade, C.F. Oxidative Stress and Lung Ischemia-Reperfusion Injury. **2015**, 2015.
106. Costa Paes, E.R. da; Braz, M.G.; Lima, J.T. de; Gomes da Silva, M.R.; Bentes de Sousa, L.; Lima, E.S.; Carvalho de Vasconcellos, M.; Cerqueira Braz, J.R. DNA damage and antioxidant status in medical residents occupationally exposed to waste anesthetic gases. *Acta Cir. Bras.* **2014**, 29, 280–286, doi:10.1590/s0102-86502014000400010.
107. Gan, T.J.; Diemunsch, P.; Habib, A.S.; Kovac, A.; Kranke, P.; Meyer, T.A.; Watcha, M.; Chung, F.; Angus, S.; Apfel, C.C.; et al. Consensus guidelines for the management of postoperative nausea and vomiting. *Anesth. Analg.* **2014**, 118, 85–113, doi:10.1213/ANE.0000000000000002.
108. Song, D.; Hamza, M.; White, P.F.; Klein, K.; Recart, A.; Khodaparast, O. The pharmacodynamic effects of a lower-lipid emulsion of propofol: a comparison with the standard propofol emulsion. *Anesth. Analg.* **2004**, 98, 687–91, table of contents, doi:10.1213/01.ane.0000103184.36451.d7.
109. El Hor, T.; Van Der Linden, P.; De Hert, S.; Melot, C.; Bidgoli, J. Impact of entropy monitoring on volatile anesthetic uptake. *Anesthesiology* **2013**, 118, 868–873, doi:10.1097/ALN.0b013e3182850c36.
110. Tewari, S.; Bhadoria, P.; Wadhawan, S.; Prasad, S.; Kohli, A. Entropy vs standard clinical monitoring using total intravenous anesthesia during transvaginal oocyte retrieval in patients for in vitro fertilization. *J. Clin. Anesth.* **2016**, 34, 105–112, doi:10.1016/j.jclinane.2016.02.029.
111. Refaat, E.K.; Yassein, T.E. Reduced sevoflurane consumption in cirrhotic compared to non-cirrhotic patients undergoing major hepatic surgery: During entropy monitored general anesthesia. *Egypt. J. Anaesth.* **2013**, 29, 61–65, doi:10.1016/j.egja.2012.07.006.