

Review

A critical review of disinfection processes to control SARS-CoV-2 transmission in the food industry

Adrián Pedreira^{1,2} , Yeşim Taşkın³ and Míriam R. García^{1*} 

¹ Bioprocess Engineering Group, IIM-CSIC, Vigo, Spain

² Lab of Recycling and Valorization of Waste Materials (REVAL), IIM-CSIC, Vigo, Spain

³ Food Engineering Department, Hacettepe University, Ankara, Turkey

* Correspondence: miriamr@iim.csic.es

Abstract: Industries of the food sector have made a great effort to control SARS-CoV-2 indirect transmission, through objects or surfaces, by updating cleaning and disinfection protocols previously focused on inactivating other pathogens, as well as food spoilage microorganisms. The information, although scarce at the beginning of the COVID-19 pandemic, has started to be sufficiently reliable to avoid over-conservative disinfection procedures. This work reviews the literature to propose a holistic view of the disinfection process where the decision variables, such as type and concentration of active substance, are optimised to guarantee the inactivation of SARS-CoV-2 and other usual pathogens and spoilage microorganisms while minimising possible side-effects on the environment and animal and human health.

Keywords: SARS-CoV-2; COVID-19 pandemic; food industry; disinfection trade-offs, one-health

1. Introduction

Efficient control of SARS-CoV-2 transmission is a high priority in the food industry. Food safety authorities agree that SARS-CoV-2, as other coronaviruses in previous outbreaks, is not transmitted through food consumption [1–3]. However, respiratory viral diseases are commonly spread via fomites (indirect route of transmission through inanimate objects or surfaces) [4,5] and different studies have proven that SARS-CoV-2 is stable for several days on different surfaces commonly used in the food industry [6–10]. The spread of the virus through this via should be prevented by properly disinfecting and cleaning of the inanimate objects.

The food industry has been implementing cleaning and disinfection measures to effectively inactivate spoilage and pathogenic microorganisms like bacteria, yeasts, molds, and viruses [11], and hence avoid food waste or poisoning outbreaks. Cleaning must be done thoroughly before the disinfection step. In general, the efficacy of all disinfectants decreases in the presence of organic matter and the goal is to remove as much organic matter as possible by eliminating the gross dirt and by cleaning with water and surfactants [12]. Disinfection aims to achieve an acceptable standard of hygiene from the microbiological point of view [13].

Quite often the industry disinfects with aggressive substances at large concentrations and contact times to ensure good hygiene standards that are more than enough to inactivate SARS-Cov-2. For example, sodium hypochlorite has been historically employed to inactivate very resistant pathogens as vegetative bacteria and bacteria spores[14] and it has also shown efficacy against viruses, including SARS-CoV-2 [15].

Therefore, given the so-far known apparent simplicity to disinfect encapsulated viruses like SARS-CoV-2 [16,17] and the current evidence of limited transmission of the virus through fomites [18,19], should the food industry update the standard disinfection protocols to guarantee the product and workers safety?

A precautionary principle, given the 2020 pandemic extend, might point to applying over-conservative disinfection protocols with large amounts of active substances and contact times or, at least, to revise and adapt the protocols to ensure inactivation of SARS-CoV-2 [20]. Crowded and cooling conditions, common in slaughterhouses and the meat processing facilities, have been related with SARS-CoV-2 outbreaks. For example, from 9–27 April 2020, COVID-19 was diagnosed in 4913 workers from 113 meat and poultry processing facilities in the USA (approximately 3% of the workforce), with

20 COVID-19-related deaths [21]. In a more recent report, including the confirmed cases up to May 31 from a total of 239 facilities, the number of cases increases to 16,233 (3.1% to 24.5% infected workers per facility) and 86 COVID-19 related deaths [22]. Risk factors included prolonged closeness to other workers for long shifts, close contact during transportation in shared vans and contact through exposure to potentially contaminated shared surfaces such as break room tables or tools [8,23].

Nevertheless, over-use of disinfection to inactivate SARS-CoV-2 disregards any other effect on microorganisms present in the food processing plant, or the impact of the residual concentration of the chemicals on the environment and human and animal health. In this regard, it is considered critical the emergence and selection of antimicrobial-resistant bacteria due to the over-use of antimicrobial soaps and disinfectant cleaners. This concern has been discussed in the clinic sector [24], but not yet in the food industry.

In this review, we propose to analyse the disinfection processes using the “one health” approach, a holistic view considering the impact of the process beyond the objective of controlling the transmission of SARS-CoV-2. The review, therefore, is organised in three sections demanding to (1) understand which disinfectants or mixtures of disinfectants, concentrations and contact times are enough to inactivate SARS-CoV-2, (2) how these disinfectants work towards inactivating other food pathogens in the food industry and (3) their side-effects in the environment and the animal and human health due to misuse or overuse of disinfectants.

2. Disinfectants, concentrations and contact times to inactivate SARS-CoV-2 in the food industry

2.1. SARS-CoV-2 stability on surfaces

Despite the zoonotic origin of SARS-CoV-2, the virus is not a foodborne pathogen and to avoid its transmission in the food industry, it is critical to control transmission through fomites or contaminated objects [25]. The virus is stable on usual surfaces in the food industry, especially on non-porous surfaces, and remains infective for 3-7 days at room temperature in smooth surfaces such as glass, plastic and stainless steel [26]. By contrast, the virus has lower stability on cardboard (up to 2 days) and paper (up to 3 hours) [15,27].

Previous reports about the prevalence of other coronaviruses on surfaces show similar behavior. hCoV 229E and SARS-CoV-1, which share similar stability kinetics with SARS-CoV-2 [27], showed a prevalence in ceramic tiles of 4-5 days [28,29]. On dispensable materials such as latex gloves, cotton gowns and cotton gauze sponges, coronaviruses have lower persistence times (≤ 48 hours) [30,31].

Food contact surfaces are typically made of materials where SARS-CoV-2 is stable for days, such as steel or plastic. Other common contact surfaces are made of materials like wood, rubber, ceramics, or glass [32]. All surfaces must be smooth, non-porous, and easily cleanable and free from large, randomly distributed irregularities such as pits, folds and crevices [33,34]. Regardless of the type of material, disinfection of objects and surfaces in contact with unprocessed foods is critical to avoid contaminated final products [35,36].

The stability time of SARS-CoV-2 on surfaces is affected by physico-chemical factors such as temperature, pH and humidity. Low temperatures extend virus viability from 7 (22°C) to 14 days (4°C), while stability is less than a day at 37°C and less than 5 minutes when exposed to 70°C [15]. SARS-CoV-2 shows a high persistence under pH ranging from 3-10 at room temperature. On the other hand, SARS-CoV-2 half-life at room temperature (24°C) decays from 18.6 hours to 6.3 hours when the relative humidity (RH) raises from 20% to 80% [37].

Table 1 summarises the stability of SARS-CoV-2, SARS-CoV-1 and other SARS-CoV surrogate coronaviruses on different surfaces at different temperatures and relative humidities. According to the literature, the stability of other coronaviruses on surfaces is analogous to SARS-CoV-2, being favored by smooth surfaces and low temperature and RH. However, the lack of a standardized methodology difficulties comparing the stability of the different viruses on same kind of surface. From the author's knowledge, to date just one work has compared the stability of SARS-CoV-2 with other coronavirus (SARS-CoV-1) under the same experimental conditions [27]. Another difficulty is the variability regarding the inoculum volume. Low volumes of virus inoculum are preferred since they probably provide a better simulation of the real mechanism of viral contamination of surfaces, such as sneezing, coughing or by touching with hands that have previously been in contact with mouth, eyes, nose or other contaminated surfaces.

Table 1 – continued from previous page

Virus	Surface / fomite	Inoculum	TCDI ₅₀	T (°C)	RH (%)	Stability	Ref.
Virus	Surface / fomite	Inoculum	TCDI₅₀	T (°C)	RH (%)	Stability	Ref.
hCoV 229E	Aluminium	10 µL	5.5·10 ⁵	21	55-70	>8 h	[31]
	Brasses >70% Cu	20 µL	1.4·10 ^{3(a)}	21	30-40	≤40 min	[29]
	Brasses ≥70% Cu	1 µL	1.4·10 ^{3(a)}	21	30-40	≤5 min	[29]
	Ceramic tile	20 µL	1.4·10 ^{3(a)}	21	30-40	>120 h	[29]
	Copper	1 µL	1.4·10 ^{3(a)}	21	30-40	≤5 min	[29]
	Copper-nickel ≥79% Cu	20 µL	1.4·10 ^{3(a)}	21	30-40	≤50 min	[29]
	Copper-nickel 70% Cu	20 µL	1.4·10 ^{3(a)}	21	30-40	≤2 h	[29]
	Cotton gauze sponges	10 µL	5.5·10 ^{5(a)}	21	55-70	≤6 h	[31]
	Glass	20 µL	1.4·10 ^{3(a)}	21	30-40	>120 h	[29]
	Latex gloves	10 µL	5.5·10 ⁵	21	55-70	≤6 h	[31]
	Plastic (PVC)	20 µL	1.4·10 ^{3(a)}	21	30-40	>120 h	[29]
	Plastic (PE)	500 µL	10 ⁷	21-25	ND	>72 h	[38]
	Plastic (PTFE)	20 µL	1.4·10 ^{3(a)}	21	30-40	>120 h	[29]
	Silicon rubber	20 µL	1.4·10 ^{3(a)}	21	30-40	≤120 h	[29]
	Steel	20 µL	1.4·10 ^{3(a)}	21	30-40	>120 h	[29]
hCoV OC43	Aluminium	10 µL	5.5·10 ⁵	21	55-70	≤3 h	[31]
	Cotton gauze sponges	10 µL	5.5·10 ⁵	21	55-70	≤1 h	[31]
	Latex gloves	10 µL	5.5·10 ⁵	21	55-70	≤1 h	[31]
MERS-CoV EMC/2012	Plastic (undefined)	100 µL	10 ⁶	20	40	≤72 h	[39]
	Plastic (undefined)	100 µL	10 ⁶	30	30	≤48 h	[39]
	Plastic (undefined)	100 µL	10 ⁶	30	80	≤24 h	[39]
	Steel	100 µL	10 ⁶	20	40	≤72 h	[39]
	Steel	100 µL	10 ⁶	30	30	≤48 h	[39]
	Steel	100 µL	10 ⁶	30	80	≤24 h	[39]
SARS-CoV-1 HKU39849	Plastic (Undefined)	10 µL	10 ⁷	22-25	80	>120 h	[40]
SARS-CoV-1 P9	Ceramic tile	300 µL	10 ⁶	20	ND	≤96 h	[28]
	Glass	300 µL	10 ⁶	20	ND	≤120 h	[28]
	Metal (undefined)	300 µL	10 ⁶	20	ND	>120 h	[28]
	Paper (filter paper)	300 µL	10 ⁶	20	ND	>120 h	[28]
	Paper (press paper)	300 µL	10 ⁶	20	ND	≤120 h	[28]
	Plastic (undefined)	300 µL	10 ⁶	20	ND	≤120 h	[28]
	Wood	300 µL	10 ⁶	20	ND	≤96 h	[28]
SARS-CoV-1 GUV6109	Cotton gown	5 µL	10 ⁴	37	ND	≤5 min	[30]
	Cotton gown	5 µL	10 ⁵	37	ND	≤1 h	[30]
	Cotton gown	5 µL	10 ⁶	37	ND	≤24 h	[30]
	Paper (request form)	5 µL	10 ⁴	37	ND	≤5 min	[30]
	Paper (request form)	5 µL	10 ⁵	37	ND	≤3 h	[30]
	Paper (request form)	5 µL	10 ⁶	37	ND	≤24 h	[30]
	Plastic disposable gown	5 µL	10 ⁴	37	ND	≤1 h	[30]
	Plastic disposable gown	5 µL	10 ⁵	37	ND	≤24 h	[30]
Plastic disposable gown	5 µL	10 ⁶	37	ND	≤48 h	[30]	
SARS-CoV-1 FFM1	Plastic (PE)	500 µL	10 ⁷	21-25	ND	≤216h	[38]
SARS-CoV-1 Tor2	Capboard	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤8h	[27]
	Copper	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤8h	[27]
	Plastic (PP)	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤72h	[27]
	Steel	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤72h	[27]

Table 1 – continued from previous page

Virus	Surface / fomite	Inoculum	TCDI ₅₀	T (°C)	RH (%)	Stability	Ref.
SARS-CoV-2 WA1-2020	Capboard	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤24h	[27]
	Copper	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤4h	[27]
	Plastic (PP)	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤72h	[27]
	Steel	50 µL	10 ⁷	RC ^(c)	RC ^(c)	≤72h	[27]
SARS-CoV-2 Unknow strain	Paper (tissue paper)	5 µL	10 ⁷ -10 ⁸	22	65	≤3h	[15]
	Paper (undefined)	5 µL	10 ⁷ -10 ⁸	22	65	≤3h	[15]
	Wood	5 µL	10 ⁷ -10 ⁸	22	65	≤3h	[15]
	Cloth	5 µL	10 ⁷ -10 ⁸	22	65	≤48h	[15]
	Glass	5 µL	10 ⁷ -10 ⁸	22	65	≤96h	[15]
	Paper (banknote)	5 µL	10 ⁷ -10 ⁸	22	65	≤96h	[15]
	Steel	5 µL	10 ⁷ -10 ⁸	22	65	≤168h	[15]
	Plastic (undefined)	5 µL	10 ⁷ -10 ⁸	22	65	≤168h	[15]
	Mask (inner layer)	5 µL	10 ⁷ -10 ⁸	22	65	≤168h	[15]
	Mask (outter layer)	5 µL	10 ⁷ -10 ⁸	22	65	>168h	[15]

Table 1. Persistence time of SARS-CoV-1, SARS-CoV-2, MERS-CoV and hCoV over different surfaces. TCDI₅₀: Median tissue culture infectious dose, RH: relative humidity, ND: not defined, PVC: polivynil chloride, PE: polyethylene, PTFE: polytetrafluoroethylene, PP: polypropylylene, RC: room conditions. ^(a): calculated from plaque forming units (PFU) following the instructions (1 TCID₅₀ ≈ 0.7 PFU) reported by the American Type Culture Collection (ATCC). ^(c): personal communication from the authors.

2.2. Approved substances for SARS-CoV-2 disinfection

The food industry should ensure good hygienic practices and food safety management systems following the advice of international authorities, such as the Food and Agriculture Organizations of the United Nations, to prevent COVID-19 transmission [1,41].

For more specific information regarding the necessary standards for chemical disinfection of SARS-CoV-2, the industry comply with regulations of their respective competent authorities. Table 2 summarizes those active substances which can be employed in the formulation of disinfectants legally available against SARS-CoV-2 in the European Union and in the USA, as well as their status in the food industry of both territories.

	SARS-2-CoV status		Food industry status		Concentration (%) in CP	
	ECHA	EPA	ECHA	EPA	EU	USA
Alcohols						
Ethanol	AP	AP	UR	AP	65.00-75.00	7.50-68.61
1-propanol	AP	NE	AP	AP	17.00-49.00	b
2-propanol	AP	AP	AP	AP	9.99	12.25-63.25
Aldehydes						
Glutaraldehyde	AP	AP	AP	AP	2.50-12.00	7.00
Glyoxal	AP	NE	AP	AP	6.00	b
Amines						
Ampholyt 20*	AP	NE	AP	NE	a	b
Glucoprotamin	AP	NE	AP	AP	a	b
N-(3-aminopropyl)- N-dodecylpropane-1,3-diamine	AP	AP	AP	AP	0.1-2.80	7.50-68.61
Biguanides						
Polyhexamethylene biguanide hydrochloride	AP	AP	AP	AP	a	0.050 - 0.089
Chlorine based compounds						

Calcium hypochlorite	AP	NE	AP	AP	a	b
Chlorine dioxide	UR	AP	UR	AP	a	0.20-5.00
Hypochlorous acid	UR	AP	UR	AP	0.017	0.017-0.046
Sodium hypochlorite	AP	AP	AP	AP	2.60-13.00	0.086-8.60
Sodium chlorite	UR	AP	UR	AP	a	0.50-30.50
Sodium dichloroisocyanurate	UR	AP	UR	AP	81.00	7.00-48.21
Tetrachlorodecaoxide complex	UR	NE	UR	NE	a	b
Tosylchloramide sodium	UR	NE	UR	AP	a	b
Trichloroisocyanuric acid	UR	NE	UR	AP	a	b
Iodine and iodophors						
Iodine	AP	NE	AP	AP	a	b
Povidone-iodine	AP	NE	AP	AP	a	b
Isothiazolinones						
Mixture of CMIT/MIT**	AP	NE	AP	AP	a	b
Organic acids						
Citric acid	AP	AP	AP	AP	a	0.60-6.00
Formic acid	UR	NE	UR	AP	a	b
Glycolic acid	UR	AP	UR	AP	a	11.19
Lactic acid	AP	AP	AP	AP	0.42-1.75	0.16-34.10
Performic acid	UR	NE	UR	AP	a	b
Peroxides and derivates						
Hydrogen peroxide	AP	AP	AP	AP	0.20-35.00	0.30-27.50
Peracetic acid	AP	AP	AP	AP	0.05-5.00	0.05-15.00
Peroxyoctanoic acid	UR	AP	UR	AP	a	0.63
Potassium peroxymonosulfate	NE	AP	UR	AP	49.70	21.41
Sodium carbonate peroxyhydrate	NE	AP	NE	AP	a	12.10-29.75
Phenolic compounds						
2-Phenylphenol	UR	AP	UR	AP	a	0.026-10.50
2-Benzyl-4-chlorophenol	NE	AP	NE	AP	a	0.023-3.03
4-tert-amylphenol	NE	AP	NE	AP	a	5.27-7.66
5-chloro-2-(4-chlorophenoxy) phenol	AP	NE	AP	NE	a	b
Biphenyl-2-ol	AP	AP	AP	AP	a	0.06
Salicylic acid	UR	NE	UR	AP	a	b
Thymol	NE	AP	NE	AP	a	0.092-0.23
Quaternary ammonium compounds						
Benzalkonium chloride	AP	AP	UR	AP	0.008-24.00	0.015-26.00
Benzalkonium saccharinate	AP	AP	UR	AP	a	0.10-0.20
Benzethonium chloride	NE	AP	NE	AP	a	0.28
Didecyldimethylammonium chloride	AP	AP	UR	AP	0.20-7.20	0.003-21.05
Didecylmethylpoly(oxethyl) ammonium propionate	AP	NE	UR	NE	a	b
Didecylmethylammonium carbonate/ bicarbonate	NE	AP	NE	AP	a	0.0369-1.38
Silver and derivates						
Silver	AP	AP	UR	AP	0.004	0.003-0.01

Silver nitrate	AP	AP	UR	AP	a	0.016
----------------	----	----	----	----	---	-------

Table 2. Status of the substances against SARS-CoV-2 according to ECHA and EPA. CP: Commercial products; AP: Approved; UR: Under review; NE: Not evaluated. a: Substance not found in the formulation of commercial biocides authorized by ECHA to be employed in the food and feed areas (PT 4). b: Substance not found in the formulation of EPA authorised commercial biocides against SARS-CoV-2 (not area distinction). Compounds tagged as "Under review" in ECHA column can be part of the formulation of authorized commercial products employed in some European countries if they are covered by their respective legislations [42]. Substances concentrations in commercial products available in the EU were obtained from the state members of Spanish [43] and Dutch [44] markets, both selected as reference markets by ECHA [42]. Substances concentrations in commercial products available in the USA was extracted from the Emergency Care Research Institute (ECRI) [45]. *N-C10-16-alkyltrimethylenedi- reaction products with chloroacetic acid. **Mixture of 5-chloro-2-methyl-2H-isothiazol-3-one (EINECS 247-500-7) and 2-methyl-2H-isothiazol-3-one (EINECS 220-239-6)

The European Chemicals Agency (ECHA) is the competent authority in the European Union for determining legal disinfection substances. ECHA reports the lists of those substances approved or under review against SARS-CoV-2 for product-type 1 (Human hygiene) and 2 (disinfectants and algicides not intended for direct application to humans or animals), but not the list of substances for use in the feed and food area (product-type 4 or PT4) [42]. Therefore, the substances for disinfection of SARS-CoV-2 in the food industry should be in (1) the list of PT4 substances and in (2) any of the lists of substances approved for SARS-CoV-2.

The lists of authorised substances for SARS-CoV-2 are not a comprehensive representation of all the disinfectant substances legally employed in the EU, since compounds tagged as "Under review" (e.g: ethanol) can be part of the formulation of authorized commercial products employed in some European countries if they are covered by their respective national legislation. ECHA also released a list of commercial products authorised under the Biocidal Products Regulation (BPR) with virucidal claims. This product list lacks information about the concentrations of active substances in the commercial products. Intending to provide a better indication of the market situation for disinfectant products, ECHA provides as example information for some member states such as Spain [43] and the Netherlands [44].

The United States Environmental Protection Agency (EPA), the competent authority in the USA, publishes the so-called "List N" gathering the authorized commercial disinfectants and active substances for use against SARS-CoV-2 [46] and their contact times. According to EPA, the authorised commercial products against SARS-CoV-2 must comply with at least one of the following criteria:

- (a) Demonstrate efficacy against SARS-CoV-2.
- (b) Demonstrate efficacy against a pathogen that is harder to inactivate than SARS-CoV-2.
- (c) Demonstrate efficacy against another type of human coronavirus similar to SARS-CoV-2.

The directions of use of disinfectants depends on the target pathogen and are specified on the product label guidance. For SARS-CoV-2 the "List N" informs if the directions to follow should be for SARS-CoV-2 or for another pathogen if it was not directly tested for SARS-CoV-2.

As in the case of ECHA, the list lacks information about the concentration of active substances in commercial products. Fortunately, this information have been gathered and published by the Emergency Care Research Institute (ECRI), a non-profit healthcare organization [45]. Figure 1 shows that most of the EPA authorised commercial products available in USA market were not tested directly for SARS-CoV-2 1. Instead, most authorised substances were tested following the b) criteria. However, the ranking of pathogens attending to the difficulty of being disinfected is based on debatable arguments, as we will discuss in section 3.

Similarly to ECHA and EPA, there are other national authorities determining the status of disinfectants for SARS-CoV-2. In Australia, for example, the Therapeutic Goods Administration (TGA) has published a list with the commercial disinfectants in the Australian Register of Therapeutic Goods (ARTG) for use against SARS-CoV-2. Again, the authorised list does not provide information about the active ingredients nor the criteria followed to select the substances [47]. In Canada, the Health Products and Food Branch's (HPFB) of the governmental department Health Canada (HC), has developed a list that is continuously updated with the likely effective commercial hard-surface disinfectants against SARS-CoV-2 and their active ingredients [48]. Substance concentrations are not reported but can be consulted in the Drug Product Database (DPD) [49]. The approval of a product is subject to the evidence provided by the manufacturers but inclusion criteria are not available for the general public.




Figure 1. Commercial products authorized to be employed in the food industry (either requiring post-rise or not) against SARS-CoV-2 according to the EPA List N grouped by their acceptance criteria.

2.3. Efficacy of authorised disinfectants

In a few months, researchers have been actively working to determine which disinfectants and concentrations are effective against SARS-CoV-2, but still more work is needed. The information is still scarce and substances are proved mostly in suspension test, instead of using dried viruses over surfaces better mimicking the conditions of the food industry [50,51].

Moreover, comparison between studies is usually difficult due to the lack of standardized testing procedures such as the high variability of contact times, and the differences in the kind and the concentration of the substances employed to simulate organic load. The initial viral concentration is also highly variable among studies, ranging from 10^5 [52] to 10^8 [15], and sometimes it is even omitted. This concentration is important since the minimum inhibitory concentration (MIC) may change with the number of organisms inoculated for different viruses, the so-called inoculum effect [53–55]. From the author's knowledge, only one work has tested the virucidal activity of chemicals to SARS-CoV-2 using the international standards for disinfection on surfaces (ASTM E1052-20) and suspension (EN 14476:2013) [56].

2.3.1. Disinfectants tested against SARS-CoV-2 or similar coronavirus

The list of tested disinfectants against SARS-CoV-2 and SARS-Cov-2 surrogates changes continuously. We have gathered the information published until September 2020 in a table that can be consulted in the following link <https://doi.org/10.5281/zenodo.4297322>. This data-sheet covers those disinfectants tested against SARS-CoV-2 or other coronaviruses. Data were extracted from several research articles indicated in the reference row. The data-sheet comprises a total of 11 fields with info regarding the virus (virus and strain/isolate names), formulation (substance(s) and its concentration in percentage) and test characteristics (suspension or surface tested, kind of surface, use dilution, disinfectant and inoculum volumes, organic load type and concentrations and contact time) as well as their results, normalized in terms of \log_{10} viral infectivity reduction.

Just a few common disinfectants have been proven effective against SARS-CoV-2 at short contact times: 30-80% ethanol, 30-75% propanol, 0.45- 7.5% povidone-iodine and 5-6% sodium hypochlorite (household bleach) (See table in [10.5281/zenodo.4297322](https://doi.org/10.5281/zenodo.4297322)). All of them are allowed in the food industry and have been successfully tested against SARS-CoV-2,

149 reaching a fast reduction on viral infectivity of more than 4 orders of magnitude (99.99% reduction), the level suggested as
150 effective by the German Association for the Control of Virus Diseases and the Robert Koch Institute [57].

151 Other common disinfectants such as 0.1% benzalkonium chloride (BAC) were effective against SARS-CoV-2, but at the
152 expense of large contact times [56]. In fact, previous tests with quaternary ammonium compounds (QACs) such as BAC, and
153 other coronaviruses also showed the need for large contact times [58,59]. More specifically, two commercial disinfectant
154 products were tested: (1) Mikrobac Forte composed of BAC and dodecylbispropylene triamine and (2) Kohrsolin FF with
155 BAC, didecyltrimethylammonium chloride (DDAC) and glutaraldehyde as active ingredients. Results showed that, after 30
156 minutes, the disinfection was not as effective as using 80% ethanol with only 30 seconds of exposure [38]. Tests using BAC to
157 disinfect SARS-CoV-1 surrogate coronaviruses (such as human coronavirus causing common cold [59,60], canine coronavirus
158 and mouse hepatitis virus [58]) showed similar results to ethanol and propanol, although again with exposition times of
159 minutes (5-10 minutes).

160 More controversial is the efficacy of other common disinfectants in the food industry such as hydrogen peroxide.
161 Although enveloped viruses are more sensitive to hydrogen peroxide than non-enveloped ones [61], all seem to point to a low
162 performance of hydrogen peroxide against coronaviruses. Hydrogen peroxide is minimally effective against SARS-CoV-2,
163 reaching just a poor viral infectivity reduction of 1-1.8 \log_{10} in a work concentration of 1-6% after 30 seconds of exposition
164 [52]. Previous reports showing a good performance of hydrogen peroxide against coronaviruses are subordinate to long
165 exposure times or the presence of other active ingredients in the formulation. In hCoV 229e, a satisfactory $\geq 4.00 \log_{10}$ viral
166 infectivity reduction was achieved after 60 seconds of exposition to 0.5% hydrogen peroxide [62]. However, an unknown
167 percentage of non reported food-grade surfactants were also present in the formulation. In TGEV, a viral infectivity reduction
168 $\sim 5 \log_{10}$ was achieved employing vaporized 35% hydrogen peroxide [63]. Unfortunately, the long contact time (2-3 hours)
169 makes its application very cost-time expensive. The World Health Organization (WHO) Formulations I and II, which contains
170 0.125% hydrogen peroxide in their composition besides 80% ethanol (Formulation I) and 70% 2-propanol (Formulation II),
171 have very good results against coronaviruses but don't allow to ensure the hydrogen peroxide performance when it acts
172 alone. In fact, both 30% ethanol and 30% propanol, used independently reached the same inactivation levels [64].

173 It should be stressed that there are also substances legally commercialized against SARS-CoV-2 without proven efficacy.
174 For example, 2% polyhexamethylene biguanide (PHMB) [65] and salicylic acid 0.10% [56], have failed in SARS-CoV-2
175 inactivation, not reaching the recommended \log_{10} viral reduction ≥ 4.00 . However, PHMB is found in commercial products
176 against SARS-CoV-2 authorized by the EPA and it has been classified as authorized substance with the same purpose in the
177 ECHA list, which also includes salicylic acid.

178 Interestingly, disinfectants such as 4.7% chloroxynol [56] and 0.05% chlorhexidine [15] show good performance against
179 SARS-CoV-2, but have not been approved by EPA nor ECHA to SARS-CoV-2 disinfection. A reason for this contradiction
180 could be that, for example, chlorhexidine is not efficient in the inactivation of other coronaviruses [58].

181 So far, no other substances have been tested against SARS-CoV-2 and their use is just supported by previous reports
182 showing antiviral activity against other viruses from the Coronaviridae family. This is the case of DDAC, glutaraldehyde and
183 phenolic compounds. DDAC has never been tested against coronaviruses as a single active ingredient, but in mixtures with
184 other disinfectants [66]. Glutaraldehyde 0.5% gets a $\geq 4.01 \log_{10}$ reduction in SARS-CoV-1 after 120 seconds of exposition
185 [38]. Other studies employing glutaraldehyde 2% reported viral reductions $> 3.00 \log_{10}$ after 60 seconds in canine coronavirus
186 [67] and $5 \log_{10}$ after 24 hours in hCoV [60]. Several phenolic formulations reached a $> 3.00 \log_{10}$ viral reduction against
187 hCoV [60] and a mixture comprising o-phenylphenol 9.09% and p-tertiary amylphenol 7.66% obtained a reduction of 2.03
188 \log_{10} in transmissible gastroenteritis virus and 1.33 \log_{10} in mouse hepatitis virus [68].

189 2.3.2. Disinfectants tested against viruses different from coronavirus

190 There is a concerning large number of authorized commercial products for SARS-CoV-2 disinfection which contains
191 active ingredients that have not been tested against the COVID-19 causing virus nor other coronaviruses. This case includes
192 organic acids as well as thymol.

193 Organic acids have not been tested against SARS-CoV-2, with the exception of salicylic acid explained in previous
194 section [56], or other coronaviruses, and their virucidal effect against other viruses is controversial. Formic acid showed to be
195 ineffective against the non enveloped viruses mammalian orthoreovirus type 1 and bovine adenovirus type 1 [69]. In another
196 study with enveloped viruses, formic and citric acids were not effective against the bovine viral diarrhea virus, although
197 showed good results in the case of vaccinia virus, both enveloped viruses [70]. It should be noted that in these studies the

exposure times were long (10-30 min), avoiding to know its cost-time efficiency. Moreover, taking into account that the action mechanism of organics acids is related to the decrease in the extra-viral pH which alters the steric disposition of specific receptors on the surface [71], the high stability of SARS-CoV-2 in acidic conditions points to a subsequent resistance to organic acids. Despite all this, organic acids as citric, lactic, formic, and performic acid are active ingredients of SARS-CoV-2 disinfectants authorized by the EPA and the ECHA.

The use of thymol is authorized by the EPA but not by the ECHA. The reports about its virucidal effectiveness are scarce and restricted to herpes virus [72,73], without any reference against other viruses. Among the authorized commercial products against SARS-CoV-2, substances mixtures are very common. Although a synergy or additive effect could be expected in a components mixture, this fact should be previously evaluated [20]. As example, WHO Formulations I and II, which contains 0.125% hydrogen peroxide besides 80% ethanol (Formulation I) and 70% 2-propanol (Formulation II), have very good results against coronaviruses (See table in [10.5281/zenodo.4297322](https://doi.org/10.5281/zenodo.4297322)), but 30% ethanol and 30% propanol used independently reached the same inactivation levels [64].

3. Potential of SARS-CoV-2 disinfectants to inactivate other pathogens or spoilers in the food industry

The food industry should guarantee that other pathogens, or even food spoilage microorganisms, in addition to SARS-CoV-2 are inactivated. Adequate precautions should be taken to prevent food from being contaminated by disinfectants, during cleaning or disinfection of rooms, equipment or utensils [74]. Thus, the concentration of the active substance, as well as dilutions (if needed) and exposure times, have to be calculated taking into account the presence of other microbes.

According to the modern interpretation of the Spaulding classification (Figure 2) [16,17], enveloped viruses such as coronaviruses are weak and easy to eradicate in comparison to other pathogens because the disruption of their lipidic envelope is considered enough to render them non-infectious.




Figure 2. Susceptibility level against disinfection of some foodborne pathogens. Adapted from [17]

The Spaulding classification provides a very useful overview but disregards many factors affecting disinfection depending on the type of microorganisms. Some factors are the organic load concentration and type of surfaces and the

220 clumping and biofilm formation of virus and bacteria, respectively [17]. In addition, useful parameters to consider when
 221 disinfecting, such as the persistence time on surfaces, do not follow the ranking in Spaulding classification. For example,
 222 SARS-CoV-2 persistence on certain surfaces is higher or similar than the persistence of harder-to-eradicate gram-positive
 223 bacteria (*Vibrio cholerae* 1-7 days, *Helicobacter pylori* \leq 90 minutes) and non-enveloped viruses (norovirus 8 hours - 7 days,
 224 echovirus 7 days and papovavirus 8 days) [11].

225 Even if those affecting factors are not in place, for the optimal disinfection of different usual microorganisms in the food
 226 industry together with SARS-CoV-2 the Spaulding classification is only a first approximation and more information is usually
 227 needed. On the one hand, and in agreement with the classification, disinfectants might only work on more susceptible
 228 groups. For example, 0.1% BAC inactivates SARS-CoV-2 but is not sufficiently effective for usual resistant bacterial strains
 229 found in the food industry such as *Staphylococcus* spp., *Klebsiella* spp. and *Escherichia coli* [15,56]. However, the use of strong
 230 disinfectants to kill hard-to-eradicate groups, does not always assure the disinfection of susceptible groups as Spaulding
 231 classification has many exceptions. For example, hydrogen peroxide, considered a high-level common disinfectant [17], is
 232 not very effective against SARS-CoV-2 [52]

233 Attending to the information gathered, those disinfectants that have been clearly proven effective against SARS-CoV-2,
 234 i.e. ethanol, propanol, povidone-iodine and sodium hypochlorite, have also been historically employed for other pathogens
 235 as vegetative bacteria, bacteria spores, fungi, enveloped and non enveloped viruses (including different coronaviruses)
 236 with remarkably efficacy [14]. Therefore a standard two-stage disinfection comprising a first cleaning stage with detergent
 237 followed by the application of a disinfectant should be enough to inactivate SARS-CoV-2 if proper disinfectant, concentration
 238 and exposure times are employed.

239 On the other hand, alternative substances that have not been tested on coronaviruses or show a poor performance
 240 against SARS-CoV-2, such as organic acids, thymol and PHMB, have also a controversial efficacy against other pathogens and
 241 are not recommended. The activity of organic acids is highly dependent of the pH and they are considered as bacteriostatic/
 242 fungistatic substances instead of bactericide/ fungicide [75]. In addition, acidophilic organism as acetic bacteria has a high
 243 tolerance to organic acids and some bacteria, fungi and yeasts can even employ them as carbon source. Likewise, thymol
 244 seems not to be active against *Pseudomonas* spp., a food-spoilage and pathogen bacteria genus commonly present in food
 245 industry [76–78] and PHMB is able to inactivate vegetative bacteria and yeast but its ehas not been demonstrated sporicide
 246 nor sporostatic activity [79].

247 4. Disinfection under the paradigm of the "one-health" approach

248 Although necessary, disinfection has side-effects that should be minimized by proper selection of disinfectants and
 249 concentrations. Understanding disinfection trade-offs is particularly relevant within the urgency of a pandemic where
 250 over-conservative and aggressive disinfection may be preferred due to the lack of solid evidence about the new pathogen in
 251 the literature.

252 The majority of chemicals used in disinfection are harmful or corrosive, being dangerous its application [80,81]. As a
 253 matter of fact, on the first month of 2020 the number of daily exposures to cleaners and disinfectants reported to U.S. poison
 254 centers increased substantially [82]. Workers in the food industries should be properly trained in this regard. However, under
 255 the high demand for disinfectants, there might be a shortage of required active compounds [83], leading to the employment
 256 of more aggressive substances.

257 Moreover, non-volatile disinfectants might reach the environment and cause harm at different levels. Inactivation of
 258 bacteria involved in relevant transformations may disrupt ecosystems, such as the case of bacteria transforming nitrogenous
 259 compounds [84]. Non-volatile compounds may remain on solid surfaces and, after water rinsing or natural precipitation,
 260 may contaminate food [85] or move to soils or water [81]. QACs, for example, are toxic to aquatic organisms [86] and
 261 have been related with a decrease of mouse fertility [87]. Chlorinated disinfectant residues may be also lethal for aquatic
 262 organisms, but also toxic to terrestrial animals (birds and mammals) [81]. Silver exposition may result in toxic effects mostly
 263 for skin and liver, given sufficient dosage and lengths of exposition [88,89].

264 Disinfection by-products (DBPs) may be also a concern. Due to water sanitation, chlorination by-products, such as
 265 chloroform, trihalomethanes and haloacetic acids, have been extensively studied with many proven associated risks [84].

266 Last but not least overuse or misuse of certain disinfectants in the food industry may promote resistance acquisition or
 267 selection, which constitutes a major threat to human and animal health [90,91]. Phenolics ([92,93]), glutaraldehyde ([94,95])

268 and silver ([89,96]), for example, are allowed substances to inactivate SARS-Cov-2 with proven risks of promoting resistance
269 and cross-resistance.

270 QACs, however, are probably the most alarming substances in the food industry because their widespread employment
271 and potential to generate resistance or even cross-resistance with antibiotics [91,97,98]. Resistance to benzalkonium chloride,
272 the most common QAC, has been measured in 57 bacteria species [99]. Certain strains even have MIC values several times
273 higher (1000-3000 mg/L) than the BAC concentration in commercial disinfectants covered by ECHA and EPA, making them
274 ineffective in their recommended dosage. Although other mechanism can be involved, it is assumed that QAC resistance is
275 mainly mediated by *qac* genes, responsible of the synthesis of efflux pumps. These multidrug efflux pumps are non-specific
276 detoxification mechanism and, thus, sub-MIC QAC exposition can also lead to the promotion of cross-resistance to dissimilar
277 biocides and antibiotics. In relation to the food industry, *qac* genes have been found in several *E.coli* strains isolated from
278 retail meats in the USA [100] and China [101]. In addition, BAC exposition increased its tolerance in 76 bacterial strains
279 isolated from food, reducing their susceptibility to other biocides (hexachlorophene, DDAC, triclosan and chlorhexidine) and
280 antibiotics such as ampicillin, sulfamethoxazol and cefotaxime [102], and conferred different degrees of resistance against
281 oxytetracycline, amoxicillin, ampicillin, levofloxacin and gentamicin in *Salmonella* sp. isolated from supermarket meat [103].

282 Fortunately, no evidences of genetic resistance have been reported to ethanol, propanol, povidone-iodine and sodium
283 hypochlorite, the most effective disinfectants against SARS-CoV-2. Moreover, broad sense resistance to these substances is,
284 when existing, derived from biofilm formation and thus easily avoidable if proper cleaning is carried out before disinfection
285 [75,104]. However, they have also disadvantages. A repeated use of alcohols over certain inanimate surfaces can damage the
286 material. Iodine can cause brown color stains on certain porous materials such plastics and clothing. Free chlorine is an
287 aggressive chemical that can promote corrosion of metal surfaces, especially at higher concentrations. Chlorine gas can be
288 released from chlorine solutions such as sodium hypochlorite solutions when exposed to heat or acid substances commonly
289 found in domestic and industrial cleaners[14].

290 5. Conclusions

291 At the beginning of the pandemic, some works proposed to update the disinfection protocols in the food industry
292 to control the new pathogen by considering aggressive chemical treatments. As more evidence became available, this
293 necessity was not clear due to (1) the limited evidence of transmission through fomites, (2) the proven efficacy of the standard
294 disinfection protocols in the food industry to inactivate SARS-CoV-2, and (3) the possible impact of aggressive disinfection
295 on the environment and human and animal health, especially because of the potential emergence of bacterial resistance.

296 The following substances, based on current literature and regulations, are recommended for disinfection of SARS-CoV-2:
297 ethanol, propanol, povidone-iodine and sodium hypochlorite. They are the most effective disinfectants against SARS-CoV-2
298 showing, at the same time minor side-effects. Other disinfectants regulated against SARS-CoV-2, such as some QACs,
299 required longer contact times and may induce bacterial resistance and cross-resistance or being fluxed to the environment
300 causing harms to different ecosystems.

301 More research is needed, however, to optimally select the type of disinfectant, its concentration and its contact time
302 to inactivate major pathogens, including SARS-CoV-2, while minimizing the impact on the environment and animal and
303 human health. The effective concentrations of disinfectants are difficult to assess due to the lack of standardization among
304 different studies. On the other hand, only the impact of the disinfectants used on water treatment are well studied, but not
305 the active compounds used to disinfect objects or surfaces in the food industry.

306 **Author Contributions:** Conceptualization MRG; investigation AP and YT; writing-original draft AP, YT and MRG; writing-review &
307 editing, AP and MRG. All authors have read and agreed to the published version of the manuscript.

308 **Funding:** This research was funded by RTI2018-093560-J-I00 (MCIU/AEI/FEDER, UE), IN606A-2020/028 (Xunta de Galicia predoctoral
309 grant) and EIT Food RIS Fellowships.

310 **Acknowledgments:** We thank Jesus Valcarcel for useful discussions regarding chemical properties of active substances.

311 **Conflicts of Interest:** The authors declare no conflict of interest

312 References

- 313 1. Codex alimentarius. COVID-19 | CODEXALIMENTARIUS FAO-WHO. [http://www.fao.org/fao-who-codexalimentarius/
314 thematic-areas/COVID-19/en/](http://www.fao.org/fao-who-codexalimentarius/thematic-areas/COVID-19/en/), accessed on 2020-07-31.

- 315 2. EFSA. Coronavirus: no evidence that food is a source or transmission route. [https://www.efsa.europa.eu/en/news/coronavirus-](https://www.efsa.europa.eu/en/news/coronavirus-no-evidence-food-source-or-transmission-route)
316 [no-evidence-food-source-or-transmission-route](https://www.efsa.europa.eu/en/news/coronavirus-no-evidence-food-source-or-transmission-route), accessed on 2020-07-29.
- 317 3. WHO. COVID-19 and Food Safety: Guidance for food businesses: Interim guidance. *COVID-19 and Food Safety: Guidance for food*
318 *businesses: Interim guidance* 2020. doi:10.4060/ca8660en.
- 319 4. Boone, S.A.; Gerba, C.P. Significance of fomites in the spread of respiratory and enteric viral disease. *Applied and Environmental*
320 *Microbiology* 2007, 73, 1687–1696. doi:10.1128/AEM.02051-06.
- 321 5. Otter, J.A.; Yezli, S.; Salkeld, J.A.; French, G.L. Evidence that contaminated surfaces contribute to the transmission of hospital
322 pathogens and an overview of strategies to address contaminated surfaces in hospital settings. *American Journal of Infection Control*
323 2013, 41, S6. doi:10.1016/j.ajic.2012.12.004.
- 324 6. Olaimat, A.N.; Shahbaz, H.M.; Fatima, N.; Munir, S.; Holley, R.A. Food Safety During and After the Era of COVID-19 Pandemic.
325 *Frontiers in Microbiology* 2020, 11, 1854. doi:10.3389/fmicb.2020.01854.
- 326 7. Ceylan, Z.; Meral, R.; Cetinkaya, T. Relevance of SARS-CoV-2 in food safety and food hygiene: potential preventive measures,
327 suggestions and nanotechnological approaches, 2020. doi:10.1007/s13337-020-00611-0.
- 328 8. Zuber, S.; Brüssow, H. COVID 19: challenges for virologists in the food industry. *Microbial Biotechnology* 2020, 13, 1689–1701.
329 doi:10.1111/1751-7915.13638.
- 330 9. Roy, A.; Parida, S.P.; Bhatia, V. Role of disinfection and hand hygiene: a COVID-19 perspective. *International Journal Of Community*
331 *Medicine And Public Health* 2020, 7, 2845. doi:10.18203/2394-6040.ijcmph20203025.
- 332 10. Han, J.; Zhang, X.; He, S.; Jia, P. Can the coronavirus disease be transmitted from food? A review of evidence, risks, policies and
333 knowledge gaps, 2020. doi:10.1007/s10311-020-01101-x.
- 334 11. Kramer, A.; Schwebke, I.; Kampf, G. How long do nosocomial pathogens persist on inanimate surfaces? A systematic review, 2006.
335 doi:10.1186/1471-2334-6-130.
- 336 12. Holah, J.T. Cleaning and disinfection practices in food processing. In *Hygiene in Food Processing: Principles and Practice: Second*
337 *Edition*; Elsevier Inc., 2013; pp. 259–304. doi:10.1533/9780857098634.3.259.
- 338 13. Kakurinov, V. Food Safety Assurance Systems: Cleaning and Disinfection. In *Encyclopedia of Food Safety*; Elsevier, 2014; Vol. 4, pp.
339 211–225. doi:10.1016/B978-0-12-378612-8.00356-5.
- 340 14. McDonnell, G.; Russell, A.D. Antiseptics and disinfectants: activity, action, and resistance. *Clinical microbiology reviews* 1999,
341 12, 147–79. doi:10.1128/cmr.12.1.147.
- 342 15. Chin, A.W.H.; Chu, J.T.S.; Perera, M.R.A.; Hui, K.P.Y.; Yen, H.L.; Chan, M.C.W.; Peiris, M.; Poon, L.L.M. Stability of SARS-CoV-2 in
343 different environmental conditions. *The Lancet Microbe* 2020, 1, e10. doi:10.1016/s2666-5247(20)30003-3.
- 344 16. SPAULDING.; EH.. Chemical disinfection and antisepsis in the hospital. *J Hosp Res.* 1972, 9, 5–31.
- 345 17. McDonnell, G.; Burke, P. Disinfection: Is it time to reconsider Spaulding?, 2011. doi:10.1016/j.jhin.2011.05.002.
- 346 18. Goldman, E. Exaggerated risk of transmission of COVID-19 by fomites, 2020. doi:10.1016/S1473-3099(20)30561-2.
- 347 19. Mondelli, M.U.; Colaneri, M.; Seminari, E.M.; Baldanti, F.; Bruno, R. Low risk of SARS-CoV-2 transmission by fomites in real-life
348 conditions, 2020. doi:10.1016/S1473-3099(20)30678-2.
- 349 20. Quevedo-León, R.; Bastías-Montes, J.M.; Espinoza-Tellez, T.; Ronceros, B.; Balic, I.; Muñoz, O. Inactivation of Coronaviruses in food
350 industry: The use of inorganic and organic disinfectants, ozone, and UV radiation, 2020. doi:10.17268/SCI.AGROPECU.2020.02.14.
- 351 21. Dyal, J.W.; Grant, M.P.; Broadwater, K.; Bjork, A.; Waltenburg, M.A.; Gibbins, J.D.; Hale, C.; Silver, M.; Fischer, M.; Steinberg, J.;
352 Basler, C.A.; Jacobs, J.R.; Kennedy, E.D.; Tomasi, S.; Trout, D.; Hornsby-Myers, J.; Oussayef, N.L.; Delaney, L.J.; Patel, K.; Shetty, V.;
353 Kline, K.E.; Schroeder, B.; Herlihy, R.K.; House, J.; Jervis, R.; Clayton, J.L.; Ortbahn, D.; Austin, C.; Berl, E.; Moore, Z.; Buss, B.F.;
354 Stover, D.; Westergaard, R.; Pray, I.; DeBolt, M.; Person, A.; Gabel, J.; Kittle, T.S.; Hendren, P.; Rhea, C.; Holsinger, C.; Dunn, J.;
355 Turabelidze, G.; Ahmed, F.S.; DeFijter, S.; Pedati, C.S.; Rattay, K.; Smith, E.E.; Luna-Pinto, C.; Cooley, L.A.; Saydah, S.; Preacely, N.D.;
356 Maddox, R.A.; Lundeen, E.; Goodwin, B.; Karpathy, S.E.; Griffing, S.; Jenkins, M.M.; Lowry, G.; Schwarz, R.D.; Yoder, J.; Peacock, G.;
357 Walke, H.T.; Rose, D.A.; Honein, M.A. COVID-19 Among Workers in Meat and Poultry Processing Facilities — 19 States, April 2020.
358 *MMWR. Morbidity and Mortality Weekly Report* 2020, 69. doi:10.15585/mmwr.mm6918e3.
- 359 22. Waltenburg, M.A.; Victoroff, T.; Rose, C.E.; Butterfield, M.; Jervis, R.H.; Fedak, K.M.; Gabel, J.A.; Feldpausch, A.; Dunne, E.M.;
360 Austin, C.; Ahmed, F.S.; Tubach, S.; Rhea, C.; Krueger, A.; Crum, D.A.; Vostok, J.; Moore, M.J.; Turabelidze, G.; Stover, D.; Donahue,
361 M.; Edge, K.; Gutierrez, B.; Kline, K.E.; Martz, N.; Rajotte, J.C.; Julian, E.; Diedhiou, A.; Radcliffe, R.; Clayton, J.L.; Ortbahn, D.;
362 Cummins, J.; Barbeau, B.; Murphy, J.; Darby, B.; Graff, N.R.; Dostal, T.K.H.; Pray, I.W.; Tillman, C.; Dittrich, M.M.; Burns-Grant, G.;
363 Lee, S.; Spieckerman, A.; Iqbal, K.; Griffing, S.M.; Lawson, A.; Mainzer, H.M.; Bealle, A.E.; Edding, E.; Arnold, K.E.; Rodriguez, T.;
364 Merkle, S.; Pettrone, K.; Schlanger, K.; LaBar, K.; Hendricks, K.; Lasry, A.; Krishnasamy, V.; Walke, H.T.; Rose, D.A.; Honein, M.A.;
365 Amoroso, K.; Diallo, Y.; Fazekas, K.; Finley, P.J.; Fuld, J.; Guest, J.L.; Herstein, J.J.; Kennedy, E.D.; Lawler, J.V.; Lowe, J.J.; Neifert, A.;
366 Schwedhelm, M.M.; Medicine, N.; Steinberg, J.M.; Trout, D.B.; Zarate-Bermudez, M. Update: COVID-19 Among Workers in Meat

- and Poultry Processing Facilities — United States, April–May 2020. *MMWR. Morbidity and Mortality Weekly Report* **2020**, *69*, 887–892. doi:10.15585/mmwr.mm6927e2.
23. Stephenson, J. COVID-19 Outbreaks Among Food Production Workers May Intensify Pandemic's Disproportionate Effects on People of Color. *JAMA Health Forum* **2020**, *1*, e200783. doi:10.1001/jamahealthforum.2020.0783.
24. Murray, A.K. The Novel Coronavirus COVID-19 Outbreak: Global Implications for Antimicrobial Resistance. *Frontiers in Microbiology* **2020**, *11*, 1020. doi:10.3389/fmicb.2020.01020.
25. Duda-Chodak, A.; Lukasiewicz, M.; Zięć, G.; Florkiewicz, A.; Filipiak-Florkiewicz, A. Covid-19 pandemic and food: Present knowledge, risks, consumers fears and safety, 2020. doi:10.1016/j.tifs.2020.08.020.
26. Aboubakr, H.A.; Sharafeldin, T.A.; Goyal, S.M. Stability of SARS-CoV-2 and other coronaviruses in the environment and on common touch surfaces and the influence of climatic conditions: A review, 2020. doi:10.1111/tbed.13707.
27. van Doremalen, N.; Bushmaker, T.; Morris, D.H.; Holbrook, M.G.; Gamble, A.; Williamson, B.N.; Tamin, A.; Harcourt, J.L.; Thornburg, N.J.; Gerber, S.I.; Lloyd-Smith, J.O.; de Wit, E.; Munster, V.J. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New England Journal of Medicine* **2020**, *382*, 1564–1567. doi:10.1056/NEJMc2004973.
28. Duan, S.M.; Zhao, X.S.; Wen, R.F.; Huang, J.J.; Pi, G.H.; Zhang, S.X.; Han, J.; Bi, S.L.; Ruan, L.; Dong, X.P.; Team, S.R. Stability of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomedical and Environmental Sciences* **2003**, *3*, 246–255.
29. Warnes, S.L.; Little, Z.R.; Keevil, C.W. Human Coronavirus 229E Remains Infectious on Common Touch Surface Materials. *mBio* **2015**, *6*. doi:10.1128/mBio.01697-15.
30. Lai, M.Y.Y.; Cheng, P.K.C.; Lim, W.W.L. Survival of Severe Acute Respiratory Syndrome Coronavirus. *Clinical Infectious Diseases* **2005**, *41*, e67–e71. doi:10.1086/433186.
31. Sizun, J.; Yu, M.W.; Talbot, P.J. Survival of human coronaviruses 229E and OC43 in suspension and after drying on surfaces: A possible source of hospital-acquired infections. *Journal of Hospital Infection* **2000**, *46*, 55–60. doi:10.1053/jhin.2000.0795.
32. Skåra, T.; Rosnes, J.T. Emerging Methods and Principles in Food Contact Surface Decontamination/Prevention. In *Innovation and Future Trends in Food Manufacturing and Supply Chain Technologies*; Elsevier Inc., 2016; pp. 151–172. doi:10.1016/B978-1-78242-447-5.00006-X.
33. Holah, J.T.; Thorpe, R.H. Cleanability in relation to bacterial retention on unused and abraded domestic sink materials. *Journal of Applied Bacteriology* **1990**, *69*, 599–608. doi:10.1111/j.1365-2672.1990.tb01554.x.
34. Verran, J. Testing surface cleanability in food processing. In *Handbook of Hygiene Control in the Food Industry*; Elsevier Inc., 2005; pp. 556–571. doi:10.1533/9781845690533.3.556.
35. Hirneisen, K.A.; Markland, S.M.; Kniel, K.E. Ozone inactivation of norovirus surrogates on fresh produce. *Journal of Food Protection* **2011**, *74*, 836–839. doi:10.4315/0362-028X.JFP-10-438.
36. Mullis, L.; Saif, L.J.; Zhang, Y.; Zhang, X.; Azevedo, M.S. Stability of bovine coronavirus on lettuce surfaces under household refrigeration conditions. *Food Microbiology* **2012**, *30*, 180–186. doi:10.1016/j.fm.2011.12.009.
37. Biryukov, J.; Boydston, J.A.; Dunning, R.A.; Yeager, J.J.; Wood, S.; Reese, A.L.; Ferris, A.; Miller, D.; Weaver, W.; Zeitouni, N.E.; Phillips, A.; Freeburger, D.; Hooper, I.; Ratnesar-Shumate, S.; Yolitz, J.; Krause, M.; Williams, G.; Dawson, D.G.; Herzog, A.; Dabisch, P.; Wahl, V.; Hevey, M.C.; Altamura, L.A. Increasing temperature and relative humidity accelerates inactivation of SARS-COV-2 on surfaces. *mSphere* **2020**, *5*. doi:10.1128/mSphere.00441-20.
38. Rabenau, H.F.; Cinatl, J.; Morgenstern, B.; Bauer, G.; Preiser, W.; Doerr, H.W. Stability and inactivation of SARS coronavirus. *Medical Microbiology and Immunology* **2005**, *194*, 1–6. doi:10.1007/s00430-004-0219-0.
39. van Doremalen, N.; Bushmaker, T.; Munster, V.J. Stability of middle east respiratory syndrome coronavirus (MERS-CoV) under different environmental conditions. *Eurosurveillance* **2013**, *18*, 20590. doi:10.2807/1560-7917.ES2013.18.38.20590.
40. Chan, K.H.; Peiris, J.S.M.; Lam, S.Y.; Poon, L.L.M.; Yuen, K.Y.; Seto, W.H. The Effects of Temperature and Relative Humidity on the Viability of the SARS Coronavirus. *Advances in Virology* **2011**, *2011*, 1–7. doi:10.1155/2011/734690.
41. Codex Alimentarius Commission. Food hygiene. Basic texts., 2009.
42. ECHA. COVID-19 lists of disinfectant active substances and products - Datasets. <https://data.europa.eu/euodp/en/data/dataset/biocidal-products-lists-of-disinfectant-active-substances-and-products>, accessed on 2020-10-14.
43. Ministerio de Sanidad. VIRICIDAL PRODUCT AUTHORISED IN SPAIN. https://echa.europa.eu/documents/10162/29202019/spain_disinfection_products_pt1-pt2_art89_en.pdf/aba3d2dc-67d1-ddeb-dc77-2e778ef918c0, accessed on 2020-08-13.
44. Ctgb (Board for the Authorisation of Plant Protection Products and Biocides in the Netherlands). Vrijstelling desinfectiemiddelen oppervlakken COVID-19 2020' | Besluit | College voor de toelating van gewasbeschermingsmiddelen en biociden. <https://www.ctgb.nl/documenten/besluiten/2020/03/31/desinfectiemiddelen-oppervlakken-covid-19>, accessed on 2020-10-14.
45. ECRI. Disinfectant Concentrations, Contact Times, and Use Settings for EPA's List of Products Effective against SARS-CoV-2, the Cause of COVID-19, 2020.

- 420 46. EPA (United States Environmental Protection Agency). List N Tool: COVID-19 Disinfectants | US EPA. <https://cfpub.epa.gov/giwiz/disinfectants/index.cfm>, accessed on 2020-10-14.
- 422 47. Australian Government Department of Health Therapeutic Goods Administration. Disinfectants for use against COVID-19 in the ARTG for legal supply in Australia. <https://www.tga.gov.au/disinfectants-use-against-covid-19-artg-legal-supply-australia>, accessed on 2020-11-03.
- 424 48. Government of Canada. Hard-surface disinfectants and hand sanitizers (COVID-19): List of disinfectants with evidence for use against COVID-19. <https://www.canada.ca/en/health-canada/services/drugs-health-products/disinfectants/covid-19/list.html#tbl1>, accessed on 2020-11-16.
- 426 49. Drug Product Database: Access the database - Canada.ca. <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>, accessed on 2020-11-03.
- 429 50. Zeitler, B.; Rapp, I. Surface-dried viruses can resist glucoprotamin-based disinfection. *Applied and Environmental Microbiology* **2014**, *80*, 7169–7175. doi:10.1128/AEM.02462-14.
- 431 51. Kindermann, J.; Karbiener, M.; Leydold, S.M.; Knotzer, S.; Modrof, J.; Kreil, T.R. Virus disinfection for biotechnology applications: Different effectiveness on surface versus in suspension. *Biologicals* **2020**, *64*, 1–9. doi:10.1016/j.biologicals.2020.02.002.
- 433 52. Bidra, A.S.; Pelletier, J.S.; Westover, J.B.; Frank, S.; Brown, S.M.; Tessema, B. Comparison of In Vitro Inactivation of SARS CoV-2 with Hydrogen Peroxide and Povidone-Iodine Oral Antiseptic Rinses. *Journal of Prosthodontics* **2020**, *29*, 599–603. doi:10.1111/jopr.13220.
- 435 53. Marks, M.I. Variables influencing the in vitro susceptibilities of herpes simplex viruses to antiviral drugs. *Antimicrobial agents and chemotherapy* **1974**, *6*, 34–38. doi:10.1128/AAC.6.1.34.
- 437 54. Hayden, F.G.; Douglas, R.G.; Simons, R.; Vanvoris, P.; Betts, R.B.; Hayden, F.G.; Christman, W.A.; Douglas, R.G. Enhancement of Activity Against Influenza Viruses by Combinations of Antiviral Agents. *Antimicrobial Agents and Chemotherapy* **1980**, *18*, 536–541. doi:10.1128/aac.18.4.536.
- 439 55. Bruce Burlington, D.; Meiklejohn, G.; Mostowt, S.R. Anti-Influenza A Virus Activity of Amantadine Hydrochloride and Rimantadine Hydrochloride in Ferret Tracheal Ciliated Epithelium. *Antimicrobial Agents and Chemotherapy* **1982**, *21*, 794–799. doi:10.1128/aac.21.5.794.
- 441 56. Khalid Ijaz DVM, M.; Whitehead, K.B.; Srinivasan, V.M.; McKinney, J.; Rubino BA, J.R.; Ripley, M.; Jones, C.; Nims, R.W.; Charlesworth, B. Microbicidal actives with virucidal efficacy against SARS-CoV-2. *American Journal of Infection Control* **2020**, *48*. doi:10.1016/j.ajic.2020.05.019.
- 443 57. Rabenau, H.F.; Schwebke, I.; Blümel, J.; Eggers, M.; Glebe, D.; Rapp, I.; Sauerbrei, A.; Steinmann, E.; Steinmann, J.; Willkommen, H.; Wutzler, P. Guideline for testing chemical disinfectants regarding their virucidal activity within the field of human medicine : as of December 1st, 2014 Prepared by the German Association for the Control of Virus Diseases (DVV) and the Robert Koch Institute (RKI), 2020. doi:10.1007/s00103-020-03115-w.
- 445 58. Saknimit, M.; Inatsuki, I.; Sugiyama, Y.; Yagami, K. Virucidal efficacy of physico-chemical treatments against coronaviruses and parvoviruses of laboratory animals. *Jikken dobutsu. Experimental animals* **1988**, *37*, 341–345. doi:10.1538/expanim1978.37.3{_}341.
- 447 59. Wood, A.; Payne, D. The action of three antiseptics/disinfectants against enveloped and non-enveloped viruses. *Journal of Hospital Infection* **1998**, *38*, 283–295. doi:10.1016/S0195-6701(98)90077-9.
- 449 60. Sattar, S.A.; Springthorpe, V.S.; Karim, Y.; Loro, P. Chemical disinfection of non-porous inanimate surfaces experimentally contaminated with four human pathogenic viruses. *Epidemiology and Infection* **1989**, *102*, 493–505. doi:10.1017/S0950268800030211.
- 451 61. Eterpi, M.; McDonnell, G.; Thomas, V. Virucidal Activity of Disinfectants against Parvoviruses and Reference Viruses. *Applied Biosafety* **2010**, *15*, 165–171. doi:10.1177/153567601001500402.
- 453 62. Omidbakhsh, N.; Sattar, S.A. Broad-spectrum microbicidal activity, toxicologic assessment, and materials compatibility of a new generation of accelerated hydrogen peroxide-based environmental surface disinfectant. *American Journal of Infection Control* **2006**, *34*, 251–257. doi:10.1016/j.ajic.2005.06.002.
- 455 63. Goyal, S.M.; Chander, Y.; Yezli, S.; Otter, J.A. Evaluating the virucidal efficacy of hydrogen peroxide vapour. *Journal of Hospital Infection* **2014**, *86*, 255–259. doi:10.1016/j.jhin.2014.02.003.
- 457 64. Kratzel, A.; Kratzel, A.; Todt, D.; V'kovski, P.; V'kovski, P.; Steiner, S.; Steiner, S.; Gultom, M.; Gultom, M.; Thao, T.T.N.; Thao, T.T.N.; Ebert, N.; Ebert, N.; Holwerda, M.; Holwerda, M.; Steinmann, J.; Steinmann, J.; Niemeyer, D.; Dijkman, R.; Dijkman, R.; Kampf, G.; Drosten, C.; Steinmann, E.; Thiel, V.; Thiel, V.; Pfaender, S. Inactivation of Severe Acute Respiratory Syndrome Coronavirus 2 by WHO-Recommended Hand Rub Formulations and Alcohols. *Emerging Infectious Diseases* **2020**, *26*, 1592–1595. doi:10.3201/eid2607.200915.
- 459 65. Welch, S.R.; Davies, K.A.; Buczkowski, H.; Hettiarachchi, N.; Green, N.; Arnold, U.; Jones, M.; Hannah, M.J.; Evans, R.; Burton, C.; Burton, J.E.; Guiver, M.; Cane, P.A.; Woodford, N.; Bruce, C.B.; Roberts, A.D.G.; Killip, M.J. Inactivation analysis of SARS-CoV-2 by specimen transport media, nucleic acid extraction reagents, detergents and fixatives. *Journal of Clinical Microbiology* **2020**. doi:10.1128/JCM.01713-20.
- 472

- 473 66. Rabenau, H.; Kampf, G.; Cinatl, J.; Doerr, H. Efficacy of various disinfectants against SARS coronavirus. *Journal of Hospital Infection* **2005**, *61*, 107–111. doi:10.1016/j.jhin.2004.12.023.
- 474
- 475 67. Pratelli, A. Canine coronavirus inactivation with physical and chemical agents. *The Veterinary Journal* **2008**, *117*, 71–79.
- 476 doi:10.1016/j.tvjl.2007.03.019.
- 477 68. Hulkower, R.L.; Casanova, L.M.; Rutala, W.A.; Weber, D.J.; Sobsey, M.D. Inactivation of surrogate coronaviruses on hard surfaces
- 478 by health care germicides. *American Journal of Infection Control* **2011**, *39*, 401–407. doi:10.1016/j.ajic.2010.08.011.
- 479 69. Yilmaz, A.; Kaleta, E.F. Evaluation of virucidal activity of three commercial disinfectants and formic acid using bovine enterovirus
- 480 type 1 (ECBO virus), mammalian orthoreovirus type 1 and bovine adenovirus type 1. *Veterinary Journal* **2003**, *166*, 67–78.
- 481 doi:10.1016/S1090-0233(02)00269-1.
- 482 70. Heinzl, M.; Kvas, A.; Weide, M.; Breves, R.; Bockmühl, D.P. Evaluation of the virucidal performance of domestic laundry
- 483 procedures. *International Journal of Hygiene and Environmental Health* **2010**, *213*, 334–337. doi:10.1016/j.ijheh.2010.06.003.
- 484 71. Poli, G.; Biondi, P.A.; Uberti, F.; Ponti, W.; Balsari, A.; Cantoni, C. Virucidal activity of organic acids. *Food Chemistry* **1979**, *4*, 251–258.
- 485 doi:10.1016/0308-8146(79)90012-8.
- 486 72. Lai, W.L.; Chuang, H.S.; Lee, M.H.; Wei, C.L.; Lin, C.F.; Tsai, Y.C. Inhibition of herpes simplex virus type 1 by thymol-related
- 487 monoterpenoids. *Planta Medica* **2012**, *78*, 1636–1638. doi:10.1055/s-0032-1315208.
- 488 73. Sharifi-Rad, J.; Salehi, B.; Baghalpour, N.; Kobarfard, F.; Sharifi-Rad, M.; Mohammadzade, M. Antiviral activity of monoterpenes
- 489 thymol, carvacrol and p-cymene against herpes simplex virus in vitro. *Pharmacy Updates* **2018**, *1*, 73–73. doi:10.22037/IPA.V11I1.20032.
- 490 74. Codex Alimentarius Commission. Guidelines on the application of general principles of food hygiene to the control of viruses in
- 491 food, 2012.
- 492 75. Kampf, G. *Antiseptic stewardship: Biocide resistance and clinical implications*; Springer International Publishing, 2018; pp. 1–694.
- 493 doi:10.1007/978-3-319-98785-9.
- 494 76. Cosentino, S.; Tuberoso, C.I.; Pisano, B.; Satta, M.; Mascia, V.; Arzedi, E.; Palmas, F. In-vitro antimicrobial activity
- 495 and chemical composition of Sardinian Thymus essential oils. *Letters in Applied Microbiology* **1999**, *29*, 130–135.
- 496 doi:10.1046/j.1472-765X.1999.00605.x.
- 497 77. Walsh, S.E.; Maillard, J.Y.; Russell, A.D.; Catrenich, C.E.; Charbonneau, D.L.; Bartolo, R.G. Development of bacterial
- 498 resistance to several biocides and effects on antibiotic susceptibility. *Journal of Hospital Infection* **2003**, *55*, 98–107.
- 499 doi:10.1016/S0195-6701(03)00240-8.
- 500 78. Noma, Y.; Asakawa, Y. Biotransformation of monoterpenoids. In *Comprehensive Natural Products II: Chemistry and Biology*; Elsevier
- 501 Ltd, 2010; Vol. 3, pp. 669–801. doi:10.1016/b978-008045382-8.00742-5.
- 502 79. ECHA. Biocidal Products Committee (BPC) Opinion on the application for approval of the active substance: PHMB (1415;
- 503 4.7) Polyhexamethylene biguanide hydrochloride with a mean number- average molecular weight (Mn) of 1415 and a mean
- 504 polydispersity (PDI) of 4.7). <https://echa.europa.eu/fr/addressing-chemicals-of-concern/biocidal-products->, accessed on
- 505 2020-09-07.
- 506 80. Bonin, L.; Vitry, V.; Olivier, M.G.; Bertolucci-Coelho, L. Covid-19: effect of disinfection on corrosion of surfaces. *Corrosion Engineering,*
- 507 *Science and Technology* **2020**, *55*, 693–695. doi:10.1080/1478422X.2020.1777022.
- 508 81. Nabi, G.; Wang, Y.; Hao, Y.; Khan, S.; Wu, Y.; Li, D. Massive use of disinfectants against COVID-19 poses potential risks to urban
- 509 wildlife. *Environmental Research* **2020**, *188*, 109916. doi:10.1016/j.envres.2020.109916.
- 510 82. Chang, A.; Schnall, A.H.; Law, R.; Bronstein, A.C.; Marraffa, J.M.; Spiller, H.A.; Hays, H.L.; Funk, A.R.; Mercurio-Zappala, M.;
- 511 Calello, D.P.; Aleguas, A.; Borys, D.J.; Boehmer, T.; Svendsen, E. Cleaning and Disinfectant Chemical Exposures and Temporal
- 512 Associations with COVID-19 — National Poison Data System, United States, January 1, 2020–March 31, 2020. *MMWR. Morbidity*
- 513 *and Mortality Weekly Report* **2020**, *69*, 496–498. doi:10.15585/mmwr.mm6916e1.
- 514 83. Takeda, Y.; Uchiumi, H.; Matsuda, S.; Ogawa, H. Acidic electrolyzed water potently inactivates SARS-CoV-2 depending on the
- 515 amount of free available chlorine contacting with the virus. *Biochemical and Biophysical Research Communications* **2020**, *530*, 1–3.
- 516 doi:10.1016/j.bbrc.2020.07.029.
- 517 84. Chu, W.; Fang, C.; Deng, Y.; Xu, Z. Intensified Disinfection Amid COVID-19 Pandemic Poses Potential Risks to Water Quality and
- 518 Safety, 2020. doi:10.1021/acs.est.0c04394.
- 519 85. Ölmez, H.; Kretzschmar, U. Potential alternative disinfection methods for organic fresh-cut industry for minimizing water
- 520 consumption and environmental impact. *LWT - Food Science and Technology* **2009**, *42*, 686–693. doi:10.1016/j.lwt.2008.08.001.
- 521 86. Zhang, C.; Cui, F.; Zeng, G.m.; Jiang, M.; Yang, Z.z.; Yu, Z.g.; Zhu, M.y.; Shen, L.q. Quaternary ammonium compounds (QACs): A
- 522 review on occurrence, fate and toxicity in the environment, 2015. doi:10.1016/j.scitotenv.2015.03.007.
- 523 87. Melin, V.E.; Potinini, H.; Hunt, P.; Griswold, J.; Siems, B.; Werre, S.R.; Hrubec, T.C. Exposure to common quaternary ammonium
- 524 disinfectants decreases fertility in mice. *Reprod Toxicol* **2014**, *50*, 163–170. doi:10.1016/j.reprotox.2014.07.071.
- 525 88. WHO. Silver as a drinking-water disinfectant. <http://apps.who.int/bookorders.>, accessed on 2020-09-16.

- 526 89. Mijndonckx, K.; Leys, N.; Mahillon, J.; Silver, S.; Van Houdt, R. Antimicrobial silver: Uses, toxicity and potential for resistance. *BioMetals* **2013**, *26*, 609–621. doi:10.1007/s10534-013-9645-z.
- 527
- 528 90. Condell, O.; Iversen, C.; Cooney, S.; Power, K.A.; Walsh, C.; Burgess, C.; Fanning, S. Efficacy of biocides used in the modern food
529 industry to control *Salmonella enterica*, and links between biocide tolerance and resistance to clinically relevant antimicrobial
530 compounds. *Applied and environmental microbiology* **2012**, *78*, 3087–3097.
- 531 91. Capita, R.; Alonso-Calleja, C. Antibiotic-Resistant Bacteria: A Challenge for the Food Industry. *Critical Reviews in Food Science and*
532 *Nutrition* **2013**, *53*, 11–48. doi:10.1080/10408398.2010.519837.
- 533 92. Thorrold, C.A.; Letsoalo, M.E.; Dusé, A.G.; Marais, E. Efflux pump activity in fluoroquinolone and tetracycline resistant *Salmonella*
534 and *E. coli* implicated in reduced susceptibility to household antimicrobial cleaning agents. *International Journal of Food Microbiology*
535 **2007**, *113*, 315–320. doi:10.1016/j.ijfoodmicro.2006.08.008.
- 536 93. Abdel Malek, S.M.; Badran, Y.R. *Pseudomonas aeruginosa* PAO1 Adapted to 2-Phenoxyethanol Shows Cross-Resistance to
537 Dissimilar Biocides and Increased Susceptibility to Antibiotics. *Folia Microbiologica* **2010**, *55*, 588–592. doi:10.1007/s12223-010-0094-6.
- 538 94. Nomura, K.; Ogawa, M.; Miyamoto, H.; Muratani, T.; Taniguchi, H. Antibiotic susceptibility of glutaraldehyde-tolerant
539 *Mycobacterium chelonae* from bronchoscope washing machines. *American Journal of Infection Control* **2004**, *32*, 185–188.
540 doi:10.1016/j.ajic.2003.07.007.
- 541 95. Vikram, A.; Bomberger, J.M.; Bibby, K.J. Efflux as a glutaraldehyde resistance mechanism in *Pseudomonas fluorescens* and
542 *Pseudomonas aeruginosa* biofilms. *Antimicrobial Agents and Chemotherapy* **2015**, *59*, 3433–3440. doi:10.1128/AAC.05152-14.
- 543 96. Percival, S.L.; Salisbury, A.M.; Chen, R. Silver, biofilms and wounds: resistance revisited. *Critical Reviews in Microbiology* **2019**,
544 *45*, 223–237. doi:10.1080/1040841X.2019.1573803.
- 545 97. Langsrud, S.; Sundheim, G.; Borgmann-Strahsen, R. Intrinsic and acquired resistance to quaternary ammonium compounds in
546 food-related *Pseudomonas* spp. *Journal of Applied Microbiology* **2003**, *95*, 874–882. doi:10.1046/j.1365-2672.2003.02064.x.
- 547 98. García, M.; Cabo, M. Optimization of *E. coli* inactivation by benzalkonium chloride reveals the importance of quantifying the
548 inoculum effect on chemical disinfection. *Frontiers in Microbiology* **2018**, *9*. doi:10.3389/fmicb.2018.01259.
- 549 99. Kampf, G. Adaptive microbial response to low-level benzalkonium chloride exposure, 2018. doi:10.1016/j.jhin.2018.05.019.
- 550 100. Zou, L.; Meng, J.; McDermott, P.F.; Wang, F.; Yang, Q.; Cao, G.; Hoffmann, M.; Zhao, S. Presence of disinfectant resistance
551 genes in *Escherichia coli* isolated from retail meats in the USA. *Journal of Antimicrobial Chemotherapy* **2014**, *69*, 2644–2649.
552 doi:10.1093/jac/dku197.
- 553 101. Zhang, A.; He, X.; Meng, Y.; Guo, L.; Long, M.; Yu, H.; Li, B.; Fan, L.; Liu, S.; Wang, H.; Zou, L. Antibiotic and Disinfectant
554 Resistance of *Escherichia coli* Isolated from Retail Meats in Sichuan, China. *Microbial Drug Resistance* **2016**, *22*, 80–87.
555 doi:10.1089/mdr.2015.0061.
- 556 102. Gadea, R.; Fernández Fuentes, M.A.; Pérez Pulido, R.; Gálvez, A.; Ortega, E. Effects of exposure to quaternary-ammonium-based
557 biocides on antimicrobial susceptibility and tolerance to physical stresses in bacteria from organic foods. *Food Microbiology* **2017**,
558 *63*, 58–71. doi:10.1016/j.fm.2016.10.037.
- 559 103. Deng, W.; Quan, Y.; Yang, S.; Guo, L.; Zhang, X.; Liu, S.; Chen, S.; Zhou, K.; He, L.; Li, B.; Gu, Y.; Zhao, S.; Zou, L. Antibiotic
560 Resistance in *Salmonella* from Retail Foods of Animal Origin and Its Association with Disinfectant and Heavy Metal Resistance.
561 *Microbial Drug Resistance* **2018**, *24*, 782–791. doi:10.1089/mdr.2017.0127.
- 562 104. Bigliardi, P.L.; Alsagoff, S.A.L.; El-Kafrawi, H.Y.; Pyon, J.K.; Wa, C.T.C.; Villa, M.A. Povidone iodine in wound healing: A review of
563 current concepts and practices, 2017. doi:10.1016/j.ijsu.2017.06.073.