

Article

16-week supplementation with a Mediterranean diet-inspired ingredient rich in a diversity of polyphenols improves health-related quality of life of overweight and obese volunteers: a randomized, double-blind, parallel clinical trial

Cindy Romain ¹, Linda H. Chung ², Elena Marín-Cascales ², Jacobo A. Rubio-Arias ², Sylvie Gaillet ³, Caroline Laurent ³, Juana M. Morillas-Ruiz ⁴, Alejandro Martínez-Rodríguez ², Pedro E. Alcaraz ² and Julien Cases ^{2,*}

¹ Fytexia – ZAE via Europa – 3 rue d'Athènes, 34350 Vendres, France; cromain@fytextia.com; jcases@fytextia.com

² Research Center for High Performance Sport – UCAM Universidad Católica de Murcia – Murcia, Spain; lhchung@ucam.edu; emarin@ucam.edu; jararias@ucam.edu; amrodriguez@ucam.edu; palcaraz@ucam.edu

³ UMR 204 Nutripass – Institut de Recherche pour le Développement – Université de Montpellier, 34095 Montpellier, France; sylvie.gaillet-foulon@univ-montp2.fr; caroline.laurent@univ-montp2.fr

⁴ Department of Food and Nutrition Technology - Universidad Católica de Murcia – Murcia, Spain; jmmorillas@ucam.edu

* Correspondence: jcases@fytextia.com; Tel.: +33-467-219-098

Abstract: Overweight and obesity adversely affect health-related quality of life (HRQOL) through day-to-day impairments of both mental and physical functioning. It is assumed that polyphenols within the Mediterranean diet may contribute to improve HRQOL. This investigation aimed at studying the effects of a polyphenol-rich ingredient on HRQOL in overweight and obese but otherwise healthy individuals. A randomized, double-blind, placebo-controlled study including 72 volunteers was conducted. Subjects were randomly assigned to receive for a 16-week period either 900 mg/day of the supplement or a placebo. Dietary recommendations were individually determined, and intakes were recorded; daily physical mobility was monitored. Improvement of HRQOL was set as the primary outcome and assessed at baseline and at the end of the investigation, using the Short-Form 36 (SF-36) Health survey. Body composition was analyzed using dual-energy X-ray absorptiometry (DXA). Physical activity level was calculated using International Physical Activity Questionnaire (IPAQ). After 16 weeks, despite there was no adherence to the Mediterranean Diet Serving Score (MDSS), supplemented individuals experienced significant HRQOL improvement (+5.3%; $P=0.001$), including enhanced perceived physical (+11.2%; $P=0.002$) and mental health (+4.1%; $P=0.021$) components; bodily pain, vitality, and general health, being the greatest contributors. Besides, body fat mass significantly decreased (-1.2 kg; $P=0.033$), mainly within trunk area (-1.0 kg; $P=0.002$). Engagement in physical activity significantly increased (+1308 Met-min/week; $P=0.050$). Hence, chronic supplementation with a nutritional diversity and dose of a Mediterranean diet-inspired polyphenol-rich ingredient resulted in a significant amelioration in both perceived physical and mental health, concomitant with the improvement of body composition, in healthy subjects with excessive adiposity.

Keywords: Health-related quality of life; Vitality; Body composition; Phenolic compounds; Mediterranean diet

1. Introduction

The World Health Organization (WHO) defines quality of life (QOL) as “an individual’s perception of their position in life in the context of culture and value systems in which they live, and

in relation to their goals, expectations, standards, and concerns" [1]. Such conception of QOL is subjective, multidimensional, and encompasses a broad range of life domains among which health is one of the most important determinants.

In addition to be an objective medical evaluation, the health-related quality of life (HRQOL) which typically combines physical, psychological, and social domains of health [2], is more and more recognized as a central outcome in healthcare strategies, highlighting the fact that health is deeply intertwined with patient's perspective. Accordingly, the use of HRQOL assessment is particularly relevant and is increasingly widespread in clinical practices [3], predominantly for ageing populations and for the inherent expansion in the prevalence of non-communicable diseases (NCDs) that include cardiovascular diseases, diabetes mellitus, and hypertension. NCDs are currently the leading cause of mortality in the modern world, contributing to 38 million deaths each year [4]. According to WHO, the global burden of NCDs is imputable to ageing, rapid urbanization, and to globalization of unhealthy lifestyles.

Among modifiable risk factors, overweight and obesity are significant contributors to the high prevalence rate of NCDs, making them the priority target in various public health programs [5]. A major cause of overweight and obesity is the excessive accumulation of body fat due to an imbalance between energy consumption and expenditure, particularly in populations with sedentary behaviors. The accumulation of body fat, predominantly within the abdominal region, is clearly associated with a chronic low-grade inflammatory state and an impaired redox status [6], both being principal pathological pathways involved in the development of NCDs. It is noteworthy that such disorders are also identified as validated metabolic indicators of ageing [7], suggesting that overweight and obesity are conditions that could potentially accelerate ageing and exacerbate the risk of ageing-linked NCDs [8].

In addition to their role in the etiology of these common medical conditions, overweight and obesity have profound adverse physical, social, and psychological consequences that can negatively affect HRQOL, and impair everyday life [9,10], which appears to be an increasingly important outcome for patients and clinicians, and for policymakers. As a result, QOL has become a valuable endpoint assessed in both epidemiological and interventional weight management studies in overweight and obese volunteers [11-13]. Measuring HRQOL is the subjective perception of one's health, where patients are asked to rate several aspects of their life. Numerous studies have demonstrated that self-rated health measures are important predictors of mortality in various population [14].

The Short-Form 36 Health survey (SF-36), a commonly used measure in HRQOL research, is a generic methodology developed and validated in the Medical Outcomes Study [15] that assesses eight important HRQOL domains that encompass health-related social, physical, and mental dimensions. The reliability of this questionnaire has been validated both in the healthy population [16], and among people with chronic and acute health conditions, but also when comparing between different groups of patients [10]. Furthermore, several authors have demonstrated a negative correlation between BMI and SF-36 scores [17,18], and also an improvement of HRQOL correlated with weight loss in overweight people [13].

Among the dietary patterns which are mostly studied for their health effects, it appears that adherence to the Mediterranean diet has been correlated to a lower risk for NCDs [19]; it is assumed that particular bioactive constituents of the Mediterranean diet, namely polyphenols, significantly contribute to the reported health-promoted effects [20]. Moreover, along with their widely studied antioxidant properties, recent studies demonstrate the modulatory effects of phenolic compounds on various cellular signaling pathways and responses, such as inflammation and energy metabolism [21], highlighting the complex mode of their individual mechanism of action in preventing NCDs. Emerging large cohort studies that investigated the Mediterranean diet, specifically the regular consumption of a diversity of its main polyphenol content, have shown a positive correlation to HRQOL [22,23], suggesting that such bioactive constituents might be beneficial in improving overweight- and obesity-linked impaired HRQOL.

The aim of this clinical study was to investigate whether a 16-week long supplementation with an accurately characterized ingredient formulated from extracts of some fruit and vegetables

commonly consumed within the Mediterranean diet, could improve HRQOL in overweight and obese, but otherwise healthy, subjects.

2. Materials and Methods

2.1. Subjects

Ninety-two healthy overweight and obese subjects were recruited through advertisements in the region of Murcia in southern Spain. Both men and women, between the ages of 25-55 years, as well as being overweight to obese (BMI:25-40 kg/m²) but otherwise healthy, were included in the study. Subjects were excluded if they: had a metabolic and/or chronic disease; had an allergy to carrot, grape, grapefruit, green tea, caffeine, or to guarana; were involved at the time of recruitment or within the previous 6 months in a chronic supplementation program, engaged smoking cessation and or high alcohol consumption; were pregnant or were breastfeeding; were in menopause; were suffering depression; were involved in physical activity more than twice a week.

The study was approved by the UCAM (Universidad Católica San Antonio de Murcia, Spain) Ethics Committee and conducted per the guidelines laid out in the Declaration of Helsinki [24] and in compliance with Good Clinical Practices defined in the ICH Harmonized Tripartite Guideline [25]. All participants were informed about the study procedures and signed written informed consent before entering the study. This trial was registered at clinicaltrials.gov as NCT03423719.

2.2. Test supplement

Fiit-ns[®], developed by FYTEXIA (France), is principally obtained by alcohol and water extraction of grapefruit (*Citrus paradisi* Macfad), grape (*Vitis vinifera* L.) and guarana seed (*Paullinia cupana* Kunth); by water extraction of green tea (*Camellia sinensis* L. Kuntze) and black carrot (*Daucus carota* L.). Fiit-ns[®] provides bioactive compounds, specifically polyphenols from the flavonoid family, and natural components of the methylxanthine family from an extract of guarana seeds, as well as vitamin B3. The placebo product was 100% maltodextrin, which is polyphenol-, methylxanthine- and vitamin B3-free. Both Fiit-ns[®] and placebo were supplied in 450-mg capsules of identical appearance and flavor.

Supplement was analysed by means of High-Performance Liquid Chromatography (HPLC). An Agilent HPLC 1260 apparatus (software Openlab CDS chemstation edition) coupled with diode array detector was used. Separations were carried out by means of a Zorbax Stablebond SB-C18 column (4.6 x 2 mm; 5 µm particle size). To detect different phenolic classes, two different analytical method were adopted: one for bioflavonoids and caffeine and one for anthocyanins.

For flavonoids and caffeine, mobile phase A consisted in water, mobile phase B was acetic acid and mobile phase C was 100% acetonitrile. The linear gradient program was used as follows: (a) 0 to 5 min 94% A and 6% B; (b) 5 to 10 min 82.4% A, 5.6% B and 12% C; (c) 10 to 15 min 76.6% A, 5.4% B and 18% C; (d) 15 to 25 min 67.9% A, 5.1% B and 27% C; (e) 25 to 30 min 65% A, 5% B and 30% C; (f) 30 to 35 100% C; (g) 35 to 40 min 100% C; (h) 40 to 45 min 64% A and 6% B. Monitoring was performed at 280 nm at a flow rate of 1 mL/min and injection volume of 25 µL. Flavanones, flavanols, and caffeine, were respectively expressed as naringin, catechin, and caffeine.

Regarding anthocyanins, mobile phase A was water, mobile phase B consisted in formic acid and mobile phase C was acetonitrile. The gradient program used is described as follows: (a) 0 to 5 min 84.18% A, 10% B and 5.82% C; (b) 5 to 20 min 77.6% A, 10% B and 12.4% C; (c) 20 to 35 min 68.2% A, 10% B and 21.8% C; (d) 35 to 40 min 58.8% A, 10% B and 31.2% C; (e) 40 to 45 min 44.7% A, 10% B and 45.3% C; (f) 45 to 50 min 44.7% A, 10%B and 45.3% C; (g) 50 to 60 min 40% A, 10% B and 50% C; (h) 60 to 65 min 84.18% A, 10% B and 5.82% C. Monitoring was performed at 520 nm at a flow rate of 0.8 mL/min and injection volume was 10 µL. Anthocyanins were expressed as kuromanin equivalent.

Naringin, catechin, and caffeine standards were purchased from Sigma-Aldrich Co. (St. Louis, M.O., U.S.A.), and kuromanin standard from Extrasynthese (Genay, France).

2.3. Study design and interventions

The study was designed as a 16-week, randomized, double-blinded, and placebo-controlled clinical trial. Eligible participants were randomized using a simple block randomization of 1:1 with an additional stratification for sex (40% minimum and 60% maximum each sex) with a separated randomization list. Once enrolled, subjects received either the supplement (n=43) or a visually identical placebo (n=49). They were instructed to take two capsules daily for 16 weeks, one in the morning at breakfast, and one at lunchtime.

Throughout the course of the study, volunteers were instructed by a dietitian to consume a normal calorie and balanced diet corresponding to their individual needs by determining their specific Resting Energy Expenditure (REE), calculated from the revised Harris Benedict equation and adjusted per individual level of physical activity [26]. At baseline (W_1), volunteers performed a 24-hour diet recall interview corresponding to the consumption of two days during the previous week and one day during the previous weekend, in order to evaluate their usual dietary habits. At the end of the studied period, the same interview was performed to check compliance to dietary instructions. A difference of $\pm 10\%$ between reported and recommended intakes at the end of the study was considered as satisfactory. Moreover, general adherence to the Mediterranean dietary pattern was assessed using the Mediterranean Diet Serving Score (MDSS) [27]. This score ranges from 0 to 24 with an optimal cut-off point of 13.5 that discriminates adherent and non-adherent individuals.

Subjects were also encouraged to maintain their usual level of physical activity throughout the 16-week long intervention period. The subjects were provided with a pedometer (HJ-321, Omron Healthcare), which was worn at the hip, to record the physical mobility as the number of daily steps. Subjects reported their daily level of activity in a diary.

Subjects reported to the UCAM Research Center for 5 visits: (i) pre-inclusion visit at week 0 (W_0) to verify subject's eligibility, to assess anthropometrics, and to collect blood samples for the evaluation of safety parameters; (ii) baseline visit (W_1); (iii) follow-up visits (W_4 , W_8 and W_{12}); final visit (W_{16}). During each visit, subjects returned their physical activity diary and the unused investigational supplements, and they were questioned about possible occurrence of adverse events before they were provided with a new pill dispenser for the 4 following weeks.

2.4. Measurements

2.4.1. Health-related quality of life

HRQOL was measured at baseline (W_1) and at the end of the intervention period (W_{16}) using the Spanish version of the 36-item Short Form (SF-36) Health Survey [28]. This generic instrument assesses participants' self-reported HRQOL across physical and mental components. Questions pertained to the individuals' typical day during the past four weeks and usual experiences. The 36 questions were distributed across eight subscales: physical function (PF); role-physical (RP) limitations caused by physical problems; bodily pain (BP); general health (GH) perception; vitality (VT); social functioning (SF); role-emotional (RE) limitations caused by emotional problems; emotional well-being (EWB). The eight dimensions range in score from 0 to 100, with higher scores indicating better HRQOL. The SF-36 also included one Physical Component Summary (PCS) score and one Mental Component Summary (MCS) score, as well as an overall score of quality of life.

2.4.2. Body composition

At baseline (W_1) and at the end of the study period (W_{16}), body composition was assessed in the morning with volunteers in a fasted state and wearing light clothing and no shoes.

Body weight (kg) was measured with calibrated weighing scales (TBF-300MA, Tanita Corporation, I.L., U.S.A.). Waist circumference (cm) was measured at the narrowest point between the lowest rib and the iliac crest using a non-stretchable tape. The Index of Central Obesity (ICO) was calculated as the waist-to-height ratio.

Body fat mass was determined using Dual-energy X-ray Absorptiometry (DXA)-scan of the whole body (XR-46; Norland Corp., Fort Atkinson, W.I., U.S.A.). Discrimination of whole-body fat mass (FM) and body trunk fat mass (TFM) was performed with a computerized software (Software

Illuminatus DXA 4.4.0, Visual MED, Inc. and Norland CooperSurgical Company) using standardized procedures.

2.4.3. Self-reported physical activity

The self-reported International Physical Activity Questionnaire (IPAQ) instrument was used to determine global physical activity levels [29]. This self-administered long form questionnaire consisted of 27 items that covered four different domains of physical activity (working, transportation, housework, and gardening & leisure-time) that occurred during the previous seven days. The results were presented as an estimation of energy expenditure in metabolic equivalent-minutes per week (Met-min/week), and a categorical score was calculated to classify volunteers as inactive, moderately, or highly, active. Volunteers completed IPAQ questionnaire in the presence of the investigator at W_1 and W_{16} .

2.4.4. Safety parameters

Safety parameters were assessed before inclusion into the study (W_0) and at the end of the intervention period (W_{16}), in order to verify and confirm healthiness of the volunteers. Safety parameters included liver function parameters (alanine transaminase (ALT), aspartate aminotransferase (AST), gamma-glutamyltransferase (GGT)), renal function parameters (urea, creatinine, sodium (Na), potassium (K)) and heart rate.

2.5. Statistical analysis

Data sets were analysed using XLSTAT-Biomed software (Version 2017.6 for Mac, Addinsoft, Paris, France). The data are expressed as mean \pm standard deviation (SD). At baseline, distribution was considered as normal. Changes within and between groups at W_1 and W_{16} were analyzed using paired and unpaired Student's *t*-test, respectively. To compare baseline differences between the SF-36 scales and population norms, one sample *t*-test was used. A minimum value of $p \leq 0.05$ was selected as the threshold for statistical significance.

The primary outcome addressed in this study was the difference in SF-36 total score after the 16-week intervention period. The power calculation was based on previous results of a pilot study conducted with Fiit-ns® [30] [$\alpha = 0.05$, power $(1 - \beta) = 0.8$] and was performed based on an expected clinical difference in SF-36 total score between W_1 and W_{16} , within the supplemented group, of a +5% benefit minimum, to determine the targeted final sample size ($n=28$ per group). Considering a drop-out of 20% and failure rate risk of 20%, inclusion of 92 subjects was recommended.

3. Results

3.1. Characterization of the supplement

The total bioactive content corresponds to 29.27 g/100 g dry matter with a total flavonoid content measured at 24.75 g/100 g. Flavanol content corresponds to 15.67 g/100 g and includes catechin, epigallocatechin gallate, epicatechin and epicatechin gallate respectively measured at 1.47, 9.55, 2.37 and 2.28 g/100 g. Flavanone content corresponds to 8.91 g/100 g among which isonaringin, naringin, hesperidin and neohesperidin content were 0.54, 7.65, 0.03 and 0.13 g/100 g, respectively, whereas total unidentified flavanone was evaluated as naringin equivalent at 0.56 g/100 g. Total anthocyanin content corresponded 0.17 g/100 g as kuromanin equivalent. Caffeine content was measured at 4.52 g/100 g, and a third-party laboratory measured vitamin B3 content at 2.02 g/100 g (**Table 1**).

Table 1. Characterization of phenolic compounds and of caffeine, present in the supplement

Compound	Rt (min)	Λ max (nm)	Content (mg/100g)	
			Mean	SD

Caffeine	13.5	273	4523.2	450.4
Catechin	9.0	278	1472.9	85.1
Epigallocatechin Gallate	11.9	274	9549.9	282.9
Epicatechin	12.8	280	2373.0	416.6
Epicatechin Gallate	15.8	278	2279.3	372.9
Flavanone-like 1	16.1	284/323	97.6	27.7
Isonaringin	18.6	284/330	540.8	156.1
Naringin	19.5	284/330	7646.9	66.3
Hesperidin	20.3	284/328	25.4	24.3
Neohesperidin	21.2	284/328	134.2	86.8
Flavanone-like 2	27.6	284/329	263.5	4.0
Flavanone-like 3	30.4	289/328	196.4	15.3
Cyanidin-3-xylosylglucosylgalactoside	6.2	516	9.5	2.0
Cyanidin-3-xylosylgalactoside	7.9	518	39.3	7.4
Cyanidin-3-sinapoylxylosylglucosylgalactoside	11.2	531	13.8	1.5
Cyanidin-3-feruloylxylosylglucosylgalactoside	12.4	528	106.1	13.1
Cyanidin-3-pcoumarylxylosylglucosylgalactoside	14.3	528	3.0	1.6

3.2. Baseline characteristics

From the 92 individuals who were randomly allocated to either the supplement (n=43) or the Placebo (n=49), 78 subjects completed the 16-week intervention (85% of the randomly assigned subjects); after having started the intervention, a total of 14 volunteers dropped out, including 6 within the supplemented group and 8 within the placebo group, for personal reasons. Moreover, at the end of the study period, 6 subjects were excluded from final analysis because of protocol deviation, including 2 subjects within the supplemented group and 4 within the placebo group who either did not complete SF-36 questionnaire or who were non-compliant to the protocol. Finally, 72 volunteers were included in the analysis, 35 and 37 individuals in the supplemented and in the Placebo group, respectively (**Figure 1**). Baseline data of the study population are presented in **Table 2**. The two groups were similar with respect to age, height, body weight and SF-36 total score. At baseline, the placebo group had a significantly higher BMI compared with the supplemented group.




Figure 1. CONSORT flow diagram of study

Table 2. Baseline characteristics of the study population

Variable	Whole population		Placebo group		Supplemented group	
Subjects, n (M/F)	72 (34/38)		37 (16/21)		35 (18/17)	
	Mean	SD	Mean	SD	Mean	SD
Age, y	40	6	41	6	39	7
Height, m	1.69	0.09	1.68	0.08	1.70	0.11
Body weight, kg	87.0	12.6	89.3	12.2	84.5	12.6
BMI, kg/m ²	30.5	3.5	31.6	4.0	29.3*	2.5
SF-36 score, pts	81.6	9.2	81.7	8.2	81.5	10.3

* Significant at $p \leq 0.05$ level

3.3. Health-related quality of life

Regarding the whole population at baseline (Table 3), SF-36 subscales regarding vitality, emotional well-being, and mental component score, were significantly lower than the age-specific populations norms taken from the Spanish population reference values [31]. At W_1 , placebo and supplemented groups exhibited similar SF-36 scores, in both individual domains and summary scores (Table 4). After 16 weeks of supplementation, the supplemented group experienced a significant +5.3% increase ($p = 0.001$) (Figure 2) in total SF-36 score while no change was observed in the total score of the placebo group. The supplemented group showed statistically significant improvements in five out of eight domains of the health-related quality of life. Respective improvements were observed for physical component summary (PCS; +11.2%, $p = 0.002$), including

physical functioning (PF; +5.5%, $p = 0.006$), bodily pain (BP; +11.2%, $p = 0.028$), and general health (GH; +7.2%, $p = 0.010$), as well as for mental component summary, (MCS; +4.1%, $p = 0.021$) in which were included vitality (VT; +7.8%, $p = 0.006$), and emotional well-being (EWB; +5.2%, $p = 0.021$). No statistically significant changes were shown within the placebo group after the 16 week-intervention.

Table 3. Mean score on SF-36 eight subscales at baseline in the study and in the Spanish population ages 25-54

Scale	Baseline	Age specific Population Norm [§]
Physical component score	83.8	83.0
Physical functioning	89.5	91.9
Role-physical	93.8	87.5
Bodily pain	79.0	82.6
General health	73.0	72.7
Mental component score	79.4 *	82.3
Role-emotional	92.4	91.3
Vitality	62.8 *	70.2
Emotional well-being	70.6 *	74.9
Social functioning	91.8	92.9

[§] Combined score for men and women ages 25-54 as the age-specific population norm

* A significant difference at baseline from the age-specific population norm for the scale at $p \leq 0.05$ level

Table 4. Mean score on SF-36 eight subscales, both physical and mental components, and total health-related quality of life at baseline (W₁) and at the end of the study (W₁₆)

Scale	Placebo group				Supplemented group			
	W1		W16		W1		W16	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
SF-36 Total score	81.7	8.2	82.2	8.2	81.5	10.3	85.8 ^{a*}	6.4
Physical component score	84.3	7.6	83.8	8.9	83.4	12.6	88.7 ^{a*}	7.6
Physical functioning	88.6	9.4	90.6	8.4	90.4	12.7	95.4 ^{a*}	7.0
Role-physical	94.3	7.3	93.6	8.1	93.4	11.6	95.7	6.6
Bodily pain	80.7	18.0	78.3	23.3	77.1	22.9	85.8 *	14.5
General health	73.5	13.3	72.8	11.7	72.5	16.0	77.8 *	13.7
Mental component score	79.1	10.8	80.7	10.5	79.7	10.7	82.9 *	8.0
Role-emotional	91.2	14.8	93.5	12.6	93.6	12.0	94.9	9.2
Vitality	62.4	11.1	63.1	12.2	63.1	13.4	68.0 ^{a*}	11.1
Emotional well-being	70.9	11.1	73.5	10.9	70.3	12.9	73.9 *	10.0
Social functioning	91.9	15.9	92.6	14.9	91.8	12.1	95.4	9.1

* An intragroup difference between baseline (W₁) and end of the study (W₁₆) at $p \leq 0.05$ level

^a An intergroup difference at the end of the study (W₁₆) at $p \leq 0.05$ level




Figure 2. % change from baseline (W₁) to the end of the study (W₁₆) in individual SF-36 domains as well as PCS (Physical Component Score), MCS (Mental Component Score), and total HRQOL score in placebo and in supplemented subjects. Arrows indicate clinically significant difference according to Samsa *et al.* [31]. * indicates an intragroup difference between baseline (W₁) and end of the study (W₁₆) at $p \leq 0.05$ level. PF: Physical Functioning; RP: Role Physical; BP: Bodily Pain; GH: General Health; RE: Role Emotional; VT: Vitality; EWB: Emotional Well-Being; SF: Social Functioning.

3.4. Body composition

At baseline, all fat mass-related variables (FM, TFM, Index of Central Obesity (ICO) and BMI) were significantly higher in the placebo group (Table 5). Such a discrepancy is explained by the higher number of obese individuals that completed the clinical investigation in the placebo group. After 16 weeks of supplementation, volunteers from the placebo group did not experience any significant change in body composition. The supplemented group showed an improvement in anthropometrics after 16 weeks with a statistically significant decrease in body weight by -1.3 kg ($p = 0.013$) and in BMI by -0.4 points ($p = 0.012$). Waist size significantly decreased by -1.1 cm ($p = 0.017$), consequently lowering the ICO by -1.3% ($p = 0.018$). Supplemented volunteers significantly lost -1.2 kg of FM ($p = 0.033$) of which -1.0 kg was fat lost only from TFM ($p = 0.002$).

Table 5. Body weight, BMI, waist circumference, ICO, total body fat mass and total trunk fat mass at baseline (W₁) and at the end of the study (W₁₆)

	Placebo group				Supplemented group			
	W1		W16		W1		W16	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Body weight, kg	89.3	12.2	89.1	12.5	84.5	12.6	83.2 ^{b*}	12.6
BMI, kg/m ²	31.6	4.0	31.5	3.9	29.3 ^a	2.5	28.9 ^{b*}	2.8
Waist circumference, cm	94.2	9.7	94.6	9.3	90.7	8.8	89.6 ^{b*}	9.3
ICO	0.560	0.05	0.563	0.05	0.533 ^a	0.04	0.526 ^{b*}	0.04
Total body fat mass, kg	35.2	10.4	35.2	10.8	30.5 ^a	7.4	29.3 ^{b*}	7.9
Total trunk fat mass, kg	18.1	5.4	18.1	5.9	15.7 ^a	4.1	14.7 ^{b*}	4.2

* An intragroup difference between baseline (W₁) and end of the study (W₁₆) at $p \leq 0.05$ level

^{a,b} An intergroup difference at baseline (W₁) and at the end of the study (W₁₆) respectively, at $p \leq 0.05$ level

3.5. Self-reported physical activity and average daily steps recording

At baseline, both groups showed similar self-reported level of physical activity. While it did not significantly change within the placebo population ($p = 0.280$), the supplemented subjects showed an increase of +1308 Met-min/week ($p = 0.05$) after 16 weeks of supplementation (**Table 6**). Regarding categorical scores at baseline, the rate of volunteers within each category (i.e., inactive, moderately active, and highly active) was similar between groups. After 16 weeks, the rate of inactive people remained the same in both groups; within the placebo group the rate of highly active subjects decreased by -43% while the rate of moderately active individuals increased by +14%. In contrast, within the supplemented population the rate of moderately active subjects decreased by -14% but the number of highly active individuals increased by +43%. The number of average daily steps was significantly different at baseline between placebo and supplemented subjects ($p = 0.028$). The placebo group did not experience any significant change in average daily steps monitored after 16 weeks while the supplemented subjects significantly decreased their average rate by -678 steps ($p = 0.019$) to reach a similar level than to the placebo population.

Table 6. Mean total score of self-reported physical activity (IPAQ) and daily number of steps (pedometer) at baseline (W₁) and at the end of the study (W₁₆)

	Placebo group				Supplemented group			
	W1		W16		W1		W16	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
IPAQ score (Met-min/week)	4798	4740	4231	4190	4766	4721	6074 [*]	6631
Inactive (%)	15.2		15.2		12.5		12.5	
Moderately active (%)	63.6		72.7		65.6		56.3	
Highly active (%)	21.2		12.1		21.8		31.2	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Daily steps	6770	2239	7186	2679	8169 ^a	2797	7491 [*]	2964

* An intragroup difference between baseline (W₁) and end of the study (W₁₆) at $p \leq 0.05$ level

^a An intergroup difference at baseline (W₁), at $p \leq 0.05$ level

3.6. Recommended and reported dietary intake

Recommended intake at baseline did not differ between the two groups ($p = 0.770$) (**Table 7**). When recommended intake was compared with reported intake at baseline, the difference was -13.7% and -7.8% for the placebo and for the supplemented groups, respectively. After 16 weeks, the difference between recommended and reported intake in both groups was lower than 10% (-8.8% and -9.0% for the placebo and for the supplemented groups, respectively). Mediterranean Diet Serving Scores (MDSS) were similar between both groups, 8.4 and 8.6 for placebo and supplemented population, respectively, indicating a non-adherence to the Mediterranean diet pattern during the intervention period.

Table 7. Recommended and reported dietary intake at baseline (W_1) and at the end of the study (W_{16})

	Placebo group				Supplemented group			
	W_1		W_{16}		W_1		W_{16}	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Recommended intake (Kcal)	2074	273	2084	281	2096	360	2039	342
Reported intake (Kcal)	1789	471	1899 *	502	1933	463	1855	392
MDSS score	Mean 8.4		SD 3.7		Mean 8.6		SD 4.2	

* An intragroup difference between baseline (W_1) and end of the study (W_{16}) at $p \leq 0.05$ level

3.7. Safety

After 16 weeks, both liver and renal function parameters were within the healthy range in both groups, suggesting that no health impairment occurred throughout the course of the study. Moreover, heart rate stayed stable throughout the course of the study. No adverse events, nor side effects, linked to the supplement were reported during the course of the study.

Table 8. Clinical safety values at baseline (W_1) and at the end of the study (W_{16})

Parameters (normal range)	Placebo group				Supplemented group			
	W_1		W_{16}		W_1		W_{16}	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Liver function								
ALT (7-55 U/L)	21.4	9.1	21.0	8.5	25.0	15.9	25.1	14.7
AST (8-48 U/L)	20.2	5.5	20.1	5.3	22.4	6.3	23.3	10.0
GGT (6-48 U/L)	19.1	11.7	19.9	11.9	23.1	13.4	24.7	14.0
Kidney function								
Urea (15-46 mg/dL)	35.4	9.1	33.0	7.7	31.9	7.8	30.1	7.4
Creatinine (0.6-1.3 mg/dL)	0.79	0.18	0.74 *	0.16	0.74	0.15	0.76	0.16
Na (135-145 mmol/L)	141.2	1.2	141.2	2.1	141.7	1.7	141.1	1.7
K (3.6-5.2 mmol/L)	4.3	0.3	4.3	0.2	4.3	0.2	4.3	0.3
Heart rate (bpm)	71.8	10.2	72.1	8.6	71.3	11.2	70.4	16.4

* An intragroup difference between baseline (W_1) and end of the study (W_{16}) at $p \leq 0.05$ level

4. Discussion

The main results of this study demonstrate that a 16-week long supplementation period with an ingredient formulated with a blend of various botanical extracts, rich in a diversity of polyphenols, that is usually consumed as part of the typical Mediterranean diet, is associated with significant improvements of both the physical and the mental components of the HRQOL, in healthy overweight and obese, but otherwise healthy, subjects of both sex.

At baseline, volunteers showed an impaired HRQOL, namely in vitality and emotional well-being subscales, for which values were below the Spanish age-specific population reference norms [31]. Although similar studies have previously reported impairment across all of the SF-36 subscales, most of them were conducted either with a population displaying a significantly higher grade of obesity or with an additional manifestation of comorbidities [10,17]. Here, baseline impairments observed for vitality and emotional well-being are in line with the work of Blissmer *et al.* [11] who have found similar decrements in a highly comparable population of healthy overweight and moderately obese subjects, indicating higher feelings of tiredness and anxiety.

Following a 16-week intervention period associated with a normal-caloric diet, both the physical and the mental components of the HRQOL significantly improved in volunteers supplemented with the polyphenol-rich ingredient compared to the placebo group. Net improvements were seen by subjective ameliorations in bodily pain > general health > vitality > physical functioning > emotional well-being. It is noteworthy that after the 16-week long period of supplementation, both the vitality and the emotional well-being values improved to achieve the level of the reference norms of the Spanish age-specific population. Improvements in these different subscales and in both the physical- and the mental-component scores must be considered as clinically significant, as it has been stated that absolute differences of 3-5 points is clinically relevant [32].

In addition to these improvements, the 16-week chronic polyphenolic supplementation induced a significant body weight loss with an average difference between both groups of 1.1 kg. It is noteworthy that this decrease is essentially driven by an 86% fat mass reduction for which 89% is located within the trunk area, pointing out a particularly beneficial effect on body composition. Such improvement may, in some extent, positively impact the HRQOL. Indeed, some authors have demonstrated that weight loss was associated with improvement of both physical and mental health dimensions in several intervention trials [11,13,33-35]. Moreover, the amount of weight loss and the level of HRQOL improvement may be directly interconnected [35,36]. Nevertheless, here, we did not demonstrate a significant correlation between weight or fat loss and HRQOL improvement (data not included), hypothesizing that weight loss could be an indirect consequence of HRQOL improvement, as it has been recently demonstrated with a bi-directional relationship between both parameters [37]. Moreover, catechins from green tea, have previously demonstrated anti-obesity effects [38] through various mechanisms of action such as the inhibition of pancreatic lipase [39] as well as through the regulation of obesity-related genes and proteins [40]. However, it is important to highlight that these interventional studies used significantly higher amounts of green tea catechins; whereas in the current supplement it only corresponded to one cup of green tea daily. Moreover, caffeine content and flavanones from grapefruit extract could also potentiate the decrease in body fat mass, as it has previously been demonstrated an enhanced lipolysis conducting to body weight and fat mass decrease in overweight and obese subjects supplemented with such kind of bioactive compounds [41].

As body composition improved, the level of physical activity, assessed through the IPAQ questionnaire, significantly intensified ($p = 0.05$) after 16 weeks of supplementation. Thus, while 69% of volunteers from supplemented group maintained their usual category of physical activity, 25% moved into a higher category compared with the placebo population, for which only 9% of volunteers improved their level of physical activity. Contradictorily, at the same time, the supplemented group showed a decrease in daily steps as assessed with a pedometer, while no change was observed in the placebo population. This discrepancy may be explained by the fact that pedometer is not suitable for the measurement of certain types of physical activity such as swimming, cycling or heavy lifting, that are otherwise assessed through the IPAQ questionnaire, making these both subjective and objective measurements, two complementary tools in physical activity assessment. As volunteers were encouraged to maintain their usual physical activity level throughout the course of the study, it can be hypothesized that the significantly higher physically active lifestyle reported within the supplemented group is not the result of a conditioned mental engagement only. Indeed, the increase of HRQOL, and namely of the feeling of more vitality, may explain such a rise on physical activity. A recent review that aimed at examining the link between physical activity and HRQOL concluded that there is a consistent cross-sectional association between physical activity level and HRQOL,

namely in the vitality and in the physical functioning domains, but the finding could not confirm a causal relationship, *i.e.*, “higher HRQOL leading to higher level of physical activity, or vice versa, or mutual influence” [42]. Nevertheless, the engagement in a more active lifestyle within the supplemented population may also have a positive effect on body composition improvement, as discussed above.

Besides positive effects on body composition and engagement in physical activity, it appears that phenolic compounds may induce, through other various mechanisms, observable effects on HRQOL improvement. Accordingly, adherence to a Mediterranean dietary pattern, characterized by wide consumption of fruit & vegetables, cereals, fish, olive oil and red wine, has been directly associated with better QOL in an analysis including more than 11000 participants that belonged to the SUN cohort [43,44]. While several nutrients and micronutrients may contribute to this effect, phenolic compounds have been suggested to be the main mediators; a large cross-sectional study demonstrated a direct relationship between antioxidant content of the Mediterranean diet, including flavonoid content, and HRQOL [23]. In addition, in another recent study including more than 13000 women, higher flavonoid intake at midlife was associated with increased odds of healthy ageing, based on higher survival at older ages free of chronic diseases and maintenance of midlife HRQOL (assessed by the SF-36 survey) [45]. Here, despite that the studied population is Spanish and is particularly prone to comply with the Mediterranean diet, the MDSS did not demonstrate any significant adherence to this pattern in both groups for which, namely, the consumption of fruit and vegetables, main sources of flavonoids, was below the recommendations of the last updated version of the Mediterranean Diet Pyramid [46]. Thus, it can be hypothesized that a regular basic diet has no or only minor impact on HRQOL, since there were no improvements within the placebo population, whereas the supplemented subjects that covered the gap of phenolic micronutrients significantly improved their HRQOL.

Bioactive compounds occurring in the supplement may positively impact physiological functions related to both physical and mental health status, mainly vascular inflammation, coagulation factors, and endothelial function [23], that are all described to be impaired during overweight/obesity [47]. The aptitude of certain phenolic compounds in improving vascular health has been demonstrated both *in vitro* and *in vivo* [48]. Catechins from green tea positively impact vascular functions through various complementary mechanisms linked to their anti-oxidative and anti-inflammatory properties, as well as their capacity in activating endothelial NO synthase [49]. Similarly, grape polyphenols also demonstrated an aptitude in improving vascular impairments through similar molecular mechanisms [50], all contributing to a better peripheral and central blood flow that in turn may positively affect physical and mental health status [51].

While modulation of both oxidative stress and inflammatory parameters, main contributors in the improvement of vascular function and blood flow, has previously been demonstrated with the current supplement in a study involving obese subjects [30], specific mechanisms in flow-mediated dilation improvements and subsequent blood flow increase have not yet been investigated. Moreover, as the beneficial effect of the supplement on HRQOL has been demonstrated, further investigations will have to be conducted in attempts to confirm the causal relationship between bioavailability and pharmacokinetics of the polyphenols metabolites and the mechanisms involved to improve vascular function.

Beyond the limitations addressed, results of the present study reveal the beneficial and systemic effects of phenolic compounds on subjective physical and mental symptoms linked to overweight/obesity. The study was designed to minimize bias, and thus, individualized calorie intake recommendations and diet interviews, as well as monitoring of daily steps, were identified as possible confounding factors. Despite that the studied population were recruited in a Mediterranean region, none of the two groups adhered to the typical regional diet, which strengthens the hypothesis that phenolic compounds certainly contribute to subjective health as previously proposed by others [23,45].

In conclusion, this study demonstrated that the 16-week long consumption of an ingredient obtained from polyphenol-rich fruit and vegetables extracts is associated with improvements in HRQOL, specifically in both mental and physical subjective feelings. In addition, the decrease in body

fat mass and the significantly increased engagement in physical activity, probably established a virtuous cycle between body composition, physical activity, and perceived HRQOL. Mechanisms of action likely involve improvements in vascular function via well-known anti-oxidative and anti-inflammatory properties of phenolic compounds. Such beneficial effects may be extended to other situations where HRQOL is impaired, particularly during the ageing process where an imbalance of body composition, a loss of vitality, and of physical functioning, associated to a more sedentary lifestyle are commonly observed.

Author Contributions: Conceptualization, C.R. and J.C.; Methodology, P.E.A. and L.H.C.; investigation, P.E.A., L.H.C., J.A.R.A., E.M.C., J.M.M.R. and A.M.R.X.X.; writing—original draft preparation, C.R.; writing—review and editing, all authors..

Funding: This research received no external funding

Conflicts of Interest: Fytexia is involved in the research & development and marketing & sales of polyphenol extracts-based ingredients for food and nutraceutical industries. Therefore, Fytexia has a commercial interest in this publication. UCAM and UMR 204 Nutripass were paid by Fytexia to perform and report the scientific work that formed the basis of this publication. Fytexia, UCAM, UMR 204 Nutripass and all authors declare that the data in this report represent a true and faithful representation of the work that has been performed. The financial assistance of Fytexia is gratefully acknowledged.

References

1. Whoqol Group. The World Health Organization Quality of Life assessment (WHOQOL): position paper from the World Health Organization. *Soc Sci Med* **1995**, *41*, 1403–1409.
2. Testa, M.; Simonson, D. Current concepts: Assessment of quality-of-life outcomes. *N Eng J Med* **1996**, *334*, 835–840.
3. Fayers, P.M.; Machin, D. Quality of Life: Assessment, Analysis and Interpretation. Chichester: John Wiley and Sons, 2000.
4. WHO. Noncommunicable diseases. World Health Organization. Available online: <http://www.who.int/mediacentre/factsheets/fs355/en/> (accessed March 2018)
5. Gortmaker, S.L.; Swinburn, B.A.; Levy, D.; Carter, R.; Mabry, P.L.; Finegood, D.; Huang, T.; Marsh, T.; Moodie, M. Changing the future of obesity: science, policy, and action. *Lancet* **2011**, *378*, 838–847.
6. Fernández-Sánchez, A.; Madrigal-Santillán, E.; Bautista, M.; Esquivel-Soto, J.; Morales-González, A.; Esquivel-Chirino, C.; Durante-Montiel, I.; Sánchez-Rivera, G.; Valadez-Vega, C.; Morales-González, J.A. Inflammation, oxidative stress, and obesity. *Int J Mol Sci* **2011**, *12*, 3117–3132.
7. Rahman, I.; Bagchi, D. Inflammation, advancing age and nutrition. Rahman I, Bagchi D, editors. Academic Press **2013**.
8. Ahima, R.S. Connecting obesity, aging and diabetes. *Nat Med* **2009**, *15*, 996–997.
9. Fontaine, K.R.; Barofsky, I. Obesity and health-related quality of life. *Obes Rev* **2001**, *2*, 173–182.
10. Slagter, S.N.; Van Vliet-Ostapchouk, J.V.; Van Beek, A.P.; Keers, J.C.; Lutgers, H.L.; van der Klauw, M.M.; Wolffenbuttel, B.H.R. Health-Related quality of life in relation to obesity grade, type 2 diabetes, metabolic syndrome and inflammation. *PLoS One* **2015**, *10*, 1–17.
11. Blissmer, B.; Riebe, D.; Dye, G.; Ruggiero, L.; Greene, G.; Caldwell, M. Health-related quality of life following a clinical weight loss intervention among overweight and obese adults: Intervention and 24 month follow-up effects. *Health Qual Life Outcomes* **2006**, *4*, 43.
12. Ford, E.S.; Moriarty, D.G.; Zack, M.M.; Mokdad, A.H.; Chapman, D.P. Self-reported body mass index and health-related quality of life: findings from the Behavioral Risk Factor Surveillance System. *Obes Res* **2001**, *9*, 21–31.
13. Fine, J.T. A Prospective Study of Weight Change and Health-Related Quality of Life in Women. *Jama* **1999**, *282*, 2136.
14. Idler, E.L.; Benyamini, Y. Self-rated health and mortality: a review of twenty-seven community studies. *J Health Soc Behav* **1997**, *38*, 21–37.

15. Ware, J.E.; Snow, K.K.; Kosinski, M.; Gandek, B. SF-36 Health Survey Manual and Interpretation Guide. Bost New Engl Med Cent **1993**.
16. Obidoa, C.; Reisine, S.; Cherniack, M. How does the SF 36 perform in healthy populations? A structured review of longitudinal studies. *J Soc Behav Heal* **2010**, *4*, 30–48.
17. Doll, H.A.; Petersen, S.E.; Stewart-Brown, S.L. Obesity and physical and emotional well-being: associations between body mass index, chronic illness, and the physical and mental components of the SF-36 questionnaire. *Obes Res* **2000**, *8*, 160–170.
18. Marchesini, G.; Solaroli, E.; Baraldi, L.; Natale, S.; Migliorini, S.; Visani, F.; Forlani, G.; Malchionda, N. Health-related quality of life in obesity: the role of eating behaviour. *Diabetes Nutr Metab* **2000**, *13*, 156–164.
19. Caretto, A.; Lagattolla, V. Non-communicable diseases and adherence to Mediterranean diet. *Endocr Metab Immune Disord Drug Targets* **2015**, *15*, 10–7.
20. Tresserra-Rimbau, A.; Rimm, E.B.; Medina-Remón, A.; Martínez-González, M.A.; López-Sabater, M.C.; Covas, M.I.; Corella, D.; Salas-Salvadó, J.; Gómez-Gracia, E.; Lapetra, J.; Arós, F.; Fiol, M.; Ros, E.; Serra-Majem, L.; Pintó, X.; Muñoz, M.A.; Gea, A.; Ruiz-Gutiérrez, V.; Estruch, R.; Lamuela-Raventós, R.M.; and on behalf of the PREDIMED Study Investigators. Polyphenol intake and mortality risk: a re-analysis of the PREDIMED trial. *BMC Med* **2014**, *12*, 77.
21. Crozier, A.; Jaganath, I.B.; Clifford, M.N. Dietary phenolics: chemistry, bioavailability and effects on health. *Nat Prod Rep* **2009**, *26*, 1001–1043.
22. Veronese, N.; Stubbs, B.; Noale, M.; Solmi, M.; Luchini, C.; Maggi, S. Adherence to the Mediterranean diet is associated with better quality of life: Data from the Osteoarthritis Initiative. *Am J Clin Nutr* **2016**, *104*, 1403–1409.
23. Bonaccio, M.; Di Castelnuovo, A.; Bonanni, A.; Costanzo, S.; De Lucia, F.; Pounis, G.; Zito, F.; Donati, M.B.; de Gaetano, G.; Iacoviello, L.; on behalf of the Moli-sani project investigators. Adherence to a Mediterranean diet is associated with a better health-related quality of life: a possible role of high dietary antioxidant content. *BMJ Open* **2013**, *3*, 1–11.
24. World Health Organisation. Declaration of Helsinki World Medical Association Declaration of Helsinki Ethical Principles for Medical Research Involving Human Subjects. *J Am Med Assoc* **2013**, *310*, 2191–4.
25. Vijayanathan, A.; Nawawi, O. The importance of Good Clinical Practice guidelines and its role in clinical trials. *Biomed Imaging Interv* **2008**, *4*, e5.
26. Roza, A.; Shizgal, H. The Harris Benedict energy requirements equation reevaluated : resting and the body cell mass. *Am J Clin Nutr* **1984**, *40*, 168–182.
27. Monteagudo, C.; Mariscal-Arcas, M.; Rivas, A.; Lorenzo-Tovar, M.; Tur, J.A.; Olea-Serrano, F. Proposal of a Mediterranean Diet Serving Score. *PLoS One* **2015**, *10*, e0128594.
28. Vilagut, G.; Ferrer, M.; Rajmil, L.; Rebollo, P.; Permanyer-Miralda, G.; Quintana, J.M.; Santed, R.; Valderas, J.M.; Ribera, A.; Domingo-Salvany, A.; Alonso, J. The Spanish version of the Short Form 36 Health Survey: a decade of experience and new developments. *Gac Sanit* **2005**, *19*, 135–150.
29. Craig, C.L.; Marshall, A.L.; Sjöström, M.; Bauman, A.E.; Booth, M.L.; Ainsworth, B.E.; Pratt, M.; Ekelund, U.; Yngve, A.; Sallis, J.F.; Oja, P. International physical activity questionnaire: 12-Country reliability and validity. *Med Sci Sports Exerc* **2003**, *35*, 1381–1395.
30. Cases, J.; Romain, C.; Dallas, C.; Gerbi, A.; Cloarec, M. Regular consumption of Fiitns, a polyphenol extract from fruit and vegetables frequently consumed within the Mediterranean diet, improves metabolic ageing of obese volunteers: a randomized, double-blind, parallel trial. *Int J Food Sci Nutr* **2015**, *66*, 120–125.
31. Alonso, J.; Regidor, E.; Barrio, G.; Prieto, L. Valores poblacionales de referencia de la versión española del Cuestionario de Salud SF 36. *Med Clin (Barc)* **1998**, *36*, 1–10.
32. Samsa, G.; Edelman, D.; Rothman, M.L.; Williams, G.R.; Lipscomb, J.; Matchar, D. Determining clinically important differences in health status measures: a general approach with illustration to the Health Utilities Index Mark II. *Pharmacoeconomics* **1999**, *15*, 141–155.
33. Rippe, J.M.; Price, J.M.; Hess, S.A.; Kline, G.; DeMers, K.A.; Damitz, S.; Kreidieh, I.; Freedson, P. Improved psychological well-being, quality of life, and health practices in moderately overweight women participating in a 12-week structured weight loss program. *Obes Res* **1998**, *6*, 208–218.
34. Kolotkin, R.L.; Fujioka, K.; Wolden, M.L.; Brett, J.H.; Bjorner, J.B. Improvements in health-related quality of life with liraglutide 3.0mg compared with placebo in weight management. *Clin Obes* **2016**, *6*, 233–242.
35. Ross, K.M.; Milsom, V.A.; Rickel, K.A.; DeGraganza, N.; Gibbons, L.M.; Murawski, M.E.; Perri, M.G. The Contributions of Weight Loss and Increased Physical Fitness to Improvements in Health-Related Quality of Life. *Eat Behav* **2009**, *10*, 84–88.

36. Samsa, G.P.; Kolotkin, R.L.; Williams, G.R.; Nguyen, M.H.; Mendel, C.M. Effect of moderate weight loss on health-related quality of life: an analysis of combined data from 4 randomized trials of sibutramine vs placebo. *Am J Manag Care* **2001**, *7*, 875–883.
37. Cameron, A.J.; Magliano, D.J.; Dunstan, D.W.; Zimmet, P.Z.; Hesketh, K.; Peeters, A.; Shaw, J.E. A bi-directional relationship between obesity and health-related quality of life: evidence from the longitudinal AusDiab study. *Int J Obes* **2012**, *36*, 295–303.
38. Suzuki, T.; Pervin, M.; Goto, S.; Isemura, M.; Nakamura, Y. Beneficial effects of tea and the green tea catechin epigallocatechin-3-gallate on obesity. *Molecules* **2016**, *21*, 5–8.
39. Grove, K.A.; Sae-Tan, S.; Kennett, M.J.; Lambert, J.D. (–)-Epigallocatechin-3-gallate inhibits pancreatic lipase and reduces body weight gain in high fat-fed obese mice. *Obesity* **2012**, *20*, 2311–2333.
40. Yang, C.S.; Zhang, J.; Zhang, L.; Huang, J.; Wang, Y. Mechanisms of body weight reduction and metabolic syndrome alleviation by tea. *Mol Nutr Food Res* **2016**, *60*, 160–174.
41. Dallas, C.; Gerbi, A.; Elbez, Y.; Caillard, P.; Zamaria, N.; Cloarec, M. Clinical study to assess the efficacy and safety of a citrus polyphenolic extract of red orange, grapefruit, and orange (Sinetrol-xpur) on weight management and metabolic parameters in healthy overweight individuals. *Phyther Res* **2014**, *28*, 212–218.
42. Bize, R.; Johnson, J.A.; Plotnikoff, R.C. Physical activity level and health-related quality of life in the general adult population: a systematic review. *Prev Med (Baltim)* **2007**, *45*, 401–415.
43. Ruano, C.; Henriquez, P.; Martínez-González, M.Á.; Bes-Rastrollo, M.; Ruiz-Canela, M.; Sánchez-Villegas, A. Empirically Derived Dietary Patterns and Health-Related Quality of Life in the SUN Project. *PLoS One* **2013**, *8*, 1–10.
44. Henríquez Sánchez, P.; Ruanon C.; de Irala, J.; Ruiz-Canela, M.; Martínez-González, M.Á.; Sánchez-Villegas, A. Adherence to the Mediterranean diet and quality of life in the SUN Project. *Eur J Clin Nutr* **2012**, *66*, 360–368.
45. Samieri, C.; Sun, Q.; Townsend, M.K.; Rimm, E.B.; Grodstein, F. Dietary flavonoid intake at midlife and healthy aging in women. *Am J Clin Nutr* **2014**, *100*, 1489–97.
46. Bach-Faig, A.; Berry, E.M.; Lairon, D.; Reguant, J.; Trichopoulos, A.; Dernini, S.; Medina, F.; Battino, M.; Belahsen, R.; Miranda, G.; Serra-Majem, L.; Aranceta, J.; Atinmo, T.; Barros, J.; Benjelloun, S.; Bertomeu-Galindo, I.; Burlingame, B.; Cballero-Bartoli, M.; Clapés-Badrinas, C.; Couto, S.; Elmadfa, I.; Estruch, R.; Faig, A.; Fidanza, F.; Franceschi, S.; Hautvast, J.; Helsing, E.; Julia-Llobet, D.; La Vecchia, C.; Lemtouni, A.; Mariné, A.; Martinez-Gonzalez, M.; Mokni, R.; Mombiola, F.; Noain, I.; Obrador, B.; Pekcan, G.; Piscopo, S.; Raido-Quintana, B.; Ros, E.; Saez-Almendros, S.; Salas-Salvado, J.; Sensat, F.; Trichopoulos, D.; Tur, J.; Vaz Da Almeida, M.; Willett, W.; Amiot-Carlin, M.; Bellio, A.; Cannella, C.; Capone, R.; Cassi, D.; Donini, L.; Lacirignola, C.; Maiani, G.; Mancini, M.; Merendino, N.; Padilla, M.; Padulosi, S. Mediterranean diet pyramid today. Science and cultural updates. *Public Health Nutr* **2011**, *14*, 2274–2284.
47. Perticone, F.; Ceravolo, R.; Candigliota, M.; Ventura, G.; Iacopino, S.; Sinopoli, F.; Mattioli, P.L. Obesity and body fat distribution induce endothelial dysfunction by oxidative stress: protective effect of vitamin C. *Diabetes* **2001**, *50*, 159–165.
48. De Pascual-Teresa, S.; Moreno, D.A.; García-Viguera, C. Flavanols and anthocyanins in cardiovascular health: a review of current evidence. *Int J Mol Sci* **2010**, *11*, 1679–1703.
49. Moore, R.J.; Jackson, K.G.; Minihane, A.M. Green tea (*Camellia sinensis*) catechins and vascular function. *Br J Nutr* **2009**, *102*, 1790–1802.
50. Dohadwala, M.M.; Vita, J.A. Grapes and Cardiovascular Disease. *J Nutr* **2009**, *139*, 1788S–1793S.
51. Watson, R.R.; Preedy, V.R.; Zibadi, S. Polyphenols in human health and disease. **2014**, Elsevier, Vol. 1.