

Review

Thyroid dysfunction: risk and management in dentistry

Georges Aoun

Department of Oral Medicine and Maxillofacial Radiology, Faculty of Dental Medicine, Lebanese University, ORCID ID: <https://orcid.org/0000-0001-5073-6882>

Correspondence: aoungorges@yahoo.com

Abstract: The thyroid gland secretes essential hormones for the growth regulation and the metabolic activity of the body. Thyroid dysfunction includes many conditions that can indirectly affect dental treatments. Therefore, the practitioner must be aware of these pathologies and their signs and symptoms in order to elaborate specific dental treatment plans, thus avoiding any per- and/or postoperative complications. Moreover, the dentist can help in the screening of undiagnosed thyroid dysfunction among patients.

Keywords: Dentistry; hyperthyroidism; hypothyroidism; management; oral manifestations; thyroid gland.

1. Introduction

The thyroid is an endocrine gland located on the anterior midline of the neck below the inferior part of the larynx at the superior segment of the trachea. It consists of two oval shaped lobes connected by an isthmus. The mean lobe dimensions and weight are 5 x 2.5 x 2.5 cm and 25 mg, respectively. Microscopically, the thyroid gland is composed of spherical follicles bordered by follicular cells and containing a substance called colloid which is thyroglobulin. This thyroglobulin along with the circulating iodine form monoiodotyrosine and diiodotyrosine, precursors of the thyroid hormones, tetraiodothyronine known as thyroxine (T4) and triiodothyronine (T3) (1).

Between the thyroid follicles or inside their wall are located multiple small cells, the C-cells (parafollicular cells), responsible for another thyroid hormone known as calcitonin (1).

The secretion of thyroid hormones, mainly T3 and T4, is directly under the control of the hypothalamus via the TSH releasing hormone –TRH and the pituitary gland via the thyroid stimulating hormone –TSH. This is done through a negative feedback mechanism (2).

As for calcitonin secretion, it is mostly stimulated by the increase of serum calcium concentration (3).

The thyroid hormones are essential for the metabolic activity of all the body; they are essential to the normal functioning of most of the organs. Some of these functions are summarized in **Table 1**.

They assist in the growth, development, and differentiation of all the body cells (4, 5).
They are a key regulator of the basal metabolic rate (BMR) which is one of the major components of energy expenditure. In humans, BMR decrease can lead to obesity (1, 6).
They play an important role in calcium homeostasis (calcitonin which is stimulated by hypercalcemia, fights against it) (1, 3), and bone mineral homeostasis and density (deposition of calcium and phosphate in the bones) (1).
They are vital for the development of central nervous system in children and brain maturation and function for life (1, 7, 8).
They stimulate somatic and psychic growth (1) and influence the mood and behavior (mental and behavioral disorders may occur with thyroid hormones imbalance) (9).
They play a main role in the normal function of the heart (rate and contraction) and vascular physiology (10, 11).
They control carbohydrate, fat, protein, and vitamins metabolism (12).
They regulate the body temperature (increased level of thyroid hormones rises body temperature, and the opposite is true) (1).
They play a vital role in maintaining electrolyte balance (1).
They increase oxygen consumption by the cells and tissues (13).
They stimulate gastrointestinal motility (14).
They improve the sensitivity of the β -adrenergic receptors to catecholamines (1).
They improve mitochondrial metabolism (1).
They regulate erythropoiesis (15).

Table 1: Some of the thyroid functions (1)

2. Thyroid dysfunction

Thyroid dysfunction is quite common in the general population, especially among women; moreover, it has been suggested that the number of undetected cases of thyroid dysfunction may be twice compared to the detected ones (16). Typically, thyroid dysfunction is divided into hyperthyroidism and hypothyroidism.

2.1. Hypothyroidism

Hypothyroidism is a decrease in thyroid gland function and thyroid hormone secretion. It is classified into 3 categories: a) primary (autoimmune, iatrogenic, transient, and iodine deficiency), b) secondary (pituitary failure and tumor), and c) tertiary (hypothalamic failure) (17). In the United States, Hashimoto thyroiditis (autoimmune disease) is the most frequent cause of hypothyroidism, but, globally, nutritional iodine deficiency is the most common cause (18). Some of hypothyroidism causes are summarized in **Table 2**.

Causes of hypothyroidism
<p>Primary:</p> <ul style="list-style-type: none"> - Autoimmune disease: <ul style="list-style-type: none"> • Atrophic thyroiditis • Hashimoto's thyroiditis (18) - Iatrogenic: <ul style="list-style-type: none"> • Radioactive iodine therapy (19) • Thyroidectomy • Drugs (anti-thyroid, amiodarone, thalidomide, lithium...) (20) - Transient: <ul style="list-style-type: none"> • Subacute thyroiditis • Postpartum thyroiditis (18) - Iodine deficiency (18) <p>Secondary hypothyroidism:</p> <ul style="list-style-type: none"> - Pituitary failure - Pituitary tumor <p>Tertiary hypothyroidism:</p> <ul style="list-style-type: none"> - Hypothalamic failure

Table 2: Main causes of hypothyroidism (17)

Clinically, the signs and symptoms can be mild and nonspecific and differs from one patient to another (18). When severe hypothyroidism occurs in childhood, it is referred to as cretinism; the major cause of cretinism is chronic iodine deficiency and it is, usually, presented as following: a) short stature/poor growth, b) decreased activity, b) frontal bossing, c) tongue protrusion, d)

hoarse voice, e) hypertelorism, f) muscles weakness (hypotonia), g) dry skin, and h) alopecia (21).

In adults, typical features of hypothyroidism include enlarged gland, weight gain, depression, anxiety, psychosis, memory loss, dysarthria and voice changes, bradykinesia, lethargy, dry skin and hair, pallor and jaundice, blank facial expressions, macroglossia, bradycardia, constipation, asthenia and muscle cramps, cold intolerance, sleep disorders, menstrual cycle irregularities, galactorrhea (18, 22). Patient can also present sleep apnea, hyponatremia, hypercholesterolemia, congestive heart failure, pericardial effusion and prolonged QT interval (22).

It is important to note that myxedema coma, an exceptional lethal condition, is a result of severely advanced hypothyroidism with loss of the adaptive mechanism to maintain homeostasis (23).

Diagnosis of hypothyroidism is based on signs and symptoms, a low serum level of T4, and a high serum TSH level (22).

Hypothyroidism is primarily treated with levothyroxine monotherapy (24). Therapy should be monitored based on TSH serum levels which must be tested every 4 to 8 weeks until target is achieved; when stable, then the monitoring interval can be extended to 6 to 12 months (22).

2.2. Hyperthyroidism

Hyperthyroidism is a syndrome related to thyroid hormone production excess. Graves's disease and toxic multinodular goiter are the more frequent causes of hyperthyroidism; however, other etiologies may include: toxic adenomas, subacute thyroiditis (de Quervain thyroiditis), iodine-induced hyperthyroidism (Jod-Basedow phenomenon), and (subacute thyroiditis) (17, 25) (**Table 3**).

Causes of hyperthyroidism
<ul style="list-style-type: none"> -Autoimmune <ul style="list-style-type: none"> • Grave's disease -Toxic nodular goiter -Toxic adenoma -Subacute thyroiditis -Iodine-induced hyperthyroidism -Drug-induced (amiodarone, lithium)

Table 3: Main causes of hyperthyroidism (17)

Hyperthyroidism pathophysiology depends on the etiology of the condition; in Grave's disease, the original cause is autoimmune where thyroid-stimulating immunoglobulins are produced and went to bind to the TSH receptor, thus mimicking the TSH effects. On the other hand, the toxic multinodular goiter which presents with palpable thyroid nodules lead to the production of excess thyroid hormone from autonomous ectopic tissue (25).

Unlike the toxic multinodular goiter which can be formed by multiple nodules, toxic adenoma classically presents with a solitary papillary nodule that also causes hyperthyroidism (25).

Clinically, hyperthyroidism signs and symptoms reflect the state of increased metabolic activity including weight loss despite an augmented appetite, heart palpitation, anxiety, tremors, weakness, diarrhea, heat intolerance, atrial fibrillation, congestive heart failure, and shortness of breath (25).

Physical examination may reveal an enlarged gland in the case of goiter or one or more palpable nodules (26).

Diagnosis of hyperthyroidism is based on TSH, T4, and T3 serum levels tests. Due to the negative feedback that T3 and T4 apply on the pituitary gland, elevated T3 and/or elevated T4 will cause decreased TSH production (27).

Treatment of hyperthyroidism depends on the underlying cause. Usually 2 types of treatment are considered, symptomatic and basic. In the former, the physician prescribes drugs depending on the symptoms; e.g. beta-adrenergic antagonist or calcium channel blocker to control the palpitations, anxiety, and tremor (28). As for the basic treatment, radioactive iodine therapy, thionamide therapy, and subtotal thyroidectomy can be used. It is important to note that all of them predispose the patient to possible long-term hypothyroidism. Thus, clinical assessment and T4 monitoring are crucial for patients who undergo any of these treatments (25).

It is important to note that untreated hyperthyroidism can lead to a major, potentially life-threatening complication referred to as thyroid storm (thyrotoxic crisis). In this condition reflecting the advanced state of hyperthyroidism, and requiring immediate attention, the patient suffers from tachycardia, increased gastrointestinal motility, diaphoresis, anxiety, and fever (25).

3. Dental management of patients with thyroid dysfunction

In dentistry, the risk in treating patients with controlled thyroid dysfunction is not elevated. Contrariwise, undiagnosed or uncontrolled thyroid conditions are risky (29). For that, the dentist should be alert for any warning signs of such dysfunction and then refer the patient for medical consultation before any dental treatment.

Following are common oral manifestations found in thyroid disorders as well as the precautions dental practitioners should take to avoid potential complications during or after dental treatments.

3.1. Hypothyroidism

Frequent oral findings in hypothyroidism include macroglossia, glossitis, dysgeusia, compromised periodontal health, salivary gland enlargement, and delayed dental eruption and wound healing (**Table 4**) (21, 30).

Before treating patients with hypothyroidism, a specialized consultation is mandatory for more information on their medical conditions (e.g. cardiovascular status, lethargy, etc.) and the treatment will be provided accordingly. Well-controlled patients require no special precautions while in uncontrolled patients elective treatments should be deferred (31).

Moreover, dentists should be aware that patients under levothyroxine may experience an increased response to drugs affecting the central nervous system function such as narcotics, tranquilizers, and barbiturates. Their use should be avoided (21, 31).

3.2. Hyperthyroidism

Oral manifestations in hyperthyroidism may include increased susceptibility to caries, severe periodontal disease, extraglandular thyroid tissue enlargement (mostly in the posterior dorsum of the tongue, known as ectopic or lingual thyroid), maxillary or mandibular osteoporosis, accelerated dental eruption/premature loss of temporary teeth, and burning mouth syndrome (**Table 4**) (21).

Hypothyroidism	Hyperthyroidism
<ul style="list-style-type: none"> -Macroglossia -Glossitis -Dysgeusia -Compromised periodontal health -Salivary gland enlargement -Delayed dental eruption -Delayed wound healing 	<ul style="list-style-type: none"> -Increased susceptibility to caries -Severe periodontal disease -Extraglandular thyroid tissue enlargement (mainly lingual thyroid) -Maxillary or mandibular osteoporosis -Accelerated dental eruption/premature loss of temporary teeth -Burning mouth syndrome

Table 4: Oral manifestations of hypothyroidism and hyperthyroidism

As patients with hyperthyroidism are susceptible to cardiovascular diseases such as atrial dysrhythmias, it is essential that dentists address these patients' cardiac history by consulting their physicians before any invasive procedures. These procedures should be deferred in case of symptomatic patients (tachycardia, irregular pulse, sweating, hypertension, tremor, etc.) and uncontrolled status (21).

The use of vasoconstrictors (epinephrine) deserves special consideration when treating patients with hyperthyroidism. Epinephrine can exacerbate symptoms of tachycardia, dyspnea, and fatigue (31). Moreover, dentists should use a stress reduction protocol; they can prescribe drugs to control anxiety but some oral sedatives may potentiate antithyroid drugs (31).

Finally, dentists should be prepared to take emergency measures while waiting the medical support if the thyrotoxic crisis clinical signs and symptoms occur. For that the following must be done: a) prevent hyperthermia by using cold towels to keep the patient cool, b) administer intravenous hydrocortisone (100-300 mg) and hypertonic glucose if available, and c) monitor vital signs and be prepared for cardiopulmonary resuscitation if necessary (31).

Table 5 resumes dental treatments modifications in case of thyroid dysfunction as proposed by Pinto and Glick (21).

<p>Before treatment:</p> <ul style="list-style-type: none"> - Confirm the type of thyroid dysfunction - Symptomatic thyroid dysfunction: postpone elective treatment and consult the physician - Uncontrolled thyroid dysfunction: consult physician - In case of cardiovascular disease: assess cardiovascular status - Patient under anticoagulation therapy: make proper treatment modifications - Assess drugs interactions - Take blood pressure and heart rate: if abnormal, postpone elective treatment and consult the physician
<p>During treatment:</p> <ul style="list-style-type: none"> - Oral examination: check for oral manifestations (enlarged salivary glands, others) - Controlled thyroid dysfunction: no contraindication to local anesthetic with vasoconstrictor - Patient under non selective β-blockers: avoid vasoconstrictors - Uncontrolled thyroid dysfunction: <ul style="list-style-type: none"> • Only emergencies should be performed • Avoid vasoconstrictors • Minimize stress– short appointments • Discontinue treatment if any symptoms related to the thyroid dysfunction occur
<p>After treatment:</p> <ul style="list-style-type: none"> - Control pain - Patients with hypothyroidism: sensitivity to central nervous system depressants and barbiturates - Patients with hyperthyroidism: Use with precaution non-steroidal anti-inflammatory drugs and avoid aspirin - Continue hormone replacement therapy or antithyroid drugs as prescribed

Table 5: *Thyroid dysfunction-considerations for dental treatments (21)*

4. Conclusion

Thyroid disorders are serious conditions. Dental practitioners should be familiar with their oral and general manifestations so they can recognize any complication and evaluate their control level. In case of an undiagnosed/uncontrolled dysfunction, all elective dental treatment should be postponed until a full medical assessment is performed. Moreover, dentists should be prepared to take emergency measures and ask for medical support if the clinical manifestations of the thyrotoxic crisis occur.

References

1. Khan YS, Farhana A. Histology, Thyroid Gland. [Updated 2020 Jul 3]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK551659/>
2. Ortiga-Carvalho TM, Chiamolera MI, Pazos-Moura CC, Wondisford FE. Hypothalamus-pituitary-thyroid axis. *Compr Physiol*. 2016; 6(3): 1387-1428. doi: 10.1002/cphy.c150027.
3. Felsenfeld AJ, Levine BS. Calcitonin, the forgotten hormone: does it deserve to be forgotten? *Clin Kidney J*. 2015; 8(2): 180-187. doi: 10.1093/ckj/sfv011.
4. Hofstee P, Bartho LA, McKeating DR, Radenkovic F, McEnroe G, Fisher JJ, et al. Maternal selenium deficiency during pregnancy in mice increases thyroid hormone concentrations, alters placental function and reduces fetal growth. *J Physiol*. 2019; 597(23): 5597-5617. doi: 10.1113/JP278473.
5. Talat A, Khan AA, Nasreen S, Wass JA. Thyroid screening during early pregnancy and the need for trimester specific reference ranges: a cross-sectional study in Lahore, Pakistan. *Cureus*. 2019; 11(9): e5661. doi: 10.7759/cureus.5661.
6. Taousani E, Savvaki D, Tsiros E, Poulakos P, Mintzioti G, Zafrakas M, et al. Regulation of basal metabolic rate in uncomplicated pregnancy and in gestational diabetes mellitus. *Hormones (Athens)*. 2017; 16(3): 235-250. doi: 10.14310/horm.2002.1743.
7. Zhu B, Zhao G, Yang L, Zhou B. Tetrabromobisphenol A caused neurodevelopmental toxicity via disrupting thyroid hormones in zebrafish larvae. *Chemosphere*. 2018; 197: 353-361. doi: 10.1016/j.chemosphere.2018.01.080.
8. Bernal J. Thyroid hormones in brain development and function. [Updated 2015 Sep 2]. In: Feingold KR, Anawalt B, Boyce A, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK285549/>
9. Acar H, Ulgen A. Relationship between thyroid hormone levels and crime type: a controlled study in prisoners. *Int J Endocrinol*. 2020; 2020: 9172134. doi: 10.1155/2020/9172134.
10. Delitala AP, Scuteri A, Maioli M, Mangatia P, Vilardi L, Erre GL. Subclinical hypothyroidism and cardiovascular risk factors. *Minerva Med*. 2019; 110(6): 530-545. doi: 10.23736/S0026-4806.19.06292-X.
11. Udovcic M, Pena RH, Patham B, Tabatabai L, Kansara A. Hypothyroidism and the heart. *Methodist Debakey Cardiovasc J*. 2017; 13(2): 55-59. doi: 10.14797/mdcj-13-2-55.
12. Damiano F, Rochira A, Gnoni A, Siculella L. Action of thyroid hormones, T3 and T2, on hepatic fatty acids: differences in metabolic effects and molecular mechanisms. *Int J Mol Sci*. 2017; 18(4): 744. doi: 10.3390/ijms18040744.
13. McAninch EA, Miller BT, Ueta CB, Jo S, Kim BW. Thyroid hormone at near physiologic concentrations acutely increases oxygen consumption and extracellular acidification in LH86 hepatoma cells. *Endocrinology*. 2015; 156(11): 4325-4335. doi: 10.1210/en.2015-1221.
14. Lee JH, Kwon OD, Ahn SH, Choi KH, Park JH, Lee S, et al. Reduction of gastrointestinal motility by unilateral thyroparathyroidectomy plus subdiaphragmatic vagotomy in rats. *World J Gastroenterol*. 2012; 18(33): 4570-4577. doi: 10.3748/wjg.v18.i33.4570.
15. Park S, Han CR, Park JW, Zhao L, Zhu X, Willingham M, et al. Defective erythropoiesis caused by mutations of the thyroid hormone receptor α gene. *PLoS Genet*. 2017; 13(9): e1006991. doi: 10.1371/journal.pgen.1006991.
16. Gessl A, Lemmens-Gruber R, Kautzky-Willer A. Thyroid disorders. *Handb Exp Pharmacol*. 2012; (214): 361-386. doi: 10.1007/978-3-642-30726-3_17.

17. Jefferys A, Vanderpump M, Yasmin E. Thyroid dysfunction and reproductive health. *Obstet Gynaecol.* **2015**; 17(1): 39–45. doi:10.1111/tog.12161
18. Patil N, Rehman A, Jialal I. Hypothyroidism. [Updated **2020** Nov 21]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK519536/>
19. Huseni MA. The incidence of hypothyroidism following the radioactive iodine treatment of Graves' disease and the predictive factors influencing its development. *World J Nucl Med.* **2016**; 15(1): 30-37. doi: 10.4103/1450-1147.167582.
20. Taylor PN, Albrecht D, Scholz A, Gutierrez-Buey G, Lazarus JH, Dayan CM, et al. Global epidemiology of hyperthyroidism and hypothyroidism. *Nat Rev Endocrinol.* **2018**; 14(5): 301-316. doi: 10.1038/nrendo.2018.18.
21. Pinto A, Glick M. Management of patients with thyroid disease: oral health considerations. *J Am Dent Assoc.* **2002**; 133(7): 849-858. doi: 10.14219/jada.archive.2002.0299.
22. Garber JR, Cobin RH, Gharib H, Hennessey JV, Klein I, Mechanick JI, et al. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the american association of clinical endocrinologists and the american thyroid association. *Thyroid.* **2012**; 22(12): 1200-1235. doi: 10.1089/thy.2012.0205. Erratum in: *Thyroid.* 2013; 23(2): 251. Erratum in: *Thyroid.* 2013; 23(1): 129.
23. Elshimy G, Correa R. Myxedema. [Updated **2020** Oct 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK545193/>
24. Jonklaas J, Bianco AC, Bauer AJ, Burman KD, Cappola AR, Celi FS, et al. Guidelines for the treatment of hypothyroidism: prepared by the american thyroid association task force on thyroid hormone replacement. *Thyroid.* **2014**; 24(12): 1670-1751. doi: 10.1089/thy.2014.0028.
25. Mathew P, Rawla P. Hyperthyroidism. [Updated **2020** Nov 21]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK537053/>
26. De Leo S, Lee SY, Braverman LE. Hyperthyroidism. *Lancet.* **2016**; 388(10047): 906-918. doi: 10.1016/S0140-6736(16)00278-6.
27. Menconi F, Marcocci C, Marinò M. Diagnosis and classification of Graves' disease. *Autoimmun Rev.* **2014**; 13(4-5): 398-402. doi: 10.1016/j.autrev.2014.01.013.
28. Ross DS, Burch HB, Cooper DS, Greenlee MC, Laurberg P, Maia AL, et al. 2016 American thyroid association guidelines for diagnosis and management of hyperthyroidism and other Causes of thyrotoxicosis. *Thyroid.* **2016**; 26(10): 1343-1421. doi: 10.1089/thy.2016.0229. Erratum in: *Thyroid.* 2017; 27(11):1462.
29. Malamed SF. Medical emergencies in the dental surgery. Part 1: Preparation of the office and basic management. *J Ir Dent Assoc.* **2015**; 61(6): 302-308.
30. Young ER. The thyroid gland and the dental practitioner. *J Can Dent Assoc.* **1989**; 55(11): 903-907.
31. Patton LL, Glick M. The ADA practical guide to patients with medical conditions. 2nd ed. Hoboken (New Jersey): John Wiley and Sons; **2016**. pp. 89-97.