

Type of the Paper (Article)

Comparison of Two Commercial Recirculated Aquacultural Systems and Their Microbial Potential in Plant Disease Suppression

Sammar Khalil^{1,*}, Preeti Panda², Farideh Ghadamgahi³, AnnaKarin Rosberg¹ and Ramesh R Vetukuri³

¹ Swedish University of Agricultural Sciences, Department of Biosystems and Technology, Box 103, 230 53 Alnarp; sammar.khalil@slu.se & annakarin@rosberg@slu.se

² Plant and Food Research, Plant Protection, Lincoln, New Zealand; preeti.panda@plantandfood.co.nz

³ Swedish University of Agricultural Sciences, Department of Plant Breeding, Box 102, 230 53 Alnarp; ramesh.vetukuri@slu.se & farideh.ghadamgahi@slu.se

* Correspondence: sammar.khalil@slu.se ; Tel.: +4640415335

Abstract: Aquaponics are food production systems advocated for food security and health. Their sustainability from a nutritional and plant health perspective is, however, a significant challenge. Recirculated aquaculture systems (RAS) form a major part of aquaponic systems but knowledge about their potential to benefit plant growth and suppress their pathogens is limited. The current study tested if the diversity and function of microbial communities in two commercial RAS were specific to the fish species used (Tilapia or Clarias) and sampling site (fish tanks and wastewaters), and whether they confer benefits to plants and have *in vitro* antagonistic potential towards plant pathogens. Microbial diversity and composition was found to be dependent on fish species and sample site. The Tilapia RAS hosted higher bacterial diversity than the Clarias RAS; but the latter hosted higher fungal diversity. Both Tilapia and Clarias RAS hosted bacterial and fungal communities that promoted plant growth, inhibited plant pathogens and encouraged biodegradation. The production of extracellular enzymes, related to nutrient availability and pathogen control, by bacterial strains isolated from the Tilapia and Clarias systems, makes them a promising tool in aquaponics and in their system design.

Keywords: Aquaponics; Tilapia; Clarias; *Bacteroidetes*; *Proteobacteria*; *Actinobacteria*; extracellular enzymes; *Pseudomonas fluorescens*; *Pseudomonas veronii*; plant growth promotion; *In vitro* antagonistic: Plant pathogens

1. Introduction

Largescale challenges such as climate change, increased world population, limited availability of natural resources, and the spread of pandemics pose risks to food security [1]. Based on the United Nations sustainable development goals, the need to shift to sustainable food production systems to face these challenges is of great importance [2, 3]. Aquaculture is one of the largest food industries worldwide with respect to the production of animal proteins. The future expansion of this industry will focus primarily on land-based recirculated aquaculture systems (RAS). The environmental profiles of RAS are characterized by their ability to reduce water consumption, allow for better control of rearing conditions, and to reduce significantly the release of nutrients (organic matter, nitrogen and phosphorous) into lakes, rivers and the sea [4]. However, the accumulation of nitrates harmful to fish and the environment in RAS remains a challenge [5].

Integrating RAS and hydroponic systems to build so-called aquaponic systems for plant cultivation has been promoted as a solution to deal with the nitrate challenge, since the nitrate content can be reduced by plant uptake. This integration makes the aquaponic system an ideal candidate as

a future food production system with a robust environmental profile and the potential to enhance food security. However, the sustainability of aquaponic systems still needs further improvement. Aquaponics are complex systems and we still lack knowledge concerning the ideal plant nutrient balance in relation to the amount and type of fish feed, system design, and plant and fish diseases [6]. These areas are mainly related to conditions in RAS. For example, the lack or low availability of some essential elements in fish water, that are required for plant growth - such as phosphorous and iron - remains a challenge and limits the productive efficiency of aquaponic systems [7]. These nutrients are therefore currently maintained at required levels by the addition of extra phosphorus and iron to the system. In addition, stabilization of the RAS element of the aquaponic system, with respect to water quality parameters such as temperature and pH, is crucial in order to achieve a balance between fish, plant and bacterial requirements [8, 9] and thereby a good plant growth.

From a plant pathology perspective, the aquaponic system is further challenged by a need for harmless and sustainable solutions for controlling levels of fish and plant pathogens [6]. Root diseases caused by fungal and oomycete pathogens such as *Pythium* spp., *Phytophthora* spp., *Fusarium* spp., *Verticillium* spp., or bacterial pathogens such as *Ralstonia* and *Xanthomonas*, are commonly found in aquatic environments such as hydroponic systems and consequently occur in aquaponic systems [10-13]. Control of these pathogens in hydroponic systems by using biological and non-biological tools is widely investigated and has been covered by other literature [14-17]. The ability for naturally occurring microbial communities to suppress pathogens in hydroponic systems has also been considered in different studies [18, 19]. However, the potential suppressive effect of natural microbial communities in aquaponic systems continues to require further research and investigation. This important area of research needs to deal with the complexity of the various components and aspects of the aquaponic system - from water quality and fish feed in the RAS system, through the biofilter component to the hydroponic unit - in and through which disease organisms may enter and grow to excess if not controlled. Restrictions governing pesticides to control plant diseases or antibiotics for the fish diseases emphasize our need to provide solutions that enhance the sustainability of aquaponic systems.

The presence of microbial communities can have a suppressive effect in the system against plant pathogens. For example, through different modes of action including competition, production of secondary metabolites such as antibiotics and extracellular enzymes, induced resistance and growth-promoting effects in plants, microorganisms can have various beneficial roles in maintaining a healthy environment [20]. In aquaponic systems, the role of microbial communities in nutrient recycling [21] and promoting plant growth [22] has been confirmed. Other studies have also shown that rhizobacteria inhabiting the root environments of aquaponic systems have antagonistic potential against pathogens such as *Pseudomonas* and *Flavobacterium* [23].

The RAS is a significant unit in the aquaponic system and has received much attention as a microbial habitat and a tool against fish diseases [24]. However, more research into the role of microbes in RAS is needed in order to deepen our understanding of how they might be optimized for plant growth and against plant diseases. Targeting those RAS factors that promote the anti-pathogenic potential and improve plant growth through the action of naturally occurring microbial communities is thus of great concern. With this in mind, the objective of the current study was to investigate microbial diversity in a commercial RAS (with no aquaponic connection) and elucidate its potential to promote plant growth and act against plant pathogens. We examined how fish species and sampling site in the system, namely the water tank compared with the fish biosolids or wastewaters, affected the results. We also investigated the microbial communities' function regarding the production of extracellular enzymes, which are known for their mode of action against plant pathogens as well as towards nutrient solubility. Our hypotheses are: (i) that a RAS (without any aquaponic connection) hosts microbial communities beneficial to plants and antagonistic to pathogens, and that these communities develop in ways that are characteristically specific to fish species used and site sampled in the system; and (ii) that the production of extracellular enzymes and the antagonistic potential to control plant pathogens *in vitro* are fish species-specific and site-specific.

2. Materials and Methods

2.1. Sample collection

Two commercial warm and fresh water RAS populated with Tilapia (*Oreochromis niloticus*) or Clarias (*Clarias gariepinus*) as the fish species and no aquaponic connection were used as the experimental units. Three independent replicates (each 5L water samples) were collected at two different sites, viz. from the fish water tank and from the wastewater where the fish feces were accumulated. Both RAS with each respective fish type were similarly sampled. The collection site was considered as a treatment. Thus, in total, three fish water tank samples (biological replicates) and three wastewater samples (biological replicates) were randomly collected from the Tilapia and Clarias system respectively. The samples were then transferred to the laboratory for subsequent microbial analyses.

2.2. Viable count and microbial enumeration

The viable count method was used to quantify the microbiota in each treatment. Dilution series and enumeration on selective agar media were applied [25]. From each replicate, 1 mL was serially diluted in 0.85% NaCl. From this dilution stock, 200 μ L aliquots were spread, in triplicate, on the following media: i) 0.1% Tryptic soya agar (TSA, DIFCO 0369-17-6) complemented with cycloheximide (100 μ L mL⁻¹) to enumerate the general bacterial flora; ii) 0.5% malt extract agar (MA, DIFCO 0186-17-7) to enumerate the general fungal flora; and iii) King Agar B (KB) with cycloheximide (100 μ g mL⁻¹) to enumerate the fluorescent pseudomonads. The MA plates were incubated at room temperature for seven days and the TSA and KB plates were incubated for 24 hours at 25 °C.

2.3. Microbial community analyses

2.3.1. Samples preparation

The total microbiome analyses started by sterile filtration of 1L of the collected samples through 0.2 μ m filters using bottle-top vacuum filtration systems, PES (WVR- Sweden, 514-0332). The filtration unit filter was then transferred to a 50 mL tube, washed with 50 mL sterile autoclaved water, followed by vigorous vortexing for 2 min. The collected material was then centrifuged at 5000 rpm for 10 min and the pellets were stored at -80 °C.

2.3.2. DNA extraction

The DNA extraction was performed using EnzyMo DNA preparation kit (D 4300, Sigma Aldrich) following the manufacturer's recommendations.

2.3.3. Illumina sequencing

The bacterial and fungal communities were sequenced with an Illumina MiSeq (2 × 300 bp) at LGC Genomics GmbH (Berlin, Germany) [26] using Illumina bcl2fastq 2.17.1.14 software. The bacterial 16S ribosomal gene was targeted using the forward primer 341F (5'-CCTACGGGNGGCWGCAG-3') and the reverse primer 785R (5'-GACTACHVGGGTATCTAATCC-3'). The fungal forward primer ITS7F (5'-GTG ART CAT CGA ATCTTTG GTT G-3') and the reverse primer (5'-TCC TCC GCT TAT TGA TAT GC-3') were used to target the ITS region for fungal assessment. Data pre-processing and OTU picking from amplicons were performed using MOTHUR pipelines (version 1.35.1). Reads with a final length of <100 bases were discarded and primer, barcode sequences as well as chimeras were removed. For taxonomical classification, alignment against 16S Mothur-Silva SEED r119 reference was performed and sequences from other domains of life were removed. Assignment of OUTs was performed at the 97% identity level using the cluster split method. The fast Tree method was used to generate the phylogenetic trees for 16S and ITS,

respectively. The sequenced data were submitted to Genbank database under the bio project submission number SUB8558843

2.3.4 Statistical analyses

All statistical analyses were performed within Microbiome Analyst v1.0 [27]. OTUs were pre-filtered before conducting any statistical analysis by retaining those present in at least two samples. The data were normalized using the total sum scaling method (counts per million normalization) to address variability in sampling depth. Alpha diversity profiling was conducted using Shannon and Chao1 diversity indices and statistical significance testing using analysis of variance (ANOVA). To examine differences between the different groups in community compositions, beta diversity was calculated using unweighted Unifrac Distance and statistical comparisons performed with ANalysis of SIMilarities (ANOSIM). Patterns of sample dissimilarity were visualized using Principal Coordinate Analysis (PCoA) plots. In addition, hierarchical clustering and heatmaps were constructed using the Ward clustering algorithm and Euclidean distance measure to examine the robustness of sample clustering and relative abundances. The results from the microbial enumeration were log-transformed to meet the assumptions of homogeneity and normality. The significance effect of treatment was tested by ANOVA followed by Tukey's multiple comparison test ($p < 0.05$) using Minitab v16. To explore how top taxa differed between the different cultivation/waste types, classical univariate statistical comparisons were inferred using t-test/ANOVA method with an adjusted p-value cutoff of 0.05.

2.4. Microbial activities

2.4.1 Bacterial pure cultures

For each treatment, two single colonies from the TSA plates and two from the KB plates were selected and transferred to be grown on broth media of tryptic soya broth (TSB) and King B broth (KBB), respectively. One loopful of culture was inoculated into 15 mL of the broth media and incubated at 25 °C with shaking (140 rpm) for 24 h. Bacterial DNA was extracted using the Quick-DNA Bacterial Microprep Kit according to the manufacturer's recommendations (Zymo Research, USA). The DNA yield and integrity was assessed using a NanoDrop Micro Photometer (NanoDrop Technologies, UK), and agarose gel electrophoresis, respectively. The 16s rRNA region of all bacterial isolates was PCR amplified individually with the universal primer pairs, 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and 907R (5'-CCGTCAATTCMTTTRAGTTT-3') [28]. PCRs were performed using ten ng of DNA with the following temperature parameters: initial denaturation step at 94 °C for 3 minutes, followed by 35 cycles at 94 °C for 45 s, 50 °C for 30 s, and 72 °C for 30 s, followed by a final extension step of 72 °C for 5 min. The PCR products were purified using the Qiagen QIAquick PCR Purification Kit (Qiagen, UK). Sanger sequencing for species identification was carried out at the GATC biotech AG sequencing facility (Germany) using 27F and 907R primers. DNA star software was used (DNASTAR, USA) to analyze and edit nucleotide sequences obtained from the sequencing platform manually. Resulting sequences with 16s region were searched for matching hits against the National Center for Biotechnology Information (NCBI) GenBank non-redundant nucleotide database (BLASTn; [29]). Search hits to sequences from records in the database were evaluated for coverage and identity and the best matched NCBI accession was recorded.

2.4.2 Enzyme activities

Assessment of the functional characters of the isolates was performed with respect to enzymatic activities. The isolated bacterial colonies were screened for the production of amylase, protease, phosphatase and cellulase on functional media using a plate assay [30]. The minimal medium (M9, Minimal Salts (VWR- Sweden) was used as a base medium. For amylase activity, the isolates were spotted on M9 amended with 1% (w/v) starch. After incubation for 24 - 72 hours at 20 ± 2 °C, the

plates were flooded with Lugol iodine; the appearance of a zone of clearance towards the dark background is a positive indication of amylase production.

For the protease activity, M9 plates amended with skimmed milk (20 mL L⁻¹) were used and incubated for 24 - 72 hours at 20 ± 2 °C. The formation of a halo zone around the colonies indicated a positive protease production. The same indication assessment and incubation conditions were used for cellulase production. The M9 plates amended with carboxymethyl cellulose were used and after incubation, they were flooded with Congo Red solution (0.2% w/v) for 30 min and washed with 1 M NaCl solution.

For the phosphatase activity, Tryptose phosphate agar supplemented with Methyl Green (0.05 mg mL⁻¹) was used. The plates were incubated for 5 days at 20 ± 2 °C; the development of green color indicated positive phosphatase activity. The siderophore production was also assessed using Chromazurol S agar plates spotted with bacterial isolates and incubated for three days at 20 ± 2 °C. The positive result was indicated by the formation of an orange halo around the colonies.

2.4.3 *In vitro* antagonistic assay

The selected bacterial isolates were screened for antagonistic activity against the oomycete/fungal pathogens *Phytophthora cactorum* and *Verticillium dahlia*. The assay was performed on agar plates by transferring a mycelial plug (1×1cm) from an actively growing 10-day-old V8 culture of the pathogens to a corn meal agar (CMA) plate. A loopful of the bacterial isolate was then streaked in parallel lines, 3 cm apart, on each side of the pathogen plug and incubated for ten days at 20 °C. Three replicate plates were maintained for each isolate. Plates inoculated with only the test pathogens were used as controls. The radial mycelial growth in the direction of the antagonist was measured and the Growth Inhibition Percentage (GIP) was calculated according to the following formula:

$$(\%) = \frac{(RC - RT)}{RC} \times 100$$

where RC is the radial growth of the pathogen in the control plate (cm) and RT is the radial growth of the pathogen (cm) in the treated plate [31].

2.4.4 *Detached leaf assay (DLA)*

The inhibitory effect of bacterial isolates against the strawberry pathogen *P. cactorum*, was further evaluated in a detached leaf assay. The bacterial isolates with high antagonistic activity from the *in vitro* studies against the pathogens were selected for the DLA assay.

For the bacterial suspension, the bacteria were grown on 15 mL TSB media in falcon tubes (50 mL) followed by overnight incubation at 28 °C on a shaker at 220 rpm. The tubes were centrifuged for 10 min at 4200 rpm and the pelleted cells were resuspended in sterile distilled water. The bacterial density was measured by OD at 600 nm and adjusted to OD = 0.1.

P. cactorum strain RV4 was cultured on V8 carrot media and incubated for 7 days at 21 °C. Three agar pieces from the colony's periphery were transferred to a petri dish with 20 mL of V8 broth and incubated in the dark at room temperature for three days to induce zoospore production. To promote sporangia development from the mycelia cultured in the dark, the mycelial mats were washed three times in autoclaved distilled water followed by the addition of 20 mL of autoclaved soil extract solution. The plates were incubated for 24 hours under fluorescent light at 21 °C. The soil extract solution was removed, and the mycelia were covered with autoclaved distilled water (20 mL) and incubated under the same conditions as mentioned above for another 24 hours.

To promote zoospore release, the plates were subjected to a cold shock in a refrigerator at eight °C for 1 hour and then transferred to room temperature for another hour. The zoospore samples were collected and counted using a hemocytometer to adjust the inoculum to 2 × 10⁴ zoospores per mL.

Young strawberry leaflets (cv. Sonata) were used and the adaxial leaf surface was inoculated on either side of the leaf midrib with 20 µL droplets of the mixture of zoospore and bacterial suspension at a final concentration of 2 × 10⁴ mL⁻¹ and of OD = 0.1, respectively. The positive control consisted of

leaflets inoculated with the *P. cactorum* zoospore suspension only, and negative controls with sterile distilled water. Each treatment was evaluated with six replicates and the experiment was repeated thrice.

Leaflets were placed on grids in plastic boxes with moist filter paper, sealed and stored at room temperature. Four days after inoculation, the leaves were scored for disease progression, and the severity of infection was assessed using the image-processing program ImageJ.

3. Results

3.1 RAS conditions

Conditions at the sample collection time in the two RAS systems with different fish species were similar regarding water quality parameters such as pH, temperature, conductivity, and ammonium and nitrate content (Table 1). The two fish species differed in their combined weight per tank. Tilapia being lower than Clarias. Both fish species were fed with the same amount of a commercial feed (<https://www.skretting.com/en/>).

Table 1. Growth conditions in each recirculated aquaculture system (RAS) populated with either Tilapia or Clarias

Cultivation factors	RAS with Tilapia	RAS with Clarias
Temperature	21 °C	20.9 °C
pH	7.7	8.5
Conductivity**	130 mS cm ⁻¹	180 mS cm ⁻¹
Ammonium content	2.1 mg / liter	1.8 mg / liter
Nitrate content	280 mg / liter	188 mg / liter
Fish weight	40-600g per tank	100-4000g per tank
Feed content	37% protein + 10% fat	44% protein + 12% fat

3.2 Microbial amount

Microbial enumeration on selective media indicated more general bacterial flora and *Pseudomonas fluorescens* in samples collected from the Tilapia RAS system than the Clarias RAS system (Figure 1). With respect to sampling site, we found no significant differences in the amount of general bacterial flora and *P. fluorescens* between the samples collected from water tanks and waste samples in either Tilapia or Clarias RAS. However, there were differences in the amount of general fungal flora. The highest amount of the general fungal flora was found in samples collected from Clarias wastes than in Clarias water tank or any samples from the Tilapia RAS.




Figure 1. Microbial colonies isolated from water and waste samples from Tilapia and Clarias RAS and enumerated on 0.1% Tryptic soya agar (TSA) complemented with cycloheximide ($100 \mu\text{L mL}^{-1}$) for enumeration of the general bacterial flora; 0.5% malt extract agar (MA) for enumeration of the general fungal flora; and on King Agar B (KB) with cycloheximide ($100 \mu\text{g mL}^{-1}$) for enumeration of fluorescent pseudomonads. Letters above the bars indicate the significant differences between the treatments.

3.3 Diversity of the microbial communities

A total of 4,558 bacteria OTUs and 405 fungal OTUs were generated after processing the Illumina MiSeq sequencing data. The total read counts for the bacterial dataset were 647,232 and 530,351 for the fungal dataset. Alpha diversity of bacterial communities was significantly higher in Tilapia water and wastes samples and in Clarias waste samples (Shannon, $p = 0.021$) compared to Clarias water samples (Figure 2a). In relation to the Shannon indices, Clarias water tank samples returned the lowest alpha diversity. Furthermore, similarities in alpha diversity were seen within Tilapia samples and between Tilapia samples and Clarias waste. The Chao1 indices also revealed a significant increase (Chao1, $p = 0.012$) in alpha diversity in samples collected from Tilapia water tank and wastes and from Clarias wastes compared with Clarias water tank. Clarias water tank showed the lowest alpha diversity with respect to Chao1 indices (Figure 2b). Similarities in the Chao1 indices were also shown within Tilapia samples and between Tilapia samples and Clarias wastes.




Figure 2. Alpha diversity of bacterial communities in the fish water tank and the waste water sampling sites of the Tilapia and Clarias RAS as judged by the diversity indices (a) Shannon and (b) Chao1. Letters above the boxplots indicate the significant differences between the treatments.

With respect to the amount of *Pseudomonas*, we found no significant differences in the log-transformed count of *Pseudomonas*. The amount of *Pseudomonas* in the Tilapia RAS was similar to that of the Clarias RAS (Figure 3). The samples collected from water tanks and wastes samples were also similar.




Figure 3. Log-transformed counts of *Pseudomonas* communities in the water tank and wastewater site from the Tilapia and Clarias RAS. Letters above the boxplots indicate the significant differences between the treatments.

However, although we found no significant differences in fungal community alpha diversity between the treatments in terms of the Shannon indices ($p = 0.211$) (Figure 4a), there were significant differences evident between the treatments in terms of the Chao1 indices ($p = 0.008$). There was higher fungal diversity in the Clarias waste and water tank and in Tilapia wastes compared to Tilapia water tank, which had the lowest fungal diversity (Figure 4b).




Figure 4. Alpha diversity of fungal communities from the water tank and the fish waste sample sites in the recirculated aquaculture systems populated with two different fish species -Tilapia or Clarias, as judged by the diversity indices (a) Shannon and (b) Chao1. Letters above the boxplots indicate the significant differences between the treatments.

Dendrogram analyses demonstrated that the bacterial and fungal communities also differed between the Tilapia and Clarias cultivation systems (Figure 5). The RAS populated with Tilapia indicated an apparent difference in the bacterial communities between the samples collected from the water tank and the waste (Figure 5a). However, these samples also shared similarities in their bacterial communities. Tilapia water tank and waste samples also differed in their fungal communities (Figure 5b). By contrast, there was no clear difference between samples collected from Clarias systems regarding either their bacterial or fungal communities.




Figure 5. The dendrogram analyses of (a) bacterial and (b) fungal communities in samples collected from the RAS populated with either Tilapia or Clarias. The samples were collected from the Tilapia water tank (TT), Tilapia waste (TW), Clarias water tank (CT), and Clarias waste (CW).

The beta diversity metrics clearly distinguished the samples of the bacterial (Figure 6) and fungal (Figure 7) communities with respect to the fish species cultivated in the system and the sampling site. The first axis differentiated between the fish species (Figure 6a) and the second differentiated between the sampling sites in the system (Figure 6b). Clear differentiation with respect to the fish types and the bacterial communities could be shown. The differentiation with respect to the sampling site was more evident for samples from the Tilapia RAS than the Clarias RAS. No differences were found concerning the sampling sites in the Clarias system. Similar observations were evident regarding the fungal communities (Figure 7).




Figure 6. PCoA plots on beta diversity of the bacterial communities distinguished by (a) the cultivation system (ANOSIM; $R = 0.92$ and $p < 0.003$) or (b) sampling site (ANOSIM; $R = 0.81$ and $p < 0.001$) in the system using Tilapia or Clarias as fish species.




Figure 7. PCoA plots on beta diversity of the fungal communities distinguished by (a) the cultivation system (ANOSIM; $R = 0.05$ and $p < 0.003$) or (b) sampling site (ANOSIM; $R = 0.81$ and $p < 0.001$) in the system using Tilapia or Clarias as fish species.

Analyses of the relative abundance of the bacterial communities showed that the phylum *Fusobacteria* was significantly abundant (50%) in the samples collected from the Tilapia wastes, followed by the phyla *Bacteroidetes* (30%) and *Proteobacteria* (20%) (Figure 8). Tilapia water tank samples were dominated by the phylum *Proteobacteria* (60%) followed by *Bacteroidetes* (20%), *Fusobacteria* and *Actinobacteria*. (10%). *Actinobacteria* and *Proteobacteria* dominated samples collected from Clarias systems. The dominance of *Actinobacteria* was higher in the water tank (75%) than the waste samples (60%). *Bacteroidetes* and *Fusobacteria* were more dominant in Clarias wastes than Clarias water tank samples. However, compared with the Clarias RAS, *Actinobacteria* in the Tilapia system were either less abundant or completely absent.




Figure 8. Relative abundance in (%) of dominant bacterial communities at phylum level in water tank, or fish wastes samples collected from RAS populated with Tilapia or Clarias.

The bacterial genera *Microbacterium*, *Cetobacterium* and *Chryseobacterium* dominated the samples collected from the Clarias system (Figure 9). The genus *Microbacterium* dominated both water and waste samples. Tilapia waste samples were dominated by the genera *Cetobacterium* and *Flavobacterium*, followed by *Chryseobacterium*, *Pseudomonas* and *Bacteroides*. The samples from the

Tilapia water tank included more genera and were dominated by *Janthinobacterium*, *Flavobacterium*, *Chryseobacterium*, *Pseudomonas*, *Pseudorhodobacter*, *Bacteroides*, *Sorangium* and *Hydrotalea*.




Figure 9. Relative abundance (%) of dominant bacterial communities at the genus level in water tank, or fish waste samples collected from RAS populated with either Tilapia or Clarias as fish species.

As indicated by the heatmap, the genera *Sorangium*, *Pseudomonas*, *Pseudorhodobacter*, *Acinetobacter*, *Simplicispira*, *Rhodococcus*, *Pedobacter*, *Rhodobacter*, *Undibacterium*, *Hydrogenobacter*, *Phenyloacterium*, *Alpinimonas*, *Acidovorax*, *Paucibacter*, *Legionella*, *Leisona*, *Duganella* were notably present in Tilapia tank water (Figure 10). Tilapia wastes included the genera *Janthinobacterium*, *Cetobacterium*, *Pseudomonas*, *Aremonas*, *Arcobacterium*, *Clostridium*, *Paludibacter*, *Sulfurospirium*, *Dechloromonas*, *Arcobacter*, *Propionivibrio*, *Limnohabitans*, *Paludibacter* and *Macellibacteroides*. The genera *Cryseobacterium*, *Cetobacterium*, *Simplicispira*, *Thermomonas*, *Rhodanobacter*, *Ottowia*, *Bergeyella*, *Prevotella* and *Clostridium* were present in Clarias wastes. The genera *Cryseobacterium*, *Microbacterium*, *Macellibacteroides* and *Azospirillum* were predominant in Clarias water tank samples.




Figure 10. The heatmap shows the abundance of dominant bacterial communities at the genus level in water tank and fish waste samples collected from the Tilapia water tank (TT), Tilapia waste (TW), Clarias water tank (CT), and Clarias waste (CW).

Analyses of the relative abundance of the fungal communities also revealed differences in the dominant communities depending on the fish species and the sampling site (Figure 11). The majority (> 70%) of the fungal communities were unclassified in the samples collected from the Tilapia cultivation system, both from water tanks and Tilapia waste. However, the phylum *Ascomycetes* was well represented in samples from the Tilapia water tanks compared with the samples from the Tilapia wastes, in which the phylum *Basidiomycetes* was more abundant.

By contrast, samples collected from Clarias water tanks and wastes had fewer unclassified fungal communities than the Tilapia samples. However, the relative abundance of the fungal communities differs between samples collected from Clarias water tanks and wastes. The waste samples were highly dominated by the phyla *Ascomycetes*, *Basidiomycetes* and unclassified phyla within the kingdom Protista. The water samples in Clarias system contained *Basidiomycetes* and *Ascomycetes* but were dominated by an unclassified phylum within the kingdom Protista (Figure 11).




Figure 11. Relative abundance (%) of dominant fungal communities at phylum level in samples collected from water tank and fish waste sites in RAS populated with Tilapia or Clarias.

The unclassified fungi also dominated at the genus level (70%) in Tilapia and Clarias RAS (Figure 12). *Trichosporon* spp. dominated (25%) in Clarias water and wastes and in Tilapia wastes. Unclassified genera within the kingdom Protista were only found in the Clarias system. The genus *Yarrowia* was exclusively found in a Tilapia water tank, while the genus *Trichomonascus* was only detected in the Clarias system.




Figure 12. Relative abundance (%) of dominant fungal communities at the genus level in samples collected from water tank and fish wastes in RAS populated with Tilapia or Clarias.

The fungal communities' heatmap showed that the phylum *Basidiomycetes* in Clarias water tank was populated by the genera *Trichosporon*, *Trichomanoascus*, *Phodotonula*, *Malaseezia*, *Mortierella*, *Ascochyta*, *Mucor*, *Cladosporium*, *Wallemia*, *Trichoderma*, *Fusarium* and *Deharyomyces* (Figur 13). The genera *Aspergillus*, *Debaryomyces*, *Wallemia*, *Sterigmatomyces*, *Exobasidium*, *Trichosporon* and *Pleospora* were found in Clarias wastes. The Tilapia waste contained more genus profiles than Tilapia water tank samples and was dominated by the genera *Candida*, *Trichosporon*, *Bettsia*, *Sporobolomyces*, *Penicillium*, *Entomocariticium*, *Cryptococcus*, *Preussia*, *Macrophomina*, *Sterigmatomyces*, *Microascuss* and *Enyodonium*. The Tilapia water tank was dominated by the genera *Yarrowia*, *Acremonium*, *Harposporium*, *Aureobasidium* and *Phaeoacremonium*.




Figure 13. The heatmap presents the abundance of dominant fungal communities at the species level in samples collected from water tanks and fish wastes in RAS populated with Tilapia or Clarias.

3.4. Identification of pure cultures and their enzyme activity potential

In total, nine isolates were collected and identified as described in Table 2. *Pseudomonas fluorescens* was the most dominant isolate identified in Tilapia water tanks and wastes and Clarias wastes. The strains *P. veronii* and *Variovorax paradoxus* were identified in Tilapia wastes. *Variovorax* spp. were also identified in Clarias wastes.

With respect to the enzyme activities, cellulase activity was indicated in some of the isolated strains from both Tilapias and Clarias waste samples (Table 2). The bacterial strains identified from the Clarias wastes were actively producing protease and phosphatase. Amylase activities were positively indicated by all the strains isolated from Tilapia wastes, although not all of these strains indicated positive activities with respect to protease and phosphatase. Strains isolated from Tilapia tanks indicated positive protease and phosphatase activities and one of these strains showed amylase activity. Further, siderophore production was actively indicated by most of the strains isolated from Clarias wastes and by some of the strains isolated from Tilapia water tank and Tilapia wastes.

Table 2. Identification and enzyme activities of the pure bacterial isolates isolated from samples collected from water tanks and wastes in RAS populated with Tilapia or Clarias. (+) indicates positive and (-) indicates negative enzyme activities.

Number of the isolate	Identification	Collection site	Amylase activity	Protease activity	Phosphatase activity	Cellulase activity	Siderophore production
1	<i>Pseudomonas fluorescens</i>	Tilapia water tank	+	+	+	-	+
2	<i>Pseudomonas fluorescens</i>	Tilapia water tank	-	+	+	-	-
3	<i>Pseudomonas fluorescens</i>	Tilapia wastes	+	-	+	+	-
4	<i>Pseudomonas veronii</i>	Tilapia wastes	+	-	+	+	+
5	<i>Variovorax paradoxus</i>	Tilapia wastes	+	-	-	-	-
6	<i>Pseudomonas fluorescens</i>	Clarias water tank	+	+	+	+	+
7	<i>Pseudomonas fluorescens</i>	Clarias wastes	-	+	+	+	+
8	<i>Pseudomonas fluorescens</i>	Clarias wastes	-	+	+	-	+
9	<i>Variovorax</i> spp.	Clarias wastes	-	-	-	-	-

3.5. In vitro antagonistic activity of bacterial isolates against plant pathogens

In vitro antagonistic tests against the plant pathogens, *P. cactorum* (Pc) and *Verticillium dahliae* (Vd) indicated inhibition activities by the isolated strains against both Pc and Vd (Figure 14). Isolate number 6 from the Clarias water tank showed better antagonistic potential against Pc and Vd than

the other isolates. However, the antagonistic activity of number 6 was higher against Vd compared with Pc. Apart from number 6, isolate numbers 3 and 4 from Tilapia wastes indicated higher antagonistic activities against Pc and Vd than all other isolates. Isolate number 2 from Tilapia water tank and number 8 from Clarias wastes had higher antagonistic activities against Vd than Pc. By contrast, isolate number 5 from Tilapia wastes was more antagonistic to Pc than Vd.




Figure 14. Inhibition (%) of radial growth of the plant pathogens *P. cactorum* and *V. dahliae* by the bacterial isolates 1-2 from Tilapia water tanks, isolates 3-5 from Tilapia wastes, isolates 6 from Clarias water tanks and isolates 7-9 from Clarias wastes.

3.7 Detached leaf assay

Two of the best performing biocontrol bacteria isolates (number 4 and 6) were tested for their biocontrol potential against *P. cactorum* on strawberry leaves. In the current study, strawberry leaves (*Fragaria × ananassa* cv. Sonata) were used in a detached leafy assay (DLA). The strawberry leaves inoculated with Pc started showing symptoms of infection four days after inoculation, while the leaves inoculated with sterile distilled water remained symptomless, confirming the viability of the zoospore suspension and no prior pathogen presence on the leaflets. Furthermore, the treatment with isolate number 6, combined with Pc showed potent disease inhibition. However, the bacterial isolate number 4 combined with Pc was less effective in controlling disease progression (Figure 15).



Figure 15. Infection of detached leaflets of strawberry cv. Sonata by *P. cactorum* isolate RV4 with and without potential antagonistic bacterial isolates. Treatments from top to bottom rows: **A:** water only; **B:** *P. cactorum*; **C:** *P. cactorum* combined with bacteria isolate no. 6 (isolated from Clarias wastes); **D:** *P. cactorum* combined with bacteria isolate no. 4 (isolated from Tilapia wastes). Wild type RV4 formed large disease lesions on treatment B. Treatment C, with bacteria 6, and treatment D with bacteria 4 exhibited no or minimal disease. Leaves were photographed four days after inoculation.

4. Discussion

Our study has provided novel insights into the diversity of microbial communities in commercial RAS, their beneficial effects on plants, and their antagonistic potential against plant pathogens. We found that microbial diversity varied depending on the fish species populating the system and from where in the system the samples were taken. The shift in microbial composition and richness as affected by the fish species was supported by the beta diversity analyses, which indicated significant differences and unique composition of the bacterial (Figure 6) and the fungal (Figure 7) communities between the Tilapia RAS and Clarias RAS. Fish species in the system also had significant effect in the water tank environment, which, with Tilapia, was richer with bacteria (Figure 2), while with Clarias it hosted more fungi than bacteria (Figure 4). These findings are in line with other studies [32, 33] indicating how fish species develop their own characteristic microbial communities. The results also agree with [33], who found higher fungal diversity in a Clarias than in a Tilapia system.

By contrast, sampling site had a significant effect on the evenness and richness of bacterial communities in the Clarias RAS (Figure 2), and of fungal communities in the Tilapia system (Figure 4). The site effect was also evident in the composition of the bacterial (Figure 6) and the fungal (Figure 7) communities in the Tilapia RAS. Tilapia wastes hosted a richer fungal diversity than the water tank, while in the Clarias system bacterial diversity was richer in the wastewater than in the tank water. Moreover, the dendrogram analyses (Figure 5) clearly distinguished between the bacterial and fungal communities based on fish species indicating that Tilapia bacterial and fungal communities differed from the Clarias. Differences based on sampling sites in the Tilapia RAS were also evident and could be related to the microbial community's richness and composition. The shift and increased richness in microbial communities in the waste samples might be due to the conditions in the waste environment as affected by the gut microbial and nutrient content [34]. Microbial diversity is also a function of water quality and its chemical properties such as pH, temperature, feed, and nutrient content [35-36]. In the current study, both fish species experienced closely similar cultivation conditions except for their density and weight. Tilapia biomass per tank was lower than that of

Clarias (Table 1). This might have influenced condition of the wastewater in the current study due to differences in the quantity of gut contents excreted into it.

The Tilapia RAS was dominated by bacteria and contained the bacterial phyla *Fusobacteria*, *Proteobacteria* and *Bacteroidetes* (Figure 8). These phyla were present in different relative abundances in the water tank and the waste environment primarily due to the dominance of *Proteobacteria* in the former and of *Fusobacteria* in the latter (Figure 8). The identified phyla are common in fresh water systems and have been previously indicated in Tilapia systems [34]. Our results are also in line with another study that has shown the dominance of *Proteobacteria* in biofilm in an aquaculture system [37]. *Proteobacteria* are characterized as r-strategists with an important function in nutrient recycling [38]. Their higher abundance in the Tilapia water tank might be due to the nutrient conditions in that environment. The high dominance of *Fusobacteria* in the waste of Tilapia is in accord with another study that has shown the relation of *Fusobacteria* to the microbiota in the gut where nutrient breakdown occurs [34]. Our results revealed that *Actinobacteria* was the most dominant phylum in the RAS environment with Clarias species (Figure 8). *Actinobacteria* have received much recent attention for their potential role as probiotic bacteria in marine and fresh water aquaculture [39, 40], further stressing the importance of the antimicrobial activities of these beneficial bacteria for fish and plant health.

Furthermore, microbial patterns in RAS with and without an aquaponic connection have been shown to be rich with bacteria that have the potential to promote plant growth [22]; while others have demonstrated that a Tilapia rearing tank can also host bacterial genera with a similar potential [41]. Our results are in line with these two studies [22, 41] and highlight the potential of the Clarias system to host, both in its tank water and in its waste water, bacterial genera with plant growth promoting characters such as *Microbacterium* and *Cetobacterium* (Figure 9), which are common rhizobacteria known to have beneficial effects on plant growth [22]. The Tilapia system in our study, and in contrast to the Clarias system, was richer in the genera *Pseudomonas*, *Cetobacterium*, *Flavobacterium*, *Sorangium*, *Pseudorhodobacter* and *Bacteroides*, which play a prominent role in promoting plant growth and protecting plants against root pathogens [22]. In addition, the water tank of Tilapia water (Figure 10) hosted genera of great importance in biodegradation and alleviating abiotic stress such as *Simplicispira*, *Rhodobacter* and *Acidovorax* [42, 43], conditions of high relevance in aquaponic cultivation systems. The identified genera in Tilapia wastes (Figure 10) indicate the shift in bacterial composition between the waste site and the water tank. The genera such as *Clostridium*, *Cryseobacterium*, *Janthinobacterium*, *Cetobacterium* and *Pseudomonas* identified in the waste environment also possess plant growth-promoting characters [22]. Moreover, the genera *Propionivibrio*, *Sulfurospirium* and *Dechloromonas* are involved in nutrient breakdown, or the removal of certain compounds, such as phosphorous, also by *Propionivibrio* [44]. The waste site also included genera involved in biodegradation and abiotic stress such as *Limnohabitans* and *Paludibacter* [45]. However, the occurrence of *Legionella* genera in the water tank and of *Aremonas* genera in Tilapia wastes is an area that requires further investigation. These genera are opportunistic pathogens occurring in aquatic environments and are of concern in terms of food safety and human health [46]. On a more positive note, the efficiency of *P. fluorescens* in controlling the pathogen *Aremonas* has been confirmed [24]. The genus of *Pseudomonas* is presented in both Tilapia and Clarias RAS (Figure 3). The bacterial species *P. fluorescens* is a rhizobacteria in the genus *Pseudomonas* which has been identified in our samples from both RAS (Table 2). Optimizing conditions to favor beneficial microbes in a RAS and in the root environment in aquaponic systems might thus be used as a strategy to control fish diseases caused by pathogens in such aquaponic systems.

With regard to the fungal communities in our sampled RAS, we found that they included genera with known important roles as promoters of plant growth and antagonists against plant pathogens. These genera include *Cladosporium*, *Mortierella* [47] and *Trichoderma* [48] in Clarias water, *Pleospora* [49] in Clarias wastes; and *Acremonium* [50] in Tilapia water and *Penicillium* [47] in Tilapia waste samples (Figures 12 and 13). However, these systems also contained genera with pathogenic potential concerning human and plant health, such as *Fusarium* [50], *Candida* and *Cryptococcus*, which pose a more significant challenge that needs further investigation. In addition, further investigations on

fungus primers are also needed and important in order to enable the identification of those fungi in the current study that were unclassifiable beyond the level of phylum and genus.

The strains of bacteria isolated from the Tilapia system demonstrated biocontrol potential against plant pathogens hinting at their having a potential for an applied role in aquaponic systems. The only isolated bacterial strain that was positively identified and present in all samples was *P. fluorescens*. However, the log transformation of the *Pseudomonas* data in our study indicated the presence of these genera in all the investigated samples (Figure 3) and thereby strengthens our findings concerning the isolated strains. Others less ubiquitous taxa were *P. veronii* and *Vaiovorax paradoxus* in Tilapia wastes, and *Variovorax* spp. in Clarias wastes (Table 2). *Pseudomonas* spp. are essential microorganisms that can produce antimicrobial substances that control plant pathogens and promote plant growth [18, 20]. *P. veronii* is a species with bioremediation potential and is found in soil and water environments [51]. *Vaiovorax paradoxus* and *Variovorax* spp. are also known to have the ability to promote plant growth [52].

Enzyme production is one of the modes of action used by microorganisms to combat pathogen attack [20]. In aquaculture, the microbial communities of probiotic nature such as *Actinobacteria*, *Proteobacteria* and *Bacteroidetes* use different enzyme activities such as protease, chitinase, glucanase, amylase, cellulase, and phosphatase to break down different nutritional components in their substrates [53]. Our results in the current study indicate the production of protease, phosphatase, cellulase and siderophore by most of the strains isolated from the Tilapia system (Table 2). *P. veronii* (number 4) isolated from Tilapia wastes exhibited an ability to produce all of the above-mentioned enzymes. *P. fluorescens* isolates from Clarias water tanks (number 6) and Tilapia wastes (number 3) were also able to produce most of the enzymes. *Pseudomonas* spp. isolated from both RAS systems are well-known for siderophore production in plant roots that improves the availability of iron by aiding its uptake by plant roots [54]. Thus, the isolates in our study can be good candidates for use in aquaponic systems to increase the availability of phosphorous and iron for plant growth.

In vitro investigations into the antagonistic abilities of the isolated strains highlighted differences in their inhibitory effects on the growth of pathogens. *P. fluorescens* (number 6), *P. fluorescens* (number 3) and *P. veronii* (number 4) suppressed the growth of the plant root pathogen, *Phytophthora cactorum* (Pc) more than the other strains (Figure 14). This difference might be attributed to the production of cellulase by these strains, which not only breaks down a component in the feed, but can also break down cell walls of Pc. However, an *in vivo* antagonistic effect is needed in aquaponics in order for these results to have practical application. Our detached leaf assay provides strong evidence of the antagonistic potential of *P. veronii* (number 4) and *P. fluorescens* (number 6) against Pc (Figure 15).

5. Conclusions

The current study has contributed new knowledge concerning the role of microbial tools in commercially based recirculated aquaculture systems (RAS) as promoters of plant growth and suppressors of disease. This knowledge strengthens the potential application of RAS as a part of aquaponic systems, which currently face challenges regarding plant nutrients and pathogens. The assemblages of microbial taxa at the level of phyla and genera both in Tilapia and Clarias RAS suggest the suitability of these systems to be used in aquaponic cultivation. However, in terms of promoting plant growth, plant protection and biodegradation, the characteristics of the richness and composition of the microbial communities in the Tilapia system make it the better choice for application in aquaponic systems. The commercial Clarias RAS could also be applied in aquaponic systems, but principally from a probiotics perspective due to the dominance of *Actinobacteria* in this system. *Pseudomonas* spp. from both Tilapia tank water and waste samples, and Clarias wastes, are good candidates with the potential to produce extracellular enzymes that enhance nutrient uptake. Although the results suggest considerable potential for using microbial communities to manage and control certain aspects of aquaponic systems, our findings need to be strengthened with *in vivo* studies to explore further the inhibition of plant pathogens and positive effects on plant nutrition. Risks arising from the presence of pathogens also need further investigation. However, our results can still be used as a foundation for the design of aquaponic systems populated with either Tilapia

or Clarias. Still, these results need further investigations considering the microbial patterns in the RAS of Tilapia and Clarias in relation to the abiotic and biotic factors in the system.

Author Contributions: Sammar Khalil: Funding acquisition; design, and performance of the experiments with respect to microbial assessments and enzyme activities; writing - original draft preparation. Preeti Panda: Metagenomic analyses; writing - review and editing the manuscript. Ramesh Vetukuri: Strain identification; *in vitro* investigations; writing - review and editing the manuscript. Farida Ghadamgahi: *In vitro* and DLA investigations; writing - review and editing the manuscript. AnnaKarin Rosberg: DNA extraction, comments on the manuscript.

Acknowledgments: Richterska Stiftelsen funded this research at <https://www.sweco.se/om-oss/priser-och-stipendier/richterska-stiftelsen/>, which is greatly acknowledged. The staff at the fish company where the samples have been collected are also acknowledged for their help.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kumm, M.; De Moel, H.; Porkka, M.; Siebert, S.; Varis, O.; Ward, P.J. Lost food, wasted resources: Global food supply chain losses and their impacts on freshwater, cropland, and fertilizer use. *Sci. Total Environ.* **2012**, *438*, 477–489.
2. The State of Food and Agriculture: Leveraging Food Systems for Inclusive Rural Transformation; Food and Agriculture Organization of the United Nations: Roma, Italy, **2017**, ISBN 9251098735.
3. Ga, U.N. Transforming our World: The 2030 Agenda for Sustainable Development; Division for Sustainable Development Goals: **2015**, New York, NY, USA.
4. FAO (Food and Agriculture Organization of the United Nations). The State of World Fisheries and Aquaculture. **2016**, Available online: <http://www.fao.org/3/a-i5555e.pdf>.
5. Martins, C.I.M.; Eding, E.H.; Verdegem, M.C.J.; Heinsbroek, L.T.N.; Schneider, O.; Blancheton, J.P.; Roque d'Orbcastel, E.; Verreth, J.A.J. New developments in recirculating aquaculture systems in Europe: a perspective on environmental sustainability. *Aquacul. Engin.* **2010**, *43* (3), 83–93.
6. Goddek, S.; Delaide, B.; Mankasingh, U.; Ragnarsdottir, K.V.; Jijakli, H.; Thorarinsdottir, R. Challenges of sustainable and commercial aquaponics. *Sustain.* **2015**, *7*, 4199–4224. <https://doi.org/10.3390/su7044199>
7. Endut, A.; Jusoh, A.; Ali, N.; WanNik, W.B.; Hassan, A. A. Study on the optimal hydraulic loading rate and plant ratios in recirculation aquaponic system. *Bioresour Technol.* **2010**, *101*, 1511–1517.
8. Yildiz, H.Y.; Robaina, L.; Pirhonen, J.; Mente, E. Fish Welfare in Aquaponic Systems: Its Relation to Water Quality with an Emphasis on Feed and Faeces—A Review. *Water.* **2017**, *9*, 13, doi:10.3390/w9010013
9. Domínguez, D.; Parisi, G.; Monsees, H.; Kloas, W.; Wuertz, S. Decoupled systems on trial: eliminating bottlenecks to improve aquaponic processes. *PLoS One.* **2017**, *12*(9), <https://doi.org/10.1371/journal.pone.0183056>
10. Postma, J.; van Os, E.; Bonants, P.J.M. Ch 10 – pathogen detection and management strategies in soilless plant growing system. In: *Soilless culture: theory and practice*. Elsevier B.V., Amsterdam, **2008**, 425–457. <https://doi.org/10.1016/B978-0-444-52975-6.50012-5>
11. Alhussaen, K. *Pythium* and *Phytophthora* associated with root disease of hydroponic lettuce. University of Technologie Sydney Faculty of Science. **2006**, <https://opus.lib.uts.edu.au/handle/10453/36864>
12. Fujiwara, K.; Aoyama, C.; Takano, M.; Shinohara, M. Suppression of *Ralstonia solanacearum* bacterial wilt disease by an organic hydroponic system. *J Gen Plant Pathol.* **2012**, <https://doi.org/10.1007/s10327-012-0371-0>.
13. Ito, S.; Takikawa, Y.; Tairako, K. Black rot of pak-choi in hydroponics caused by *Xanthomonas campestris* pv. *Campestris*. *Japan. J. of phytopath.* **2003**, DOI: 10.3186/JPHYTOPATH.69.407.
14. Calvo-bado, L.A.; Pettitt, T.R.; Parsons, N.; Petch, G.M.; Morgan, J.A.W.; Whipps, J.M. Spatial and Temporal Analysis of the Microbial Community in Slow Sand Filters Used for Treating Horticultural Irrigation Water. *Appl. Envir. Microbiol.* **2003**, *69*, 2116–2125.
15. Chatterton, S.; Sutton, J.C.; Boland, G. J. Timing *Pseudomonas chlororaphis* applications to control *Pythium aphanidermatum*, *Pythium dissotocum*, and root rot in hydroponic peppers. *Biol Cont.* **2004**, *30*, 360–373.

16. Cherif, M.; Tirilly, Y.; Belanger, R.R. Effect of oxygen concentration on plant growth, lipid peroxidation, and receptivity of tomato roots to *Pythium* under hydroponic conditions. *Eur J Plant Pathol.* **1997**, *103*, 255–264.
17. Ehret, D.L.; Alsanius, B.; Wohanka, W.; Menzies, J.G.; Utkhede, R. Disinfestation of recirculating nutrient solutions in greenhouse horticulture. *Agronomie* **2001**, *21*, 323–339.
18. Renault, D.; Déniel, F.; Benizri, E.; Sohier, D.; Barbier, G.; Rey, P. Characterization of *Bacillus* and *Pseudomonas* strains with suppressive traits isolated from tomato hydroponic-slow filtration unit. *Can J Microbiol.* **2007**, *53*, 784–797.
19. Renault, D.; Vallance, J.; Déniel, F.; Wery, N.; Godon, J.J.; Barbier, G.; Rey, P. Diversity of bacterial communities that colonize the filter units used for controlling plant pathogens in soilless cultures. *Microb Ecol.* **2012**, *63*, 170–187.
20. Mazzola, M. Assessment and management of soil microbial community structure for disease suppression. *An. Rev. Phytopath.* **2004**, *42*, 35–59.
21. Zeng, Q.; Tian, X.; Wang, L. Genetic adaptation of microbial populations present in high-intensity catfish production systems with therapeutic oxytetracycline treatment. *Sci. Rep.*, **2017**, *7*, 1–13.
22. Eck, M.; Sare, A.R.; Massart, S.; Schmutz, Z.; Junge, R.; Smits, T. H. M.; Jijakli, M. H.. Exploring Bacterial Communities in Aquaponic Systems. *Water.* **2019**, *11*, 260, doi:10.3390/w11020260.
23. Schmutz, Z.; Graber, A.; Jaenicke, S.; Goesmann, A.; Junge, R.; Smits, T. H. M. Microbial diversity in different compartments of an aquaponics system. *Arch of Microbiol.* **2016b**, 613–620.
24. Eissa, N.; Abou El-Ghiet, E.N. Efficacy of *Pseudomonas fluorescens* as biological control agents against *Aeromonas hydrophila* infection in *Oreochromis niloticus*. *World J. Fish Mar. Sci.* **2012**, *3*, 564–569.
25. Khalil, S.; Alsanius, B.W. Dynamics of the indigenous microflora inhabiting the root zone and the nutrient solution of tomato in a commercial closed greenhouse system. *Gartenbauwissen* **2001**, *66*, 188–198.
26. Darlison, J.; Mogren, L.; Rosberg, A.; Grudén, M.; Minet, A.; Line, C.; Mieli, M.; Bengtsson, T. Leaf mineral content govern microbial community structure in the phyllosphere of spinach (*Spinacia oleracea*) and rocket (*Diplotaxis tenuifolia*). *Scien. Total. Envir.* **2019**, 675, DOI: 10.1016/j.scitotenv.2019.04.254
27. Dhariwal, A.; Chong, J.; Habib, S.; King, I.L.; Agellon, L.B.; Xia, J. Microbiome Analyst: a web-based tool for comprehensive statistical, visual and metaanalysis of microbiome data. *Nucleic Acids Res.* **2017**, *45*, W180eW188. <https://doi.org/10.1093/nar/gkx295>.
28. Morales, S.E.; Holben, W.E. Empirical testing of 16S rRNA gene PCR primer pairs reveals variance in target specificity and efficacy not suggested by in silico analysis. *Appl. Envir. Microbiol.* **2009**, *75*, 2677–2683.
29. Altschul, S.F.; Madden, T.L.; Schäffer, A.; et al. Gapped BLAST and PS I-BLAST: a new generation of protein database search programs. *Nucl. Acid. Res.* **1997**, *25*, 3389–3402. doi: 10.1093/nar/25.17.3389
30. Choudhary, D. K.; Agarwal, P. K.; Johri, B. N. Characterization of functional activity in composted casing amendments used in cultivation of *Agaricus bisporus* (Lange) Imbach. *Ind. J. of Biotech* **2009**, *8*(1), 97–109.
31. Rahman, M. A.; Begum, M. F.; Alam, M. F. Screening of *Trichoderma* Isolates as a Biological Control Agent against *Ceratocystis paradoxa* Causing Pineapple Disease of Sugarcane. *Mycobiol.* **2009**, *37*(4), 277. doi: 10.4489/MYCO.2009.37.4.277.
32. Jimoh, W.A.; Sulyman, T.; Taiwo, A.T. Microbial load and diversity in the gastro-intestinal tract of cultured Nile tilapia (*Oreochromis niloticus*) and hybrid catfish (*Clarias gariepinus* ♀ x *Heterobranchius bidorsalis* ♂) in Ilorin Metropolis, Nigeria. *Int. J. Aquatic Biol*, **2019**, *7*(5), 271–279, DOI: <https://doi.org/10.22034/ijab.v7i5.672>
33. Minich, J. J.; Zhu, O.; Xu, Z.Z.; Amnon, A.; Ngochera, M.; Simwaka, M.; Allen, E.E.; Zidana, H.; Knight, R. Microbial effects of livestock manure fertilization on freshwater aquaculture ponds rearing Tilapia (*Oreochromis shiranus*) and North African catfish (*Clarias gariepinus*). *Microbiol Open*, **2018**, <https://doi.org/10.1002/mbo3.716>
34. Giatsis, C.; Sipkema, D.; Smidt, H.; Heilig, H.; Benvenuti, G.; Verreth, J.; Verdegem, M. The impact of rearing environment on the development of gut microbiota in tilapia larvae. *Scientific Reports*, **2015**, *5*, 18206.
35. Liu, S.; Ren, H.; Shen, L.; Lou, L.; Tian, G.; Zheng, P.; Hu, B. pH levels drive bacterial community structure in sediments of the Qiantang River as determined by 454 pyrosequencing. *Front Microbiol* **2015**, *6*: 285.
36. Campbell, B.J.; Kirchman, D.L. Bacterial diversity, community structure and potential growth rates along an estuarine salinity gradient. *SME J.* **2013**, *7*, 210–220.
37. Schreier, H.J.; Mirzoyan, N.; Saito, K. Microbial diversity of biological filters in recirculating aquaculture systems. *Cur. Opin. Biotech.* **2010**, *21*, 318–325.

38. Garcia, J. C.; Ketover, R. D. J.; Loh, A. N.; Parsons, M. L.; Urakawa, H. Influence of freshwater discharge on the microbial degradation processes of dissolved organic nitrogen in a subtropical estuary. *Antonie Van Leeuw.* **2015**, 107, 613–632.
39. Das, S.; Ward, L.R.; Burke, S. Prospects of using marine actinobacteria as probiotics in aquaculture. *Appl Microbiol Biotechnol* **2008**, 81:419–429 DOI 10.1007/s00253-008-1731-8.
40. Jami, M.; Ghanbari, M.; Kneifel, W.; Domig, K.J. Phylogenetic diversity and biological activity of culturable Actinobacteria isolated from freshwater fish gut microbiota. *Microbiol. Res.* **2015**, 175, 6-15.
41. Sanchez, F.A.; Vivian-Rogers, V.R.; Urakawa, H. Tilapia recirculating aquaculture systems as a source of plant growth promoting bacteria. *Aquacul Res.* **2019**, 50, 2054–2065, DOI: 10.1111/are.14072.
42. Sirakov, I.; Lutz, M.; Graber, A.; Mathis, A.; Staykov, Y.; Smits, T.H.M.; Junge, R. Potential for combined biocontrol activity against fungal fish and plant pathogens by bacterial isolates from a model aquaponic system. *Water* **2016** 8:518.
43. Guo, X.; Miao, Y.; Wu, B.; Ye, L.; Yu, H.; Liu, S.; Zhang, X-x. Correlation between microbial community structure and biofouling as determined by analysis of microbial community dynamics. *Bioresource Techn.* **2015**, 197, 99–105.
44. Albertsen, M.; McIlroy, S.J.; Stokholm-Bjerregaard, M.; Karst, S.M.; Nielsen, P.H. Candidatus *Propionivibrio aalborgensis*: A Novel Glycogen Accumulating Organism Abundant in Full-Scale Enhanced Biological Phosphorus Removal Plants. *Front. in Microb.* **2017**, 1033, doi: 10.3389/fmicb.2016.01033.
45. Shu, D.; He, Y.; Yue, H.; Wangd, O. Microbial structures and community functions of anaerobic sludge in six full-scale wastewater treatment plants as revealed by 454 high-throughput pyrosequencing. *Bioresource Techn.* **2015**, 186, 163–172.
46. Martins, P.; Cleary, D. F. R.; Pires, A. C. C.; Rodrigues, AM.; Quintino, V.; Calado, R.; Gomes, N.C. M. Molecular Analysis of Bacterial Communities and Detection of Potential Pathogens in a Recirculating Aquaculture System for *Scophthalmus maximus* and *Solea senegalensis*. *Plos One.* **2013**, 8 (11), e80847.
47. Tagawa, M.; Tamaki, H.; Manome, A.; Koyama, O.; Kamagata, Y. Isolation and characterization of antagonistic fungi against potato scab pathogens from potato field soils. *FEMS Microbiol Lett.* **2010**, 305, 136–142.
48. Vinalea, F.; Sivasithamparamb, K.; Ghisalberti, E.L.; Marraa, R.; Wooa, S.L.; Lorito, M. Trichoderma-plant-pathogen interactions. *Soil Biol. & Bioch.* **2008**, 40, 1–10.
49. Bailey, B.A.; Apel-Birkhold, P.C.; Akingbe, O. O.; Ryan, J. L.; O'Neill, N. R.; Anderson, J.D. Nep1 Protein from *Fusarium oxysporum* Enhances Biological Control of Opium Poppy by *Pleospora papaveracea*. *Phytopathol.* **2000**, 90 (8), 812-818.
50. Auer, S.; Ludwig-Müller, J. Biological control of clubroot (*Plasmodiophora brassicae*) by the endophytic fungus *Acremonium alternatum*. *J. Endocyt. and Cell Res.* **2015**, 26, 43-49.
51. Vullo, D.L.; Ceretti, H.M.; Daniel, M.A.; Ramírez, S.A.M.; Zalts, A. Cadmium, zinc and copper biosorption mediated by *Pseudomonas veronii* 2E. *Bioresour Technol* **2008**, 99: 5574–5581.
52. Han, J.I.; Choi, H.K.; Lee, S.W.; Orwin, P.M.; Kim, J.; LaRoe, S.L.; Kim, T.G.; O'Neil, J.; Leadbetter, J.R.; Lee, S.Y.; Hur, C.-G.; Spain, J.C.; Ovchinnikova, G.; Goodwin, L.; Han, C. Complete Genome Sequence of the Metabolically Versatile Plant Growth-Promoting Endophyte *Variovorax paradoxus* S110. *J. of bacterial.* **2011**, 193 (5), 1183–1190, doi:10.1128/JB.00925-10.
53. Dean, R.A.; Timberlake, W.E. Production of Cell Wall-Degrading Enzymes by *Aspergillus nidulans*: A Model System for Fungal Pathogenesis of Plants. *The Plant Cell* **1989**, 1, 265-273.
54. Djibaoui, R.; Bensoltane, A. Effect of iron and growth inhibitors on siderophores production by *Pseudomonas fluorescens*. *Afric J. Biotechnol.* **2005**, 4 (7), 697-702.