

Case Report

How fear of COVID-19 can affect treatment choices for Anaplastic Large Cell Lymphomas ALK+ therapy: a case report

Antonello Sica^{1, ¶*}, Caterina Sagnelli^{2, ¶}, Beniamino Casale³, Gino Svanera⁴, Massimiliano Creta⁵, Armando Calogero⁶, Renato Franco⁷, Evangelista Sagnelli² and Andrea Ronchi⁷

¹ Department of Precision Medicine, University of Campania "Luigi Vanvitelli", 80131 Naples, Italy; antonello.sica@fastwebnet.it; antonellosica@gmail.com.

² Department of Mental Health and Public Medicine, University of Campania "Luigi Vanvitelli", 80138 Naples, Italy; caterina.sagnelli@unicampania.it, evangelista.sagnelli@unicampana.it.

³ Pain Department, AORN Dei Colli - V. Monaldi, 80131 Naples, Italy; benny.casale@hotmail.com

⁴ Department of Medical Area ASLNA2 NORTH, 80014 Giugliano, Naples, Italy; gsvanemato@inwind.it

⁵ Department of Neurosciences, Reproductive Sciences and Odontostomatology, University of Naples "Federico II", 80131 Naples, Italy; massimiliano.creta@unina.it.

⁶ Department of Advanced Biomedical Sciences, University of Naples Federico II, 80131 Naples, Italy; armando.calogero2@unina.it.

⁷ Pathology Unit, Department of Mental and Physical Health and Preventive Medicine, University of Campania "Luigi Vanvitelli", 80138 Naples, Italy; renato.franco@unicampania.it, andrea.rochi@unicampania.it.

* Correspondence: Dr. Antonello Sica (A.S), Department of Precision Medicine, University of Campania Luigi Vanvitelli, 80131 Naples, Italy, Tel: +393332253315, +3908119573375, e-mail: antonello.sica@fastwebnet.it; antonellosica@gmail.com

Abstract: Background: The t (2; 5) chromosomal rearrangement of ALK gene with Nucleophosmin1 gene (NPM1), resulting in NPM1-ALK fusion, was first demonstrated in 1994 in anaplastic large cell lymphoma, (ALCL), a T-cell lymphoma responsive to cyclophosphamide, abirblastine, vincristine and prednisone in approximately 80% of cases; refractory cases usually respond favorably to brentuximab-vedotin. These treatments are regarded as a bridge to allogeneic hematopoietic stem cell transplantation (allo-SCT). Nowadays, transplant procedures and monitoring of chemotherapy patients proceed very slowly because the SARS-CoV-2 pandemic has heavily clogged the hospitals in all countries.

Results: A 40-year-old Caucasian woman was first seen at our clinical center in June 2020. She had ALCL ALK +, a history of failure to two previous therapeutic lines and was in complete remission after 12 courses of Brentuximab, still pending allo-SCT after two failed donor selection. Facing of a new therapeutic failure, we requested the Italian drug regulatory agency, and obtained the authorization, to administer 250 mg twice a day of Crizotinib, a drug incomprehensibly not registered for ALCL ALK +. **Conclusions:** The response to Crizotinib was optimal, since no adverse event occurred, and CT-PET persisted negative; this drug has proved to be a valid bridge to allo-SCT.

Keywords: Crizotinib; Anaplastic Large Cell Lymphomas ALK+, bridge therapy in NHL ALK+, ALK+ patients; anticancer therapy

1. Introduction

The t (2; 5) chromosomal rearrangement and the resulting nucleophosmin (NPM1)-ALK fusion was first described in anaplastic large cell lymphoma (ALCL) in 1994 and further studies have shown that ALK rearrangements are present in approximately 55% of ALCL cases and in almost all affected

children. Considering that, it would be natural that ALK inhibitors were registered for this disease, but the Italian drug regulatory agency (Agenzia Italiana del Farmaco-AIFA) authorized these drugs only for treating Non-Small Cell Lung Cancer (NSCLC)[2]. The enormous efficacy of ALK inhibitors in the treatment of NSCLC has shown that these mutations behave like real oncogenes and therefore have completely replaced the chemotherapy previously practiced in these neoplasms. For ALCL, conventional chemotherapy has shown better efficacy than ALK inhibitors in all studies performed, which probably explains why AIFA has not submitted ALCL for approval. ALCL is a T-cell lymphoma that responds to CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) in about 80% of treated patients, while the anti-CD30 monoclonal antibody brentuximab-vedotin is usually beneficial in refractory cases [3-6]. These treatments are used as a bridge to hematopoietic allogeneic stem cell transplantation (allo-SCT) which is considered the definitive therapy for refractory cases.

Nowadays, due to SARS-CoV-2 pandemic, the strategy leading to allo-SCT or chemotherapy have changed in clinical practice [7]. In fact, the clogging of health facilities caused by this pandemic has slow down the procedures to access the allo-SCT and made complex the follow up of patients, either transplanted or under chemotherapy. These slowdowns require alternative therapeutic strategies to block the progression of the disease. Waiting for a suitable donor, we treated with crizotinib, a drug with has a specific action against ALK + and a proven specific efficacy, a 40-year-old Caucasian woman with ALCL and a history of failure to a first- and a second-line therapy [8-9]. This drug is not registered in Italy for this type of lymphoma, but its use for our patient was authored by AIFA. In addition to NPM1, many other fusion partners have been described: TPM3, TFG, ATIC, CLTC, TPM4, MSN, ALO17, MYH9, TRAF1 and TPM3 [10,11]. Current knowledge suggests that the presence of oncogenes prevails over the phenotype/ diagnosis. As with non-small cell lung cancer (NSCLC), fusion partner can influence the response to different ALK-TKIs, but we still don't know how it works in other clinical situations with ALK rearrangement [12]. Several clinical trials are currently underway: alectinib for relapsed or refractory anaplastic lymphoma kinase-positive anaplastic large cell lymphoma [13], lorlatinib for refractory ALK positive ALCL (NCT03505554) and for refractory neuroblastoma (NCT03107988) and brigatinib for ALK positive ALCL (NCT03719898). These studies will lead to better understanding how these ALK-TKIs work in ALCL or other solid tumors with ALK rearrangement.

Current knowledge indicates that Crizotinib can be effective in pediatric patients with ALK positive ALCL [14] and in infantile hemispheric glioma with ALK -rearrangement [15]; this drug was also effective in an adult ALK-positive-patient with advanced stage ganglioneuroblastoma. Such favorable effects of the ALK inhibitor crizotinib led us to choose this drug to create a safe bridge to allo-SCT for our adult patient with ALCL ALK+ no more responder to several therapies; this was an excellent choice.

2. Case report

A 40-year-old Caucasian woman with Anaplastic Large Cell Lymphoma ALK+ (ALCL ALK+) in complete remission (CR) after 16 cycles of therapy with Brentuximab (1,8mg/kg) was first observed at our clinical center in June 2020, a treatment scheduled pending allo-SCT.

The diagnosis of ALCL ALK+, was made with the microscopic examination of a latero-cervical lymph node surgically removed on October 2016, also using immunohistochemistry and fluorescence in situ hybridization (FISH). Immunohistochemistry showed intense and diffuse positivity in neoplastic cells, allowing the diagnosis of anaplastic large cell lymphoma, ALK-positive. Moreover, we indagated the specific fusion partner using a Two Color, Two Fusion Translocation Probe, designed to detect the translocation between the ALK gene located at 2p23 and the NPM1 gene located at 5q35. The patient was stage IV according to the Ann Arbor classification (Figure 1A), and was treated with CEOP (cyclophosphamide, epirubicin, vincristine, prednisone) for 6 cycles. In March 2017, as soon as these cycles finished, she had an inguinal recurrence with presence of blasts of meningeal lymphoma. From April to August 2017, she received a second-line therapy with MTX-ARAC-THIOthepa (MAT: methotrexate, cytarabine, Thiothepa) for 4 cycles plus 6 rachicentesis with infusion of MTX, ARAC and steroids. Reaching remission, she underwent peripheral blood stem cells

(PBSC) mobilization chemotherapy according to the idarubicin, ARA-C, filgrastim scheme plus Plerixafor. In October 2017, she practiced PBSC apheresis and in November 2017 therapeutic consolidation with autologous stem cell transplantation (ASCT), after high-dose chemotherapy (HDT) with thiotepa-busulfan (Thio-Bu) regimen ([tab.1](#)), with complete remission (Figure 1 B). A voluminous lymphadenopathy appeared in the right armpit at the end of May 2019, greatly reduced with paracetamol, we interpreted as an early sign of progression, oddly improved with NSAID. In July 2019, the patient developed mild systemic low-grade irregular fever and slight weight loss; she performed all blood chemistry tests that showed: ESR 20. Ferritin 58; β 2-microglobulin 1.32; LDH 366; Hgb 12.5 gr/dL; PLT $198,000 \times 10^3/\text{mmc}$; GB $4,800 \times 10^3/\text{mmc}$; 49%; Leucocyte 37%; Monocyte 9%. As the axillary lymphadenopathy persisted, a biopsy was performed, and the histological examination showed a new relapse of the disease (Figure 1 C).

Consequently, in July 2019 she was treated with Brentuximab, performed for 16 cycles until June 2020 (Figure 1 D). Before starting this therapy, it was necessary to add tramadol 150 mg per day orally as analgesic therapy, because she reported continuous pain resistant to paracetamol in the right armpit [17-18].

Since CR was achieved after 6 months of this third line of treatment, the donor selection for allo-SCT began. A haploidentical male cousin was summoned in December 2019 to undergo the first level investigations to be admitted to donation: HCV-Ab, HBsAg, HBcAb, HIV-Ab, blood count, creatinine, BUN, AST, ALT, amylase, lipase, bilirubin, electrolytes, triglyceride, cholesterol and glycemy, serum protein electrophoresis, LDH [19-24].

Unfortunately, he was found EBV IgG positive, with anti-HLA, donor mismatched on locus A and B (6/8) and therefore not completely suitable according to the criteria of the Italian Bone Marrow Donor Registry. During the SARS-CoV-2 pandemic, the search for donors has slowed down a lot, mostly due to the fear of donors and patients accessing hospitals. In this period, aplasticizing therapies for transplants were considered extendable up to 8-12 weeks for non-aggressive or non-life-threatening diseases. These extensions have been suggested in the hope of a future strong reduction in the circulation of the virus. A lower likelihood of SARS-CoV-2 infection is certainly desirable for patients for whom strong immunosuppressive therapy is required. In subjects awaiting allogeneic transplantation, conditioning chemotherapy should be started only after the arrival of the donor cells and cryopreservation. As the waiting time for allogeneic transplantation, however indispensable for our patient, was extended, all possible courses of brentuximab were performed. Of note, the schedule for administering brentuximab to patients with stabilized or improving disease suggests a maximum of 16 cycles to be practiced in approximately one year. The attempt to select a suitable donor has continued, but with no success, due to the significant shortage of donors in this prolonged SARS-CoV-2 pandemic, when many people, including organ donors, fearing contagion avoid entering hospitals. In the, meanwhile, AIFA approval was requested to administer Crizotinib 250 mg twice daily, a drug not registered for treating ALCL ALK+ in Italy. The approval was given on February 2020 and treatment began one months after the last dose of brentuximab. The response to the new treatment was optimal since no side effect have occurred and the [18f] FDG-Positron Emission Tomography/Computed Tomography ([18f] FDG-PET-CT) performed in September 2020 was negative (Figure 1 D).

A haploidentical female cousin was summoned to undergo the first level investigations (HIV-Ab, CMV-Ab, HCV-Ab, HBsAg, HBcAb, EBV-Ab, blood count, creatinine, BUN, AST, ALT, amylase, lipase, bilirubin, electrolytes, triglyceride, cholesterol and glycemy, serum protein electrophoresis) to be admitted to donation in October 2020 [25-28]. Unfortunately, she was EBV-Ab IgM positive and therefore suspended from donations. Continuing the complete remission, it was decided to attempt the collection of autologous peripheral stem cells free from disease, for a possible second approach with HDT, an attempt however unsuccessful.

The potential donor (the female cousin) was again summoned for the end of November 2020. Meanwhile, Crizotinib therapy was continued and AIFA's approval requested for its continuation. Pending the authorization of AIFA, we plan to continue crizotinib therapy until allo-SCT will be performed, provided that the response to this drug continues to be good. Although we do not have

any data on allo-SCT after this targeted therapy with ALK-TKI, we also do not have data that advise against this treatment before allo-SCT. Therefore it seems the best strategy that can be used right now.

3. Statement of Ethics

All procedures performed were in accordance with the international guidelines, with the Helsinki Declaration of 1975, revised in 1983 and the roles of the Italian laws of privacy. Crizotinib treatment was approved by AIFA (Agenzia Italiana del Farmaco) on 07 February 2020.

Patient signed an anonymous informed consent for the use of his data for anonymous clinical investigations and scientific publications.

4. Discussion

The time we live in is particularly dramatic worldwide due to the SARS-CoV-2 pandemic. Hospital assistance is in trouble everywhere and everyday hospital structures previously used for the various sectors of medicine are adapted into centers for the reception of patients affected by SARS-CoV-2. There are severe limitations to outpatient and day hospital activities and in many local situations the access to hospitals for neoplastic or otherwise immunosuppressed patients is prevented or severely slowed for fear of contagion. Equally slowed down is the process for accessing organ transplantation and transplant operations have become infrequent and limited to situations of emergency [29-30].

In Italy, these conditions are aggravated by a prolonged organizational inefficiency of the healthcare governing institutions, which have caused a substantial reduction in the number of doctors, nurses and all other categories of health workers and a marked reduction in the number of beds, conditions known from years, but dramatically enhanced by the SARS-CoV-2 pandemic. In these conditions it is complex to put into practice the indications of international guidelines and to avoid that the infection by SARS-CoV-2 increases the risk of mortality in neoplastic patients.

After two relapses, our patient, achieved complete remission of the disease with Brentuximab used as a third-line therapy. According to international guidelines, patients with ALCL ALK +, once a CR at a rescue therapy has been obtained, must undergo allo-SCT which could have a high chance of success [31-35]. The young age of the patient and her refractory to the first- and second-line treatments prompted us to proceed toward allo-SCT, so far without success due to the unsuitability of the contacted donors [36-39]. Although in fair condition after 16 cycles of Brentuximab, fearing that even this third-line treatment would fail while waiting for a suitable donor, we decided to use Crizotinib [40-43], a drug that in Italy is not registered for this type of lymphoma, but with a specific action against ALK + and effective in published anecdotal cases.


Already in 2013, Gambacorti Passerini described encouraging results obtained with crizotinib in patients affected by ALCL ALK + and relapsed in multiple therapeutic lines. Since then, however, the trend was to continue using the drugs used before for other neoplasms and only recently few cases of ALCL ALK + treated with crizotinib have been described [44-45].

5. Conclusions

In Conclusion, as regards our patient, the fear of a failure of the third-line therapy (Brentuximab), the difficulties encountered in identifying a compatible donor and the poor stability of the hospital network during this pandemic led us to treat our patient with crizotinib 500 mg/day. Excellent results have been obtained, a persistent CR and an excellent clinical condition both witnessed by a perfectly normal [18f] FDG-PET-CT, and an excellent psycho-emotional condition that sustains her while waiting for all-SCT.

Our judgment on the use of crizotinib therapy in ALCL ALK + treated is excellent and we believe it should be included in guidelines as a bridge to allo-SCT.

6. Figure legend



A) [18f] FDG-PET-CT scans performed at October 2016: the examination showed pathologies with a high glucose metabolism in correspondence with multiple lymph node packets in the left supraclavicular region, retroclavare left and right, left retrocrural, lombo-aortica, intercavale, paracavale, the iliac bifurcation and along the path of the internal inguinal vessels and right exteriors.

B) [18f] FDG-PET-CT scans performed at January 2018: the examination don't show pathologies with a high glucose metabolism.

C) [18f] FDG-PET-CT scans performed at July 2019: the examination show pathologies with a high glucose metabolism in correspondence of numerous lymphadenomegalies in the right axillary region and of the lymph nodes in the right lung perillary area.

D) [18f] FDG-PET-CT scans performed at June 2020: the examination don't show pathologies with a high glucose metabolism.

E) [18f] FDG-PET-CT scans performed at September 2020: the examination don't show pathologies with a high glucose metabolism.

Table.1: Treatment timeline

PERIOD	THERAPY
From October 2016 to February 2017	CEOP 6 cycles
From April 2017 to August 2017	MTX-ARAC-THIOthepa 4 cycles plus 6 rachi-centesis with infusion of MTX, ARAC and steroids
From October 2017 to November 2017	ASCT after HDT with Thio-Bu
From July 2019 to June 2020	Brentuximab 16 cycles
From July 2020 to present	Crizotinib 250 mg twice daily

Author Contributions: Conceptualization, methodology, investigation, formal analysis, writing—original draft preparation, writing—review and editing, validation, visualization, S.A., C.S., and E.S.; data curation, B.C, G.S., M.C, A.C, R.F, and A.R.; validation, visualization, B.C, G.S., M.C, A.C, R.F, and A.R.; supervision, S.A., C.S., and E.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: All the authors of the manuscript declare they have no conflict of interest in connection with this paper.

References

1. Sica, A.; Vitiello, P.; Ronchi, A.; Casale, B.; Calogero, A.; Sagnelli, E.; Costa Nachtigal, G.; Troiani, T.; Franco R.; Argenziano G.; Moscarella E.; Sagnelli, C. Primary Cutaneous Anaplastic Large Cell Lymphoma (pcALCL) in the Elderly and the Importance of Sport Activity Training. *Int J Environ Res Public Health*. **2020**;17(3):839. doi: 10.3390/ijerph17030839.
2. Fiorelli, A.; D'Andrilli, A.; Carlucci, A.; Vicidomini, G.; Loizzi, D.; Ardò, NP.; Marasco, RD.; Ventura, L.; Ampollini, L.; Carboognani, P.; Bocchialini G.; Lococo, F.; Paci M.; Di Stefano, TS.; Ardissonne, F.; Gagliasso, M.; Mendogni, P.; Rosso, L.; Mazzucco, A.; Vanni, C.; Marinucci BT.; Potenza, R.; Matricardi, A.; Ragusa, M.; Tassi, V.; Anile, M.; Poggi, C.; Serra, N.; Sica, A.; Nosotti, M.; Sollitto, F.; Venuta, F.; Rendina, E.A.; Puma, F.; Santini, M. Prognostic factors of lung cancer in lymphoma survivors (the LuCiLyS study). LuCiLyS study group. *Transl Lung Cancer Res*. **2020**;9(1):90-102. doi: 10.21037/tlcr.2019.12.28.
3. Sica, A.; Vitiello, P.; Sorriento, A.; Ronchi, A.; Calogero, A.; Sagnelli, C.; Troiani, T.; Fasano, M.; Dodaro, C.A.; Franco, R.; Casale, B.; Santangelo, M.; Ciccozzi, M.; Ciardiello, F.; Argenziano, G.; Moscarella E. Lymphomatoid papulosis. *Minerva Med*. **2020**;111(2):166-172. doi: 10.23736/S0026-4806.19.06395-X.
4. Caccavale, S.; Vitiello, P.; Franco, R.; Panarese, I.; Ronchi, A.; Sica, A.; Jurakic Toncic, R.; Alfano, R.; Argenziano, G. Dermoscopic characterization of folliculotropic mycosis fungoides selectively localized on trunk and limbs. *Int J Dermatol*. **2019**;58(10):e187-e189. doi: 10.1111/ijd.14490.
5. Sica, A.; Casale, B.; Spada, A.; Di Dato Teresa M.; Sagnelli, C.; Calogero, A.; Buonavolontà, P.; Salzano, A.; Martinelli, E.; Saracco, E.; Troiani, T.; Dodaro C.A.; Tammamaro, D.; Luisa De Rimini, M.; Ciardiello, F.; Papa, A. Differential Diagnosis: Retroperitoneal Fibrosis and Oncological Diseases. *Open Med (Wars)*. **2019**;15:22-26. doi: 10.1515/med-2020-0005.
6. Cascone, R.; Sica, A.; Sagnelli, C.; Carlucci, A.; Calogero, A.; Santini, M.; Fiorelli, A. Endoscopic Treatment and Pulmonary Rehabilitation for Management of Lung Abscess in Elderly Lymphoma Patients. *Int J Environ Res Public Health*. **2020**;17(3):997. doi: 10.3390/ijerph17030997.
7. Sica A.; Casale, D.; Rossi, G.; Casale, B.; Ciccozzi, M.; Fasano, M.; Ciotti, M.; Sagnelli, E.; Papa, A.; Sagnelli, C. The impact of the SARS-CoV-2 infection; with special reference to the haematological setting. *J Med Virol*. **2020**. doi: 10.1002/jmv.26197.
8. Gambacorti Passerini et Al. Crizotinib in Advanced; Chemoresistant Anaplastic Lymphoma Kinase – Positive Lymphoma Patients. (JNCI Natl Cancer Inst (2014) 106 (2): djt378 doi: 10.1093/jnci / djt378; Long-term effects of crizotinib in ALK-positive tumors (excluding NSCLC): A phase 1b open-label study. *Am J Hematol*. **2018**; 93: 607-614.
9. Calogero, A.; Sagnelli, C.; Carlomagno, N.; Tammamaro, V.; Candida, M.; Vernillo, A.; Peluso, G.; Minieri, G.; Santangelo, M.; Dodaro, C.A.; et al. Familial polyposis coli: The management of desmoid tumor bleeding. *Open Med (Wars)*. **2019**;14:572-576. DOI: 10.1515/med-2019-0064.
10. Tsuyama, N.; et al. Anaplastic large cell lymphoma: pathology, genetics, and clinical aspects. *Journal of clinical and experimental hematopathology*. **2017**,57(3), 120-142.
11. Hallberg, B.; & Palmer, R.H.; The role of the ALK receptor in cancer biology. *Annals of Oncology* 27 (Supplement 3): iii4–iii15, 2016, doi:10.1093/annonc/mdw301.
12. Childress, MA.; Himmelberg, S.M.; Chen, H.; Deng, W.; Davies, M.A.; Lovly, C.M. ALK Fusion Partners Impact Response to ALK Inhibition: Differential Effects on Sensitivity, Cellular Phenotypes, and Biochemical Properties. *Mol Cancer Res*. **2018**;16(11):1724-1736. doi: 10.1158/1541-7786.MCR-18-0171.
13. Fukano, R.; Mori, T.; Sekimizu, M.; Choi, I.; Kada, A.; Saito, A.M.; Asada, R.; Takeuchi, K.; Terauchi, T.; Tateishi, U.; Horibe, K.; Nagai, H. Alectinib for relapsed or refractory anaplastic lymphoma kinase-positive anaplastic large cell lymphoma: An open-label phase II trial. *Cancer Sci*. **2020**;111(12):4540-4547. doi: 10.1111/cas.14671.
14. Mossé, Y.P.; Voss, S.D.; Lim, M.S.; Rolland, D.; Minard, C.G.; Fox, E.; Adamson, P.; Wilner, K.; Blaney, S.M.; Weigel, B.J. Targeting ALK With Crizotinib in Pediatric Anaplastic Large Cell Lymphoma and Inflammatory Myofibroblastic Tumor: A Children's Oncology Group Study. *J Clin Oncol*. **2017**;35(28):3215-3221. doi: 10.1200/JCO.2017.73.4830.
15. Guerreiro Stucklin, A.S.; Ryall, S.; Fukuoka, K.; et al. Alterations in ALK/ROS1/NTRK/MET drive a group of infantile hemispheric gliomas. *Nat Commun* **2019**;10,4343. doi: 10.1038/s41467-019-12187-5.
16. Risum, S.; Knigge, U.; Langer, S.W.; Hitherto unseen survival in an ALK-positive-patient with advanced stage adult ganglioneuroblastoma treated with personalized medicine. *Clin Case Rep*. **2017**;5(12):2085-2087. doi: 10.1002/ccr3.1262.

17. Sica, A.; Casale, B.; Dato, M.T.D.; Calogero, A.; Spada, A.; Sagnelli, C.; Santagata, M.; Buonavolontà, P.; Fiorelli, A.; Salzano, A.; Dodaro, C.A.; Martinelli, E.; Saracco, E.; Troiani, T.; Tammaro, D.; Ciardiello, F.; Papa, A. Cancer- and Non-cancer Related Chronic Pain: From the Physiopathological Basics to Management. *Open Med (Wars)*. 2019;14:761-766. doi: 10.1515/med-2019-0088.
18. Sica, A.; Casale, B.; Sagnelli, C.; Di Dato, M.T.; Buonavolontà, P.; Salzano, A.M.; et Al. All In One spinal cord stimulation in lymphoproliferative diseases. *Front Neurol*. 2020;11:550554. doi: 10.3389/fneur.2020.550554.
19. Merli, M.; Frigeni, M.; Alric, L.; Visco, C.; Besson, C.; Mannelli, L.; Di Rocco, A.; Ferrari, A.; Farina, L.; Pirisi, M.; Piazza, F.; Loustaud-Ratti, V.; Arcari, A.; Marino, D.; Sica, A.; Goldaniga, M.; Rusconi, C.; Gentile, M.; Cencini, E.; Benanti, F.; Rumi, M.G.; Ferretti, V.V.; Grossi, P.; Gotti, M.; Sciara, R.; Tisi, MC.; Cano, I.; Zuccaro, V.; Passamonti, F.; Arcaini, L. Direct-Acting Antivirals in Hepatitis C Virus-Associated Diffuse Large B-cell Lymphomas. *Oncologist*. 2019;24(8):e720-e729. doi: 10.1634/theoncologist.2018-0331.
20. Guastafierro, S.; Sica, A.; Parascandola, R.R.; Ferrara, M.G.; Di Martino, A.; Pezone, L.; Falcone, U. Clinical significance of serum triple monoclonal components: a report of 6 cases and a review of the literature. *Leuk Res*. 2014;38(2):166-9. doi: 10.1016/j.leukres.2013.10.020.
21. Guastafierro, S.; Sica, A.; Parascandola, R.R.; Ferrara, MG; Di Martino, A.; Pezone, L.; Falcone, U.; Clinical significance of serum triple monoclonal components: a report of 6 cases and a review of the literature. *Leuk Res*. 2014;38(2):166-9. doi: 10.1016/j.leukres.2013.10.020.
22. Merli, M.; Defrancesco, I.; Visco, C.; Besson, C.; Di Rocco, A.; Arcari, A.; Sica, A.; Cencini, E.; Tisi, MC.; Frigeni, M.; Grossi, P.; Bianchi, B.; Mora, B.; Bertù, L.; Bruno, R.; Passamonti, F.; Arcaini, L. Direct-acting antivirals in relapsed or refractory hepatitis C virus-associated diffuse large B-cell lymphoma. *Leuk Lymphoma*. 2020:1-7. doi: 10.1080/10428194.2020.1755859.
23. Tonziello, G.; Pisaturo, M.; Sica, A.; Ferrara, M.G.; Sagnelli, C.; Pasquale, G.; Sagnelli, E.; Guastafierro, S.; Coppola, N. Transient reactivation of occult hepatitis B virus infection despite lamivudine prophylaxis in a patient treated for non-Hodgkin lymphoma. *Infection*. 2013;41(1):225-9. doi: 10.1007/s15010-012-0305-y.
24. Guastafierro, S.; Ferrara, M.G.; Sica, A.; Parascandola, R.R.; Santangelo, S.; Falcone, U. Serum double monoclonal components and hematological malignancies: only a casual association? Review of 34 cases. *Leuk Res*. 2012;36(10):1274-7. doi: 10.1016/j.leukres.2012.05.008.
25. Pisaturo, M.; Guastafierro, S.; Filippini, P.; Tonziello, G.; Sica, A.; Di Martino, F.; Sagnelli, C.; Ferrara, M.G.; Martini, S.; Cozzolino, D.; Sagnelli, E.; Coppola, N. Absence of occult HCV infection in patients experiencing an immunodepression condition. *Infez Med*. 2013;21(4):296-301.
26. Coppola, N.; Pisaturo, M.; Guastafierro, S.; Tonziello, G.; Sica, A.; Sagnelli, C.; Ferrara, M.G.; Sagnelli, E. Absence of occult hepatitis C virus infection in patients under immunosuppressive therapy for oncohematological diseases. *Hepatology*. 2011;54(4):1487-9. doi: 10.1002/hep.24436.
27. Coppola, N.; Pisaturo, M.; Guastafierro, S.; Tonziello, G.; Sica, A.; Iodice, V.; Sagnelli, C.; Ferrara, M.G.; Sagnelli, E. Increased hepatitis C viral load and reactivation of liver disease in HCV RNA-positive patients with onco-hematological disease undergoing chemotherapy. *Dig Liver Dis*. 2012;44(1):49-54. doi: 10.1016/j.dld.2011.07.016.
28. Bagaglio, S.; Uberti-Foppa, C.; Sagnelli C.; Lai, A.; Hasson, H.; Salpietro, S.; Messina, E.; Morsica, G.; Zaffina, C.; Sica, A.; Lazzarin, A.; Angeletti, S.; Sagnelli, E.; Foley, B.; Ciccozzi, M. HIV-1 recombinant forms in immigrants regularly residing in Milan; northern Italy. *Infection*. 2020. doi: 10.1007/s15010-020-01434-3.
29. Vivarelli, S.; Falzone, L.; Grillo, C.M.; Scandurra, G.; Torino, F.; Libra, M. Cancer Management during COVID-19 Pandemic: Is Immune Checkpoint Inhibitors-Based Immunotherapy Harmful or Beneficial? *Cancers (Basel)*. 2020;12(8):2237. doi: 10.3390/cancers12082237.
30. Al-Quteimat, O.M.; Amer, A.M. The Impact of the COVID-19 Pandemic on Cancer Patients. *Am J Clin Oncol*. 2020;43(6):452-455. doi: 10.1097/COC.0000000000000712.
31. Sica, A.; Vitiello, P.; Papa, A.; Calogero, A.; Sagnelli, C.; Casale, D.; Mottola, M.; Svanera, G.; Dodaro C.A.; Martinelli, E.; Troiani, T.; Ciardiello, F.; Casale, B. Use of Rituximab in NHL Malt Type Pregnant in I^o Trimester for Two Times. *Open Med (Wars)*. 2019;14:757-760. doi: 10.1515/med-2019-0087.
32. Falcone, U.; Parascandola, R.R.; Ferrara, M.G.; Sica, A.; Di Martino, A.; Guastafierro, S. Bilateral lacrimal gland involvement with mantle cell lymphoma. *Am J Med Sci*. 2015;349(3):268. doi: 10.1097/MAJ.0b013e3182a58ae0.

33. Reginelli, A.; Belfiore, M.P.; Russo, A.; Turriziani, F.; Moscarella, E.; Troiani, T.; Brancaccio, G.; Ronchi, A.; Giunta, E.; Sica, A.; Iovino, F.; Ciardiello, F.; Franco, R.; Argenziano, G.; Grassi, R.; Cappabianca, S. A Preliminary Study for Quantitative Assessment with HFUS (High-Frequency Ultrasound) of Nodular Skin Melanoma Breslow Thickness in Adults Before Surgery: Interdisciplinary Team Experience. *Curr Radiopharm.* **2020**;13(1):48-55. doi: 10.2174/1874471012666191007121626.
34. Sica, A.; Vitiello, P.; Caccavale, S.; Sagnelli, C.; Calogero, A.; Doraro, C.A.; Pastore, F.; Ciardiello, F.; Argenziano, G.; Reginelli, A.; Cappabianca, S.; Franco, R.; Ronchi, A. Primary Cutaneous DLBCL Non-GCB Type: Challenges of a Rare Case. *Open Med (Wars).* **2020**;15:119-125. doi: 10.1515/med-2020-0018.
35. Sica, A.; Sagnelli, C.; Papa, A.; Ciccozzi, M.; Sagnelli, E.; Calogero, A.; Martinelli, E.; Casale, B. An Anecdotal Case Report of Chronic Lymphatic Leukemia with del(11q) Treated with Ibrutinib: Artificial Nourishment and Physical Activity Program. *Int J Environ Res Public Health.* **2020**;17(6):0. doi: 10.3390/ijerph17061929.
36. Vitiello, P.; Sica, A.; Ronchi, A.; Caccavale, S.; Franco, R.; Argenziano, G. Primary cutaneous B cell lymphomas: an update. *Front. Oncol.* **2020**;10:651. doi: 10.3389/fonc.2020.00651.
37. Ronchi, A.; Zito Marino, F.; Vitiello, P.; Caccavale, S.; Argenziano, G.; Crisci, S.; Franco, R.; Sica, A. A case of primary cutaneous B-cell lymphoma with immature features in an old man. Diffuse Large B-cell Lymphoma with immature features or B-cell Lymphoblastic Lymphoma? *J Cutan Pathol.* **2020**. doi: 10.1111/cup.13795.
38. Visco, C.; Di Rocco, A.; Evangelista, A.; Quaglia, F.M.; Tisi, M.C.; Morello, L.; Zilioli, V.R.; Rusconi, C.; Hohaus, S.; Sciarra, R.; Re, A.; Tecchio, C.; Chiappella, A.; Marin-Niebla, A.; McCulloch, R.; Gini, G.; Perrone, T.; Nassi, L.; Pennese, E.; Stefani, P.M.; Cox, M.C.; Bozzoli, V.; Fabbri, A.; Polli, V.; Ferrero, S.; Celis, M.I.A.; Sica, A.; Petrucci, L.; Arcaini, L.; Rule, S.; Krampera, M.; Vitolo, U.; Balzarotti, M. Outcomes in first relapsed-refractory younger patients with mantle cell lymphoma: results from the MANTLE-FIRST study. *Leukemia.* **2020**. doi: 10.1038/s41375-020-01013-3.
39. Reginelli, A.; Urraro, F.; Sangiovanni, A.; Russo, G.M.; Russo, C.; Grassi, R.; Agostini, A.; Belfiore, M.P.; Cellina, M.; Floridi, C.; Giovagnoni, A.; Sica, A.; Cappabianca, S. Extranodal Lymphomas: a pictorial review for CT and MRI classification. *Acta Biomed.* **2020**;91(8-S):34-42. doi: 10.23750/abm.v91i8-S.9971.
40. Dumont, D.; Dô, P.; Lerouge, D.; Planchard, G.; Riffet, M.; Dubos-Arvis, C.; Danhier, S.; Gervais, R. Off-Label Use of Crizotinib as a Neoadjuvant Treatment for a Young Patient When Conventional Chemotherapy Gave No Benefits in Stage IIIA Non-Small Cell Lung Cancer. *Am J Case Rep.* **2017**;18:890-893. doi: 10.12659/ajcr.903528.
41. Falzone, L.; et al. Evolution of Cancer Pharmacological Treatments at the Turn of the Third Millennium. *Front Pharmacol.* **2018**. 10.3389/fphar.2018.01300.
42. Guo, L.; Zhang, H.; Shao, W.; Chen, B. Crizotinib as a personalized alternative for targeted anaplastic lymphoma kinase rearrangement in previously treated patients with non-small-cell lung cancer. *Drug Des Devel Ther.* **2015**;9:5491-7. doi: 10.2147/DDDT.S91988. eCollection 2015.
43. Qian, H.; Gao, F.; Wang, H.; Ma, F. The efficacy and safety of crizotinib in the treatment of BMC Cancer. anaplastic lymphoma kinase-positive non-small cell lung cancer: a meta-analysis of clinical trials. **2014**;14:683. doi: 10.1186/1471-2407-14-683.
44. Viscardi, G.; Zanaletti, N.; Ferrara, M.G.; Sica, A.; Falcone, U.; Guastafierro, S.; Bracale, U.; Ribero, D.; Fasano, M.; Napolitano, S.; Vitale, P.; De Falco, V.; Giunta, E.F.; Martinelli, E.; Ciardiello, D.; Ciardiello, F.; Troiani, T. Atypical haemolytic-uraemic syndrome in patient with metastatic colorectal cancer treated with fluorouracil and oxaliplatin: a case report and a review of literature. *ESMO Open.* **2019**;4(5):e000551. doi: 10.1136/esmoopen-2019-000551.
45. Capece, M.; Creta, M.; Calogero, A.; La Rocca, R.; Napolitano, L.; Barone, B.; Sica, A.; Fusco, F.; Santangelo, M.; Dodaro, C.; Sagnelli, C.; Carlomagno, N.; Crocetto, F.; Califano, G.; Mangiapia, F.; Longo, N. Does Physical Activity Regulate Prostate Carcinogenesis and Prostate Cancer Outcomes? A Narrative Review. *Int J Environ Res Public Health.* **2020**;17(4):1441. doi: 10.3390/ijerph17041441.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

