

## Article

# Male sexual preference for female swimming activity in the guppy (*Poecilia reticulata*)

David Bierbach <sup>1,2,3 \*</sup>, Ronja Wenzel <sup>1</sup>, Stefan Gehrig <sup>1</sup>, Serafina Wersing <sup>1</sup>, Olivia L. O'Connor <sup>2,3</sup> and Jens Krause <sup>1,2,3</sup>

<sup>1</sup> Department of Biology and Ecology of Fishes, Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Müggelseedamm 310, 12587 Berlin, Germany; RW: [ronja.wenzel@fu-berlin.de](mailto:ronja.wenzel@fu-berlin.de); SG: [stefan-gehrig@t-online.de](mailto:stefan-gehrig@t-online.de); SW: [serafina.wersing@igb-berlin.de](mailto:serafina.wersing@igb-berlin.de); JK: [j.krause@igb-berlin.de](mailto:j.krause@igb-berlin.de)

<sup>2</sup> Faculty of Life Sciences, Humboldt University, Invalidenstrasse 42, 10115 Berlin, Germany; DB: [bierbach@hu-berlin.de](mailto:bierbach@hu-berlin.de); OLO: [oconnorolivia30@gmail.com](mailto:oconnorolivia30@gmail.com)

<sup>3</sup> Science of Intelligence (SCIOI), Technische Universität Berlin, Marchstraße 23, 10587 Berlin, Germany

\* Correspondence: [bierbach@hu-berlin.de](mailto:bierbach@hu-berlin.de)

**Simple Summary:** In a first experiment, we found that male and female guppies differ consistently in their swimming activity both when alone and when together in pairs and that males regardless of their own activity show more sexual behaviours to more active females. In a second experiment, we gave males the choice between a slow and a fast moving virtual female and found that males prefer to associate with the fast moving female. Both experiments together provide evidence that guppy males prefer faster moving females as mating partners over slower moving ones. Higher activity patterns may indicate health, fitness and receptivity and a mating preference for more active females can help males to increase their own reproductive success.

**Abstract:** Mate choice that is based on behavioural traits is a common feature in the animal kingdom. Using the Trinidadian guppy, a species with mutual mate choice, we investigated whether males use female swimming activity – a behavioural trait known to differ consistently among individuals in many species – as a trait relevant for their mate choice. In a first experiment, we assessed male and female activity in an open field test alone (two repeated measures) and afterwards in heterosexual pairs (two repeated measures). In these pairs, we simultaneously assessed males' mating efforts by counting number of sexual behaviours (courtship displays and copulation). Male and female guppies showed consistent individual differences in their swimming activity when tested both alone and in a pair, and these differences were maintained across both test situations. When controlling for male swimming behaviour and both male and female body size, males performed more courtship displays towards females with higher swimming activity. In a second experiment, we tested for a directional male preference for swimming activity by presenting males video animations of low and high active females in a dichotomous choice test. In congruence with experiment 1, we found males to spend significantly more time in association with the high active female stimulus. Both experiments thus point towards a directional male preference for higher activity levels in females. We discuss the adaptive significance of this preference as activity patterns might indicate individual female quality, health or reproductive state while, mechanistically, females that are more active might be more detectable to males as well.

**Keywords:** Animal personality, swimming activity, male mate choice, mating preferences, *Poecilia reticulata*

## 1. Introduction

Males just like females can exercise mate choice [1, 2]. Male mate choice is not only evident in species with reversed sex roles and male parental care [3, 4] but also in species where male investment is low but females differ in quality [5-7]. Female quality in such species is often approximated by fecundity, and, as a correlate, female body size [2, 8]. In addition, males also choose their mating partners on the base of coloration [9], actual mating status [10] or quality of parental care [11]. However, some studies on female mate choice report on sexual preferences for behaviours not related directly to reproduction (see [12, 13]) like feeding rates [14], exploration tendencies [15, 16] or predator inspection behaviour [17].

Similarly, males may choose females based on non-sexually motivated behaviours if these behaviours are consistent indicators of quality (see [13, 18, 19]). Alternatively, males may benefit when mating with females that match their own behavioural phenotype [13, 20]. In zebra finches, females prefer males that are similar in their exploration tendency to themselves [15] and pairs with similar behavioural type have increased parental care success [21]. In guppies, females paired with males who differed from them in risk-taking behaviour had a lower parturition success than females who mated with males with a similar degree of risk-taking [22].

In the current study, we ask whether males use female activity – a behavioural trait known to differ consistently among individuals in many species [23-25] – as a criterion in mate choice. Activity patterns can indicate female quality as it often positively correlates with metabolism [26-28], growth rate [29, 30], general health [31], as well as survival [18, 19, 32]. In addition, activity patterns may even indicate reproductive status, especially in livebearers where females carry developing embryos and eggs and thus may lose movement agility when pregnant or gravid [33, 34]. Mechanistically, more active females might be more conspicuous to males as found for prey that is more visible to predators when moving more (see [35, 36]).

We decided to use the live-bearing Trinidadian guppy (*Poecilia reticulata*) as model organism as both males and females are known to exercise mate choice [37] and individuals often differ consistently in their activity levels (with males being on average more active than females [38, 39]). In this species, higher activity levels in females correlate positively with risk-taking behaviours [40], and negatively with gestation period [34] and body size [40].

We hypothesized that males should show a general preference for more active females ('directional preference hypothesis') as found for example in lizards [16]. As an alternative hypothesis we explored whether males prefer females with an activity types that is similar to their own ('phenotypic matching hypothesis') as found in zebra finches [21]. To provide evidence for either hypothesis, we assessed in a first experiment male and female activity in an open field test alone (non-sexual context treatment, two repeated measures) and afterwards in heterosexual pairs (sexual context treatment, two repeated measures). In these pairs, we simultaneously assessed males' mating efforts by counting the number of sexual behaviours (courtship displays and copulation). Sexual behaviours exhibited by male guppies are a direct measure of sexual preference [5, 10, 41]. For the directional preference hypothesis, we predicted that males should show largest number of sexual behaviours towards the most active females while for the phenotypic matching hypothesis, males should direct most sexual behaviours to females with activity levels similar to their own.

As the first experiment will provide purely correlational evidence, we performed a second experiment testing specifically the directional preference hypothesis using a dichotomous choice design. Males could choose to associate with computer-animated females that were either fast or slow swimming. Here the prediction was that males should spend more time close to the faster animated females.

2. Materials and Methods

Study organism

Test fish were laboratory-reared descendants of wild-caught Trinidadian guppies that originate from the Arima river in Trinidad (low predation site; see [37]). They came from large, randomly outbred, mixed-sex stocks reared under constant physical conditions (25°C, 12 hours of light). The fish were fed three times per day with commercially available flake food (TetraMin® Tetra GmbH); water was exchanged once a week. Prior to the experiments, males and females were separated by sex for one week to increase sexual motivation [5].

1st Experiment

All tests were conducted in a standard tank (35 cm x 20 cm) that was filled with water from the holding tanks to a level of 5 cm. The walls were covered to avoid disturbances from the outside and observation was done via a webcam fixed at 50 cm above the tank. Male activity was measured by introducing a male into a Plexiglas cylinder inside the test tank. After a habituation of 5 min, the cylinder was lifted and the activity of the test fish was videotaped for another five minutes. This testing was repeated for all individuals after 48 hours (figure 1a). After the first test, males were housed separately in small tanks of a circulating water system and thus had experienced both visual as well as olfactory cues of other conspecifics.




Figure 1: Schematic overview of the experimental timeline for experiment 1 (a) and 2 (b).

Female activity was measured as described for males but each 5 min activity test was followed by another phase in which we presented females with a male (Figure 1). To do so, females were placed back into their cylinder. Then, a randomly selected male from our pool of males was introduced into the test tank. After five minutes the cylinder was removed and both fish were videotaped for five minutes. As for the male activity test, this two-phase test was repeated after 48 hours with another male.

From the videos, we extracted the total distance moved by each fish during the five minutes tested alone and in dyads as a proxy for individual activity [42]. In the dyads, we further counted the number of copulation attempts as well as courtship displays (see [5] for a description) by the male directed to the female as a proxy for male sexual effort. To extract the distance, videos were analysed using the tracking software EthoVision 10 XT (Noldus Information Technology) with 15 frames per second and the smoothing profile 2mm MDM based on five samples before and after every sample point. The standard length (head to base of tail) of each fish was measured to the nearest millimetre via photos using ImageJ software (ImageJ 1.46a). In total, 82 females (standard length:  $18.78 \pm 0.37$  mm, mean  $\pm$  SEM) and 52 males (SL:  $13.81 \pm 0.31$  mm) have been tested and their size range is comparable to natural observations [43].

We first checked whether males and females showed consistent individual differences in their activity levels when tested alone (non-sexual context treatment). To do so, we calculated the behavioural repeatability with linear mixed models (LMMs) following the protocol by [44]. We included 'trial' as a fixed factor to account for average differences in activity between first and second personality assessment (Figure 1) and individual 'body size' (SL) as a covariate to evaluate whether activity correlates with body size. Fish ID was specified as a random factor to allow the decomposition of the variance into a within and an among-individual component and subsequent calculation of behavioural repeatability. This 'conditional repeatability' is a measure for consistent differences among individuals and its p-values derive from LRT tests against  $\chi^2$  distributions with 1 degree of freedom [45].

Then, we asked whether females differ also consistently in their activity when being together with a male (sexual context). Thus, we calculated the repeatability of female activity in dyads using a similar LMM as described above. As swimming performance is known to be influenced by social partners [46], we included both activity of the male partner and its body size as covariates in the model. To compare average activity levels across treatments, we run a LMM with 'trial' and 'treatment' (indicating sexual or non-sexual context) as well as their interaction as fixed factors. Female ID was used as random factor.

To see whether females differ consistently across treatments (alone and in a pair), we calculated the among-individual correlation of female activity using a bivariate LMM. To do so, we first standardized female activity within both treatments and measures to a mean of 0 and a SD of 1 (z-transformation). Thus, no average differences between the treatments and measures may affect the among-individual correlation. In a second step, activity levels of females alone and in the dyads were used as dependent variable with Female ID as a random factor. We allowed within-individual as well as among-individual variances to differ between treatments as well as to correlate within treatments (see [40, 47] for similar approaches). As for repeatability, LRT tests were employed to check whether the inclusion of within-and among-individual correlations significantly improved the model.

Our main question was whether males directed more sexual effort towards more actively swimming females (directional preference) or whether male sexual effort depended on both male and female activity levels (phenotypic matching). We thus used 'received male sexual efforts' (sum of copulation attempts + courtship displays per pair) as dependent variable in a generalized linear mixed model (GLMM) with Poisson error-distribution and Log-link function as for count data. As fixed factors, we included 'trial', 'female activity' (within the dyad), 'male activity' (within the dyad; residuals from linear regression with female activity as both variables were highly correlated,

see Results) as well as male and female absolute body size ('SL female'; 'SL male'). We further included also the interaction between 'female activity' (within the dyad) and 'male activity' (within the dyad). Female ID was used as a random factor. A significant effect of the term 'female activity' would indicate a directional male preference, while a significant effect of 'male activity' would indicate that male activity determined their sexual efforts. If the terms 'female activity' and 'male activity' together were significant and positive, more active males would have directed more sexual efforts to more active females. A significant interaction term of both effects would indicate a mismatching sexual preference as less active males would direct more sexual efforts to high active females. However, we removed the interaction term from the final model as there was little evidence supporting that it was different from zero in either direction (see Results).

## 2nd Experiment

Using a dichotomous choice design (see [48]), we evaluated male preferences (total length (TL  $\pm$  SEM):  $20.4 \pm 0.5$  mm;  $n = 17$ ) for two virtual stimulus female types (sized on screen set to 30.0 mm TL each) by showing each focal male video animations of the same virtual female either swimming at 0.7 cm/s or with 2.7 cm/s and measured association preferences for 10 min (with switched side assignment after 5 min). The used swimming speeds were the lower and upper 10% percentiles of females from experiment 1. This technique is well established to evaluate mating preferences in fish per se [49-53] and has been used successfully by the authors in previous studies ([48, 54].

To produce the video animations, digital images of 5 females were taken with grey background colour using a Canon 400D digital camera (see Supplemental Information S1). From each resulting picture, the image of the female was extracted from the background using the "magic wand" selection tool in Adobe® Photoshop CS5. Each picture was then embedded into a Microsoft PowerPoint presentation and a straight movement of the picture from left to right and right to left was generated in front of a uniformly light grey background. For each picture of a female, we generated a slow and a fast moving version. Simultaneous playback was performed using two identical computer monitors (Dell P2419H) as well as two laptops with identical MS Office versions (Office 365+). The monitors' refresh rate was 60 Hz and a sample animation (.mp4) is available as a supplement (movie SI).

For the preference tests, monitors were placed on either side of a test tank (40 cm length x 25 cm width x 25 cm depth) that was visually divided into three sections: two preference zones (5 cm length) adjacent to the monitors and a central neutral zone (30 cm length). Both long sides of the tank were covered by white Perspex, so that the focal male did not see the experimenter in the room. Water level was kept at 15 cm, which was also the height of the monitors. Water temperature was maintained at 25°C, and illumination was provided through two LED stripes on the ceiling of the experimental room. The focal male was observed via a Basler acA1920-155um USB3 camera in Full HD with 30 fps and the resulting videos were analysed using Ethovision software.

To initiate a trial, we introduced a male into the test tank. After a habituation period of 5 minutes (showing only a grey video screen), we started a 5-minute observation period during which we measured association times, i.e., times spent in each preference zone, near the monitor showing the animation of the fast or slow moving female. To account for potential side-biases, we interchanged the stimulus animations and repeated measurement of male preferences after another 5 minutes of habituation. We decided a priori to assume a side preference when a male spent 80% of its association time during both test units in one of the two preference zones (i.e., at only one side of the test tank). Based on this criterion, 6 out of 23 trials had to be discarded from the final analysis resulting in a final sample size of 17 males. We summed the times spent near either female animation during the two test units and further extracted male swimming velocity and body size from the videos. Association preference are well established to translate into reproductive outcome in poeciliid fishes [55]



We first compared the time males spent near each stimulus using paired- samples t-tests. In order to include male body size, male swimming activity as well as stimulus females' picture ID in our analysis, we calculated an SOP (strength of preference score) as the time spent near the fast swimming animation minus the time spent near the slow swimming animation divided by the total time spent choosing. Thus, an SOP score of +1 reflects a maximal preference for the fast stimulus animation and - 1 maximal preference for the slow stimulus animation. SOP scores were used as dependent variable in an LMM with male body size, male swimming activity as covariates well as stimulus females' picture ID as random factor. An additional LMM on SOP score was fit as a robustness check for the paired t-test on raw time spent near stimulus mentioned above. Since this LMM contained no fixed effects, the intercept was an estimate for the marginal strength of preference for females, adjusted for stimulus females' picture ID.

### 3. Results

#### 3.1. 1<sup>st</sup> Experiment

We found significant among-individual variation in male and female activity when fish were observed alone ( $Repeatability_{males}=0.76$ ,  $p<0.001$ ;  $Repeatability_{females}=0.63$ ,  $p<0.001$ ). On average, activity significantly decreased with repeated testing in both sexes (males:  $F_{1,52}=78.06$ ,  $p<0.001$ ; females:  $F_{1,81}=67.60$ ,  $p<0.001$ ). In females, activity negatively and significantly covaried with body size ( $F_{1,80}=4.81$ ,  $p=0.031$ ) meaning that larger females were less active (*posthoc* Pearson's correlation:  $r_p=-0.201$ ;  $p=0.010$ ).

With males females showed consistent among-individual variation ( $Repeatability=0.45$ ,  $p<0.001$ ) and average female activity covaried positively and significantly with the male partner's activity ( $F_{1,149}=230.85$ ,  $p<0.001$ ; *posthoc* Pearson's correlation:  $r_p=0.80$ ;  $p<0.001$ ). There was some evidence that females, on average, decreased their activity with repeated testing ( $F_{1,74}=3.7$ ,  $p=0.057$ ). However, there was no significant effect of female or male body size on female activity levels (female body size:  $F_{1,81}=0.39$ ,  $p=0.845$ ; male body size:  $F_{1,149}=2.51$ ,  $p=0.115$ ).

When comparing average female activity among treatments, we found females to be less active when tested with a male than tested alone (Figure 2a; treatment:  $F_{1,243}=276.20$ ,  $p<0.001$ ). Similar to above model results, females reduced their activity with repeated testing also when pooling data from both treatments (Figure 2a; trial:  $F_{1,243}=61.80$ ,  $p<0.001$ ). However, the decrease in activity from non-sexual to sexual context treatment was more pronounced in the first personality assessment (Figure 2a; 'treatment  $\times$  trial':  $F_{1,243}=11.92$ ,  $p=0.001$ ). Despite these differences, we found strong support for among-individual correlation in activity across treatments ( $r_{among}=0.73$ ,  $p<0.001$ , Figure 2b). There was also less pronounced within-individual correlation ( $r_{within}=0.25$ ,  $p=0.021$ ). Thus, females that were on average more active alone were also more active when tested together with a male (Figure 2b, among-individual correlation). Furthermore, females that were more variable between repeated trials in their activity alone were also more variable in activity when with a male (significant within-individual correlation).




Figure 2: (a) Distance moved by females. Shown are mean distances for both trials (1<sup>st</sup> and 2<sup>nd</sup>) as well as for the treatments (female alone or with a male partner). (b) Consistency in female swimming activity across treatments. Shown is a scatter plot with individual-level error bars and a linear line of best fit (with 95% CI, dotted lines) for female distances moved in an open field arena both alone and with a male partner. Error bars represent SEM (a) and SD (b).

Our GLMM with received sexual efforts as dependent variable found a strong positive relationship between male sexual efforts and female activity ('female activity (dyad)':  $F_{1,157}=9.51$ ,  $p=0.002$ ). This suggests that more actively swimming females received more sexual behaviours than less active ones (Figure 3). None of the other predictors had a significant effect ('male activity' (residualised):  $F_{1,157}=0.19$ ,  $p=0.661$ ; 'female SL':  $F_{1,157}=0.31$ ,  $P=0.578$ ; 'male SL':  $F_{1,157}=1.57$ ,  $p=0.21$ ; 'trial':  $F_{1,157}=2.60$ ,  $p=0.109$ ; interaction 'female activity by male activity' (not in the final model):  $F_{1,156}=1.20$ ,  $p=0.275$ ). Thus, there was no evidence that male sexual effort was related to body size of the involved fish, males' swimming activity or to the interaction of male with female activity.




Figure 3: Male sexual activity (sum of courtship displays and copulation attempts) and female swimming activity (distance moved in an open field tank with a male partner). The regression line refers to the final GLMM model predictions (with 95% CI). Error bars represent SD.

### 3.1. 2<sup>nd</sup> Experiment

We found males to spend significantly more time near fast moving female animations as compared to slow moving ones ( $t_{16}=2.209$ ;  $p=0.0421$ , Figure 4). The strength of these preference (SOP score) appeared to be independent of male body size (LMM:  $F_{1,14}=0.08$ ;  $p=0.79$ ) as well as swimming speed during the mate choice trials (LMM:  $F_{1,14}=0.94$ ;  $p=0.35$ ). Analysing SOP scores in an LMM without any fixed effects (but a random effect for stimulus females' picture ID) confirmed the overall preference for fast moving females. The model intercept was positive and significantly different from zero (estimate: 0.226, SE: 0.100,  $t_{16} = 2.263$ ,  $p=0.038$ ).






Figure 4: Results of dichotomous mate choice tests to assess directional mating preferences of male guppies for female swimming activity. Shown are the mean ( $\pm$  SE) times males spent near fast (2.7 cm/s) and slow (0.7 cm/s) moving female animations. An asterisk indicates significant difference ( $p < 0.05$ ) in paired-samples  $t$ -test.

#### 4. Discussion

We found male and female guppies to show consistent individual differences in their swimming activity when tested alone as well as in pairs. In support for the directional preference hypothesis, more actively swimming females received higher numbers of sexual behaviours by males, regardless of female or male body size or male swimming activity. Our 2<sup>nd</sup> experiment confirmed the correlational evidence for a male preference for female swimming activity from experiment 1 by showing that males spent more time near fast moving female animations compared to slow moving ones in dichotomous mate choice tests.

Levels of individual swimming activity in an open field arena were found to be consistently different among individuals (i.e. there was considerable and significant repeatability) when males and females were tested both alone and together (= significant among-individual correlation). These results are consistent with previous studies showing significant repeatability in activity levels in the guppy [38, 56-58] even when interacting with (artificial) conspecifics (see [40, 59, 60]). Those individual differences in swimming patterns can be maintained even in larger groups as shown for mosquitofish (*Gambusia holbrooki*, [46]) and sticklebacks [61]. Thus, swimming activity fulfils a prerequisite of a trait that could be used as a cue in mate choice [62, 63]. For male mate choice, we assume that activity patterns can indicate female quality as it often positively correlates with metabolism, growth rate, general health, as well as survival (see introduction section). The fact that more active individuals are also more visible to predators [35] may render high activity levels a costly trait that could fulfil the requirements of the “handicap principle” (after [64]). Following the

same logic and besides the aforementioned indication of quality, higher activity levels may increase an individual's visibility towards the opposite sex or even stimulate more exaggerated sexual responses ('arousal hypothesis'). It is further known that gestation statuses in live-bearing animals where developing eggs or embryos are carried by females hamper the female's agility and mobility [33, 34]. Female guppies are most receptive a few days post-partum [65] and males typically need to assess females' reproductive status through nipping at the female's gonopore [66] or by observing and imitating the mate choice of other males (mate choice copying, [67]). Both is assumed to be costly for the males in terms of opportunity costs [68]). It is thus possible that male guppies use female activity patterns to select females that gave birth recently and are thus most receptive.

As predicted by the 'directional preference hypothesis' males in general increased their sexual efforts when paired with females that were more actively swimming (experiment 1). Contrary to the 'phenotypic matching hypothesis', this preference was independent of male activity (and its interaction with female activity) as well as body size of males and females. Although guppy males can perform courtship prior to copulation attempts, their mating systems is assumed to be coercive [37, 69, 70] and forced copulations and sexual harassment are common features both in the lab and in the wild [71-74]. Thus, a phenotypic matching on activity with potential reproductive benefits like, as an example, synchrony during mating or egg/sperm release, may not play a significant role in the guppy, compared to other species [21, 75]. This view is further supported by our second experiment where males spent significantly more time in association with the more active female animation regardless of own size or swimming activity. Future work on post-copulatory effects of mating partners' swimming activity and overall reproductive success is needed to investigate the possibility of such a pattern in more detail. Nevertheless, if female swimming activity indicated overall quality or reproductive status (as our experiments suggest), a mating preference that leads to phenotypic matching for activity in pairs might not be beneficial in this species.

Female activity was lower in the second trial than in the first one and lower in the sexual than in the non-sexual context treatment.. Reducing swimming activity in the presence of males could be an adaptive strategy to avoid male harassment if females anticipate increased sexual efforts by males as a response towards high swimming activity. A study by [71] found, however, that females that experienced high male sexual harassment (many males in holding tank; sex ratio 1:1) were more active than females that experienced only low levels of harassment (few males in holding tank; sex ratio 1:5.7). Also, females from the high harassment treatment swam more efficiently (less oxygen needed at a given speed), spending less time performing costly pectoral fin-assisted swimming compared to low harassment females. In contrast to our study, these authors recorded activity in the holding tanks and their study lacked a treatment with no males present. We thus argue that the activity reduction in response to a male stimulus which we found could represent a short-term behavioural adaptation to male harassment. On the long run, increased male harassment could result in more efficient swimming performance by females due to experience [71] and exercise [76], ultimately driven by the opportunity costs associated with male harassment. The general reduction in swimming activity from trial one to trial two in our study is most likely an effect of habituation to the experimental setup and the test environment (see [77]).

Why was there no evidence for a preference for large female body size in our study? Similar experiments in the mosquitofish (*G. holbrooki*) revealed an increased sexual activity towards larger females [41]. In the Trinidadian guppy, [10] found males of one guppy population to show such preference (lower Aripo, high predation site) as well, but not the males of another one (upper Aripo, low predation site). Interestingly, our study population derived from a low predation habitat too, similar to the population that did not show a preference for large females in the study by [10]. Thus, in accordance with their interpretations, guppy males from low-predation sites might have a reduced preference for large female body size. We therefore encourage future

experimentation on population differences in male preferences for female activity and body size (see [78]) for a review on variation in female preferences among populations).

## 5. Conclusions

In conclusion, our study provides first evidence for a male sexual preference for high female activity levels in the guppy. Activity differs consistently among females and may indicate both fitness and reproductive status in live-bearing animals. Increased activity levels further enhance visibility or arouse the mating partner. All this can be predicted to render a directional preference for higher activity levels an adaptive strategy for males in guppies but also other species. Future work on male mating preferences for activity levels in other species as well as the post-copulatory effects of such preferences on reproductive fitness may well be recommended, in particular when assuming activity differences to play a ubiquitous role in the animal kingdom.

**Supplementary Materials:** S1: Pictures of female guppies (*Poecilia reticulata*) used to generate animations in experiment 2. Video S1: Example of an animated fast moving female guppy (*Poecilia reticulata*). Data experiment 1 and 2.

**Author Contributions:** DB and JK conceived the study and designed the experiments. DB, RW and SG performed experiment 1. DB, SW and OLO performed experiment 2. DB, RW and SG analysed the data. DB wrote the manuscript with input and approval from all authors.

**Funding:** The Leibniz Competition ("B-Types project"; SAW-2013-IGB-2) as well as the Excellence Cluster "Science of Intelligence" (Germany's Excellence Strategy, EXC 2002/1, project 390523135) financed this study.

**Institutional Review Board Statement:** The reported experiments comply with the current German law approved by LaGeSo Berlin (GO124/14 to D. Bierbach).

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** All data are available as supplemental material.

**Acknowledgments:** We thank Marcus Ebert, David Lewis and Christopher Schutz for help with animal care. We further thank Julianne Lukas for providing us with pictures of guppies for animations in experiment 2.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. Andersson, M., *Sexual selection*. 1994, Princeton: Princeton University Press
2. Edward, D.A. and T. Chapman, *The evolution and significance of male mate choice*. Trends in Ecology & Evolution, 2011. **26**(12): p. 647-654.
3. Berglund, A., *Many Mates Make Male Pipefish Choosy*. Behaviour, 1995. **132**(3): p. 213-218.
4. Scherer, U. and W. Schuett, *No male mate choice for female boldness in a bi-parental West African cichlid, the rainbow krib (*Pelvicachromis pulcher*)*. PeerJ, 2018. **6**: p. e5373.
5. Bierbach, D., et al., *Casanovas are liars: behavioral syndromes, sperm competition risk, and the evolution of deceptive male mating behavior in live-bearing fishes [version 3; peer review: 3 approved]*. F1000Research, 2013. **2**(75).
6. Plath, M., et al., *Male fish deceive competitors about mating preferences*. Current Biology, 2008. **18**(15): p. 1138-41.
7. Dosen, L. and R. Montgomerie, *Mate preferences by male guppies (*Poecilia reticulata*) in relation to the risk of sperm competition*. Behavioral Ecology and Sociobiology, 2004. **55**(3): p. 266-271.
8. Bonduriansky, R., *The evolution of male mate choice in insects: a synthesis of ideas and evidence*. Biological Reviews, 2001. **76**(3): p. 305-339.
9. Amundsen, T. and E. Forsgren, *Male mate choice selects for female coloration in a fish*. Proceedings of the National Academy of Sciences, 2001. **98**(23): p. 13155-13160.
10. Ojanguren, A.F. and A.E. Magurran, *Uncoupling the links between male mating tactics and female attractiveness*. Proceedings of the Royal Society of London B: Biological Sciences, 2004. **271**(Suppl 6): p. S427-S429.

11. Pampoulie, C., K. Lindström, and C.M. St. Mary, *Have your cake and eat it too: male sand gobies show more parental care in the presence of female partners*. Behavioral Ecology, 2004. **15**(2): p. 199-204.
12. Byers, J., E. Hebets, and J. Podos, *Female mate choice based upon male motor performance*. Anim Behav, 2010. **79**(4): p. 771-778.
13. Schuett, W., T. Tregenza, and S.R.X. Dall, *Sexual selection and animal personality*. Biological Reviews, 2010. **85**(2): p. 217-246.
14. Snowberg, L.K. and C.W. Benkman, *Mate choice based on a key ecological performance trait*. Journal of Evolutionary Biology, 2009. **22**(4): p. 762-9.
15. Schuett, W., J.-G.J. Godin, and S.R.X. Dall, *Do Female Zebra Finches, Taeniopygia guttata, Choose Their Mates Based on Their 'Personality'?* Ethology, 2011. **117**(10): p. 908-917.
16. Teyssier, A., et al., *Partners' personality types and mate preferences: predation risk matters*. Behavioral Ecology, 2014.
17. Godin, J.G. and L.A. Dugatkin, *Female mating preference for bold males in the guppy, Poecilia reticulata*. Proceedings of the National Academy of Sciences, 1996. **93**(19): p. 10262-10267.
18. Smith, B.R. and D.T. Blumstein, *Fitness consequences of personality: a meta-analysis*. Behavioral Ecology, 2008. **19**(2): p. 448-455.
19. Moiron, M., K.L. Laskowski, and P.T. Niemelä, *Individual differences in behaviour explain variation in survival: a meta-analysis*. Ecology Letters, 2020. **23**(2): p. 399-408.
20. Scherer, U., J.-G.J. Godin, and W. Schuett, *Do female rainbow kribis choose males on the basis of their apparent aggression and boldness? A non-correlational mate choice study*. Behavioral Ecology and Sociobiology, 2020. **74**(3): p. 34.
21. Schuett, W., S.R.X. Dall, and N.J. Royle, *Pairs of zebra finches with similar 'personalities' make better parents*. Anim Behav, 2011. **81**(3): p. 609-618.
22. Ariyomo, T.O. and P.J. Watt, *Disassortative mating for boldness decreases reproductive success in the guppy*. Behavioral Ecology, 2013. **24**(6): p. 1320-1326.
23. Réale, D., et al., *Evolutionary and ecological approaches to the study of personality*. Philosophical Transactions of the Royal Society B: Biological Sciences, 2010. **365**(1560): p. 3937-3946.
24. Réale, D., et al., *Integrating animal temperament within ecology and evolution*. Biological Reviews, 2007. **82**(2): p. 291-318.
25. Bell, A.M., S.J. Hankison, and K.L. Laskowski, *The repeatability of behaviour: a meta-analysis*. Anim Behav, 2009. **77**.
26. Careau, V. and T. Garland, *Performance, Personality, and Energetics: Correlation, Causation, and Mechanism*. Physiological and Biochemical Zoology, 2012. **85**(6): p. 543-571.
27. Careau, V., et al., *Energy metabolism and animal personality*. Oikos, 2008. **117**(5): p. 641-653.
28. Polverino, G., et al., *Body length rather than routine metabolic rate and body condition correlates with activity and risk-taking in juvenile zebrafish Danio rerio*. Journal of Fish Biology, 2016. **89**(5): p. 2251-2267.
29. Stamps, J.A., *Growth-mortality tradeoffs and 'personality traits' in animals*. Ecology Letters, 2007. **10**(5): p. 355-363.
30. Biro, P.A. and J.A. Stamps, *Are animal personality traits linked to life-history productivity?* Trends Ecol Evol, 2008. **23**.
31. Cavigelli, S.A., *Animal personality and health*. Behaviour, 2005. **142**(9-10): p. 1223-1244.
32. Smith, B.R. and D.T. Blumstein, *Behavioral types as predictors of survival in Trinidadian guppies (Poecilia reticulata)*. Behavioral Ecology, 2010.
33. Plaut, I., *Does pregnancy affect swimming performance of female Mosquitofish, Gambusia affinis?* Functional Ecology, 2002. **16**(3): p. 290-295.
34. Ghalambor, Cameron K., David N. Reznick, and Jeffrey A. Walker, *Constraints on Adaptive Evolution: The Functional Trade-Off between Reproduction and Fast-Start Swimming Performance in the Trinidadian Guppy (Poecilia reticulata)*. The American Naturalist, 2004. **164**(1): p. 38-50.
35. Krause, J. and J.-G.J. Godin, *Predator preferences for attacking particular prey group sizes: consequences for predator hunting success and prey predation risk*. Anim Behav, 1995. **50**(2): p. 465-473.
36. Ruell, E.W., et al., *Fear, food and sexual ornamentation: plasticity of colour development in Trinidadian guppies*. Proceedings of the Royal Society B: Biological Sciences, 2013. **280**(1758).
37. Magurran, A.E., *Evolutionary ecology: The Trinidadian guppy* Oxford Series in Ecology and Evolution. 2005, Oxford: Oxford University Press.

38. Burns, J.G., *The validity of three tests of temperament in guppies (<em>Poecilia reticulata</em>)*. Journal of Comparative Psychology, 2008. **122**(4): p. 344-356.
39. Harris, S., et al., *Picking personalities apart: estimating the influence of predation, sex and body size on boldness in the guppy Poecilia reticulata*. Oikos, 2010. **119**(11): p. 1711-1718.
40. Lukas, J., et al., *Consistent behavioural syndromes across seasons in an invasive freshwater fish*. bioRxiv, 2020: p. 2020.03.03.974998.
41. Head, M.L., et al., *Male mate choice and insemination success under simultaneous versus sequential choice conditions*. Anim Behav, 2015. **103**: p. 99-105.
42. Bierbach, D., K.L. Laskowski, and M. Wolf, *Behavioural individuality in clonal fish arises despite near-identical rearing conditions*. 2017. **8**: p. 15361.
43. Reznick, D., et al., *Life-history evolution in guppies (Poecilia reticulata)*. Differential mortality as a mechanism for natural selection. Evolution, 1996. **50**: p. 1651 - 1660.
44. Dingemanse, N.J. and N.A. Dochtermann, *Quantifying individual variation in behaviour: mixed-effect modelling approaches*. Journal of Animal Ecology, 2013. **82**(1): p. 39-54.
45. Nakagawa, S. and H. Schielzeth, *Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists*. Biological Reviews, 2010. **85**(4): p. 935-956.
46. Herbert-Read, J.E., et al., *The role of individuality in collective group movement*. Proceedings of the Royal Society of London B: Biological Sciences, 2012. **280**(1752).
47. Bierbach, D., et al., *Personality affects mate choice: bolder males show stronger audience effects under high competition*. Behavioral Ecology, 2015.
48. Bierbach, D., et al., *Homosexual behaviour increases male attractiveness to females*. Biology Letters, 2013. **9**(1): p. 20121038.
49. Chouinard-Thuly, L., et al., *Technical and conceptual considerations for using animated stimuli in studies of animal behavior*. Current Zoology, 2017. **63**(1): p. 5-19.
50. Gierszewski, S., et al., *The virtual lover: variable and easily guided 3D fish animations as an innovative tool in mate-choice experiments with sailfin mollies-II. Validation*. Current Zoology, 2017. **63**(1): p. 65-74.
51. Gierszewski, S., et al., *Using the FishSim Animation Toolchain to Investigate Fish Behavior: A Case Study on Mate-Choice Copying In Sailfin Mollies*. JoVE, 2018(141): p. e58435.
52. Fisher, H.S. and G.G. Rosenthal, *Male swordtails court with an audience in mind*. Biology Letters, 2007. **3**(1): p. 5-7.
53. Schlupp, I., et al., *A novel sexually selected trait in poeciliid fishes: Female preference for mustache-like, rostral filaments in male Poecilia spheonops*. Behavioral Ecology and Sociobiology, 2010. **64**: p. 1849-1855.
54. Sommer-Trembo, C., et al., *Context-dependent female mate choice maintains variation in male sexual activity*. Royal Society Open Science, 2017. **4**(7).
55. Walling, C., et al., *Do female association preferences predict the likelihood of reproduction?* Behavioral Ecology and Sociobiology, 2010. **64**(4): p. 541-548.
56. Brown, C. and E. Irving, *Individual personality traits influence group exploration in a feral guppy population*. Behavioral Ecology, 2014. **25**(1): p. 95-101.
57. Trompf, L. and C. Brown, *Personality affects learning and trade-offs between private and social information in guppies, Poecilia reticulata*. Anim Behav, 2014. **88**(0): p. 99-106.
58. Budaev, S.V., *"Personality" in the guppy (<em>Poecilia reticulata</em>): A correlational study of exploratory behavior and social tendency*. Journal of Comparative Psychology, 1997. **111**(4): p. 399-411.
59. Jolles, J.W., et al., *Group-level patterns emerge from individual speed as revealed by an extremely social robotic fish*. Biology Letters, 2020. **16**(9): p. 20200436.
60. Bierbach, D., et al., *Using a robotic fish to investigate individual differences in social responsiveness in the guppy*. bioRxiv, 2018.
61. Jolles, J.W., et al., *Consistent Individual Differences Drive Collective Behavior and Group Functioning of Schooling Fish*. Current Biology, 2017. **27**(18): p. 2862-2868.e7.
62. Candolin, U., *The use of multiple cues in mate choice*. Biological Reviews, 2003. **78**(4): p. 575-595.
63. Partan, S. and P. Marler, *Issues in the Classification of Multimodal Communication Signals*. The American Naturalist, 2005. **166**(2): p. 231-245.
64. Zahavi, A., *Mate selection—A selection for a handicap*. Journal of Theoretical Biology, 1975. **53**(1): p. 205-214.
65. Houde, A.E., *Sex, color, and mate choice in guppies*. 1997, Princeton: Princeton University Press.



66. Liley, N.R., *Ethological isolating mechanisms in four sympatric species of Poeciliid fishes*. Behaviour, 1966. **13 (suppl.)**: p. 1 -197.
67. Schlupp, I. and M.J. Ryan, *Male Sailfin mollies (Poecilia latipinna) copy the mate choice of other males*. Behavioral Ecology, 1997. **8(1)**: p. 104-107.
68. Plath, M. and D. Bierbach, *Sex and the public: Social eavesdropping, sperm competition risk and male mate choice*. Communicative and Integrative Biology, 2011. **4(3)**: p. 276-80.
69. Magurran, A.E., *Sexual coercion*, in *Ecology and evolution of poeciliid fishes*, J.P. Evans, A. Pilastro, and I. Schlupp, Editors. 2011, Chicago University Press: Chicago. p. 209-217.
70. Evans, J.P. and A.E. Magurran, *Multiple benefits of multiple mating in guppies*. Proceedings of the National Academy of Sciences, 2000. **97(18)**: p. 10074-10076.
71. Killen, S.S., et al., *Male sexually coercive behaviour drives increased swimming efficiency in female guppies*. Functional Ecology, 2015: p. n/a-n/a.
72. Dadda, M., *Female social response to male sexual harassment in poeciliid fish: A comparison of six species*. Frontiers in Psychology, 2015. **6**.
73. Gasparini, C., A. Devigili, and A. Pilastro, *Cross-generational effects of sexual harassment on female fitness in the guppy*. Evolution, 2011: p. no-no.
74. Evans, J.P., A. Pilastro, and I.W. Ramnarine, *Sperm transfer through forced matings and its evolutionary implications in natural guppy (Poecilia reticulata ) populations*. Biological Journal of the Linnean Society, 2003. **78(4)**: p. 605-612.
75. Laubu, C., et al., *Mismatched partners that achieve postpairing behavioral similarity improve their reproductive success*. Science Advances, 2016. **2(3)**: p. e1501013.
76. Sinclair, E.L.E., et al., *Exercise changes behaviour*. Functional Ecology, 2014. **28(3)**: p. 652-659.
77. Brown, C., *Familiarity with the test environment improves escape responses in the crimson spotted rainbowfish, Melanotaenia duboulayi*. Animal Cognition, 2001. **4(2)**: p. 109-113.
78. Jennions, M.D. and M. Petrie, *Variation in mate choice and mating preferences: A review of causes and consequences*. Biological Reviews, 1997. **72(2)**: p. 283-327.