

High Frequency of Viral Co-Detections in Acute Bronchiolitis

Authors:

Hortense PETAT^{1,2,3}, Vincent GAJDOS⁴, François ANGOULVANT^{5,6}, Pierre-Olivier VIDALAIN^{7,8}, Sandrine CORBET^{1,2}, Christophe MARGUET^{1,3}, Jacques BROUARD^{1,9}, Astrid VABRET^{1,2}, Meriadeg AR GOUILH^{1,2}.

1 Groupe de Recherche sur l'Adaptation Microbienne (GRAM 2.0), Normandie Univ, UNICAEN, UNIROUEN, EA2656, Caen F-14033, France

2 Laboratoire de Virologie, Centre Hospitalo-Universitaire, Caen F-14033, France

3 Département de pédiatrie médicale, Centre hospitalier universitaire de Rouen, EA2656 université de Normandie, UNIRouen, F-7600 Rouen, France.

4 Antoine Béclère University Hospital, APHP, Paris, France

5 Assistance Publique - Hôpitaux de Paris, Paediatric Emergency Department, Necker-Enfants Malades University Hospital, Université de Paris, Paris, France

6 INSERM, Centre de Recherche des Cordeliers, UMRS 1138, Sorbonne Université, Université de Paris, Paris, France.

7 Equipe Chimie et Biologie, Modélisation et Immunologie pour la Thérapie (CBMIT), Université Paris Descartes, CNRS UMR 8601, 75006 Paris, France

8 Equipe VIRIMI - Infections Virales, Métabolisme et Immunité, Centre International de Recherche en Infectiologie (CIRI), INSERM U1111, CNRS UMR5308, Université Lyon 1, ENS de Lyon, France

9 Service de Pédiatrie Médicale, CHU Caen, France

Corresponding author : Hortense Petat, hortense.petat@univ-rouen.fr

Keywords: Respiratory Syncytial Virus, Molecular Diagnostic, Respiratory Viruses, Coronavirus, Bronchiolitis

Abstract :

Over two years (2012-2014), 719 nasopharyngeal samples were collected from 6 weeks to 12 months old infants presenting at emergency department with moderate to severe acute bronchiolitis. Viral testing was performed and we found 98% positive samples including 90% Respiratory Syncytial Virus, 34% Human Rhino Virus, and 55% viral co-detections with predominance of RSV/HRV co-infections (30%). Interestingly, we found that the risk of being infected by HRV is higher in the absence of RSV, suggesting interferences or exclusion mechanisms between these two viruses. Conversely, Coronavirus infection had no impact on the likelihood of co-infection involving HRV and RSV. Bronchiolitis is the leading cause of hospitalizations in infants before 12 months of age, and many questions about its role in the later chronic respiratory diseases (asthma and chronic obstructive pulmonary disease) do exist. Role of virus detection and burden of viral codetections need to be further explored, in order to understand the physiopathology of chronic respiratory diseases, a major public health issue.

Article:

1. Introduction

Acute bronchiolitis is the most common respiratory disease in infants under 2 years of age, and the leading cause of hospitalizations in infants before 12 months of age (1). Respiratory Syncytial Virus (RSV) is the most frequently identified virus in bronchiolitis, detected in 41 to 83% of patients (2) (3). RSV is responsible for 34 millions new cases of lower respiratory tract infections, and 2,4 millions hospitalizations of infants all over the world with 199 000 deaths per year, mostly in developing countries (4). Many other viruses are found in bronchiolitis, including human rhinoviruses (HRV),

Metapneumovirus (hMPV), coronaviruses (CoV), bocaviruses (BoV), influenza viruses, adenoviruses (ADV) and parainfluenza viruses (PIF) (2)(5). In few years, PCR methods and especially multivalent techniques have become instrumental to detect viruses associated to acute respiratory infections. Many studies evaluated the virus-dependent risk of outcomes, and the impact of viral etiology on severity and length of stay has been established (2) (3) (5) (6). Viral co-infections remain poorly investigated and available studies are scarce (2). The aim of this study was to evaluate the burden of viral co-detections in infants with acute bronchiolitis.

2. Methods

Study design, Patient Population, sampling and statistical analysis

In a prospective, multicenter, double-blinded, randomized prospective study called GUERANDE which aimed to test the efficacy of 3% hypertonic saline nebulization in acute viral bronchiolitis, respiratory samples were collected from 719 patients (7). Twenty-four French hospital centers included patients between October 15, 2012 and April 15, 2014. Infants from 6 weeks to 12 months old who were taken to emergency departments (ED) for a first episode of moderate to severe bronchiolitis were included. Nasopharyngeal swab samples were obtained for viral testing, and sent to the virology laboratory of the University Hospital Center of Caen (Normandy, France). Results from the two PCR technics were compared using Kappa's coefficient, and the risk of co-infection was calculated with Fisher's test. Written informed consent was obtained from the parents' patients. The Saint-Germain-en-Laye Ethics' Committee approved the study (reference 12020).

Sample processing and PCR assay

Swab material was resuspended into 3ml viral transport medium, divided into aliquots and stored at -80°C until processed. Total nucleic acids were extracted from 300 µL of sample and eluted in 100 µL of TAE buffer using the QiaSymphony apparatus. First aliquot was analyzed by real-time duplex RT-


qPCR with Taqman® hydrolysis probes in order to type RSV-A and RSV-B. Second aliquot was analyzed by multiplex RT-qPCR (Luminex NxTAG®RPP) targeting 18 viruses : RSV-A and B, hMPV, Influenza virus A (H1v/H3) and B, Rhinovirus/enterovirus (HRV), PIF 1-4, CoV 229E, NL63, OC43, HKU1, ADV and BoV), and 3 intracellular bacteria (*Mycoplasma pneumoniae*, *Chlamydiae pneumoniae* and *Legionella pneumophila*).

3. Results

Patients' median age was 3 months (IQR[2-5]) and 719 samples were collected in the study. According to the duplex real-time RT-PCR targeting RSV, 88% (n=633) samples were positive. RSV-A and RSV-B positive samples account for 52% (n=374) and 34% (n=244) respectively. Co-detections (RSV-A and RSV-B) account for 2% (n=12).


In comparison, the multiplex RT-PCR found only 2% (n=12) of virus negative samples (Figure 1), while 90% (n=647) were RSV positive, and 34% (n=246) were HRV positive. Co-detections (2 or more viruses) were found in 55% (n=396) of samples. The most abundant co-detection was RSV-HRV (30%, n=218), with or without another virus. The RSV-HRV co-detection rate without another virus was 21% (n=149).

Figure 1



Concerning the multiplex technic, we reported 49% (n=352) and 29% (n=206) RSV-A positive and RSV-B positive samples respectively (Figure 2). Twelve percent (n=89) samples were positive for both RSV-A and B. Among the RSV-A positive samples, 37% were mono-infected, 20% were co-infected with HRV, 10% were co-infected with RSV-B, 12% were co-infected with another virus, 14% were co-infected with 2 other viruses, and 4% were co-infected with 3 other viruses. Among the RSV-B positive samples, 37% were mono-infected, 13% were co-infected with HRV, 11% were co-infected with another virus, 10% were co-infected with 2 other viruses, and 5% was co-infected with 3 other viruses (Figure 2). Among the RSV-A and B positive samples, 52% were co-infected by another virus.

Figure 2



Forty-four (6,1%) discrepancies between the two PCR technics used were found (Figure 3). On the one hand, duplex PCR found more positives than multiplex technique for 10 RSV-A and 6 RSV-B. On the other hand, the multiplexed technique reported more positives than the duplex for 20 RSV-A, 5 RSV-B, 3 RSV A+B. Kappa's tests between the two PCR technics show a moderate accordance for RSV-A ($K=0,47$) and a good accordance for RSV-B ($K=0,66$).

Figure 3

Discrepancies

Duplex RSV-A/B	RSV-A	Multiplex		10
		+	-	
Duplex RSV-A/B	+	∅	23*	10
	-	23*	∅	6
Duplex RSV-A/B	RSV-B	∅	8*	∅
	-	8*	∅	6

*included 3 samples RSV-A and RSV-B positive in Multiplex and negative in Duplex

We compared the proportion of co-detections HRV/RSV with the proportion of mono-detections HRV: the risk of being infected by HRV is significantly higher in the absence of RSV detection ($p<0,001$, Odds Ratio= 0,09, IC_{95%} [0,0537 ; 0,1457]). The risk of being co-infected by a Coronavirus and HRV or RSV is not significantly different than being not co-infected by these viruses.

4. Discussion

Our study enrolled 719 infants aged 6 weeks to 12 months and presenting an acute moderate to severe first episode of bronchiolitis at the emergency departments in 24 different hospitals across France. Nasopharyngeal swabs were collected and virus detection was realized by RT-qPCR.

One limitation of the study consists in the selection of the cohort's participants and the resulting homogeneity of the cohort. Indeed, all patients were less than one year old, without bronchiolitis or wheezing history and their inclusion took place at the ED, where the most serious cases are usually observed. Moreover, there was no control group, in which we could estimate the rate of asymptomatic viral infections. In 2016, Self *et al.* found 24% of virus detection by asymptomatic children (8), but these controls were hospitalized for a programmed Ear, Nose and Throat (ENT) intervention, and we know that this population is characterized by frequent infections and may be not fully representative of the general population.

We found only 2% negative samples when analyzed by multiplex PCR, consistently with the literature. Mansbach *et al.* estimate this rate at 6% in 2012 in hospitalized patients with severe bronchiolitis in USA (5). A low level of negative samples in children cohort is commonly seen, especially in symptomatic infants.

A high amount of viral co-detections is found in our cohort (55%), that stands between the 34% in

the US EPIC study (6), which studied viral detection in children with pneumonia under 5 years of age, and 61% according to the ORAACLE Study Group (which was a Norwegian clinical cohort studying length of stay of hospitalized patients with bronchiolitis according to viruses found in PCR) (9). These differences could be explained by the accuracy of the new molecular detection methods used here. Nevertheless, we still ignore consequences of these co-detections, and whether this impacts the development of illnesses, particularly for future asthma or COPD (Chronic Obstructive Pulmonary Disease). HRV, which is found in 34% cases in our cohort, is known to be associated with a later asthma development (10), but no study has investigated the potential role of co-infections in the development of respiratory chronic respiratory diseases.

Several new antiviral molecules are under development (11), and their interest remains unclear, because it is not known when to use them, and how (indications, prophylaxis). Few studies focusing on the burden of viral load exist and investigate if the viral load could be a marker to follow infections and to evaluate antiviral efficacy (12). The severity of bronchiolitis is associated with a high RSV viral load (9), resulting in a longer length of stay and an increased frequency of oxygen and ventilation support. In the same way, severe ADV pneumonia have also been associated with a persistently high viral load (13). This suggests that severe illnesses exhibit a high viral replication, whereas a decline of the viral load has been observed in healing patients. We need to follow-up the viral load over acute clinical infections, for providing more data. Multiplex PCR is a qualitative but not quantitative method, and viral load cannot be calculated. Studies often rely on semi-quantitative PCR methods based on the number of Cycle Thresholds (CT), which are sometimes correlated to a viral load in copies/mL. However, these methods are not always reproducible between different commercial machines or kits, and between different centers.

Seventeen percent of cases are RSV/HRV co-detections, representing 38% of total co-detections. We still ignore the sequence of events leading to these co-detections. The potential exclusion or facilitating effects need to be further explored. For example, we do not know if RSV infection may promote HRV infection, and if co-infections occur sequentially or simultaneously. Interestingly, we

found that the risk of being infected by HRV is higher in the absence of RSV, suggesting exclusion mechanisms between these two viruses. Conversely, Coronavirus infection had no impact on the likelihood of co-infection involving HRV and RSV. To better characterize these potential interactions, it would be interesting to include mild bronchiolitis, which are cared by the general practitioners and represent the majority of bronchiolitis, and to precisely focus on the onset-to sampling delay. We propose also to collect several samples for each patient, in order to study the longitudinal history of natural infections. It would be interesting to determine the viral loads in co-detections, to know which virus(es) is/are currently active and responsible for the symptoms of our patients.

5. Patents

Funding Statement:

The funders has no role in the design of the study. The Guérande study has been funded by a grant from the French Health Ministry (P110143/IDRCB2012-A00228-35). Molecular diagnostics were supported by the University Hospital of Caen. The first author's thesis fellowship was supported by GenMark Diagnostics, Inc.

Informed Consent Statement:

Written informed consent was obtained from the parents' patients. The Saint-Germain-en-Laye Ethics' Committee approved the study (reference 12020).

Acknowledgments:

The authors wish to acknowledge the hospital centers and all staff involved in the Guérande project, the Virology unit and all contributors to the study of the University Hospital of Caen.

Conflicts of Interest:

None declared.

References

1. Florin TA, Plint AC, Zorc JJ. Viral bronchiolitis. *The Lancet*. 2017 Jan 14;389(10065):211–24.
2. Marguet C, Lubrano M, Gueudin M, Roux PL, Deschildre A, Forget C, et al. In Very Young Infants Severity of Acute Bronchiolitis Depends On Carried Viruses. *PLOS ONE*. 2009 Feb 25;4(2):e4596.
3. Miller EK, Gebretsadik T, Carroll KN, Dupont WD, Mohamed YA, Morin L, et al. Viral Etiologies of Infant Bronchiolitis, Croup and Upper Respiratory Illness During 4 Consecutive Years. *Pediatr Infect Dis J*. 2013 Sep 1;32(9):950–5.
4. Nair H, Simões EA, Rudan I, Gessner BD, Azziz-Baumgartner E, Zhang JSF, et al. Global and regional burden of hospital admissions for severe acute lower respiratory infections in young children in 2010: a systematic analysis. *The Lancet*. 2013 Apr 20;381(9875):1380–90.
5. Mansbach JM, Piedra PA, Teach SJ, Sullivan AF, Forgey T, Clark S, et al. Prospective Multicenter Study of Viral Etiology and Hospital Length of Stay in Children With Severe Bronchiolitis. *Arch Pediatr Adolesc Med*. 2012 Aug 1;166(8):700–6.
6. Jain S, Williams DJ, Arnold SR, Ampofo K, Bramley AM, Reed C, et al. Community-acquired pneumonia requiring hospitalization among U.S. children. *N Engl J Med*. 2015 Feb 26;372(9):835–45.
7. Angoulvant F, Bellétre X, Milcent K, Teglas J-P, Claudet I, Guen CGL, et al. Effect of Nebulized Hypertonic Saline Treatment in Emergency Departments on the Hospitalization Rate for Acute Bronchiolitis: A Randomized Clinical Trial. *JAMA Pediatr*. 2017 Aug 1;171(8):e171333–e171333.

8. Self WH, Williams DJ, Zhu Y, Ampofo K, Pavia AT, Chappell JD, et al. Respiratory Viral Detection in Children and Adults: Comparing Asymptomatic Controls and Patients With Community-Acquired Pneumonia. *J Infect Dis.* 2016 Feb 15;213(4):584–91.
9. Skjerven HO, Megremis S, Papadopoulos NG, Mowinckel P, Carlsen K-H, Carlsen L, et al. Virus Type and Genomic Load in Acute Bronchiolitis: Severity and Treatment Response With Inhaled Adrenaline. *J Infect Dis.* 2016 Mar 15;213(6):915–21.
10. Jartti T, Kuusipalo H, Vuorinen T, Söderlund-Venermo M, Allander T, Waris M, et al. Allergic sensitization is associated with rhinovirus-, but not other virus-, induced wheezing in children. *Pediatr Allergy Immunol Off Publ Eur Soc Pediatr Allergy Immunol.* 2010 Nov;21(7):1008–14.
11. Tripp RA, Power UF, Openshaw PJM, Kauvar LM. Respiratory Syncytial Virus: Targeting the G Protein Provides a New Approach for an Old Problem. *J Virol.* 2018 Jan 2;92(3):e01302-17.
12. Scagnolari C, Midulla F, Selvaggi C, Monteleone K, Bonci E, Papoff P, et al. Evaluation of viral load in infants hospitalized with bronchiolitis caused by respiratory syncytial virus. *Med Microbiol Immunol (Berl).* 2012 Aug 1;201(3):311–7.
13. Gu L, Qu J, Sun B, Yu X, Li H, Cao B. Sustained Viremia and High Viral Load in Respiratory Tract Secretions Are Predictors for Death in Immunocompetent Adults with Adenovirus Pneumonia. *PLOS ONE.* 2016 Aug 17;11(8):e0160777.