

Article

Breastfeeding practices among adolescent mothers and associated factors in Bangladesh (2004-2014)

Kingsley Emwinyore Agho^{1,3,7*} (K.gho@westernsydney.edu.au), Tahmeed Ahmed² (tahmeed@icddr.org), Catharine Fleming¹ (Catharine.Fleming@westernsydney.edu.au), Mansi Vijaybhai Dhami³ (18807282@student.westernsydney.edu.au), Chundung Asabe Miner⁴ (miner@unijos.edu.ng), Raphael Torome⁵ (rtorome@barmeramedical.com.au), Felix Akpojene Ogbo^{3,6} (F.Ogbo@westernsydney.edu.au), and on behalf of the Global Maternal and Child Health Research Collaboration (GloMACH)[†]

¹School of Health Sciences, Western Sydney University, Campbelltown Campus, Penrith NSW 2571, Australia

² International Centre for Diarrhoeal Disease Research, Bangladesh (Icddr).

³Translational Health Research Institute (THRI), School of Medicine, Western Sydney University, Campbelltown Campus, Penrith NSW 2571 Australia

⁴Department of Community Medicine, Faculty of Clinical Sciences, College of Health Sciences, University of Jos, Nigeria

⁵Barmera Medical Clinic [Lake Bonney Private Medical Clinic], 24 Hawdon Street, Barmera SA 5345 Australia

⁶General Practice Unit, Prescott Specialist Medical Centre, Welfare Quarters, Makurdi, Benue State 972261, Nigeria

⁷African Vision Research Institute (AVRI), University of KwaZulu-Natal, Durban, 4041, South Africa

* Correspondence: k.gho@westernsydney.edu.au Tel: 612- 46203635

Abstract: Optimal breastfeeding practices among mothers have been proven to have health and economic benefits, but evidence on breastfeeding practices among adolescent mothers in Bangladesh is limited. Hence, this study aims to estimate breastfeeding indicators and factors associated with selected feeding practices. The sample included 2554 children aged 0-23 months of adolescent mothers aged 12-19 years from four Bangladesh Demographic and Health Surveys collected between 2004 and 2014. Breastfeeding indicators were estimated using World Health Organization indicators. Selected feeding indicators were examined against potential confounding factors using univariate and multivariate analyses. Only 42.2% of adolescent mothers initiated breastfeeding within the first hour of birth, 53% exclusively breastfed their infants, predominant breastfeeding was 17.3%, and 15.7% bottle-fed their children. Parity (2-3 children), older infants and adolescent mothers who made postnatal check-up after two days were associated with increased exclusive breastfeeding (EBF) rates. Adolescent mothers aged 12-18 years and who watched television were less likely to delay breastfeeding initiation within the first hour of birth. Adolescent mothers who delivered at home (adjusted OR=2.63, 95% CI:1.86, 3.74) and made postnatal check-up after two days (adjusted OR=1.67, 95% CI: 1.21, 2.30) were significantly more likely to delay initiation breastfeeding within the first hour of birth. Younger infants, adolescent mothers living in Barisal region and who listened to radio reported increased odds of predominant breastfeeding and increased odds for bottle-feeding included male infants, an infant aged 0-5 months, Adolescent mothers who had eight or more antenatal clinic visits and richest wealth quintiles. In order for Bangladesh to meet the Sustainable Development Goals (SDGs) 2 and 3 by 2030, breastfeeding promotion programmes should discourage bottle-feeding among adolescent mothers from richest households and promote early initiation of breastfeeding especially among adolescent mothers who delivered at home and had late postnatal check-up after delivery.

Keywords: Breastfeeding; Infants; Bangladesh; Morbidity; Adolescent mothers; Mortality.

1. Introduction

In many low- and middle-income countries (LMICs), approximately 13 million adolescent girls give birth annually [1, 2], and Bangladesh has the highest adolescent pregnancy rate in Asia. An estimated 12% of adolescent Bangladeshi girls fall pregnant before the age of 19 years, and this is largely driven by high rate of child marriage [3]. This early age of marriage and subsequent pregnancy has huge implications for both the

mother and baby, as pregnancy and childbirth-related issues were the leading causes of death in adolescent girls in 2017 [4, 5], while babies born to adolescent mothers have increased risk of dying compared to those born to older mothers [6].

Appropriate breastfeeding practices increase child survival and maternal health. The mechanism, as well as the benefits of how breastfeeding increases child survival, are well documented in the scientific literature [7]. Breastfeeding prevents the introduction of potentially contaminated prelacteal foods, provides newborns with colostrum (rich in a variety of nutrients and immunoglobulins) and protects against diarrheal diseases and respiratory tract infection [7-12] – two leading causes of preventable under-five deaths in Bangladesh [13]. Numerous research from Bangladesh has identified the determinants of breastfeeding (BF) practices among mothers of all age groups [14-18]. However, specific studies that focus on breastfeeding practices of adolescent mothers are limited. Key factors associated with appropriate breastfeeding practices among Bangladeshi mothers of all age groups included non-primordial feeding practices [15], a lack of intimate partner violence and/or child abuse [14], access to mass media, home birthing [16], non-caesarean birthing, and receipt of breastfeeding counselling, antenatal and postnatal cares [16, 18]

In 2016, the World Health Organisation (WHO) and partners called for an accelerated global action to improve adolescent health, with subsequent impact on infants of adolescent mothers [4, 19]. In relation to breastfeeding practices in adolescent mothers, evidence from both high-income countries (such as Australia [20-22]) and LMICs (e.g., India [23], Brazil [24] and Nigeria [25]) has indicated that adolescent mothers are less likely to breastfeed their infants compared to older mothers. These studies found that barriers such as socioeconomic disadvantaged and a lack of prenatal attitude towards breastfeeding in Australia, and low socioeconomic status and poor health service contact (e.g., no antenatal care visits) prevented the uptake of optimal breastfeeding behaviours adolescent mothers. This specific information can help policy decision-makers and health practitioners to implement targeted interventions to increase breastfeeding in this age-group. The research also suggested that breastfeeding support and education for adolescent mothers is needed to increase their breastfeeding participation. However, the utility of this evidence to inform breastfeeding interventions among the Bangladeshi adolescent mothers is limited due to varied socio-economic, health and political contexts.

Limited studies have examined breastfeeding and complementary feeding practices among adolescent mothers in Bangladesh. For example, Qualitative research that investigated knowledge, attitudes and perceptions towards infant and young child feeding (IYCF) among adolescent mother in two rural regions in north-west Bangladesh found that knowledge of IYCF among adolescent mothers was generally low [26]. Another study that examine trends and determinants of EBF among adolescent mothers using data from both Health and Demographic Surveillance (HDSS) system areas of the International Centre for Diarrheal Disease Research, Bangladesh (icddr,b) service area (ISA) and government service area (GSA) in rural Bangladesh found that EBF prevalence was 43% on average, with no significant associated noted between the study factors and EBF in multivariate analyses [27]. Limitations of these studies are that adolescent mothers who lived in urban, semi-urban and other rural areas were excluded, and the study findings may not be generalisable to the wider Bangladeshi population. Therefore, this present study aimed to investigate the prevalence and factors associated with breastfeeding practices (i.e., early initiation of breastfeeding, exclusive breastfeeding, predominant breastfeeding and bottle feeding) among adolescent mothers in Bangladesh using nationally representative data, to identify a specific population to provide targeted breastfeeding education and support services.

2. Materials and Methods

We used the combined datasets for the years 2004, 2007, 2011 and 2014 from the Bangladesh Demographic and Health Survey (BDHS). A detail description of the survey methodology, sampling procedure, and questionnaires used for data collection is provided elsewhere [28]. BDHS datasets were collected by Population Research and Training (NIPORT) of the Ministry of Health and Family Welfare. The survey was implemented by Mitra and Associates, a Bangladeshi research firm located in Dhaka. ORC Macro of Calverton, Maryland, provided technical assistance to the project as part of its international Demographic and Health Surveys program, the BDHS covered the entire population residing in private dwellings units in the country, but 2004 and 2007 BDHS was administratively divided into six divisions while 2011 and 2014 BDHS was divided into seven divisions with Rangpur included. The BDHS collects socio-demographic, maternal and child health information from a nation-wide representative sample of households, using a three-stage sampling strategy.

A weighted total sample of 2554 adolescent mothers aged 12-19 years were included in the final analysis (2004: n=614; 2007: n=521; 2011: n=693; and 2014: n=726). The average response rate for the four surveys was 98%. For the purpose of this study, we restricted our analysis to the last-born child under 24 months, alive and living with the respondent. A weighted total of 2554 children born to adolescent mothers were assessed for the World Health Organization (WHO) defined breastfeeding indicators.

2.1. Outcome and confounding factors

In this study, the selected BF variables examined were early initiation of breastfeeding within one hour of birth (EIBF), exclusive breastfeeding (EBF), predominant breastfeeding (PBF) and bottle-feeding (BotBF) because early initiation of breastfeeding and exclusive breastfeeding plays an important role in reducing infant mortality and morbidity [29] while, predominant breastfeeding and bottle-feeding are risk factors for diarrhoeal and respiratory illness [30]. The selected breastfeeding indicators, assessed based on the WHO definitions for assessing IYCF practices [31]:

- EBF was defined as the proportion of infants 0–5 months of age who received breast milk only and no other solids or liquids except for vitamins, minerals, medicines or oral rehydration solution;
- PBF was defined as the proportion of infants 0–5 months of age who received breast milk and water, water-based liquids such as sugar water and juices but not infant formula or milk;
- EIBF was defined as the proportion of children aged 0–23 months who were put to the breast within one hour of birth and,
- BotBF was defined as the proportion of children aged 0–23 months who received any food or liquid including non-human milk and formula, any liquid (including breast milk) or semi-solid food from a bottle with nipple/teat.

Confounding factors included in the analysis were guided by previous studies [32–34] and were classified into four levels (see supplementary Table 1 for categorisation details): individual, exposure to media, household and community level. Individual-level factors relate directly to the 'mother-infant dyad' described by a previous group [35]. The attributes of the infant included age, a number of children, sex of the baby, while the characteristic of the adolescent mother included mother's age, working status, Husband education and occupation, religion, education level, maternal body mass index (BMI) and marital status. Factors related to delivery included (antenatal visits, combined place and mode of delivery, type of delivery assistance; birth order and postnatal contact) were classified under attributes of the mother-infant dyad.

Household wealth index and women's role in household decisions were included as household-level variables. The household wealth index was constructed using methods recommended by the World Bank Poverty Network and United Nations International Children's Emergency Fund (UNICEF) and described by Filmer and Pritchett [36]; and UNICEF, 2005 [37]. The principal components statistical procedure was used to determine the weights for the wealth index based on information collected about 22 household assets and facilities in order to estimate the household wealth index factor score. The household wealth index factor score (hv271) was constructed by the BDHS for each year of the survey. The household wealth index factor score for the datasets was pooled and was divided into quintiles. We then created a new variable (called the household wealth index) for the quintiles and the bottom 20% of households was arbitrarily referred to as poorest households, the next 20% as the poorer households and the top 20% as richest households as used in past studies [23, 38] (Supplementary Table 1).

In BDHS surveys, women's decision-making autonomy was assessed by information on women's participation in three different types of decisions: person who usually decides on respondent's health care, person who usually decide on making large household purchases and person who usually decides on visits to family or relatives. We categorised each of the three different types of decisions as category 1 for 'respondent alone' and 'respondent and husband/partner' and category 0 for 'husband/partner alone', 'respondent and other person' and someone else. We then created a composite score ranging from 0–3 points which was then categorized into 3 namely no decision (0 score), some decisions (1–2 scores) and all decisions (3 scores). Exposure to media variables included Newspapers, Radio and Television and the community level factors include the residential region of residence and geographical region.

2.2. Statistical Analysis

First, the selected BF indicators were dichotomized as category 1 for EBF, BotBF and PBF, and category 0 for not EBF, BotBF and PBF while, delayed initiation of breastfeeding was categorised as 1 and EIBF as 0.

Second, the sampling weights were different by survey and hence we re-normalised the sampling weights for each year of the survey to add up to 1 by computing the total sum of weights for each survey round and divide each year of survey sampling weights with the total sum of weights. This was followed by frequency tabulations to describe the characteristics of the study population and descriptive analysis for estimating 95% confidence intervals (CI) around prevalence estimates for all the breastfeeding indicators and linear interpolation was used to obtain the median duration of any breastfeeding and EBF in month.

Third, for the selected BF indicator variables, univariate logistic regression analysis was used to present as unadjusted OR (95% CI) for each confounding variables while multivariate logistic regression was used to identify factors associated with the selected breastfeeding indicators.

In the multivariate logistic regression analysis, four-stage modelling was employed. In the first stage, the individual level factors were entered into the first stage model (model 1). We conducted a manually executed elimination method to determine factors associated with each selected breastfeeding indicator ($p < 0.05$). The significant factors in Model 1 were then added to the exposure to media factors in the second stage model (model 2); this was then followed by a manually executed elimination procedure. The significant factors for combined Models 1 and 2 were added to household factors in the third model (Model 3). We used a similar approach for community-level factors in the fourth stages (Model 4), respectively. Factors associated were presented as adjusted OR (95% CI) for the variables retained in the final modelling step. These analyses were performed using Stata 'svy' commands that allow for adjustments of cluster sampling design and weights. STATA V.14.1 (STATA Corporation, College Station, Texas, USA, 2015) was employed for all the study analyses.

3. Results

3.1. Characteristics of the sample.

Table 1 lists the individual, household and community-level characteristics of the children included in this analysis. Nearly one-tenth (9.2%) of adolescent mothers had worked in the past 12 months prior to the survey, out of which, more than a third (34.9%) had a secondary or higher level of education. Approximately one in four adolescent mothers (39.5%) were from the poorer or poorest household wealth quintiles, and the majority (80%) was from rural areas. Of the respondents, 86% never read newspapers, 76% never listened to the radio, and 41% never watched television. Low BMI ($< 18.5 \text{ kg/m}^2$) was observed in 30%, one third (29.9%) had no antenatal clinic visit during pregnancy, 75.6% were home deliveries, 21.5% had health professionals available at delivery, and 12.1% were delivered by caesarean section. The frequency distribution of the individual, exposure to media, household and community level characteristics of children 0-23 months of age by year of survey are presented (see, Supplementary Table 2).

Table 1.

Individual, exposure to media, household and community level characteristics of children 0-23 months of age, Bangladesh 2004-2014 (n=2554)

Characteristic	n	%
Individual-level factors		
Mother's religion (n= 2553)		
Islam	2374	93.0
Others [§]	179	7.0
Maternal working status		
Non-working	2320	90.8
Working (past 12 months)	234	9.2
Maternal education		
No education	268	10.5
Primary	811	31.8
Secondary and higher	1475	57.8
Partner's education		
No education	596	23.3
Primary	1064	41.7
Secondary and higher	892	34.9
Husband's occupation (n= 2552)		
Non-agricultural	1582	61.9
Agricultural	667	26.1
Not working	305	12.0
Mother's age		
12- 18 years	1062	41.6
18-19 years	1492	58.4
Marital status		
Currently married	2527	99.0
Formerly married [^]	26	1.0
Birth order		

First-born	2083	81.6
2 nd -4 th	471	18.4
Preceding birth interval		
No previous birth	2083	81.6
Yes	467	18.3
Sex of baby		
Male	1303	51.0
Female	1251	49.0
Number of living children		
1	2157	84.5
2-4	396	15.5
Age of child (in months)		
0-5	745	29.2
6-11	717	28.1
12-17	628	24.6
18-23	464	18.2
Combined mode and place of delivery (n= 2548)		
Caesarean	299	12.1
Vaginal	289	11.9
Home	1955	75.7
Type of delivery assistance (n= 2543)		
Health professional	549	21.2
Non-health professional	2005	78.4
Antenatal Clinic visits		
8+	104	4.1
4-7	508	19.9
1-3	1179	46.2
None	763	29.9
Postnatal check-up		
0-2 days	631	24.7
After 2 days	295	11.6
No postnatal check-up	1628	63.8
Mother's BMI (n=2521)		
<18.5	754	29.5
18.5-24.9	1675	65.6
25+	92	3.6
Exposure to Media		
Mother reading newspapers (n= 2552)		
Not at all	2196	86.0
Yes [‡]	355	13.9
Mother listening to radio		
Not at all	1950	76.3
Yes [‡]	604	23.7
Mother watching television (n= 2552)		
Not at all	1056	41.4
Yes [‡]	1496	58.6
Household-level factors		
Household wealth Index		
Richest	497	19.5
Richer	542	21.2
Middle	507	19.9
Poorer	503	19.7
Poorest	505	19.8
Decision-making (Autonomy)		
No Decision (0 scores)	1061	41.5
Some Decisions (1-2 scores)	801	31.4

All Decisions (3 scores)	693	27.1
Community-level factors		
Residence		
Urban	516	20.2
Rural	2038	79.8
Geographical Region without Rangpur		
Barisal	145	6.1
Chittagong	534	22.5
Dhaka	791	33.3
Khulna	266	11.2
Rajshahi	494	20.8
Sylhet	145	6.1
Geographical Region with Rangpur		
Barisal	145	5.7
Chittagong	534	20.9
Dhaka	791	31.0
Khulna	266	10.4
Rajshahi	494	19.4
Sylhet	145	5.7
Rangpur	178	7.0

^divorced/separated/widow, §Hinduism/ buddhism/ Christianity/others; #less than once a week/at least once a week/almost every day. Total count 2554 unless otherwise given in brackets

3.2 Breastfeeding and infant feeding indicators

As shown in Table 2, of the total 2554 children aged 0-23 months, the ever-breast-fed rate was universal (99.7%) and, 99.7% were currently breastfed during the previous day or night. The breastfeeding continuation in the first year was 96.2% and in the second year 91.4%. EIBF was observed in only 42.2% and the exclusive breastfeeding rate in children less than 6 months of age was 53.0%. BotBF in the last 24 hours was noted in 15.7%, and only 64.9% were introduced to complementary feeds early. The median duration of exclusive breastfeeding 0-6 months was 3.6 months and for any breastfeeding was 23 months among the children aged 0-23 months. Key breastfeeding indicators among children 0-23 months of age examined as the outcome variables by year of survey are presented (see, Supplementary Table 3).

Table 2.

Breastfeeding indicators among children 0-23 months of age, Bangladesh 2004-2014 (n=2554)

Breastfeeding indicators	N*	n*	Rate (95% CI)
Early initiation of breastfeeding rate^a			
Yes	2554	1078	42.2 (39.7,44.8)
Ever breastfed rate^a			
Yes	2554	2546	99.7 (99.3, 99.9)
Bottle-feeding rate^{a^}			
Yes	2032	319	15.7 (13.7, 17.9)
Current breastfeeding rate^a			
Yes	2554	2546	99.7 (99.3, 99.9)
Age-appropriate breastfeeding			
Yes	2554	1938	75.9 (73.8,77.8)
Exclusive breastfeeding rate^b			
Yes	745	395	53.0 (48.6, 57.3)
Predominant breastfeeding rate^b			
Yes	745	129	17.3 (14.4,20.6)
Continued breastfeeding rate (1 year)^c			
Yes	413	397	96.2 (93.1, 97.9)
Continued breastfeeding rate (2 year)^d			

Yes	284	260	91.4 (86.6, 94.6)
Early introduction of complementary feeding rate^e			
Yes	361	234	64.9 (59.0, 70.4)
Median duration of any breastfeeding a (months)	23.0		
Median duration of exclusive breastfeeding a (months)	3.6		

N* = weighted population; n* = weighted sample; ^ bottle feeding not correct in 2007 DHS; a - children under 24 months; b - infants below 6 months; c - children 12-15 months; d - children 20-23 months; e - infants 6-8 months

Figure 1 showed the proportion of children born to Adolescent mothers by breastfeeding status according to age from 2004-2014. As illustrated in Figure 1, the EBF at birth was above 87%, and declined rapidly to 48% by 4 months of age. At birth, infants' proportion of breastfed plus water was 2.7% and increased to 14% at 2 months. Similarly, breastfed plus other milk was 7.2% at birth and increased to 11.7% by 2 months of age.




Figure 1.
Distribution of children born to Adolescent mothers by breastfeeding (BF) status, according to age in Bangladesh 2004-2014 (n=2554)

3.3. Determinants of selected feeding indicators

Unadjusted and adjusted odds ratios (OR) were calculated to estimate the strength of association between independent variables and four infant feeding outcomes: (1) delayed initiation of breastfeeding; (2) BotBF, (3) EBF and, (4) PBF. As shown in tables 3 and 4.

3.3.1. Factors associated with delayed initiation of breastfeeding and bottle-feeding

As shown in Table 3, adolescent mothers who delivered at home (adjusted OR=2.60, 95%CI: 1.84, 3.69; $p < 0.001$), had no ANC during pregnancy (adjusted OR=1.83, 95%CI: 1.13, 2.97; $p=0.014$) and, had postnatal check-up two days after delivery (adjusted OR=1.70, 95%CI: 1.24, 2.35; $p=0.001$) were more likely to delay initiation of breastfeeding. Adolescent mothers aged 18-19 years were less likely to delay initiation of breastfeeding (adjusted OR= 0.78, 95%CI: 0.65, 0.94; $p=0.008$ for adolescent mothers aged 18- 19 years). Delayed

initiation of breastfeeding was lower among adolescent mothers who watched TV (adjusted OR=0.76, 95%CI: 0.63, 0.93; $p=0.006$) and lived Rangpur region (adjusted OR=0.54, 95%CI: 0.36, 0.82; $p=0.004$).

Adolescent mothers who had higher secondary or more educational status (adjusted OR=0.57, 95%CI: 0.32, 0.99; $p=0.050$) were less likely to bottle feed their children (Table 3). Being female infant (adjusted OR=0.70 95%CI: 0.51, 0.96; $p=0.026$) and children aged 18-23 months of age (adjusted OR=0.61 95%CI: 0.38, 0.98; $p=0.043$) were less likely to bottle-fed. Adolescent mothers who did not have any antenatal care visit during pregnancy reported a decreased the likelihood of bottle feeding (adjusted OR=0.38, 95%CI: 0.18, 0.80 $p=0.011$) compared to adolescent mothers who had 8 or more antenatal care visits during pregnancy. Compared to adolescent mothers from richest wealth quintiles, adolescent mothers from middle (adjusted OR=0.57 95%CI: 0.36, 0.90; $p=0.015$), poor (adjusted OR=0.38 95%CI: 0.22, 0.64; $p<0.001$) and poorest (adjusted OR=0.25 95%CI: 0.14, 0.44; $p<0.001$) wealth quintiles had decreased the likelihood of bottle feeding (see columns 14-17 in table 3 for details)

Table 3

Survey logistic modelling of a child bottle-fed and delayed initiation of BF - unadjusted and adjusted Odds Ratios, Bangladesh 2004-2014

Characteristic	Delayed initiation of BF							BotBF								
	Unadjusted				Adjusted			Unadjusted			Adjusted					
	OR	[95% CI]	<i>p</i>		OR	[95% CI]	<i>p</i>	OR	[95% CI]	<i>p</i>	OR	[95% CI]	<i>p</i>			
Individual-level factors																
Mother's religion																
Others ^s (Islam, OR=1)	0.81	0.57	1.16	0.247					0.62	0.32	1.18	0.144				
Maternal working status																
Working (Non-working, OR=1)	0.87	0.64	1.17	0.359					0.90	0.52	1.57	0.703				
Marital status																
Formerly married^(Currently married, OR=1)	0.88	0.39	1.99	0.760					1.01	0.20	4.97	0.991				
Maternal education																
Primary (No education, OR=1)	0.80	0.57	1.12	0.197					0.92	0.54	1.60	0.776	0.65	0.36	1.15	0.139
Secondary and above	0.82	0.59	1.13	0.214					1.25	0.73	2.12	0.417	0.57	0.32	1.00	0.049
Partner's education																
Primary (No education, OR=1)	0.97	0.71	1.32	0.826					0.86	0.48	1.53	0.606				
Secondary and above	0.72	0.55	0.94	0.015					1.30	0.76	2.21	0.334				
Husband's occupation																
Agricultural (Non-agricultural, OR=1)	0.99	0.80	1.23	0.939					0.72	0.43	1.20	0.212				
Not working	0.77	0.58	1.03	0.080					0.96	0.63	1.46	0.851				
Mother's age																
18- 19 years (12- 18 years, OR=1)	0.81	0.68	0.98	0.028	0.79	0.66	0.95	0.014	1.00	0.73	1.37	0.995				
Birth order																
2 nd _4 th (First-born, OR=1)	0.99	0.77	1.27	0.948					0.66	0.43	1.01	0.054				
Preceding birth interval																
Yes (No previous birth, OR=1)	0.98	0.76	1.26	0.862					0.60	0.39	0.94	0.024				
Sex of baby																
Female (Male, OR=1)	1.09	0.91	1.32	0.354					0.70	0.51	0.96	0.026	0.70	0.51	0.96	0.026
Number of children																
2-4 (I child, OR=1)	1.01	0.77	1.33	0.937					0.56	0.35	0.89	0.014				
Age of child (months)																
6-11 (0-5 months, OR=1)	0.93	0.69	1.24	0.606					1.31	0.83	2.05	0.244	1.30	0.85	1.99	0.233
12-17	1.07	0.84	1.37	0.560					0.85	0.58	1.27	0.432	0.88	0.58	1.33	0.545

18-23	0.95	0.72	1.26	0.730						0.63	0.40	0.99	0.047	0.61	0.38	0.98	0.043
Combined Mode and Place of delivery																	
Vaginal (Caesarean, OR=1)	0.83	0.64	1.08	0.163	1.07	0.79	1.45	0.683		1.29	0.83	2.00	0.257				
Home	1.93	1.42	2.63	<0.001	2.60	1.86	3.69	<0.001		1.93	1.31	2.85	0.001				
Type of delivery assistance																	
Non-health professional (Health professional, OR=1)	0.79	0.64	0.98	0.029						0.62	0.45	0.86	0.004				
Antenatal Clinic visits																	
4-7 (8+, OR=1)	1.30	0.82	2.06	0.259	1.52	0.96	2.41	0.076		0.89	0.41	1.94	0.763	0.99	0.46	2.11	0.978
1-3	1.18	0.75	1.85	0.482	1.34	0.85	2.12	0.204		0.60	0.30	1.20	0.150	0.72	0.36	1.42	0.341
None	1.60	1.00	2.57	0.050	1.83	1.13	2.97	0.014		0.28	0.13	0.58	0.001	0.38	0.18	0.80	0.011
Postnatal check-up																	
After 2 days (0-2 days, OR=1)	1.61	1.17	2.20	0.003	1.70	1.24	2.35	0.001		0.91	0.56	1.48	0.707				
No postnatal check-up	1.12	0.89	1.40	0.337	1.26	0.96	1.64	0.091		0.62	0.44	0.89	0.008				
Mother's BMI																	
18.5-24.9 (<18.5, OR=1)	1.03	0.65	1.65	0.885						0.75	0.41	1.36	0.336				
25+	1.16	0.71	1.91	0.548						0.53	0.28	1.00	0.052				
Exposure to Media																	
Mothers reading Newspapers																	
Yes# (Not at all, OR=1)	1.07	0.80	1.43	0.661						1.60	1.10	2.34	0.014				
Mothers listening to the radio																	
Yes# (Not at all, OR=1)	1.28	1.02	1.60	0.033						1.30	0.92	1.83	0.137				
Mothers watching TV																	
Yes# (Not at all, OR=1)	0.81	0.67	0.98	0.028	0.76	0.63	0.94	0.006		1.72	1.15	2.57	0.009				
Household-level factors																	
Household Wealth Index																	
Richer (Richest, OR=1)	1.14	0.86	1.50	0.359						0.68	0.43	1.08	0.104	0.75	0.47	1.20	0.225
Middle	1.07	0.78	1.45	0.685						0.48	0.31	0.75	0.001	0.57	0.36	0.90	0.015
Poorer	1.05	0.80	1.38	0.729						0.32	0.19	0.53	<0.001	0.38	0.22	0.64	<0.001
Poorest	1.00	0.72	1.37	0.976						0.22	0.13	0.39	<0.001	0.25	0.14	0.44	<0.001
Decisions women have final say																	
Some Decisions (No Decision, OR=1)	1.06	0.85	1.32	0.581						0.97	0.69	1.36	0.877				
All Decisions	0.82	0.65	1.03	0.081						0.82	0.54	1.21	0.313				
Residence																	
Rural (Urban, OR=1)	0.96	0.77	1.19	0.700						0.63	0.46	0.86	0.003				
Geographical Region without Rangpur																	
Chittagong (Barisal, OR=1)	1.03	0.71	1.48	0.890						0.84	0.46	1.53	0.564				

Dhaka	0.91	0.64	1.29	0.584					2.21	1.27	3.86	0.005
Khulna	0.68	0.46	0.98	0.041					0.72	0.38	1.35	0.308
Rajshahi	0.83	0.59	1.19	0.310					1.29	0.71	2.34	0.401
Sylhet	0.77	0.53	1.11	0.166					1.21	0.64	2.28	0.560
Geographical Region with Rangpur												
Chittagong (Barisal, OR=1)	1.03	0.71	1.48	0.890	1.12	0.77	1.63	0.567	0.84	0.46	1.53	0.564
Dhaka	0.91	0.64	1.29	0.584	0.98	0.69	1.40	0.913	2.21	1.27	3.86	0.005
Khulna	0.68	0.46	0.98	0.041	0.73	0.51	1.08	0.120	0.72	0.38	1.35	0.308
Rajshahi	0.83	0.59	1.19	0.310	0.92	0.64	1.12	0.656	1.29	0.71	2.34	0.401
Sylhet	0.77	0.53	1.12	0.168	0.79	0.54	1.16	0.211	1.21	0.64	2.28	0.559
Rangpur	0.49	0.33	0.74	0.001	0.54	0.36	0.82	0.004	0.72	0.31	1.71	0.461

^divorced/separated/widow, \$Hinduism/ buddhism/ Christianity/others; #less than once a week/at least once a week/almost every day

3.3.2. *Factors associated with exclusive breastfeeding and predominant breastfeeding*

As expected, EBF was more likely in younger children (Table 4) because the child's age increases, as EBF decreases. Adolescent mothers who had more than one child reported higher EBF and those adolescent mothers **who had postnatal care after 2 days** had higher likelihood of EBF. The infant from Adolescent mothers who resides in Chittagong, Khulna, Sylhet and Rangpur significantly had a positive impact on EBF.

Child's age increases as PBF increases. Adolescent mothers who reside in Chittagong, Dhaka, Rajshahi and Rangpur significantly reported lower PBF and adolescent mothers **who listened to radio** had a greater likelihood of PBF.

Table 4 Survey logistics modelling of a child being breastfed and PBF - unadjusted and adjusted Odds Ratios, Bangladesh 2004-2014

Characteristic	EBF								PBF							
	Unadjusted				Adjusted				Unadjusted				Adjusted			
	OR	[95% CI]	p		OR	[95% CI]	p		OR	[95% CI]	p		OR	[95% CI]	p	
<i>Individual-level factors</i>																
Mother's religion																
Others ^a (Islam, OR=1)	1.74	0.84	3.58	0.133					0.80	0.23	2.86	0.735				
Maternal working status																
Working (Non-working, OR=1)	0.64	0.30	1.39	0.262					2.53	1.02	6.29	0.045				
Marital status																
Formerly married ^a (Currently married, OR=1)	*	*	*	*					*	*	*	*				
Maternal education																
Primary (No education, OR=1)	0.70	0.37	1.32	0.263					1.07	0.46	2.48	0.876				
Secondary and above	0.76	0.44	1.31	0.326					1.74	0.80	3.78	0.160				
Partner's education																
Primary (No education, OR=1)	1.24	0.78	1.97	0.360					0.87	0.50	1.51	0.615				
Secondary and above	1.48	0.92	2.38	0.102					0.84	0.47	1.50	0.566				
Husband's occupation																
Agricultural (Non-agricultural, OR=1)	0.98	0.64	1.50	0.921					1.37	0.80	2.35	0.252				
Not working	1.13	0.64	2.00	0.669					1.09	0.58	2.05	0.791				
Mother's age																
18- 19 years (12- 18 years, OR=1)	1.30	0.89	1.88	0.170					1.07	0.70	1.64	0.758				
Birth order																
2 nd -4 th (First-born, OR=1)	1.38	0.89	2.15	0.149					0.82	0.49	1.36	0.439				
Preceding birth interval																
Yes (No previous birth, OR=1)	1.38	0.89	2.15	0.149					0.82	0.49	1.36	0.439				
Sex of baby																
Female (Male, OR=1)	1.23	0.85	1.79	0.278					0.69	0.45	1.05	0.082				
Number of children																
2-4 (1 child, OR=1)	1.62	1.02	2.59	0.041	1.72	1.00	2.97	0.050	0.72	0.41	1.26	0.254				
Age of child (months)																
	0.57	0.51	0.65	<0.001	0.55	0.48	0.62	<0.001	1.17	1.03	1.33	0.018	1.16	1.02	1.315	0.029

Exposure to Media													
Mother reading Newspapers													
Yes# (Not at all, OR=1)	0.81	0.52	1.28	0.367		1.43	0.83	2.48	0.196				
Mother listening to the radio													
Yes# (Not at all, OR=1)	0.71	0.47	1.07	0.100		1.61	1.02	2.54	0.040	1.71	1.07	2.73	0.024
Mother watching television													
Yes# (Not at all, OR=1)	0.83	0.57	1.21	0.336		1.15	0.74	1.77	0.531				
Household-level factors													

Chittagong (Barisal, OR=1)	3.14	1.70	5.82	<0.001						0.46	0.23	0.93	0.030				
Dhaka	1.53	0.83	2.82	0.171						0.32	0.16	0.66	0.002				
Khulna	3.10	1.66	5.78	<0.001						0.53	0.25	1.10	0.089				
Rajshahi	1.42	0.78	2.59	0.254						0.41	0.20	0.84	0.015				
Sylhet	4.13	2.00	8.56	<0.001						0.58	0.26	1.29	0.178				
Geographical Region with Rangpur																	
Chittagong (Barisal, OR=1)	3.14	1.70	5.82	<0.001	3.19	1.59	6.41	0.001		0.46	0.23	0.93	0.030	0.45	0.22	0.92	0.029
Dhaka	1.53	0.83	2.82	0.171	1.33	0.71	2.50	0.372		0.32	0.16	0.66	0.002	0.31	0.15	0.64	0.002
Khulna	3.10	1.66	5.78	<0.001	3.62	1.80	7.29	<0.001		0.53	0.25	1.10	0.089	0.50	0.24	1.03	0.061
Rajshahi	1.42	0.78	2.59	0.254	1.34	0.69	2.62	0.389		0.41	0.20	0.84	0.015	0.35	0.18	0.80	0.011
Sylhet	4.13	1.97	8.65	<0.001	5.11	2.24	11.63	<0.001		0.58	0.26	1.30	0.184	0.57	0.25	1.28	0.173
Rangpur	6.42	2.71	15.22	<0.001	6.10	2.20	16.86	0.001		0.19	0.06	0.54	0.002	0.20	0.07	0.58	0.003

^divorced/separated/widow, \$Hinduism/ buddhism/ Christianity/others; #less than once a week/at least once a week/almost every day, * values and 95%CI too wide due to small sample size.

4. Discussion

Our results indicated that EBF and EIBF rates among adolescent mothers were sub-optimal because less than half of adolescent mothers had EIBF, and that only half of the adolescent mothers EBF their infants aged less than 6 months and need further improvement in order to gain the full benefits of breastfeeding including health and nutritional status of their children. Multivariate analyses showed that child age in months, parity, exposure to media (radio), postnatal check-up after 2 days and geographical region were associated with EBF and PBF. EIBF was greater among adolescent mothers aged 18-19 years and exposure to media (Television). Infant delivered at home, had no ANC and had postnatal check-up after two days after delivery reported a decreased likelihood of EIBF. Children from adolescent mothers who had secondary or more educational status, female infants, children aged 18-23 months, children from adolescent mothers who had no ANC and children from the middle, poor and poorest wealth quintiles had decreased likelihood of bottle feeding.

Our study found that adolescent mothers aged <18 years reported a decreased likelihood of EIBF compared to those aged 18-19 years. This finding was consistent with a cross-sectional study conducted in Nigeria which found that EIBF among adolescent mothers aged 12-18 years reduced by the crude odds of 36% compared with young women (20-24.9 years) [39]. Younger mothers lack experience, education, attitude and the necessary social support required for the appropriate breastfeeding practices [40]. We found the relationship between exposure to media (television) and EIBF. A study conducted in 30 countries in Sub-Saharan Africa revealed that women who watched television had a higher likelihood to make useful decision about breastfeeding [41]. Additionally, the association could also be attributed to the fact that adolescent mothers had a higher likelihood of being exposed to media campaigns and public health messaging regarding appropriate IYCF practices [40] because in South Asia, higher economic status strongly correlates with access to media [42],

It was observed that adolescent mothers who had no ANC during pregnancy reported delayed initiation of breastfeeding. This finding supported a cross-sectional study that no ANC during pregnancy was less likely to EIBF conducted in Bangladesh [43] and India [44] but contradicts a cross-sectional study that examined trends and factors associated with breastfeeding practices among adolescent and younger mothers in Nigeria that found adolescent mothers who had ANC were less likely to delayed [45]. A past study conducted in Bangladesh indicated that among women that had ANC, about half of them were accompanied by their partners/husbands to the health facility [46]. Partners/husbands play a significant role in improving breastfeeding, including EIBF [47]. BF information to partners should be timely at antenatal and postnatal periods, and partner-to-partner BF support groups using social media such as Facebook are also needed to improve BF including EIBF among adolescent mothers in Bangladesh.

Delayed breastfeeding initiation is more prevalent among adolescent mothers who delivered at home and those who had postnatal care visit 2 days after delivery. These findings are consistent with a cross-sectional study conducted in Nigeria, and secondary analysis of the WHO Global Survey conducted in 24 countries which found that adolescent mothers who utilise health service (delivered at the health facility and early postnatal care) were more likely to initiate breastfeeding within one hour of birth and late or absence postnatal care affect EIBF [48]. Although the finding that adolescent mothers who delivered at the health facility were more likely to initiate breastfeeding within one hour of birth seems counterintuitive because the previous study in Bangladesh revealed that mothers aged 18-49 years were more likely to initiate breastfeeding within one hour of birth if they give birth home than in health facility [49]. However, community-based breastfeeding counselling intervention including health education messages similar to those received through the Baby-Friendly Hospital Initiative (BFHI) may be needed to increase the likelihood of EIBF among those adolescent mothers who delivered at home.

Adolescent mothers who lived Khulna and Rangpur division were more likely to initiate early breastfeeding compared with adolescent mothers who lived in Barisal division. The finding that Rangpur division was more likely to initiate EIBF supported a previous study on the prevalence and determinants of EIBF in Bangladesh conducted in 2019 [43] but our findings could be attributed to the differences in the proportion of adolescent women who initiated breastfeeding early. However, further research is needed to explore the differences in region-specific cultural practices which could be associated with EIBF to improve the effectiveness of EIBF program among adolescent mothers in the country.

Adolescent mothers who had higher education were less likely to bottle-feed their children than those with no schooling. This finding was supported by a community-based cross-sectional study conducted among economically disadvantaged mothers in Karachi, Pakistan found that illiterate and less educated mothers were significantly more likely to bottle-feed their children than more educated mothers [50]. The increased likelihood of bottle-feeding among adolescent mothers with no schooling may be due to limited economic resources and higher maternal education increases opportunities to make informed child health-related decisions. These decisions include the uptake and practice of appropriate IYCF, which can improve a mother's attitude towards seeking appropriate child health support for appropriate IYCF [51]. Additionally, our study found that female

infants were less likely to be bottle-fed than male infants and adolescent mothers from poor and poorest wealth quintiles were less likely to bottle-feed their children compared to those from richest wealth quintiles. These findings were inconsistent with a cross-sectional study conducted on bottle-feeding in Indonesia which found that richest family reported an increased likelihood of bottle feeding and female child were less likely to bottle-feed than their counterparts [52]. The significantly lower odds of bottle feeding practice among poor mothers reported in this study may be attributed to the inability to finance or have access to infant formula or costly complementary substitutes to breast milk, which may result in poor adolescent mothers depending solely on breastfeeding [53].

Older infant aged 18-23 months were less likely to receive BotBF, but the main effect of children age across the categories did not differ statistically (Adjusted Wald $\chi^2 = 2.22$, $P = 0.085$). This finding contradicts previous studies which reported that increased age of infant was associated with bottle-feeding [52, 54]. There is also a possibility that this finding could be affected by the number of children bottle-fed in different age categories with about one-third of children aged 0-17 months were bottle-fed compared to 16.2% of children aged 18-23 months. We found that adolescent mothers who had no antenatal clinic (ANC) visits during pregnancy were less likely to bottle-feed their children compared to those adolescent mothers who had eight or more ANC visits during pregnancy. The finding was supported by a cross-sectional study conducted in Pakistan which reported that the increased likelihood bottle-feeding among adolescent mothers who had adequate (4 or more) ANC visits during pregnancy [55]. However, it was unexpected to find that mothers who had adequate ANC visits during pregnancy were more likely to bottle-feed their children than those who had no visits because of the breastfeeding counselling and health education messages received through the Baby-Friendly Hospital Initiative (BFHI). Hence, there is a need for community-based health education interventions (such as father/partners involvement in breastfeeding [56]) and nutritionally appropriate information sessions for new adolescent mothers to have better opportunities to make informed decisions about their child health.

In our study, adolescent mothers who had two or more children, younger infants, living in Chittagong, Khulna, Sylhet and Rangpur divisions had higher odds of EBF. These findings are consistent with a past cross-sectional study conducted in Bangladesh, and 13 Economic Community of West African States (ECOWAS) countries which found that higher birth order was associated with increased exclusive breastfeeding rates [16, 32, 57]. The finding that adolescent mothers who had older infants reported lower odds of EBF could be attributed to the duration of maternity leave because infant adolescent mothers may be forced to return to full-time work with a shorter breastfeeding span given Bangladesh recent impressive economic growth and changes in the organisation of work, such as part-time working which has been reported to increase breastfeeding span [58]. These findings may be related to inexperience regarding breastfeeding among first time adolescent mothers including inadequate use of maternal health services, lack of effective breastfeeding intervention programs among adolescent mothers who worked at the factory and previous research found an association between mothers age and inadequate use of maternal health services [59].

PBF was less likely in younger infants than older infants. This finding was consistent with a cross-sectional study conducted in Malawi, which revealed that the odds of PBF was higher in older infants than younger infants under 6 months of age [60] and PBF (BF plus water) should be discouraged at 6 months because it is a risk factor for diarrhoeal and respiratory illness in infants while, EIBF and EBF should be encouraged because they play a vital role in protecting infants against diarrhoeal diseases and reduce mortality [61]. EBF increased among adolescent mothers who had postnatal check-up after 2 days and a prospective cohort study on the effect of timing of first postnatal care home visit on neonatal mortality conducted in Bangladesh revealed that neonatal mortality reduced by about 21% among women who had postnatal check-up after 2 days after delivery compared to those with no postnatal check-up [62].

Our study found that PBF increased among adolescent mothers who listened to radio than those who did not listen to the radio. This finding supported a study conducted in Bangladesh which suggested the need to reinforce breastfeeding messages through various mass media and backed these messages during ANC visits [63]. The survey conducted by Faruque et al. [63] further suggested the need to closely monitor those breastfeeding messages to ensure that they are exact and should reach adolescent mothers with low literacy level, limited exposure to electronic and print media. Adolescent mothers living in Chittagong, Dhaka, Rajshahi and Rangpur divisions reported increased odds of predominant breastfeeding and predominant breastfeeding was less likely in younger infants, and the past study revealed that Chittagong, Dhaka and Rajshahi reported the lower prevalence of PBF than Sylhet [57].

This study has several strengths and limitations. First, this was the first study to examine breastfeeding practices among adolescent mothers and associated factors in Bangladesh using nation-wide population-based surveys because appropriate breastfeeding practices contributed to two Sustainable Development Goals (SDGs)—SDG 2 and SDG 3, which are the combination of improved nutrition and increase maternal and child health. Second, the study has great statistical power because four datasets (2004, 2007, 2011, 2014) were combined

to create a large sample size and to detect statistical differences. Third, our analysis was restricted to alive and living with the respondent to reduce adolescent mothers recall bias and are less likely to be influenced by selection bias due to an extremely high response rate and the data collection was done over a validated standardised questionnaire. However, this study also has some limitations. First, the cross-sectional study design limits causal inference and recall bias may have influenced our findings because self-reported data were collected and analysed. Second, there are chances that there could have been some misclassification bias and, that may have led to an overestimation or underestimation of the relevant breastfeeding indicators. Third, we did not account for all the confounding factors (e.g. cultural beliefs, birth weight, influences of food safety measures) may have impacted factors associated with breastfeeding practices among adolescent mothers are not included in the analysis. Fourth, bottle-feeding was not collected in the 2007 BDHS. Fifth, it is difficult to establish a clear temporal association between the factors associated and the breastfeeding practices due to the cross-sectional data of the study. Lastly, some maternal and infant health factors (such as gestational age, Apgar score, birth weight and maternal gynecological history, e.g., miscarriage) may be important variables for breastfeeding outcomes, but were not examined in this study due to data availability.

6. Conclusion

The present study suggests considerable variations in the prevalence of the key breastfeeding indicators among adolescent mothers in Bangladesh. Factors associated with the breastfeeding indicators also varied across the individual, community, and the national level. At the individual, BF information to adolescent mothers and their partners should be early and consistent at hospitals, birthing units, community midwives and health visitors from the Upazila. At the household levels, increasing education for adolescent mothers on the best BF practices, appropriate BF education to partners and their mothers-in-law of adolescent mothers about breastfeeding and improving health service use for both the adolescent mother and her baby should be encouraged. At the community level, improving geography-specific access to breastfeeding messages will improve breastfeeding practices of adolescent mothers.

Author Contributions: “conceptualization, K.E.A. and F.A.O.; methodology, K.E.A. and M.V.D.; software, K.E.A.; validation, K.E.A, M.V.D and F.A.O.; formal analysis, K.E.A, F.A.O.; investigation, T.A., C.F.; resources, T.A., R.T; data curation, K.E.A, F.A.O., C.A.M and M.V.D; writing—original draft preparation, K.E.A and F.A.O.; writing—review and editing, C.F, C.A.M, R.T, T.A.; visualization, M.V.D., C.F; supervision, K.E.A., F.A.O. and T.A.; project administration, C.A.M., M.V.D. and K.E.A.

Funding: This research received no external funding.

Acknowledgments: The authors are grateful to Measure DHS, ICF International, Rockville, Maryland, USA for providing the data for this study.

GloMACH: members are Kingsley E Agho, Felix A Ogbo, Thierno Diallo, Osita E Ezeh, Osuagwu L Uchechukwu, Pramesh R Ghimire, Blessing Akombi, Paschal Ogeleka, Tanvir Abir, Abukari I Issaka, Kedir Yimam Ahmed, Nilu Nagdev and Mansi Dhami.

Conflicts of Interest: The authors declare no conflict of interest. K.E.A is a member of the Editorial Board for the *Nutrients* and does not have any role in the journal review and decision-making process for this manuscript

References

1. UNFPA: **Girlhood, not motherhood: Preventing adolescent pregnancy**. New York:: UNFPA; 2020.
2. Darroch J, Woog V, Bankole A, Ashford L.S: **Adding it up: Costs and benefits of meeting the contraceptive needs of adolescents**. New York: Guttmacher Institute; 2016.
3. **Young People** [<https://bangladesh.unfpa.org/en/topics/young-people-10>]
4. World Health Organization: **Global Accelerated Action for the Health of Adolescents (AA-HA!): guidance to support country implementation**. Geneva: World Health Organization; 2017.
5. Reiner RC, Olsen HE, Ikeda CT, Echko MM, Ballestreros KE, Manguerra H, Martopullo I, Millier A, Shields C, Smith A: **Diseases, injuries, and risk factors in child and adolescent health, 1990 to 2017: findings from the Global Burden of Diseases, Injuries, and Risk Factors 2017 Study**. *JAMA pediatrics* 2019, **173**:e190337-e190337.
6. WHO, UNICEF, Mathers C: **Global strategy for women's, children's and adolescents' health (2016-2030)**. Geneva: World Health Organization; 2016.
7. Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, Murch S, Sankar MJ, Walker N, Rollins NC: **Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect**. *The Lancet* 2016, **387**:475-490.
8. Ogbo FA, Page A, Idoko J, Claudio F, Agho KE: **Diarrhoea and Suboptimal Feeding Practices in Nigeria: Evidence from the National Household Surveys**. *Paediatric and Perinatal Epidemiology* 2016, **30**:346-355.
9. Ogbo FA, Agho K, Ogeleka P, Woolfenden S, Page A, Eastwood J: **Infant feeding practices and diarrhoea in sub-Saharan African countries with high diarrhoea mortality**. *Plos One* 2017, **12**:e0171792.
10. Turfkruyer M, Verhasselt V: **Breast milk and its impact on maturation of the neonatal immune system**. *Current opinion in infectious diseases* 2015, **28**:199-206.
11. Wang H, Bhutta ZA, Coates MM, Coggeshall M, Dandona L, Diallo K, Franca EB, Fraser M, Fullman N, Gething PW: **Global, regional, national, and selected subnational levels of stillbirths, neonatal, infant, and under-5 mortality, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015**. *The Lancet* 2016, **388**:1725-1774.
12. Ahmed KY, Page A, Arora A, Ogbo FA, Maternal G, collaboration CHR: **Associations between infant and young child feeding practices and acute respiratory infection and diarrhoea in Ethiopia: A propensity score matching approach**. *PloS one* 2020, **15**:e0230978.
13. Vos T, Lim SS, Abbafati C, Abbas KM, Abbasi M, Abbasifard M, Abbasi-Kangevari M, Abbastabar H, Abd-Allah F, Abdelalim A: **Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019**. *The Lancet* 2020, **396**:1204-1222.
14. Islam MJ, Baird K, Mazerolle P, Broidy L: **Exploring the influence of psychosocial factors on exclusive breastfeeding in Bangladesh**. *Archives of women's mental health* 2017, **20**:173-188.
15. Nguyen PH, Kim SS, Tran LM, Menon P, Frongillo EA: **Early breastfeeding practices contribute to exclusive breastfeeding in Bangladesh, Vietnam and Ethiopia**. *Maternal & Child Nutrition* 2020:e13012.
16. Hossain M, Islam A, Kamarul T, Hossain G: **Exclusive breastfeeding practice during first six months of an infant's life in Bangladesh: a country based cross-sectional study**. *BMC pediatrics* 2018, **18**:93.
17. Rahman MA, Khan MN, Akter S, Rahman A, Alam MM, Khan MA, Rahman MM: **Determinants of exclusive breastfeeding practice in Bangladesh: Evidence from nationally representative survey data**. *Plos one* 2020, **15**:e0236080.
18. Karim F, Khan ANS, Tasnim F, Chowdhury MAK, Billah SM, Karim T, Arifeen SE, Garnett SP: **Prevalence and determinants of initiation of breastfeeding within one hour of birth: An**

- analysis of the Bangladesh Demographic and Health Survey, 2014. *PloS one* 2019, **14**:e0220224.
19. Patton GC, Sawyer, S. M., Santelli, J. S., Ross, D. A., Afifi, R., Allen, N. B., Arora, M., Azzopardi, P., Baldwin, W., Bonell, C., Kakuma, R., Kennedy, E., Mahon, J., McGovern, T., Mokdad, A. H., Patel, V., Petroni, S., Reavley, N., Taiwo, K., Waldfogel, J., ... Viner, R. M.: **Our future: a Lancet commission on adolescent health and wellbeing.** *Lancet (London, England)* 2016, **387**:2423–2478.
 20. Ogbo FA, Eastwood J, Page A, Arora A, McKenzie A, Jalaludin B, Tennant E, Miller E, Kohlhoff J, Noble J: **Prevalence and determinants of cessation of exclusive breastfeeding in the early postnatal period in Sydney, Australia.** *International Breastfeeding Journal* 2017, **12**:16.
 21. Ogbo FA, Ezech OK, Khanlari S, Naz S, Senanayake P, Ahmed KY, McKenzie A, Ogunsiji O, Agho K, Page A: **Determinants of exclusive breastfeeding cessation in the early postnatal period among culturally and linguistically diverse (CALD) Australian mothers.** *Nutrients* 2019, **11**:1611.
 22. Mossman M, Heaman M, Dennis C-L, Morris M: **The influence of adolescent mothers' breastfeeding confidence and attitudes on breastfeeding initiation and duration.** *Journal of Human Lactation* 2008, **24**:268-277.
 23. Senanayake P, O'Connor E, Ogbo FA: **National and rural-urban prevalence and determinants of early initiation of breastfeeding in India.** *BMC Public Health* 2019, **19**:896.
 24. Luthje EH, Mainardi TEB, Luthje GMH, Lopez EA: **Prevalence of Exclusive Breastfeeding and Factors Associated with Exclusive Breastfeeding in Adolescent Mothers in an Upper Middle Income Country.** *Am Acad Pediatrics*; 2020.
 25. Agho KE, Ogeleka P, Ogbo FA, Ezech OK, Eastwood J, Page A: **Trends and predictors of prelacteal feeding practices in Nigeria (2003–2013).** *Nutrients* 2016, **8**:462.
 26. Hackett KM, Mukta US, Jalal CS, Sellen DWJM, nutrition c: **Knowledge, attitudes and perceptions on infant and young child nutrition and feeding among adolescent girls and young mothers in rural Bangladesh.** 2015, **11**:173-189.
 27. Rahman A, Nomani D, Taneepanichskul S: **Trends and Determinants of EBF among Adolescent Children Born to Adolescent Mothers in Rural Bangladesh.** *Int J Environ Res Public Health* 2020, **17**:9315.
 28. National Institute of Population and Training Mitra and Associates D [Bangladesh], ICF International: **Bangladesh Demographic and Health Survey 2013.** (National Institute of Population and Training ed. Maryland, USA: National Institute of Population and Training and ICF International,; 2014.
 29. Victor R, Baines SK, Agho KE, Dibley MJJoB: **Determinants of breastfeeding indicators among children less than 24 months of age in Tanzania: a secondary analysis of the 2010 Tanzania Demographic and Health Survey.** 2013, **3**.
 30. Dhami MV, Ogbo FA, Diallo TM, Agho KEJJoER, Health P: **Regional analysis of associations between infant and young child feeding practices and Diarrhoea in Indian children.** 2020, **17**:4740.
 31. WHO., UNICEF, USAID., FANTA., AED., UC., DAVIS., IFPI.: **Indicators for assessing infant and young child feeding practices I.** Geneva, Switzerland: World Health Organization; 2008.
 32. Agho KE, Ezech OK, Ghimire PR, Uchechukwu OL, Stevens GJ, Tannous WK, Fleming C, Ogbo FA, Maternal GJN: **Exclusive Breastfeeding Rates and Associated Factors in 13 "Economic Community of West African States"(ECOWAS) Countries.** 2019, **11**:3007.
 33. Dhami MV, Ogbo FA, Osuagwu UL, Agho KE: **Prevalence and factors associated with complementary feeding practices among children aged 6-23 months in India: a regional analysis.** *BMC Public Health* 2019, **19**:1034.

34. Ogbo FA, Dhami MV, Awosemo AO, Olusanya BO, Olusanya J, Osuagwu UL, Ghimire PR, Page A, Agho KE: **Regional prevalence and determinants of exclusive breastfeeding in India.** *International breastfeeding journal* 2019, **14**:20.
35. Hector D, King L, Webb K, Heywood PJNSWphb: **Factors affecting breastfeeding practices. Applying a conceptual framework.** 2005, **16**:52-55.
36. Filmer D, Pritchett LH: **Estimating wealth effects without expenditure data—or tears: An application to educational enrollments in states of India** *Demography* 2001, **38**:115-132.
37. **UNICEF Statistics, Multiple Indicator Cluster Surveys (MICS) - Assessing the economic status of households, 2005.** [https://www.unicef.org/statistics/index_24302.html]
38. Ogbo FA, Dhami MV, Awosemo AO, Olusanya BO, Olusanya J, Osuagwu UL, Ghimire PR, Page A, Agho KE: **Regional prevalence and determinants of exclusive breastfeeding in India.** *Int Breastfeed J* 2019, **14**:20.
39. Benova L, Siddiqi M, Abejirinde I-OO, Badejo OJBgh: **Time trends and determinants of breastfeeding practices among adolescents and young women in Nigeria, 2003–2018.** 2020, **5**:e002516.
40. Hackett KM, Mukta US, Jalal CS, Sellen DW: **Knowledge, attitudes and perceptions on infant and young child nutrition and feeding among adolescent girls and young mothers in rural Bangladesh.** *Maternal & child nutrition* 2015, **11**:173-189.
41. Seidu A-A, Ahinkorah BO, Hagan Jr JE, Ameyaw EK, Abodey E, Odoi A, Agbaglo E, Sambah F, Tackie V, Schack TJFiP: **Mass media exposure and women's household decision-making capacity in 30 sub-Saharan African countries: Analysis of demographic and health surveys.** 2020, **11**.
42. Senarath U, Gunawardena NSJAPJoPH: **Women's autonomy in decision making for health care in South Asia.** 2009, **21**:137-143.
43. Karim F, Khan ANS, Tasnim F, Chowdhury MAK, Billah SM, Karim T, Arifeen SE, Garnett SPJPo: **Prevalence and determinants of initiation of breastfeeding within one hour of birth: An analysis of the Bangladesh Demographic and Health Survey, 2014.** 2019, **14**:e0220224.
44. Senanayake P, O'Connor E, Ogbo FAJBPH: **National and rural-urban prevalence and determinants of early initiation of breastfeeding in India.** 2019, **19**:1-13.
45. Benova L, Siddiqi M, Abejirinde I-OO, Badejo O: **Time trends and determinants of breastfeeding practices among adolescents and young women in Nigeria, 2003–2018.** *BMJ global health* 2020, **5**:e002516.
46. Rahman AE, Perkins J, Islam S, Siddique AB, Moinuddin M, Anwar MR, Mazumder T, Ansar A, Rahman MM, Raihana S, childbirth: **Knowledge and involvement of husbands in maternal and newborn health in rural Bangladesh.** *BMC pregnancy* 2018, **18**:1-12.
47. Hackett KM, Mukta US, Jalal CS, Sellen DWJBPH: **A qualitative study exploring perceived barriers to infant feeding and caregiving among adolescent girls and young women in rural Bangladesh.** 2015, **15**:771.
48. Takahashi K, Ganchimeg T, Ota E, Vogel JP, Souza JP, Laopaiboon M, Castro CP, Jayaratne K, Ortiz-Panozo E, Lumbiganon PJSR: **Prevalence of early initiation of breastfeeding and determinants of delayed initiation of breastfeeding: secondary analysis of the WHO Global Survey.** 2017, **7**:44868.
49. Karim F, Billah SM, Chowdhury MAK, Zaka N, Manu A, Arifeen SE, Khan ANSJPo: **Initiation of breastfeeding within one hour of birth and its determinants among normal vaginal deliveries at primary and secondary health facilities in Bangladesh: a case-observation study.** 2018, **13**:e0202508.
50. Shamim S, Jamalvi SW, Naz FJJoAMCA: **Determinants of bottle use amongst economically disadvantaged mothers.** 2006, **18**.
51. Naik SN, Verma A, Dhumal GG: **Knowledge, attitudes and practices regarding breastfeeding in postpartum mothers at a tertiary care institute during a public health awareness campaign.** *Breastfeeding Review* 2017, **25**:7.

52. Nasrul N, Hafid F, Ramadhan K, Suza DE, Efendi FJC, Review YS: **Factors associated with bottle feeding in children aged 0–23 months in Indonesia.** 2020, **116**:105251.
53. Flacking R, Nyqvist KH, Ewald UJEJoPH: **Effects of socioeconomic status on breastfeeding duration in mothers of preterm and term infants.** 2007, **17**:579-584.
54. Dibley MJ, Roy SK, Senarath U, Patel A, Tiwari K, Agho KE, Mhrshahi S, Food SAIFRNJ, bulletin n: **Across-country comparisons of selected infant and young child feeding indicators and associated factors in four South Asian countries.** 2010, **31**:366-379.
55. Hazir T, Akram, Dure-Samin, Nisar, Yasir B., Kazmi, N., Agho, Kingsley E., Abbasi, S., Khan, Amira M., Dibley, Michael J.: **Determinants of suboptimal breast-feeding practices in Pakistan.** *Public Health Nutrition* 2013, **16**:659-672.
56. Ogbo FA, Akombi BJ, Ahmed KY, Rwabilimbo AG, Ogbo AO, Uwaibi NE, Ezech OK, Agho KE: **Breastfeeding in the Community—How Can Partners/Fathers Help? A Systematic Review.** *International Journal of Environmental Research and Public Health* 2020, **17**:413.
57. Mhrshahi S, Kabir I, Roy SK, Agho KE, Senarath U, Dibley MJ, Food SAIFRNJ, bulletin n: **Determinants of infant and young child feeding practices in Bangladesh: secondary data analysis of Demographic and Health Survey 2004.** 2010, **31**:295-313.
58. Gila-Díaz A, Carrillo GH, López de Pablo ÁL, Arribas SM, Ramiro-Cortijo DJIJoER, Health P: **Association between maternal postpartum depression, stress, optimism, and breastfeeding pattern in the first six months.** 2020, **17**:7153.
59. Hackett K, Lenters L, Vandermorris A, LaFleur C, Newton S, Ndeki S, Zlotkin S: **How can engagement of adolescents in antenatal care be enhanced? Learning from the perspectives of young mothers in Ghana and Tanzania.** *BMC Pregnancy and Childbirth* 2019, **19**:184.
60. Chipojola R, Lee GT, Chiu H-Y, Chang P-C, Kuo S-YJIH: **Determinants of breastfeeding practices among mothers in Malawi: a population-based survey.** 2020, **12**:132-141.
61. Black RE, Allen LH, Bhutta ZA, Caulfield LE, De Onis M, Ezzati M, Mathers C, Rivera J, Maternal, lancet CUSGJT: **Maternal and child undernutrition: global and regional exposures and health consequences.** 2008, **371**:243-260.
62. Baqui AH, Ahmed S, El Arifeen S, Darmstadt GL, Rosecrans AM, Mannan I, Rahman SM, Begum N, Mahmud AB, Seraji HR: **Effect of timing of first postnatal care home visit on neonatal mortality in Bangladesh: a observational cohort study.** *Bmj* 2009, **339**.
63. Faruque ASG, Ahmed AS, Ahmed T, Islam MM, Hossain MI, Roy S, Alam N, Kabir I, Sack DAJJoh, population,, nutrition: **Nutrition: basis for healthy children and mothers in Bangladesh.** 2008, **26**:325.