

Effects and risk assessment of the polycyclic musk compounds Galaxolide® and Tonalide® on early life stages of marine organisms

Friday Ojie Ehiguese^a, M. Judit González-Delgado^a, Carmen Garrido-Perez^b, Cristiano V.M. Araújo^c and M. Laura Martín-Díaz^a

^a Physical Chemistry Department, University Institute of Marine Research (INMAR), International Campus of Excellence of the Sea (CEI.MAR), University of Cádiz, 11510 Puerto Real (Cádiz), Spain.

^b Environmental Technology Department, University Institute of Marine Research (INMAR), International Campus of Excellence of the Sea (CEI.MAR), University of Cádiz, 11510 Puerto Real (Cádiz), Spain.

^c Department of Ecology and Coastal Management, Institute of Marine Sciences of Andalusia (ICMAN-CSIC), 11510 Puerto Real, Cádiz, Spain.

Abstract

The current research investigated the environmental risk of the polycyclic musk compounds, Galaxolide® (HHCB) and Tonalide® (AHTN), in the marine environments. These substances are lipophilic, bioaccumulated and potentially biomagnified in aquatic organisms. To understand the toxicity of HHCB and AHTN, we performed acute toxicity tests by exposing marine microalgae (*Phaeodactylum tricornutum*, *Tetraselmis chuii* and *Isochrysis galbana*), crustaceans (*Artemia franciscana*), echinoderms (*Paracentrotus lividus*), bivalves (*Mytilus galloprovincialis*), fish (*Sparus aurata*) and a candidate freshwater microalga (*Raphidocelis subcapitata*) to environmentally relevant concentrations (0.005 - 5 µg/L) following standardized protocols. The effects of both substances on microalgae growth were incipient and only *I. galbana* was sensitive to HHCB and AHTN, with IC₁₀ values of 5.22 µg/L and 0.328 µg/L, respectively. Significant ($p < 0.01$) concentration dependent responses were measured in *P. lividus* and *M. galloprovincialis* larvae developments as well as *S. aurata* mortality tested with HHCB. The effect of HHCB on *P. lividus* larvae development was the most sensitive endpoint recorded, producing an EC₅₀ value of 4.07 µg/L. Our results show that HHCB represents a high risk to *P. lividus* larvae development for early life stages in marine environments.

Keywords: environmental risk assessment, fragrances, acute toxicity, growth inhibition, larvae development.

1. Introduction

Polycyclic musk compounds (PMCs) have been included in the priority lists of the European Commission existing substances regulation (OSPAR, 2004). PMCs are synthetic chemicals, produced in large quantities as a replacement for nitro musks, which have been banned because of their environmental persistence and adverse effects on humans and the environment (Gao et al., 2016a). They have wide applicability in household and personal care products such as detergents, shampoos, lotions, perfumes, as well as additives in cigarettes and fish baits (Gao et al., 2016b; Heberer, 2002). Due to their high water solubility, inherent lipophilicity and biological stability, coupled with external application (they do not undergo

biotransformation), it is not surprising to find them as contaminants in aquatic ecosystems at concentrations ranging from ng/L to µg/L (Dietrich and Hitzfeld, 2004).

In particular, the PMCs marketed as Galaxolide® (HHCB) and Tonalide® (AHTN) represent about 95% of total fragrance materials in the perfume industry (Balk and Ford, 1999a) and are the most commonly detected PMCs in environmental compartments and biological tissues (Bester et al., 1998; Díaz-Garduño et al., 2017; Fromme et al., 2001; Kannan et al., 2005; Nakata et al., 2007; Sumner et al., 2010; Villa et al., 2012). Fromme et al. (2001) detected HHCB and AHTN in surface water in Berlin (Germany) at concentration values ranging from 70 – 1590 ng/L and 20 – 530 ng/L, respectively. Similarly, Heberer (2002) also reported high levels of HHCB and AHTN in surface water in Berlin at concentrations ranging from 20 – 12500 ng/L and 30 – 1100 ng/L, respectively. Moreover, these substances have been detected in surface waters in almost every country in Europe (Dsikowitzky et al., 2002; Fromme et al., 2001; Villa et al., 2012). Although most studies were for the freshwater environment, there are data confirming their presence in the marine environment (Bester et al., 1998; Díaz-Garduño et al., 2017; Pintado-Herrera et al., 2013; Sumner et al., 2010), and reported values over 5 µg/L depending on the proximity to a sewage treatment plants (STP). Sumner et al. (2010) studied the transport of PMCs from an STP effluent into coastal waters and reported that the concentrations of HHCB and AHTN in the open sea depends on the distance from STP and the dilution power towards the open sea. For instance, the authors measured the concentration of HHCB in STP effluent ranging from 987 – 2098 ng/L, being diluted towards the sea over 2 km, to 6 – 28 ng/L. However, this level could be higher near urban coasts where STP effluents are directly discharged into coastal waters, which could represent a potential threat to the survival of nearshore organisms, an important part of the marine ecosystems.

Bioaccumulation of PMCs, particularly HHCB and AHTN, have also been reported in marine organisms including crustaceans, bivalves, fish, marine birds and mammals (Kannan et al., 2005; Moon et al., 2012; Nakata et al., 2007; Rüdel et al., 2006) at levels of environmental concern. The concern over the potential effects are not only related to the environment, but also the impacts on food safety and, consequently, public health. Therefore, Vandermeersch et al. (2015) reviewed emerging contaminants in seafood, acknowledging that HHCB and AHTN were the most commonly detected PMCs, with concentrations reaching 160 and 45 µg/g lipid weight in mollusks and fish, respectively (Gatermann et al., 2002). Similarly, in the framework of the FP7 ECsafefood project, where 62 commercial seafood samples (mackerel, tuna, salmon, seabream, cod, monkfish, crab, shrimp, octopus, perch and plaice) in the European Union were analysed for residues from personal care products, HHCB was detected at concentrations ranging from 2.5 – 414.4 µg/kg dry weight, with the highest concentration measured in fish (sole) and AHTN found at concentrations ranging from 2.5 – 12.2 µg/kg dry weight (Cunha et al., 2018). Furthermore, these compounds have the potential to elicit adverse effects in marine organisms due, respectively, to the bio-concentration factors (based on lipid content) of 3504 and 5017 (Fromme et al., 2001) and high octanol water partition coefficient (LogKow) of 5.9 and 5.7 (Balk and Ford, 1999a).

Data on the acute, sub-chronic and chronic aquatic toxicity of HHCB and AHTN are available for algae, crustaceans, mollusks, bivalves and various fish (Breitholtz et al., 2003; Carlsson et al., 2000; Carlsson and Norrgren, 2004; Gooding et al., 2006; Liu et al., 2011; Luckenbach and Epel, 2005; Parolini et al., 2015; Pedersen et al., 2009; Wollenberger et al., 2003; Yamauchi et al., 2008). Although the majority of the studies were focused on freshwater ecosystems, Breitholtz et al. (2003) and Wollenberger et al. (2003) investigated the acute, sub-lethal and lethal effects of these substances on the marine copepods, *Nitocra spinipes* and *Acartia tonsa*, respectively. While Wollenberger et al. (2003) concluded that they inhibited larval development in *A. tonsa* at low concentration and should be considered very toxic, Breitholtz et al. (2003) reported low adverse effects in *N. spinipes*.

Several attempts have been made to assess the environmental risk of HHCB and AHTN in the aquatic environment. Balk and Ford (1999b) provided an insight by using acute and chronic toxicity data from freshwater and terrestrial organisms to estimate the environmental risk, but with no reference to the marine

environment. Other ERAs performed so far (European Commission, 2008a, 2008b; HERA, 2004; OSPAR, 2004) revealed a paucity of ecotoxicity data for the marine environment needed to effectively assess the risk of these substances.

The aim of this research was to examine the toxicity of HHCB and AHTN in the marine environment by exposing organisms from different trophic levels at early life stages to environmental concentrations of HHCB and AHTN and to estimate the level of risk they might exert on the marine ecosystems. Therefore, we performed acute toxicity using marine organisms - microalgae (*Phaeodactylum tricornutum*, *Tetraselmis chuii* and *Isochrysis galbana*), crustaceans (*Artemia franciscana*), echinoderms (*Paracentrotus lividus*), bivalves (*Mytilus galloprovincialis*) and fish (*Sparus aurata*); and a candidate freshwater alga – *Raphidocelis subcapitata*. Furthermore, understanding the toxicity of HHCB and AHTN in lower trophic organisms will help to predict possible bottom-up and top-down effects, which could lead to a functional and structural disruption of the ecosystems (De Lange et al., 2010; Fleeger et al., 2003; Johnston and Roberts, 2009; Van Den Brink, 2008). The species used in the current study have been used to examine the effects of effluents from sewage treatment plants (Diaz-Garduno et al., 2016; Meric et al., 2005), metals (Moreno-Garrido et al., 2005; Yu et al., 2007), organic solvents (Okumura et al., 2001) and inorganic chemicals (Latała et al., 2009; Mariani et al., 2006) in marine environments. Also, these organisms have been endorsed by international organisations for ERA of contaminants due to their sensitivity (OECD, 2013; Environment Canada, 2011; European Commission Joint Research Centre, 2003; US EPA, 2002, 1992).

2. Materials and Methods

2.1 Chemical selections

Analytical grade HHCB and AHTN were purchased from Sigma Aldrich, Spain. The physiochemical properties, preparations in organic solvent and determination of exposure concentrations followed details found in Ehigues et al. (2019). In brief, HHCB and AHTN were dissolved in dimethyl sulfoxide (DMSO) (0.001% v/v) in glass vials to form a stock solution. Concentrations (0.005, 0.05, 0.5 and 5.0 µg/L) of each substance were prepared by diluting the stock solutions in Nanopure water.

2.2 Acute Toxicity Test

2.2.1 Microalgae growth inhibition test

Microalgae growth inhibition tests were performed using three marine and one freshwater species, following the procedure reported by Garrido-Perez et al., (2008). Inocula of *Phaeodactylum tricornutum*, *Tetraselmis chuii* and *Isochrysis galbana* (marine species) were provided by the Laboratory of Marine Culture of the University of Cádiz, Spain, and *Raphidocelis subcapitata* (strain 61.81) was provided by the Culture Collection of Algae at Goettingen University, Germany. Seawater microalgae culture media was prepared by adding nutrients (macro- and micronutrients and vitamins) according to f/2 medium (Guillard and Ryther 1962) to synthetic seawater according to the formula from USEPA (2002a). For the freshwater species, the culture media was prepared according to the concentrations of COMBO Media (Kilham et al. 1998) in Nanopure water. Before exposure to HHCB and AHTN, an inoculum from each species of microalgae was cultured in fresh media and in the same chamber of assay (19±1 °C; 11000 lux; photoperiod 14/10 light/day). Inocula were maintained for three days to reach the exponential growth phase, in order to provide acclimatised and healthy cells for the tests.

An aliquot of 0.25 mL of each inoculum was added to 3.75 mL of the culture media spiked with different concentrations of contaminants. The exposure was performed in triplicates including two controls: (1)

culture media without contaminants (normal growing of the microalga in a fresh medium), and (2) culture media without contaminants but including DMSO (to test the toxic effects of the organic solvent). The addition time of the microalgae was as short as possible, to ensure that they had similar biomass at the initial time. After 30 min inoculation, the initial biomass (B_0) was measured, representing time zero (T_0). The biomass was measured indirectly through absorbance at 680 nm (maximum chlorophyll peak in a fresh culture of microalga), with a TECAN 2000 micro-well plate reader. This measurement was repeated at 24, 48 and 72 h. The endpoint observed in this test was the inhibition of biomass growth at 72 h compared to the control. The minimal growth acceptable for the control was 16 times higher than the initial density.

2.2.2 *Artemia* toxicity test

The cysts of *A. franciscana* are certified biological material (AF450, INVE) and were acquired from Acuazul, S.C. (Spain). Before starting the trial, dehydrated artemia cysts were hatched. In a one-litre Erlenmeyer flask containing 300 mL of natural seawater (35 g/L salinity), approximately 100 mg of cysts were resuspended and left with aeration and continuous illumination throughout the hatching time, which occurs within 24–36 h. Once the artemia were hatched, the aeration was removed. The hatched nauplii were placed in clean seawater to avoid moving empty or unhatched cysts. Therefore, all nauplii used in the test belong to the same cohort. The test was carried out in triplicates using glass Petri dishes of 60 mm diameter. On each plate 10 nauplii were placed using a glass Pasteur pipette to avoid stress to the nauplii. Subsequently, seawater spiked with each contaminant was added to a total volume of 10 mL. The test was conducted at 20 ± 0.4 °C and in the dark to minimize the swimming of the nauplii and the consumption of energy. The test lasted for 72 h and records were taken at 24, 48 and 72 h. The *Artemia* were counted using a stereomicroscope at a magnification of 2x and 4x, identifying dead specimens (mortality was recorded when they showed no sign of any movement of their limbs for 30 seconds) and those that presented motility problems such as movements in circles, asynchronous, among others. The test was considered acceptable when survival of over 80% was recorded in the control group after 72 h.

2.2.3 *Sea urchin* toxicity test

Sea urchin (*P. lividus*) fertilization and larval development tests were performed following procedures described by Fernandez and Beiras (2001), and Environment Canada (2011). Individuals were collected from the uncontaminated rocky subtidal environment off the Bay of Cádiz (Spain) at 1.5 – 2 m depth. They were immediately transported to the laboratory in a cooler box. Matured individuals were dissected, and eggs and sperm were collected using a micropipette. For the fertilization test, 10 μ L of sperm were added to the aliquots containing 10 mL of the test concentrations arranged in sequence for 10 min; then, 1 mL of eggs was added to each test tube, swirled gently and allowed to proceed for 10 min. The endpoint for fertilization success was the presence of a fertilization membrane. A larval development test was performed by adding 1 mL of fertilized eggs to beakers containing test solutions in triplicate, including the controls (seawater and DMSO), in dark conditions for 48 h. The test was considered valid when development in the control was $\geq 80\%$ and the result expressed as percentage of normal pluteus stage, normalized to the corresponding seawater control. Both fertilization and larval development tests were conducted in a controlled chamber at a temperature of 20 °C.

2.2.4 *Mussels* larvae development test

The mussels *M. galloprovincialis* were purchased from an aquaculture farm in north-western Spain. The depurated mussels were transported to the laboratory at 8 °C. Animals with matured gonads were held at 9 °C for two weeks to acclimate to the laboratory condition. The toxicity test was performed following the

ASTM protocol for acute toxicity of saltwater bivalves (ASTM, 2012). Mussels were placed each in a beaker containing filtered seawater and induced to spawn by thermal stimulation at 19 °C. Eggs and sperm were filtered to remove debris using 75 µm and 37 µm screens, respectively. Before fertilization, the egg and sperm quality and density were evaluated under a microscope. Aliquots of 10 µL of sperm were added to the eggs for fertilization (10⁶/egg) and fertilization success was assessed under the microscope. Fertilized eggs were added to test solutions at 50 eggs/ml and incubated for 48 h at 16±0.2 °C. The test was performed in triplicate including seawater and solvent controls (DMSO). Samples were fixed with 40% formalin at the end of the test and 100 larvae were counted under the microscope (X 40 magnification) per replicate distinguishing between normal developed larvae (D-shaped) and malformed larvae. Test results were accepted when normal developed larvae in control was ≥80%.

2.2.5 Fish larva mortality test

The effects of environmental concentrations of HHCB and AHTN on the survival of seabream (*Sparus aurata*) larvae was tested following OECD guidelines for fish early-life stage toxicity test (OECD, 2013). Yolk-sac larvae (3 – 5 h post hatch) of *S. aurata* were obtained from the laboratory of Marine Culture, Faculty of Marine and Environmental Sciences, University of Cádiz, Spain. Individuals (n = 70) of the yolk-sac larvae were added to each beaker (Pyrex[®]) containing 600 ml of seawater spiked with the concentrations of contaminants, and each treatment was run in triplicate. Seawater and solvent (DMSO) controls were also tested; the latter was to ensure that there were no solvent driven effects. The exposure proceeded for 96 h, physiochemical parameters were monitored using the CRISON CM35+ and 40MM+ multiparameter probes (Crison-Hachs Lange S.L.U., Spain) and values recorded during the exposure were: Temperature (16.01±0.2 °C), salinity (34.2±0.3), pH (7.7±0.2) and oxygen (>5 mg/L). The test was considered valid if mortality did not exceed 10% in the control group. Mortality in each treatment group was recorded and data were expressed as the percentage of survived larvae at each experimental condition.

2.3 Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics, version 24.0. Significant differences between organisms exposed to the different concentrations of PMCs or to the organisms from control were checked using a one-way ANOVA followed by multiple comparisons of Turkey's or Dunnett's test. Statistically significant differences were set at $p < 0.05$. Spearman's rank order of correlation was calculated between the measured effects and the exposure concentrations. In addition, the ICPIN statistical programme, provided by USEPA on its website, was used for the analysis of toxicity data and to calculate the IC_x/EC_x (concentrations that cause growth inhibition or any effect, respectively, to x% of the population).

The Risk Quotient (RQ) was derived from the ratio of the measured environmental concentrations (MECs) listed in Table 1 to the PNEC (predicted no effect concentration). PNEC is the effect concentrations (EC) or inhibition concentration (IC) obtained from the tests divided by an assessment factor (European Commission Joint Research Centre, 2003).

$$RQ = \frac{MEC}{PNEC} = \frac{MEC}{EC/1000} \text{ or } \frac{MEC}{IC/1000}$$

Where:

If $RQ < 1$ = no risk expected, and no further evaluation required

If $RQ > 1$ = potential risk and further evaluation is required

3. Results

3.1 Microalgae growth inhibition tests

Data of growth inhibition for microalgae are presented in Figure 1. There was no critical difference between microalgae exposed to seawater control and DMSO. In brief, it is possible to observe that toxicity of both compounds was incipient for the four microalgae, although AHTN seemed to be slightly more toxic. For *P. tricornutum*, growth inhibition for both compounds were observed in some concentrations. However, significant differences ($p < 0.05$) in growth in relation to the control only occurred with exposure to AHTN at 0.005 $\mu\text{g/L}$ by 16.2% (Figure 1A). On the one hand, *T. chuii* growth was inhibited by AHTN and the decrease in biomass was significantly ($p < 0.01$) concentration dependent (Table S1 – Supplementary Materials). On the other hand, the exposure to HHCB produced a stimulation (hormesis) in the growth (Figure 1B) and similar biphasic responses was seen in *I. galbana* biomass after exposure to both musk compounds (Figure 1C). The freshwater microalgae, *R. subcapitata* growth decreased after exposure to HHCB and AHTN. In the case of AHTN, significant ($p < 0.05$) concentration dependent growth inhibition of *R. subcapitata* was observed (Table S1 – Supplementary Materials), and the highest inhibition occurred at 5 $\mu\text{g/L}$ by 23.5%. In contrast, *R. subcapitata* growth inhibition by HHCB was more severe at 0.05 $\mu\text{g/L}$ by 14.5% in relation to the control (Figure 1D).

3.2 Invertebrate and fish early life stage toxicity tests

For all the organisms tested for fertilization, larvae development, motility and mortality, the threshold values for the control experiments (seawater control and solvent control – DMSO) were not exceeded.

Table 1: Measured Environmental Concentrations (MEC) of Galaxolide (HHCB) and Tonalide (AHTN) in seawater from different location. Measured values are in ng/L. Maximum values (bold) were used in calculating risk quotient (RQ)

Country/Location	HHCB	AHTN	Reference
Germany (North Sea)	0.09 - 4.8	0.08 - 2.6	Bester et al., 1998
Germany (Elbe Estuary)	95 - 136	65 - 200	Bester et al., 1998
United Kingdom (Tamar Estuarine - Plym Sound)	6.00 - 30	3.00 - 15	Sumner et al., 2010
Spain (Bay of Cadiz)	230±0.1	NA	Pintado-Herrera et al, 2013
Singapore (Coastal water)	1.66 - 21.8	0.244 - 1.85	Zhang et al., 2015

NA = Not applicable




Figure 1A-1D: Microalgae exposed to galaxolide and tonalide for growth inhibition tests. % Enrichment of (A) *P. tricornotum*, (B) *T. chuii*, (C) *I. galbana* and (D) *R. subcapitata* (negative values corresponding to % growth inhibition). Asterisks (*) indicate significant differences ($p < 0.05$) in relation to control.

Moreover, there was no significant difference ($p < 0.05$) in responses detected in organisms exposed to seawater control and DMSO.

The effects of HHCB and AHTN on survival and motility of *A. franciscana* are shown in Figure 2A and 2B, respectively. Both responses showed no significant difference ($p < 0.05$) compared to the control. For the *A. franciscana* mortality test, percentage mortalities in the controls were 3.33% and 6.67% for seawater and DMSO, respectively. The effect of HHCB on the survival of *A. franciscana* was very minimal with the highest mortality of 10% recorded at a 0.5 µg/L HHCB. The effect of AHTN on *A. franciscana* mortality was also very low, with only 3.33% mortality at 0.05 and 0.5 µg/L. Other concentrations of AHTN tested had no effect on *A. franciscana* survival. However, Artemia motility test was not sensitive to either compound (Figure 2B).

The effects of HHCB and AHTN on *P. lividus* fertilization tests are presented in Figure 2C and 2D. The percentage of sea urchin that were able to fertilize in the controls were 89.50% and 90.00% for seawater and DMSO, respectively. Percentage fertilization of sea urchin tested with 0.005 µg/L, 0.05 µg/L, 0.5 µg/L

and 5 µg/L HHCB were 86.33%, 82.33%, 81.00% and 82.33%, respectively. Under exposure to AHTN, sea urchin fertilization success recorded at 0.005 µg/L, 0.05 µg/L, 0.5 µg/L and 5 µg/L were 89.00%, 83.33%, 86.33% and 86.33%, respectively. Although the effect is incipient, *P. lividus* fertilization failure induced by both substances was more pronounced for HHCB with significant effects at 0.05 – 5 µg/L, while a significant difference ($p < 0.05$) was only observed at 0.05 µg/L AHTN compared to control (Figure 2C). The results for the sea urchin larvae development tested with HHCB and AHTN are presented in Figure 2D. The percentages of larvae development in the seawater and solvent controls were 83.50% and 81.50%, respectively. The number of oocytes that were able to develop to pluteus stage was significantly ($p < 0.01$) concentration dependent (Table S2 – Supplementary Materials) and a significant ($p < 0.05$) decrease in larvae development was recorded at 5 µg/L HHCB compared to the control (Figure 2D). The percentage of larvae that were able to develop from morula stage to normal pluteus stage for 0.005 µg/L, 0.05 µg/L, 0.5 µg/L and 5 µg/L HHCB were 85.00%, 85.00%, 68.00% and 44.33%, respectively, while similar concentrations of AHTN reduced larvae development by 63.00%, 75.33%, 54.67% and 62.33%, respectively.

The results for the larval development toxicity test for *M. galloprovincialis* exposed to HHCB and AHTN are presented in Figure 2E. The number of fertilized eggs that developed after 48 h to D-veliger stage in the control was 98%. However, the embryotoxicity of HHCB to mussels was significantly ($p < 0.01$) concentration dependent (Table S2 – Supplementary Materials). From 0.05 – 5 µg/L HHCB, the percentage of abnormal larvae development was significantly ($p < 0.05$) different from the control (Figure 2E) and the percentage effect increased to 19.88% in the highest concentration tested (5 µg/L). Similarly, significant toxicity of AHTN to embryo of *M. galloprovincialis* was observed in 0.5 µg/L and 5.0 µg/L with percentages of abnormal larvae of 8.36% and 11.63%, respectively (Figure 2E).

The results of the effect of HHCB and AHTN on the survival of yolk-sac larvae of *S. aurata* larvae after 96 h exposure was significantly ($p < 0.01$) concentration dependent (Table S2 – Supplementary Materials) and the measured effect is presented in Figure 2F. After 96 h, the percentage mortality of yolk-sac larvae of *S. aurata* in seawater and DMSO controls was 5.33% each. Percentages of mortality in fish exposed to 0.005 µg/L, 0.05 µg/L, 0.5 µg/L and 5 µg/L of HHCB were 8.67%, 12.00%, 10.67% and 13.33%, respectively. On the other hand, the percentage mortality of fish tested with the same range of concentrations of AHTN were 10.00%, 7.33%, 12.67% and 13.33%, respectively.




Figure 2: *A. franciscana* mortality and motility, *P. lividus* fertilization and larval development, *M. galloprovincialis* larval development and *S. aurata* larval mortality tests exposed to galaxolide and tonalide. Asterisks (*) indicate significant differences ($p < 0.05$) in relation to control.

3.3 Risk Quotient (RQ)

Quantitative risk estimation of chemicals in the environment is achieved using monitoring data of measured environmental concentration (MEC) and predicted no effect concentration (PNEC), giving a risk quotient necessary for risk characterisation. A literature survey of MEC for both polycyclic musk compounds were undertaken, and the values are presented in Table 1. Due to incipient effects detected in this study for all the species of microalgae and animals, it was not possible to calculate the EC₅₀ values and their respective confident intervals, except for *P. lividus* larvae development tested with HHCB, producing an EC₅₀ value of 4.07 µg/L. For aquatic environments (freshwater and marine), PNEC is estimated by dividing the EC₅₀ value by an assessment factor of 1000 (EC TGD, 2003). Where possible, IC₁₀/EC₁₀ was evaluated and only *I. galbana* produced an IC₁₀ values for HHCB and AHTN (5.22 and 0.33 µg/L, respectively) that fall within the range of the concentration tested. The IC₁₀ values were also considered for risk estimation and the estimated risk for each substance tested are presented in Table 2. Our results indicated that HHCB posed a high risk to *P. lividus* larvae development with RQ between 1.18 – 56.53 and both contaminants were potential threat to *I. galbana* growth with RQ of 0.92 – 44.06 and 5.64 – 609.76 for HHCB and AHTN, respectively.

Table 2: EC₅₀ and IC₁₀ with estimated risk quotients of Galaxolide (HHCB) and Tonalide (AHTN) as the MEC (*maximum value per location*)/PNEC ratio

Species/Endpoint	substance	
	HHCB	AHTN
<i>P. lividus</i> larvae development EC ₅₀ (µg/L)	4.07	NA
MEC _{max. value per location} (ng/L)	4.8 - 230	1.85 - 200
PNEC (ng/L)	4.07	NA
RQ (MEC _{max. value per location} /PNEC)	1.18 - 56.53	NA
Risk	Yes	NA
<i>I. galbana</i> growth inhibition IC ₁₀ (µg/L)	5.22	0.328
MEC _{max. value per location} (ng/L)	4.8 - 230	1.85 - 200
PNEC (ng/L)	5.22	0.328
RQ (MEC _{max. value per location} /PNEC)	0.92 - 44.06	5.64 - 609.77
Risk	Yes	Yes

NA = Not applicable

4. Discussion

The aim of this study was to evaluate the impacts of environmentally relevant concentrations of HHCB and AHTN on marine organisms after short-term exposure and, where possible, characterise the risk following

the EU technical guidance document for ERA (European Commission Joint Research Centre, 2003). Several *in vitro* and *in vivo* toxicity tests have been performed with these compounds using freshwater species (Balk and Ford, 1999b; Carlsson and Norrgren, 2004; Gooding et al., 2006; Parolini et al., 2015; Schreurs et al., 2002; Składanowski et al., 2005) and the only tests with marine organisms used high concentrations (Breitholtz et al., 2003; Wollenberger et al., 2003), making it difficult to compare such results with actual environmental impacts. Furthermore, for evaluation of toxicity of substances, it is imperative to use a varied battery of tests because organisms are not equally susceptible to the same toxic substance.

The effects of HHCB and AHTN on *P. tricornutum*, *T. chuii*, *I. galbana* and *R. subcapitata* showed that both compounds have limited effects on microalgae growth (Figure 1). Microalgae have been used in water quality assessments as *in-situ* bio-monitors because they are primary producers at the base of the ecological trophic arrangement and the basic supplier of oxygen in aquatic ecosystems (Sibila et al., 2008). Microalgae toxicity tests are useful in ERA and have gained international recognition leading to development of test guidelines for reliable and relevant toxicity data (OECD, 2002). The potential of HHCB and AHTN to inhibit the growth of microalgae in the aquatic ecosystem have been incipiently reported. Previous studies revealed that significant acute toxicity of microalgae occurred at concentration greater than 100 µg/L or 500 µg/L depending on the species (Balk and Ford, 1999b; Ding et al., 2020) but our data suggest that even at low concentrations microalgae growth might be inhibited.

The microalgae growth inhibition tests were mildly sensitive to HHCB and AHTN; IC₁₀ values for most of the microalgae were out of range of the concentrations tested, except for *I. galbana* with IC₁₀ values of 5.220 and 0.328 µg/L for HHCB and AHTN, respectively (Table 2). In contrast, previous studies have reported higher EC₅₀ values of 0.050 mg/L for *Navicula* spp. and 0.336 mg/L for *Scenedesmus quadricauda* exposed to HHCB (Ding et al., 2020). This is because microalgae are not equally sensitive to contaminants. For example, in all the microalgae tested, we found that differential sensitivity was exhibited, the order of sensitivity to HHCB was *P. tricornutum* > *I. galbana*, and to AHTN was *T. chuii* > *P. tricornutum* > *I. galbana*. Similarly, differential sensitivity of two microalgae to HHCB was recently reported, whilst the microalgae, *Navicula* spp. are more sensitive to HHCB than *S. quadricauda* (Ding et al., 2020).

Artemia was not sensitive to either substance since the endpoints measured were not critically different from the control (Figure 2: A & B). This is not surprising because *Artemia* is known to be fairly resistant to toxic substances in comparison to other invertebrates and they respond differently to different environmental contaminants (Ferreira et al., 2007; Hadjispyrou et al., 2001).

HHCB significantly reduced the fertilization success of *P. lividus* as the concentrations increased. Meanwhile, AHTN had no significant effects on sea urchin fertilization success (Figure 2C). Sea urchin fertilization success depends on sperm fitness, motility potentiated by the axonemal engine in the flagellum, morphology and chemotactic navigation (Hussain et al., 2016). For organoleptic substances, such as HHCB and AHTN, the latter might be the most compelling factor inducing reduction in sea urchin fertilization success since the exposure procedure involved prior treatment of sperm with the contaminants before the introduction of eggs. Between the two contaminants tested, HHCB had more critical effects on *P. lividus* fertilization by 19% reduction at 0.5 µg/L, and we observed that only 0.05 µg/L AHTN significantly reduced sea urchin fertilization by 16.67% (Figure 2C). Importantly, the effect exerted by both contaminants on sea urchin fertilization success was below 20%, presenting less toxic effects compared to other contaminants of emerging concern. For example, 500 ng/L propanol, 500 ng/L 17α-ethinylestradiol and 5000 ng/L gemfibrozil reduced sea urchin fertilization success by 24.1%, 36.9% and 26.9%, respectively (Capolupo et al., 2018). Similarly, other contaminants of emerging concern belonging to pharmaceutical and personal care products have been reported to significantly affect sea urchin fertilization success at concentrations detected in the environment (Aguirre-Martínez et al., 2015; Paredes et al., 2014).

Ecotoxicity studies with early life stages of aquatic organisms have been recommended as a faster and more cost-effective means of examining chemicals and environmental samples, because newly hatched larvae are sensitive to exogenous substances as the embryos lose their protective membranes and are fully exposed

to potential xenobiotics (Beiras et al., 2003). *P. lividus* and *M. galloprovincialis* are well recognized in toxicity bioassays and are applied globally for the evaluation of toxicity of marine contaminants by the exposure of gametes to aqueous phases, such as surface waters and pore waters (Aguirre-Martínez et al., 2015; Bellas et al., 2005) and elutriates (Losso et al., 2007). Critical effects were detected in *P. lividus* and *M. galloprovincialis* larvae exposed to HHCB and AHTN compared to the control (Figure 2D and E). The effects of HHCB on sea urchin larvae development was significantly ($p < 0.01$) concentration dependent (Table S2 – Supplementary Materials) and significantly different ($p < 0.05$) compared to the control, with only 44.33% of larvae able to develop to pluteus stage after 48 h exposure to 5 $\mu\text{g/L}$ HHCB. The sensitivity of sea urchins to environmental chemicals is widely reported and significant evidence showed that they represent an important biomonitoring tool for ecosystems health. Similarly, HHCB and AHTN significantly ($p < 0.05$) affected the development of *M. galloprovincialis* larvae when compared to the control (Figure 2E). However, when considered in relation to the number of oocytes exposed to each contaminant, the effects were incipient with the highest percentage of deformed and undeveloped oocytes being 19.88% and 17.60% for HHCB and AHTN, respectively. Although *P. lividus* and *M. galloprovincialis* larvae development tests were similar, the effects recorded were more pronounced in sea urchin larvae than mussels.

Significant ($p < 0.01$) concentration dependent responses (Table S2 – Supplementary Materials) were observed in *S. aurata* mortality tests with HHCB and AHTN (Table S2 – Supplementary Materials). Notwithstanding, the percentage mortality of *S. aurata* exposed to both fragrances were below the toxic threshold, the highest effect being 13% for HHCB and AHTN at the highest concentration of 5 $\mu\text{g/L}$. Although this species and endpoints proved to be sensitive to other contaminants at low concentrations (Capolupo et al., 2018; Díaz-Garduño et al., 2016), the impacts of environmental concentrations of the tested fragrances in this study were low.

Analysing the RQs, HHCB represented potential high risk for the marine environment based on the EC_{50} estimated for *P. lividus* larvae development. *P. lividus* seems to be very sensitive to chemical exposure because previous studies of environmental contaminants in coastal waters have reported significant toxicity of industrial and domestic effluent (Díaz-Garduño et al., 2016), pharmaceutically active products (Aguirre-Martínez et al., 2015; Capolupo et al., 2018), UV-filters (Paredes et al., 2014) and organic pollutants (Bellas et al., 2005; Mariani et al., 2006) to sea urchin larvae development. The quantitative risk estimate of HHCB obtained in this study for *P. lividus* (1.18 - 56.53) was comparable to that of propanol (0.02 – 17.29), previously reported by Capolupo et al. (2018). We also found that both compounds posed high risk to *I. galbana* based on IC_{10} values recorded (Table 2). Microalgae have broadly been used in evaluation of ERA for other emerging contaminants. For example, *I. galbana* have been reported to be significantly affected by UV filters and pharmaceutical active ingredients (Aguirre-Martínez et al., 2015; Paredes et al., 2014). Fragrances have been demonstrated to pose high risk to microalgae in the marine environment, of which HHCB and AHTN were more toxic than musk xylene and musk ketone (Díaz-Garduño et al., 2017). The adversity of measured environmental concentrations of HHCB and AHTN to *I. galbana* deserve attention because microalgae occupy the lowest trophic level of the marine food chain and, therefore, serve as food for higher trophic organisms, supporting a balance in the ecosystem. Again, HHCB and AHTN inhibition of microalgae growth represents a bottom-up effect that might result in structural and functional disruption of the ecosystems (Fleege et al., 2003).

Due to the instability of these substances under laboratory exposure, it is difficult to estimate the exact concentration causing toxic effect. Although we could not measure the concentrations of the exposure water and bioaccumulation because of practical constraints, studies have shown that after 3 h of exposure, over 30% was lost and this reduced to ca. 80% after 96 h (Breitholtz et al., 2003; Ehigues et al., 2019; Tumová et al., 2019; Wollenberger et al., 2003). Again, there are currently no techniques to measure the concentrations of these substances *in-situ*; therefore, there is a probability that reported environmental concentrations are a fraction of the actual concentration in the marine environments eliciting biological effects, given that some amount could be lost before or during sample analysis in the laboratory. As a result,

the present study should be considered relevant for marine protection goals due to the evidence of potential adversity of low concentrations of HHCB and AHTN, especially to *I. galbana* biomass and *P. lividus* larvae development.

5. Conclusion

This study evaluated the toxicity of comparable measured environmental concentrations of HHCB and AHTN to microalgae and early life stages of marine organisms, adding to the dearth of information regarding the adverse effects of fragrances in the marine environment. For all the species of microalgae, invertebrates and fish tested, differential sensitivity was observed. Although the effects of both fragrances were generally incipient, *P. lividus* larvae development was the most sensitive species/endpoint. From our data, the environmental risk of HHCB and AHTN was characterized as high for *P. lividus* larvae development and *I. galbana* growth, suggesting the necessity for further research into possible sublethal effects of both substances in the marine environment.

Acknowledgements

This research was supported by the Spanish Ministry of Economy and Competitiveness (Ref. CTM2012-37591), ERDF support from European Union and also partially funded by the Spanish Ministry of Science, Innovation and Universities through the CaPETHaS project (#CGL2017-92160-EXP). CVM Araújo is grateful for the Ramón y Cajal contract (RYC-2017-22324) from the Spanish Ministry of Science and Innovation.

Conflict of Interest

The authors declare that there is no conflict of interest for the publication of this present work and are willing to provide any clarification whenever called upon to do so.

References

- Aguirre-Martínez, G. V., Owuor, M.A., Garrido-Pérez, C., Salamanca, M.J., Del Valls, T.A., Martín-Díaz, M.L., 2015. Are standard tests sensitive enough to evaluate effects of human pharmaceuticals in aquatic biota? Facing changes in research approaches when performing risk assessment of drugs. *Chemosphere* 120, 75–85. <https://doi.org/10.1016/j.chemosphere.2014.05.087>
- Aguirre-Martínez, G. V., Delvalls, A.T., Martín-Díaz, M.L., 2015. Yes, Caffeine, ibuprofen, carbamazepine, novobiocin and tamoxifen have an effect on *Corbicula fluminea* (Müller, 1774). *Ecotoxicol. Environ. Saf.* 120, 142–154. <https://doi.org/10.1016/j.ecoenv.2015.05.036>
- ASTM International, 2012. Standard guide for conducting acute toxicity test starting with embryos of four species of saltwater bivalve molluscs. E724 – 98 98, 21. <https://doi.org/10.1520/E0724-98R12.2>
- Balk, F., Ford, R.A., 1999a. Environmental risk assessment for the polycyclic musks AHTN and HHCB in the EU. I. Fate and exposure assessment. *Toxicol. Lett.* 111, 57–79. [https://doi.org/10.1016/S0378-4274\(99\)00169-1](https://doi.org/10.1016/S0378-4274(99)00169-1)
- Balk, F., Ford, R.A., 1999b. Environmental risk assessment for the polycyclic musks, AHTN and HHCB II. Effect assessment and risk characterisation. *Toxicol. Lett.* 111, 81–94.
- Beiras, R., Fernández, N., Bellas, J., Besada, V., González-Quijano, A., Nunes, T., 2003. Integrative assessment of marine pollution in Galician estuaries using sediment chemistry, mussel bioaccumulation, and embryo-larval toxicity bioassays. *Chemosphere* 52, 1209–1224.

- [https://doi.org/10.1016/S0045-6535\(03\)00364-3](https://doi.org/10.1016/S0045-6535(03)00364-3)
- Bellas, J., Granmo, Å., Beiras, R., 2005. Embryotoxicity of the antifouling biocide zinc pyrithione to sea urchin (*Paracentrotus lividus*) and mussel (*Mytilus edulis*). Mar. Pollut. Bull. 50, 1382–1385. <https://doi.org/10.1016/j.marpolbul.2005.06.010>
- Bester, K., Hühnerfuss, H., Lange, W., Rimkus, G.G., Theobald, N., 1998. Results of non target screening of lipophilic organic pollutants in the german bight ii : polycyclic musk fragrances. Water Res. 32, 1857–1863.
- Breitholtz, M., Wollenberger, L., Dinan, L., 2003. Effects of four synthetic musks on the life cycle of the harpacticoid copepod *Nitocra spinipes*. Aquat. Toxicol. 63, 103–118.
- Capolupo, M., Díaz-Garduño, B., Martín-Díaz, M.L., 2018. The impact of propranolol, 17 α -ethinylestradiol, and gemfibrozil on early life stages of marine organisms: effects and risk assessment. Environ. Sci. Pollut. Res. 25, 32196–32209. <https://doi.org/10.1007/s11356-018-3185-6>
- Carlsson, G., Norrgren, L., 2004. Synthetic Musk Toxicity to Early Life Stages of Zebrafish (*Danio rerio*). Arch. Environ. Contam. Toxicol. 46, 102–105. <https://doi.org/10.1007/s00244-003-2288-2>
- Carlsson, G., Örn, S., Andersson, P.L., Söderström, H., Norrgren, L., 2000. The impact of musk ketone on reproduction in zebrafish (*Danio rerio*). Mar. Environ. Res. 50, 237–241.
- Cunha, S.C., Trabalón, L., Jacobs, S., Castro, M., Fernandez-Tejedor, M., Granby, K., Verbeke, W., Kwadijk, C., Ferrari, F., Robbens, J., Sioen, I., Pocurull, E., Marques, A., Fernandes, J.O., Domingo, J.L., 2018. UV-filters and musk fragrances in seafood commercialized in Europe Union: Occurrence, risk and exposure assessment. Environ. Res. 161, 399–408. <https://doi.org/10.1016/j.envres.2017.11.015>
- De Lange, H.J., Sala, S., Vighi, M., Faber, J.H., 2010. Ecological vulnerability in risk assessment - A review and perspectives. Sci. Total Environ. 408, 3871–3879. <https://doi.org/10.1016/j.scitotenv.2009.11.009>
- Díaz-Garduño, B., Pintado-Herrera, M.G., Biel-Maeso, M., Rueda-Márquez, J.J., Lara-Martín, P.A., Perales, J.A., Manzano, M.A., Garrido-Pérez, C., Martín-Díaz, M.L., 2017. Environmental risk assessment of effluents as a whole emerging contaminant: Efficiency of alternative tertiary treatments for wastewater depuration. Water Res. 119, 136–149. <https://doi.org/10.1016/j.watres.2017.04.021>
- Díaz-Garduño, B., Rueda-Márquez, J.J., Manzano, M.A., Garrido-Pérez, C., Martín-Díaz, M.L., 2016. Are combined AOPs effective for toxicity reduction in receiving marine environment? Suitability of battery of bioassays for wastewater treatment plant (WWTP) effluent as an ecotoxicological assessment. Mar. Environ. Res. 114, 1–11. <https://doi.org/10.1016/j.marenvres.2015.12.011>
- Dietrich, D.R., Hitzfeld, B.C., 2004. Bioaccumulation and ecotoxicity of synthetic musks in the aquatic environment. Handb. Environ. Chem. 3, 233–244. <https://doi.org/10.1007/b14125>
- Ding, T., Li, W., Cai, M., Jia, X., Yang, M., Yang, B., Li, J., 2020. Algal toxicity, accumulation and metabolic pathways of galaxolide. J. Hazard. Mater. 384, 121360. <https://doi.org/10.1016/j.jhazmat.2019.121360>
- Dsikowitzky, L., Schwarzbauer, J., Littke, R., 2002. Distribution of polycyclic musks in water and particulate matter of the Lippe River (Germany). Org. Geochem. 33, 1747–1758.
- Ehiguese, F.O., Fernandez, M. del C.C., Lara-Martín, P.A., Martín-Díaz, M.L., Araújo, C.V.M., 2019. Avoidance behaviour of the shrimp *Palaemon varians* regarding a contaminant gradient of galaxolide and tonalide in seawater. Chemosphere 232, 113–120. <https://doi.org/10.1016/j.chemosphere.2019.05.196>
- European Comission Joint Research Centre, 2003. Technical guidance document on risk assessment in support of Commission Directive 93/67/ EEC on risk assessment for new notified substances and Commission Regulation (EC) No. 1488/94 on risk assessment for existing substances. Part II. EUR 20418 EN/2. Eur. Chem. Bur. Part II, 7–179.
- European Commission, 2008a. European Union Risk Assessment Report on AHTN. Rep. AHTN. <https://doi.org/10.2788/40195>
- European Commission, 2008b. European Union Risk Assessment Report on HHCB. Rep. HHCB.

- <https://doi.org/10.2788/40195>
- Ferreira, C.S.G., Nunes, B.A., Henriques-Almeida, J.M. de M., Guilhermino, L., 2007. Acute toxicity of oxytetracycline and florfenicol to the microalgae *Tetraselmis chuii* and to the crustacean *Artemia parthenogenetica*. *Ecotoxicol. Environ. Saf.* 67, 452–458.
<https://doi.org/10.1016/j.ecoenv.2006.10.006>
- Fleeger, J.W., Carman, K.R., Nisbet, R.M., 2003. Indirect effects of contaminants in aquatic ecosystems. *Sci. Total Environ.* 317, 207–233. [https://doi.org/10.1016/S0048-9697\(03\)00141-4](https://doi.org/10.1016/S0048-9697(03)00141-4)
- Fromme, H., Otto, T., Pilz, K., 2001. Polycyclic musk fragrances in different environmental compartments in Berlin. *Water Res.* 35, 121–128.
- Gao, Y., Ji, Y., Li, G., Mai, B., An, T., 2016a. Bioaccumulation and ecotoxicity increase during indirect photochemical transformation of polycyclic musk tonalide: A modeling study. *Water Res.* 105, 47–55. <https://doi.org/10.1016/j.watres.2016.08.055>
- Gao, Y., Ji, Y., Li, G., Mai, B., An, T., 2016b. Bioaccumulation and ecotoxicity increase during indirect photochemical transformation of polycyclic musk tonalide: A modeling study. *Water Res.* 105, 47–55. <https://doi.org/10.1016/j.watres.2016.08.055>
- Gatermann, R., Biselli, S., Hu, H., Rimkus, G.G., Hecker, M., Karbe, L., 2002. Synthetic musks in the environment. Part I: Species-dependent bioaccumulation of polycyclic and nitro musk fragrances in freshwater fish and mussels. *Arch. Environ. Contam. Toxicol.* 42, 437–446.
<https://doi.org/10.1007/s00244-001-0041-2>
- Gooding, M.P., Newton, T.J., Bartsch, M.R., Hornbuckle, K.C., 2006. Toxicity of synthetic musks to early life stages of the freshwater mussel *Lampsilis cardium*. *Arch. Environ. Contam. Toxicol.* 51, 549–558. <https://doi.org/10.1007/s00244-005-0223-4>
- Hadjispyrou, S., Kungolos, A., Anagnostopoulos, A., 2001. Toxicity, bioaccumulation, and interactive effects of organotin, cadmium, and chromium on *Artemia franciscana*. *Ecotoxicol. Environ. Saf.* 49, 179–186. <https://doi.org/10.1006/eesa.2001.2059>
- Heberer, T., 2002. Occurrence, fate, and assessment of polycyclic musk residues in the aquatic environment of urban areas – A review. *Acta Hydrochim. Hydrobiol.* 30, 227–243.
- Human & Environmental Risk Assessment on ingredients of Household Cleaning Products (HERA), 2004. Polycyclic musks AHTN (CAS 1506-02-1) and HHCB (CAS 1222-05-05). Scenario.
- Hussain, Y.H., Guasto, J.S., Zimmer, R.K., Stocker, R., Riffell, J.A., 2016. Sperm chemotaxis promotes individual fertilization success in sea urchins. *J. Exp. Biol.* 219, 1458–1466.
<https://doi.org/10.1242/jeb.134924>
- Johnston, E.L., Roberts, D.A., 2009. Contaminants reduce the richness and evenness of marine communities: A review and meta-analysis. *Environ. Pollut.* 157, 1745–1752.
<https://doi.org/10.1016/j.envpol.2009.02.017>
- Kannan, K., Reiner, J.L., Se, H.Y., Perrotta, E.E., Tao, L., Johnson-Restrepo, B., Rodan, B.D., 2005. Polycyclic musk compounds in higher trophic level aquatic organisms and humans from the United States. *Chemosphere* 61, 693–700. <https://doi.org/10.1016/j.chemosphere.2005.03.041>
- Latała, A., Marcin, N., Stepnowski, P., 2009. Acute toxicity assessment of perfluorinated carboxylic acids towards the Baltic microalgae 28, 167–171. <https://doi.org/10.1016/j.etap.2009.03.010>
- Liu, S., Zhou, Q., Wang, Y., 2011. Ecotoxicological responses of the earthworm *Eisenia fetida* exposed to soil contaminated with HHCB. *Chemosphere* 83, 1080–1086.
<https://doi.org/10.1016/j.chemosphere.2011.01.046>
- Losso, C., Picone, M., Novelli, A.A., Delaney, E., Ghetti, P.F., Ghirardini, A.V., 2007. Developing toxicity scores for embryotoxicity tests on elutriates with the sea urchin *Paracentrotus lividus*, the oyster *Crassostrea gigas*, and the mussel *Mytilus galloprovincialis*. *Arch. Environ. Contam. Toxicol.* 53, 220–226. <https://doi.org/10.1007/s00244-006-0136-x>
- Luckenbach, T., Epel, D., 2005. Nitromusk and polycyclic musk compounds as long-term inhibitors of cellular xenobiotic defense systems mediated by multidrug transporters. *Environ. Health Perspect.* 113, 17–24. <https://doi.org/10.1289/ehp.7301>
- Mariani, L., Pascale, D. De, Faraponova, O., Tornambe, A., Sarni, A., Giuliani, S., Ruggiero, G., Onorati,

- F., Magaletti, E., 2006. The Use of a Test Battery in Marine Ecotoxicology : The Acute Toxicity of Sodium Dodecyl Sulfate 373–379. <https://doi.org/10.1002/tox>
- Meriç, S., Nicola, E. De, Iaccarino, M., Gallo, M., Di, A., Morrone, G., Warnau, M., Belgiorno, V., Pagano, G., 2005. Toxicity of leather tanning wastewater effluents in sea urchin early development and in marine microalgae. *Chem. Biodivers.* 61, 208–217. <https://doi.org/10.1016/j.chemosphere.2005.02.037>
- Moon, H.B., Lee, D.H., Lee, Y.S., Kannan, K., 2012. Occurrence and accumulation patterns of polycyclic aromatic hydrocarbons and synthetic musk compounds in adipose tissues of Korean females. *Chemosphere* 86, 485–490. <https://doi.org/10.1016/j.chemosphere.2011.10.008>
- Moreno-Garrido, I., Campana, O., Lubián, L.M., Blasco, J., 2005. Calcium alginate immobilized marine microalgae: Experiments on growth and short-term heavy metal accumulation. *Mar. Pollut. Bull.* 51, 823–829. <https://doi.org/10.1016/j.marpolbul.2005.06.008>
- Nakata, H., Sasaki, H., Takemura, A., Yoshioka, M., Tanabe, S., Kannan, K., 2007. Bioaccumulation, temporal trend, and geographical distribution of synthetic musks in the marine environment. *Environ.Sci.Technol.* 41, 2216–2222.
- OECD, 2013. Test No. 210: Fish, Early-Life Stage Toxicity Test. OECD Guidel. Test. Chem. Sect. 2, OECD Publ. Paris 1–24. <https://doi.org/10.1787/9789264070103-en>
- OECD, 2002. Proposal for updating guideline 201 freshwater alga and cyanobacteria, growth inhibition test. OECD Guidel. Test. Chem. 1–21.
- Okumura, Y., Koyama, J., Takaku, H., Satoh, H., 2001. Influence of organic solvents on the growth of marine microalgae. *Arch. Environ. Contam. Toxicol.* 128, 123–128. <https://doi.org/10.1007/s002440010229>
- OSPAR, 2004. Musk xylene and other musks, hazardous substances series.
- Paredes, E., Perez, S., Rodil, R., Quintana, J.B., Beiras, R., 2014. Ecotoxicological evaluation of four UV filters using marine organisms from different trophic levels *Isochrysis galbana*, *Mytilus galloprovincialis*, *Paracentrotus lividus*, and *Siriella armata*. *Chemosphere* 104, 44–50. <https://doi.org/10.1016/j.chemosphere.2013.10.053>
- Parolini, M., Magni, S., Traversi, I., Villa, S., Finizio, A., Binelli, A., 2015. Environmentally relevant concentrations of galaxolide (HHCB) and tonalide (AHTN) induced oxidative and genetic damage in *Dreissena polymorpha*. *J. Hazard. Mater.* 285, 1–10. <https://doi.org/10.1016/j.jhazmat.2014.11.037>
- Pedersen, S., Selck, H., Salvito, D., Forbes, V., 2009. Effects of the polycyclic musk HHCB on individual- and population-level endpoints in *Potamopyrgus antipodarum*. *Ecotoxicol. Environ. Saf.* 72, 1190–1199. <https://doi.org/10.1016/j.ecoenv.2008.10.012>
- Pintado-Herrera, M.G., González-Mazo, E., Lara-Martín, P.A., 2013. Environmentally friendly analysis of emerging contaminants by pressurized hot water extraction-stir bar sorptive extraction-derivatization and gas chromatography-mass spectrometry. *Anal. Bioanal. Chem.* 405, 401–411. <https://doi.org/10.1007/s00216-012-6453-1>
- Rüedel, H., Böehmer, W., Schröeter-Kermani, C., 2006. Retrospective monitoring of synthetic musk compounds in aquatic biota from German rivers and coastal areas. *J. Environ. Monit.* 8, 812–823. <https://doi.org/10.1039/b602389b>
- Schreurs, R.H.M.M., Quaedackers, M.E., Seinen, W., Van Der Burg, B., 2002. Transcriptional activation of estrogen receptor ER α and ER β by polycyclic musks is cell type dependent. *Toxicol. Appl. Pharmacol.* 183, 1–9. <https://doi.org/10.1006/taap.2002.9458>
- Sibila, M.A., Garrido, M.C., Perales, J.A., Quiroga, J.M., 2008. Ecotoxicity and biodegradability of an alkyl ethoxysulphate surfactant in coastal waters. *Sci. Total Environ.* 394, 265–274. <https://doi.org/10.1016/j.scitotenv.2008.01.043>
- Składanowski, A.C., Stepnowski, P., Kleszczy, K., Dmochowska, B., 2005. AMP deaminase in vitro inhibition by xenobiotics A potential molecular method for risk assessment of synthetic nitro- and polycyclic musks, imidazolium ionic liquids and N -glucopyranosyl ammonium salts. *Environ. Toxicol. Pharmacol.* 19, 291–296. <https://doi.org/10.1016/j.etap.2004.08.005>

- Sumner, N.R., Guitart, C., Fuentes, G., Readman, J.W., 2010. Inputs and distributions of synthetic musk fragrances in an estuarine and coastal environment ; a case study. *Environ. Pollut.* 158, 215–222. <https://doi.org/10.1016/j.envpol.2009.07.018>
- Tumová, J., Šauer, P., Golovko, O., Koba Učun, O., Grabic, R., Máchová, J., Kocour Kroupová, H., 2019. Effect of polycyclic musk compounds on aquatic organisms: A critical literature review supplemented by own data. *Sci. Total Environ.* 651, 2235–2246. <https://doi.org/10.1016/j.scitotenv.2018.10.028>
- US EPA, 2002. EPA/812/R-02-012: Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms.
- US EPA, 1992. Guidelines for Exposure Assessment. *Risk Assess. Forum* 57, 22888–22938. <https://doi.org/EPA/600/Z-92/001>
- Van Den Brink, P.J., 2008. Ecological risk assessment: From book-keeping to chemical stress ecology. *Environ. Sci. Technol.* 42, 8999–9004. <https://doi.org/10.1021/es801991c>
- Vandermeersch, G., Lourenço, H.M., Alvarez-Muñoz, D., Cunha, S., Diogène, J., Cano-Sancho, G., Sloth, J.J., Kwadijk, C., Barcelo, D., Allegaert, W., Bekaert, K., Fernandes, J.O., Marques, A., Robbens, J., 2015. Environmental contaminants of emerging concern in seafood - European database on contaminant levels. *Environ. Res.* 143, 29–45. <https://doi.org/10.1016/j.envres.2015.06.011>
- Villa, S., Assi, L., Ippolito, A., Bonfanti, P., Finizio, A., 2012. First evidences of the occurrence of polycyclic synthetic musk fragrances in surface water systems in Italy: Spatial and temporal trends in the Molgora River. *Sci. Total Environ.* 416, 137–141. <https://doi.org/10.1016/j.scitotenv.2011.11.027>
- Wollenberger, L., Breitholtz, M., Ole, K., Bengtsson, B., 2003. Inhibition of larval development of the marine copepod *Acartia tonsa* by four synthetic musk substances. *Sci. Total Environ.* 305, 53–64.
- Yamauchi, R., Ishibashi, H., Hirano, M., Mori, T., 2008. Effects of synthetic polycyclic musks on estrogen receptor, vitellogenin, pregnane X receptor, and cytochrome P450 3A gene expression in the livers of male medaka (*Oryzias latipes*). *Aquat. Toxicol.* 90, 261–268. <https://doi.org/10.1016/j.aquatox.2008.09.007>
- Yu, Y., Kong, F., Wang, M., Qian, L., Shi, X., 2007. Determination of short-term copper toxicity in a multispecies microalgal population using flow cytometry. *Ecotoxicol. Environ. Saf.* 66, 49–56. <https://doi.org/10.1016/j.ecoenv.2005.10.014>