

Surgical Mask use in Physical Exercise in young healthy subjects

SUME Trial

Café, HM^{1,3} MD; Leitão, M¹ CT; Freitas, A² RN; Marreiros, A³ PhD

1- Hospital Particular do Algarve, Gambelas, Portugal

2 - Faro Health Center

3 - Medicine and Biomedical Sciences Faculty, University of Algarve, Portugal

Corresponding author: Hugo Mendonça Café, hfcafe@ualg.pt

Abstract

In the context of the COVID-19 Pandemic, the use of surgical masks has become the new normal. The use of these devices in exercise and medical situations has been advocated with the purpose of reducing contagions, but some concerns exist regarding its safety. We performed maximal treadmill stress tests in 12 healthy young subjects, with and without surgical mask use, and measured exercise capacity, oxygen saturation (rest, peak exercise and post-exercise) and electrocardiographic changes. Exercise capacity and Oxygen saturation levels decreased in peak exercise vs rest in a statistically significant manner when mask was used. ECG changes, although not significant, were present in 3 subjects when mask was used and disappeared when the test was made unmasked. We conclude that masked exercise has the potential to cause decreased exercise load and oxygen saturation and potentially cause diagnostic errors in medical exams.

KeyWords: Surgical mask, exercise, treadmill test, stress test, Oxygen saturation, Covid19, ECG

Introduction

In the present SARS-COV-2 pandemic, one of the measures that has been progressively implemented by the various nations is related to the mandatory use of a mask in order to reduce the spread of the pathology^{1,2}. The locations in which the use of a mask is mandatory vary, with some nations having mandatory use in any public space and others only in non-ventilated places³

An issue that proves to be important has to do with physical exercise. Given that exercise has the potential to generate high levels of aerosols, it has been advocated that a use of a mask should be kept during all activity in order to potentially reduce the risk of viral transmission⁴. The effect of such mask in physical activity, however, is not clear. A study by Fikenzer *et al*⁵ showed significant negative changes in ventilation parameters in an cycle ergometer test in healthy subjects wearing surgical and FFP2 masks. During our study, a paper published by Shaw et Al⁶ showed that time to exhaustion during a cycle ergometer test was not different with or without surgical mask use, neither was arterial oxygenation.

On the other hand, in the context of the COVID-19 Pandemic, specific exams in Cardiology such as the Electrocardiogram (ECG), stress Test (treadmill or cycle ergometer), are sometimes done with masked patients in order to avoid possible viral spread, given that this test produces significant amounts of aerosols. We do not know, however, whether the use of a mask causes significant changes in cardiovascular parameters in order to interfere in the interpretation of the exam.

We, therefore, aimed to evaluating the impact of the use of a standard surgical mask in a Treadmill stress test, in terms of ECG changes, oxygen saturation and total exercise capacity.

Methodology

The study protocol was presented to the Ethics Commission of Hospital Particular do Algarve, that approved its methodology and objectives and confirmed that the study was in accordance with the Declaration of Helsinki. The approval has the number 06-2020 and was finalized in July 27, 2020.

12 self reported healthy volunteers (medical students and nurses), 4 male and 8 female, with average characteristics presented in table 1, were asked to perform a Treadmill Stress Test (Bruce protocol) while wearing a standard surgical mask in which total physical capacity, ECG changes and oxygen saturation were evaluated. The tests were stopped after the subjects considered that they had reached exhaustion and therefore requested termination of the exam. Subjects were asked for a personal opinion if they believed the use of surgical mask interfered with their performance. Two weeks after the first exam, a second one was done, this time without surgical mask (two days before they were required to have a negative SARS-COV-2 test).

The Treadmill test was performed in a BTL-770M device, with Blood Pressure measurements done with a Heine Gamma XXL LF device. Oxygen Saturation was evaluated by pulse oximetry with a Sangool FS20D device. A Cardiologist and a Cardiology Technician were present at all times during the tests.

In terms of ECG changes, they were interpreted by a Cardiologist in accordance with ACC guidelines⁷. Arterial saturation was measured directly in one of the fingers of the subject before exercise, at peak exercise and post-exercise (5 minutes after the end of the exam), and total physical capacity estimated from the total time of exercise. Since different stages are involved in Bruce Protocol, we created a MET-Score in which the result was obtained by multiplying the seconds of exercise done by the METs associated with the stage in which the exercise was done, then we made the sum of all stages, therefore corresponding to an estimate of the full load of exercise performed. This was done because there is a clear difference in exercise load by doing 10 seconds of exercise in *e.g.* stage 2 or in stage 5.

All data was delivered to a data manager for post-processing.

All subjects received a document explaining the study and signed a consent form authorizing use of their data and agreeing to participate in the study.

In order to maintain consistency, we provided surgical masks to all patients. The mask was an OP-MASK B, a type IIR medical device.

	Male	Female	Total	p-Value
n	4	8	12	-
Age (mean \pm SD) (median \pm IQR)	30.5 \pm 5.69 30.50 \pm 11.00	27.3 \pm 3.80 26.50 \pm 7.00	28.3 \pm 4.60 27.5 \pm 8.00	.283
BMI (mean \pm SD) (median \pm IQR)	24.9 \pm 3.22 24.72 \pm 5.96	22.0 \pm 1.95 21.91 \pm 3.91	23.0 \pm 2.70 22.67 \pm 3.50	.214
Weight (mean \pm SD) (median \pm IQR)	78.0 \pm 7.44 79.50 \pm 14.00	57.6 \pm 5.00 58.00 \pm 9.50	64.4 \pm 11.50 62.50 \pm 20.25	.004 < .05
Height (mean \pm SD) (median \pm IQR)	1.8 \pm 0.10 174.50 \pm 17.25	1.62 \pm 0.02 162.00 \pm 4.50	1.67 \pm 0.09 164.00 \pm 11.50	.004 < .05
BSA (mean \pm SD) (median \pm IQR)	1.10\pm0.12 1.96\pm0.23	1.6\pm0.07 1.62\pm0.14	1.7\pm0.19 1.67\pm0.30	.004 < .05

SD- Standard Deviation

IQR - InterQuartilic Range

Table 1 – General characteristics of the sample

Statistical Analysis

We performed a statistical analysis using inter-quartilic range , mean, standard deviation and median. Hypotheses tests were executed for independent samples - Mann-Whitney U test - and for paired samples - Wilcoxon signed rank test and Chi squared test to test association between categorical variables

IBM SPSS version 27 software was used for the statistical analysis.

Results

Results are presented in table 2

	Mask	No Mask	p-Value
MET-SCORE (mean \pm SD) (median \pm IQR)	5831.92 \pm2041.38 5825.50\pm2972.00	6570.58\pm2787.53 6059.50\pm4580.25	.023 < .05
Basal SAT (mean \pm SD) (median \pm IQR)	98.67 \pm 1.67 99.00 \pm 2.00	98.00 \pm 1.954 98.50 \pm 1.750	.062
Peak SAT (mean \pm SD) (median \pm IQR)	95.92\pm2.81 96.00\pm4.00	97.50\pm2.468 98.50\pm3.000	.049 < .05
Post SAT (mean \pm SD) (median \pm IQR)	97.00 \pm 2.17 97.00 \pm 3.50	97.42 \pm 1.782 97.50 \pm 1.750	.357
ΔSAT1 (mean \pm SD) (median \pm IQR)	2.75\pm2.22 3.00\pm3.75	0.67\pm1.435 0.00\pm1.750	.021 < .05
Δ SAT2 (mean \pm SD) (median \pm IQR)	1.08 \pm 2.15 2.00 \pm 2.00	-0.08 \pm 1.165 0.00 \pm 2.000	.549
ECG changes (n.%) Yes No	3.25% 9.75%	0.0% 0.0%	.064
Discomfort (n,%) Yes No	6, 50% 6, 50%	0,0% 0,0%	.005

SD- Standard Deviation

IQR - InterQuartilic Range

Table 2 – Results

All but one of the subjects had a higher MET-Score while exercising without a mask. Comparing groups, there was a slight but statistically significant increase in MET-Score when subjects performed without a mask, showing that the subjects were able to increase exercise load when the mask was removed (figure 1).




Figure 1 - Increased MET-Score in No Mask group. $p = .023$

Half of the participants declared feeling a sensation of discomfort while wearing the mask. This was also a statistically significant result.

Saturation levels were evaluated in rest, peak exercise and post-exercise (5 minutes after stopping the test). We compared basal and peak levels in one variable (ΔSAT1 , difference between basal saturation and peak exercise saturation) and peak and post-exercise levels in another (ΔSAT2 , difference between peak exercise saturation and post-exercise saturation). We found that the decrease of ΔSAT1 was more apparent in the masked group, and this finding had statistical significance (figure 2). This was closely related with a statistically significant difference in peak exercise saturation (figure 3). ΔSAT2 was not statistically significant.




Figure 2 - Clear difference between Δ SAT1 in both groups (lower the better). $p = .021$




Figure 3 - Lower saturations observed in peak exercise when the subjects wore mask (higher the better). $p = .049$

ECG changes were evaluated by a trained cardiologist. In the masked group 3 subjects, all female, had ECG changes that albeit not diagnostic for coronary ischemia, were at least doubtful. Two subjects had slight ST depression (one in inferior leads, the other in V4-V6 leads, neither reaching 1mm depression but over 0.5mm) and the other had premature ventricular beats (figures 4, 5 and 6). These changes were not the typical "upslope ST segment" expected in exercise, but a more horizontal change with less than 10mm/second upstroke. Also relevant, none of these changes appeared in peak exercise (two in recovery, another at medium level of effort). When the subjects repeated the test unmasked, these alterations were no longer present.




Figure 4 - Premature ventricular beats in recovery phase (masked group)




Figure 5 - Slight ST depression V4-V6 leads during exercise (masked group)




Figure 6 - Slight ST depression in Inferior leads, recovery phase (masked group)

Discussion

The current study provides valuable insights regarding use of surgical masks in exercise. In our study subjects were not required to control their average physical activity, to abstain from exercise in specific periods or to control their eating habits. This had the purpose of not interfering with the daily routines of our subjects and, since they were acting as their own control, obtain a result with higher external validity. The study by Fikenzer *et al*⁵ did showed changes in respiratory parameters like forced expiratory volume and peak expiratory flow, but neither oxygen saturation nor physical activity performed was evaluated. Shaw *et al*⁶ however, did test oxygen saturation levels between rest and peak exercise and total physical activity performed. In this study neither of these parameters was statistically significant. In our study this was not the case. Both the physical load performed (estimated by our MET-Score) and Oxygen saturation (O2Sat) has clear differences when subjects wore masks. Physical capacity and O2Sat at peak exercise were decreased when subjects wore a mask compared to not wearing one, showing that indeed the use of mask can cause measurable respiratory and exercise capacity changes even in healthy volunteers doing

exercise. When we compare our data with the data from Shaw *et al*⁶, it is possible that the fact that we used a treadmill test and not a cycle ergometer might have affected the results regarding O2Sat. Since higher muscle mass is used in a treadmill than in a cycle ergometer⁸, this could theoretically cause a higher oxygen demand. The notion is not without its merits, if we consider that at least for patients with chronic obstructive pulmonary disease a treadmill test has been described as eliciting lower O2Sat levels at peak exercise *vs* a cycle ergometer⁹. It has been shown that the Maximal Oxygen consumption (VO_2 max) is higher in treadmill stress tests compared with cycle ergometer stress tests in trained individuals¹⁰, which in theory would convey a higher aerobic capacity and therefore a possible higher exercise capacity in that time of activity. Although our methods are different, we clearly had a difference with the use of a mask. We preferred a treadmill because it's more commonly used in our country and we believe replicates more accurately the type of leisure exercise most healthy subjects do, so we believe our results can be more relevant to the general population, notwithstanding the small numbers used.

A quite surprising result obtained, although not reaching statistical significance, was the appearance of non diagnostic but doubtful ECG changes in subjects while doing the test with a mask that disappeared when the test was repeated without mask. Since all subjects were healthy, asymptomatic, and the unmasked exam was done two weeks later, without any medication or change in habits, we do not believe this was related to ischemia. While we could speculate that an unusually high change in O2Sat between rest and peak could be that cause of this, that did not happen in our subjects, since the changes were slight. Also, it is questionable that such hypoxic effect could cause such changes unless it's quite significant^{11,12}. These subjects will perform further evaluations to rule-out the possibility of other alterations.

Since the beginning of this pandemic period, it has been regular practice to perform treadmill tests for diagnosis of coronary artery disease with the patients wearing masks. Considering our findings regarding the ECG changes, the possibility of diagnostic interference with mask use exists and has to be taken into account, and cardiologists should be aware of this while interpreting studies.

Another relevant issue is, since we did detect significant changes in exercise capacity,

oxygen saturation and reported discomfort with mask use, if this might impact the amount and level of exercise performed by the everyday subject if a mask is mandatory in these situations: While for the trained athlete the decrease in exercise load is the main issue (this might be relevant also for the casual subject that does leisure physical activity, since it might demoralize someone that's starting to exercise), the discomfort and the decrease in O2Sat might be much more significant for the casual person, and might cause an increased abandonment of exercise.

From our results, if viable from the epidemiological perspective, exercise should be done without a mask, and consideration for exercise stress tests without masks has to be considered.

Conclusion:

Our study shows that young healthy subjects have significant decreases in exercise capacity and oxygen saturation when performing exercise using surgical masks, besides reporting discomfort with its use. Also, there were slight ECG changes only present with mask use, which could cause difficulties in diagnosis of ischemia. A study with a higher number of subjects might clarify these findings, but we believe exercise should be done without a mask and consideration for exercise stress tests without masks should be discussed in the Medical Community.

Author contribution:

Café, HM is a Medical Doctor, Cardiologist and Professor of Medicine. He designed the study and provided cardiological support and interpretation of clinical results

Leitão, M is a Cardiology Technician. She provided the technical support for the realization of the stress tests

Freitas, A is a Registered Nurse. She was responsible for data managing, insight and planning and proper methodology delineation

Marreiros, A is a Professor of Medicine, Vice-Dean of the Medical Degree in University of Algarve and PhD in Applied Mathematics. She was responsible for the statistical analysis of the study

Bibliography

1. CDC (2020) Recommendation regarding the use of cloth face coverings, especially in areas of significant community-based transmission
2. ECDC (2020) Using face masks in the community—reducing COVID-19 transmission from potentially asymptomatic or presymptomatic people through the use of face masks
3. Greenhalgh T. Face coverings for the public: Laying straw men to rest. *J Eval Clin Pract.* 2020.
4. Chandrasekaran, B.; Fernandes, S. “Exercise with Facemask; Are We Handling a devil’s sword?”—A Physiological Hypothesis. *Med. Hypothesis* 2020, 144, 110002
5. Fikenzer S, Uhe T, Lavall D, et al. Effects of surgical and FFP2/N95 face masks on cardiopulmonary exercise capacity. *Clin Res Cardiol.* 2020;109(12):1522-1530.
6. Shaw K, Butcher S, Ko J, Zello GA, Chilibeck PD. Wearing of Cloth or Disposable Surgical Face Masks has no Effect on Vigorous Exercise Performance in Healthy Individuals. *Int J Environ Res Public Health.* 2020;17(21):8110. Published 2020 Nov 3. doi:10.3390/ijerph17218110
7. Gibbons RJ, Balady GJ, Beasley JW, Bricker JT, Duvernoy WF, Froelicher VF, Mark DB, Marwick TH, McCallister BD, Thompson PD, Winters WL Jr, Yanowitz FG, Ritchie JL, Cheitlin MD, Eagle KA, Gardner TJ, Garson A Jr, Lewis RP, O'Rourke RA, Ryan TJ. ACC/AHA guidelines for exercise testing: executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines

(Committee on Exercise Testing). *Circulation*. 1997 Jul 1;96(1):345-54.

8. Kisan R At Al. Treadmill and Bicycle Ergometer Exercise: Cardiovascular Response comparison; *Global Journal of Medical research* 2012, 12 (5)

9. Hsia D, Casaburi R, Pradhan A, Torres E, Porszasz J. Physiological responses to linear treadmill and cycle ergometer exercise in COPD. *Eur Respir J*. 2009 Sep;34(3):605-15

10. Basset FA, Boulay MR. Specificity of treadmill and cycle ergometer tests in triathletes, runners and cyclists. *Eur J Appl Physiol*. 2000

11. Entwistle MD, Sommerville D, Tandon AP, Jones JG. Effect of hypoxaemia on the resting electrocardiogram (ECG) in patients with cardiac ischaemia. *Ann Acad Med Singap*. 1994 Jul;23(4):460-4

12. Coustet B, Lhuissier FJ, Vincent R, Richalet JP. Electrocardiographic changes during exercise in acute hypoxia and susceptibility to severe high-altitude illnesses. *Circulation*. 2015 Mar 3;131(9):786-94