Type of the Paper (Article.)

A solution of the cosmological constant, using multiverse version of Penrose CCC cosmology, and Enhanced Quantization compared

Andrew Beckwith ¹

1 Physics Department, Chongqing University, College of Physics, Chongqing University Huxi Campus, No. 44 Daxuechen Nanlu, Shapinba District, Chongqing 401331, People's Republic of China, rwill9955b@gmail.com

Abstract:

We reduplicate the Book “Dark Energy” by M. Li, X-D. Li, and Y. Wang, zero-point energy calculation with an unexpected “length’ added to the ‘width’ of a graviton wavefunction just prior to the entrance of ‘gravitons’ to a small region of space-time prior to a nonsingular start to the universe. We compare this to a solution worked out using Klauder Enhanced quantization, for the same given problem. The solution of the first Cosmological Constant problem relies upon the geometry of the multiverse generalization of CCC cosmology which is explained in this paper. The second solution, used involves Klauder enhanced quantization. We look at energy given by our methods and compare and contrast it with the negative energy of the Rosen model for a mini sub universe and estimate GW frequencies

Keywords: Minimum scale factor, cosmological constant, space-time bubble, bouncing cosmologies

1. Introduction

We bring up this study due to the general failure of field theory methods to obtain a working solution to the Cosmological Constant problem. One of the reasons for this work is that Quintessence studies involving evolution of the Cosmological Constant over time have routinely failed in match ups with Early supernova data. The working hypothesis is that if Dark Energy is commensurate with the Cosmological Constant that the Cosmological Constant, and Dark energy would be the same from the start of inflation. The traditional model of inflation also involves due to the Penrose Singularity theorems [1] a singular point for initial expansion of the Universe. Our study clearly initiates a Cosmological constant which is invariant over time. We digress from using the Penrose Singularity theorem [1] in terms of a Cosmic bounce[2] style nonsingular initial bubble of space-time for reason we state in the manuscript. The importance we have to our work is that we are convinced that the Cosmological Constant [3][4] was set initially for conditions in the interior of a nucleated space-time bubble, with a wavefunction style calculation for collapsing infill into the space-time bubble with that wavefunction of Dark energy [5][6] to be at a minimum 10 to the 30 power times Planck length. We conclude that Polarization states for obtaining early universe high Frequency Gravitational waves detected and analyzed may be more favorable for higher dimensional generalizations of our Dark Energy generating procedures, with Klauder Enhanced Quantization[7],[8] limited to 3+1 Geometry whereas our Vacuum energy[9][10] calculation may be amendable to higher frequency calculation incantations. While considering [11],[12],[13],[14],[15],[16],[17],[18],[19] issues. In particular [19] gives us future goals to consider. As an example, could multiverse CCC allow incorporating [19] with [13]?

2. Methods

We will first start off with the redone calculation as to the Vacuum energy as given in [1] and how we rescale them to be in sync as to the observed experimental value for vacuum energy which is of the present era. This methodology is consistent with the Zero-point energy calculation, we start off with the following as given by [11]

$$\frac{1}{2} \cdot \sum_i \omega_i \equiv V(\text{volume}) \cdot \int_0^{\hat{\lambda}} \sqrt{k^2 + m^2} \frac{k^2 dk}{4\pi^2} \approx \frac{\hat{\lambda}^4}{16\pi^2} \quad (1)$$

$$\xrightarrow{\hat{\lambda}=M_{\text{Planck}}} \rho_{\text{boson}} \approx 2 \times 10^{71} \text{ GeV}^4 \approx 10^{119} \cdot \left(\rho_{DE} = \frac{\Lambda}{8\pi G} \right)$$

In stating this we have to consider that $\rho_{DE} = \frac{\Lambda}{8\pi G} \approx \hbar \cdot \frac{(2\pi)^4}{\lambda_{DE}^4}$, so then that

the equation we have to consider is a wavelength $\lambda_{DE} \approx 10^{30} \ell_{\text{Planck}}$ which is about 10^{30} times a Plank length radius of a space-time bubble [13] as a nonsingular expansion point for Cosmology, at the start of inflation with the space-time bubble of about a Plank length radius in size. . Having said that , how do we get having the Penrose multiverse condition in this problem, for

$\lambda_{DE} \approx 10^{30} \ell_{\text{Planck}}$ just before the near singularity?

2.1 Looking now at the Modification of the Penrose CCC (Cosmology)

We now outline the generalization for Penrose CCC(Cosmology) just before inflation which we state we are extending Penrose's suggestion of cyclic universes, black hole evaporation, and the embedding structure our universe is contained within, This multiverse has BHs and may resolve what appears to be an impossible dichotomy. The following is largely from [13] and has serious relevance to the final part of the conclusion .That there are N universes undergoing Penrose 'infinite expansion' (Penrose) [18] contained in a mega universe structure. Furthermore, each of the N universes has black hole evaporation, with Hawking radiation from decaying black holes. If each of the N universes is defined by a partition function, called $\{\Xi_i\}_{i=1}^N$, then there exist an information ensemble of mixed

minimum information correlated about $10^7 - 10^8$ bits of information per partition function in the set $\{\Xi_i\}_{i=1}^N$ _{before}, so minimum information is conserved between a set of partition functions per universe [13]

$$\left\{ \Xi_i \right\}_{i=1}^N \Big|_{\text{before}} \equiv \left\{ \Xi_i \right\}_{i=1}^N \Big|_{\text{after}} \quad (2)$$

However, there is non-uniqueness of information put into partition function $\{\Xi_i\}_{i=1}^N$. Also

Hawking radiation from black holes is collated via a strange attractor collection in the mega

universe structure to form a new inflationary regime for each of the N universes represented

Our idea is to use what is known as CCC cosmology[13],[18], which can be thought of as the following. First. Have a big bang (initial expansion) for the universe which is represented by $\{\Xi_i\}_{i=1}^i$. Verification of this mega structure compression and expansion of information with stated non-uniqueness of information placed in each of the N universes favors ergodic mixing of initial values for each of N universes expanding from a singularity beginning. The n_f stated

value, will be using (Ng, 2008) $S_{entropy} \sim n_f \cdot [20]$. How to tie in this energy expression, as in

Eq.(12) will be to look at the formation of a nontrivial gravitational measure as a new big bang for each of the N universes as by $n(E_i) \cdot$ the density of states at energy E_i for partition function[13],[14].

$$\{\Xi_i\}_{i=1}^i \propto \left\{ \int_0^\infty dE_i \cdot n(E_i) \cdot e^{-E_i} \right\}_{i=1}^i \quad (9)$$

Each of E_i identified with Eq.(9) above, are with the iteration for N universes [13],and [18](Penrose, 2006) Then the following holds, by asserting the following claim to the universe, as a mixed state, with black holes playing a major part, i.e.

CLAIM 1

See the below[13] representation of mixing for assorted N partition function per CCC cycle

$$\frac{1}{N} \cdot \sum_{j=1}^N \Xi_j \Big|_{j\text{-before-nucleation-regime}} \xrightarrow{\text{vacuum-nucleation-transfer}} \Xi_i \Big|_{i\text{-fixed-after-nucleation-regime}} \quad (10)$$

For N number of universes, with each $\Xi_j \Big|_{j\text{-before-nucleation-regime}}$ for j = 1 to N being the partition function of each universe just before the blend into the RHS of Eq. (10) above for our present universe. Also, each independent universes as given by $\Xi_j \Big|_{j\text{-before-nucleation-regime}}$ is constructed by the absorption of one to ten million black holes taking in energy. **I.e. (Penrose)** [18]. Furthermore, the main point is done in [13] in terms of general ergodic mixing [21][22][23]

Claim 2

$$\Xi_j \Big|_{j\text{-before-nucleation-regime}} \approx \sum_{k=1}^{Max} \tilde{\Xi}_k \Big|_{black\text{-holes-}jth\text{-universe}} \quad (5)$$

What is done in **Claim 1 and 2** [13] is to come up as to how a multi dimensional representation of black hole physics enables continual mixing of spacetime [13],[18],[22],[24] largely as a way to avoid the Anthropic principle[13][25], as to a preferred set of initial conditions

2.2 Looking at the Modification of the Penrose CCC (Cosmology)

We argue this modification is mandated by having the initial DE wavefunction set as

$$\lambda_{DE} \approx 10^{30} \ell_{Planck} \quad (6)$$

After having set this in place we will be comparing this method to the Klauder argument

2.3 Looking at Klauder Enhanced Quantization for Cosmological Constant:

We use the Padmanabhan 1st integral [26] of the form , with the third entry of Eq. (1) having a Ricci scalar defined via [27] [28]and usually the curvature $\mathfrak{K}$ [13] set as extremely small, with

general relativity[13][28] action of

$$\begin{aligned} S_1 &= \frac{1}{2\kappa} \int \sqrt{-g} \cdot d^4x \cdot (\mathfrak{R} - 2\Lambda) \\ \mathfrak{g} - g &= -\det g_{uv} \\ \mathfrak{g}\mathfrak{R} &= 6 \cdot \left(\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 + \frac{\mathfrak{K}}{a^2} \right) \end{aligned} \quad (7)$$

2.3. Next for the idea from Klauder

We are going to go to page 78 by Klauder [8][13] of what he calls on page 78 a restricted

Quantum action principle which he writes as: S_2 where we write a 1-1 equivalence as in

[8][13] , which is

$$S_2 = \int_0^T dt \cdot \left[p(t) \dot{q}(t) - H_N(p(t), q(t)) \right] \approx S_1 = \frac{1}{2\kappa} \int \sqrt{-g} \cdot d^4x \cdot (\mathfrak{R} - 2\Lambda) \quad (8)$$

Our assumption is that Λ is a constant[13], hence we assume then use the approximation ,

from [8], and [13]

$$\begin{aligned}
\frac{p_0^2}{2} &= \frac{p_0^2(N)}{2} + N; \text{ for } 0 < N \leq \infty, \text{ and } q = q_0 \pm p_0 t \\
V_N(x) &= 0; \text{ for } 0 < x < 1 \\
V_N(x) &= N; \text{ otherwise} \\
H_N(p(t), q(t)) &= \frac{p_0^2}{2} + \frac{(\hbar \cdot \pi)^2}{2} + N; \text{ for } 0 < N \leq \infty
\end{aligned} \tag{9}$$

Our innovation is to then equate $q = q_0 \pm p_0 t \sim \phi$ and to assume small time step values.

Then[13]

$$\Lambda \approx \frac{-\left[\frac{V_0}{3\gamma-1} + 2N + \frac{\gamma \cdot (3\gamma-1)}{8\pi G \cdot \tilde{t}^2}\right]}{\frac{1}{\kappa} \int \sqrt{-g} \cdot d^3x} + \left(6 \cdot \left(\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2\right)\right) \Bigg|_{t=\tilde{t}} \tag{10}$$

If we assume that Eq. (10) was formed within a cosmic bubble of space-time ‘[12]

$$\left(\frac{\dot{a}}{a}\right)^2 = H_{\text{bounce-cos}}^2 = \frac{8\pi}{3M_P^2} \cdot \left(\rho - \frac{\rho^2}{2|\sigma|}\right) \tag{11}$$

Here, we have that ρ is a space-time density function, whereas σ is the tension of a space-time bubble presumably of the order of a Planck radius.

Also within the bubble of space-time $\left(\frac{\ddot{a}}{a}\right) \approx \varepsilon^+ \simeq 0$ and $a(t) = a_{\min} t^\gamma$ at the surface of the space-time bubble. Hence

$$\Lambda \approx \frac{-\left[\frac{V_0}{3\gamma-1} + 2\tilde{N} + \frac{\gamma(3\gamma-1)}{8\pi G t_{\text{planck}}}\right]}{\frac{1}{\kappa} \int \sqrt{-g} d^3x} + c_1 \frac{16\pi}{M_P^2} \cdot \left(\rho - \frac{\rho^2}{2|\sigma|}\right) \tag{12}$$

c_1 is to be determined, whereas the inflaton[13],[26].

$$\phi = \sqrt{\frac{\gamma}{4\pi G}} \ln \left\{ \sqrt{\frac{8\pi G V_0}{\gamma(3\gamma-1)}} \cdot t \right\} \tag{13}$$

And what we will use the “inflaton potential” we write as[13],[26]

$$V = V_0 \cdot \left\{ \sqrt{\frac{8\pi G V_0}{\gamma(3\gamma-1)}} \cdot t \right\}^{\sqrt{\frac{\gamma}{4\pi G}} - \sqrt{\frac{8\pi G}{\gamma}}} \tag{14}$$

Also

$$\rho \approx \frac{\dot{\phi}^2}{2} + V(\phi) \equiv \frac{\gamma}{8\pi G} \cdot t^2 + V_0 \cdot \left\{ \sqrt{\frac{8\pi G V_0}{\gamma(3\gamma-1)}} \cdot t \right\}^{\sqrt{\frac{\gamma}{4\pi G}} - \sqrt{\frac{8\pi G}{\gamma}}} \quad (15)$$

3. Results: comparing Eq (1) implications with Eq (12)

The significance of the procedure is that with tweaking we may be seeing the actual realization of classical gravity as an Eikonal Approximation to quantum theory[29][30] as what was brought up by Horowitz and Oron in a highly nonstandard way as seen in reference [30] which gives a different interpretation to Eq. (15) above,. The term we refer to as Eikonal approximation to a quantum state has the form of the following decomposition. From Powell and Crasemann,[31] we have the decomposition of Geometric style decomposition of the optical wave equation of

$$\nabla^2 \psi - \frac{1}{v_{\text{Velocity}}^2} \cdot \frac{d^2 \psi}{dt^2} = 0 \quad (16)$$

Whereas we use the common $\lambda \cdot \omega = 2\pi v_{\text{Velocity}}$, and if we substitute in an angular frequency dependence of $e^{-i\omega t}$ in ψ , we get from (16) that if $\lambda = h/p$ and $p = \sqrt{2m \cdot (E - V(\text{potential}))}$, Then we have a transformation of Eq. (17) to Eq.(18)

$$\nabla^2 \psi + \frac{4\pi^2}{\lambda^2} \cdot \psi = 0 \quad (17)$$

Leads to a Schrodinger equation of the form given by

$$\nabla^2 \psi + \frac{8\pi^2}{h^2} \cdot (E - V(\text{potential})) \cdot \psi = 0 \quad (18)$$

Eq. (17) has[32]

$$\psi = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} (a_{\ell m} j_{\ell}(kr) + b_{\ell m} y_{\ell}(kr)) \cdot Y_{\ell}^m(\theta, \phi) \quad (19)$$

Where we could get away with making a substitution of[31]

$$k = \frac{2\pi}{\lambda} = \frac{2\pi}{h} \cdot (E - V(\text{potential}))^{1/2} \quad (20)$$

Where the first and second Bessel Equation solutions[32] are of the form

$$j_{\ell}(kr = x) = (-x)^{\ell} \left(\frac{1}{x} \frac{d}{dx} \right)^{\ell} \frac{\sin x}{x} \quad (21)$$

$$y_{\ell}(kr = x) = -(-x)^{\ell} \left(\frac{1}{x} \frac{d}{dx} \right)^{\ell} \frac{\cos x}{x} \quad (22)$$

Then we have via making use of Kieffer's result of variance of the energy [33] is

$$\Delta E = \frac{\sqrt{2\tilde{n}+1}}{\lambda} \approx \frac{\hbar\sqrt{2\tilde{n}+1} \cdot (E - V(\text{potential}))^{1/2}}{2\pi} \equiv \hbar\sqrt{2\tilde{n}+1} \cdot \sqrt{\frac{V_{\text{volume}}}{8\pi G}} \cdot t \quad (23)$$

We get via [33] on page 239 of this reference, Eq.(23). after we use (23), then

$$\Delta E \Delta t \approx \sqrt{2\tilde{n}+1} \cdot \sqrt{\frac{V_{\text{volume}}}{8\pi G}} \cdot t \cdot \Delta t \cdot \hbar \approx \hbar \quad (24)$$

If so, then we have that if so by use of [34] since this is also applicable to black holes

$$\begin{aligned} t \propto t_{\text{Planck}} \propto \Delta t \Rightarrow \Delta E \Delta t \approx \sqrt{2\tilde{n}+1} \cdot \sqrt{\frac{V_{\text{volume}}}{8\pi G}} \cdot (\Delta t)^2 \cdot \hbar \propto \hbar \\ \Rightarrow (\Delta t)^2 \propto \frac{1}{\sqrt{2\tilde{n}+1}} \cdot \sqrt{\frac{8\pi G}{V_{\text{volume}}}} \end{aligned} \quad (25)$$

Using Planck units, what this means is that if $G = \hbar = \ell_{\text{Planck}} = k_B = 1 = \text{Planck time}$

$$(\Delta t)^2 \propto \frac{1}{\sqrt{2\tilde{n}+1}} \cdot \sqrt{\frac{8\pi G}{V_{\text{volume}}}} \equiv \sqrt{\frac{8\pi}{2\tilde{n}+1}} \cdot (t_{\text{Planck}})^2 \quad (26)$$

If $\tilde{n} = \text{quantum-number}$, then for Δt being proportional to $t_{\text{Planck}} \propto \Delta t$, we have definite restrictions on $\tilde{n} = \text{quantum-number}$, i.e. of the type given by

$$\sqrt{\frac{8\pi}{2\tilde{n}+1}} \leq 1 \quad (27)$$

Whereas the Kieffer Wavefunction commensurate as to Eq. (23) for a quantum Dust universe initially is given as [33] on page 239 of this reference

$$\psi_{\tilde{n},\lambda}(t,r) \equiv \frac{1}{\sqrt{2\pi}} \cdot \frac{\tilde{n}! (2\lambda)^{\tilde{n}+1/2}}{\sqrt{(2\tilde{n})!}} \cdot \left[\frac{1}{(\lambda + i \cdot t + i \cdot r)^{\tilde{n}+1}} - \frac{1}{(\lambda + i \cdot t - i \cdot r)^{\tilde{n}+1}} \right] \quad (28)$$

Here the time t would be proportional to Planck time, and r would be proportional to Planck length, whereas we set

$$\lambda \approx \sqrt{\frac{8\pi G}{V_{\text{volume}} \hbar^2 t^2}} \xrightarrow{G=\hbar=\ell_{\text{Planck}}=k_B=1} \sqrt{\frac{8\pi}{t^2}} \equiv \frac{\sqrt{8\pi}}{t} \quad (29)$$

Then a preliminary emergent space-time wavefunction would take the form of

$$\begin{aligned} \psi_{\tilde{n},\lambda}(\Delta t, r) \\ \equiv \frac{1}{\sqrt{2\pi}} \cdot \frac{\tilde{n}! \left(2 \cdot \sqrt{8\pi} \cdot (\Delta t)^{-1}\right)^{\tilde{n}+1/2}}{\sqrt{(2\tilde{n})!}} \cdot \left[\frac{1}{\left(\sqrt{8\pi} \cdot (\Delta t)^{-1} + i \cdot \Delta t + i \cdot r\right)^{\tilde{n}+1}} - \frac{1}{\left(\sqrt{8\pi} \cdot (\Delta t)^{-1} + i \cdot \Delta t - i \cdot r\right)^{\tilde{n}+1}} \right] \end{aligned} \quad (30)$$

Just at the surface of the bubble of space-time, with $t_{Planck} \propto \Delta t$, and $r \propto \ell_{Planck}$

This is from a section, page 239 of the 3rd edition of Kieffer's book, as to a quantum theory of collapsing dust shells, and is a way to use what is referred to as what Isham in 1984 referred to as "group quantization", as we need to remember that we are assuming $G = \hbar = \ell_{Planck} = k_B = t_{Planck} = 1$. Having said that, let us look at Eq.(1) and Eq. (12) again

3.1 We compare Eq. (1) and Eq. (12) with the above new information

$$\begin{aligned} \Lambda \approx \frac{-\left[\frac{V_0}{3\gamma-1} + 2\tilde{N} + \frac{\gamma(3\gamma-1)}{8\pi G t_{Planck}}\right]}{\frac{1}{\kappa} \int \sqrt{-g} d^3x} + c_1 \frac{16\pi}{M_P^2} \cdot \left(\rho - \frac{\rho^2}{2|\sigma|}\right) \\ \Leftrightarrow \rho_{DE} = \frac{\Lambda}{8\pi G} \Leftrightarrow \Lambda \approx \hbar \cdot 8\pi G \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30} \ell_{Planck}} \equiv 8\pi \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30}} \end{aligned} \quad (31)$$

To compare these two values we can state that within the bubble, that just before the bubble boundary, we have $\tilde{N} = 0$, hence we look at, initially

$$\begin{aligned} \Lambda \approx \frac{-\left[\frac{V_0}{3\gamma-1} + \frac{\gamma(3\gamma-1)}{8\pi G t_{Planck}}\right]}{\frac{1}{\kappa} \int \sqrt{-g} d^3x} + c_1 \frac{16\pi}{M_P^2} \cdot \left(\rho - \frac{\rho^2}{2|\sigma|}\right) \\ \Leftrightarrow \Lambda \approx \hbar \cdot 8\pi G \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30} \ell_{Planck}} \equiv 8\pi \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30}} \end{aligned} \quad (32)$$

i.e. we pick c_1 so that the two are equivalent in value,

Note this is just before a very large value set to $\tilde{N}$ at the boundary of the initial space-time bubble. Hence we also would be looking at

$$\begin{aligned}
\Lambda &\approx \gamma \cdot t^2 + 8\pi V_0 \cdot \left\{ \sqrt{\frac{8\pi V_0}{\gamma(3\gamma-1)}} \cdot t \right\} \bigg|_{t \approx t_{\text{Planck}}=1}^{\sqrt{\frac{\gamma}{4\pi}} - \sqrt{\frac{8\pi}{\gamma}}} \\
&\approx \frac{-\left[\frac{V_0}{3\gamma-1} + \frac{\gamma(3\gamma-1)}{8\pi} \right]}{\frac{1}{\kappa} \int \sqrt{-g} d^3x} + c_1 \frac{16\pi}{M_P^2} \cdot \left(\rho - \frac{\rho^2}{2|\sigma|} \right) \\
&\approx \hbar \cdot 8\pi G \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30} \ell_{\text{Planck}}} = 8\pi \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30}}
\end{aligned} \tag{33}$$

In Eq. (31) we will also make the following identification $\rho \approx \rho_{DE} \approx \hbar \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \bigg|_{\lambda_{DE} \approx 10^{30}}$

I.e. just before the bubble, we will factor in a very large bubble tension, σ

Doing so would be to have an optimal c_1 value, and that this choice of c_1 would allow us to make the identification of the choice of wavefunction for our model which would according to the Eq.(28) model have a quantum flavor, while at the same time in adherence be similar to the Eq.(31) values of Dark Energy density

3.2 What we obtain if we model the cosmological constant this way?

First of all, as noted by Kieffer, the wavefunction in Eq. (28) vanishes if $r=0$. We avoid having a vanishing wavefunction by asserting that we will be making an evaluation of it in the neighborhood of a Planck Length. Secondly in making this assumption we are at least proceeding in a direction where we can ditch the Anthropic principle, which I consider a MAJOR blight on cosmology. I also too though this this procedure as outline is experimentally verifiable, once we understand what may constitute embedding of Quantum Mechanics within a deterministic structure. The entire referral of this problem can be contrasted with the solution given in [19] by Rosen, which is in the mode of Quantum mechanics for a single particle in an isotropic universe,

The primary difference between the Rosen approach, [19] and what is done in this document is that the mass of a graviton, is proportional to the square of the Cosmological constant, an assumption which Rosen does not make. In addition the Rosen document assumes that " The situation is somewhat analogous to the quantum mechanics of a single particle as compared to quantum field theory" and assumes an isotropic background

We do not confine ourself to Isotropic backgrounds. We also do share with Rosen [19] a procedure as to rendition a linkage between classical to quantum mechanical treatment of our problem, but our work is more in tune with using what Powell and Craseman in their book on quantum mechanics used, [] which is similar to a limiting case of geometric optics.

I wish to thank Christian Corda as to alerting me to the Rosen article which is also applicable to a Dust Cosmology. I also wish to thank Fangyu Li , Wen Hao, and others for alerting me as to the crucial role of polarization. As it is, I presented part of this idea in Zeldovich 4, in the ICRANET on line conferences (a small part of it) and found that the

audience and the reviewers did not understand the Klauder treatment of the cosmological constant via Klauder Enhanced quantization.

Hence this expanded treatment whereas the Rosen treatment as Corda outlined is most effective for quantum black holes

3.3 Final reference to high frequency gravitational waves

$$(1 + z_{\text{initial-era}}) \equiv \frac{a_{\text{today}}}{a_{\text{initial-era}}} \approx \left(\frac{\omega_{\text{Earth-orbit}}}{\omega_{\text{initial-era}}} \right)^{-1} \quad (34)$$

$$\Rightarrow (1 + z_{\text{initial-era}}) \omega_{\text{Earth-orbit}} \approx 10^{25} \omega_{\text{Earth-orbit}} \approx \omega_{\text{initial-era}}$$

Equation (34) can be seen in [35] as well as [36]

Whereas We postulate that we specify an initial era frequency via dimensional analysis which is slightly modified by Maggiore for the speed of a graviton[[35] whereas

$$c(\text{light-speed}) \approx \omega_{\text{initial-era}} \cdot (\lambda_{\text{initial-post-bubble}} = \ell_{\text{Planck}}) \quad (35)$$

and that dimensional comparison with initially having a temperature built up so as

$$\Delta E \approx \hbar \omega_{\text{initial-era}} \quad (36)$$

where $T_{\text{universe}} \approx T_{\text{Planck-temperature}} = 1.22 \times 10^{19} \text{ GeV}$. If so then the Planck era

temperature would be extremely high leading to a change in temperature from the

Pre Planckian conditions to Planck era leading to

$$\Delta E = \frac{d(\text{dim})}{2} \cdot k_B \cdot T_{\text{universe}} \quad (37)$$

In doing so, be assuming

$$\omega_{\text{initial-era}} \approx \frac{c}{\ell_{\text{planck}}} \leq 1.8549 \times 10^{43} \text{ Hz} \quad (38)$$

Where we would be assuming $\omega_{\text{initial-era}} \approx \frac{c}{\ell_{\text{planck}}} \leq 1.8549 \times 10^{43} \text{ Hz}$ so then we

would be looking at frequencies on Earth from gravitons of mass m(graviton) less

than of equal to

$$\omega_{\text{Earth-orbit}} \leq 10^{-25} \omega_{\text{initial-era}} \quad (39)$$

And this partly due to the transference of cosmological ‘information’ as given in [13]

for a phantom bounce type of construction .Further point that since we have that gravitons travel at nearly the speed of light [36], that gravitons are formed from the surface of a bubble of space-time up to the electroweak era that mass values of the order of 10^{-65} grams (rest mass of relic gravitons) would increase due to extremely

high velocity would lead to enormous $\Delta E \approx \hbar \omega_{\text{initial-era}}$ values per graviton,

which would make the conflation of ultrahigh temperatures with gravitons traveling at

nearly the speed of light as given in Eq.(39) as compared with $\Delta E \approx \hbar \omega_{\text{initial-era}}$

We can in future work compare this with the Rosen[19] value of energy for a mini universe of (from a Schrodinger equation) with ground state mass of

$m = \sqrt{\pi} M_{\text{Planck}}$ and an energy of

$$E_n = \frac{-Gm^5}{2\pi^2 \hbar^2 n^2} \quad (40)$$

Our preliminary supposition is that Eq. (40) could represent the initial energy of a Pre Planckian Universe and that Eq.(37) be the thermal energy dumped in due to the use of Cyclic Conformal cosmology (maybe in multiverse form) so that if there is a build up of energy greater than Eq.(40) due to thermal build up of temperature due to infall of matter-energy, we have a release of Gravitons in great number which would commence as a domain wall broke down about in the Planckian era with a temperature of the magnitude of Planck Energy for a volume of radius of the order of Planck Length. This will be investigated in detailed future calculations . All this should be in fidelity, in experimental limits to [37], as well as looking at ideas about Quantum tunneling we may gain from [38] as to understand the transition from Pre Plankian to Planckian physics [39] [40]

acknowledgement

This work is supported in part by National Nature Science Foundation of China grant No.

11375279

References

1. Hawking, Stephen & Ellis, G. F. R. (1973). *The Large Scale Structure of Space-Time*. Cambridge: Cambridge University Press
2. [Shin'ichi Nojiri](#), [S.D. Odintsov](#), [V.K. Oikonomou](#), [Tanmoy Paul](#), "Nonsingular bounce cosmology from Lagrange multiplier $F(R)$ gravity", Phys. Rev. D **100**, 084056 – Published 25 October 2019; <https://arxiv.org/abs/1910.03546>
3. Wang, Qingdi; Zhu, Zhen; [Unruh, William G.](#) (2017). "How the huge energy of quantum vacuum gravitates to drive the slow accelerating expansion of the Universe". *Physical Review D*. **95**(10): 103504. [arXiv:1703.00543](#)
4. Adler, Ronald J.; Casey, Brendan; Jacob, Ovid C. (1995). "Vacuum catastrophe: An elementary exposition of the cosmological constant problem". *American Journal of Physics*. **63**(7): 620–626. [Bibcode:1995AmJPh..63..620A](#)

5. Astier, Pierre ([Supernova Legacy Survey](#)); Guy; Regnault; Pain; Aubourg; Balam; Basa; Carlberg; Fabbro; Fouchez; Hook; Howell; Lafoux; Neill; Palanque-Delabrouille; Perrett; Pritchett; Rich; Sullivan; Taillet; Aldering; Antilogus; Arsenijevic; Ballard; Baumont; Bronder; Courtois; Ellis; Filiol; et al. (2006). "The Supernova legacy survey: Measurement of Ω_m , Ω_Λ and W from the first year data set". *Astronomy and Astrophysics*. **447** (1): 31–48. [arXiv:astro-ph/0510447](#)
6. Frieman, Joshua A.; Turner, Michael S.; Huterer, Dragan (1 January 2008). "Dark Energy and the Accelerating Universe". *Annual Review of Astronomy and Astrophysics*. **46** (1): 385–432. [arXiv:0803.0982](#)
7. [John R. Klauder](#) "Enhanced Quantization: The Right way to Quantize Everything", <https://arxiv.org/abs/1702.04713>
8. J.R. Klauder, *Enhanced Quantization: Particles, Fields & Gravity*, (World Scientific, Singapore, 2015)
9. [Joan Sola](#), "Cosmological constant and vacuum energy: old and new ideas", J.Phys.Conf.Ser. 453 (2013) 012015, <https://arxiv.org/abs/1306.1527>
10. Poincaré Seminar, Duplantier, B., & Rivasseau, V. (2003). "Poincaré Seminar 2002: vacuum energy-renormalization". *Progress in mathematical physics*, v. 30. Basel: Birkhäuser Verlag.
11. Miao Li (Author), Xiao-Dong Li (Contributor), Shuang Wang (Contributor), Yi Wang (Contributor), "Dark Energy", Peking University Press, World Scientific, Singapore, Republic of Singapore, 2015
12. Katherine Freeze, Mathew Brown, and William Kinney, "The Phantom Bounce, A new proposal for an oscillating Cosmology", pp 149-156, of "The Arrows of Time, A debate in cosmology", Fundamental theories in physics, Volume 172, with Laura Mercini-Houghton, and Rudy Vaas, editors, Springer Verlag, Heidelberg, Federal Republic of Germany, 2012
13. Andrew Walcott Beckwith, "Using "Enhanced Quantization" to Bound the Cosmological Constant, (for a Bound on Graviton Mass), by Comparing Two Action Integrals (One Being from General Relativity) at the Start of Inflation", pp 21-36, of "Fundamental Physics and Physics Education Research", editors of Burra G. Sidharth • Jesús Carnicer Murillo • Marisa Michelini • Carmen Perea, Published by Springer Nature Switzerland AG 2021, in Gewerbestrasse 11, 6330 Cham, Switzerland
14. Poplawski, N. (2011) Cosmological Constant from QCD Vacuum and Torsion. *Annals of Physics (Berlin)*, 523, 291. <https://doi.org/10.1002/andp.201000162>
15. Saunders, S., & Brown, H. R. (1991). *The Philosophy of Vacuum*. Oxford [England]: Clarendon Press
16. [Steven D. Bass](#), "Vacuum energy and the cosmological constant", <https://arxiv.org/abs/1503.05483>
17. Klaus Kiefer, "Quantum Gravity, 3rd edition", Oxford Science Publications, Oxford University Press, Oxford, United Kingdom, 2012
18. Roger Penrose, (2006). ["Before the Big Bang: An Outrageous New Perspective and its Implications for Particle Physics"](#) (PDF). *Proceedings of the EPAC 2006, Edinburgh, Scotland*: 2759–2762.
19. Nathan Rosen, "Quantum Mechanics of a Miniuniverse", *International Journal of Theoretical Physics*, Vol. 32, No. 8, 1993
20. Ng, Y. Jack Entropy 2008, 10(4), 441-461; DOI: [10.3390/e10040441](#)
21. P. Halmos, *Ergodic theory*, Publ. Math. Soc. Japan, No. 3, Math. Soc. Japan, Tokyo, 1956
22. V. I. Arnold and A. Avez, *Ergodic Problems of Classical Mechanics*, (1968) W. A. Benjamin, Inc.
23. Omri Sarig, "Lecture Notes on Ergodic Theory", March 8, 2020, <http://www.weizmann.ac.il/math/sarigo/sites/math.sarigo/files/uploads/ergodicnotes.pdf>

24. LECTURES BY MARYAM MIRZAKHANI NOTES BY TONY FENG, "INTRODUCTION TO ERGODIC THEORY", https://www.mit.edu/~fengt/ergodic_theory.pdf
25. [Barrow, John D.](#); [Tipler, Frank J.](#) (1986). [The Anthropic Cosmological Principle](#) 1st edition 1986 (revised 1988). [Oxford University Press](#).
26. Padmanabhan, Thanu, "An Invitation to Astrophysics", World Press Scientific, World Scientific Series in Astronomy and Astrophysics: Volume 8, Singapore, Republic of Singapore, 2006
27. Geoffrey Compère, "Advanced Lectures on General Relativity", arXiv:1801.07064v4 [hep-th] 8 Feb 2019
28. Weinberg, Stephen, "Gravitation And Cosmology: Principles And Applications Of The General Theory Of Relativity", John Wiley and Sons, Cambridge, Massachusetts, USA, 1972
29. M. Lévy, J. Sucher (1969). ["Eikonal Approximation in Quantum Field Theory"](#). *Phys. Rev* (Maryland, USA). doi:10.1103/PhysRev.186.1656
30. Lawrence Horwitz, and Ori Oron, "Classical Gravity as an Eikonal Approximation to a Manifestly Lorentz Covariant Quantum Theory with Brownian Motion", Chapter 5, pp 143-190, of "Quantum Gravity Research Trends, editor Albert Reimer, Noval Science Publishers, Inc., 2006, New York City, New York, USA
31. John L. Powell, Bernd Crasemann, "Quantum Mechanics", Addison and Westley Company Inc. Reading Massachusetts, USA, 1961
32. George Arfken, Hans Weber, and Frank Harris, "Mathematical Methods for Physicists, A Comprehensive Guide", 7th edition, Academic Press, Elsevier, 2012, San Francisco, California, USA
33. Claus Kiefer, "Quantum Gravity, Third Edition", Oxford Science Publications, Oxford University Press, Oxford, United Kingdom, 2012
34. Claus Kiefer, "Path Integrals in Quantum Cosmology, Pages 729-40, In" Fluctuating Paths and Fields "(editor W. Janke, et.al., World Press Scientific, Singapore, Republic of Singapore, 2001
35. Maggiore, Michele, "Gravitational waves, Volume 1, theory and Experiment", Oxford University Press, New York City, New York, USA, 2008
36. Landau, L. D.; Lifshitz, E. M. (1975). *Classical Theory of Fields* (Fourth Revised English ed.). Oxford: Pergamon.
37. Misner, Charles; Thorne, Kip S.; Wheeler, John Archibald "Gravitation". San Francisco: W. H. Freeman., San Francisco, California, USA, 1973
38. Dilip K. Roy "A Quantum Measurement Approach to Tunnelling", World Press Scientific, Singapore, Republic of Singapore, 1993
39. Vasileiou, Vlasios; Granot, Jonathan; Piran, Tsvi; Amelino-Camelia, Giovanni (2015). ["A Planck-scale limit on spacetime fuzziness and stochastic Lorentz invariance violation"](#). *Nature Physics*. **11** (4): 344–346
40. Perlman, Eric S.; Rappaport, Saul A.; Christensen, Wayne A.; Jack Ng, Y.; DeVore, John; Pooley, David (2014). "New Constraints on Quantum Gravity from X-ray and Gamma-Ray Observations". *The Astrophysical Journal*. **805**: 10. [arXiv:1411.7262](#)