

# The concurrent detection of chelonid alphaherpesvirus 5 and *Chelonia mydas* papillomavirus 1 in tumoured and non-tumoured green turtles

Mashkour N, Jones K, Wirth W, Burgess G and Ariel E

College of Public Health, Medical and Veterinary Sciences, James Cook University,  
Townsville, 4814, Queensland, Australia.

## Abstract

Characterised by the growth of benign tumours, fibropapillomatosis (FP) is a debilitating disease that predominantly afflicts the endangered green turtle (*Chelonia mydas*). A growing body of histological and molecular evidence has consistently associated FP tumours with Chelonid alphaherpesvirus 5 (ChHV5), leading this virus to be considered the most likely aetiological agent of FP. However, a recent study which detected both ChHV5 and *Chelonia mydas* papillomavirus 1 (CmPV1) DNA in FP tumour tissues has challenged this hypothesis. The present study aimed to establish the wider prevalence of CmPV1 and co-occurrence with ChHV5 in marine turtles in waters adjacent to the east coast of Queensland, Australia. This comprehensive molecular survey screened a total of 353 samples from 275 foraging turtles using probe-based qPCR. Three sample categories were used in this study: Group A (FP tumours), Group B (non-tumoured skin from turtles with FP tumours) and Group C (non-tumoured skin from turtles without FP tumours). Concurrent detection of ChHV5 and CmPV1 DNA is reported for all three categories, with the highest rate of concurrent detection reported for Group A samples (43.5%). Collectively, these results pivot the way we think about FP; as an infectious disease where two separate viruses may be at play.

**Keywords:** Sea turtles; tumour; fibropapillomatosis; Chelonid alphaherpesvirus 5; *Chelonia mydas* papillomavirus 1

## 1. Introduction

The endangered green turtle (*Chelonia mydas*) faces many threats, each of which need to be effectively managed in order to conserve this vulnerable species. Although the impact of disease in wild populations is largely unknown [1], green turtles are particularly susceptible to a tumour-forming disease [2]. Fibropapillomatosis (FP) is a neoplastic disease characterised by the formation of benign tumours. Fibrosarcomas, fibropapillomas, papillomas, fibromas and myxofibromas have all been associated with FP, with lesion type thought to be linked to different stages of development [3-6]. Depending on the location of growth, these tumours can impair feeding, vision, and locomotion of affected turtles [3,4,7]. Turtles with FP tumours are also more vulnerable to chronic stress and secondary infections [8,9]. Although this disease has been reported in all species of marine turtle, it predominantly affects green turtles [2].

While the aetiological agent of FP is yet to be confirmed, early studies provided evidence for a viral aetiological agent of FP [10,11]. A body of subsequent research has identified a range of potential candidates; this includes papillomaviruses, polyomaviruses, retroviruses and herpesviruses [4,12-16]. However, histological and molecular evidence has consistently linked FP to a herpesviral infection [10,11,13,14,17-29].

Chelonid alphaherpesvirus 5 (ChHV5) is considered to be the most likely aetiological agent of this disease. Elements of ChHV5 DNA have been amplified from FP tumours and from turtles without tumours, which is consistent with the latent nature of herpesviral infections [17,18,23,24,29-31]. However, despite the development of more sensitive assays, varying rates of ChHV5 detection in FP tumours have been reported [23,25,28,32-34]. It is possible that this variable rate of detection is the result of the different methods used in these studies. Alternatively, these results may simply be accurate for each population studied. It has been suggested that the inconsistency of ChHV5 detection in FP tumours indicates the presence of other co-factors contributing to disease manifestation [28].

Based on the morphology and histopathology of FP tumours, an alternative aetiological agent could be papillomavirus. The epidermal hyperplasia and mesenchymal proliferation observed in FP is similar to equine sarcoids, a neoplastic disease of horses caused by bovine papillomavirus type 1 or type 2 [35]. The characteristics of the dermal proliferation is also similar to those observed in cattle fibropapillomas and deer fibromas [11], which are both caused by papillomaviruses [36,37].

Moreover, transmission electron microscopy studies on samples from FP tumours have reported virions similar in morphology to papillomavirus [16]. These histopathological indicators triggered

molecular investigations of a relationship between FP tumours and papillomaviruses a few decades ago, but they were not conclusive [16,38]. Subsequent research on links between papillomaviruses and FP were therefore abandoned until more recently, when a viral discovery project used primary cell lines from green turtles to screen a number of tissues from deceased turtles as well as FP tumour biopsies [39]. The cytopathic effect in cell cultures inoculated with material from the tumours was further investigated with current molecular techniques, which facilitated the identification of a papillomavirus. Mashkour *et al.* utilised species-specific assays for detection of virus [39], which likely contributed to the successful detection of both CmPV1 and ChHV5 in eight FP tumour samples collected from green turtles with FP, thereby challenging the perception that FP is associated with just herpesvirus.

It is clear that the role of papillomaviruses in FP and the fact that viruses are the least studied pathogens in sea turtles [1] warrant further investigation. The overall objective of the current study was to conduct a comprehensive molecular survey to better understand the disease ecology. This objective was met through the following aims: 1) Investigate presence of ChHV5 and CmPV1 DNA in a larger sample collection from Australian turtles both with and without FP tumours and 2) Establish the viral load of ChHV5 and/or CmPV1 present in these samples.

## 2. Materials and methods

### 2.1. Sample collection

A total of 275 marine turtles were sampled at six locations along the Great Barrier Reef. This included 89 turtles with FP tumours, 87 of which were green turtles. The remaining turtles with FP tumours consisted of one loggerhead (*Caretta caretta*) turtles and one green/hawksbill (*Eretmochelys imbricata*) hybrid turtle. Where possible, paired samples were collected from FP tumours and non-tumoured skin. In addition, multiple samples of FP tumours from individual turtles were taken opportunistically. Samples from a further 186 non-tumoured turtles were also collected from three of the six study sites. These samples were grouped into three categories: Group A (FP Tumours), Group B (Non-tumoured skin from turtles with FP tumours) and Group C (Skin from non-tumoured turtles) (Table 1). The total dataset consisted of samples from waters adjacent to Cairns (n=2), Townsville (n=80), Bowen (n=130), Airlie Beach (n=1), Gladstone (n=54) and Brisbane (n=8).

**Table 1.** The number of turtles used in this study categorised by study site and sample type.

Study Site	Total number of turtles	Number of turtles with FP tumours	Number of samples		
			Group A (FP Tumours)	Group B (Non-tumoured skin from turtles with FP tumours)	Group C (Skin from non-tumoured turtles)
Cairns	2	2	3	-	-
Townsville	80	24	52	13	56
Bowen	130	50	53	15	80
Airlie Beach	1	1	4	-	-
Gladstone	54	4	4	4	50
Brisbane	8	8	15	4	-
Total	275	89	131	36	186

Most turtles sampled in this study were captured using the rodeo capture technique [40] and flipper-tagged with a unique alpha-numeric inscribed titanium tag (Stockbrands Company, Pty. Ltd., Perth, Western Australia). The remaining tumour samples were collected during necropsy and others were donated (see Supplementary Tables S1, S2, S3). The curved carapace length (CCL) of each turtle in this study was measured using a flexible tape measure ( $\pm 2$  mm).

All samples were collected using fresh, sterile, disposable scalpel blades and stored in cryovials containing 90% ethanol. Any paired skin samples were collected from non-tumoured sites on the trailing edge of the front flipper of each turtle. Samples were stored at 4°C prior to DNA extraction.

All live turtles were sampled under permits from James Cook University Animal Ethics Committee (A1501 and A1971), Department of Environment and Science (WISP06619309 and WISP13754613) and Great Barrier Reef Marine Park Authority (G12/35326.1, G10/33220.1 and G36593.1).

## 2.2. DNA Extraction

DNA was extracted using Promega Wizard1 SV Genomic DNA Purification System according to the manufacturer's instructions with an additional 10µL of proteinase K used per reaction. DNA concentration was assessed by spectrophotometric analysis (NanoDrop™ 2000 Spectrophotometer), and DNA purity was determined using OD<sub>260/280</sub>. Extracted DNA was stored in -20 pre-analysis.

## 2.3. Primer and probe design

PCR primers and probes were designed and used to specifically amplify conserved regions of CmPV1 and ChHV5: the E1 gene and DNA polymerase, respectively. The green turtle Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) gene was selected to quantify a single-copy host target as a proxy to host cell copy number. These targets will hereby be referred to as CmPV1-E1, ChHV5-DNApol and GAPDH. In addition, one set of primers was designed to target the E2 gene from CcPV1 from loggerhead turtles (CcPV1-E2).

Primers and/or probes for ChHV5-DNApol and GAPDH targets were designed in this study, while CmPV1-E1 target was amplified using the Cm-Pap-109 primer and probe set described in Mashkour *et al.* (2018) (Table 2). Primer and probe design for the remaining targets was based on sequences available from GenBank (National Center for Biotechnology Information; NCBI, Bethesda, Maryland). Accession numbers HQ878327.2 and FJ234450.1 served as the main base for the as ChHV5-DNApol and GAPDH primers respectively. To increase the specificity of the assay, ChHV5-specific primers were designed based on the consensus sequences of DNA polymerase sequences isolated from green turtles only. Cc-Pap-99 primer set was designed to specifically target CcPV1-E2 gene. The E2 genes of CmPV1 (Genbank accession no. EU493091.1) and CcPV1 (GenBank accession no EU493092.1) were aligned to identify the site of highest variability. This section of the CcPV1-E2 was then used to design specific PCR primers to detect only CcPV1. All primers and probes were designed using AlleleID version 7.7 (Premier Biosoft International, Palo Alto, California) and synthesised by Macrogen Inc. (Seoul, South Korea) (Table 2).

**Table 2.** The sequences of the PCR primers and probes along with the PCR cycling protocols used in the PCR to amplify the target genes from cloned plasmids (standards), positive controls and test samples. Abbreviations: BHQ1, black hole quencher-1; FAM, 6-carboxyfluorescein; F, forward primer; R, reverse primer; P, probe.

	Primers and probes	Sequence	Target gene	Amplicon length	Cycling conditions	Reference
Cloned plasmids for standard assays	Cm-Dpol-864	F: 5'-ATG ACG GAC GGA CAA CAG-3'	ChHV5-DNApol	864 bp	2 min at 95°C; 35 cycles (95°C for 10s, 57 °C for 15s, and 72°C for 55s)	The present study
		R: 5'-GGA GAT GAC GGC TGC TAA-3'				
	GAPDH-475	F: 5'-CCT TTA ATG CGG GTG CTG-3'	GAPDH	475 bp	2 min at 95°C; 35 cycles (95°C for 10s, 57 °C for 15s, and 72°C for 30s)	The present study
		R: 5'-CAC GGT TGC TGT ATC CAA-3'				
Screening assays	Cm-Pap-109	F: 5'-GCC GAT GAT GTC CAC TTA T-3'	CmPV1-E1	109 bp	2 min at 95°C; 40 cycles (95°C for 10s, 60 °C for 20s, and 72°C for 30s)	Mashkour <i>et al.</i> (2018)
		R: 5'-GCT GAA TCC ACA GAG GTA G-3'				
	Cm-Pap-109	P: 5'-FAM CGA CCC ATG AAG CCG CTG T BHQ1-3'	CmPV1-E1	109 bp	2min at 95°C; 50 cycles (95°C for 10s, 60°C for 10s)	Mashkour <i>et al.</i> (2018)
	Dpol-ChHV5-82	F: 5'-CTA CCT TGT CTG GAG GTG GC-3'	ChHV5-DNApol	82 bp	2min at 95°C; 50 cycles (95°C for 10s, 60°C for 10s)	The present study
		R: 5'-GGG TGT GAA TAA AAT CCC GCG-3'				
		P: 5'-FAM TAG GGC GCG ACA TGC TTC BHQ1-3'				
	GAPDH-83	F: 5'-CTG GTC TCC TGG TAT GGA-3'	GAPDH	83 bp	2min at 95°C; 50 cycles (95°C for 10s, 60°C for 10s)	The present study
		R: 5'-CAT GGA CTC CCA ACC TAT C-3'				
		P: 5'-FAM AAA CCA CCC TCC AAA TCT GGC BHQ1-3'				
	Cc-Pap-99	F: 5'-AAA GGG CAG TGG GAA ATC TC-3'	CcPV1-E2	99 bp	2 min at 95°C; 40 cycles (95°C for 10s, 58 °C for 20s, and 72°C for 30s)	The present study
		R: 5'-TGT GAT GGC GAC GAT GTG-3'				

GoTaq® Probe qPCR reactions (20µl) were made with 10µl of GoTaq® Probe qPCR Master Mix (Promega), 1.6µl of each primer (0.8µM), 0.2 µl of probe, ~2µl of template DNA (~80 ng) and 4.6µl nuclease free water. All GoTaq® Probe qPCRs were carried out using a Magnetic Induction Cyclers (MIC) qPCR machine (Bio Molecular Systems). The thermocycling conditions for these assays were as follows: 2min at 95°C followed by 50 cycles (95°C for 10s, 60°C for 10s). The dye-based qPCR reactions (20µl) were made with 10µl of 5X GoTaq® qPCR Hot Start Master Mix (Promega), 1.6µl of each primer (0.8µM), ~2µl of template DNA (~80 ng) and 4.8µl nuclease free water. The PCR was carried out on a Rotor-Gene 6000 thermocycler machine. The results of these assays were visualised in real time with the thermocycling conditions provided in Table 1.

A green turtle embryo fibroblast cell line, CMEM which previously tested negative for herpesviral and papillomaviral DNA using published PCR primers (Mashkour *et al.*, 2018), was used as the negative biological control for these assays.

#### 2.4. Producing standard curves for ChHV5, CmPV1 and green turtle genomic DNA

To be able to report viral copy numbers, robust standards were required from each of the viruses and the host samples. For ChHV5-DNApol and GAPDH targets, Cm-Dpol-864 and GAPDH-475 assays were used to generate amplicons for use as plasmid inserts (Table 2). At the time, the Australian CmPV1 strain wasn't fully sequenced and to avoid possible SNPs and missing samples, for CmPV1-E1 target the Cm-Pap-109 primers were used from Mashkour *et al.* [39].

After successful polymerisation, the amplicons CmPV1-E1, ChHV5-DNApol and GAPDH were cloned separately into pGEM®-T Easy Vectors using the TA blue/white cloning system according to pGEM®-T Easy Vector Systems protocol (Promega). To confirm the presence of desired inserts the plasmid DNAs were sequenced along both strands using a universal M13 primer set (Macorgen Inc., Seoul, Korea).

A supercoiled circular plasmid can suppress the PCR amplification and result in an error as big as seven copies of the target gene per cell [41]. To avoid misinterpretation of viral loads the cloned plasmids were treated with restriction enzymes. To select an enzyme which only cut the pGEM®-T Easy Vector once, without digesting the target genes, the vector and the cloned sequences were analysed via Geneious v10.1 [42], subsequently SacI and PstI were chosen for treatment. The linear plasmids were eluted in DNase free water and stored at -20°C.



## 2.5. Calibration curves and real-time quantitative PCR

For the setup of a dilution series, the mass of one copy of the cloned CmPV1-E1, ChHV5-DNApol and GAPDH DNA plasmid were calculated using the mass of 1bp. The gram per mol of each amplicon was used to determine the appropriate amount needed for the initial dilution steps. The measurement was obtained using a Quantus™ Fluorometer using QuantiFluor® dsDNA Systems.

Tenfold serial dilutions were prepared in nuclease free water to produce a titration series from  $10^8$  to  $10^1$  of each plasmid in 2µl/reaction (as the GoTaq® Probe qPCR was undertaken using 2µl of template DNA in a 20µl PCR). From this, calibration curves for pCmPV1-E1, pChHV5-DNApol and pGAPDH were produced. To make the standard curves, each dilution point was assayed in triplicate and the highest efficiencies were considered for the following calculations. Calibration curves for pCmPV1-E1, pChHV5-DNApol and pGAPDH were constructed by plotting cycle numbers versus the log concentrations using the MIC software (see Figure S1).

## 2.6. Viral and host qPCR reaction conditions and data interpretation

Linear pCmPV1-E1, pChHV5-DNApol and pGAPDH were run independently in singleplex qPCR reactions. Samples were tested in duplicates and in each run a negative biological control (green turtle primary culture), a no template control (NTC; nuclease free water) and triplicate standards were included. The MIQE guidelines [43] were followed to avoid assay contamination by genomic or plasmid DNA. To ensure repeatability, a single person ran the qPCRs and measures of intra and inter-assay variation were used. Care was taken to provide a wide range standard curve and cover the copy number range observed in “unknown” samples. Samples were considered positive if the emitted fluorescence of both duplicates exceeded the threshold limit of detection (LOD) and were within 0.5  $C_t$  of each other and negative otherwise. Lower LOD was determined by carrying out qPCR on 24 replicates of low copy number per reaction for each plasmid (5, 10, 50 and 100 target copies per 2 µL). The 50% and 100% LOD were calculated for linear pCmPV1-E1, pChHV5-DNApol and pGAPDH.

The runs were considered valid if the efficiency was between 90-110%, NTC and negative control were negative and if the MIC standardised value would let import the normalised standard curve for the run. Data were analysed in the MIC software where an adaptive baseline threshold was automatically generated for each assay. PCR efficiencies were calculated using the formula:  $E = (10^{-1/\text{slope}} - 1) \times 100$ .

The viral copy number of each virus per cell was calculated according to CmPV1-E1 and/or ChHV5-DNApol copies per host cell using the following equation:

$$= \frac{\text{CmPVE1 and/or ChHV5DNApol quantity}}{\frac{1}{2} \text{GAPDH quantity}}$$

For all samples, viral load per diploid genome was determined by dividing CmPV1-E1 and ChHV5-DNApol copy numbers by half of the GAPDH copy number.

## 2.7. Statistical analysis

The resulting copy numbers of group A and B were compared using unpaired t-test with a 95% confidence interval. The two-tailed P-value was also calculated. Dealing with non-parametric groups of results, the Kruskal–Wallis test by ranks was done on samples from different regions to investigate the distribution of each virus in six regions covered by this study. The rate of detection was calculated using Chi-square and Fisher's exact test of independence and was used to compare the portion of animals, tumours, non-tumoured skin from turtles with FP tumours, and skin from non-tumoured turtles, with each virus in each region. Values with P-value<0.05 were considered statistically significant. The statistical analyses were conducted using IBM SPSS Statistics 25. The mean copy number of ChHV5 in each sample group in the present study was also compared with results in other ChHV5 studies [24,44,45]. Unit conversions were done where appropriate.

### 3. Results

#### 3.1. qPCR validation and optimisation

In silico evaluation of the designed primers and probes confirmed the specificity of these oligonucleotides. Initial amplified products and plasmid amplicons were sequenced and had 100% pairwise identity to the desired target genes. The known negative controls (both NTC and biological control) did not react in any assay.

The standard curves were successfully constructed and used for the qPCRs and subsequently the calculation of copy numbers per cells of each virus. The standard curves for linear pGAPDH, pCmPV1 and pChHV5 are shown in supplements (Figures S1). Absolute quantification and the standard curves for three cloned plasmids were plotted based on the cycles (Cq values) and the log of concentration (10 to 10<sup>8</sup> copies per reaction). The data points represent three replicates of each dilution. The R<sup>2</sup> values, the efficiency of each qPCR and the equation of the standard curves are as follows: pGAPDH R<sup>2</sup> 0.950; Efficiency 0.99; Equation  $y = -3.35x + 40.76$ ; pChHV5 Dpol R<sup>2</sup> 0.993; Efficiency 0.92; Equation  $y = -3.54x + 40.49$ ; pCmPV1-E1 R<sup>2</sup> 0.974; Efficiency 1.05; Equation  $y = -3.21x + 40.48$ .

The qPCR sensitivity was determined by 50% certainty cut-off for LOD as described by World Organization for Animal Health [46]. In this study, 10 copies per reaction were detected in 75% of the replicates for CmPV1-E1 and ChHV5-DNApol while 50 copies per reaction were detected for GAPDH. The 100% LOD was 100 copies per reaction for all the three plasmid DNAs.

#### 3.2. Viral DNA detection

A total of 353 samples from 275 foraging turtles were screened in this study using probe-based qPCR. Of the turtles sampled in this study, 89 individual turtles were afflicted with FP tumours. On an individual turtle basis, the detection of CmPV1 and/or ChHV5 amplicons varied from 0 to 100% in turtles with FP tumours. If one sample from one animal reacted in the PCR, the animal was considered positive for the presence of the tested virus. Of these 89 turtles, 69 (77.3%) tested positive for ChHV5, 46 (51.7%) for CmPV1, 40 (44.9%) for both viruses and samples from 14 (15.7%) turtles tested negative for both.

Viral DNA detection of both ChHV5 and CmPV1 varied between sample types. Group A had the highest rates of viral detection and within this group ChHV5 was detected in 86.3% of samples (Table 3). This is almost double the rate of detection of CmPV1 in the sample type. Moreover, 57 of the 62 samples in which CmPV1 was detected also reacted to the ChHV5 assay. Therefore, ChHV5 DNA was

more predominant in FP tumours but was often detected alongside CmPV1. Only 9.2% of the Group A samples did not react in either assay (Table 3). Interestingly, a higher rate of detection for both viruses was observed in the Group C samples compared to Group B, despite the Group C samples being sourced from turtles without FP tumours (Table 3). Similarly, of the samples from which no viral DNA was detected Group B samples had the highest rate (Table 3).

**Table 3.** The results from the 353 samples screened in this study for the presence of chelonid alphaherpesvirus 5 (ChHV5) and *Chelonia mydas* papillomavirus 1 (CmPV1). The results are categorised by sample types: Group A (FP Tumours), Group B (Non-tumoured skin from turtles with FP tumours) and Group C (Skin from non-tumoured turtles). A summary of the total detections is also provided.

Sample Type	Viral DNA Detections			
	ChHV5	CmPV1	Both ChHV5 and CmPV1	Neither ChHV5 and CmPV1
Group A	113/131 (86.3%)	62/131 (47.3%)	57/131 (43.5%)	12/131 (9.2%)
Group B	4/36 (11.1%)	9/36 (25.0%)	1/36 (2.8%)	24/36 (66.7%)
Group C	39/186 (21.0%)	68/186 (36.6%)	8/186 (4.3%)	85/186 (45.7%)
Overall	156/353 (44.2%)	139/353 (39.4%)	66/353 (18.7%)	121/353 (34.3%)

Using the specific primer sets for loggerhead papillomavirus (Cc-Pap-99 PCR primers), CcPV1 was detected in six tumour samples of loggerhead turtles. The tumour tissues of loggerhead samples did not react in CmPV1-specific assays and the green samples did not react in CcPV1-specific assays. This provides strong support for the already defined species barrier between the CmPV1 and CcPV1 viruses.

### 3.2.1. Viral loads of CmPV1 and ChHV5

The mean viral loads detected in each of the three sampling groups are reported in Table 4. The ChHV5 load in Group A ranged from 0 to 226.1 (mean: 15.635; SD: 30.152) copies in each cell and the CmPV1 load range was 0 to 2.54 (mean: 0.0814 ( $\approx 1$  copy in 12 cells); SD: 0.357). The raw data is provided in Supplementary Tables S1, S2, S3. With respect to ChHV5, the highest viral load was detected in the Group A samples (FP tumours). Interestingly, the mean ChHV5 load for Group C samples was slightly higher than that for Group B despite the fact that Group B samples were collected from turtles afflicted with FP. The viral load per cell of ChHV5 in Group A samples is significantly higher than Group B samples ( $P$ -value<0.0001). The mean viral load of CmPV1 varied amongst the sample groups, with the highest load detected in Group B (Table 4). The FP tumour samples (Group A) had a lower mean load of CmPV1 than Group B, this difference is statistically

significant (P-value=0.0138). Overall, the highest mean viral load detected was ChHV5 in Group A samples (Table 4).


**Table 4.** Mean viral load (reported as copy number per cell ± standard deviation) detected in the three different biological sample types screened in this study; Group A (FP Tumours), Group B (Non-tumoured skin from turtles with FP tumours) and Group C (Skin from non-tumoured turtles).

	ChHV5	CmPV1
	(copy number per cell)	(copy number per cell)
Group A	15.635± 30.152	0.0814± 0.357
Group B	0.0204± 0.0379	0.855± 2.351
Group C	0.100 ± 0.606	0.005 ± 0.023

The mean copy number of ChHV5 in each sample group in the present study was found to be consistent with results in other ChHV5 studies (See supplementary table S4). No similar undertaking could be done for the CmPV1 results as there are no comparable datasets in sea turtles. However, the results are consistent with human cervical intraepithelial neoplasia (CIN) I, II, III caused by human papillomavirus (HPV) 16, 18, 33 and 58 [47-49].

3.3. Geographic distribution of ChHV5 and CmPV1

Viral detection varied between the six regions in this study, both between regions and within regions but between viral detections (Figure 1). In Group A samples from both Townsville, Brisbane, and Bowen, ChHV5 was detected in significantly higher rates than CmPV1 ( $P<0.05$ ) (Table 5). This is also true of Group C samples from Gladstone ( $P<0.001$ ) (Table 5). However, in Group C samples collected from Bowen, significantly higher rates of CmPV1 were detected ( $P<0.001$ ) (Table 5).



**Figure 1.** The distribution of chelonid alphaherpesvirus 5 (ChHV5) and *Chelonia mydas* papillomavirus 1 (CmPV1) in FP tumour samples (Group A) collected from green turtles at six regions along the Queensland coast. A total of 131 FP tumour tissues were tested. Of the assay samples 62 reacted in the qPCR for CmPV1, 113 for ChHV5, and 12 samples did not react in either assay. 57 tumor samples reacted in both viruses. The viral detection at each site is categorised into three categories: ChHV5 only (red), CmPV1 only (green), both ChHV5 and CmPV1 detected in the same

sample (beige). The total number of sample screened (n) and the total number of negatives (neg) are shown for each location.


**Table 5.** Number of viral DNA detections and mean viral load (reported as copy number per cell  $\pm$  standard deviation) of the samples which tested positive in the three different biological sample types screened in this study; Group A (FP Tumours), Group B (Non-tumoured skin from turtles with FP tumours) and Group C (Skin from non-tumoured turtles). The number of detections within each group were compared using Chi square and Fisher’s exact test, respectively, with the results also reported here. Statistically significant differences are highlighted in bold. Unavailable samples are denoted by a hyphen.

		Group A			Group B			Group C		
		ChHV5	CmPV1	P-value	ChHV5	CmPV1	P-value	ChHV5	CmPV1	P-value
Cairns	No. (%) positive	3/3 (100.0%)	3/3 (100.0%)	1.000	-	-	-	-	-	-
	Average copy number/cell	39.300 $\pm$ 56.788	0.002 $\pm$ 0.002		-	-	-	-	-	-
Townsville	No. (%) positive	51/52 (98.1%)	26/52 (50.0%)	<b>&lt;0.00001</b>	2/13 (15.4%)	2/13 (15.4%)	1.000	5/56 (8.9%)	11/56 (19.6%)	0.105
	Average copy number/cell	11.364 $\pm$ 16.120	0.010 $\pm$ 0.030		0.039 $\pm$ 0.053	0.023 $\pm$ 0.033		0.758 $\pm$ 1.695	0.009 $\pm$ 0.024	
Bowen	No. (%) positive	39/53 (73.6%)	26/53 (49.1%)	0.0095	1/15 (6.7%)	6/15 (40.0%)	0.0309	17/80 (21.25%)	52/80 (65%)	<b>&lt;0.00001</b>
	Average copy number/cell	16.456 $\pm$ 28.911	0.088 $\pm$ 0.257		-			0.231 $\pm$ 0.918	0.005 $\pm$ 0.024	
Airlie Beach	No. (%) positive	4/4 (100%)	2/4 (50.0%)	0.429	-	-	-	-	-	-
	Average copy number/cell	3.554 $\pm$ 0.835	0.001 $\pm$ 0.0004		-	-	-	-	-	-
Gladstone	No. (%) positive	2/4 (50.0%)	2/4 (50.0%)	1.000	0/4 (0.0%)	1/4 (25.0%)	1.000	22/50 (44.0%)	6/50 (12.0%)	<b>0.0003</b>
	Average copy number/cell	2.061 $\pm$ 2.900	0.003 $\pm$ 0.003		-	-	-	0.000004 $\pm$ 0.00001	0.0003 $\pm$ 0.0008	
Brisbane	No. (%) positive	14/15 (93.3%)	4/15 (26.6%)	<b>&lt;0.001</b>	1/4 (25.0%)	0/4 (0.0%)	1.000	-	-	-
	Average copy number/cell	29.220 $\pm$ 58.466	0.634 $\pm$ 1.267		-	-	-	-	-	-




### 3.3.1. Geographic distribution of viral loads

The distribution of ChHV5 viral load is comparable in Group A samples across the six regions tested, as Kruskal–Wallis test by ranks resulted in a non-significant difference between the ChHV5 viral load in tumour samples from these six regions (ChHV5<sub>GroupA</sub> P value=0.100). Group C turtles from these regions also don't have significantly different loads of ChHV5 (ChHV5<sub>GroupC</sub> P value=0.122). The viral load distributions in samples that contained ChHV5 DNA from six regions of Queensland can be seen in Figure 2.



**Figure 2.** Boxplots of ChHV5 viral loads (log scale) in different sample types from six regions of Queensland. Group A (FP Tumours), Group B (Non-tumoured skin from turtles with FP tumours) and Group C (Skin from non-tumoured turtles). The Box plots of viral loads include the median (centre horizontal line), interquartile range (box), minimum and maximum (whiskers), and outliers (diamonds).

The distribution of CmpV1 load is the same in the tumours sampled across the six regions tested, as Kruskal–Wallis test by ranks resulted in a non-significant difference between the CmpV1 viral load in tumour samples from these six regions (CmpV1<sub>GroupA</sub> P value=0.317). This trend can be seen in Figure 3.



**Figure 3.** Box-plots of CmPV1 viral loads (log scale) in different sample types from six regions of Queensland. Group A (FP Tumours), Group B (Non-tumoured skin from turtles with FP tumours) and Group C (Skin from non-tumoured turtles). The Box plots of viral loads include the median (centre horizontal line), interquartile range (box), minimum and maximum (whiskers), and outliers (diamonds).

#### 4. Discussion

Although papillomavirus was proposed as an aetiological agent of FP in the 1990s [4] and this hypothesis was supported by transmission electron microscopy evidence [16], papillomaviral DNA was not successfully detected in FP tumour tissue in studies conducted around this time [16,38]. Conversely, a body of histological and molecular evidence emerged which led to herpesviral infection to be considered the primary candidate for an aetiological agent of FP [10,11,13,14,18-29,32]. Despite the subsequent discovery and description of CmPV1 and CcPV1 [50,51], these papillomaviruses were not linked to FP lesions. However, following the initial detection of both ChHV5 and CmPV1 DNA in a small sample size of FP tumour tissue [39], the current study set out to test the wider prevalence of CmPV1 and co-occurrence with ChHV5 in marine turtles in waters adjacent to the east coast of Queensland, Australia. Three sample categories were used in this study: Group A (FP tumours), Group B (non-tumoured skin from turtles with FP tumours) and Group C (non-tumoured skin from turtles without FP tumours). Concurrent detection of ChHV5 and CmPV1 DNA is reported for all three categories which challenges previous hypotheses of ChHV5 as the leading etiological agent of FP. This study is the first to report concurrent detection of viral DNA from ChHV5 and CmPV1 in non-tumoured skin samples collected from marine turtles. This study is also the first

comprehensive molecular survey of CmpV1 utilising species-specific primers, allowing for a more in-depth investigation of the results in Mashkour *et al.* [39]. However, the results of this study also provide further evidence toward the association between ChHV5 and FP, where higher detection rates of ChHV5, compared to CmpV1, are reported in FP tumour samples (Group A). Furthermore, regional differences in the rate of detection and viral loads in Group A samples support the hypothesis of FP tumour “hot-spots” in Queensland [52]. Collectively, these results pivot the way we think about FP; as an infectious disease where two separate viruses may be at play.

### Rate of detection

Overall, the rate of detection for ChHV5, CmpV1 and the combination of both was highest for Group A (Table 3). These results are consistent with the theory that ChHV5 is associated with FP tumours [2,10,14,24,53], and open a new line of scientific inquiry regarding the role of CmpV1 in FP tumour development. However, if a simple cause-and-effect relationship is at play, a detection rate of 100% in FP tumour samples would provide strong support of this. Although the rates of detection of these viruses were highest in FP tumour samples, these rates were still variable; ChHV5 was detected in 86.3% of FP tumour samples, while CmpV1 was detected in less than half (47.3%) of these samples. Although there is no comparable data available for CmpV1, this variable rate of ChHV5 detection is consistent with similar studies [17,23,25,28,54]. However, the detection of CmpV1 in Group A samples provides yet another small piece in the puzzle for the trigger factors involved in FP tumour development. Taken together, these results add weight to the theory that there are other factors at play in FP development [2,28,55].

This study also serves as a validation for newly developed qPCR assays for the detection of ChHV5, CcPV1 and GAPDH described herein. Of note is the improved sensitivity of the Dpol-ChHV5-82 assay in comparison to conventional PCR assays previously used in this region by Jones *et al.* [28] to facilitate longer amplicon length for subsequent phylogenetic analyses. Using a combination of conventional PCR assays ChHV5 DNA was detected in 62 of the 89 Group A individual turtles [28]. In the present study, ChHV5 DNA was detected in 69 Group A turtles using a single qPCR assay (Dpol-ChHV5-82), which emphasizes the need to match the assay selection with end-point goals.

Consistent with the known advantages of qPCR [56], our results show that qPCR assays are suitable in studies which aim to determine presence or absence of viral DNA, due to the enhanced sensitivity of these assays over conventional PCR. Moreover, the benefits of qPCR assays can extend beyond simple presence/absence tests into viral load assessment for positive samples.

## Mean viral load

Here, we reported viral loads for both ChHV5 and CmPV1 across three sample types (Groups A, B and C). The highest mean load of ChHV5 was found in Group A samples, adding to the body of evidence that this virus is closely associated with FP tumours. The highest mean load of CmPV1 was detected in Group B samples, but the significance of this result is unclear. The small sample size comprising this Group (n=36), in contrast to Group A and C (n= 131 and 186 respectively), limit the conclusions that can be drawn about this result. Further analysis on a larger set of samples of this type (non-tumoured skin from turtles with FP tumours) would undoubtedly add resolution to this result. Other than the case of ChHV5 in Group A samples, the mean viral load in all other sample types for both viruses was less than one copy per cell indicating a low level infection in these samples.

While the mean ChHV5 load results obtained in this study were consistent with other reports [24,44,45], such comparisons could not be made for CmPV1 as there are no other comparable datasets in marine turtles. However, the CmPV1 viral loads are in the same range reported for human cervical malignancies caused by HPV 16, 18, 33 and 58 where the mean HPV copies per 10,000 cells were reported to be  $\sim \log 3-5$  in CIN I, II and III [47-49]. Additional molecular surveys targeting both ChHV5 and CmPV1 from other regions would be hugely beneficial to understanding the role of CmPV1 in FP, and in marine turtle health as a whole.

This study assessed both the rate of detection and viral load of ChHV5 and CmPV1 and, in some cases, there was disparity between the rate of detection and viral load. For example, CmPV1 was detected in 100% of Group A samples from Cairns yet the viral load was relatively low ( $0.002 \pm 0.002$  copies per cell). This is in comparison to a ChHV5 load of  $39.300 \pm 56.788$  copies per cell in the same samples with the same rate of detection adding to the complexity of the picture rather than a presence/absence result.

## Concurrent detection

Concurrent infections of herpesvirus and papillomavirus have been described extensively in humans (see [57]) and in Atlantic bottlenose dolphins [58]. Our study reports the concurrent presence of herpesviral and papillomaviral DNA in both tumoured and non-tumoured marine turtles. The presence of these viruses in non-tumoured skin of turtles with FP tumours (Group B) and those free of clinical disease (Group C) are consistent with previous studies on these viruses [22,45,59] and are in line with the ubiquity and commensal nature of both viruses [45,60]. These results lend further weight to the hypothesis that FP manifestation relies on more than just exposure to viral infection alone [2].

## Locations

Differences in both viral load and rate of detection of both ChHV5 and CmPV1 were found across the six study sites. The highest mean viral load was detected in Cairns. This group was comprised of three samples from two turtles which were both severely afflicted by FP tumours. Higher rates of detection and viral loads are reported in Group A samples from Townsville, Bowen and Brisbane. These results are consistent with previous suggestions that these regions are FP “hot-spots” in Queensland [52]. However, when examining the data beyond mean loads, it is clear that higher loads of ChHV5 were detected in Group A samples at each study site. This suggests that while there may indeed be FP tumour hot-spots in Queensland, this relationship is not directly linked with viral load. Bowen is of particular interest, being the only study site where some Group A samples contained CmPV1 exclusively. In all other study sites, when CmPV1 was detected from Group A samples it was in conjunction with ChHV5. Moreover, Group C samples from Bowen had a higher rate of CmPV1 detection compared to ChHV5 (P-value<0.001) indicating a high prevalence of papillomavirus in clinically healthy turtles in this region, which would make it an excellent study site for more in-depth investigations on CmPV1.

## Limitations

In a field that relies on opportunistic sampling of free ranging marine wildlife, it was a major effort to collect samples (n=353) from a total of 275 individual turtles, especially those with FP. Although more samples distributed evenly between sample groups and study sites may have provided more confidence in the conclusions that can be drawn regarding these categories, such sample sets are not always realistic in wildlife disease investigations [28]. Therefore, in the interest of elucidating the potential role of papillomavirus in FP as identified by Mashkour *et al.* [39], we have reported the findings for all sites and sample types. Assay sensitivity and specificity is always a consideration and as tests are constantly evolving, future development may include more sensitive methods and targets such as viral mRNA as an indicator of viral expression. Despite these limitations, the results reported in the present study show some clear and reliable trends which we hope will be further explored on a global scale.

Our understanding of FP, and marine turtle virology as a whole, is riddled with knowledge gaps [1] which need to be addressed in order to develop comprehensive management plans for these endangered species. The results from this and other studies have revealed that viral DNA is present in animals that do not show clinical signs, which questions their role in tumour development and

highlights the need for further understanding of co-factors in disease development and why some turtles do not develop tumours despite the presence of viral DNA. Diseases in terrestrial animals are often tracked by serological methods, and such tests could be more widely implemented to determine whether turtles have mounted an immune response to ChHV5, CmPV1 and/or CcPV1 which would negate the need to rely on clinical signs alone for exposure status of an animal [9].

The results of the present study provide valuable insight into some of these gaps and with improved detection of papillomavirus in FP tumour tissue, it also challenges long held perceptions of the role of herpesvirus as the sole infectious agent in FP tumours.

## 6. Acknowledgement

The authors gratefully acknowledge Ms. Johanna Leonhardt and Dr. Ian Bell for their efforts in sample collection of the Group C Gladstone samples.

## 7. References

1. Mashkour, N.; Jones, K.; Kophamel, S.; Hipolito, T.; Ahasan, S.; Walker, G.; Jakob-Hoff, R.; Whittaker, M.; Hamann, M.; Bell, I., et al. Disease risk analysis in sea turtles: A baseline study to inform conservation efforts. *PLOS ONE* **2020**, *15*, e0230760, doi:10.1371/journal.pone.0230760.
2. Jones, K.; Ariel, E.; Burgess, G.; Read, M. A review of fibropapillomatosis in Green turtles (*Chelonia mydas*). *The Veterinary Journal* **2016**, *212*, 48-57, doi:<http://dx.doi.org/10.1016/j.tvjl.2015.10.041>.
3. Work, T.M.; Balazs, G.H.; Rameyer, R.A.; Morris, R.A. Retrospective pathology survey of green turtles *Chelonia mydas* with fibropapillomatosis in the Hawaiian Islands, 1993-2003. *Diseases of aquatic organisms* **2004**, *62*, 163-176.
4. Herbst, L.H. Fibropapillomatosis of marine turtles. *Annual Review of Fish Diseases* **1994**, *4*, 389-425, doi:10.1016/0959-8030(94)90037-X.
5. Norton, T.; Jacobson, E.; Sundberg, J. Cutaneous Fibropapillomas and Renal Myxofibroma in a Green Turtle, *Chelonia mydas*. *Journal of wildlife diseases* **1990**, *26*, 265-270, doi:10.7589/0090-3558-26.2.265.
6. Kang, K.; Torres-Velez, F.; Zhang, J.; Moore, P.; Moore, D.; Rivera, S.; Brown, C. Localization of fibropapilloma-associated turtle herpesvirus in green turtles (*Chelonia mydas*) by in-situ hybridization. *Journal of comparative pathology* **2008**, *139*, 218-225.
7. Flint, M.; Limpus, C.; Patterson-Kane, J.; Murray, P.; Mills, P. Corneal fibropapillomatosis in green sea turtles (*Chelonia mydas*) in Australia. *Journal of Comparative Pathology* **2010**, *142*, 341-346.
8. Aguirre, A.A.; Balazs, G.H.; Spraker, T.R.; Gross, T.S. Adrenal and hematological responses to stress in juvenile green turtles (*Chelonia mydas*) with and without fibropapillomas. *Physiological Zoology* **1995**, *68*, 831-854.
9. Work, T.M.; Rameyer, R.A.; Balazs, G.H.; Cray, C.; Chang, S.P. Immune status of free-ranging green turtles with fibropapillomatosis from Hawaii. *Journal of wildlife diseases* **2001**, *37*, 574-581.

10. Herbst, L.H. The etiology and pathogenesis of green turtle fibropapillomatosis. 1995.
11. Jacobson, E.; Mansell, J.; Sundberg, J.; Hajjar, L.; Reichmann, M.; Ehrhart, L.; Walsh, M.; Murru, F. Cutaneous fibropapillomas of green turtles (*Chelonia mydas*). *Journal of comparative pathology* **1989**, *101*, 39-52.
12. Casey, R.N.; Quackenbush, S.L.; Work, T.M.; Balazs, G.H.; Bowser, P.R.; Casey, J.W. Evidence for retrovirus infections in green turtles *Chelonia mydas* from the Hawaiian islands. *Diseases of Aquatic Organisms* **1997**, *31*, 1-7, doi:10.3354/dao031001.
13. Jacobson, E.R.; Buergett, C.; Williams, B.; Harris, R.K. Herpesvirus in cutaneous fibropapillomas of the green turtle *Chelonia mydas*. *Diseases of Aquatic Organisms* **1991**, *12*, 1-6.
14. Quackenbush, S.L.; Bowser, P.R.; Work, T.M.; Balazs, G.H.; Casey, R.N.; Casey, J.W.; Rovnak, J.; Chaves, A.; duToit, L.; Baines, J.D., et al. Three Closely Related Herpesviruses Are Associated with Fibropapillomatosis in Marine Turtles. *Virology* **1998**, *246*, 392-399, doi:10.1006/viro.1998.9207.
15. Herbst, L.; Ene, A.; Su, M.; Desalle, R.; Lenz, J. Tumor outbreaks in marine turtles are not due to recent herpesvirus mutations. *Current Biology* **2004**, *14*, R697-R699.
16. Lu, Y.; Aguirre, A.A.; Work, T.M.; Balazs, G.H.; Nerurkar, V.R.; Yanagihara, R. Identification of a small, naked virus in tumor-like aggregates in cell lines derived from a green turtle, *Chelonia mydas*, with fibropapillomas. *Journal of Virological Methods* **2000**, *86*, 25-33, doi:[http://dx.doi.org/10.1016/S0166-0934\(99\)00175-5](http://dx.doi.org/10.1016/S0166-0934(99)00175-5).
17. Alfaro-Nunez, A.; Bertelsen, M.F.; Bojesen, A.M.; Rasmussen, I.; Zepeda-Mendoza, L.; Olsen, M.T.; Gilbert, M.T.P. Global distribution of Chelonid fibropapilloma-associated herpesvirus among clinically healthy sea turtles. *BMC Evolutionary Biology* **2014**, *14*, doi:10.1186/s12862-014-0206-z.
18. Lackovich, J.K.; Brown, D.R.; Homer, B.L.; Garber, R.L.; Mader, D.R.; Moretti, R.H.; Patterson, A.D.; Herbst, L.H.; Oros, J.; Jacobson, E.R. Association of herpesvirus with fibropapillomatosis of the green turtle *Chelonia mydas* and the loggerhead turtle *Caretta caretta* in Florida. *Diseases of aquatic organisms* **1999**, *37*, 89-97.
19. Lu, Y.; Wang, Y.; Aguirre, A.; Zhao, Z.; Liu, C.; Nerurkar, V.; Yanagihara, R. RT-PCR detection of the expression of the polymerase gene of a novel reptilian herpesvirus in tumor tissues of green turtles with fibropapilloma. *Archives of virology* **2003**, *148*, 1155-1163.
20. Nigro, O.; Aguirre, A.A.; Lu, Y. Nucleotide sequence of an ICP18. 5 assembly protein (UL28) gene of green turtle herpesvirus pathogenically associated with green turtle fibropapilloma. *Journal of virological methods* **2004**, *120*, 107-112.
21. Nigro, O.; Yu, G.; Aguirre, A.A.; Lu, Y. Sequencing and characterization of the full-length gene encoding the single-stranded DNA binding protein of a novel Chelonian herpesvirus. *Archives of virology* **2004**, *149*, 337-347.
22. Page-Karjian, A.; Norton, T.M.; Ritchie, B.; Brown, C.; Mancía, C.; Jackwood, M.; Gottdenker, N.L. Quantifying chelonid herpesvirus 5 in symptomatic and asymptomatic rehabilitating green sea turtles. *Endangered Species Research* **2015**, *28*, 135-146, doi:10.3354/esr00687.
23. Page-Karjian, A.; Torres, F.; Zhang, J.; Rivera, S.; Diez, C.; Moore, P.A.; Moore, D.; Brown, C. Presence of chelonid fibropapilloma-associated herpesvirus in tumored and non-tumored green turtles, as detected by polymerase chain reaction, in endemic and non-endemic aggregations, Puerto Rico. *SpringerPlus* **2012**, *1*, 1-8, doi:10.1186/2193-1801-1-35.
24. Quackenbush, S.L.; Aguirre, A.A.; Spraker, T.R.; Horrocks, J.A.; Vermeer, L.A.; Balazs, G.H.; Casey, J.W.; Casey, R.N.; Murcek, R.J.; Paul, T.A., et al. Quantitative analysis of herpesvirus sequences from normal tissue and fibropapillomas of marine turtles with real-time PCR. *Virology* **2001**, *287*, 105-111, doi:10.1006/viro.2001.1023.
25. Rodenbusch, C.R.; Baptistotte, C.; Werneck, M.R.; Pires, T.T.; Melo, M.T.D.; de Ataíde, M.W.; Testa, P.; Alieve, M.M.; Canal, C.W. Fibropapillomatosis in green turtles *Chelonia mydas* in



- Brazil: characteristics of tumors and virus. *Diseases of aquatic organisms* **2014**, *111*, 207-217, doi:10.3354/dao02782.
26. Yu, Q.; Hu, N.; Lu, Y.; Nerurkar, V.R.; Yanagihara, R. Rapid acquisition of entire DNA polymerase gene of a novel herpesvirus from green turtle fibropapilloma by a genomic walking technique. *Journal of virological methods* **2001**, *91*, 183-195.
  27. Yu, Q.; Lu, Y.; Nerurkar, V.; Yanagihara, R. Amplification and analysis of DNA flanking known sequences of a novel herpesvirus from green turtles with fibropapilloma. *Archives of virology* **2000**, *145*, 2669-2676.
  28. Jones, K.; Burgess, G.; Budd, A.M.; Huerlimann, R.; Mashkour, N.; Ariel, E. Molecular evidence for horizontal transmission of chelonid alphaherpesvirus 5 at green turtle (*Chelonia mydas*) foraging grounds in Queensland, Australia. *PloS one* **2020**, *15*, e0227268, doi:10.1371/journal.pone.0227268.
  29. Lu, Y.; Wang, Y.; Yu, Q.; Aguirre, A.; Balazs, G.; Nerurkar, V.; Yanagihara, R. Detection of herpesviral sequences in tissues of green turtles with fibropapilloma by polymerase chain reaction. *Archives of virology* **2000**, *145*, 1885-1893.
  30. McGeoch, D.J.; Gatherer, D. Integrating reptilian herpesviruses into the family herpesviridae. *Journal of virology* **2005**, *79*, 725-731.
  31. Page-Karjian, A.; Gottdenker, N.L.; Whitfield, J.; Herbst, L.; Norton, T.M.; Ritchie, B. potential noncutaneous sites of chelonid herpesvirus 5 persistence and shedding in green sea turtles *Chelonia mydas*. *Journal of aquatic animal health* **2017**, *29*, 136-142.
  32. Alfaro-Núñez, A.; Gilbert, M.T.P. Validation of a sensitive PCR assay for the detection of Chelonid fibropapilloma-associated herpesvirus in latent turtle infections. *Journal of virological methods* **2014**, *206*, 38-41.
  33. Ariel, E.; Nainu, F.; Jones, K.; Juntunen, K.; Bell, I.; Gaston, J.; Scott, J.; Trocini, S.; Burgess, G. Phylogenetic variation of chelonid alphaherpesvirus 5 in green turtle (*Chelonia mydas*) populations along the Queensland Coast, Australia. *Journal of Aquatic Animal Health* **2017**, <http://dx.doi.org/10.1080/08997659.2017.1330783>, doi:<http://dx.doi.org/10.1080/08997659.2017.1330783>.
  34. Lawrance, M.F.; Mansfield, K.L.; Sutton, E.; Savage, A.E. Molecular evolution of fibropapilloma-associated herpesviruses infecting juvenile green and loggerhead sea turtles. *Virology* **2018**, *521*, 190-197.
  35. Ugochukwu, I.C.I.; Aneke, C.I.; Idoko, I.S.; Sani, N.A.; Amoché, A.J.; Mshiela, W.P.; Ede, R.E.; Ibrahim, N.D.G.; Njoku, C.I.O.; Sackey, A.K.B. Bovine papilloma: aetiology, pathology, immunology, disease status, diagnosis, control, prevention and treatment: a review. *Comparative Clinical Pathology* **2019**, *28*, 737-745.
  36. Garcês, A.; Pires, I.; Savini, F.; Scagliarini, A.; Gallina, L. Cutaneous Fibropapilloma in a Red Deer (*Cervus elaphus*) Associated with *Cervus elaphus* Papillomavirus in Portugal. *Journal of wildlife diseases* **2020**, *56*, 636-639.
  37. Munday, J.S.; Cullum, A.A.; Thomson, N.A.; Bestbier, M.; McCormack, T.; Julian, A.F. Anal fibropapillomas containing bovine papillomavirus type 2 DNA in two groups of heifers. *New Zealand veterinary journal* **2018**, *66*, 267-271, doi:10.1080/00480169.2018.1479317.
  38. Brown, D.; Lackovich, J.; Klein, P. Further evidence for the absence of papillomaviruses from sea turtle fibropapillomas. *Veterinary record: journal of the British Veterinary Association* **1999**.
  39. Mashkour, N.; Maclaine, A.; Burgess, G.W.; Ariel, E. Discovery of an Australian *Chelonia mydas* papillomavirus via green turtle primary cell culture and qPCR. *J Virol Methods* **2018**, *10.1016/j.jviromet.2018.04.004*, doi:10.1016/j.jviromet.2018.04.004.
  40. LIMPUS, C.J.; REED, P.C. The green turtle, *Chelonia mydas*, in Queensland: population structure in a coral reef feeding area. In *Biology of Australian Frogs and Reptiles*, Grigg, G., Shine, R., Ehmann, H., Eds. Surrey Beatty and Sons: Sydney, Australia, 1985; pp. 47-52.



41. Hou, Y.; Zhang, H.; Miranda, L.; Lin, S. Serious overestimation in quantitative PCR by circular (supercoiled) plasmid standard: microalgal pcna as the model gene. *PloS one* **2010**, *5*, e9545.
42. Kearse, M.; Moir, R.; Wilson, A.; Stones-Havas, S.; Cheung, M.; Sturrock, S.; Buxton, S.; Cooper, A.; Markowitz, S.; Duran, C., et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)* **2012**, *28*, 1647-1649, doi:10.1093/bioinformatics/bts199.
43. Bustin, S.A.; Benes, V.; Garson, J.A.; Hellemans, J.; Huggett, J.; Kubista, M.; Mueller, R.; Nolan, T.; Pfaffl, M.W.; Shipley, G.L., et al. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clinical chemistry* **2009**, *55*, 611-622, doi:10.1373/clinchem.2008.112797.
44. Greenblatt, R.J.; Balazs, G.H.; Casey, J.W.; Work, T.M.; Dutton, P.; Sutton, C.A.; Spraker, T.R.; Casey, R.N.; Diez, C.E.; Parker, D., et al. Geographic variation in marine turtle fibropapillomatosis. *Journal of Zoo and Wildlife Medicine* **2005**, *36*, 527-530, doi:10.1638/04-051.1.
45. Alfaro-Núñez, A.; Bojesen, A.M.; Bertelsen, M.F.; Wales, N.; Balazs, G.H.; Thomas P. Gilbert, M. Further evidence of Chelonid herpesvirus 5 (ChHV5) latency: High levels of ChHV5 DNA detected in clinically healthy marine turtles. *PeerJ* **2016**, *2016*, e2274, doi:10.7717/peerj.2274.
46. Díaz, F. OIE Standard on principles and methods of validation of diagnostic assays for infectious diseases.
47. Sun, Z.; Zhang, R.; Liu, Z.; Liu, C.; Li, X.; Zhou, W.; Yang, L.; Ruan, Q.; Zhang, X. Development of a fluorescence-based multiplex genotyping method for simultaneous determination of human papillomavirus infections and viral loads. *BMC cancer* **2015**, *15*, 1-11.
48. Oyervides-Muñoz, M.A.; Pérez-Maya, A.A.; Sánchez-Domínguez, C.N.; Berlanga-Garza, A.; Antonio-Macedo, M.; Valdéz-Chapa, L.D.; Cerda-Flores, R.M.; Trevino, V.; Barrera-Saldaña, H.A.; Garza-Rodríguez, M.L. Multiple HPV infections and viral load association in persistent cervical lesions in Mexican women. *Viruses* **2020**, *12*, 380.
49. Duan, L.; Du, H.; Wang, C.; Huang, X.; Qu, X.; Shi, B.; Liu, Y.; Zhang, W.; Duan, X.; Wei, L. The application of BMRT-HPV viral load to secondary screening strategies for cervical cancer. *PloS one* **2020**, *15*, e0232117.
50. Manire, C.A.; Stacy, B.A.; Kinsel, M.J.; Daniel, H.T.; Anderson, E.T.; Wellehan, J.F. Proliferative dermatitis in a loggerhead turtle, *Caretta caretta*, and a green turtle, *Chelonia mydas*, associated with novel papillomaviruses. *Veterinary microbiology* **2008**, *130*, 227-237.
51. Herbst, L.H.; Lenz, J.; Van Doorslaer, K.; Chen, Z.; Stacy, B.A.; Wellehan Jr, J.F.X.; Manire, C.A.; Burk, R.D. Genomic characterization of two novel reptilian papillomaviruses, *Chelonia mydas* papillomavirus 1 and *Caretta caretta* papillomavirus 1. *Virology* **2009**, *383*, 131-135, doi:<http://dx.doi.org/10.1016/j.virol.2008.09.022>.
52. Jones, K. Environmental influences on the epidemiology of fibropapillomatosis in green turtles (*Chelonia mydas*) and consequences for management of inshore areas of the Great Barrier Reef. James Cook University, 2019.
53. Lackovich, J.K.; Jacobson, E.R.; Curry, S.S.; Klein, P.A.; Brown, D.R.; Homer, B.L.; Garber, R.L.; Mader, D.R.; Moretti, R.H.; Patterson, A.D., et al. Association of herpesvirus with fibropapillomatosis of the green turtle *Chelonia mydas* and the loggerhead turtle *Caretta caretta* in Florida. *Diseases of aquatic organisms* **1999**, *37*, 89-97, doi:10.3354/dao037089.
54. Ariel, E.; Nainu, F.; Jones, K.; Juntunen, K.; Bell, I.; Gaston, J.; Scott, J.; Trocini, S.; Burgess, G.W. Phylogenetic Variation of Chelonid Alpha herpesvirus 5 (ChHV5) in Populations of Green Turtles *Chelonia mydas* along the Queensland Coast, Australia. *Journal of Aquatic Animal Health* **2017**, *29*, 150-157, doi:10.1080/08997659.2017.1330783.
55. Greenblatt, R.J.; Work, T.M.; Balazs, G.H.; Sutton, C.A.; Casey, J.W.; Casey, R.N. The *Ozobranchus leech* is a candidate mechanical vector for the fibropapilloma-associated turtle

- herpesvirus found latently infecting skin tumors on Hawaiian green turtles (*Chelonia mydas*). *Virology* **2004**, 321, 101-110, doi:10.1016/j.virol.2003.12.026.
56. Smith, C.J.; Osborn, A.M. Advantages and limitations of quantitative PCR (Q-PCR)-based approaches in microbial ecology. *FEMS microbiology ecology* **2009**, 67, 6-20, doi:10.1111/j.1574-6941.2008.00629.x.
57. Guidry, J.T.; Scott, R.S. The interaction between human papillomavirus and other viruses. *Virus research* **2017**, 231, 139-147, doi:10.1016/j.virusres.2016.11.002.
58. Cruz, D.; Rodríguez, M.; Kouri, V.; Soto, Y.; Zamora, L.; Rodríguez, D.; Barrera, M.; Rehtanz, M. Concurrent papillomavirus- and herpesvirus-infection in Atlantic bottlenose dolphins (*Tursiops truncatus*) inhabiting the Cuban coast. *Marine mammal science* **2014**, 30, 1564-1572, doi:10.1111/mms.12124.
59. Kullander, J.; Handisurya, A.; Forslund, O.; Geusau, A.; Kirnbauer, R.; Dillner, J. Cutaneous human papillomavirus 88: remarkable differences in viral load. *International journal of cancer* **2008**, 122, 477-480.
60. Antonsson, A.; Forslund, O.; Ekberg, H.; Sterner, G.; Hansson, B.G. The ubiquity and impressive genomic diversity of human skin papillomaviruses suggest a commensalic nature of these viruses. *Journal of virology* **2000**, 74, 11636-11641.