

Article

Feasibility of Virtual Research on the Psychological Status and Group Exercises among Community-Dwelling Older Adults during the COVID-19 Pandemic

Amirah Ibrahim ¹, Mei Chan Chong ², Selina Khoo ³, Li Ping Wong ⁴, Ivy Chung ⁵ and Maw Pin Tan ^{1,6,7*}

¹ Ageing and Age-Associated Disorders Research Group, University of Malaya; ami.fatin@gmail.com

² Department of Nursing Science, Faculty of Medicine, University of Malaya

³ Centre for Sport and Exercise Sciences, University of Malaya

⁴ Department of Social and Preventive Medicine, Faculty of Medicine, University of Malaya

⁵ Department of Pharmacology, Faculty of Medicine, University of Malaya

⁶ Department of Medicine, Faculty of Medicine, University of Malaya

⁷ Department of Medical Sciences, Faculty of Healthcare and Medical Sciences, Sunway University

* Correspondence: mptan@ummc.edu.my

Abstract: Social isolation magnified by the restriction of movement order during the COVID-19 pandemic may lead to negative psychosocial health impact among community-dwelling older adults. We evaluated the feasibility of virtual research methods and virtual group exercises among individuals aged 60 years and over in Malaysia. Participants were recruited from the Promoting Independence in Seniors with Arthritis (PISA) pilot cohort through social media messages. A four-week course of virtual group exercise was offered. Anxiety and depression were assessed with the Hospital Anxiety and Depression Scale (HADS) during the last attended follow-up of the cohort study (pre-pandemic), pre-intervention, and post-intervention. Exercise adherence was recorded using diary with daily entries and attendance to the virtual group exercise sessions were captured daily. The outcomes of interest were changes of anxiety and depression scores from baseline to pre-intervention (pandemic-related) and post-intervention (virtual exercise related). Forty-three individuals were recruited. A significant increase in anxiety scores from baseline to pre-intervention was observed. Comparisons using repeated-measures analysis of variance between those who attendance ≥ 14 and < 14 group exercise sessions revealed no between-within subjects differences in depression scores. There was a 23% dropout rate in the post intervention survey and 60.5% of diaries were returned. This study demonstrated the feasibility of conducting research entirely virtually among older persons during the peak of global first wave of a pandemic. The pandemic has led to increased anxiety among community-dwelling older adults.

Keywords: Older adults, Physical exercise, Virtual exercise, COVID-19

1. Introduction

Social isolation is associated with increased risk of poor health and death, with the magnitude of increase in risk reported to be as high as cigarette smoking [1]. However, in order to curb the spread of the Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), the virus which is responsible for the current Coronavirus Disease 2019 (COVID-19), many countries have had to resort to variable degrees of lockdown measures. Even when lockdowns are relaxed, social distancing practices are still advocated as the new norm until a vaccine finally becomes available [2]. In Malaysia, individuals aged 60 years and above make up an estimated 60% of all deaths, further fuelling government recommendations for older persons to stay at home as much as possible, relying on the

community to provide for them. The country continues to observe various degrees of restrictions in response to changes in the number of new cases, with the return to the Conditional Movement Control Order (CMCO) from the Recovery Movement Control Order occurring in many states in response to the third wave of the pandemic in Malaysia [3].

The first case of COVID-19 entered Malaysia when tourists from Wuhan crossed the Singapore-Malaysia border [4]. Strict testing, isolation and contact tracing procedures ensured containment until a super-spreader event occurred in March 2020 during a religious mass gathering which led to a Movement Control Order (MCO) on the 18th of March, 2020 under the Prevention and Control of Infectious Diseases Act 1988 and the Police Act 1967 [5]. The MCO prohibited all travel in and out of the country, closing all schools and businesses apart from essential services, limiting interstate travel and confining all residents indoors apart from for accessing essential services [5]. The police, with the assistance of military personnel, guarded and set-up roadblocks across borders between districts and cordoned off several localized areas with exceptionally high transmission rates in an effort to restrict travel [6,7]. Public announcements were made through loudspeakers from police patrol cars and public places advising people to stay inside their own home or within their own compound.

While these measures were effective in controlling viral transmission, it is expected to have a negative impact on the psychosocial health of the community [8-10]. The heightened risk of severe COVID-19 illness for older persons further increased fear among older people. Therefore, community-dwelling older people living in urban areas stayed at home and became socially isolated as a result of both pandemic response measures as well as anxiety-induced self-imposed isolation. We, therefore, evaluated the feasibility of online survey data collection and a virtual exercise intervention among individuals aged 60 years and over in Malaysia, an upper middle-income developing nation in South-East Asia.

2. Materials and Methods

2.1. Sample population

The Caring For Mental Health During the Covid Pandemic (CaMHeP) study was initiated as part of a university-wide initiative to mitigate potential psychological effects of the pandemic. Participants were recruited from a pre-existing pilot longitudinal cohort study, Promoting Independence in our Seniors with Arthritis (PISA) which was recruited in 2015 (Figure 1). A total of 251 individuals aged 55 years and over with and without osteoarthritis were recruited and followed-up annually, with the final follow-up appointment occurring in 2019. Therefore, up to four follow-up visits were achieved per participant. Ethics approval had been obtained prior and written informed consent was obtained from participants prior to recruitment (UM.TNC2/UMREC - 622). Individuals who were unable to perform physical exercise, had ongoing untreated heart conditions, and those unable to participate in telephone or video interviews were excluded.

2.2. Recruitment Procedure

The researcher communicated with participants via social media messaging (WhatsAppTM, Facebook Inc., USA), telephone calls and video conferencing (Google MeetTM, Google Inc., USA). Participants' information was extracted from the PISA database and those with mobile phone numbers were approached first via a WhatsApp bulk messaging application with text invitation and a short teaser video. Participants who replied to the text or simply replied "yes" were contacted via telephone calls to provide further detailed explanations. Details provided included information on exercises on offer and a link to complete the mental health assessment and Google MeetTM installation. Participants who needed hard copies of the materials were identified and the material was delivered via courier. Participants were included in a conditional WhatsappTM Group chat, in which only the researcher had access to post to the group. Participants were advised to send message privately to the researcher for any queries.




Figure 1. CONSORT flow diagram

2.3. Measures

Mental well-being was measured at baseline (pre-pandemic), at recruitment to the intervention study, and again at the end of the exercise intervention. Baseline scores were obtained from participants' last visit recorded between one to two years prior to this study. The pre and post-intervention scores were assessed between May and June 2020. Additional measures obtained from the PISA dataset were physical activity, social participation, and social network.

2.3.1 Mental Health

Mental well-being was measured using the Hospital Anxiety and Depression Scale (HADS) [11]. HADS was specifically designed for non-psychiatric practice as a self-assessment mood scale. The short time (2 to 5 minutes) required to complete the scale [12] is an advantage. The scale has 14 items related to anxiety and depression divided equally. The scale scores range from 0 to 21 for each sub-scale, which in this study were analyzed separately. For each sub-scale, the score of 0 to 7 is regarded as normal, 8 to 10 is suggestive of the presence of a mood disorder, and higher than 11 indicates the probable presence of a mood disorder [12].

2.3.2. Physical Activity

To measure physical activity level, the International Physical Activity Questionnaire – Short Form (IPAQ-SF) was administered. The IPAQ-SF is a self-administered questionnaire that evaluated physical activity in the last seven days. It comprised seven items that enquired about the presence and time spend performing vigorous activities, performing moderate activities, walking and sitting. The levels of physical activity (low, medium and high) were then determined using a published analysis protocol [13,14].

2.3.3. Social Participation

To measure the occurrence of participation restriction, the Keele's Assessment of Participation (KAP) was used. KAP is an 11-item self-administered questionnaire. Four levels of restrictions (no restrictions, minimal, moderate and substantial) are then identified available published cut-offs [15].

2.3.4. Social Network

The Lubben Social Network Scale (LSNS-6) was used to evaluate participants' social network. Total scores ranged from 0 to 30 [16]. Participants with scores below 12 were considered at risk of social isolation.

2.4. Intervention

A quasi-experimental study comparing two virtual exercise groups determined through number of attendances to the virtual exercise sessions was conducted (Figure 2). All participants were invited to take part in a four-week course of daily virtual group exercises. Participants with virtual group exercises attendance of 14 or more were included in Group A while those attended fewer than 14 sessions were included in Group B. The cut-off of 14 session was determined using the median attendance level. Exercise adherence was recorded using a daily exercise diary. Both the booklet and virtual exercises focused on muscle strengthening by gradually increasing repetitions, coordination and fine motor training. The study protocol had been registered with the National Malaysian Research Registry (NMRR-20-1629-55960). The four-week virtual exercise was tailor-designed for older persons by the authors and a physiotherapist. Recruited participants were invited to take part in a 30-minute daily virtual exercise class. The installing instructions of Google Meet were sent via text and video demonstration. A 2-hour trial run or rehearsal session was set up two days before the actual start of the program. The researcher continued to provide technical support from the time of the trial run up to the start of the program and throughout all exercise sessions through the social media messaging group set up for this purpose.

The virtual exercise class started with deep breathing and correction of posture. Exercises for neck, chest, shoulder, upper back, lower back, abdomen, thigh and ankle rotation were conducted in the seated position. Lower back and abdominal strengthening exercise were conducted in the supine position. Hip abduction and adduction, calf and ankle exercises were then conducted in the standing position. Coordination and fine motor training included face muscles exercise, fingers movement exercise, alternating hand exercise, and towel toe curl. Additionally, ankle alphabet exercise were performed to improve flexibility. The sessions ended with deep breathing and a reminder to fill-in their exercise diary. The range of exercises explained above were performed alternately and gradually increments in repetitions which were limited largely by time availability.

Participants who were unable to attend the virtual group exercise, were encouraged to conduct individual home-based exercises guided by an electronic exercise booklet and video. The "PISA exercises for older adults" video and booklet had been developed in 2016 as part of a public engagement activity of the original research program. Those who did not participate in virtual exercises were reminded to exercise daily and to fill-in their diary, either via the WhatsApp™ group chat or private messages.

2.4.2. Adherence

Adherence to both virtual groups and home-based individual exercises were measured using self-administered exercise diaries with daily entries. This collected information on which days of the week the participant had conducted their exercises. Exercise diaries were either sent out electronically or through couriers as per participant preference. In addition, attendances were taken for each virtual group exercise sessions as an additional measure of adherence.



Figure 2. Study framework

2.5. Data analysis

Data analysis was performed using the IBM SPSS Statistics 21 and a p value of less than .05 was considered statistically significant. Descriptive statistics were presented in frequencies with percentages and mean with standard deviation. The mixed between-within subjects ANOVA were used to investigate the effect of time (baseline, pre and post intervention), intervention and interactions between time and groups. Missing data were analyzed using Little's MCAR test and multiple imputation were performed.

3. Results

3.1. Recruitment

Two hundred and fifty-one PISA participants were approached via the social media messaging application WhatsApp™, of whom 45 agreed to participate with two dropouts before the exercise program started (Figure 1). A 23% drop-out was recorded for the online post-intervention survey. Two reminders were sent out via WhatsApp™ in group and personal text messages.

3.2. Groups comparison

The two groups were homogenous in years-of-education, gender, ethnicity, marital status, person(s) dwelling with participants, self-rated health condition and health compared to others ($p > 0.05$). The mean age for participants in Group A were significantly higher than those in Group B. Participants in Group B had lower self-rated health compared to others (Table 1).

Table 1. Demographic characteristics of participants in Group A and Group B.

Variable	Mean \pm Standard Deviation		p -Value			
	Group A (n=21) (14 or >14 sessions attendance to virtual exercises)	Group B (n=22) (<14 sessions attendance to virtual exercises)				
Age (years)	65.1 \pm 4.81	61.8 \pm 5.15	Df = 41 $t = 2.16$ $p = .037^*$			
Years of education	15.79 \pm 2.42	14.48 \pm 2.92	Df = 38 $t = 1.54$ $p = .132$			
Attendance to online sessions	12.43 \pm 2.16	4.36 \pm 4.44	Df = 41 $t = 11.4$ $p < .001^*$			
Variable	Sub-variable	Group A		Group B		p -Value
		n	%	n	%	
Gender	Male	4	19	6	28.6	$p = .717$

	Female	17	81	15	71.4	
Ethnicity	Malay	3	14.3	2	9.1	$p = .542$
	Chinese	17	81	15	68.2	
	Indian	0	0	5	22.7	
	Others	1	4.8	0	0	
Marital status	Single	1	4.8	3	13.6	$p = .155$
	Married	19	90.5	15	68.2	
	Widowed	1	4.8	2	9.1	
	Partner	0	0	2	9.1	
	Divorced	0	0	2	9.1	
Dwelling	Independent	0	0	7	31.8	$p = .070$
	Spouse	17	81	11	50	
	Children	3	14.3	4	18.2	
	Parent	1	4.8	0	0	
Health condition (self-rated)	Poor	0	0	2	9.1	$p = .085$
	Fair	2	9.5	6	27.3	
	Good	14	66.7	12	54.5	
	Very Good	5	23.8	2	9.1	
Health compared to others	Poor	0	0	1	4.5	$p = .002^*$
	Fair	0	0	9	40.9	
	Good	11	52.4	7	31.8	
	Very Good	9	42.9	4	18.2	
	Excellent	1	4.8	1	4.5	

A paired-samples t-test was conducted to evaluate the impact of the pandemic on anxiety and depression scores of all participants. There was a statistically significant increase in mean anxiety scores from baseline ($M = 3.5$, $SD = 3.5$) to pre-intervention ($M = 6.02$, $SD = 3.97$, $t(41) = -3.75$, $p = .001$). No significant mean difference in depression scores from baseline ($M = 4.98$, $SD = 3.92$) to pre-intervention ($M = 5.57$, $SD = 3.66$, $t(41) = -1.02$, $p = .31$).

A complete case, mixed between-within subjects analysis of variance showed no difference between groups and across time in anxiety scores (Wilks' Lambda = .96, $F(2, 30) = .61$, $p = .55$, partial eta squared = .04), and depression scores (Wilks' Lambda = .89, $F(2, 30) = 1.80$, $p = .18$, partial eta squared = .11). With multiple imputation on post-intervention anxiety scores, there was a substantial main effect for time ($p < .005$) suggesting a significant increase in anxiety scores across time between baseline, pre- and post-intervention. Post-hoc test (Bonferroni) showed significant increase in anxiety scores between baseline anxiety scores and both pre- and post-intervention. There was no significant difference between before and after intervention in anxiety scores (Supplementary Table S1).

Depression scores do not show any significant differences across time or between groups for both original and imputed data analysis. Changes in anxiety and depression scores over time for all participants and the two intervention groups are also summarized in Figure 3 and Figure 4.




Figure 3. Anxiety scores at baseline, pre and post intervention.




Figure 4. Depression scores at baseline, pre and post intervention.

3.3. Adherence

3.3.1. Attendance

Overall mean attendance was 10.26 (SD=7.02) sessions. Mean attendance for Group A was 16.43 (SD 2.16) sessions, minimum attendance was 14 and maximum attendance was 20 sessions. Most participants attended at least 14 sessions (n=8) and one attended all the 20 sessions. Individuals in Group B had mean attendance of 4.36 (SD= 4.44) sessions.

3.3.2. Exercise Diary

60.5% (n=26) participants returned their exercise diaries. 85.7% (n=18) of participants in Group A returned the exercise diary compared to 36.4% (n=8) from participants in Group B ($p=.002$) (Table

2). There was a significant difference in mean attendance between participants in both groups who returned and did not returned the diary $t(41)=4.79, p<.001$, two-tailed. No difference in age, years of education, gender, HADS score at baseline, physical activities, co-morbidities, KAP and LSNS-6 ($p > .05$) between participants who returned and did not returned the diary.

On a sub-group analysis of both groups, there was no significance difference in age, years of education, gender, HADS score at baseline, physical activities, co-morbidities, KAP and LSNS-6 ($p > .05$). In Group B, there was a significance mean difference in attendance for participants who returned ($M = 6.88, SD = 4.94$) and did not return the diary ($M = 2.93, SD = 3.54$) ($t(20) = 2.18, p = .041$).

Table 2. Attendance summary

Variables	All	Group A (n=21) (14 or >14 sessions attendance to virtual exercises)	Group B (n=22) (<14 sessions attendance to virtual exercises)
Number of returned diaries	26	18	8
Total exercises sessions attended	666	464	202
Total exercise attended per person	25.6	25.8	25.3
Total virtual group exercises	441	345	96
Total virtual group exercises per person	10.3	19.2	12
Total individual home exercises	225	119	106
Total individual home exercises per person	8.7	6.6	13.3

Only 5 out of 26 participants (19.2%) recorded the same amount of attendance in their diary as captured by Google Meet attendance. 20 (76.9%) participants recorded higher number of attendances in their diary compared to captured by Google Meet attendance. A Cohen's kappa agreement analysis between Google Meet captured attendance and diary attendance for virtual exercises session showed slight agreement, $\kappa = .137, p = .004$.

3.3.3. Technical issues

Of the 45 recruited participants, 10 individuals successfully joined the initial trial run which occurred two days before the start of the program. Text messages (n=9) and phone calls (n=6) were received between the two days on issues including inability to download the application, unable to see or turn on video/audio, forgetting the time of the session, and device memory is full. Queries were attended on an individual basis. A further three trial runs were conducted separately in between the two days for three individuals. Two participants used social messaging platform using their computers. Real-time step-by-step for demonstration of the web browsing procedure was helpful. Text or voice instructions alone were insufficient. The use of two devices (laptop and smart phone) at the same time was essential for the researcher to this program. Participants were mostly considerate and avoided contacting the researcher in the evenings. The researcher established ground rules and only responded to queries from 7am until 9pm. Due to the overwhelming number of calls and texts, a separate number was established. To ensure authenticity, the social media messaging group icon displayed the project logo.

4. Discussion

Compared to baseline data, participants had increased anxiety scores pre-intervention, which coincided with the first movement control order of the COVID-19 pandemic. Following the intervention phase, anxiety scores remained. No difference in depression was observed between the three time points in both groups. Around 20% of our overall PISA cohort were willing to participate in the intervention study with an eventual dropout rate of 23%. Individuals attended over 50% of the 20 virtual exercise sessions on offer.

This study was conducted at the peak of the first wave of the COVID-19 pandemic worldwide. Growing anxiety among older people is likely to be linked to the higher mortality rates recorded among people aged 60 years and older from COVID-19. A study conducted in the UK found that 12.8% of participants reporting worse depression scores while 12.3% felt worse on the anxiety scores using the HADS scale [17]. On the contrary, our participants did not demonstrate any deterioration in depression scores. Both studies, nevertheless, has clearly demonstrated a worrying trend towards increase in psychological symptoms associated with the COVID-19 pandemic among older adults.

Low recruitment rates for this study had occurred as the recruitment strategy was through social media messaging in view of time and human resource constraints. Some cohort participants felt they were unable to participate due to their arthritis or difficulty accessing any internet connection, while many did not respond to our messages despite reminders. Those who agreed to participate, however, demonstrated an acceptable dropout. In face to face exercise intervention studies, dropout rates of 14% to 31.9% have been reported [18-21] while online dropout rates of 7.5% [22] in studies with sample less than 200 and a large study (n=36,373) recorded 48% non-adherence [23]. Adherence was 93% for a telephone-guided exercise program [24] and 70% for community-based face-to-face programs [25].

As this study was primarily intended to evaluate the feasibility of virtual studies within an older population in a developing country, process evaluation is more important than outcomes for which this study was not powered for. There is a perception that older people are unable to use technology and often pester younger family members and acquaintances for help with gadgets. This may be reflected upon by the low recruitment rate of 45 people out of 251 who responded to social media messaging invitations. There were no comparison groups from younger populations from which such a conclusion may be drawn. However, social media messaging apps are widely used among older persons. The recruitment strategy only employed one social media messaging platform which is the most popular platform, therefore, some older persons may have used alternative platforms. Nevertheless, this study has demonstrated that it is possible to recruit through social media messaging. In addition, those who responded and participated were able to operate gadgets/devices independently, indicating that such a strategy does work for at least a segment of the older population in our setting. Prior research work on social media use in research recruitment has centred on young adults or hard-to-reach populations such as those with HIV [26-28]. Low accrual rates are in fact expected for social media recruitment and it is recommended that multiple platforms are used. However, as this study employed an existing dataset with availability of mobile phone numbers, it was still possible to use social media messaging on a solitary commonly used platform to recruit.

The inconsistency of recorded attendance between participants' diary against researcher's captured Google Meet™ attendance could be due to several personal and technical issues. Participants were inclined to fill in diaries during the final day or week of the exercise program, once researcher asked for the diaries to be returned. The researcher provided daily reminders after each virtual session to fill in the diaries daily. There were participants who waited several days of exercises before bundling their entries. The researcher used Google Meet™ attendance Chrome extension by claycodes.org, the Beta version, which may have had technical glitches in collecting participants' attendance. Nevertheless, we were able to demonstrate differences in self-reported and objective recording of attendance, cautioning against the accuracy of self-administered exercise diaries in future studies.

The process of preparing the participants for the online program is time consuming which has replaced logistics issues associated with face-to-face programs. The older participants needed time to

process new information on the use of online applications to conduct exercise session. Prior to the intervention, only a handful had used other video conferencing platforms for communication, and none has been using Google Meet™ which was the platform selected here. Therefore, intensive support from the researcher was required to enable the participants who signed up to eventually successfully join the virtual exercise sessions. Participants were not able to sign on to the video conferencing platforms using instruction provided through social media text messaging alone or recorded voice instructions alone. Instead support using telephone instructions and video calls with social media applications were used. Nevertheless, once the older persons had mastered the procedures required, they were able to continue to log in to the platform using the link provided for the rest of the study. Prior pre-pandemic efforts at engaging older persons in telehealth using videoconferencing has yielded mixed results [29]. Movement restrictions incurred throughout the pandemic and the need to shield our older population has, however, limited the older person's choice, and therefore potentially altered their determination to engage through videoconferencing. Of note, not only were the older adults able to engage with virtual exercises using videoconferencing, they were also able to self-administer the pre-intervention and post-intervention questionnaires through an online platform which was deliberately limited to just the HADS to enhance completion rates, since older adults were expected to encounter technical issues with online questionnaire completion and were expected to require a longer time and some assistance to self-administer the questionnaire online compared to previous printed versions, which they frequently required assistance to complete [30].

This study is limited by the non-randomized allocation to groups. It was a pragmatic decision, since the priority was to test the virtual assessment and interventions, and potential communication issues associated with virtual contact precluded the possibility of detailed explanations that is necessary before randomization can proceed. Allocation of groups to the participants may hinder their motivation for those who will not be chosen to be in the virtual group, as our main aim is to have as many older persons involved in virtual meetings. Therefore, the decision was to allow recruitment through convenience sampling and for individuals to select their preferred mode of exercise. However, this has resulted in contamination between the groups, because everyone has the links to access the virtual exercise groups. This suggests that individual randomization in future such studies may experience the same issues, and hence cluster randomization with safeguards against contamination will be necessary. Ultimately the study was able to demonstrate the feasibility of conducting virtual interventions and data collection in a developing nation setting where the older population may have limited information technology literacy, as well as highlight potential limitations in recruitment, conduct and representativeness of such a study. Nevertheless, during a pandemic when older adults would otherwise be completely excluded due to the need to physical distance, shield and obey lockdown measures, conducting online surveys and virtual interventions will at least allow some inclusion.

5. Conclusions

Our study has demonstrated the feasibility of collecting data using online surveys and delivering interventions through videoconferencing among individuals aged 60 years and over in a developing nation. The delivery of the intervention, however, requires investment in training and extensive technical support which needs to be delivered through a variety of methods. We also established significant increased anxiety symptoms during the peak of the first wave globally when lockdown measures were in place. While limitations exist with regards to recruitment and treatment allocation, it is still important to include older adults in clinical research using virtual communications, to ensure that older adults are not completely excluded from ongoing studies.

Supplementary Materials: Table S1: Comparison of anxiety and depression scores at baseline (pre-pandemic), before and after intervention (post-pandemic) between groups.

Author Contributions: Conceptualization, M.P.T., and I.C.; methodology, A.I., C.M.C., W.P.L., I.C., and M.P.T.; formal analysis, A.I., and M.P.T.; investigation, A.I., and S.K.; resources, A.I., S.K., C.M.C., L.P.W., and I.C.;

writing—original draft preparation, A.I., and M.P.T.; writing—review and editing, C.M.C., S.K., L.P.W., and I.C.; supervision, C.M.C., and M.P.T.; funding acquisition, L.P.W., and I.C. All authors have read and agreed to the published version of the manuscript.”

Funding: This research was funded by University of Malaya, UM Grand Challenge, GC002C-14 HTM and the UM COVID grant, RG564-2020HWB.

Acknowledgments: We would like to express our appreciation to other investigators and researchers involved in the CAMHEP study. Special thanks to Sun Zek Jong who assisted with exercise sessions.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Holt-Lunstad, J.; Smith, T.B.; Layton, J.B. Social relationships and mortality risk: A meta-analytic review. *Public Library of Science*: 2010; Vol. 7.
- World Health, O. Coronavirus disease (COVID-19) advice for the public. 2020.
- MALAYSIAN NATIONAL SECURITY COUNCIL. SOP PERINTAH KAWALAN PERGERAKAN. Available online: <https://www.mkn.gov.my/web/ms/sop-pkp-pemulihan/> (accessed on 1 Dec 2020).
- Abdullah, J.M.; Wan Ismail, W.F.N.M.; Mohamad, I.; Ab Razak, A.; Harun, A.; Musa, K.I.; Lee, Y.Y. A critical appraisal of COVID-19 in Malaysia and beyond. *Malaysian Journal of Medical Sciences* **2020**, *27*, 1-9, doi:10.21315/mjms2020.27.2.1.
- Prime Minister's Office of, M. The Prime Minister's Special Message on COVID-19 - 16 March 2020 - Prime Minister's Office of Malaysia. 2020.
- Prime Minister's Office of, M. Kenyataan Media MKN: Perincian Perintah Kawalan Pergerakan - Prime Minister's Office of Malaysia. 2020.
- Prime Minister's Office of, M. Kenyataan Media: Penguatkuasaan PKP Dengan Bantuan ATM - Prime Minister's Office of Malaysia. 2020.
- Delmastro, M.; Zamariola, G. The Psychological Effect of COVID-19 and Lockdown on the Population: Evidence from Italy. *researchsquare.com* **2020**, 10.21203/RS.3.RS-82494/V1, doi:10.21203/RS.3.RS-82494/V1.
- Pandey, D.; Bansal, S.; Goyal, S.; Garg, A.; Sethi, N.; Pothiyill, D.I.; Sreelakshmi, E.S.; Sayyad, M.G.; Sethi, R. Psychological impact of mass quarantine on population during pandemics—The COVID-19 Lock-Down (COLD) study. *PLOS ONE* **2020**, *15*, e0240501, doi:10.1371/journal.pone.0240501.
- Sugaya, N.; Yamamoto, T.; Suzuki, N.; Uchiumi, C. A real-time survey on the psychological impact of mild lockdown for COVID-19 in the Japanese population. *Scientific Data* **2020**, *7*, 1-6, doi:10.1038/s41597-020-00714-9.
- Zigmond, A.S.; Snaith, R.P. The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica* **1983**, *67*, 361-370, doi:10.1111/j.1600-0447.1983.tb09716.x.
- Snaith, R.P. The hospital anxiety and depression scale. *Health and Quality of Life Outcomes* **2003**, *1*, 29-29, doi:10.1186/1477-7525-1-29.
- Chu, A.H.Y.; Moy, F.M. Reliability and validity of the malay international physical activity questionnaire (IPAQ-M) among a malay population in Malaysia. *Asia-Pacific Journal of Public Health* **2015**, *27*, NP2381-NP2389, doi:10.1177/1010539512444120.
- Ekelund, U.; Sepp, H.; Brage, S.; Becker, W.; Jakes, R.; Hennings, M.; Wareham, N.J. Criterion-related validity of the last 7-day, short form of the International Physical Activity Questionnaire in Swedish adults. *Public Health Nutrition* **2006**, *9*, 258-265, doi:10.1079/phn2005840.

15. Wilkie, R.; Peat, G.; Thomas, E.; Hooper, H.; Croft, P.R. The Keele Assessment of Participation: A new instrument to measure participation restriction in population studies. Combined qualitative and quantitative examination of its psychometric properties. *Quality of Life Research* **2005**, *14*, 1889-1899, doi:10.1007/s11136-005-4325-2.
16. Lubben, J.; Blozik, E.; Gillmann, G.; Iliffe, S.; Von Kruse, W.R.; Beck, J.C.; Stuck, A.E. Performance of an abbreviated version of the lubben social network scale among three European community-dwelling older adult populations. *Gerontologist* **2006**, *46*, 503-513, doi:10.1093/geront/46.4.503.
17. Robb, C.E.; de Jager, C.A.; Ahmadi-Abhari, S.; Giannakopoulou, P.; Udeh-Momoh, C.; McKeand, J.; Price, G.; Car, J.; Majeed, A.; Ward, H., et al. Associations of Social Isolation with Anxiety and Depression During the Early COVID-19 Pandemic: A Survey of Older Adults in London, UK. *Frontiers in Psychiatry* **2020**, *11*, 991-991, doi:10.3389/fpsy.2020.591120.
18. Heinrich, K.M.; Patel, P.M.; O'Neal, J.L.; Heinrich, B.S. High-intensity compared to moderate-intensity training for exercise initiation, enjoyment, adherence, and intentions: An intervention study. *BMC Public Health* **2014**, *14*, doi:10.1186/1471-2458-14-789.
19. Pfaff, J.J.; Alfonso, H.; Newton, R.U.; Sim, M.; Flicker, L.; Almeida, O.P. ACTIVEDEP: A randomised, controlled trial of a home-based exercise intervention to alleviate depression in middle-aged and older adults. *British Journal of Sports Medicine* **2014**, *48*, 226-232, doi:10.1136/bjsports-2013-092510.
20. Schoene, D.; Valenzuela, T.; Toson, B.; Delbaere, K.; Severino, C.; Garcia, J.; Davies, T.A.; Russell, F.; Smith, S.T.; Lord, S.R. Interactive Cognitive-Motor Step Training Improves Cognitive Risk Factors of Falling in Older Adults – A Randomized Controlled Trial. *PLOS ONE* **2015**, *10*, e0145161-e0145161, doi:10.1371/journal.pone.0145161.
21. Shimada, H.; Makizako, H.; Doi, T. Effects of Combined Physical and Cognitive Exercises on Cognition and Mobility in Patients With Mild Cognitive Impairment: A Randomized Clinical Trial Prevalence of frailty in Japan: a systematic review and meta-analysis View project. *Article in Journal of the American Medical Directors Association* **2017**, 10.1016/j.jamda.2017.09.019, doi:10.1016/j.jamda.2017.09.019.
22. Baez, M.; Far, I.K.; Ibarra, F.; Ferron, M.; Didino, D.; Casati, F. Effects of online group exercises for older adults on physical, psychological and social wellbeing: A randomized pilot trial. *PeerJ* **2017**, *2017*, doi:10.7717/peerj.3150.
23. Liu, M.; Miyawaki, C.E. What Types of Physical Function Predict Program Adherence in Older Adults? *Rehabilitation nursing : the official journal of the Association of Rehabilitation Nurses* **2020**, *45*, 279-286, doi:10.1097/rnj.000000000000209.
24. Vallerand, J.R.; Rhodes, R.E.; Walker, G.J.; Courneya, K.S. Feasibility and preliminary efficacy of an exercise telephone counseling intervention for hematologic cancer survivors: a phase II randomized controlled trial. *Journal of Cancer Survivorship* **2018**, *12*, 357-370, doi:10.1007/s11764-018-0675-y.
25. Farrance, C.; Tsofliou, F.; Clark, C. Adherence to community based group exercise interventions for older people: A mixed-methods systematic review. Academic Press Inc.: 2016; Vol. 87, pp 155-166.
26. Arigo, D.; Pagoto, S.; Carter-Harris, L.; Lillie, S.E.; Nebeker, C. Using social media for health research: Methodological and ethical considerations for recruitment and intervention delivery. *DIGITAL HEALTH* **2018**, *4*, 205520761877175-205520761877175, doi:10.1177/2055207618771757.
27. Iribarren, S.J.; Ghazzawi, A.; Sheinfil, A.Z.; Frasca, T.; Brown, W.; Lopez-Rios, J.; Rael, C.T.; Balán, I.C.; Crespo, R.; Dolezal, C., et al. Mixed-Method Evaluation of Social Media-Based Tools and Traditional Strategies to Recruit High-Risk and Hard-to-Reach Populations into an HIV Prevention Intervention Study. *AIDS and Behavior* **2018**, *22*, 347-357, doi:10.1007/s10461-017-1956-6.

28. Topolovec-Vranic, J.; Natarajan, K. The use of social media in recruitment for medical research studies: A scoping review. *Journal of Medical Internet Research*: 2016; Vol. 18.
29. Banbury, A.; Chamberlain, D.; Nancarrow, S.; Dart, J.; Gray, L.; Parkinson, L. Can videoconferencing affect older people's engagement and perception of their social support in long-term conditions management: a social network analysis from the Telehealth Literacy Project. *Health and Social Care in the Community* **2017**, *25*, 938-950, doi:10.1111/hsc.12382.
30. Tan, M.P.; Ho, Y.Y.; Chin, A.V.; Saedon, N.I.I.; Abidin, I.Z.; Chee, K.H.; Khor, H.M.; Goh, C.H.; Hairi, N.N.; Othman, S., et al. Ethnic differences in lifetime cumulative incidence of syncope: the Malaysian elders longitudinal research (MELoR) study. *Clinical Autonomic Research* **2020**, *30*, 121-128, doi:10.1007/s10286-019-00610-2.