

## **Disability, Caste and Intersectionality: Does co-existence of disability and caste compound marginalization for women seeking maternal healthcare in southern Nepal?**

Hridaya Raj Devkota<sup>1</sup>, Andrew Clarke<sup>2</sup>, Emily Murray<sup>3</sup>, Maria Kett<sup>4</sup>, Nora Groce<sup>5</sup>

---

<sup>1</sup>Department of Epidemiology and Public Health, University College London (UCL), UK. 1 – 19 Torrington Place London WC1E 6BT; Email: [hridayadevkota@hotmail.com](mailto:hridayadevkota@hotmail.com) (Corresponding author).

<sup>2</sup> Global Programmes, Save the Children UK, London; Division of health research, Lancaster University, Lancaster, UK; Email: [andrewcws@gmail.com](mailto:andrewcws@gmail.com)

<sup>3</sup> Department of Epidemiology and Public Health, University College London (UCL), UK. 1 – 19 Torrington Place London WC1E 6BT; Email: [emily.murray@ucl.ac.uk](mailto:emily.murray@ucl.ac.uk)

<sup>4</sup> Leonard Cheshire Disability and Inclusive Development Centre, Department of Epidemiology and Public Health, University College London (UCL), UK. 1 – 19 Torrington Place London WC1E 6BT; Email: [m.kett@ucl.ac.uk](mailto:m.kett@ucl.ac.uk)

<sup>5</sup> Leonard Cheshire Disability and University College London (UCL), UK. 1 – 19 Torrington Place London WC1E 6BT; Email: [nora.groce@ucl.ac.uk](mailto:nora.groce@ucl.ac.uk)

## ABSTRACT

**Background:** Disability and caste are two different forms of oppressions, however Dalits and people with disabilities commonly face similar types of marginalities. Dalit women with disabilities may experience double discrimination because of the intersectionality of disability and caste. This study examines whether the disability and caste identity of women together affects and compounds the utilization of maternal healthcare services.

**Methods:** A cross-sectional survey was conducted using a semi-structured questionnaire among a total of 354 Dalit and non-Dalit women, with and without a disability aged between 15 – 49 years. Maternal healthcare service utilization was assessed by Ante-Natal Care (ANC), health facility (HF) delivery and Post-Natal Care (PNC) during the last pregnancy. Logistic regression was performed to detect the predictors of service utilization, and identify whether disability and caste were associated with service utilization. First, disability and caste were fitted separately in models. Secondly, intersectionality of disability and caste was tested by inclusion of disability\*caste interaction term. Finally, the confounding effect of socio-demographic factors was investigated.

**Results:** Out of surveyed women, 73% had 4+ ANC visits, 65% had HF delivery and 29% had a PNC visit during their last pregnancy. Women with a disability had lower odds of HF delivery (OR 0.50, CI 0.30 – 0.84) and PNC (OR 0.47, CI 0.25 – 0.88) than women without a disability. Adjustment for women's age and household wealth explained associations in HF delivery by women with disabilities. There was no association between caste and service utilization. Disability overrode caste and there was no other evidence of effect modification by women's caste status in the utilization. However, a weak interaction effect in the utilization of ANC services was found in the caste group by their education (OR 0.19, 95% CI 0.05 – 0.74).

**Conclusions:** Disabled women – whether Dalit or non-Dalit - had lower rates of utilizing all maternal healthcare services than non-disabled women. However, Dalit women with disabilities were more likely to receive PNC than non-Dalit women with disabilities. Increasing equity in maternal healthcare service utilization requires that traditional approaches to service development and program intervention to be re-examined and more nuanced interventions considered to ensure improved access and outcome among all vulnerable groups.

**Keywords:** Disability, Caste, Intersectionality, Maternal healthcare, Utilization, Nepal

## BACKGROUND

Despite Nepal's good progress in the past two decades to reduce maternal mortality, it remains still high at 238 per 100,000 live births (1). Disparities in health and healthcare access, delays in seeking and obtaining care, and poor quality of care due to lack of skilled health professionals are key contributing factors in maternal deaths (2,3). Studies also show significant differences in the utilization of maternal healthcare services and health outcomes among social groups (4,5). Socially marginalized populations, such as disabled and lower caste groups (Dalit) women are less likely to utilize maternal healthcare services (5–8).

### Disability and Maternal Healthcare Service Utilization

The UN Convention on the Rights of Persons with Disabilities (2006) defines disability as "... those who have long-term physical, mental, intellectual or sensory impairments which in interaction with various barriers may hinder their full and effective participation in society on an equal basis with others" (9). About 15% of the world's population is estimated to have some form of disability; with prevalence higher among females (10). Reported rates of disability in Nepal vary widely. The Nepal census 2011 found less than 2% of the population with disabilities (11) whereas other surveys have reported prevalence rates ranging from 1 to 13% (12) and up to 29% for women with children (6,13).

Around the world, women with disabilities face numerous barriers in accessing maternal healthcare services (14–16). Findings in Nepal are similar to those found in other low and middle-income countries (17,18). Poor quality of care and health worker attitudes, physical barriers, cost and lack of family support as well as poverty all are barriers to accessing maternal healthcare services by disabled women (6,19,20).

### Caste and Maternal Healthcare Service Utilization

Out of 125 ethnic and caste groups in Nepal, the 2010 census listed 28 social groups as Dalits ('untouchables'), comprising 13.6% of the population (11,21). 'Dalits' are at the bottom of the caste hierarchy and the word literally means "oppressed". The caste system and discrimination linked to this – manifests as a rigid form of exploitation and oppression in South Asia that has left some groups, such as the Dalits, marginalized and vulnerable (22). The literature demonstrates that caste has a strong influence on maternal healthcare service utilization including the quality of care (23–25). Dalit women endure worse health than their upper caste counterparts across almost all indicators in Nepal (26). However, one study showed no effect of caste in the use of healthcare services (27). The contradictory findings suggest a more complex picture and that whilst caste is a factor, caste alone may not be the

sole determinant in women's health indicators and in accessing healthcare services; other factors may influence women's healthcare access. Few papers make direct reference to the combined issues of disability and caste. Based on the literature it can be hypothesized that the co-existence of gender discrimination together with caste and disability could have a negative impact on disabled and Dalit women's maternal healthcare experiences.

### **Intersectionality between disability and caste**

Intersectionality is broadly understood here as the intersection between disability and caste identity, influencing individual lives, social practices, institutional arrangements and cultural ideologies with the power outcomes (28) that ultimately impact in the service use by those population groups. The interactions that take place between and within multiple dimensions and power structures, and social systems, lead to various forms of oppressive experiences. Disability and the caste system are two different forms of oppressions in South Asia including in Nepal, however, disabled and Dalit people face many similar types of oppression. Little is known however about the combined effect of disability and caste in the context of maternal health. In this study, we set out to understand whether the combined effect of disability and caste would further compound marginalisation and impact on access to care of disabled women and their new-borns.

This study hypothesized that healthcare services may be least accessible to disabled, lower caste women. It also theorized that higher caste non-disabled and disabled women have greater access to healthcare services because of their higher status in the society, higher level of education, better economic status, and greater participation in and access to community resources. Further, higher caste disabled women's access to healthcare might be lower than that of their non-disabled peers, but would still be better than that of either Dalit women or disabled Dalit women. This has implications for women in Nepal, where different forms of social oppression such as gender inequality, disability and caste continue, despite measures to address these (29,30). No studies to date have explored the intersection of caste and disability in the context of maternal healthcare service access and utilization in Nepal. To better understand the complex relationship between disability and caste in the utilization of maternal healthcare services, the study aims to identify the key predictors of maternal healthcare services by women with disabilities, and examine whether the disability and caste identity of women affects and compounds the utilization of maternal healthcare services.

## METHOD

### Setting

The study was conducted in Rupandehi, a southern district of Nepal with a population of 880,196 of which 50.89% are female (11). Just over 25% of the population are Janajati (indigenous) followed by 21% Brahmin and Chhetri caste and 12% are from the more disadvantaged Dalit castes. 1.12% of the district population are reported to have a disability (11).

Primary healthcare services including maternal care in the district are delivered through five Primary Healthcare Centers (PHCC), six Urban Health Clinics (UHC), six Health Posts (HP) and 58 Sub-Health Posts (SHP). One district hospital and one zonal hospital (covering six districts) provide secondary care services. In addition to the government health services, there is a wide network of NGOs and private sector services delivered by private hospitals, nursing homes, clinics and pharmacies/drug shops in the district (31).

### Study design

A cross-sectional survey was conducted among married and unmarried women aged between 15 – 49 years, who had been pregnant within the last five years. Women with two or more pregnancies or deliveries were instructed to refer to the most recent pregnancy or childbirth. The sample was drawn from four different categories: Dalit and non-Dalit women with a disability, Dalit women without a disability and non-Dalit women without a disability. While disabled and Dalit women were of particular interest in the study, non-Dalit non-disabled women were included for comparison. Women with a severe intellectual disability, unable to provide responses and those with a hearing impairment with whom it was not possible to communicate by verbal and or non-verbal means, such as Nepali Sign Language, were excluded from the study.

### Sample

The study used a non-probability, multi-stage sampling method for selection and recruitment of participants. In the first stage, disabled women within the criteria were identified using the most recent census and with the help of local disabled people's organizations (DPOs). Out of 119 disabled women identified, 79 (66.4%) women were interviewed. Exclusions included 30 women not located, four that did not meet the disability criteria of the UN Washington Group questionnaire ([www.washingtongroup-disability.com](http://www.washingtongroup-disability.com)) and six who could not be interviewed due to communication difficulties. Based on the number of disabled women, the number of non-disabled Dalit and non-disabled non-Dalit women for interview was determined using a 1:2:2 ratios resulting in a total sample size of 354 for study.

## The tool and data collection

A structured questionnaire with 95 questions covering demographics, maternal healthcare and service utilization was adapted from previously developed standard questions used in Nepal (such as the NDHS questionnaire), with additional new questions relating to disability and empowerment. The questionnaire was field tested, refined and validated before use in the interviews (32). Twelve trained female field researchers collected the data. The data was checked to ensure completeness of information; for any found incomplete, participants were revisited to complete/confirm the information.

## Measures

### Outcome Measures

Three measures of maternal healthcare service utilization were used: Antenatal Care (ANC), Delivery and Post-Natal Care (PNC). The study considered four or more ANC visits during the last pregnancy as adequate utilization (33). Utilization for delivery was assessed as delivery in any type of health facility - private or public birthing centres, health posts, primary healthcare centres or hospitals. Adequate Post-Natal Care utilization was determined as '*at least one distinct visit to a health facility for post-natal care*' made within one week of delivery. Table 1 shows the outcome measure variables, exposures and covariates used in the study.

**Table 1: Variables and their description measuring maternal healthcare service utilization**

Measure	Definition /Coding	Level of Measurement
<b>Outcome Variables</b>		
ANC Visit	Attendance at a health facility (HF) by a pregnant woman for antenatal care during last pregnancy (0 = <4 visit, 1 = 4+ visit)	Binary
HF Delivery	Most recent childbirth at a health facility (HF) designated for safe delivery. (0 = Home delivery, 1 = HF delivery)	Binary
PNC Visit	Attendance at a HF for postnatal care by a woman within 7 days of last childbirth. (0 = No, 1 = Yes)	Binary
<b>Exposure Variables</b>		
Disability	Disability status of respondent women meeting Washington Group Questionnaire criteria. (1 = Non-disabled, 2 = disabled)	Binary
Caste	Self-reported caste of respondent woman. (1 = Dalit, 2 = Non-Dalit)	Binary
<b>Covariates</b>		
Place of residence	At the time of survey, respondent living in VDC are considered rural and those living in Municipality are urban. (1 = rural, 2 = urban)	Binary
Age	Completed age of women at the time of survey in years. (1=15 -24, 2 =25-34, 3 =35- 49)	Categorical
Parity	Number of pregnancy or childbirths in lifetime. (1=Primipara, 2 = Multipara)	Binary
Respondent's education	Number of years of school education completed by the women. (1 = Illiterate, 2 = Literate/Primary education up to class 5, 3 = Secondary education or above)	Categorical
Knowledge and awareness Index	Knowledge of danger signs spontaneously cited relating to pregnancy, childbirth and postpartum (An additive index). (0 = No knowledge at all, 1 = Little knowledge or low level of awareness, 2 = Enough knowledge – high level of awareness)	Ordinal
Empowerment Index	An index developed using 24 questionnaire items at individual, family and community level indicators. (1 = low, 2 = medium, 3 = high)	Ordinal
Household Wealth Index	Wealth index developed based on socio-economic indicators and household assets reported at the time of survey. (1=low, 2 = Middle, 3 = high)	Ordinal

**Household Wealth Index**

The household wealth index variable was constructed using 22 indicators of household possessions having a total score of 26. Household possession or dwelling characteristics were chosen from the Nepal Demographic Health Survey 2011. Except for Roofing Materials and Source of water, all variables were dichotomized assigning the code 0 or 1. Roofing Materials was assigned the weight 1 with natural/non-durable roofing materials, 2 with durable roofing materials and 3 for long durable roofing such as cement and concrete. Households using surface water/shallow wells were given the score 1, those using tube wells were assigned 2, and those using piped water were given 3. Women were ranked according to the score of their households. For the purpose of analysis, the lowest two and the highest two groups were collapsed to make three categories (low, medium and high) of wealth ranking.

**Women's Empowerment Index**

Four dimensions: economic, socio-cultural, familial/interpersonal and political were used to construct the women's empowerment index (34,35). These dimensions were adapted to reflect the local context at individual/family (micro) and community (meso) level indicators with 24 questionnaire items to derive a score. All the questionnaire items were equally weighted, dichotomized, and a composite measure was created using the sum (maximum score of 24).

**Knowledge and awareness Index**

The knowledge and awareness index was created by asking two questions. Firstly, respondents were asked: 'Are you aware of any problems that are potentially dangerous to the mother during pregnancy and childbirth?' Those who answered affirmatively were asked the second question: 'What are the danger signs to mothers during pregnancy, childbirth and after delivery?' Women who answered "No" for the first question were given a score of 0 and women stating one to three correct danger signs to the second question were given a score of 1. Those stating four or more signs scored 2.

**Statistical Analysis**

Data was entered using EPI-Info version 3.4.1, then transferred into SPSS (version 16.0 for Windows) for analysis. The associations between the measures of utilization – Ante-natal Care (any check-up and frequency of check-up), Delivery (HF vs home delivery) and Post-natal check-up (yes/no) and disability/Dalit status were examined using cross tabulations and chi-square tests. Logistic regression analysis was carried out to test whether disability or Dalit status was related to utilization measures once socio-demographic, empowerment and awareness characteristics of individuals had been taken into account. Confounding was investigated by multivariate analysis. Further, disability and caste were fitted

separately in models and intersectionality of disability and caste was tested by inclusion of a disability\*caste interaction term.

## RESULTS

### Socio-demographic characteristics of survey participants

Table 2 shows the characteristics of survey participants. Of 354 women surveyed, 40% were non-disabled non-Dalits, about 38% were non-disabled Dalits, 17% disabled non-Dalits and 5% were Disabled-Dalit. Over 80% participants lived in rural areas. Approximately half of the respondents were 25 – 34 years, 38% were 15 – 24 years and the remaining were between 35 – 49 years of age. Nearly one-third of women reported having no education or were illiterate, and one fifth had no knowledge at all about danger signs in pregnancy. The majority of participants (56%) were primiparous. Over 19% participants were assessed as having higher levels of empowerment and autonomy, 58.8% with medium levels, and 21.8% with low levels. Similarly, one fifth of women were ranked as having higher levels of household wealth, 56.5% with middle levels, and 22.9% with lower levels.

**Table 2: Characteristics of study participants**

Background Characteristics	n=354 (%)
<b>Participant Type</b>	
Non-disabled non-Dalit	142 (40.1%)
Non-disabled Dalit	133 (37.6%)
Disabled non-Dalit	61 (17.2%)
Disabled Dalit	18 (5.1%)
<b>Place of Residence</b>	
Rural	284 (80.2%)
Urban	70 (19.8%)
<b>Respondent's Age</b>	
15 – 24 Years	134 (37.9%)
25 – 34 Years	172 (48.6%)
35 – 49 Years	48 (13.6%)
<b>Education</b>	
Illiterate	116 (32.8%)
Literate/Primary (up to 5 Class)	117 (33.1%)
Secondary (6 class & above)	121 (34.2%)
<b>Knowledge &amp; Awareness</b>	
No Knowledge at All	71 (20.1%)
Little Knowledge (Low level of Awareness)	138 (39.0%)
Enough Knowledge (High level of Awareness)	145 (41.0%)
<b>Parity</b>	
Primipara	198 (55.9%)
Multipara	156 (44.1%)
<b>Women Empowerment Index</b>	

Low	77 (21.8%)
Medium	208 (58.8%)
High	69 (19.5%)
<b>Household Wealth Index</b>	
Low (<40% in ranking)	81 (22.9%)
Middle (40 – 60% in ranking)	200 (56.5%)
High (>60% in ranking)	73 (20.6%)

### Maternal Health Care Service Utilization

Over 73% of women had attended the recommended four or more ANC visits in their last pregnancy. Less than two-thirds of participants reported that they had a health facility (HF) delivery and only 29% received postnatal care. Among the four groups, non-disabled non-Dalit women reported the highest percentage of utilization of all services followed by non-disabled Dalits. Disabled Dalits had the least utilization ANC and HF delivery. However, the percentage of disabled Dalits using postnatal care services was higher than disabled non-Dalit women (28% vs 18%). (Table 3).

**Table 3: Maternal Health Care Service Utilization by different groups of study participants**

Outcome Variables	Disabled Dalit	Disabled non-Dalit	Non-disabled Dalit	Non-disabled non-Dalit	Total	P - Value
	n = 18	n = 51	n = 133	n = 142	n = 354	
Utilization of 4+ ANC	10 (56%)	37 (73%)	101 (76%)	111 (78%)	259 (73%)	0.136
Health Facility Delivery	9 (50%)	31 (61%)	84 (63%)	105 (74%)	229 (65%)	0.008
Utilization of PNC services	5 (28%)	9 (18%)	40 (30%)	49 (35%)	103 (29%)	0.017

Pearson Chi-square test of significance

### Bivariate Analysis Result

Table 4 illustrates that there is no significant relationship between an individual's disability and caste status and attendance for the full-recommended number of antenatal care visits. ( $P>0.5$ ). However, education, empowerment and household wealth factors showed some relationship with 4 or more ANC visits ( $P<0.05$ ). Compared to illiterate women, there was higher odds of 4 or more ANC visits with increasing levels of women's education (OR 2.27, 95% CI 1.22 – 4.23 for literate or primary level education; OR 3.61, 95% CI 1.84 – 7.08 for secondary or above education). It was also evident that women with greater levels of empowerment/autonomy had higher odds of using ANC services (OR 1.91, 95% CI 1.05 – 3.49 for medium, and OR 4.40, 95% CI 1.74 – 11.10 for high level empowerment). Also, women with higher household wealth index (OR 1.98, 95% CI 1.05 – 3.72) were more likely to receive the full 4 ANC services than those from households with a low index.

The respondents' disability status showed significant associations with health facility delivery ( $P < 0.05$ ); whereas the respondent's caste did not show any association ( $P > 0.05$ ). The odds of health facility delivery for disabled non-Dalit and disabled Dalits women was 60% (OR 0.40, 95% CI 0.21 – 0.77) and 65% (OR 0.35, 95% CI 0.13 – 0.96) lower respectively than non-disabled non-Dalits. Urban women were also twice as likely to have a HF delivery compared to rural women (OR 2.02, 95% CI 1.10 – 3.72).

Women's education, age and parity were strong factors influencing whether to deliver at home or HF. Women with secondary level or higher education had greater odds than those who were illiterate or had no education (OR 14.63, 95% CI 6.94 – 30.83). The odds ratio of HF delivery for the women age between 25 – 34 years was 0.42, 95% CI 0.25 – 0.71; and 35 – 49 years had OR 0.21, 95% CI 0.10 – 0.42. Multiparous women had lower odds (OR 0.23, 95% CI 0.14 – 0.36) than primiparous. Women's knowledge and awareness of pregnancy danger signs and household wealth, were also found to be strong predictors for giving birth at a health facility (OR 2.05, 95% CI 1.12 – 3.75 for adequate knowledge and awareness and OR 5.84, 95% CI 2.77 – 12.30 for highest wealth ranking groups). (Table 4).

Women's disability and some socio-demographic factors such as respondent's education, age at marriage and parity were associated with PNC service utilization ( $P < 0.05$ ). Women with secondary or higher education had greater odds (OR 1.87, 95% CI 1.05 – 3.33) of utilizing PNC services than those with no education. Women with two or more pregnancies were 50% less likely to use PNC than those with their first pregnancy (OR 0.54, 95% CI 0.33 – 0.86). No relationship was found between the respondent's caste and PNC service utilization ( $P > 0.05$ ) (Table 4).

**Table 4: Bivariate Analysis**

Variables	ANC (4+)			HF Delivery			PNC (At least 1)		
	OR	95% CI	P - Value	OR	95% CI	P - Value	OR	95% CI	P - Value
<b>Disability</b>									
Non-disabled	1			1			1		
Disabled	0.63	(0.34 - 1.16)	0.138	0.50	(0.30 - 0.84)	0.009	0.47	(0.25 - 0.88)	0.019
<b>Caste</b>									
Non-Dalit	1			1			1		
Dalit	0.81	(0.48 - 1.37)	0.440	0.77	(0.49 - 1.20)	0.243	0.93	(0.60 - 1.52)	0.839
<b>Place of Residence</b>									
Rural	1			1			1		
Urban	1.92	(0.90 - 4.09)	0.093	2.02	(1.10 - 3.72)	0.023	0.54	(0.28 - 1.01)	0.056
<b>Age</b>									
15 - 24 yrs	1			1			1		
25 - 34 yrs	0.64	(0.36 - 1.14)	0.127	0.42	(0.25 - 0.71)	0.001	1.03	(0.63 - 1.69)	0.896
35 - 49 yrs	0.48	(0.21 - 1.12)	0.088	0.21	(0.10 - 0.42)	0.000	0.44	(0.19 - 1.03)	0.059
<b>Education</b>									
Illiterate	1			1			1		
Primary Education	2.27	(1.22 - 4.23)	0.010	2.27	(1.34 - 3.84)	0.002	1.54	(0.86 - 2.77)	0.150
Secondary or higher	3.61	(1.84 - 7.08)	0.000	14.63	(6.94 - 30.83)	0.000	1.87	(1.05 - 3.33)	0.034

<b>Parity</b>								
Primi	1		1		1			
Multi	0.60 (0.35 - 1.01)	0.054	0.23 (0.14 - 0.36)	0.000	0.54 (0.33 - 0.86)	0.011		
<b>Knowledge &amp; awareness</b>								
No knowledge	1		1		1			
Little	1.07 (0.53 - 2.14)	0.859	1.04 (0.58 - 1.87)	0.891	0.87 (0.45 - 1.65)	0.663		
Enough	1.51 (0.74 - 3.06)	0.258	2.05 (1.12 - 3.75)	0.020	1.25 (0.67 - 2.33)	0.483		
<b>Empowerment</b>								
Low	1		1		1			
Medium	1.91 (1.05 - 3.49)	0.034	1.04 (0.60 - 1.80)	0.887	1.01 (0.56 - 1.82)	0.980		
High	4.40 (1.74 - 11.10)	0.002	1.69 (0.83 - 3.44)	0.150	1.37 (0.68 - 2.78)	0.381		
<b>Household Wealth</b>								
Low	1		1		1			
Middle	1.98 (1.05 - 3.72)	0.034	2.76 (1.62 - 4.70)	0.000	1.50 (0.82 - 2.75)	0.191		
High	1.63 (0.76 - 3.47)	0.208	5.84 (2.77 - 12.30)	0.000	1.83 (0.90 - 3.74)	0.097		

\*P<0.05, \*\*P<0.01, \*\*\*P<0.001

### Multi-variate Analysis

Table 5 shows odds ratios for disability status with all three service utilization outcomes, adjusted for socio-demographic and household factors theorized to be a potential confounders of associations.

Overall, disabled respondents reported less odds of utilization in all three outcomes, compared to non-disabled. In the unadjusted model, there was no association between disability and the 4 recommended ANC visits, and After adjusting for potential confounding variables, the data did not show a significant relationship between disability and ANC visits.

Disabled women had lower odds of HF delivery (OR 0.48, 95% CI 0.26 – 0.86) and PNC (OR 0.47, 95% CI 0.25 – 0.88) compared to non-disabled women. After adjustment for women's age, there was no association between disability and HF delivery. Adjustment for place of residence (location) and parity reduced the association. In the full model, there was not a significant association between disability and HF delivery. The same group of confounders affected the relationship between disability and PNC in the same way as in HF delivery.

**Table 5: Unadjusted and adjusted odds ratio of disability with maternal health care utilization**

Factors	ANC 4+ visit			HF Delivery			PNC visit		
	OR	95% CI		OR	95% CI		OR	95% CI	
<b>Disability Only</b>	<b>0.63</b>	<b>0.34 - 1.16</b>		<b>*0.48</b>	<b>0.26 - 0.86</b>		<b>*0.47</b>	<b>0.25 - 0.88</b>	
<b>Disability + Caste</b>	<b>0.58</b>	<b>0.31 - 1.09</b>		<b>**0.45</b>	<b>0.26 - 0.77</b>		<b>*0.46</b>	<b>0.24 - 0.88</b>	
Non-Dalit	1			1			1		
Dalit	0.74	0.43 - 1.27		0.66	0.41 - 1.05		0.93	0.58 - 1.50	
<b>Disability + Education</b>	<b>0.68</b>	<b>0.36 - 1.28</b>		<b>*0.54</b>	<b>0.32 - 0.91</b>		<b>*0.46</b>	<b>0.26 - 0.92</b>	
Illiterate/No education	1			1			1		
Primary education	*2.18	1.17 - 4.08		**2.12	1.25 - 3.61		1.44	0.79 - 2.60	
Secondary or higher	***3.55	1.81 - 6.98		***14.73	6.94 - 31.26		*1.81	1.01 - 3.24	
<b>Disability + Location</b>				<b>*0.54</b>	<b>0.32 - 0.91</b>				

Rural	1					
Urban	*1.85	1.01 - 3.43				
<b>Disability + Women's Age</b>	<b>0.72</b>	<b>0.41 - 1.26</b>				
15 - 24 yrs	1					
25 - 34 yrs	**0.44	0.26 - 0.74				
35 - 49 yrs	***0.25	0.12 - 0.52				
<b>Disability + Parity</b>	<b>*0.61</b>	<b>0.35 - 1.06</b>	<b>*0.52</b>	<b>0.27 - 0.98</b>		
Primi	1		1			
Multi	***0.24	0.15 - 0.39	*0.58	0.35 - 0.93		
<b>Disability + Knowledge/Awareness</b>	<b>*0.54</b>	<b>0.32 - 0.91</b>				
No knowledge	1					
Little	1.00	0.55 - 1.81				
Enough	*1.89	1.02 - 3.48				
<b>Disability + Empowerment</b>	<b>0.64</b>	<b>0.34 - 1.19</b>				
Low	1					
Medium	*1.91	1.05 - 3.49				
High	**4.37	1.23 - 11.05				
<b>Disability + Wealth Index</b>	<b>0.70</b>	<b>0.37 - 1.31</b>	<b>0.68</b>	<b>0.39 - 1.19</b>		
Low	1		1			
Middle	1.85	0.98 - 3.52	**2.54	1.47 - 4.39		
High	1.48	0.68 - 3.23	***5.21	2.43 - 11.17		
<b>Full Model</b>	<b>0.71</b>	<b>0.37 - 1.38</b>	<b>0.76</b>	<b>0.38 - 1.50</b>	<b>*0.52</b>	<b>0.27 - 0.99</b>

\*P&lt;0.05, \*\*P&lt;0.01, \*\*\*P&lt;0.001

Table 6 and 7 show the models with interaction terms that look at the effect of interactions between exposures and key predictors of service utilization that included age, education, parity and wealth factors and their association in utilization for disabled and non-disabled, and Dalit and non-Dalit women were examined. Given the data, in most cases there was insufficient evidence of interaction effect between disability and women's age, education, parity and household wealth at 5% significant level. (Table 6). However, a weak interaction effect in the utilization of recommended ANC services found in the caste group by their education (OR 0.19, 95% CI 0.05 - 0.74). (Table 7).

**Table 6: Models with interaction terms analysing the effects of interactions between key predictors and service utilization by disability**

Predictors	ANC4+			HF Delivery			PNC		
	OR	95% CI	P Value	OR	95% CI	P Value	OR	95% CI	P Value
<b>Disability + Age</b>									
Disabled	0.83	0.20 - 3.43	0.800	0.65	0.20 - 2.06	0.462	1.23	0.26 - 5.88	0.794
Non-disabled									
<b>Age</b>									
15 - 24 years	2.02	0.65 - 6.34	0.226	4.41	1.64 - 11.88	<b>0.003</b>	2.98	0.83 - 10.76	0.096
25 - 34 years	1.27	0.42 - 3.85	0.674	1.55	0.60 - 3.97	0.364	2.83	0.79 - 10.18	0.110
35 - 49 years									
<b>Disabled</b>									
15 - 24 years	0.61	0.08 - 4.50	0.626	0.41	0.08 - 2.15	0.290	0	0	0.999

25 - 34 years	0.95	0.18 - 4.89	0.947	1.69	0.42 - 6.79	0.459	0.56	0.10 - 3.31	0.524
35 - 49 years									
<b>Disability + Education</b>									
Disabled	0.71	0.29 - 1.76	0.462	0.63	0.27 - 1.48	0.292	0.27	0.08 - 0.99	<b>0.048</b>
Non-disabled									
<b>Education</b>									
Illiterate									
Primary Education	1.94	0.97 - 3.85	0.060	2.28	1.25 - 4.14	<b>0.007</b>	1.29	0.68 - 2.44	0.440
Secondary and above	4.58	1.99 - 10.53	<b>0.000</b>	21.06	7.76 - 57.17	<b>0.000</b>	1.56	0.82 - 2.95	0.176
<b>Disabled</b>									
Illiterate									
Primary Education	2.49	0.41 - 15.11	0.323	0.75	0.21 - 2.78	0.671	2.01	0.35 - 11.45	0.434
Secondary and above	0.425	0.10 - 1.84	0.253	0.36	0.08 - 1.76	0.208	2.42	0.48 - 12.07	0.282
<b>Disability + Parity</b>									
Disabled	0.67	0.29 - 1.52	0.334	0.58	0.28 - 1.17	0.126	0.44	0.17 - 1.13	0.089
Non-disabled									
<b>Parity</b>									
Primipara	1.60	0.87 - 2.91	0.129	4.03	2.35 - 6.92	<b>0.000</b>	1.65	0.97 - 2.80	0.065
Multipara									
<b>Disabled</b>									
Primipara	1.03	0.30 - 3.60	0.963	1.17	0.38 - 3.67	0.783	1.37	0.38 - 4.99	0.631
Multipara									
<b>Disability + Wealth</b>									
Disabled	0.82	0.15 - 4.54	0.823	1.36	0.15 - 12.36	0.786	0.73	0.13 - 4.07	0.720
Non-disabled									
<b>Wealth Index</b>									
Low wealth group	0.80	0.32 - 1.97	0.626	0.26	0.11 - 0.60	<b>0.002</b>	0.70	0.31 - 1.58	0.389
Middle wealth group	1.22	0.60 - 2.50	0.584	0.49	0.24 - 1.01	<b>0.050</b>	0.89	0.49 - 1.63	0.700
High wealth group									
<b>Disabled</b>									
Low wealth group	0.58	0.08 - 4.36	0.600	0.28	0.03 - 3.10	0.301	0.64	0.08 - 5.04	0.671
Middle wealth group	1.05	0.15 - 7.33	0.958	0.68	0.07 - 7.01	0.743	0.68	0.10 - 4.70	0.695
High wealth group									

**Table 7: Models with interaction terms analysing the effects of interactions between key predictors and service utilization by caste**

Predictors	ANC4+			HF Delivery			PNC		
	OR	95% CI	P Value	OR	95% CI	P Value	OR	95% CI	P Value
<b>Caste + Age</b>									
Dalit	3.11	0.67 - 14.44	0.147	1.28	0.40 - 4.09	0.683	0.32	0.07 - 1.56	0.159
Non-Dalit									
<b>Age</b>									
15 - 24 years	5.00	1.57 - 15.94	<b>0.007</b>	6.10	2.35 - 15.84	<b>0.000</b>	1.30	0.41 - 4.12	0.653
25 - 34 years	2.30	0.84 - 6.34	<b>0.106</b>	2.83	1.22 - 6.60	<b>0.016</b>	1.17	0.37 - 3.67	0.792
35 - 49 years									
<b>Dalit</b>									
15 - 24 years	0.15	0.02 - 0.89	<b>0.037</b>	0.58	0.14 - 2.45	0.460	2.91	0.51 - 16.53	0.228
25 - 34 years	0.25	0.05 - 1.38	0.112	0.44	0.12 - 1.66	0.226	3.71	0.67 - 20.47	0.132
35 - 49 years									
<b>Caste + Education</b>									
Dalit	1.79	0.79 - 4.05	0.161	1.01	0.48 - 2.11	0.990	1.14	0.47 - 2.76	0.769
Non-Dalit									
<b>Education</b>									
Illiterate									
Primary Education	3.66	1.45 - 9.23	<b>0.006</b>	2.06	0.99 - 4.28	<b>0.054</b>	1.91	0.82 - 4.46	0.135

Secondary and above	7.27	2.84 - 18.62	<b>0.000</b>	20.83	7.42 - 58.46	<b>0.000</b>	2.11	0.81 - 5.46	0.125
<b>Dalit</b>									
Illiterate									
Primary Education	0.38	0.11 - 1.37	0.140	1.21	0.42 - 3.46	0.727	0.64	0.20 - 2.10	0.463
Secondary and above	0.19	0.05 - 0.74	<b>0.017</b>	0.41	0.09 - 1.83	0.240	0.82	0.25 - 2.73	0.745
<b>Caste + Parity</b>									
Dalit	1.32	0.63 - 2.78	0.469	1.11	0.59 - 2.09	0.736	1.01	0.48 - 2.14	0.984
Non-Dalit									
<b>Parity</b>									
Primipara	2.36	1.14 - 4.89	<b>0.020</b>	5.14	2.72 - 9.75	<b>0.000</b>	1.71	0.89 - 3.31	0.108
Multipara									
<b>Dalit</b>									
Primipara	0.46	0.16 - 1.32	0.147	0.69	0.27 - 1.79	0.446	1.26	0.48 - 3.32	0.638
Multipara									
<b>Caste + Wealth</b>									
Dalit	0.77	0.24 - 2.44	0.656	0.76	0.22 - 2.64	0.665	1.20	0.43 - 3.33	0.726
Non-Dalit									
<b>Wealth Index</b>									
Low wealth group	0.73	0.24 - 2.21	0.581	0.13	0.05 - 0.35	<b>0.000</b>	0.77	0.30 - 1.98	0.586
Middle wealth group	1.09	0.47 - 2.52	0.847	0.48	0.21 - 1.14	0.096	0.76	0.37 - 1.57	0.462
High wealth group									
<b>Dalit</b>									
Low wealth group	0.85	0.17 - 4.11	0.835	1.89	0.41 - 8.73	0.416	0.48	0.11 - 2.10	0.330
Middle wealth group	1.37	0.35 - 5.39	0.652	1.00	0.25 - 3.97	0.997	1.13	0.35 - 3.73	0.836
High wealth group									

## DISCUSSION

This study found that disabled women used all maternal healthcare services less than non-disabled women of all castes. The results show that it is the disability status of pregnant Nepali women, not their caste that determines the utilization of health facility delivery and post-natal care. Associations between disability and all three maternal healthcare utilization outcomes were entirely explained by the inclusion of education and household wealth; suggesting that the main reasons why disabled women were less likely to utilize maternal healthcare services was due to social and household circumstances.

There are few statistics available to compare the findings of this study within Nepal as the government has limited mechanisms for recording disability data, and most previous studies on maternal healthcare have not included disability. The few qualitative studies conducted in this field in Nepal indicate poor utilization of services by women with disabilities during their pregnancy and childbirth (6,13). This contrasts with findings from developed countries, where a more structured and intensive approach to disability pregnancy care may be offered within well-developed health systems (16).

There was no evidence of effect modification by the women's caste status. . There have been long standing and highly politicized concerns that major caste-dependent differences remain, with lower

maternal healthcare utilization among Dalits consistently reported in the literature (36–41). Increases in public awareness and policy developments resulting from these concerns may have contributed to changes in social equity and improvements in utilization across caste groups, so it is possible that differences in service utilization resulting from caste-based factors (27) could have improved in response to activism and policy attention to this disparity.

The study identified a number of individual, socio-demographic and family related factors associated with maternal healthcare service utilization by women with disabilities. Parity was associated with increased likelihood of health facility delivery if it was a first birth, corresponding with previous studies in Nepal (42,43). First time mothers may perceive more risks due to a lack of prior childbirth experience or may have other anxieties stemming from a lack of reliable information.

Women's level of education and their knowledge and awareness were also strong predictors for utilization of services, with better educated women more likely to give birth at a HF and to receive PNC services, consistent with other's findings (44–47). Educated women may have a greater understanding of the benefits of using services as well as the autonomy and confidence to use them. Education may also increase women's status in their family, enabling them to choose and access services more easily (48). Disabled women, both Dalit and non-Dalit consistently had less education than their non-disabled peers and hence this may be a significant explanatory factor for why disabled women utilized services at lower rates.

### **Limitations of the Study**

We acknowledge several constraints and limitations. This research was a part of a Safe Motherhood Project in southern Nepal; therefore, the study population was limited to the project district. Secondly, having a very specific 'hard to reach' population for the study, with narrow inclusion criteria (i.e. Dalit and women with disability having had at least one pregnancy between 2009 and 2014), limited the sample size. Furthermore, the small and disproportionate sample size had implications for data analysis, forcing the researchers to employ a logical collapse of the variables where possible to measure associations with other variables or outcomes. In several circumstances the tests for association was calculated with a small figure, which may have implications for accuracy. Given the large number of comparisons made in the quantitative section, significant associations close to the 0.05 level should be interpreted with caution, as 1 in 20 tests may have a p-value below 0.05 purely due to chance (49). Consistent relationships were seen between disability status and some maternal healthcare outcomes, suggesting that at least for disability status, findings are robust.

Language and communication barriers were another prominent constraint in the study. that was overcome by recruiting local research assistants and also using a questionnaire translated into the local languages (Nepali and Bhojpuri).

## **CONCLUSION**

The results from this study show that the disability status of pregnant Nepali women, not their caste, determines the utilization of health facility delivery and post-natal care. Associations between disability and maternal healthcare utilization outcomes were explained by socio-demographic factors. Except for a weak effect modification by education in the utilization of four recommended ANC services by Dalits, there was no evidence of effect modification in the utilization of services by women's caste. The findings of this study suggest that policy and political discourse should now be widened, to acknowledge and address the disparities and challenges faced by women with disabilities of all castes.

**List of Abbreviations:** ANC: Antenatal Care, BLF: Big Lottery Fund, CI: Confidence Interval, DPO: Disabled Person's Organization, HF: Health Facility, NDHS: Nepal Demographic Health Survey, NGO: Non-governmental Organization, OR: Odds Ratio, PNC: Post-natal Care, UN: United Nations

## Declarations

**Ethics approval and consent to participate:** Ethical permission was obtained from the Nepal Health Research Council (NHRC) – Ref. no. 1184 and UCL ethics committee project ID: 5260/001. Additionally, approval was received from the government District Public Health Office in the study district. Verbal and signed consent was taken from all participants before conducting interviews. Parental consent was also obtained for participants under 16 years old.

**Consent for publication:** Not Applicable

**Availability of data and material:** The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

**Competing interests:** The authors declare that there is no competing interest.

**Funding:** Big Lottery Fund UK through Kidasha in Nepal funded the study. There was no role of the funding body in study design, data collection, analysis or interpretation and writing the manuscript.

**Author's contribution:** HRD conceived and implemented the project. NG guided design and field implementation. MK provided input in study design, implementation and writing. EM supported in statistical analysis, interpretation and ensured the data analysis procedures. HRD and AC prepared the manuscript and NG, MK and EM commented and provided their inputs for finalization. All authors reviewed and approved the final version of the manuscript.

**Acknowledgement:** The authors acknowledge the support and contribution of Kidasha and the colleagues – Janice Miller, Shanti Shirish, Ishwar Mainali and Tanka Shrestha, for the implementation of this study, field work and data analysis. In addition, the authors wish to acknowledge the support provided by Disabled People's Organizations (DPOs) and the local partner, NAMUNA and its staff in Rupandehi. The authors are grateful to the women who shared their views and personal experiences.

**Author's information:** Hridaya Raj Devkota (PhD), Post-doctoral Fellow, Formerly Program Manager, Kidasha Nepal. Email: [hridayadevkota@hotmail.com](mailto:hridayadevkota@hotmail.com) (Corresponding author); Andrew Clarke (MPH), Senior Health Advisor, Save the Children UK, London; Honorary Researcher, Division of health research,

Lancaster University, UK. Email: [andrewcws@gmail.com](mailto:andrewcws@gmail.com); Emily Murray (PhD), Senior Research Associate, Department of Epidemiology and Public Health, University College London (UCL), UK. Email: [emily.murray@ucl.ac.uk](mailto:emily.murray@ucl.ac.uk); Maria Kett (PhD) Reader, Leonard Cheshire Disability and Inclusive Development Centre, Department of Epidemiology and Public Health, UCL. Email: [m.kett@ucl.ac.uk](mailto:m.kett@ucl.ac.uk); Nora Groce (PhD), Professor and Leonard Cheshire Chair, Leonard Cheshire Disability and Inclusive Development Centre, Division of Epidemiology and Public Health, University College London (UCL), UK. Email: [nora.groce@ucl.ac.uk](mailto:nora.groce@ucl.ac.uk).

## References

1. MoH/NewEra/ICF- International. Nepal Demographic and Health Survey 2016. Kathmandu Nepal: Ministry of Health, Nepal. 2017.
2. Thaddeus S, Maine D. TOO FAR TO WALK: MATERNAL MORTALITY IN CONTEXT. Soc Sci Med [Internet]. 1994;38(8):1091–110. Available from: [https://doi.org/10.1016/0277-9536\(94\)90226-7](https://doi.org/10.1016/0277-9536(94)90226-7)
3. ARROW. MDG 5 in Asia: Progress, Gaps and Challenges 2000 - 2010. 2010;1–8.
4. Bennett L, Dahal DR, Govindasamy P. Caste, Ethnic and Regional Identity in Nepal: Further Analysis of the 2006 Nepal Demographic and Health Survey. Calverton, Maryland, USA: Macro International Inc. 2008;
5. Kumar P, Gupta A. Determinants of Inter and Intra caste Differences in Utilization of Maternal Health Care Services in India : Evidence from DLHS-3 Survey. Int Res J Soc Sci. 2015;4(1):27–36.
6. Morrison J, Basnet M, Budhathoki B, Adhikari D, Tumbahangphe K, Manandhar D, et al. Disabled women's maternal and newborn health care in rural Nepal: A qualitative study. Midwifery [Internet]. 2014 Mar 28 [cited 2014 Sep 22];1–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24768318>
7. Pandey JP, Dhakal MR, Karki S, Poudel P, Pradhan MS. Maternal and Child Health in Nepal: The Effects of Caste, Ethnicity, and Regional Identity: Further Analysis of the 2011 Nepal Demographic and Health Survey. Kathmandu, Nepal; 2013.
8. Saroha E. Caste As a Determinant of Utilization of Maternal and Neonatal Healthcare Services in Maitha, Uttar Pradesh, India [Internet]. The University of Alabama at Birmingham, ProQuest Dissertations Publishing; 2007. Available from: <https://search.proquest.com/docview/304893270?pq-origsite=gscholar&fromopenview=true>
9. UNITED NATIONS. Convention on the Rights of Persons with Disabilities and Optional Protocol [Internet]. 2006. p. 1–37. Available from: <https://www.un.org/disabilities/documents/convention/convoptprot-e.pdf>
10. WHO, The World Bank. WORLD REPORT ON DISABILITY [Internet]. New York; 2011. Available from: [https://www.who.int/disabilities/world\\_report/2011/report.pdf](https://www.who.int/disabilities/world_report/2011/report.pdf)
11. GoN/NPC/CBS. National Population and Housing Census 2011 (National report). Vol. 01. 2012.

12. Shrestha E, Nilsson A. Mainstreaming disability in the new development paradigm: Evaluation of Norwegian support to promote the rights of persons with disabilities (Nepal country report). 2012.
13. Anon. Maternal and new-born care practices among disabled women, and their attendance in community groups in rural Makwanpur, Nepal. London; 2013. (Background Paper: 01).
14. Becker H, Stuifbergen A, Tinkle M. Reproductive health care experiences of women with physical disabilities: a qualitative study. Arch Phys Med Rehabil [Internet]. 1997 Dec;78(12 Suppl 5):S26-33. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9422004>
15. Lipson JG, Rogers JG. Pregnancy, birth, and disability: women's health care experiences. Health Care Women Int [Internet]. 2000;21(1):11–26. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11022446>
16. Redshaw M, Malouf R, Gao H, Gray R. Women with disability: the experience of maternity care during pregnancy, labour and birth and the postnatal period. BMC Pregnancy Childbirth [Internet]. 2013 Jan [cited 2014 May 3];13(1):174. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3848505&tool=pmcentrez&rendertype=abstract>
17. Smith E, Murray SF, Yousafzai a K, Kasonka L. Barriers to accessing safe motherhood and reproductive health services: the situation of women with disabilities in Lusaka, Zambia. Disabil Rehabil [Internet]. 2004 Jan 21 [cited 2013 Jun 26];26(2):121–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/14668150>
18. Trani JF, Browne J, Kett M, Bah O, Morlai T, Bailey N, et al. Access to health care, reproductive health and disability: a large scale survey in Sierra Leone. Soc Sci Med [Internet]. 2011 Nov [cited 2013 Nov 8];73(10):1477–89. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22014873>
19. Dhungana BM. The lives of disabled women in Nepal: vulnerability without support. Disabil Soc [Internet]. 2006;21(2):133–46. Available from: <https://www.tandfonline.com/doi/abs/10.1080/09687590500498051>
20. Khanal NA. Status of Reproductive Health and Experience of Motherhood of Disabled Women in Nepal. 2013;(July):1–52.
21. Gurung H. The Dalit Context. Occas Pap Sociol Anthropol [Internet]. 2008 Jun 20;9. Available from: <http://nepjol.info/index.php/OPSA/article/view/1133>
22. Nohrlind R. CASTE-BASED DISCRIMINATION IN SOUTH ASIA: SITUATIONAL OVERVIEW, RESPONSES AND WAYS FORWARD [Internet]. 2009. Available from: [https://idsn.org/wp-content/uploads/user\\_folder/pdf/New\\_files/EU/EU\\_Study\\_Caste\\_Discrimination\\_merged\\_April2010.pdf](https://idsn.org/wp-content/uploads/user_folder/pdf/New_files/EU/EU_Study_Caste_Discrimination_merged_April2010.pdf)
23. Saroha E, Altarac M, Sibley LM. Caste and Maternal Health Care Service Use Among Rural Hindu Women in Maitha, Uttar Pradesh, India. J Midwifery Womens Health. 2008;53(5):41–7.
24. Saxena D, Vangani R, Mavalankar D V, Thomsen S. Inequity in maternal health care service utilization in Gujarat: analyses of district-level health survey data. Glob Health Action [Internet]. 2013 Jan;6:1–9. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3591509&tool=pmcentrez&rendertype=abstract>

25. Devkota HR, Clarke A, Murray E, Groce N. Do experiences and perceptions about quality of care differ among social groups in Nepal?: A study of maternal healthcare experiences of women with and without disabilities, and Dalit and non-Dalit women. PLoS One [Internet]. 2017;12(12). Available from: <https://doi.org/10.1371/journal.pone.0188554>
26. MoHP/NewEra/ICF International Inc. Nepal Demographic and Health Survey 2011. Kathmandu, Nepal; 2012.
27. Niraula BB. Use of health services in Hill villages in Central Nepal. Heal Transit Rev. 1994;(4):151–66.
28. Davis K. Intersectionality as buzzword: A sociology of science perspective on what makes a feminist theory successful. SAGE Publ [Internet]. 2008;9(1):67–85. Available from: <http://ft.sagepub.com>
29. Buettgen A, Gorman R, Marcia Rioux K Das, Vinayan S. Employment , Poverty , Disability and Gender : A Rights Approach for Women with Disabilities in India, Nepal and Bangladesh. Women's Ment Heal Resist Resil community Soc. 2015;(September):3–18.
30. Bennett L. Gender, Caste and Ethnic Exclusion in Nepal: Following the Policy Process from Analysis to Action. New Front Soc Policy. 2005;1–48.
31. DPHO. Annual Report 2069/70: District Public Health Office, Rupandehi. 2014.
32. Devkota HR. Maternal Health Care Service Access to Disabled and Dalit Women in Nepal [Internet]. University College London (UCL); 2017. Available from: <http://discovery.ucl.ac.uk/1559753/>
33. GoN/MoHP/FHD. National Medical Standard for Reproductive Health; Volume III: Maternal and Neonatal Care. Kathmandu, Nepal; 2009.
34. Malhotra A, Schuler SR, Boender C. Women's empowerment as a variable in International Development. In: Background Paper Prepared for the World Bank Workshop on Poverty and Gender: New Perspectives [Internet]. 2002. p. 71–88. Available from: <http://books.google.com/books?hl=en&lr=&id=BzXyApyTGOYC&pgis=1>
35. Narayan D. Measuring Empowerment: Conceptual Framework and Methodological Challenges [Internet]. Washington, DC: The World Bank; 2005. p. 3–38. Available from: <http://www.worldbank.icebox.ingenta.com/content/wb/2029>
36. GoN/MoHP/DoHS. Annual Report 2069/70 (2012/2013). Vol. 70. Kathmandu, Nepal; 2013.
37. GoN/MoHP/PD. Nepal Population Report 2011. Kathmandu, Nepal; 2011.
38. Khanal V, Adhikari M, Karkee R, Gavidia T. Factors associated with the utilisation of postnatal care services among the mothers of Nepal: analysis of Nepal Demographic and Health Survey 2011. BMC Womens Health [Internet]. 2014 Jan [cited 2014 May 4];14(1):19. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3911793&tool=pmcentrez&rendertype=abstract>
39. MoHP/NewEra/OrgMacro. Nepal Demographic and Health Survey 2006. 2007;
40. Paudel DR, Pitakmanaket O. Utilization of Maternal Health Services in Nepal. JHAS. 2010;1(1):28–37.

41. Shah R, Rehfuess E a, Maskey MK, Fischer R, Bhandari PB, Delius M. Factors affecting institutional delivery in rural Chitwan district of Nepal: a community-based cross-sectional study. *BMC Pregnancy Childbirth* [Internet]. 2015;15:27. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4339648&tool=pmcentrez&rendertype=abstract>
42. Wagle RR, Sabroe S, Nielsen BB. Socioeconomic and physical distance to the maternity hospital as predictors for place of delivery : an observation study from Nepal. *BMC Pregnancy Childbirth* [Internet]. 2004;4(8):1–10. Available from: <http://www.biomedcentral.com/1471-2393/4/8>
43. Karkee R, Lee AH, Pokharel PK. Women’s perception of quality of maternity services: a longitudinal survey in Nepal. *BMC Pregnancy Childbirth* [Internet]. 2014 Jan;14:45. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3902186&tool=pmcentrez&rendertype=abstract>
44. Ahmed S, Creanga A a, Gillespie DG, Tsui AO. Economic status, education and empowerment: implications for maternal health service utilization in developing countries. *PLoS One* [Internet]. 2010 Jan [cited 2014 Apr 29];5(6):e11190. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2890410&tool=pmcentrez&rendertype=abstract>
45. Celik Y, Hotchkiss DR. The socio-economic determinants of maternal health care utilization in Turkey. *Soc Sci Med* [Internet]. 2000 Jun;50(12):1797–806. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10798333>
46. Pandey S, Lama G, Lee H. Effect of women’s empowerment on their utilization of health services: A case of Nepal. *Int Soc Work* [Internet]. 2011 Jul 18 [cited 2013 Jul 2];55(4):554–73. Available from: <http://isw.sagepub.com/cgi/doi/10.1177/0020872811408575>
47. Worku AG, Yalew AW, Afework MF. Factors affecting utilization of skilled maternal care in Northwest Ethiopia: a multilevel analysis. *BMC Int Health Hum Rights* [Internet]. 2013 Jan;13:20. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3639034&tool=pmcentrez&rendertype=abstract>
48. Matsumura M, Gubhaju B. Women’s status, household structure and the utilization of maternal health services in Nepal. *Asia-Pacific Popul J*. 2001;16(1):23–44.
49. Dahiru T. P - Value, A True Test of Statistical Significance? A Cautionary note. *Ann Ibadan Postgrad Med*. 2008;6(1):21–6.