

Article

High NO₂ concentrations measured by passive samplers in Czech cities: Unresolved aftermath of Dieselgate?

Michal Vojtíšek-Lom^{1,2,*}, Miroslav Šuta³, Jitka Sikorová¹ and Radim J. Sram¹

1 Department of Genetic Toxicology and Epigenetics, Institute of Experimental Medicine AS CR, Videnska 1083, 142 20 Prague, Czech Republic

2 Department of Automotive, Combustion Engine and Railway Engineering, Faculty of Mechanical Engineering, Czech Technical University of Prague, Technická 4, 166 07 Prague, Czech Republic

3 Center for Environment and Health, Thámová 1275/21, 301 00 Plzeň, Czech Republic

* Correspondence: michal.vojtisek@fs.cvut.cz, michal.vojtisek@tul.cz, tel. +420 774 262 854

Highlights

NO₂ measured by 104 passive samplers at 65 places in Prague, corrected mean 36 µg/m³

NO₂ increases with traffic intensity corrected for intersections and hills

High NO₂/NO_x ratios and excess NO_x emissions from diesel cars a culprit

Not much improvement after “Dieselgate”

Annual limit of 30 µg/m³ suggested based on health evidence literature review

Abstract: This work examines the effects of two problematic trends in diesel passenger car emissions – increasing NO₂/NO_x ratio by conversion of NO into NO₂ in catalysts and a disparity between the emission limit and the actual emissions in everyday driving – on ambient air quality in Prague. NO₂ concentrations were measured by 104 membrane-closed Palmes passive samplers at 65 locations in Prague in March-April and September-October of 2019. NO₂ concentrations measured by city stations during those periods were comparable with the 2016-2019. The average measured NO₂ concentrations at the selected locations, after correcting for the 18.5% positive bias of samplers co-located with a monitoring station, were 36 µg/m³ (range 16-69 µg/m³, median 35.3 µg/m³), with the EU annual limit of 40 µg/m³ exceeded at 32% of locations. The NO₂ concentrations have correlated well ($R^2=0.76$) with the 2019 average daily vehicle counts, corrected for additional emissions due to uphill travel and intersections. In addition to expected “hot-spots” at busy intersections in the city center, new ones were identified, i.e. along a six-lane road V Holešovičkách. Comparison of data from six monitoring stations during March 15-April 30, 2020 travel restrictions with the same period in 2016-2019 revealed an overall reduction of NO₂ and even a larger reduction of NO. The spatial analysis of data from passive samplers and time analysis of data during the travel restrictions both demonstrate a consistent positive correlation between traffic intensity and NO₂ concentrations along/near the travel path. The slow pace of NO₂ reductions in Prague suggests that stricter vehicle NO_x emission limits, introduced in the last decade or two, have so far failed to sufficiently reduce the ambient NO₂ concentrations, and there is no clear sign of remedy of Dieselgate NO_x excess emissions.

Keywords: NO₂; passive sampler; Dieselgate; Prague; traffic volume; citizen science; air quality; public policy; health effects

1. Introduction

Mobile sources, including on-road vehicles, remain to be one of the largest contributors to the air pollution in most metropolitan areas in Europe, with particulate matter and nitrogen oxides (NO_x, defined as a sum of nitric oxide NO and nitrogen dioxide NO₂) being of highest concern. Outdoor air pollution is now being considered one of the leading causes of premature death [1], with estimated tolls of approximately half a million premature deaths annually in the EU [2], and associated economic damage around 5% of GDP in Central Europe [3]. At the same time, the state-of-the art technology of the internal combustion engine has improved considerably over the last decades. Very low levels of sulfur and metals in the fuel have allowed the introduction of three-way catalysts on spark ignition engines, a common technology used throughout the U.S. over the last four decades with a somewhat delayed deployment in Europe, and the introduction of diesel particle filters on virtually all on-road diesel engines manufactured in the last decade. The emissions of nitrogen oxides, primarily NO, on engines operating with excess air remained a challenge, being ultimately resolved about a decade ago with selective catalytic reduction (SCR) systems on heavy-duty vehicles [4] and more recently also on light-duty vehicles.

In the EU, the concentrations of NO₂, deemed to be more detrimental to human health than NO, are limited and monitored in the ambient air. Overall, the concentrations of NO₂ have not been decreasing as fast as those of other key pollutants. In the Czech Republic, the concentrations of NO₂ at most air quality monitoring stations have been, according to the data in [5], decreasing by on the order of 1% a year over the last two decades. A gradual decrease of NO₂ concentrations in the overall atmosphere above the Czech Republic over the last decade has been also reported from remote sensing satellite measurements [6].

NO₂ in ambient air originates both from direct (primary) emissions and from gradual conversion of NO into NO₂ [7]. While the total emissions of NO_x have been gradually decreasing, there is no apparent trend of a decrease in NO₂ primary emissions over the last 15 years [6]. One of the culprits of high primary NO₂ emissions are diesel vehicles, which have been, over the last two decades, equipped with oxidation catalysts, which convert a considerable portion of NO into NO₂. In the U.S., average NO₂/NO_x ratio in vehicle exhaust (all vehicles, including predominantly gasoline cars and light trucks and predominantly diesel heavy trucks) was 5.3% [8], compared to approximately 15% in Europe [9].

This paper explores a hypothesis that the observed decrease in NO₂ concentrations falls short of that expected based on order-of-magnitude decrease in vehicle NO_x emissions limits, and that non-compliant diesel cars could substantially contribute to this shortfall. The underlying aspects of NO_x emissions and the adverse health effects of NO₂ are summarized. The results of a citizen science campaign aimed at monitoring NO₂ with passive samplers are reported and discussed in light of these findings. As an additional insight, the effects of coronavirus related restrictions on NO and NO₂ concentrations in Prague are reported and discussed.

2. Review of trends and shortcomings in NO₂ and NO_x emissions from vehicles

Nitrogen oxide (NO) is formed in combustion processes from atmospheric nitrogen and oxygen at high temperatures [10, 11], which are generally associated both with efficient combustion and with high thermal efficiency of the engine. Subsequent oxidation of NO in the atmosphere yields primarily nitrogen dioxide (NO₂), a brownish irritant gas. Other oxides of nitrogen – N₂O₂, N₂O₃, N₂O₄, N₂O₅ – are generated in small concentrations, are unstable and short-lived in the atmosphere. The oxides of nitrogen are summarily referred to as NO_x, although there is no precise definition. Often, NO_x is evaluated as the sum of NO and NO₂. Technically, the sum of NO_x also includes nitrous oxide (N₂O), which is, however, not hazardous to human health, but is a potent greenhouse. NO_x lead to the formation of nitrous acid (HNO₂) [12, 13], nitric acid (HNO₃), and a variety of salts such

as ammonium nitrate, present in the atmosphere as particulate matter [14]. Photodissociation of NO_2 under the presence of sunlight produces NO and atomic oxygen, which reacts with molecular oxygen to form ozone [15] a highly reactive compound generally harmful to human health, organisms and plants. NO_x and ground-level (tropospheric) ozone are, together with particulate matter, the principal part of urban air pollution.

On spark ignition engines, CO and VOC, principally a product of incomplete oxidation of fuel and to a lesser extent engine lubricating oil, and NO_x have been successfully abated by the combination of three-way catalysts [16] and by maintaining stoichiometric air-fuel ratio through closed-loop control of the quantity of fuel injected [17]. This technology has proven to be remarkably efficient.

On diesel engines, the emissions of NO_x have been, at first, controlled through delayed combustion timing and exhaust gas recirculation, both associated with a slight fuel penalty, and at a later time, with NO_x storage and reduction catalysts and selective reduction catalysts (SCR). The reduction of NO_x has historically come at an expense of both capital and operating costs, with operating costs including either fuel (notably on older vehicles using delayed combustion, exhaust gas recirculation, NO_x storage and reduction catalysts) or a reducing agent used in SCR (mostly aqueous solution of urea, known as Diesel Exhaust Fluid or „AdBlue“). These costs have motivated, over last few decades, many manufacturers and vehicle users to circumvent NO_x reduction efforts, as the savings were realized by them directly, while considerably larger overall damage to human health was born by the society, a problem known as the Tragedy of the Commons [18]. A widespread practice of dual engine mapping in the U.S. in 1990's [19, 20] has lead to the gradual extension of vehicle emissions limits to ordinary on-road operation first of heavy-duty and later of light-duty vehicles [21-23]. In the heavy-duty vehicle engine sector, many recent studies now show that on-road NO_x emissions of newer heavy-duty vehicles have been successfully reduced by an order of magnitude except for low-load operation typical for congested urban areas. Quiros et al. [24] reports NO_x emissions of 2013 and 2014 model year heavy trucks of 0.36 g/km during motorway operation in California. Jiang et al. [25] reports, for similar conditions, 0.3 g/km NO_x during extraurban and motorway operation. Grigoratos et al. [26] reports NO_x emissions during motorway operation in Europe of 0.07, 0.08, 0.17 and 0.24 g/kWh for four trucks and 0.80 g/kWh for a bus. Giechaskiel et al. [22] reports NO_x emissions of a garbage collection truck of less than 0.4 g/kWh during extraurban operation (note: for heavy vehicles, emissions per kWh roughly correspond to emissions per km).

Unfortunately, this has not been the case with light-duty vehicles with diesel engines, highly prevalent in Europe, where they account for tens of percents of vehicle registration and in Prague, for about two thirds of vehicles counted on the road [27]. Large portion of European automobile diesel engines produced over the last one to two decades have been reported to emit substantially, often by an order of magnitude, more NO_x on the road than during the type approval test [28-32]. Weiss et al. [29] reports on-road NO_x emissions factors 0.76 ± 0.12 g/km for Euro 4, 0.71 ± 0.30 g/km for Euro 5 and 0.21 ± 0.09 for Euro 6. In a more recent study by Suarez-Bertoa et al. [23], NO_x emissions from Euro 6 diesel cars varied substantially from mid tens to mid hundreds of mg NO_x per km, with median value about 0.2 g/km NO_x during city-motorway test.

At the same time, on nearly all light-vehicle diesel engines of the last decade or so, oxidation catalysts are used to convert NO into NO_2 , as higher concentrations of NO_2 , around tens of percent, are beneficial both for the combustion of soot in DPF and for the „fast“ reduction of NO_x in SCR catalysts. As a result, NO_2 from newer engines accounts for tens of percent of NO_x [33, 34]. On passenger cars and light-duty trucks, NO_2/NO_x ratios of around 10-15% up to Euro 3 and 25-30% for Euro 4 and 5 were found in a London remote sensing study [35]. In the U.S., NO_2/NO_x ratio from heavy duty diesel trucks have doubled from around 7% in 2010 (average of trucks passing on the road in a given year, not a model year of the vehicles) to around 15% in 2018 [36]. This increase, however, did not result in an absolute increase in NO_2 emissions, as total NO_x emissions have decreased dramatically thanks to the widespread use of SCR catalysts. According to Preble-

Caldecott [36], „Fleet-average NO₂ emission rates remained about the same, despite the intentional oxidation of engine-out NO to NO₂ in DPF systems, due to the effectiveness of SCR systems in reducing NO_x emissions and mitigating the DPF-related increase in primary NO₂ emissions.“

In Europe, NO_x emissions from diesel cars have not, however, decreased in proportion to the decreasing emissions limits. A recent on-road study in Prague report mean emissions of Euro 5 and 6 diesel cars and vans of over 0.1 g/km NO₂ and over 0.5 g/km NO_x [37], while a recent study of one of the most common diesel cars (Euro 6) reported about 0.15 g/km over WLTC cycle, and about 0.4 g/km over the Artemis driving cycle [38], which is more than the 0.08 g/km Euro 6 limit for total NO_x (with which the vehicle reasonably complied over the NEDC cycle).

The presumption of the regulators that increased NO₂/NO_x ratio after the oxidation catalyst and before the DPF, highly beneficial both for DPF and SCR operation, will be mitigated by the rather high efficiency of the NO_x aftertreatment, envisioned in both U.S. EPA and EU emissions standards, has been compromised by intentional acts resulting in diminished, or even zero, efficiency of NO_x aftertreatment. Examples of such acts include dual-mapping of the engines by the manufacturers (a prime example of which is „Diesel-gate“) and disabling of the SCR (and emulating its proper functioning to the on-board diagnostics by „SCR emulators“) by vehicle operators. Under such conditions, relatively high amounts of NO₂, intended to be reduced in NO_x aftertreatment, are emitted out of the tailpipe. Logically, this results in very high, and much higher than intended, primary emissions of NO₂ in the streets. This finding is consistent with the rather slow decrease in NO₂ concentrations.

To build up on this hypothesis, the measurements of NO₂ concentrations at various locations by passive samplers are examined. Some of the results were presented by Deutsche Umwelthilfe [39] as preliminary data; in this study, the results from Prague were examined in a greater detail.

3. Measurement of NO₂ in Prague by passive samplers

For passive monitoring, membrane-closed Palmes tube [40] passive samplers (Passam, Switzerland [41]) were used. Several hundreds of samplers were placed at selected locations in the Czech Republic, out of which 65 were in Prague, during spring and fall of 2019 (46 and 58 samplers, respectively, a total of 104 samplers), each time for a period of approximately one month. The placement of the tubes followed the requirements set in the EU Air Quality Directive (2008/50). The measured concentrations, along with average daily vehicle traffic counts [42], are reported in Table 1.

Table 1. Measured NO₂ concentrations and average daily vehicle counts.

NO2 measurements by passive samplers	Spring measurement period	Concentration as analyzed [ug/m ³]				Adjusted (div 1.185) concentrations			Traffic vehicles/day		Hill climb	Inter-section	>6 tons excl. zone
		Mar-Apr	Aug30-Sep29	Sep7-Oct30	Sep29-Oct30	Spring	Fall	Average	Actual	Adjusted			
31 Budějovická	Mar 9-Apr 6	33.6				28.3		28.3					1
32 třída 5. května 39	Mar 9-Apr 6	42.9				36.2	34.8	35.5	73818	110727	50%		1
33 Na Veselí	Mar 9-Apr 6	48.7			41.2	41.1	34.5	37.8	15500	31000	100%		1
34 Sokolská/Ječná	Mar 9-Apr 6	78.4			40.9	66.2	56.1	61.1	56000	280000	100%	100%	1
35 Ječná/Štěpánská	Mar 9-Apr 6	63.6	70.4		62.6	53.6	52.9	53.3	27600	138000	100%	100%	1
36 Jugoslávských partyzánů 27	Mar 9-Apr 6	34.6		62.7		29.2		29.2	16723	16723			
37 Na píškách/Evropská	Mar 9-Apr 6	52.0				43.9	47.6	45.7	40600	162400		100%	
38 Kačkova/Svatovítská	Mar 9-Apr 6	46.1		56.4		38.9	39.0	39.0	26101	104404		100%	
39 Svatovítská / tunel	Mar 9-Apr 6	31.0		46.2		26.1	28.5	27.3	36901	36901			
40 Na Ořechovce	Mar 9-Apr 6	45.3		33.8		38.2		38.2	12800	12800			
41 Dejvice train station	Mar 9-Apr 6	73.4				62.0	50.2	56.1	29200	131400	50%	100%	1
42 Hradčanská (metro station)	Mar 9-Apr 6	34.2		59.5		28.8	30.3	29.6	18409	18409			1
43 Veletržní/Sochařská	Mar 9-Apr 6	50.5		36.0		42.6	39.9	41.2	22100	99450	50%	100%	1
44 Janovského/Veletržní	Mar 9-Apr 6	40.5		47.3		34.2	28.5	31.4	19400	77600		100%	1
45 Křižovnická	Mar 9-Apr 6	40.1		33.8		33.8		33.8	21000	21000			1
46 Vinohradská/Flora	Mar 9-Apr 6	34.2				28.8	31.0	29.9	26400	26400			
47 Flora-mall (bus stop)	Mar 9-Apr 6	42.6			36.7	35.9	29.8	32.8	11312	45248		100%	
48 Bělocerkevská (bus stop)	Mar 9-Apr 6	51.4			35.3	43.4	39.2	41.3	26500	132500	100%	100%	
49 Vršovická (Slavia tram stop)	Mar 9-Apr 6	32.9			46.5	27.8	30.6	29.2	13900	55600		100%	
52 Rumunská/Sokolská	Mar 9-Apr 6	53.4			36.2	45.0		45.0	43100	129300	50%	50%	1
120 Severní Spořilov podchod	Mar 13-Apr 24	45.4				38.3		38.3	48900	73350	50%		
121 Chodov / Dálnice	Mar 13-Apr 24	54.7				46.2		46.2	118100	177150	50%		
122 Zenklova/Na Korábě	Mar 13-Apr 24	39.4				33.2	25.2	29.2	13000	13000			
123 Vychovatelna (bus)	Mar 13-Apr 24	67.0			29.8	56.5	41.0	48.7	109300	163950	50%		
124 Rokoska (podchod)	Mar 13-Apr 24	63.6			48.5	53.7	44.8	49.3	88561	132842	50%		
125 V Holešovičkách 8/10	Mar 13-Apr 24	50.7			53.1	42.8	37.8	40.3	88561	132842	50%		
126 Hotel Paw llovia	Mar 13-Apr 24	40.4			44.8	34.1	36.4	35.3	88561	88561			
127 main train station	Mar 13-Apr 24	41.9			43.2	35.3	43.4	39.4	85053	85053			1
128 Hrusická 6 (balcony)	Mar 13-Apr 24	21.2	51.4			17.9		17.9	0	0			
129 hlavní 25 (balcony)	Mar 13-Apr 24	29.1				24.6		24.6	8000	8000			
130 Havní / most	Mar 13-Apr 24	36.7				31.0		31.0	50487	75730.5	50%		
181 Kotevní 2	Mar 19-Apr 24	32.2				27.1		27.1	26500	26500			1
182 Strakonická 21/23	Mar 19-Apr 24	40.9				34.5		34.5	54753	54753			1
183 Svornosti 19a	Mar 19-Apr 24	48.2				40.6		40.6	11800	11800			1
184 Zborovská 3	Mar 19-Apr 24	48.2				40.7	37.1	38.9	14500	58000		100%	1
185 V Botanice 4 (regional government)	Mar 19-Apr 24	55.8	43.7		44.2	47.1	47.1	47.1	25028	100112		100%	1
186 V Botanice (bank)	Mar 19-Apr 24	43.3	48.7		62.9	36.5	37.1	36.8	22000	88000		100%	1
187 Ptežská 14, Hotel IBIS	Mar 19-Apr 24	49.2			43.9	41.5	35.4	38.4	32700	130800		100%	
188 Radlická 14/Anděl	Mar 19-Apr 24	47.9			41.9	40.5	40.8	40.6	25030	100120		100%	
189 Ostrovského	Mar 19-Apr 24	43.2			48.3	36.5	34.4	35.4	23190.5	92762		100%	
190 Billa Karlín	Mar 19-Apr 24	32.2	40.8			27.2	23.8	25.5					
191 Pobřeží (business center)	Mar 19-Apr 24	43.5		28.2		36.7	33.5	35.1	31200	31200			
192 Pobřeží (monitoring station)	Mar 19-Apr 24	38.3		39.7		32.3	25.7	29.0	31200	31200			
193 Negrelliho viadukt	Mar 19-Apr 24	33.3		30.5		28.1	32.8	30.4	13335	13335			
194 Florenc (bus stop)	Mar 19-Apr 24	46.4		38.8		39.2	35.6	37.4	14612	58448		100%	
195 Nám. Republiky (Kotva)	Mar 19-Apr 24	47.2		42.2		39.8		39.8	8300	33200		100%	1
Mezibraná 3	none		84.4		78.9		68.9	68.9	59645	298225	100%	100%	1
Sokolská/Ječná , Prague	none		74.0		62.6		57.6	57.6	55445	277225	100%	100%	1
Rumunská/Legerova, Prague	none		62.3		52.1		48.3	48.3	45452	181808		100%	1
Bubenská , Prague	none			48.0			40.5	40.5					
Vysočanská , Prague	none			26.1			22.0	22.0					
Vysočanská (ČHMÚ), Prague	none			37.1			31.3	31.3					
Thámová / Sokolovská , Prague	none			28.2			23.8	23.8					
Radlická (ČSOB), Prague	none			37.6			31.7	31.7					
Radlická (Kotelna Park) , Prague	none			33.4			28.2	28.2					
Resslova 1/3, Prague	none			52.4			44.2	44.2	33027	148622	50%	100%	
Spořilov 1, Prague	none				51.3		43.3	43.3					
Spořilov 2, Prague	none				33.5		28.3	28.3					
Boční / Jihovýchodní V II, Prague	none				28.1		23.7	23.7					
Pankrác 1 BAUHAUS, Prague	none				37.1		31.3	31.3				100%	
Pankrác 2 Doudlebská , Prague	none				29.2		24.7	24.7				100%	
Pankrác 3 viadukt , Prague	none				31.8		26.8	26.8				100%	
Pankrác 4 Hvězdova 35, Prague	none				31.1		26.3	26.3				100%	
Radlická / Klicperova, Prague	none				48.3		40.8	40.8	25030	100120		100%	
Suchdol AV ČR, Prague	none				19.9		16.8	16.8					
Suchdol AV ČR, Prague	none				18.7		15.7	15.7					
NO2 measurements by the national air quality monitoring network		Average of 1-hour concentrations [ug/m ³]				Average concentrations			Traffic vehicles/day		Hill climb	Inter-section	>6 tons excl. zone
Station	Mar9-Apr6	Mar19-Apr21	Aug30-Sep29	Sep7-Oct30	Sep29-Oct30	Spring	Fall	2016-2019	Actual	Adjusted			
Legerova	45.52	61.5	44.7		44.7	53.5	44.7	51.1	46300	185200		100%	1
Náměstí Republiky	29.1	34.9	26.0		35.8	32.0	30.9	29.6	10400	41600		100%	1
Kobylisy	19.9	20.9			26.4	20.4	26.4	20.0	0	0			
Průmyslová	31.3	31.8			29.7	31.6	29.7	31.4	35000	35000			
Vysočanská	28.7	37.0		31.4		32.9	31.4	34.8	37035	37035			
Karlín		31.9		26.0		31.9	26.0	29.2	31200	31200			

3.1 Validation by comparison with the air quality monitoring network.

According to [43], passive diffusion tubes for measuring NO₂ concentrations in air were originally developed in the late 1970s for personal monitoring. They have been widely used in Europe for spatial and temporal measurement of NO₂ concentrations. The method has been found to be cheap, simple, and „provides concentration data in most circumstances that are sufficiently accurate for assessing exposure and compliance with Air Quality criteria.“ [43] Reporting on a series of comparison tests, Buzica et al. [44] have concluded that „In the case of NO₂, all the results of the laboratory and field experiments respected the requirements necessary for the demonstration of equivalence.“ and that the MCPT are equivalent to the reference methods for assessment of NO₂. Passive diffusion tubes were reported to show a positive bias when sampling close to sources of NO, such as roadside or street canyons [43]. At the same time, prolonged (several weeks) sampling periods were reported to lead to negative bias [43].

To evaluate the bias, two samplers were placed at the Legerova monitoring station (urban hotspot), but both were stolen. Several samplers from nearby locations, and additional samplers from locations with similar traffic flow as on Náměstí Republiky, were compared with the online data from the Legerova and Náměstí Republiky monitoring stations (data shown in Table 1), in each case averaged over the actual sampling period of each respective sampler. The results of this comparison are given in Figure 1. Two larger points represent one sampler co-located near the Náměstí Republiky station and another one relatively close to the Legerova station. Additional samplers close to the Legerova station were closer to intersections and therefore exposed to additional cross-traffic, in addition to the increase in emissions rates in the vicinity of intersections. To evaluate the bias, two samplers were placed at the Legerova monitoring station (urban hotspot) in the spring of 2019, but both were stolen. Additional samplers were placed near the Karlín monitoring station and near the Náměstí Republiky monitoring stations, as well as in the general vicinity of the Legerova station. The NO₂ concentrations reported for the samplers were compared with the average NO₂ concentrations measured by the monitoring station, obtained by averaging data over the time the samplers were exposed on the site.

The results of this comparison are given in Figure 1. The three larger points (in red/orange) represent samplers co-located with the Karlín monitoring station over three separate one-month periods show a linear correlation with a slope of 1.185. Blue points show additional locations. Two samplers were placed at an urban background monitoring station Suchbátka, however, data from this station was not available, and the readings are compared with another background monitoring station in Kobylisy. Two samplers were placed near Náměstí Republiky monitoring station, but a few dozens meters away and near an exit/entrance ramp to a large shopping center underground parking garage. Two samplers were placed on the corner of Legerova and Rumunská, near the monitoring station but at an intersection controlled by a traffic light. The readings from these four samplers were higher than from the monitoring station, which can be reasonably expected as they were near stopped and accelerating vehicles.

Additional samplers used in the comparison were at reasonably close locations with not overly dissimilar traffic, and were not too far from the 15% tolerance reported by the Defra report [43]. It should be noted that the tolerance is applicable to the deviation of the sampler-reported and reference value, and not to the differences due to the samplers being at different locations with different emissions characteristics.

For all subsequent data analysis, the concentrations from the passive samplers were divided by the regression slope of 1.185.




Figure 1. Comparison of passive sampler reported NO₂ concentrations to the corresponding average values from corresponding monitoring stations. Larger points circled in red denote co-location of the sampler at the monitoring station.

3.2 Comparison of NO₂ during passive samplers deployment with long-term averages

The variation of climatic and weather conditions is an additional source of bias to consider when comparing passive samplers to annual mean values. Fig. 2 shows that the average values of NO₂ recorded at the monitoring stations over sampling periods of individual samplers (different four-week periods in March-April 2019) did not dramatically differ from annual means during the last four years (2016-2019), although differences in trends were observed among the stations. For example, the Legerova urban hotspot station exhibited an annual average of 51.1 µg/m³ (2016-2019), compared to 45.5 µg/m³ during the period of March 9 – April 6 and 61.5 µg/m³ during March 19 – April 24. The Náměstí Republiky urban background station had a 2016-2019 average of 29.6 µg/m³, compared to 29.1 µg/m³ during Mar 9 – Apr 6 and 34.9 µg/m³ during Mar 19 – Apr 24.




Figure 2. Comparison of monitoring station NO₂ averages during sampling periods with four-year average.

The consistency of the measurement by passive samplers during spring and fall periods is shown, along with data from the reference monitoring stations, in Fig. 3. The slope

of regression is 0.908 for the monitoring stations and 0.921 for the passive samplers, showing that the monitoring stations and the passive samplers report the same overall trends in NO₂ concentrations.




Figure 3. Comparison of spring and fall NO₂ concentrations.

3.3 Effects of traffic

For further analysis, all passive sampler measurements were divided by a factor of 1.17 (the slope of regression of passive sampler vs. reference NO₂, see Fig. 1).

The relationship between the vehicular traffic intensity and the NO₂ concentrations measured by the passive samplers is given in Fig. 4. As samplers were used over two different periods, they are plotted separately in two series, one for each period, along with the average values from Legerova and Náměstí Republiky monitoring stations. It appears that there is a moderate positive trend of NO₂ increasing with traffic. Also, samplers located next to uphill section of a divided highway (or a one-way street with the traffic going in the uphill direction) and next to an intersection tend to exhibit higher NO₂ concentrations. It also appears that the NO₂ concentrations are higher in urban canyons and congested streets of the city center and near intersections.

To assess whether high NO₂ are associated with truck traffic, samplers located in the area with limited access of vehicles over 6 tons gross weight (entry by permit only, restricted to local traffic) are plotted separately in Fig. 5 (for locations where multiple samplers were used, average values are plotted). It is clear from the figure that the highest NO₂ were measured in areas where trucks over 6 tons are mostly excluded.




Figure 4. Relationship between traffic intensity and NO2 concentrations measured by passive samplers in spring and fall of 2019 and by the national monitoring network (average of 2016-2019).




Figure 5. Relationship between traffic intensity and NO2 concentrations measured by passive samplers (average of all measurement periods) and by the national monitoring network (average of 2016-2019).

To account for additional emissions due to hills and intersections, the intensity of traffic traveling uphill was increased by 100% to account for additional fuel consumption, and for samplers located at intersections, the intensity of traffic was increased by 300% to account for fuel consumed at idle and when accelerating (where the intersection was without a major delay, such as time-synchronized signals at intersections of a larger one-way street with a side street or pedestrian crossing, the factor was reduced by one half). These adjustment factors were arbitrarily selected based on experience with vehicle emissions behaviour (additional emissions due to climbing a hill, additional emissions due to idling at intersections and acceleration from intersections) and were independent of each other. The relationship between the adjusted vehicle volume and NO₂ concentrations is plotted in Figure 6.

The relatively strong correlation between the adjusted traffic volumes and NO₂ concentrations ($R_2 = 0.78$ for September-October data and 0.76 for spring-fall averages) suggests that „local“ NO₂, comprising of primary NO₂ emitted from the tailpipe and NO₂ formed locally from NO by reaction with ozone (i.e., [45]), is a considerable and in many locations dominant source of NO₂. There is no observable difference between the sampling locations where truck traffic over 6 tons was excluded and the locations where it was not excluded. Overall, there seems to be a very strong correlation between the estimated relative intensity of mobile source emissions and the measured NO₂ concentrations. It is likely that the correlation could be further improved by taking into the account distance from the traffic, traffic on adjacent streets, tunnel exits, and other compounding factors.

A similar plot of the regression of the dependency of NO₂ on adjusted traffic volumes is plotted separately for the spring and fall campaigns in Fig. 7, with red line denoting the legal annual NO₂ limit of 40 µg/m³ and green line the recommended alternative limit of 30 µg/m³ proposed by the last author and explained later in this manuscript. The regression shows that NO₂ concentrations, in all cases, increase by 0.13 µg/m³ per 1000 vehicles daily traffic volume, adjusted for uphill and intersections, where adjusted traffic count is traffic count multiplied by a factor of $(1 + \text{fraction of vehicles travelling uphill} + 3 * \text{fraction of vehicles stopping at an intersection})$. It should be noted that the intercept of the regression (25-28 µg/m³ in Fig. 6 and 7) is higher than the „urban background“ concentrations of 15-20 µg/m³, and that even the urban background concentrations cannot be considered as NO₂ concentrations that would be theoretically be expected if no motor vehicles were operated in Prague, due to the dispersion of the pollutants.

Even at a rather conservative adjustment of the passive sampler readings (according to the regression, the sampler readings were 18% higher, however, this was, to a large extent, due to many samplers being at locations where the concentrations would reasonably be expected to be higher than at the corresponding monitoring station), it is clear from Fig. 7 that the annual average limit of 40 µg/m³ NO₂ is likely to be exceeded at numerous locations throughout Prague, generally, where the adjusted traffic volumes exceed the equivalent of 100 thousands of vehicles per day. This is, for example, the north-south passageway through the center city (Wilsonova, Sokolská and Legerova Street) with many intersections, but also roads like V Holešovičkách (a six-lane road with 85-90 thousands of vehicles per day, with a gradient of approximately 3%), a possible new hot-spot in Prague. In the worst case (intersection of two one-way streets with all vehicles traveling uphill), this limit could be reached already at 20 thousands of vehicles per day, as also apparent from Fig. 5.




Figure 6. Relationship between adjusted traffic intensity (traffic count * (1 + fraction of vehicles travelling uphill + 3 * fraction of vehicles stopping at an intersection)) and NO₂ concentrations measured by passive samplers (average of all measurement periods) and by the national monitoring network (average of 2016-2019).




Figure 7. Relationship between adjusted traffic intensity (traffic count * (1 + fraction of vehicles travelling uphill + 3 * fraction of vehicles stopping at an intersection)) and NO₂ concentrations measured by passive samplers (average of all measurement periods) and by the national monitoring network (average of 2016-2019).

4. Effects of travel restrictions on ambient NO and NO₂ concentrations

In order to assess the contribution of light and heavy vehicles to NO and NO₂ concentrations, hour-by-hour NO and NO₂ ambient air quality data from the national air quality monitoring network was analyzed for a period of March 14-April 30, 2020, during which travel restrictions were imposed, including the prohibition of all non-cargo international travel (truck traffic was exempted). For reference, the same period was assessed for four previous years.

A total of five stations in Prague were selected:

- a. Legerova street, considered an urban hotspot, with about 45 thousands of vehicles traveling daily in one direction (with similar traffic volumes in the opposite direction on a parallel street), primarily (97-98%) light-duty vehicles (trucks over 12 tons are restricted from entering inner Prague and trucks over 6 tons are restricted in the Prague historical district);
- b. Vysočanská street and Průmyslová street, two traffic stations located on heavily traveled main roads used by local as well as transit truck traffic;
- c. Náměstí Republiky, urban background station in a historical city center, on the border of pedestrian area
- d. Kobylišy, a station in a suburban residential neighborhood
- e. For comparison, a rural background station in Košetice, serving as the Czech national reference station, was used as a reference.

Arithmetic and geometric means and the NO_2/NO_x ratios are plotted, for each station and all years, in Table 2. Analysis of variance (ANOVA) was performed to compare the variances among the four reference years (2016-2019) with the variance of the year 2020 from the reference years. Differences between 2020 and the previous years are shown in the table, along with stars representing the significance level.

Table 2. Arithmetic and geometric means and the NO₂/NO_x ratios for each station and all years.

Station	March 14-April 30 year	ug/m3, arithmetic mean			ug/m3, geometric mean			ratio NO ₂ /NO _x
		NO	NO ₂	NO _x	NO	NO ₂	NO _x	
Legerova <i>type: traffic</i> <i>predominantly</i> <i>light-duty < 3.5 tons</i>	2016	43.5±47.2	55.2±27.4	122.0±95.1	24.8±3.1	48.4±1.7	91.7±2.2	55%±16%
	2017	35.4±38.6	46.5±28.3	100.8±84.7	17.1±4.0	36.8±2.1	67.4±2.7	57%±16%
	2018	44.7±46.2	59.3±29.0	128.0±94.3	24.6±3.4	51.4±1.8	95.5±2.3	57%±17%
	2019	36.9±38.2	55.0±27.2	111.7±81.2	21.6±3.1	46.8±1.9	83.9±2.3	58%±14%
	2020	21.6±27.7	43.2±21.0	76.5±59.3	12.2±2.9	38.4±1.6	60.5±2.0	66%±15%
	2020 vs. 2016-2019	- 46% ****	- 20% ****	- 34% ****	- 44% ****	- 16% ****	- 28% ****	+ 23% ****
Průmyslová <i>type: traffic</i> <i>all types, truck transit</i>	2016	24.8±40.1	34.6±19.8	72.8±77.2	9.1±4.8	29.3±1.8	48.7±2.4	64%±20%
	2017	21.9±33.2	33.4±20.2	67.0±67.9	8.1±4.8	27.3±1.9	44.3±2.5	65%±19%
	2018	21.4±35.8	31.8±22.0	64.7±72.4	6.5±5.5	24.1±2.2	38.1±2.9	67%±20%
	2019	19.7±39.0	30.6±21.3	60.8±77.3	5.8±5.2	24.3±2.0	37.1±2.6	69%±19%
	2020	16.0±29.0	27.5±19.4	52.0±60.1	5.5±4.3	21.0±2.2	31.8±2.7	69%±17%
	2020 vs. 2016-2019	- 27% ****	- 15% ****	- 22% ****	- 24% ****	- 20% ****	- 24% ****	+ 6% **
Vysočanská <i>type: traffic</i> <i>all types, truck transit</i>	2016	22.7±29.5	38.0±18.9	72.9±60.4	12.0±3.3	33.5±1.7	55.7±2.1	63%±16%
	2017	18.4±26.6	35.1±19.1	63.5±56.5	8.2±3.9	30.3±1.7	46.6±2.2	68%±17%
	2018	18.8±25.1	36.0±19.9	64.9±54.7	7.8±4.3	30.5±1.8	46.9±2.3	68%±18%
	2019	17.4±22.3	34.1±19.1	60.8±49.9	8.7±3.5	28.9±1.8	45.5±2.2	66%±16%
	2020	14.2±19.9	33.2±18.9	55.1±45.0	7.0±3.3	28.1±1.8	41.8±2.1	70%±16%
	2020 vs. 2016-2019	- 27% ****	- 7% ****	- 16% ****	- 23% ****	- 9% ****	- 14% ****	+ 8% ****
Náměstí Republiky <i>type: urban</i> <i>background</i>	2016	12.0±14.0	20.2±7.1	38.8±26.4	6.9±3.2	19.2±1.4	33.3±1.7	59%±26%
	2017	12.1±12.5	33.1±14.6	51.7±30.8	9.4±1.9	30.4±1.5	46.0±1.6	66%±15%
	2018	15.6±19.5	35.2±17.7	59.1±43.7	9.8±2.6	31.5±1.6	49.1±1.8	65%±18%
	2019	10.9±14.2	31.9±15.2	48.7±33.5	7.5±2.1	28.9±1.5	41.9±1.7	70%±13%
	2020	10.8±10.6	27.8±14.5	44.6±28.2	8.0±2.1	24.9±1.6	38.4±1.7	66%±12%
	2020 vs. 2016-2019	- 14% **	- 7% ****	- 10% ****	- 3% ****	- 8% ****	- 9% ****	+ 2%
Kobylisy <i>type: residential</i> <i>background</i>	2016	3.8±9.3	10.4±6.3	16.3±19.0	1.2±3.4	9.1±1.7	11.9±2.0	80%±16%
	2017	3.7±9.4	14.5±8.7	19.7±19.9	1.5±3.1	12.7±1.7	15.4±1.9	80%±16%
	2018	3.7±8.8	21.7±15.9	27.5±26.0	1.4±3.2	17.2±1.9	20.4±2.1	86%±11%
	2019	3.4±8.8	19.6±15.7	25.0±27.1	1.1±3.2	15.8±1.9	18.4±2.0	87%±14%
	2020	2.8±5.9	17.3±14.1	21.0±20.8	1.5±2.3	13.0±2.1	14.8±2.2	81%±14%
	2020 vs. 2016-2019	- 22% *	+ 4% **	- 5%	+ 14% **	- 2%	- 8%	- 6% ****
Košetice <i>national reference</i> <i>background</i> <i>outside of Prague</i>	2016	0.5±0.6	6.0±2.6	6.8±3.1	0.3±2.0	5.4±1.6	6.2±1.6	90%±7%
	2017	0.3±0.4	7.3±3.0	7.8±3.2	0.3±1.8	6.7±1.5	7.2±1.5	93%±5%
	2018	0.3±0.4	3.9±2.7	4.3±3.0	0.2±2.6	3.1±2.0	3.5±1.9	90%±9%
	2019	0.2±0.3	3.6±1.9	4.0±2.1	0.1±2.9	3.1±1.8	3.5±1.7	91%±9%
	2020	0.2±0.3	3.1±1.7	3.5±1.9	0.1±2.8	2.7±1.8	3.0±1.8	90%±9%
	2020 vs. 2016-2019	- 29% ****	- 40% ****	- 40% ****	- 35% ****	- 39% ****	- 38% ****	- 2.5% ****

* p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001

It is apparent from the Table 2 that NO concentrations have significantly decreased at all locations, with highest mean decrease of 46% at Legerova. The decrease in NO₂ concentrations was lower than for NO at all Prague station, highest at Legerova (15%), and even higher (40%) at the Košetice rural background station. As vehicles emit primarily NO, the NO₂/NO_x ratio tends to increase with the age of the emissions, being lowest (around 60%) at Legerova street, 65-70% at Vysočanská, Průmyslová and Náměstí Republiky, 80% at the Kobylisy residential background station, and around 90% at the rural station in Košetice. One possible interpretation of the increase in the NO₂/NO_x ratio at Legerova could be that the primary emissions of both NO and NO₂ were reduced, with lower reduction in „background“ NO₂ originating from NO_x emitted elsewhere.

It should be noted, however, that the interplay of different factors is rather complex. For example, diminished traffic volumes result in lower frequency of low-speed driving in congested areas, during which the efficiency of exhaust aftertreatment is reduced, resulting in higher overall exhaust temperatures (and thus higher production of NO₂ in oxidation catalysts), but also higher probability of SCR functionality (and thus lower NO_x emissions) – however, thanks to Dieselgate, the reality of NO_x aftertreatment efficiency is likely to be variable, questionable, and poorly known.

Also, according to [46], it appears that on-road oxidation of NO by ambient O₃ is a significant, but so far ignored, contributor to curbside and near-road NO₂. This is in agreement with on-road NO₂/NO_x ratios in U.S. being reported to be 25-35% and substantially higher than anticipated tailpipe emissions rates [47].

5. Review of the impact of NO₂ to central nervous system in children and adults

The first experimental data were obtained several decades ago, indicating that air pollution may induce behavioral changes. Singh [48] studied the effect of NO₂ exposure on pregnant mice, exposed during gestation day 7 to 18. Prenatal exposure significantly altered the righting reflex and aerial righting score. These results suggest that maternal NO₂ exposure produce deficits in the functional capability of the offsprings.

Wang et al. [49] was the first one, who studied the impact of NO₂ exposure to children's neurobehavioral changes. They studied this effect in the year 2005 on two groups of children (A N=431, B N=430) in the age 8-10 years) using neurobehavioral testing. Group A was exposed to 7 µg NO₂/m³, group B to 36 µg NO₂/m³. Children from polluted area showed poor performance in all tests: visual simple reaction time, continuous performance, digit symbol, pursuit aiming, and sign register. This study found a significant relationship between chronic low-level traffic related air pollution and neurobehavioral function in exposed children.

Guxens et al. [50] analyzed the association between prenatal exposure, diet, and infant mental development in four regions in Spain, in 1 889 children, who were exposed to 29.0±11.2 µg NO₂/m³ (20.1-36.8). Infant mental development was evaluated at 14 months by Bailey Scales of Mental Development. Exposure to NO₂ did not show a significant association with mental development. Inverse association was observed in infants whose mothers reported low intake of fruit/vegetables during pregnancy (-4.13 (-7.06, -1.21)). This study suggests that antioxidants in fruits and vegetables during pregnancy may modulate adverse effect of NO₂ on infants mental development.

Kim et al. [51] investigated the association between maternal exposure to NO₂ 49.4 µg /m³ (25.9-84.8) and neurodevelopment in children in Korea (mental development index (MDI) and psychomotor development index (PDI) by Bailey Scales of Mental Development) at ages 6, 12 and 24 months. Study completed 455 - 371 children. NO₂ exposure impaired psychomotor development (β = -1.30; p = 0.05). At 6 months NO₂ affected MDI (β = -3.12; p<0.001) as well as PDI (β = -3.01; p<0.001). These data suggest that exposure to NO₂ may delay neurodevelopment in early childhood.

Similar study was organized in Spain on 438 mother-child pairs by Lertxundi et al. [51] at 15 months of age, using Bailey Scales of Mental Development. 1 µg NO₂/m³ increase during pregnancy decreased mental score (β = - 0.29; 90% CI: -0.47; -0.11). Prenatal residential exposure to NO₂ adversely affects infant motor and cognitive development.

A prospective cohort study was conducted with 2 715 children aged 7 to 10 years in Barcelona, Spain, as a part of the BREATHE project (Brain dEvelopment and Air polluTion ultrafine particles in scHool childrEn ([53])). Children were tested each 3 months with computerized test. Cognitive development was assessed with the n-back and the attentional network test as working memory and inattentiveness. NO₂ exposure was outdoor in low traffic region 40.5±9.6 µg/m³, high traffic region 56.1±11.5 µg/m³. Children attending schools with higher NO₂ pollution had for 11.5% (95% CI 8.9%-12.5%) slower working memory and slower growth in all cognitive measurements, which means a smaller improvement in cognitive development.

Pujol et al. [54] selected from this cohort 263 children, aged 8 to 12 years, for magnetic resonance investigation (MRI) to analyze brain volumes, tissue composition, myelination, cortical thickness, neural tract architecture, membrane metabolites, functional connectivity. Outdoor NO₂ exposure was 46.8±12.0 µg/m³/year, indoor NO₂ exposure 29.4±11.7 µg/m³/year. Higher NO₂ exposure was associated with slower brain maturation with changes specifically concerning the functional domain.

Forns et al. [55] evaluated 2 897 children from the Barcelona cohort within BREATHE project. NO₂ exposure in schools was 29.82 µg/m³ (11.47-65.65), outdoor 48.46 µg/m³ (25.92-84.55). Behavioral development was assessed using the Strengths and Difficulties Questionnaire (SDQ), which was filled out by parents. NO₂ exposure was positively associated with SDQ total difficulties scores, suggesting more frequent behavioral problems. This study was understood as the first one to evaluate the impact of air pollution on behavioral development in schoolchildren using both indoor and outdoor air pollution levels measured at schools. NO₂ outdoor levels (IQR=22.26 µg/m³) significantly increased total difficulties score (1.07, 95%CI: 1.01, 1.14, $p < 0.05$). NO₂ exposure at school is associated with worse general behavioral development in schoolchildren.

Min and Min [56] studied in Korea 8 936 children born in the year 2002 and followed for next 10 years, relationship between exposure to NO₂ and attention-deficit hyperactive disorder (ADHD). They diagnosed 313 children with ADHD. With the increase in 1 µg NO₂/m³ hazard ratio (HR) was 1.03 (95% CI: 1.02-1.04). Comparing infants with lowest tertile of NO₂ exposure with the highest tertile of NO₂, HR = 2.10 (95% CI: 1.54-2.85), exposure had 2 fold increased risk of ADHD. Study showed a significant association between exposure to NO₂ and the incidence of ADHD in children.

Sentis et al. [57] evaluated prenatal and postnatal exposure to NO₂ and attentional function in children at 4-5 years of age in four regions of Spain (N=1 298). Attentional function was evaluated by the Kiddie-Conners Continuous Performance Test (K-CPT). The prenatal NO₂ level was 31.1 µg/m³ (18.4 - 37.9). Higher exposure to prenatal levels of NO₂ was associated with 1.12 ms (95% CI: 0.22, 2.02) increase in hit reaction time and 6% increase in the number of emission errors (95% CI: 1.01, 1.11) per 10 µg/m³ increase in prenatal NO₂. Higher exposure to NO₂ during pregnancy is associated with impaired attentional function, especially increased inattentiveness in children aged 4-5 years. This reduced attentional function in population could lead to poorer educational indicators. It seems to be important, that this effect was observed with NO₂ concentrations lower than EU standard 40 µg/m³.

Sunyer et al. [58] followed in 2012-2013 2 687 school children from Barcelona, assessing children's attention process 4 times every three months, using Attention Network test (ANT). NO₂ indoor pollution was 30.09±9.51 µg/m³, ambient air pollution 37.75±18.41 µg/m³. Daily ambient levels were negatively associated with all attention processes (children in the bottom quartile of daily exposure to NO₂ had a 14.8 msec (95% CI: 11.2, 18.4) faster response time than those in the top quartile, which corresponds to 1.1 month delay (95%CI: 0.84, 1.37) in natural development). Short-term exposure to NO₂ is associated with potential harmful effects on neurodevelopment.

Forns et al. [59] examined after 3.5 years the cohort of children from Barcelona (N=1 439), whose cognitive development was evaluated 4 times in the years 2012/2013 [52]. Working memory was estimated by computerized n-back test. Exposure to NO₂ was related to the slower development of working memory (β = - 4.22, 95% CI: -6.22, -2.22). These reductions corresponded to - 20% (95% CI: -30.1, -10.7) change in annual working memory development associated with one interquartile range increase in outdoor NO₂. Forns et al. [58] observed persistent negative association between NO₂ levels at school and cognitive development over a course of 3.5 years. Therefore they suggested, that highly exposed children might face obstacles to fully achieve their academic goals.

Vert et al. [60] analyzed association between exposure to NO₂ and mental disorders on 958 residents from Barcelona (45-74 years old). Long-term residential exposure (period 2009-2014) was related to patients self-reported history of anxiety and depression disorders. NO₂ exposure corresponded to 57.3 µg/m³ (50.7 - 62.7). NO₂ increased odd ratio for depression of 2.00 (95%CI: 1.37, 2.93) for each 10 µg NO₂/m³ increase. Study shows that long-term exposure to NO₂ may increase the incidence of depression.

Aleman et al. [61] analyzed on the group of children from the BREATHE project (N=1 667 at the age 7-11 years), if there is any association between traffic-related air pollution and ε4 allele of Apolipoprotein E gene, which is understood as genetic risk factor for Alzheimer's disease. NO₂ exposure at home address was 54.25±18.40 µg/m³, at

schools $47.74 \pm 12.95 \mu\text{g}/\text{m}^3$. NO₂ exposure increased behavioral problems scores (characterized by SDQ) in $\epsilon\epsilon$ 4 carriers (N=366) vs. non-carriers (N=1 223) 1.14 (95% CI: 1.04, 1.26) vs. 1.02 (95%CI: 0.95, 1.10, $p = 0.04$) and was associated with smaller caudate volume in $\epsilon\epsilon$ 4 carriers (N=37) vs. non-carriers (N=126) - 737.9 (95% CI: - 1 201.3, - 274.5) vs. - 157.6 (95%CI: -388.8, 73.6, $p=0.03$). Annual average NO₂ concentrations in children's schools were associated with smaller caudate volume and higher behavior problem scores among APOE $\epsilon\epsilon$ 4 allele carriers. It is possible that $\epsilon\epsilon$ 4 carriers are more vulnerable to neuroinflammatory and oxidative stress induced by air pollution exposure.

Carey et al. [62] investigated the incidence of dementia to residential level of NO₂ in London. Among 130 978 adults aged 50-79 years was in the period 2005-2013 2181 subjects diagnosed with dementia (39% Alzheimer's disease, 29% vascular dementia). The average annual concentrations of NO₂ was $37.1 \pm 5.7 \mu\text{g}/\text{m}^3$. Higher risk of Alzheimer's disease was observed in subjects exposed to highest concentrations of NO₂ ($> 41.5 \mu\text{g}/\text{m}^3$) vs. subjects with lowest concentrations of NO₂ ($< 31.9 \mu\text{g}/\text{m}^3$) (HR=1.40, 95%CI 1.12 - 1.74). These associations were more consistent for Alzheimer's disease than vascular dementia. Study found evidence of a positive association between residential level of NO₂ across London and being diagnosed with dementia.

Roberts et al. [63] explored the effect of NO₂ exposure to mental health problems in children in London, U.K. (N=284). Symptoms of anxiety, depression, conduct disorder, and ADHD were assessed at ages 12 and 18. NO₂ concentration in the year 2007 was $37.9 \pm 5.5 \mu\text{g}/\text{m}^3$ (IQR 34.1-41.7). They did not observe any association between NO₂ exposure in childhood and mental health problems at age 12. But they detected association between NO₂ exposure and subsequent development of symptoms and clinically diagnosable depression and conduct disorders at age 18. They demonstrated that NO₂ exposure at age 12 years was significantly associated with major depressive disorder at age 18.

Prenatal exposure to NO₂ and sex dependent infant cognitive and motor development was analyzed by Lertxundi et al. [64] in children at 4-6 years of age, in four regions in Spain (N= 1 119). Infant neuropsychological development was assessed by McCarthy scales: Verbal, Perceptive-Manipulative, Numeric, General Cognitive, Memory and Motor. NO₂ exposure during pregnancy was from $18.7 \pm 6.1 \mu\text{g}/\text{m}^3$ to $41.8 \pm 10.7 \mu\text{g}/\text{m}^3$. The majority of cognitive domains were negative for NO₂, associations were more negative for boys, statistically significant for memory, global cognition and verbal. These findings indicate a greater vulnerability of boys in domains related to memory, verbal and general cognition.

Jorcano et al. [65] assessed association between NO₂ and depressive and anxiety symptoms, and aggressive symptoms in children of 7-11 years, related to their prenatal and postnatal exposure. Data were analyzed in 13 182 children from 8 European population-based cohorts. Prenatal NO₂ levels ranged from $15.9 \mu\text{g}/\text{m}^3$ to $43.5 \mu\text{g}/\text{m}^3$, postnatal levels ranged from $14.0 \mu\text{g}/\text{m}^3$ to $43.5 \mu\text{g}/\text{m}^3$. 1 108 (8.4%) and 870 (6.6%) children were classified as having depressive and anxiety symptoms, and with aggressive symptoms. Obtained results suggest that prenatal and postnatal exposure to NO₂ is not associated with depressive and anxiety symptoms or aggressive symptoms in children of 7 to 11 years old.

Loftus et al. [66] used mother-child cohort from CANDLE study, analyzed the impact of prenatal NO₂ exposure ($22.3 \pm 7.1 \mu\text{g}/\text{m}^3$) and postnatal exposure ($16.2 \pm 4.7 \mu\text{g}/\text{m}^3$) on childhood behavior (N=975). In sample was 64% African American, 53% had a household annual income below \$ 35 000, child age was 4.3 years. Mothers completed the Child Behavior Checklist, measure of problem behaviors in the past two weeks. $4 \mu\text{g}/\text{m}^3$ higher prenatal NO₂ was positively associated with externalizing behavior (6%, 95% CI: 1, 11%), the effect of postnatal exposure was stronger (8%, 95% CI: 0, 16%). Prenatal NO₂ exposure was also associated with significant internalizing and externalizing behaviors. NO₂ exposure is positively associated with child behavior problems, African American and low SES children may be more susceptible.

Kulick et al. [67] examined in 5 330 participants from the northern Manhattan area of New York City effect of long-term exposure to NO₂ (annual estimates $57.4 \pm 22.1 \mu\text{g}/\text{m}^3$)

and PM_{2.5} (annual estimates 13.1±4.8 µg/m³), predominantly in women, median age 75.2 (±6.46) years. A + IQR increase of residential NO₂ was predictive of a 22.SD (95% CI, -0.30, -0.14) lower global cognitive score at baseline and more rapid decline (-0.06 SD; 95% CI -0.08, -0.04) in global cognitive function between biennial visits.

Erikson et al. [68] studied the association between NO₂ exposure and total gray matter and total white matter volumes in adults, using sample from UK Biobank. Participants were recruited from 2006 to 2010, Subset with magnetic-resonance brain imaging (MRI) included 18 292 participants, average age of 62 (44-80), NO₂ levels were 25.61±6.86 µg/m³. The mean total gray-matter volume was 708 111 mm³ (±47 940), the mean total white-matter volume was 708 111 mm³ (±40 696). The total gray-matter volume was inversely associated with NO₂ (b= -103, p < 0.01). The effect of NO₂ on gray-matter volume was more pronounced in females (b= 161, p < 0.05). Obtained findings suggest that NO₂ concentrations lower than EU standard could be associated with reduced total gray-matter.

All reviewed studies indicate a significant health risk of NO₂ exposure:

Concentrations of NO₂ lower than EU standard 40 µg/m³:

- Prenatal exposure impaired attentional function at the age 4-5 years
- Induce neurobehavioral changes in children at the age of 8-10 years
- Affect attention process in children aged 8-12 years, induced changes are persistent for another 3.5 years
- Increase major depressive disorder at age 18
- Increase the incidence of dementia
- Exposure to NO₂ is associated with reduced total gray-matter

All mentioned studies indicate, that EU standard for NO₂ should be decreased to 30 µg/m³/year.

6. Discussion

A detailed analysis of NO₂ concentrations measured by the passive samplers shows a clear correlation of NO₂ concentrations with daily traffic counts, adjusted for additional emissions due to uphill travel and stopping at intersections. This finding is in good agreement with the data from the monitoring stations, which, by themselves, are too sparse to make such inference. The correlation of NO₂ concentrations with vehicular traffic intensity is also apparent from the comparison of the data from state air quality monitoring stations during the period of March 14-April 30, 2020, during which travel restrictions were imposed, including the prohibition of all non-cargo international travel, with comparable periods of four previous years. Overall, the findings confirm that vehicular traffic, through primary NO₂ emissions (and possibly through fast reaction of primary NO with ozone), directly affects the NO₂ concentrations in the immediate vicinity.

This correlation, along with correlation of passive sampler readings and air quality monitoring stations, and good consistency of reported NO₂ concentrations among samplers used within the same location at different time periods, all suggest that passive samplers appear to provide, at a reasonable cost and effort, a fairly good image of the distribution of NO₂ concentrations. Judging from limited data, the passive samplers were found to measure about 18.5% higher values than the monitoring stations. Repeated – and most likely deliberate – removals of passive samplers from the immediate vicinity of the monitoring stations have prevented a more quantitative comparison. A comparison of a broader set of data reveals a slightly smaller bias, contributed to, in several cases, by the passive samplers being at more exposed locations (i.e., near the exit of a large underground parking garage) than the monitoring stations. The true bias could therefore be possibly even lower.

Since the trends are comparable within and outside the heavy truck exclusion area, this seems to be primarily an effect of cars and other lighter vehicles (per city statistics, about 90% of traffic is passenger cars, [42]). This is in line with the findings that truck NO_x emissions have decreased to a considerably higher extent than those of diesel cars in Europe.

The monitoring station at Legerova street is most likely not the absolute hot-spot – it is expected that the emissions of NO_x would be higher on the parallel street where the vehicles travel uphill (Legerova is one-way street downhill) and at nearby intersections. The street V Holešovičkách, a six-lane road which is, unlike most other roads of similar size, immediately bordered by residential neighborhoods, with a traffic intensity approaching 100 thousands vehicles per day, a major increase after the opening of a new complex of tunnels providing alternative route through congested areas, further complicated by a 3% grade, could easily be the next traffic hot-spot.

Considering the finding that about half of the vehicles traveling on the road are not older than 7 years [27], and the several-fold decrease in NO_x emissions standards over the last decade and half, a much sharper decrease of NO₂ concentrations would be expected than the approximately 1% annually reported by Hůnová [5]. Given the decrease in the limit values of roughly two thirds from Euro 3 (0.50 g/km NO_x, 2000) to Euro 5 (0.18 g/km, 2009-2010) and from Euro 4 (0.25 g/km NO_x, 2005) to Euro 6 (0.08 g/km, 2014-2015), the introduction of Euro 5 in late 2009 and Euro 6 in late 2014 should have resulted in about two thirds NO_x reduction in at least half of the vehicles, or, about one third reduction in NO_x emissions in general. As learned from the analysis of the effects of traffic restrictions, the effect on NO₂ concentrations may be different, and possibly somewhat smaller than the reduction in NO_x emissions, due to atmospheric chemistry. The effects of such decrease could also have been diminished by an increase in traffic, however, in center city, the intensity of automobile traffic has been stagnating, or even slightly decreasing.

The mediocre decrease in NO₂ concentrations, despite more dramatic reduction being expected from improving vehicle technology, is in line with earlier findings that the real NO_x emissions of diesel vehicles did not decrease despite the decreasing emissions limits. The situation should have been, however, substantially remedied by „post-Dieselgate“ vehicles and by repairs of vehicles affected by Dieselgate. Since it was not, a question therefore arises as to the possibility that Dieselgate relevant repairs were not done on a sufficient number of vehicles and/or were not sufficiently effective and/or were reversed to the „original factory conditions“ by the vehicle owners. The authors do not have any reliable statistics on this matter. Furthermore, considering that all three mentioned situations could be associated with criminal offenses and/or considerable civil penalties, detailed investigation of the matter is likely to be considerably difficult.

If there is no assurance that the NO₂ concentrations will decrease dramatically due to a radical improvement in primary NO_x emissions, the only other suitable strategy to improve the air quality is to reduce, to the extent required, the intensity of vehicular traffic. Contrary to the remote regions where automobiles are, in most cases, the only practical means of travel, Prague has an extensive network of public transit. According to the City of Prague statistics [42], only 29% trips in Prague are done by automobile; 26% of trips are by walking, and 42% of trips by public transit. Of the public transit, slightly over one third is done by subway, and another third by trams and commuter rail, which are, with the exception of a rather small number of diesel rail cars used on sparsely traveled rail lines, run on electric power, and therefore with very small effect on NO₂ emissions. The remaining third of trips is by diesel buses, the majority of which are equipped with SCR catalysts, and potentially reaching NO_x emissions not much larger (and according to measurements possibly even smaller) levels, per kilometer and vehicle, than an average diesel car. It is therefore readily apparent that shifting from an average automobile to any other means of transport is likely to reduce the NO₂ concentrations. (Shift to electric power, compressed natural gas, or other “clean” propulsion is a gradual process and is unlikely to be done, within a few years, on a sufficiently large number of vehicles to make a difference throughout the city.)

Impact of NO₂ on neurobehavioral changes in children were studied especially in Spain. Lertxundi et al. [52] analyzed the effect of prenatal NO₂ exposure on children at 15 months of age, observed adverse effect on cognitive development. Sentis et al. [57] evaluated prenatal NO₂ exposure (31.1 µg/m³ (18.4-37.9) to attention at 4-5 years of children,

exposure to NO₂ during pregnancy reduced attentional function. Loftus et al. [66] analyzed the impact of prenatal NO₂ exposure ($22.3 \pm 7.1 \text{ } \mu\text{g}/\text{m}^3$) on childhood behavior in the South of USA at child age 4.3 years. Prenatal exposure increased behavioral problems.

Carey et al. [62] investigated in London in adults aged 50-79 years the risk related to dementia. The higher risk of Alzheimer disease was observed in subjects exposed to higher concentrations of NO₂. Erikson et al. [68] in UK observed the effect of NO₂ exposure ($25.6 \pm 6.9 \text{ } \mu\text{g}/\text{m}^3$) - to reduce total gray-matter volume, especially in women.

7. Conclusions

Despite massive reductions in diesel cars NO_x emission limits, of about two thirds from Euro 3 to Euro 5 and from Euro 4 to Euro 6, NO₂ concentrations throughout the Czech Republic have been decreasing at a mediocre rate of 1% annually. To elucidate the effects of motorized traffic on NO₂ concentrations, data from 104 passive NO₂ samplers deployed at 65 locations in Prague during March-April and September-October of 2019 were examined. Comparisons with the national monitoring network show a positive bias of 18.5% for co-located samplers and 17% for samplers nearby (or in similar settings as) the monitoring stations. There was a good correlation among repeated measurements at the same locations. The data from the national air quality monitoring network show that the average concentrations in both spring and fall sampling periods were consistent with 2016-2019 averages.

The average measured NO₂ concentrations at the selected locations, after correcting for the 18.5% bias, were in the range of 16 to 69 $\mu\text{g}/\text{m}^3$, with a mean of 36 $\mu\text{g}/\text{m}^3$ and a median of 35.3 $\mu\text{g}/\text{m}^3$, and were higher than the EU and national limit (annual average) of 40 $\mu\text{g}/\text{m}^3$ at 32% of locations. The NO₂ concentrations have correlated well with the intensity of traffic (average daily vehicle counts), corrected for additional emissions due to uphill travel and due to idling at, and accelerating from, intersections. Several additional "hot-spots" were identified, in addition to the "hot-spot" monitoring station at Legerova street (2016-2019 NO₂ average of 51 $\mu\text{g}/\text{m}^3$), where the vehicles travel on a slight decline on one-way street: several intersections at Sokolská street, parallel with Legerova with uphill direction of travel, and emerging hot-spots along V Holešovičkách street, where the traffic intensity increased due to the opening of a new series of tunnels. Analysis of the effect of coronavirus related travel restrictions were evaluated by comparing the data from six monitoring stations (March 15-April 30, 2020, relative to the same period during 2016-2019) reveal a reduction of NO, NO₂ and NO_x (except for a small increase of NO₂ at one of the background stations), with NO reduction being, at high traffic locations, higher than that of NO₂. The spatial analysis of data from passive samplers and time analysis of data during the travel restrictions both demonstrate a consistent positive correlation between traffic intensity and NO₂ concentrations along/near the travel path.

It appears that decreases in vehicle NO_x emission limits, introduced in the last decade or two, have failed to sufficiently reduce the ambient NO₂ concentrations in exposed locations in Prague. This is in part due to increased fraction of NO₂ in NO_x in newer vehicles, and in part due to "a major disparity between the numerical value of the emission limit and the actual emissions in everyday driving". Further, there is no apparent sign of, and it is far from clear that, the "excess emissions" of NO_x, a problem known as Dieselgate, have been efficiently remedied.

All reviewed studies indicate a significant effect of prenatal NO₂ exposure to children's neurobehavioral development, in adults to dementia at concentrations lower than EU standards of 40 $\mu\text{g}/\text{m}^3/\text{year}$. These results should be understood as an imperative to decrease the EU standard to 30 $\mu\text{g}/\text{m}^3/\text{year}$. All presented studies prove that NO₂ can significantly deteriorate CNS and therefore this knowledge should be used to improve the quality of our lives.

Acknowledgements: The passive sampler measurements were organized and performed by Dr. Miroslav Šuta from the Center for Environment and Health, Thámova 1275/21, 301 00 Plzeň, Czech Republic. The acquisition and analysis of the passive samplers was funded by Deutsche Umwelthilfe (Environmental Action Germany), Hackescher Markt 4, 10178 Berlin, Germany, www.duh.de. The data from the national air quality monitoring network was provided by the Czech Hydrometeorological Institute. The authors thank Václav Novák, the Head of Air Quality Information System Department, for providing the data and for helpful advice. Evaluation of the passive samplers and some of the background research on emissions was done (M.V.) within the H2020 project no. 851002 uCARE, you can also reduce emissions. Review of the health effects and remaining work has been supported by the European Regional Development Fund under Grant Healthy Aging in Industrial Environment HAIE (CZ.02.1.01/0.0/0.0/16_019/0000798).

References

1. Health Effect Institute, State of Global Air. **2018**, Report, Available online: <https://www.stateofglobalair.org/sites/default/files/soga-2018-report.pdf> (accessed on 8th February 2021).
2. EEA (European Environment Agency), Air Quality in Europe – 2020 report. doi: 10.2800/786656. Available online: <https://www.eea.europa.eu/publications/air-quality-in-europe-2020-report> (accessed on 8th February 2021).
3. World Bank, The Cost of Pollution: Strengthening the Economic Case for Action. **2016**, Washington, DC, Report, Available online: <https://openknowledge.worldbank.org/bitstream/handle/10986/25013/108141.pdf?sequence=4&isAllowed=y> (accessed on 8th February 2021).
4. Hesterberg, T.W., Long, C.M., Sax, S.N., Lapin, C.A., McClellan R.O., Bunn, W.B., Valberg, P.A. (2011). Particulate matter in new technology diesel exhaust (NTDE) is quantitatively and qualitatively very different from that found in traditional diesel exhaust (TDE). *J. Air Waste Manag. Assoc.* **2011**, 61, 894-913, doi:10.1080/10473289.2011.599277.
5. Hunova, I., Baumelt, V., Modlik, M. Long-term trends in nitrogen oxides at different types of monitoring stations in the Czech Republic. *Sci. Total Environ.* **2020**, 699, 134378, doi: 10.1016/j.scitotenv.2019.134378.
6. Georgoulas, A.K., Ronald. van der A, J., Stammes, P., Boersma, K. F., Eskes, H.J. Trends and trend reversal detection in 2 decades of tropospheric NO₂ satellite observations. *Atmospheric Chem. Phys.* **2019**, 19, 6269–6294, doi: 10.5194/acp-19-6269-2019.
7. Casquero-Vera, J.A., Lyamani, H., Titos, G., Borrás E., Olmo, F.J., Alados-Arboledas L. Impact of primary NO₂ emissions at different urban sites exceeding the European NO₂ standard limit. *Sci. Total Environ.* **2019**, 646, 1117-1125, doi: 10.1016/j.scitotenv.2018.07.360.
8. Wild, R.J., Dubé, W.P., Aikin, K.C., Eilerman, S.J., Neuman, J.A., Peischl, J., Ryerson, T.B., Brown, S.S. On-road measurements of vehicle NO₂/NO_x emission ratios in Denver, Colorado, USA. *Atmos. Environ.* **2017**, 148, 182-189, doi: 10.1016/j.atmosenv.2016.10.039.
9. Grange, S.K., Lewis, A.C., Moller, S.J., Carslaw, D.C. Lower vehicular primary emissions of NO₂ in Europe than assumed in policy projections. *Nat. Geosci.* **2017**, 10, 914–918, doi: 10.1038/s41561-017-0009-0.
10. Zeldovich, Y.B. The Oxidation of Nitrogen in Combustion Explosions, *Acta Physicochimica* **1946**, 21, 577–628.
11. Lavoie G.A., Heywood J.B., Keck J.C. Experimental and Theoretical Study of Nitric Oxide Formation in Internal Combustion Engines, *Combust. Sci. Technol.* **1970**, 1:4, 313-326, doi: 10.1080/00102206908952211.
12. Gutzwiller L., Arens F., Baltensperger, U., Gaggeler H.W., Ammann M. Significance of Semivolatile Diesel Exhaust Organics for Secondary HONO Formation. *Environ. Sci. Technol.* **2002**, 36, 677-682, doi: 10.1021/es015673b.
13. Kurtenbach, R., Becker, K.H., Gomes, J.A.G., Kleffmann, J., Lorzer, J.C., Spittler, M., Wiesen, P., Ackermann, R., Geyer, A., Platt U. Investigations of emissions and heterogeneous formation of HONO in a road traffic tunnel. *Atmos. Environ.* **2001**, 35, 3385-3394, doi: 10.1016/S1352-2310(01)00138-8.

14. Heeb, N.V., Zimmerli, Y., Czerwinski, J., Schmid, P., Zennegg, M., Haag, R., Seiler, C., Wichser, A., Ulrich, A., Honegger, P., Zeyer, K., Emmenegger, L., Mosimann, T., Kasper, M., Mayer, A. Reactive nitrogen compounds (RNCs) in exhaust of advanced PM-NO_x abatement technologies for future diesel applications. *Atmos. Environ.* **2011**, *45*, 3203-3209, doi: 10.1016/j.atmosenv.2011.02.013.
15. Seinfeld J.H., Pandis S.N. *Atmospheric chemistry and physics: from air pollution to climate change*. John Wiley & Sons: Hoboken, New Jersey, USA, 1998.
16. Mooney, J.J., Thompson, C.E., Dettling, J.C. Three-Way Conversion Catalysts Part of the New Emission Control System. *SAE Transactions* **1977**, *86*, 1553-1562.
17. Falk, C.D., Mooney, J. J. Three-Way Conversion Catalysts: Effect of Closed-Loop Feed-Back Control and Other Parameters on Catalyst Efficiency. *SAE Transactions* **1980**, *89*, 1822-1832.
18. Hardin, G. (1968). The tragedy of the commons. *Science* **162**, 3859, 1243-1248, doi: 10.1126/science.162.3859.1243.
19. Thompson, G., Carder, D., Clark, N., and Gautam, M., "Summary of In-use NO_x Emissions from Heavy-Duty Diesel Engines," *SAE Int. J. Commer. Veh.* **2009**, 1(1), 162-184, <https://doi.org/10.4271/2008-01-1298>.
20. USDOJ, 2015. United States Department of Justice: Clean Air Act diesel engine cases. Available online: <https://www.justice.gov/enrd/diesel-engines> (accessed on 2nd February 2021).
21. US CFR, 2000: United States Code of Federal Regulations, Volume 40, Part § 86.1370: Not-To-Exceed test procedures. United States Federal Register, Vol. 65, No. 195, October 6, 2000, as amended by subsequent regulations.
22. Gieshaskiel, B., Gloria R., Carriero, M., Lahde, T., Forloni, F., Perujo Mateos del Parque, A., Martini, G., Bissi, L.M., Terenghi, R. Emission Factors of a Euro VI Heavy-duty Diesel Refuse Collection Vehicle. *Sustainability* **2019**, *10*, 1067, doi: 10.3390/su11041067.
23. Suarez-Bertoa, R., Valverde, V., Clairotte, M., Pavlovic, J., Giechaskiel, B., Franco, V., Kregar, Z., Astorga-Llorens, M. On-road emissions of passenger cars beyond the boundary conditions of the real-driving emissions test. *Environ. Res.* **2019**, *176*, 108572. doi 10.1016/j.envres.2019.108572.
24. Quiros, D.C., Thiruvengadam, A., Pradhan, S., Besch, M., Thiruvengadam, P., Demirgok, B., Carder, D., Oshinuga, A., Huai, T., Hu, S. Real-world emissions from modern heavy-duty diesel, natural gas, and hybrid diesel trucks operating along major California freight corridors. *Emiss. Control. Sci. Technol.* **2016**, *2*(3), 156-172, doi: 10.1007/s40825-016-0044-0.
25. Jiang, Y., Yang, J., Cocker 3rd, D., Karavalakis, G., Johnson, K.C., Durbin, T.D Characterizing emission rates of regulated pollutants from model year 2012 + heavy-duty diesel vehicles equipped with DPF and SCR systems. *Sci. Total Environ.* **2018**, *619–620*, 765-771, doi: 10.1016/j.scitotenv.2017.11.120.
26. Grigoratos, T., Fontaras, G., Giechaskiel, B., Zacharof, N. Real world emissions performance of heavy-duty Euro VI diesel vehicles. *Atmos. Environ.* **2019**, *201*, 348-359, doi: 10.1016/j.atmosenv.2018.12.042.
27. Skacel, J., Vojtisek, M., Beranek, V., Pechout, M.: Black Sheep - Detecting polluting vehicles on the road using roadside particle measurement. Proceedings of ETH Conference on Combustion Generated Nanoparticles, Zurich, Switzerland, June 2018, Available online: https://nanoparticles.ch/archive/2018_Skacel_PO.pdf (accessed on 2nd February 2021).
28. Vojtisek-Lom, M., Fenkl, M., Dufek, M., and Mareš, J. Off-cycle, Real-World Emissions of Modern Light Duty Diesel Vehicles. *SAE Tech. Pap.* **2009**, 2009-24-0148, doi: 10.4271/2009-24-0148.
29. Weiss M., Bonnel P., Kuhlwein J., Provenza, A., Lambrecht, U., Alessandrini, S., Carriero, M., Colombo, R., Forni, F., Lanappe, G., Lijour, P., Manfredi, U., Montigny, F., Sculati, M. Will Euro 6 reduce the NO_x emissions of new diesel cars? - Insights from on-road tests with Portable Emissions Measurement Systems (PEMS). *Atmos. Environ.* **2012**, *62*, 657–665, doi: 10.1016/j.atmosenv.2012.08.056.
30. Ligterink, N., Kadijk, G., Mensch, P. van, Hausberger, S., Rexeis, M., Investigations and real world emission performance of Euro 6 light-duty vehicles. **2013**, report TNO 2013 R11891, 53.

31. Franco, V., Sanchez, F.P., German, J., Mock, P., Real-word exhaust emissions from modern diesel cars a meta-analysis of PEMS emissions data from EU (EURO 6) and US (TIER 2 BIN 5/ULEV II) diesel passenger cars. **2014**, White paper, ICCT (International Council Clean on Transportation).
32. Yang, L., Franco, V., Mock, P., Kolke, R., Zhang, S., Wu, Y., German, J. Experimental Assessment of NO_x Emissions from 73 Euro 6 Diesel Passenger Cars. *Environ. Sci. Technol.* **2015**, *49*, 14409–14415, doi:10.1021/acs.est.5b04242.
33. Olsen, D.B., Kohls, M., Arney, G. Impact of oxidation catalysts on exhaust NO₂/NO_x ratio from lean-burn natural gas engines. *J. Air Waste Manag. Assoc.* **2010**, *60*, 867-874, doi:10.3155/1047-3289.60.7.867.
34. Carslaw, D.C. Evidence of an increasing NO₂/NO_x emissions ratio from road traffic emissions. *Atmos. Environ.* **2005**, *39*, 4793-4802, doi:10.1016/j.atmosenv.2005.06.023.
35. Carslaw, D., Rhys-Tyler, G. Remote sensing of NO₂ exhaust emissions from road vehicles: A report to the City of London Corporation and London Borough of Ealing. **2013**, DEFRA Project.
36. Preble, C.V., Harley, R.A., Kirchstetter, T.W., "Measuring real-world emissions from the on-road heavy-duty truck fleet. **2019**, University of California, Berkeley, USA.
37. Vojtisek-Lom, M., Beranek, V., Klir, V., Jindra, P., Pechout, M., Vorisek, T. On-road and laboratory emissions of NO, NO₂, NH₃, N₂O and CH₄ from late-model EU light utility vehicles: Comparison of diesel and CNG. *Sci. Total Environ.* **2018**, *616-617*, 774-784, doi: 10.1016/j.scitotenv.2017.10.248.
38. Pechout, M., Kotek, M., Jindra, P., Macoun, D., Hart, J., Vojtisek-Lom, M. Comparison of hydrogenated vegetable oil and biodiesel effects on combustion, unregulated and regulated gaseous pollutants and DPF regeneration procedure in a Euro6 car. *Sci. Total Environ.* **2019**, *696*, 133748, doi: 10.1016/j.scitotenv.2019.133748.
39. DUH (Deutsche Umwelthilfe e.V.), October 2019: NO₂ Report Hotspots in Germany, Czech Republic, Slovenia, Bulgaria and Serbia, https://www.duh.de/fileadmin/user_upload/download/Projektinformation/Verkehr/Abgasalarm/NO2_Report_17_10_19.pdf
40. Palmes, E.D., Gunnison, A.F., DiMattio, J., Tomczyk, C. Personal Sampler for Nitrogen Dioxide. *Am. Ind. Hyg. Assoc. J.* **1976**, *37*, 570-577, doi: 10.1080/0002889768507522.
41. Passam, NO₂ passive sampler data sheet and specifications., Switzerland. https://www.passam.ch/wp-content/uploads/2020/01/en_NO2lt.pdf (accessed on 2nd February 2021).
42. Technická správa komunikací hl. m. Prahy (TSK; City of Prague Highway Department), Prague Transportation Yearbook 2019. Online at <http://www.tsk-praha.cz/static/udi-rocenka-2019-en.pdf> (accessed on 7th February 2021).
43. Cape, J.N., Review of the use of passive diffusion tubes for measuring concentrations of nitrogen dioxide in air. Defra **2005**.
44. Buzica, D., Gerboles, M., Plaisance, H. The equivalence of diffusive samplers to reference methods for monitoring O₃, benzene and NO₂ in ambient air. *J. Environ. Monit.* **2008** *10*, 1052-1059, doi: 10.1039/b802260g.
45. Altshuller, A.P. Thermodynamic considerations in the interactions of nitrogen oxides and oxy-acids in the atmosphere. *J. Air Pollut. Control. Assoc.* **1956**, *6*, 97–100, doi: 10.1080/00966665.1956.10467740.
46. Yang, B., Zhang, K.M., Xu, W.D., Zhang, S., Batterman, S., Baldauf, R.W., Deshmukh, P., Snow, R., Wu, Y., Zhang, Q., Li, Z., Wu, X. On-Road Chemical Transformation as an Important Mechanism of NO₂ Formation. *Environ. Sci. Technol.* **2018**, *58*, 4574–4582.
47. Richmond-Bryant, J., Owen, R.C., Graham, S., Snyder, M., McDow, S., Oakes, M., Kimbrough, S. Estimation of on-road NO₂ concentrations, NO₂/NO_x ratios, and related roadway gradients from near-road monitoring data. *Air Qual. Atmos. Health* **2017**, *10*, 611-625, doi: 10.1007/s11869-016-0455-7.
48. Singh, J. Nitrogen dioxide exposure alters neonatal development. *Neurotoxicology* **1988**, *9*, 545-549.

49. Wang, S.Q., Zhang, J.L., Zeng, X.D., Zeng, Y.M., Wang, S.C., Chen, S.Y. Association of traffic-related air pollution with children's neurobehavioral functions in Quanzhou, China. *Environ. Health Perspect.* **2009**, *117*(10), 1612-1618, doi: 10.1289/ehp.0800023.
50. Guxens, M., Aguilera, I., Ballester, F., Estarlich, M., Fernandez-Somoano, A., Lertxundi, A., Lertxundi, N., Mendez, M.A., Tardon, A., Vrijheid, M., Sunyer, J. Prenatal exposure to residential air pollution and infant mental development: Modulation by antioxidants and detoxification factors. *Environ. Health Perspect.* **2012**, *120*, 144-149, doi: 10.1289/ehp.1103469.
51. Kim, E., Park, H., Hong, Y-C., Ha, M., Kim, Y., Kim, B.N., Kim, Y., Roh, Y.M., Lee, B.E., Ryu, J.M., Kim, B.M., Ha, E.H. Prenatal exposure to PM10 and NO2 and children's neurodevelopment from birth to 24 months of age: Mothers and Children Environmental Health (MOCEH) study. *Sci. Total Environ.* **2014**, *481*, 439-445, doi: 10.1016/j.scitotenv.2014.01.107.
52. Lertxundi, A., Baccini, M., Letxundi, N., Fano, E., Aranbarri, A., Martinez, M.D., Ayerdi, M., Alvarez, J., Santa-Marina, L., Dorronsoro, M., Ibarluzea, J. Exposure to fine particle matter, nitrogen dioxide and benzene during pregnancy and cognitive and psychomotor developments in children at 15 months of age. *Environ. Int.* **2015**, *80*, 33-40, doi: 10.1016/j.envint.2015.03.007.
53. Sunyer, J., Esnaola, M., Alvarez-Pedrerol, M., Forns, J., Rivas, I., Lopez-Vicente, M., Suades-Gonzalez, E., Foraster, M., Garcia-Esteban, R., Basagana, X., Viana, M., Cirach, M., Moreno, T., Alastuey, A., Sebastian-Galles, N., Nieuwenhuijsen, M., Querol, X. Association between traffic-related air pollution in schools and cognitive development in primary school children: A prospective cohort study. *PloS Med.* **2015**, *12*, e 1001792, doi:10.1371/journal.pmed.1001792.
54. Pujol, J., Martinez-Vilavella, G., Macia, D., Fenoll, R., Alvarez-Pedrerol, M., Rivas, I., Forns, J., Blanco-Hinojo, L., Capellades, J., Querol, X., Deus, J., Sunyer, J. Traffic pollution exposure is associated with altered brain connectivity in school children. *Neuroimage* **2016**, *129*, 175-184, doi: 10.1016/j.neuroimage.2016.01.036.
55. Forns, J., Dadvand, P., Foraster, M., Alvarez-Pedrerol, M., Rivas, I., Lopez-Vicente, M., Suades-Gonzalez, E., Garcia-Esteban, R., Esnaola, M., Cirach, M. Traffic-related air pollution, noise at school, and behavioral problems in Barcelona schoolchildren: A cross-sectional study. *Environ. Health Perspect.* **2016**, *124*, 529-535, doi: 10.1289/ehp.1409449.
56. Min, J., Min, K. Exposure to ambient PM10 and NO2 and the incidence of attention-deficit hyperactivity disorder in childhood. *Environ. Int.* **2017**, *99*, 221-227, doi: 10.1016/j.envint.2016.11.022.
57. Sentis, A., Sunyer, J., Dalmau-Bueno, A., Andiaarena, A., Ballester, F., Ciracha, M., Estarlich, M., Fernandez-Somoano, A., Ibarluzea, J., Iniguez, C., Lertxundi, A., Tardon, A., Nieuwenhuijsen, M., Vrijheid, M., Guxens, M.; INMA Project. Prenatal and postnatal exposure to NO2 and child attentional function at 4-5 years of age. *Environ. Int.* **2017**, *106*, 170-177, doi: 10.1016/j.envint.2017.05.021.
58. Sunyer, J., Suades-Gonzales, E., Garcia-Esteban, R., Rivas, I., Pujol, J., Alvarez-Pedrerol, M., Forns, J., Querol, X., Basagana, X. Traffic-related air pollution and attention in primary school children. Short-term association. *Epidemiology* **2017**, *28*, 181-189, doi: 10.1097/EDE.0000000000000603.
59. Forns, J., Dadvand, P., Esnaola, M., Alvarez-Pedrerol, M., Lopez-Vicente, M., Garcia-Esteban, R., Cirach, M., Basagana, X., Guxens, M., Sunyer, J. Longitudinal association between air pollution exposure at school and cognitive development in school children over a period of 3.5 years. *Environ. Res.* **2017**, *159*, 416-421, doi: 10.1016/j.envres.2017.08.031.
60. Vert, C., Sanchez-Benavides, G., Martinez, D., Gotsens, X., Gramunt, N., Cirach, M., Molinuevo, J.L., Sunyer, J., Nieuwenhuijsen, M.J., Crous-Bou, M., Gascon, M. Effect of long-term exposure to air pollution on anxiety and depression in adults: A cross-sectional study. *Int. J. Hyg. Environ. Health* **2017**, *220*, 1074-1080, doi: 10.1016/j.ijheh.2017.06.009.
61. Alemany, S., Vilor-Tejedor, N., Garcia-Esteban, R., Bustamante, M., Dadvand, P., Esnaola, M., Mortamais, M., Forns, J., van Drooge, B.L., Alvarez-Pedrerol, M., Grimalt, J.O., Rivas, I., Querol, X., Pujol, J., Sunyer, J. Traffic-related air pollution, APOE ϵ 4 status, and neurodevelopmental outcomes among school children enrolled in the BREATHE project (Catalonia, Spain). *Environ. Health Perspect.* **2018**, *126*(8), 087001, doi: 10.1289/EHP2246.

-
62. Carey, I.M., Anderson, H.R., Atkinson R.W., Beevers, S., Cook, D.G., Strachan, D.P., Dajnak, D., Gulliver, J., Kelly, F.J. Are noise and air pollution related to the incidence of dementia? A cohort study in London, England. *BMJ Open* **2018**, 8, e022404, doi:10.1136/bmjopen-2018-022404.
 63. Roberts, S., Arseneault, L., Barratt, B., Danese, A., Odgers, C.L., Moffitt, T.E., Reuben, A., Kelly, F.J., Fisher, H.L. Exploration of NO₂ and PM_{2.5} air pollution and mental health problems using high-resolution data in London-based children from a UK longitudinal cohort study. *Psychiatry Res.* **2019**, 272, 8-17, doi: 10.1016/j.psychres.2018.12.050.
 64. Lertxundi, A., Andiaarena, A., Martinez, M.D., Ayerdi, M., Murcia, M., Estarlich, M., Guxens, M., Sunyer, J., Julvez, J., Ibarluzea, J. Prenatal exposure to PM_{2.5} and NO₂ and sex-dependent infant cognitive and motor development. *Environ. Res.* **2019**, 174, 114-121, doi: 10.1016/j.envres.2019.04.001.
 65. Jorcano, A., Lubczynska, M.J., Pierotti, L., Altung, H., Ballester, F., Cesaroni, G., El Marroun, H., Fernandez-Somoano, A., Freire, C., Hanke, W., Hoek, G., Ibarluzea, J., Iñiguez, C., Jansen, P.W., Lepeule, J., Markevych, I., Polańska, K., Porta, D., Schikowski, T., Slama, R., Standl, M., Tardon, A., Vrijkotte, T.G.M., von Berg, A., Tiemeier, H., Sunyer, J., Guxens, M. Prenatal and postnatal exposure to air pollution and emotional and aggressive symptoms in children from 8 European birth cohorts. *Environ. Int.* **2017**, 131, 104927, doi: 10.1016/j.envint.2019.104927.
 66. Loftus, C.T., Ni, Y., Szpiro, A.A., Hazlehurst, M.F., Tylavsky, F.A., Bush, N.R., Sathyanarayana, S., Carroll, K.N., Young, M., Karr, C.J., LeWinn, K.Z. Exposure to ambient air pollution and early childhood behavior: A longitudinal cohort study. *Environ. Res.* **2020**, 183, 109075, doi.org/10.1016/j.envres.2019.109075.
 67. Kulick, E.R., Wellenius, G. A., Boehma, A. K., Joyce, N. R., Schupf, N., Kaufman, J. D., Mayeux, R., Sacco, R. L., Manly, J. J., Elkind, M., S., V. Long-term exposure to air pollution and trajectories of cognitive decline among older adults. *Neurology* **2020**, 94: e1782-e1792. doi: 10.1212/WNL.0000000000009314.
 68. Erickson, L.D., Gale, S.D., Anderson, J.E., Brown, B.L., Hedges, D.W. Association between exposure to air pollution and total gray matter and total white matter volumes in adults: A cross-sectional study. *Brain Sci.* **2020**, 10 (3), 164, doi: 10.3390/brainsci10030164.