

Article

Savouring as an Intervention to Decrease Negative Affect in Anxious Mothers of Children with Autism and Neurotypical Children

Ariel S. Pereira¹, Atiqah Azhari¹ , Chloe A. Hong¹, Gerin E. Gaskin³, Jessica L. Borelli⁴  and Gianluca Esposito^{1,4,5*} ¹ Psychology Program, School of Social Sciences, Nanyang Technological University, Singapore, Singapore² Division of Behavioral and Organizational Sciences, Claremont Graduate University, Claremont, California, USA³ Department of Psychological Science, University of California, Irvine, California, USA⁴ Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore⁵ Department of Psychology and Cognitive Science, University of Trento, Rovereto, Italy

* Correspondence: gianluca.esposito@ntu.edu.sg; gianluca.esposito@unitn.it

† These authors contributed equally to this work.

Abstract: Savouring is an emotion regulation strategy and intervention that focuses on the process of attending, intensifying and prolonging positive experiences and positive affect associated with these memories. Personal savouring involves a reflection on positive memories that are specific to the individual and do not involve others. In contrast, relational savouring entails reflecting on instances when people were responsive to the needs of their significant others. Such interventions hold potential to benefit parents of children with autism spectrum disorder (ASD). Mothers of children with ASD tend to experience higher anxiety, lower positive affect (PA) and more negative affect (NA) compared to mothers of neurotypical children, leading to a diminished overall well-being. Moreover, ASD has multiplied by up to four times in the recent decade. Thus, this paper investigates whether savouring may enhance the overall well-being of mothers of young children with ASD by increasing positive affect and decreasing negative affect. 52 mothers of neurotypical children and 26 mothers of children with ASD aged 3-7 years old were given a series of questionnaires and randomly assigned to either relational savouring or personal savouring conditions. In relational savouring, mothers were asked to reflect upon a shared positive experience with their child while in the personal savouring condition, a personal positive experience was recalled. Across mothers of children with ASD and neurotypical children, findings suggest that savouring leads to a decrease in NA ($p < .01$) but not increases in PA. Similarly, mothers with higher levels of anxiety experience a greater decrease in NA ($p < .001$) compared to mothers with lower levels of anxiety post-savouring. This study proposes that a brief savouring intervention may be effective among mothers of preschoolers. As lower levels of negative affect is linked to healthier psychological well-being, mothers might be able to engage in more effective and warm parenting after savouring exercises, which would cultivate positive mother-child relationships that benefit their children in the long-term.

Keywords: savouring; affect; anxiety; mothers; children with autism spectrum disorder

Citation: Ariel S. Pereira, Atiqah Azhari, Chloe A. Hong, Gerin E. Gaskin, Jessica L. Borelli and Gianluca Esposito Savouring as an Intervention to Decrease Negative Affect in Anxious Mothers of Children with Autism and Neurotypical Children. *Preprints* 2021, 1, 0. <https://doi.org/>

Received:
Accepted:
Published:

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

1. Introduction

In contemporary life, it is easy to let moments pass us by. Yet, the process of reliving and recollecting positive moments can reap benefits. Savouring, a psychological exercise first conceptualized within the field of positive psychology, is the process of attending, intensifying and prolonging positive experiences [1]. Savouring has been associated with an array of positive psychological outcomes, such as lower levels of depression, greater life satisfaction, and positive emotion [2]. People can savor experiences that have occurred in the past, experiences that are occurring in the moment, or experiences that are anticipated to occur in the future [1]. Personal savouring involves reflecting on personal positive memories. In contrast, relational savouring encompasses reflecting on moments of positive connectedness or sensitive responsiveness [3]. Past savouring interventions have found savouring to be efficacious among various populations including mothers, military spouses, and couples in long-distance relationships [4–6]. In recent years, there has been greater emphasis placed on positive aspects rather than tackling the problems faced in parenting a child with autism spectrum disorder (ASD) [7]. ASD is classified as a neurodevelopmental disability, and characteristics of the disorder may include social deficits in

communication, accompanied by repetitive behaviours and restricted interests. With communication posing a significant challenge, such deficits are more easily identifiable when children reach the age of social interaction and display marked deviation from the milestones of neurotypical peers [8], around the age of two [9]. Existing on a spectrum, persons with ASD can exhibit widely differing symptoms and require support ranging from mild to severe, with a striking male dominance [10]. This diagnosis has multiplied by up to four times in the recent decade [11,12], which is of great concern. Studies generally find that compared to mothers of neurotypical children, mothers of children with ASD experience significantly less positive affect (PA) and more negative affect (NA) [13,14]. Affect, an emotional response, can be broadly classified as PA or NA, and can be measured using the Positive Affect and Negative Affect Schedule (PANAS) scale [15]. The PANAS is an established self-report measure of PA and NA [15], designed to measure affect in various contexts such as at the present moment, the past day, week, or year. Furthermore, it is based on a two-dimensional conceptual model of mood [16], which proposes that PA and NA are seen as unique and independent constructs that are capable of being experienced simultaneously [17]. Mothers of children with ASD report strong NA when interacting with their children, which clouds their perceptions and behaviour [18]. Several studies emphasise that mothers of children with developmental disabilities are more likely to experience NA compared to counterparts with neurotypical children [19], and these NA emerge and simmer in the child's early years from the time a child is diagnosed, and onwards especially when they fail to hit their developmental milestones [20]. Brief savouring interventions have successfully improved PA and decreased NA [21,22]. Positive affect can be highly adaptive (review in [23]), serving as a psychological buffer by enabling mothers of children with ASD to cope during difficult situations [24,25]. Thus far, only one study has examined the effects of savoring on the PA and NA of mothers of neurotypical infants and toddlers, but not in mothers of children with ASD. In a seminal study on 435 parents of neurotypical children, [6] assessed a subsample of 64 parents and found that both personal and relational savoring conditions elicited heightened PA and reduced NA. Correspondingly, savouring interventions may be promising in their ability to improve well-being among parents. Since mothers are generally the primary caregivers of children with autism and face an increased risk of adverse psychological outcomes such as heightened anxiety [26–28], our current study therefore intends to build and extend on the findings of [6] to mothers of neurotypical children and mothers of children with ASD. The psychological profile of mothers of children with ASD greatly differs from that of mothers of typical children, with the former experiencing significantly higher levels of anxiety symptoms (e.g. [26,29–31]). Generalized Anxiety Disorder (GAD) is characterised by excessive anxiety and worry about various events or activities. Over half of parents of children with ASD experience severe generalized anxiety [32] which may stem from worries about confronting their child's problematic and inappropriate behaviours [30,33], managing shortfalls in time and monetary resources [26,28] and overcoming the emotional turmoil they experience such as guilt and anger on a daily basis [27]. As PA is found to be lacking among people with anxiety disorders [34], highly anxious mothers of children with ASD may also stand to gain from savouring interventions. Since the study by [6] showed that personal and relational savouring both efficaciously improved mothers' affect, the two savouring conditions would be combined in this study. The broader literature on savouring likewise points to the efficacy of either types of savouring in enhancing emotional affect (e.g. [1]). Subsequently, we embarked on this study with four hypotheses. First, similar to the findings in the study by [6], we hypothesize that a decrease in Positive and Negative Affect Schedule - Negative Affect (PANAS-NA) will be seen post-savouring. Secondly, we hypothesize that savouring will increase Positive and Negative Affect Schedule - Positive Affect (PANAS-PA) among mothers of young children due to enhanced positive perceptions, similar to the success of other positive interventions in literature as found above. In the event where PANAS-NA was reduced after savouring, or PANAS-PA was enhanced post-savouring, we embark on an exploratory hypothesis to investigate whether mothers' GAD is associated with change in affect.

2. Materials and Methods

2.1. Participants

This study obtained approval from NTU Institutional Review Board- Social Behavioural Educational Sciences (IRB; REF number: 2017-01-032) (CRC; REF number 612-2018). Participants were informed that participation was entirely voluntary and that they could withdraw at any point of time in the study. During the pre-screening call, participants were provided with the details of the study and gave their consent both during the call and before they proceeded with the online survey. Mothers of young children between the ages of three to seven residing in Singapore were invited to participate in this study. This study was distributed via advertisements on online platforms, requesting for mothers to participate in a study focusing on mother-child relationships. Some early intervention centres in Singapore also helped disseminate this information. Over the span of two years, 72 mothers of neurotypical children and over the span of three years, 73 mothers of children diagnosed with ASD participated in this study. There was a high rate of attrition among eligible participants, especially among mothers of children with ASD. Out of which, 52 mothers of neurotypical children completed the entire study (73.61% completion rate), while 26 mothers of children with ASD completed the study (35.62% completion rate). Mothers of children in both groups who included relations of others in their personal savouring were excluded from this analysis. The eventual sample in each group consisted of 52 mothers of neurotypical children and 26 mothers of children with ASD (N=78). The reasons for removal of incomplete data have been reported in the CONSORT flow chart in 1. In the ASD group, 26 mothers between the ages of 26 to 47 (M= 35.73, SD=4.84) of children who have been diagnosed with ASD between the ages of 3 to 7 (M= 5.44, SD=1.19) were in this group. The control group consisted of 52 mothers between the ages of 22 years to 44 years old (M= 35.12, SD= 4.34) of neurotypical children between the ages of 3 to 6 (M= 4.10, SD =1.19). To participate in this study, mothers in both groups were required to be 1) proficient in English as the study was conducted in English, 2) currently residing in Singapore 3) above 21 years old, 4) have a target child between the ages of 3-7 years old at the time of the survey, and 5) possess the ability to read and understand English proficiently. Furthermore, mothers in both groups should have no 1) known mental disorders and 2) severely impaired cognitive deficits that may prevent them from understanding and responding to the online questionnaires. Moreover, the target child in the ASD group should have an official ASD diagnosis. In contrast, the target child in the control group should not have any known diagnosis.

2.2. Procedure

A pre-screening call was made to interested participants to determine if they fit the criteria required for the study. Subjects were strongly encouraged to complete the study in one sitting, within the span of one week. Eligible participants were provided with a confirmation email link to the survey with their own unique study code. A brief explanation of savouring was given and consent from participants was sought prior to the administration of the study. The data was administered via an online data collection tool, Qualtrics (Qualtrics, Provo, UT). In addition to other scales, baseline measurements for PANAS [15] were taken before the random assignment of the experimental condition (of personal savouring or relational savouring) was administered. Participants in the personal savouring condition took an average of 147 seconds to submit the page on the savouring task while those in the relational savouring condition took an average of 118 seconds. The PANAS scale was re-administered to mothers in both conditions after the experimental savouring condition, and Generalised Anxiety Disorder-7 (GAD-7; [35,36]) was measured after the administration of savouring intervention. Demographic information of mothers was recorded in the last section of the survey. Upon completion of the study, a debriefing form was sent to participants, and they were given monetary compensation as a token of appreciation.




Figure 1. CONSORT Flow Chart depicting the reasons for removal of incomplete data in both mothers of neurotypical children and children with ASD.

2.3. Measures

2.3.1. Generalised Anxiety Disorder-7 (GAD-7).

The GAD-7 scale [35,36] is a 7-item self-report questionnaire that measures the degree of worry and anxiety. Items including, “Worrying too much about different things,” are rated on a response scale from 1 (not at all) to 3 (nearly every day). Items were then added to constitute a total score (range 0-21), in which higher scores represent greater severity of anxiety. The scale has been validated in 2740 primary care patients [35] and is demonstrated to have good sensitivity and specificity as a screener for GAD, panic, social anxiety, and posttraumatic stress disorder [36]. It has been proven to have excellent internal consistency ($\alpha = .92$), good test-retest reliability (intraclass correlation = 0.83) and strong criterion validity. Convergent validity was seen from the correlations with 2 anxiety measures: the Beck Anxiety Inventory ($r = 0.72$) and the anxiety subscale of the Symptom Checklist-90 ($r = 0.74$; [35]). The Cronbach’s alpha for the sample in this study was high at 0.95.

2.3.2. Positive Affect and Negative Affect Schedule (PANAS).

Mood was measured using the PANAS scale, an established self-report measure of PA and NA [15]. Mothers answered 20 questions from the general dimension of the PANAS-SF scale [37], which consisted of ten items from the general dimension of PA and 10 items which reflect negative moods as general PA and NA was our interest in this study. The PANAS scale was administered immediately preceding and succeeding the savouring condition. For each item, mothers were instructed to report their feelings on a scale of 1- 5 (1 ‘very slightly or not at all’ to 5 ‘extremely’). PA and NA are considered distinct from each other [15]. This measure has high validity with a Cronbach alpha coefficient of .89 to .85 respectively for PA and NA, and the test-retest correlation scores for both PA and NA respectively was .54 and .45 [15]. The scores for these corresponding 10 items of PA and NA were summed up pre and post-test and the changes in PA and NA were also tabulated. In this study, the Cronbach’s alpha coefficient was 0.97 for PA and 0.96 for NA.

2.4. Experimental condition

Through random assignment, mothers in the ASD and control group were placed into one of two different savouring conditions- relational and personal savouring. The tasks in these interventions involve mental reflection and are similar to the tasks in [?] and were developed in consultation with the senior author on that study.

2.4.1. Relational Savouring.

The relational savouring task prompted participants to reflect on an occasion where they felt connected, close to or in tune with their child, regardless if it was a significant milestone or a daily affair. Participants could proceed to the next page after a minute. Follow up questions involved guiding mothers to describe the positive experience and the details surrounding this event. Mothers were then encouraged to utilise the next two minutes to let their mind wander and place their attention on positive aspects of their chosen event (see [A1](#) in Appendix A).

2.4.2. Personal Savouring.

Identical to the relational savouring task, the only difference in the personal savouring condition was to exclude the presence of another in their savouring event. Mothers were therefore prompted to spend a minute reflecting on an occasion where they felt a positive and personal experience where no one else was present, it could potentially be a time where they felt a sense of achievement (see [A2](#) in Appendix A). Examples of the prompt include asking them to come up with a memory of a recent time when you have felt happy. This should be something you experienced on your own (without anybody else present) and something you enjoyed but haven't had time to really think about." Other questions included "How did you feel at the time?" and "What thoughts did you have at the time?"

2.5. Analytical plan

Sociodemographic information and descriptive statistics of mother's age and child's age would be reported for the two samples of mothers. Preliminary analyses on the types of pronouns participants used in the savouring task were conducted to determine if personal and relational savouring conditions significantly differed from each other. First-person singular pronouns ("I" and "me") and first-person plural pronouns ("us" and "we") and ("he", "she", "him" and "her") were extracted from participants' response to the savouring task, for personal and relational savouring, respectively. Chi-square tests were conducted on the frequency of pronoun category usage across the two savouring conditions. Inferential analyses addressing the three hypotheses were subsequently conducted. To test the first hypothesis, that savouring would lead to a decrease in PANAS-NA, a one-sided paired Wilcoxon signed rank test was conducted, comparing PANAS-NA post-savouring to pre-savouring levels. To test the second hypothesis, that savouring would increase PANAS-PA, a one-sided paired Wilcoxon signed rank test was also conducted, contrasting PANAS-PA post-savouring to pre-savouring scores. As there were two models to be tested, Bonferroni correction was applied onto alpha, such that $\alpha = 0.05/2 = 0.025$. To test the third exploratory hypothesis, that higher levels of GAD in the mother would result in a greater decrease in PANAS-NA, a nonparametric Kendall–Theil linear regression test was conducted where change in PANAS-NA ($PANAS-NA_{\Delta}$; Post-Pre) was the dependent variable, and GAD score was the independent factor. Subsequently, Pearson's correlation analysis would be conducted to determine the direction of effect.

3. Results

Sociodemographic information, and descriptive statistics of mother's age and child's age for the two samples of mothers are reported in [1](#).

Table 1. Sociodemographic information of participants.

	Sample 1: Mothers with Typical Children	Sample 2: Mothers with ASD Children
N	52	26
Savouring Condition	Personal: 22 Relational: 30	Personal: 10 Relational: 16
Age (Mothers)	Mean age of 35.1 years (SD = 4.34)	Mean age of 35.7 years (SD = 4.84)
Age (Children)	Mean age of 4.1 years (SD = 1.2)	Mean age of 5.4 years (SD = 1.2)
Sex (Children)	Female: 18 Male: 33	Female: 5 Male: 21
Education (Mothers)	4 GCE 'O' Levels 10 Diploma 38 Graduate Degree	3 GCE 'O' Levels School 7 Diploma 16 Graduate Degree
Employment Status	26 Employed Full-time 5 Employed Part-time 3 Self-employed 18 Unemployed	8 Employed Full-time 2 Employed Part-time 1 Self-employed 15 Unemployed
Marital Status	2 Unmarried 50 Married	26 Married
Household Income	10 \$1,000 - \$4,999 21 \$5,000 - \$8,999 11 \$9,000 - \$12,999 5 \$13,000 - \$19,999 2 > \$20,000	9 \$1,000 - \$4,999 9 \$5,000 - \$8,999 6 \$9,000 - \$12,999 1 \$13,000 - \$19,999 1 > \$20,000

3.1. Preliminary Analyses

3.1.1. Between savouring conditions.

To determine if there were significant differences across savouring conditions, chi-square conducted to compare the average usage of first-person singular pronouns ("I" and "me") and that of first-person plural pronouns ("us" and "we") and ("he", "she", "him" and "her") respectively for each savouring question. The lack of significant findings suggested that participants did not use the pronouns differently across conditions (see A3 and A4 in Appendix A). Since the two savouring conditions did not differ in the preliminary analyses, we decided to pool the two conditions together for the subsequent analyses.

3.1.2. Between groups of mothers.

Mean and standard deviation values of PA and NA for groups of mothers are reported in 2. To determine if there were significant differences across the two groups of mothers, a two-sample nonparametric Mann-Whitney test was conducted between the two groups for PANAS-PApre, PANAS-PApos, PANAS-NApre and PANAS-NApos values (see 2). Since no significant difference was found between the PA and NA values of the two groups of mothers, we decided to pool the two diagnostic groups together for the subsequent analyses. 2 shows pooled mothers' PA and NA scores before and after savouring conditions.

Table 2. Mean, standard deviations and Mann Whitney U Test of pre-savouring positive affect (PANAS-PA_{pre}), post-savouring positive affect (PANAS-PA_{pos}), pre-savouring negative affect (PANAS-NA_{pre}) and post-savouring negative affect (PANAS-NA_{pos}) for mothers of typical children and children with ASD.

	ASD		Typical		Mann-Whitney Test	
	Mean	SD	Mean	SD	U	p-value
PANAS-PA _{pre}	31.2	9.8	28	9.28	771	0.316
PANAS-PA _{pos}	29.0	10.9	29	11	666.5	0.924
PANAS-NA _{pre}	18.4	8.8	15	5.2	771.5	0.311
PANAS-NA _{pos}	16.4	7.8	14	5.4	760	0.370

3.2. Inferential Analyses

Since there was no significant difference in pronoun usage between savouring conditions, and no significant difference between the means of PA and NA between groups of mothers, inferential analyses were conducted on a pooled sample of mothers and savouring conditions. Shapiro-Wilk normality test was conducted on the pooled sample which found the pooled sample to be non-normal ($W = 0.846$, $p = 1.625e-07$). Subsequently, nonparametric tests were used for the inferential analyses. To test hypothesis 1, a one-sided paired dependent 2-group paired Wilcoxon signed rank test was conducted between PANAS-NA_{pos} and PANAS-NA_{pre} for each of the two groups of mothers, and for the pooled group of mothers (see 3), where PANAS-NA_{pos} was expected to be less than PANAS-NA_{pre}. Post savouring, PANAS-NA_{pos} was found to be significantly less than PANAS-NA_{pre} for mothers of typical children ($V = 274$, $p = 0.0132$) and for the pooled group of mothers ($V = 616$, $p = 0.00359$). Subsequently, the nonparametric Kendall–Theil linear regression test was conducted to determine the effect of GAD on PANAS-NA $_{\Delta}$. Analyses revealed a significant effect of GAD on PANAS-NA $_{\Delta}$ ($V = 449.5$, $p = 0.00733$). Pearson's correlation test later showed that savouring led to a greater decrease in PANAS-NA $_{\Delta}$ among mothers with greater GAD ($t = -3.27$, $df = 76$, $p = 0.002$; 3). To test hypothesis 2, a one-sided paired dependent 2-group paired Wilcoxon signed rank test was conducted between PANAS-PA_{pos} and PANAS-PA_{pre} for each of the two groups of mothers, and for the pooled group of mothers (see 3), where PANAS-PA_{pos} was expected to be greater than PANAS-PA_{pre}. However, no significant difference emerged between mothers' reported positive affect before and after savouring.

Table 3. Dependent 2-group paired Wilcoxon signed rank tests between pre-PA and post-PA, and pre-NA and post-NA for mothers of typical children, mothers of children with ASD and pooled mothers *Note: * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.005$.

Group	N	Paired Comparisons	Paired Wilcoxon Test	
			V	p-value
ASD	26	PANAS-PA _{pos} - PANAS-PA _{pre}	46.5	0.996
		PANAS-NA _{pos} - PANAS-NA _{pre}	71	0.0621
Typical	52	PANAS-PA _{pos} - PANAS-PA _{pre}	471	0.406
		PANAS-NA _{pos} - PANAS-NA _{pre}	274	0.0132*
Pooled	78	PANAS-PA _{pos} - PANAS-PA _{pre}	857.5	0.89
		PANAS-NA _{pos} - PANAS-NA _{pre}	616	0.00359**




Figure 2. Bar plot of PANAS-NA_{pos} and PANAS-NA_{pre} for each data index point in the pooled sample. Points below the blue line represent observations where PANAS-NA_{pre} is greater than PANAS-NA_{pos}.




Figure 3. Correlation plot of Generalized Anxiety (GAD) score against PANAS-NA Δ (Post-Pre).

4. Discussion

One primary goal of positive psychology is to improve well-being and develop scientifically sound activities to boost personal well-being. This study aimed to examine whether savouring increased positive affect and/or reduced negative affect among mothers of typical and ASD children. A secondary aim of this study was to examine if general anxiety levels influence the efficacy of a brief savouring in mothers. Supporting the first hypothesis and congruent with past literature on savouring among mothers of infants [?], this study found that there was a decrease in NA after savouring among mothers of young children in both the ASD and control groups. However, contrary to the second hypothesis, savouring failed to significantly increase PA among mothers across both ASD and control groups after this brief online savouring intervention was administered. This study further demonstrated in an exploratory analysis that a greater decrease in NA was observed after savouring among mothers with higher levels of generalised anxiety. Similar to [?], there were no significant differences found across both personal and relational savouring conditions among mothers. This may be partially explained by the demographic of the population that we are exploring - females in a collectivist culture of Singapore. Females tend to value interdependence and interconnectedness [38], and a similar trend is observed among collectivistic cultures [39]. Having been raised in a society which subscribes to these values, mothers in this study may be naturally inclined to include elements of others when savouring, even when they had been allocated to a personal savouring condition. It may also suggest that focusing on personal memories when savouring may not be natural or easy, and that when asked to focus on positive memories, many people gravitate toward the relational, in accordance with Bryant's framework [1]. Furthermore, in understanding the differences between relational and personal savouring, this may indicate a need for an in-person approach to savouring, such as having interveners to administer the intervention and guide people in engaging with the intervention as intended. In line with literature on savouring [?], both personal savouring and relational savouring resulted in significantly less NA in parents of typically developing children in comparison to controls. Contrary to the second hypothesis, this study did not find any significant changes in PANAS-PA. A cultural perspective may provide one possible explanation for this result. While savouring and affect are closely associated with happiness [1], happiness is not cross-

culturally universal. While North Americans define happiness based on personal achievement and strive for PA, East Asians tend to relate happiness with interpersonal connectedness and balancing PA and NA [40] instead of leaning towards hedonic emotions. Easterners may also be more likely to endorse dampening of positive events, which may result in fewer changes in PA [41]. Savouring also thrives under conditions whereby the person can be fully present to savour [1]. Yet, distractions such as stress and multitasking among mothers may impede savouring. Ideally, savouring would be most effective after stress has occurred or been reduced [1]. Yet, mothers experience more ongoing stress and tiredness during caregiving than fathers [42], and mothers of children with ASD endure chronic stress [43], which may hinder a mother's ability to attend closely to positive moments and may distract mothers from fully savouring. In addition, a mother's role may typically require them to multitask in order to successfully manage affairs in the family, which may divide their attention. Multitasking may make it difficult for mothers to be in the moment [1,44], which could result in dampening of PA [45]. Furthermore, theoretically, the duration of savouring may interact with the ability to fully savour [1]. As such, while this online savouring intervention may be too brief to be considered effective in increasing PA, it is possible that increases in PA savouring may not happen immediately, but are seen over a longer period of time [46]. This study provides support for the effectiveness of positive interventions on patients with anxiety [47]. The role of self-esteem and self-efficacy may be potential mechanisms which alleviates anxiety symptoms. The process of savouring encourages mothers to remember and relish their past achievements. Correspondingly, lower maternal self-efficacy has been found to be correlated to higher self-reported maternal anxiety [48]. Furthermore, relational savouring may focus on moments whereby mothers have been successful in protecting, supporting or aiding in their child's growth salient to mothers. This may, in turn, increase the self-efficacy of mothers, and hence, reduce NA of mothers. Moreover, more generally, savouring positive memories pertain to reliving one's successes, and successes have been found to increase state self-esteem [49]. The relationship between anxiety and self-esteem has been found to be symmetrically reciprocal, whereby anxiety predicts lower self-esteem, and the vice versa (see review in [50]). Since self-esteem and self-efficacy were not explicitly studied in this paper, future research can better investigate the links between savouring, self-efficacy and self-esteem. A decrease in NA experienced by mothers with higher levels of anxiety is also consistent with another study done on another positive psychology concept-mindfulness. Among those with high anxiety sensitivity but not low anxiety sensitivity, trait mindfulness was significantly related to a reduced NA when there was a brief stress inducement [51]. Given that savouring overlaps with mindfulness to the extent that it demands for people to be mindful of a particular experience [1], a savouring intervention may similarly activate the same mechanisms to reduce NA. However, the effects of savouring on PA may not be readily increased as mothers with higher levels of GAD may more readily engage in dampening of PA when it occurs [52]. There could also be a floor effect of NA among mothers with already low levels of anxiety, such that mothers who are higher in anxiety also have higher levels of NA. Therefore, partaking in savouring tasks would be more likely to be able to reduce already comparatively high levels of NA among mothers, compared to mothers with lower levels of anxiety with potentially already lower levels of NA at baseline. Although there was a significantly greater decrease in NA after savouring among mothers with higher anxiety, a significant increase in PA was not found among mothers. Several reasons could have underpinned the null findings for PA. First, this may be attributed to the idea that people with anxiety (after controlling for depressive symptoms) also tend to endorse dampening when they experience PA, and were also less likely to endorse savouring beliefs [52]. As such, being less inclined to savour coupled with dampening of positive affect may make it harder for people with anxiety to experience an increase in PA. Future studies could control for dampening of affect among mothers, and explore if there were ways to counter this mentality. Moreover, as depression is closely related to anxiety, future studies could also explore the possibility of the effects savouring have on depression. Second, the Asian culture of dampening PA may also be involved whereby mothers (regardless of anxiety levels) engage in dampening when they savour positive experiences, reducing their ability to increase PA [41]. Since dampening is a thought process considered to occur among mothers in Asian cultures and those with higher levels of anxiety, which thus might hinder their ability

to increase PA after savouring, future studies should account for the effects of dampening in savouring interventions. There are some caveats which should be taken into consideration when interpreting the results of this study. First, a control group that was present in the [6] study was missing in the present study. The presence of a control group would have allowed us to rule out extraneous factors not directly related to the savouring task, such as mothers obtaining time to themselves when using the phone to complete the online intervention, that could have reduced negative affect. Future studies should include a control condition to strengthen the argument for the efficacious effects of savouring. Second, the savouring intervention was held for only one session, and no follow-up assessment was conducted to evaluate how long the effect of the intervention lasts for. Since mothers of both neurotypical children and children with ASD may experience recurring anxiety on a daily basis, future studies should be conducted over a longer period of time to investigate the frequency with which savouring needs to be practised to produce long-term psychological wellbeing outcomes. Third, parents were also encouraged to complete the survey in one sitting; however, there were some parents who completed the PANAS before and after the savouring intervention over one week. It is possible that maturation effects could have occurred due to other situations or experiences that took place during this interval, affecting the pre- or post- PANAS scores if they were not completed immediately before and after the savouring activity. Fourth, there is also a high attrition rate of mothers who did not complete this study, and as such, their responses were removed from the analysis. It is, however, unclear why they did not continue with the study. A possibility could be that mothers who dropped out may have children with more severe needs, thus they may not be able to afford the time to complete the study. However, this bias may also be reflective of mothers who are keen on understanding and improving the relationship with their child. Upon conducting Mann-Whitney U Tests and Fisher's exact tests to determine if there were demographic differences between parents who did not complete the study and those who did, it was found that there were no significant differences between the two groups across all demographic variables ($p > .05$). Fifth, although the environmental context where mothers were doing the online intervention was not controlled for, this may increase the ecological validity of our study as in an online intervention, the environmental context may vary across mothers. Future studies may extend on this research to consider the environment of the mother as a factor that may enhance PA and NA. Moreover, this study may include a randomised control group in future to determine causality of savouring in future. This current study adds to a body of literature that savouring may be especially useful to enhance affect among mothers by decreasing their NA. Mothers with anxiety may be able to harness these benefits. Mothers of preschool-aged children may stand to decrease negative feelings by savouring past experiences of achievements that had occurred both with and without their child. This is more so among mothers who experience greater levels of anxiety. Savouring may, therefore, be a possible intervention incorporated as an emotional coping strategy, perhaps in conjunction with other problem-solving strategies among mothers and especially those with anxiety to decrease NA. In problem-coping strategies (methods which tackle the issue or alter the source of stress), the relationship between affect and learning is complex and thus, high levels of NA may hinder a person's ability to learn [53]. However, such strategies are still likely to play an important role in helping mothers cope [54]. Therefore, to enhance mothers' receptivity to these strategies, incorporating elements of emotion coping that target managing affect (such as this savouring intervention) to decrease NA, at least temporarily, may be beneficial, especially among mothers with higher levels of anxiety. Moreover, lower levels of NA is linked to better psychological health. Therefore, savouring as an intervention may be helpful to improve the mental well being of mothers of young children with ASD and neurotypical children alike.

Author Contributions: Conceptualization, G.E. and J.B.; formal analysis, A.A., C.A.H.; data curation, A.S.P., C.A.H.; writing—original draft preparation, A.A., A.S.P., C.A.H.; writing—review and editing, A.A., G.E.G., J.B., G.E.; visualization, A.A.; funding acquisition, G.E. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the NTU NAP SUG 2015.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board of Nanyang Technological University(protocol code 2017-01-032).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: In this section, please provide details regarding where data supporting reported results can be found, including links to publicly archived datasets analyzed or generated during the study. Please refer to suggested Data Availability Statements in section “MDPI Research Data Policies” at <https://www.mdpi.com/ethics>. You might choose to exclude this statement if the study did not report any data.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A.**Table A1.** Relational Savouring Memory Prompt.

Parents of young children often tell us they don't have much time to focus on positive experiences they've had with their children because they are so busy and don't have time to stop and reflect. For this task, we would like you to come up with a memory of a time when you felt extremely connected, close, or "in-sync" with your child. We are especially interested in hearing about a time when you found joy in helping your child grow, or a time when your child needed you and you were there for your child. It may be a time when you felt like you comforted, soothed, protected, or supported your child. Feel free to choose something that you felt was a milestone or something simple that happens on a daily basis. Please spend one minute focusing on this memory. You will be asked some questions about the details of this event in the following section.

1. Using as much detail as possible, describe what happened. Please enter a minimum of 100 characters.
2. What was the air like? What was the weather like? Please enter a minimum of 20 characters.
3. What time of day did the moment occur? Please enter a minimum of 20 characters.
4. What were you wearing? Please enter a minimum of 20 characters.
5. What was your child wearing? Please enter a minimum of 20 characters.
6. How did you feel at the time? (excited, proud, calm, relaxed etc.) Please enter a minimum of 100 characters.
7. What thoughts did you have at the time? About your child? About your relationship? Please enter a minimum of 100 characters.
8. What thoughts are you having now about your child and about your relationship? Please enter a minimum of 100 characters.

Mindfulness prompt: For the next two minutes, please let your mind wander in any way you'd like related to this event. You may want to think about things you were asked about earlier or you may want to think about how this memory is related to your other relationships and your life. It's normal for your mind to wander to other topics or feelings - if you notice that your mind has wandered, just gently bring it back to the positive aspects of this event. Let your mind wander in any way you'd like, but try to keep focused on the positive parts of this memory.

Table A2. Personal Savouring Memory Prompt.

Parents of young children often tell us they don't have much time to focus on themselves because they are so busy taking care of their children. The goal of this task is to help you focus on something positive that happened in your life lately. What I'd like you to do is to come up with a memory of a recent time when you have felt happy. This should be something you experienced on your own (without anybody else present) and something you enjoyed but haven't had time to really think about. The memory should also be something you would like to spend some time thinking about. It can be something as simple as enjoying a good meal, or taking a nice walk. It could also be something as major as getting a promotion or accomplishing a big task. The important thing is to come up with a time when you were feeling happy, content, or relaxed.

Please spend one minute focusing on this memory. You will be asked some questions about the details of this event in the following section.

1. Using as much detail as possible, describe what happened. Please enter a minimum of 100 characters.
2. What was the air like? What was the weather like? Please enter a minimum of 20 characters.
3. What time of day did the moment occur? Please enter a minimum of 20 characters.
4. What were you wearing? Please enter a minimum of 20 characters.
5. How did you feel at the time? (excited, proud, calm, relaxed etc.) Please enter a minimum of 100 characters.
6. What thoughts did you have at the time? Please enter a minimum of 100 characters.
7. What thoughts are you having now? Please enter a minimum of 100 characters.

Mindfulness prompt: For the next two minutes, please let your mind wander in any way you'd like related to this event. You may want to think about things you were asked about earlier or you may want to think about how this memory is related to your other relationships and your life. It's normal for your mind to wander to other topics or feelings - if you notice that your mind has wandered, just gently bring it back to the positive aspects of this event. Let your mind wander in any way you'd like, but try to keep focused on the positive parts of this memory.

Table A3. Chi-square Tests Comparing the Average Usage of First-person Singular Pronouns (“I” and “me”) and of First-person Plural Pronouns (“us” and “we”) for Each Savouring Question.

Savouring question	X ²	p
What happened?	3.07	.38
Weather?	0.37	.95
Time of day?	0.45	.93
Wearing?	1.08	.78
Feeling?	3.22	.36
Thoughts then?	3.35	.34
Thoughts now?	3.53	.32

Table A4. Chi-square Tests Comparing the Average Usage of First-person Singular Pronouns (“I” and “me”) and of First-person Plural Pronouns (“he”, “she”, “him” and “her”) for Each Savouring Question.

Savouring question	X2	p
What happened?	2.73	.44
Weather?	0.31	.96
Time of day?	0.37	.95
Wearing?	1.08	.78
Feeling?	3.22	.36
Thoughts then?	3.35	.34
Thoughts now?	2.64	.45

References

- Bryant, F.B.; Veroff, J. *Savoring: A New Model of Positive Experience*; Psychology Press, 2017.
- Bryant, F. Savoring Beliefs Inventory (SBI): A scale for measuring beliefs about savouring. *Journal of Mental Health* **2003**, *12*, 175–196.
- Borelli, J.L.; Smiley, P.A.; Kerr, M.L.; Hong, K.; Hecht, H.K.; Betsy Blackard, M.; Falasiri, E.; Cervantes, B.R.; Bond, D.K. Relational savoring: An attachment-based approach to promoting interpersonal flourishing. *Psychotherapy* **2020**, *57*, 340–351.
- Borelli, J.L.; Sbarra, D.A.; Snively, J.E.; McMakin, D.L.; Coffey, J.K.; Ruiz, S.K.; Wang, B.A.; Chung, S.Y. With or without you: Preliminary evidence that attachment avoidance predicts nondeployed spouses’ reactions to relationship challenges during deployment. *Professional Psychology: Research and Practice* **2014**, *45*, 478–487.
- Borelli, J.L.; Rasmussen, H.E.; Burkhart, M.L.; Sbarra, D.A. Relational savoring in long-distance romantic relationships. *Journal of Social and Personal Relationships* **2015**, *32*, 1083–1108.
- Burkhart, M.L.; Borelli, J.L.; Rasmussen, H.E.; Sbarra, D.A. Cherish the good times: Relational savoring in parents of infants and toddlers. *Personal Relationships* **2015**, *22*, 692–711.
- Hastings, R.P.; Taunt, H.M. Positive perceptions in families of children with developmental disabilities. *Am. J. Ment. Retard.* **2002**, *107*, 116–127.
- Volkmar, F.R.; Lord, C.; Bailey, A.; Schultz, R.T.; Klin, A. Autism and pervasive developmental disorders. *Journal of Child Psychology and Psychiatry* **2004**, *45*, 135–170.
- Lord, C.; Risi, S.; DiLavore, P.S.; Shulman, C.; Thurm, A.; Pickles, A. Autism from 2 to 9 years of age. *Arch. Gen. Psychiatry* **2006**, *63*, 694–701.
- Christensen, D.L.; Bilder, D.A.; Zahorodny, W.; Pettygrove, S.; Durkin, M.S.; Fitzgerald, R.T.; Rice, C.; Kurzius-Spencer, M.; Baio, J.; Yeargin-Allsopp, M. Prevalence and Characteristics of Autism Spectrum Disorder Among 4-Year-Old Children in the Autism and Developmental Disabilities Monitoring Network. *J. Dev. Behav. Pediatr.* **2016**, *37*, 1–8.
- Ardhanareeswaran, K.; Volkmar, F. Introduction. Focus: autism spectrum disorders. *Yale J. Biol. Med.* **2015**, *88*, 3–4.
- Baio, J.; Wiggins, L.; Christensen, D.L.; Maenner, M.J.; Daniels, J.; Warren, Z.; Kurzius-Spencer, M.; Zahorodny, W.; Robinson, C.; Rosenberg, White, T.; Durkin, M.S.; Imm, P.; Nikolaou, L.; Yeargin-Allsopp, M.; Lee, L.C.; Harrington, R.; Lopez, M.; Fitzgerald, R.T.; Hewitt, A.; Pettygrove, S.; Constantino, J.N.; Vehorn, A.; Shenouda, J.; Hall-Lande, J.; Van, K.; Naarden.; Braun.; Dowling, N.F. Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years — Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2014. *MMWR. Surveillance Summaries* **2018**, *67*, 1–23.
- Smith, L.E.; Hong, J.; Seltzer, M.M.; Greenberg, J.S.; Almeida, D.M.; Bishop, S.L. Daily experiences among mothers of adolescents and adults with autism spectrum disorder. *J. Autism Dev. Disord.* **2010**, *40*, 167–178.
- Seltzer, M.M.; Almeida, D.M.; Greenberg, J.S.; Savla, J.; Stawski, R.S.; Hong, J.; Taylor, J.L. Psychosocial and Biological Markers of Daily Lives of Midlife Parents of Children with Disabilities. *Journal of Health and Social Behavior* **2009**, *50*, 1–15.
- Watson, D.; Clark, L.A.; Tellegen, A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J. Pers. Soc. Psychol.* **1988**, *54*, 1063–1070.
- Tran, V. Positive Affect Negative Affect Scale (PANAS). *Encyclopedia of Behavioral Medicine* **2013**, pp. 1508–1509.
- Larsen, J.T.; McGraw, A.P.; Cacioppo, J.T. Can people feel happy and sad at the same time? *J. Pers. Soc. Psychol.* **2001**, *81*, 684–696.
- Zhou, T.; Yi, C. Parenting Styles and Parents’ Perspectives on How Their Own Emotions Affect the Functioning of Children with Autism Spectrum Disorders. *Family Process* **2014**, *53*, 67–79.
- Margalit, M.; Ankonina, D.B. Positive and negative affect in parenting disabled children. *Counselling Psychology Quarterly* **1991**, *4*, 289–299.
- Hodapp, R.M.; Dykens, E.M.; Evans, D.W.; Merighi, J.R. Maternal emotional reactions to young children with different types of handicaps. *J. Dev. Behav. Pediatr.* **1992**, *13*, 118–123.
- Hurley, D.B.; Kwon, P. Savoring Helps Most When You Have Little: Interaction Between Savoring the Moment and Uplifts on Positive Affect and Satisfaction with Life. *Journal of Happiness Studies* **2013**, *14*, 1261–1271.
- Lyubomirsky, S.; Layous, K. How Do Simple Positive Activities Increase Well-Being? *Current Directions in Psychological Science* **2013**, *22*, 57–62.
- Lyubomirsky, S.; King, L.; Diener, E. The benefits of frequent positive affect: does happiness lead to success? *Psychol. Bull.* **2005**, *131*, 803–855.

24. Ekas, N.V.; Whitman, T.L. Adaptation to daily stress among mothers of children with an autism spectrum disorder: the role of daily positive affect. *J. Autism Dev. Disord.* **2011**, *41*, 1202–1213.
25. Folkman, S.; Moskowitz, J.T. Positive affect and the other side of coping. *American Psychologist* **2000**, *55*, 647–654.
26. Estes, A.; Munson, J.; Dawson, G.; Koehler, E.; Zhou, X.H.; Abbott, R. Parenting stress and psychological functioning among mothers of preschool children with autism and developmental delay. *Autism* **2009**, *13*, 375–387.
27. Ghanizadeh, A.; Alishahi, M.J.; Ashkani, H. Helping families for caring children with autistic spectrum disorders. *Arch. Iran. Med.* **2009**, *12*, 478–482.
28. Hastings, R.P.; Kovshoff, H.; Ward, N.J.; degli Espinosa, E.; Brown, T.; Remington, B. Systems analysis of stress and positive perceptions in mothers and fathers of pre-school children with autism. *J. Autism Dev. Disord.* **2005**, *35*, 635–644.
29. Davis, N.O.; Carter, A.S. Parenting stress in mothers and fathers of toddlers with autism spectrum disorders: associations with child characteristics. *J. Autism Dev. Disord.* **2008**, *38*, 1278–1291.
30. Phetrasuwan, S.; Shandor Miles, M. Parenting stress in mothers of children with autism spectrum disorders. *J. Spec. Pediatr. Nurs.* **2009**, *14*, 157–165.
31. Wang, J.; Hu, Y.; Wang, Y.; Qin, X.; Xia, W.; Sun, C.; Wu, L.; Wang, J. Parenting stress in Chinese mothers of children with autism spectrum disorders. *Soc. Psychiatry Psychiatr. Epidemiol.* **2013**, *48*, 575–582.
32. Bitsika, V.; Sharpley, C.F. Stress, Anxiety and Depression Among Parents of Children With Autism Spectrum Disorder. *Australian Journal of Guidance and Counselling* **2004**, *14*, 151–161.
33. Green, J.; Charman, T.; McConachie, H.; Aldred, C.; Slonims, V.; Howlin, P.; Le Couteur, A.; Leadbitter, K.; Hudry, K.; Byford, S.; Barrett, B.; Temple, K.; Macdonald, W.; Pickles, A.; PACT Consortium. Parent-mediated communication-focused treatment in children with autism (PACT): a randomised controlled trial. *Lancet* **2010**, *375*, 2152–2160.
34. Watson, D.; Naragon-Gainey, K. On the specificity of positive emotional dysfunction in psychopathology: evidence from the mood and anxiety disorders and schizophrenia/schizotypy. *Clin. Psychol. Rev.* **2010**, *30*, 839–848.
35. Spitzer, R.L.; Kroenke, K.; Williams, J.B.W.; Löwe, B. A Brief Measure for Assessing Generalized Anxiety Disorder. *Archives of Internal Medicine* **2006**, *166*, 1092.
36. Kroenke, K.; Spitzer, R.L.; Williams, J.B.W.; Monahan, P.O.; Löwe, B. Anxiety Disorders in Primary Care: Prevalence, Impairment, Comorbidity, and Detection. *Annals of Internal Medicine* **2007**, *146*, 317.
37. Watson, D.; Clark, L.A. The PANAS-X: Manual for the Positive and Negative Affect Schedule - Expanded Form. *University of Iowa* **1994**.
38. Cross, S.E.; Madson, L. Models of the self: self-construals and gender. *Psychol. Bull.* **1997**, *122*, 5–37.
39. Ross, M.; Elaine Xun, W.Q.; Wilson, A.E. Language and the Bicultural Self. *Personality and Social Psychology Bulletin* **2002**, *28*, 1040–1050.
40. Uchida, Y.; Norasakkunkit, V.; Kitayama, S. Cultural Constructions of Happiness: Theory and Empirical Evidence. *Happiness Studies Book Series* **2013**, pp. 269–280.
41. Miyamoto, Y.; Ma, X. Dampening or savoring positive emotions: a dialectical cultural script guides emotion regulation. *Emotion* **2011**, *11*, 1346–1357.
42. McDonnell, C.; Luke, N.; Short, S.E. Happy Moms, Happier Dads: Gendered Caregiving and Parents' Affect. *Journal of Family Issues* **2019**, *40*, 2553–2581.
43. Gray, D.E. Ten years on: a longitudinal study of families of children with autism. *Journal of Intellectual & Developmental Disability* **2002**, *27*, 215–222.
44. Offer, S.; Schneider, B. Revisiting the Gender Gap in Time-Use Patterns. *American Sociological Review* **2011**, *76*, 809–833.
45. Quoidbach, J.; Berry, E.V.; Hansenne, M.; Mikolajczak, M. Positive emotion regulation and well-being: Comparing the impact of eight savoring and dampening strategies. *Personality and Individual Differences* **2010**, *49*, 368–373.
46. Seligman, M.E.P.; Steen, T.A.; Park, N.; Peterson, C. Positive Psychology Progress: Empirical Validation of Interventions. *American Psychologist* **2005**, *60*, 410–421.
47. Chakhssi, F.; Kraiss, J.T.; Sommers-Spijkerman, M.; Bohlmeijer, E.T. The effect of positive psychology interventions on well-being and distress in clinical samples with psychiatric or somatic disorders: a systematic review and meta-analysis. *BMC Psychiatry* **2018**, *18*.
48. Rezendes, D.L.; Scarpa, A. Associations between Parental Anxiety/Depression and Child Behavior Problems Related to Autism Spectrum Disorders: The Roles of Parenting Stress and Parenting Self-Efficacy. *Autism Res. Treat.* **2011**, *2011*, 395190.
49. Rosi, A.; Cavallini, E.; Gamboz, N.; Vecchi, T.; Van Vugt, F.T.; Russo, R. The Impact of Failures and Successes on Affect and Self-Esteem in Young and Older Adults. *Frontiers in Psychology* **2019**, *10*.
50. Sowislo, J.F.; Orth, U. Does low self-esteem predict depression and anxiety? A meta-analysis of longitudinal studies. *Psychol. Bull.* **2013**, *139*, 213–240.
51. Arch, J.J.; Craske, M.G. Laboratory stressors in clinically anxious and non-anxious individuals: The moderating role of mindfulness. *Behaviour Research and Therapy* **2010**, *48*, 495–505.
52. Eisner, L.R.; Johnson, S.L.; Carver, C.S. Positive affect regulation in anxiety disorders. *J. Anxiety Disord.* **2009**, *23*, 645–649.
53. Rowe, A.D.; Fitness, J. Understanding the Role of Negative Emotions in Adult Learning and Achievement: A Social Functional Perspective. *Behav. Sci.* **2018**, *8*.
54. Essex, E.L.; Seltzer, M.M.; Krauss, M.W. Differences in coping effectiveness and well-being among aging mothers and fathers of adults with mental retardation. *Am. J. Ment. Retard.* **1999**, *104*, 545–563.