

Review

Control of ABA signaling and crosstalk with other hormones by selective degradation of pathway components

Agnieszka Sirko*, Anna Wawrzyńska*, Jerzy Brzywczy, Marzena Sienko

Laboratory of Plant Protein Homeostasis, Institute of Biochemistry and Biophysics, Polish Academy of Sciences, ul. Pawinskiego 5A, 02-106 Warsaw, Poland

* Correspondence: asirko@ibb.waw.pl (AS); blaszczyk@ibb.waw.pl (AW)

Abstract: A rapid and appropriate genetic and metabolic acclimation, which is crucial for plants' survival in a changing environment, is maintained due to the coordinated action of plant hormones and cellular degradation mechanisms influencing proteostasis. The plant hormone abscisic acid (ABA) rapidly accumulates in plants in response to environmental stress and plays a pivotal role in the reaction to various stimuli. Increasing evidence demonstrates a significant role of autophagy in controlling ABA signaling. This field has been extensively investigated and new discoveries are constantly being provided. We present updated information on the components of the ABA signaling pathway, particularly on transcription factors modified by different E3 ligases. Then, we focus on the role of selective autophagy in ABA pathway control and review novel evidence on the involvement of autophagy in different parts of the ABA signaling pathway that are important for crosstalk with other hormones, particularly cytokinins and brassinosteroids.

Keywords: abscisic acid; autophagy; brassinosteroids; cytokinins; hormone crosstalk; ubiquitin

1. Introduction

Plants are often subjected to various detrimental environmental stressors. Despite their sessile lifestyle, plants are able to maintain their internal equilibrium when exposed to external challenges. This balance, called homeostasis, can be achieved by internal mechanisms that alter cell metabolism, thanks to which the whole organism can maintain relatively constant conditions. Such rapid and appropriate genetic and metabolic acclimation, which is a crucial issue for plants' survival, is maintained due to the coordinated action of plant hormones and cellular degradation mechanisms influencing proteostasis.

Protein homeostasis (or proteostasis) integrates cellular pathways that mediate biogenesis, folding, trafficking, and degradation of polypeptides to maintain the required concentrations of all proteins that compose the proteome. This field has been intensively investigated by numerous research groups [1]. Proteins are often tagged for removal by ubiquitination. Polyubiquitinated proteins and protein aggregates are degraded via the ubiquitin-proteasome system (UPS) pathway and via the autophagy-vacuolar route. The UPS pathway is restricted in its ability to degrade aggregated proteins, which are too large to pass through the narrow proteasome entrance channel [2]. Thus, autophagy represents a major mechanism in plants for degrading macromolecular ubiquitinated protein aggregates as a response to stressors [3,4]. There is growing evidence that these two degradation systems are interconnected and form an integrated quality control network in response to general cellular stress [5]. Many reviews summarize the significance of ubiquitination in plant stress responses [6–8].

Not only protein degradation mechanisms but also phytohormones play a pivotal role in plant responses to various stimuli. Although the relationships between them are poorly understood, it is generally accepted that extensive crosstalk and multiple regulatory loops are responsible for maintaining the balance of growth and stress responses [9].

Phytohormones are master regulators of plant growth, development, and stress response. Of them, abscisic acid (ABA) is rapidly accumulated in plants in response to abiotic stress [10]. Like most other hormones, ABA is involved in developmental processes and in the response to adverse environmental stressors [11]. Plants exposed to abiotic stress quickly activate the ABA signaling cascade, resulting in the activation of ABA-responsive transcription factors and the induction of ABA-responding genes. However, after achieving stress tolerance, it is necessary to terminate (or attenuate) the ABA pathway. The balance is maintained by selective degradation of specific components of the ABA signaling pathway upon their tagging for decay by ubiquitination. Conventional degradation of the tagged components takes place via the proteasome; however, increasing evidence demonstrates a significant role of autophagy in controlling ABA signaling (for recent reviews [9,12–16]). In addition to this interplay with degradation systems, an additional layer of complexity is provided by extensive crosstalk between ABA and other phytohormones (for reviews [17–20]).

The above issues have been frequently reviewed; however, the field has been extensively investigated, and new discoveries are constantly being provided. The purpose of this review was to update information on the selective degradation of components of the ABA signaling pathway, particularly on the control of ABA-responsive transcription factors and the role of selective autophagy in ABA pathway control. We also reviewed novel evidence on the involvement of autophagy in different parts of the ABA signaling pathway that are important for crosstalk with other hormones, particularly cytokinins and brassinosteroids.

2. Principles of ABA perception and signaling and their control by selective degradation systems

Reversible protein phosphorylation is a posttranslational modification catalyzed by protein kinases and phosphatases. It acts as a molecular switch for almost all key events of cell metabolism and signaling in eukaryotes. Additionally, in the field of ABA signal transduction, protein kinases and phosphatases play pivotal roles, forming a reversible protein phosphorylation cascade, including ABA receptors [pyrabactin resistance/regulatory components of ABA receptor (PYR/RCAR)], type 2C protein phosphatases (PP2Cs), protein kinases [Snf1 (sucrose-non-fermentation 1)-related kinases subfamily 2, SnRK2s], and downstream targets. The key function of RCARs is to indirectly control the activity of SnRK2s, which phosphorylate numerous stress-activated targets in the response to ABA, including ABA-responsive transcription factors (TFs) [21,22]. This is accomplished through a negative regulatory pathway, involving RCARs, PP2C, and SnRK2 proteins and E3 ubiquitin ligases (Figure 1).




Figure 1. Abscisic acid (ABA) signaling in stress and growth. In growth promoting conditions, ABA receptors (RCARs), protein kinases (SnRK2s) dephosphorylated by type 2C protein phosphatases (PP2Cs), and ABA-responsive transcription factors (TF) are marked for degradation with ubiquitin (Ub) by specific E3 ligases (E3). In stress conditions, ABA accumulates and binds to RCARs allowing for interaction with PP2Cs to inhibit phosphatase activity. SnRK2s are released to phosphorylate and control the activity of downstream targets to trigger physiological responses. Blue circles indicate ubiquitination, yellow circles indicate phosphorylation events.

When ABA accumulates in cells in response to environmental stress or developmental cues, ABA binds to RCARs and triggers a conformational change that allows the binary complex (receptor-ABA) to physically interact with PP2Cs and inhibit its phosphatase activity. The various RCAR-PP2C heterodimers preclude binding of SnRK2s to PP2Cs, thus stimulating SnRK2s kinase activity. Consequently, SnRK2s are released to phosphorylate and control the activity of downstream factors to trigger physiological responses [23].

Several studies have suggested that plant responses to ABA are complemented by the enhanced degradation of PP2Cs by different E3 ligases (i.e., U-box type, RING type, and multimeric CRL type), which can facilitate ABA signaling and play an important role to prevent excessive accumulation of PP2Cs. However, E3 ligases are not only able to enhance ABA signaling but also facilitate the resetting of ABA signaling when tolerance to the stressor has been achieved (for review [15]).

3. Role of ubiquitination and selective degradation in ABA perception and signaling

Ubiquitination is (next to phosphorylation) another reversible, post-translational modification, originally discovered as a critical element in highly regulated proteolysis and regarded as essential for many other cellular processes. Ubiquitin (Ub) serves as a reusable tag that decides the cellular fate of the proteins and is recognized by numerous Ub-binding proteins directing the target proteins for selective turnover (for recent reviews [24–26]). Ub, often in the form of polymeric chains, is covalently attached to the target proteins through a three-step E1-E2-E3 conjugation cascade (Figure 2).




Figure 2. The ubiquitination cascade. Ubiquitin-activating enzyme (E1) activates ubiquitin (Ub) in an ATP-dependent manner, and next the thioester-linked Ub is transferred to Ub-conjugating enzyme (E2). Subsequently, E2 enzymes catalyze the attachment of Ub to a substrate using Ub ligase (E3). E3s, responsible for substrate recognition, transfer Ub either directly from E2 or form an E3-Ub intermediate prior to the transfer.

The process starts with E1 (or Ub-activating enzyme) in which activated Ub in an ATP-dependent manner, through a thioester bond between the C-terminus of Ub and a cysteine residue of E1, and next the thioester-linked Ub is transferred onto a cysteine

residue of E2 (or Ub-conjugating enzyme) by transesterification to form an unstable E2–Ub intermediate. Next, E2 enzymes catalyze the attachment of Ub to lysine (Lys) residues of target proteins using an E3 (or Ub ligase). E3 enzymes are responsible for substrate recognition and either directly transfer the Ub to the substrates from E2 or form a E3–Ub intermediate prior to the transfer. In successive rounds, additional Ub moieties can be attached to one of the seven lysine residues (Lys6, Lys11, Lys27, Lys29, Lys33, Lys48, and Lys63) or the N-terminal methionine (Met1) of Ub to produce poly-Ub chains. The end-product is a Ub–protein conjugate that contains an isopeptide bond between the C-terminal glycine residue of Ub and one of the Lys residues in the substrate. These structurally distinct ubiquitylation patterns are recognized by various effector proteins harboring a ubiquitin-binding domain (UBDs) to result in diverse downstream signals. For example, poly-Ub chains linked by Lys48 target substrates to the 26S proteasome for degradation, whereas poly-Ub chains linked by Lys63 direct substrates to autophagosomes, both with the concomitant release of Ub moieties for reuse ([26] and references therein).

The E3 Ub ligases can be classified into single- and multi-subunit groups (for review [27]). The single-subunit group includes several subfamilies based upon their mechanisms of action and the presence of specific domains: HECT (homology to E6-AP C terminus), RING (really interesting new gene), and U-box type E3s. The multi-subunit group, cullin-RING box1-ligases (CRLs), are further divided into four subfamilies: SCF (S phase kinase-associated protein 1–cullin 1–F-box), BTB (Bric-a-brac–Tramtrack–Broad complex), DDB (DNA damage-binding domain-containing), and APC (anaphase-promoting complex).

Three E2 enzymes, UBC32, UBC33, and UBC34 play a negative role in the drought stress response and ABA signaling [28], and numerous E3 ligases are involved in the inhibition of ABA signaling in optimal conditions, ABA induction upon environmental stress, and ABA attenuation after achieving stress tolerance [15]. The currently known components of ABA perception and signaling and E3 ligases involved in the process are listed in Table 1.

Table 1. ABA pathway components targeted by E3 ligases.

Pathway component	Specific target	E3 ligase [Reference] (remarks)
ABA Receptors	RCAR1/PYL9	DDA1 [29]; REA1 [30]; PUB22 [31]; PUB23 [31]
	RCAR3/PYL8	RIFP1 [32]; DDA1 [29]
	RCAR10/PYL4	DDA1 [29]; RSL1 [33]; UBC26 (E2)-RFA4 (E3) [34]; RFA1 [34]
	RCAR11/PYR1	RSL1 [33]; RFA1 [34]; RFA4 [34]
Type 2C protein phosphatases (PP2C)	ABI1	PUB12 [35]; PUB13 [35]; BPM3 [36]; BPM5 [36]; UBC27/AIRP3 [37]; COP1 [38]
	ABI2	RGLG1 [39]; RGLG5 [39]
	HAB1	BPM3 [36]; BPM5 [36]
	HAB2/NHL29	RGLG1 [39]; RGLG5 [39]
	AHG1	PIR1.2 [40]; PIR2 [40]
	AHG3/PP2CA	RGLG1 [39]; RGLG5 [39]; PIR1.2 [40]; PIR2 [40]; COP1 [38]; BPM3 [36]; BPM5 [36]
	HAI1/SAG113	PIR1.2 [40]; PIR2 [40]
	HAI3	PIR1.2 [40]; PIR2 [40]
SnRK2 kinases	SnRK2.3/SRK2I	PP2-B11 [41]
	SnRK2.6/OST1	HOS15 [42]
ABA-responsive transcription factors	ABI5	DWA1 [43]; DWA2 [43]; KEG [44]; ABD1 [45]; COP9 [46]
	ABI3	AIP2 [47,48]
	ABI4	COP1 [49]

	ABF1	KEG [50,51]
	ABF2/AREB1	KEG [52]
	ABF3/AREB2	KEG [50,51]
	DREB2	DRIP1 [53]; DRIP2 [53]; BPM2 [54]
	HB6	BTB1-6/BPM1-6 [55]
	MYB96	MIEL1 [56] (Myb96 activates ABI4)
	MYB30	RHA2b [57]; MIEL1 [56,58]
	RAV1	BPM1 [59,60]
Other regulatory and signaling factors and ABA-responding genes	ADA2B	SKIP24/At1g08710 [61]
	ATP1/SDIRIP1	AIRIP2 [62,63]; AIRP1 [62,63]; SDIR1 [64]; (SDIRIP1 activates ABI5)
	ACD11	XBAT35.2 [65]; (ACD11 is one of ABA-responding genes)
	RD21	AIRP3/LOG2 [66]; (positive role in ABA responses; RD21 is Cys protease)
	CIPK26	KEG [67,68]
	PP2A	CHIP [69]

3.1. E3 in ABA perception and ABA signaling cascade

The E3 ligases involved in controlling ABA signaling pathway, as well as their targets (Table 1), are linked together creating a large network of interactions (Figure 3). When dividing this network according to a functional group of E3 targets, it can be observed that the majority of E3s are highly selective with respect to their targets and are usually limited to a single functional group of the ABA signaling pathway. This is particularly evident in the case of SnRK2s, where only two ‘E3-target’ pairs have been identified so far, namely HOS15 ubiquitinates SnRK2.6/OST1, while PP2-B11 ubiquitinates SnRK2.3. It is not clear whether so few Ub ligases controlling SnRK2s result from paucity of experimental data or if this signaling step is weakly controlled at the level of protein ubiquitination.




Figure 3. The network of E3 ligases and their targets. E3 ligases are shown in orange, and their targets in ABA signaling pathway are colored according to target protein function. Different groups of target proteins are also shaded and labeled. Activation is shown by arrows and inhibition by T-shaped bars. Black lines show interactions between proteins, while blue lines denote transcriptional control. The references are provided in Table 1.

At least eight E3 ligases were found to target at least four ABA receptors (Figure 3). Though some of them have two or three known targets, they exclusively target ABA receptors and no other element of the ABA signaling pathway.

The interaction network becomes slightly more complex for the eight PP2Cs, having known interactions with 10 E3s (Figure 3). Although all eight PP2Cs are targeted by more than one Ub ligase and most of these E3s target more than one protein in the ABA signaling pathway, only four of these E3s interact with proteins other than PP2Cs. Three E3 ligases are common for some PP2Cs and two transcription factors involved in the ABA response.

Thus far, 11 transcription factors, involved in regulation by ABA, were found to be targeted by 17 E3 ligases (Figure 3). Among them, 13 are specific to transcription factors and they usually target only one or two proteins. Transcription factor HB6 is under the control of six BPM Ub ligases (BPM1-6). Transcription factor ABI5 is controlled by ubiquitination directly by five Ub ligases (DWA1, DWA2, ABD1, COP9, and KEG). ABI5 is also controlled at the transcriptional level by RAV1 and SDIRIP1 which are controlled by ubiquitination. The Ub ligase KEG controls four transcription factors (ABI5, ABF1, ABF2, and ABF3) and kinase CIPK26, so KEG appears to be one of the most important ligases controlling the ABA signaling pathway.

3.2. E3 in ABA-responsive transcription factors

Ub-mediated protein degradation can affect the gene expression needed to mitigate the potential negative effects of environmental stress by modulating the abundance of transcription factors. Additionally, under non-stress conditions, E3 ligases may prevent transcription by targeting the transcription factor for degradation. Transcriptional changes of ABA-responsive genes are mediated by a number of transcription factors, including members of the basic leucine zipper (bZIP), AP2/ERF, R2R3, and B3 families [10]. The abundance of many of these transcription factors is strictly guarded by numerous E3 ligases (Figure 3, Table 1).

A RING-type E3 ligase, ABI3-INTERACTING PROTEIN 2 (AIP2), was the first E3 ligase to be implicated in the regulation of ABA signaling because it mediates ABSCISIC ACID INSENSITIVE 3 (ABI3) ubiquitination and its destruction through the 26S proteasome [47,48]. The abundance of AIP2 transcripts increases in response to ABA and corresponds well with the decrease in ABI3 protein levels. The *aip2* mutant was found to have higher levels of ABI3 and be hypersensitive to ABA. The ortholog of AIP2, OsDSG1 (DELAYED SEED GERMINATION 1), has been described in rice [70]. It also physically interacts with OsABI3 and has E3 Ub ligase activity, while negatively regulating tolerance to salt and drought stress.

Although ABI4 was initially demonstrated to act as a positive regulator in the ABA signaling cascade, several studies have revealed that it is involved in the control of a variety of other cascades and crosstalk among several phytohormones [71]. ABI4 either stimulates or represses transcription of its target genes, which is consistent with its diverse biological functions. Due to the ABI4 protein's instability [72], the E3 ligases that are directly responsible for its degradation have remained elusive thus far. ABI4 together with ELONGATED HYPOCOTYL 5 (HY5) have been described to form an antagonistic module that controls seedling and chloroplast development during the transition from dark to light [49]. ABI4 promotes and HY5 inhibits the expression of CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1), the main repressor of photomorphogenesis. In turn, to ensure proper crosstalk between ABI4 and HY5 during the seedling greening process, COP1 targets ABI4 and HY5 for degradation in light and dark conditions, respectively. COP1 promotes ABI4 degradation, as evidenced by in vitro pull-down and in vivo co-immunoprecipitation assays [49].

The well-studied ABI5 is a nucleo-cytoplasmic bZIP-type transcription factor that acts downstream of ABI3. A monomeric RING-type E3 and two CUL4-RING E3 ligase (CRL) complexes have been implicated in controlling ABI5 stability and attenuating ABA signaling. CRL complexes, which are the largest class of Ub-modification enzymes in Arabidopsis, consist of a scaffold protein, one cullin protein, and an adaptor protein responsible for substrate recruitment [73]. DWD HYPERSENSITIVE TO ABA1 and ABA2

(DWA1 and DWA2, respectively) and ABA-HYPERSENSITIVE DCAF 1 (ABD1), the substrate receptors in CRLs, are able to recognize ABI5 in the nucleus to negatively regulate its stability via ubiquitination and subsequent proteasomal degradation [43,45]. Both *dwa1dwa2* double mutants and *abd1* mutants displayed ABA-hypersensitive phenotypes during seed germination and seedling growth and exhibited accumulation of ABI5 following exposure to the hormone [45]. ABI5 did not accumulate in the *dwa1dwa2* mutant in the absence of ABA, which agrees with CRL being required to regulate transcription factor abundance in the presence of the hormone. Importantly, in the absence of ABD1, proteasome-dependent degradation of ABI5 after removal of ABA is slower, implying that the E3 ligase is involved in the termination of ABA responses [45]. ABI5 is also targeted by Ub-dependent proteolysis by a RING-type E3 ligase KEEP ON GOING (KEG) [44]. There is a feedback loop between ABA signaling, KEG, and ABI5 in which KEG promotes ABI5 degradation to suppress ABA signaling, while ABA accelerates KEG self-ubiquitination and degradation, resulting in ABI5 accumulation and ABA response promotion. Because KEG is localized to the trans-Golgi network/early endosome it is believed that KEG marks ABI5 for degradation before it reaches the nucleus and activates ABA responses [51]. Additionally, another element that regulates ABI5 protein turnover is ABI FIVE BINDING PROTEIN (AFP) [74]. AFP belongs to a small family of proteins, AFP1–4, that can interact with ABI5. Though AFP is not an E3 ligase per se, it facilitates ABI5 proteasomal degradation in nuclear bodies, however the precise mechanism is not known. In rice, the AFP orthologue MEDIATOR OF OsBZIP46 DEACTIVATION AND DEGRADATION (MODD) interacts with the ABI5 orthologue OsBZIP46 to promote its ubiquitination by the E3 ligase OsPUB70, stopping ABA signaling [75]. Recently, another layer of regulation was discovered for ABI5 stability. The COP9 signalosome (CSN) regulates seed germination by facilitating ABI5 degradation [46]. CSN is a multi-subunit protein complex that may either inhibit or promote CRL complex activity. The CSN5A subunit was shown to interact with ABI5; however, the precise mechanism by which CSN5A regulates ABI5 stability will be a challenge for future studies. Finally, posttranslational modifications, such as phosphorylation and sumoylation, of ABI5 may also play a role in ABI5 proteasomal turnover [74,76,77].

In addition to ABI5, KEG also targets two other bZIP transcription factors, ABRE-BINDING FACTORS (ABF) 1 and ABF3, for degradation [50]. ABF1 and ABF3, which are also involved in seed germination and stress responses, are stabilized by ABA. Other ABFs, including ABF2 and ABF4, may also be involved in the KEG-mediated degradation pathway. The phenotype of *keg* mutants is quite severe compared to other ABA mutants, and growth arrest occurs soon after germination in the absence of the hormone [67]. This lethality of the KEG mutation might be explained by the fact that KEG targets multiple components (CIPK26, ABI5, and ABF1/3) of the ABA signaling pathway. The accumulation of its substrates may lead to misregulation and activation of other pathways independent of ABA [67].

DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN 2A (DREB2A) is a key transcription factor controlling the expression of numerous genes responsive to drought, heat, and salt [78,79]. DREB2A only accumulates in transgenics capable of attaching Ub molecules to DREB2A in non-stress conditions within the nucleus [53]. DREB2A is stable in *drip1drip2* double mutants, which present significantly higher survival rates in response to dehydration [53,80]. The situation is different under stress conditions, where the nuclear-localized A20/AN1-type zinc finger protein, STRESS ASSOCIATED PROTEIN 5 (SAP5), functions as an E3 ligase that promotes the degradation of DRIP1/2 to release and stabilize DREB2A [81]. Notably, SAP5 shows a preference for adding linear and Lys63-linked compared to Lys48-linked chains [82]. This underlines the potential role of SAP5 in selected trafficking of polyubiquitinated proteins to other 26S proteasome degradation systems, such as autophagy. DREB2A abundance increases upon exposure to heat stress. However, to attenuate the stress response, the level of the transcription factor gradually decreases over time due to the interaction with

a MATH DOMAIN-CONTAINING BTB PROTEIN (BPM), which functions as a DREB2A receptor in a Cul3 BPM1-6 RING E3 ligase complex [54]. DREB2A contains a 30-amino acid serine threonine-rich region referred to as the NEGATIVE REGULATORY DOMAIN (NRD), which is necessary for Cul3 BPM1-6 RING-dependent degradation. Biochemical data proved that the NRD domain, by itself, is capable of adequately interacting with BPM proteins. Deletion of the NRD increases DREB2A stability and carries on its constitutive action in improving tolerance to drought and heat stress [54,78]. To add further complexity, a recent study discovered that SUMO has a positive regulatory role in the action of DREB2A. The SUMOylation of Lys163, an adjacent lysine to the NRD, interferes with BPM–DREB2A interaction, stabilizing DREB2A [83]. This SUMOylation process is induced by heat shock, thus tuning the degradation rate of DREB2A in response to abiotic stress.

The same Cul3-BPM1-6 RING E3 complex is also engaged in the degradation of yet another transcription factor of ABA signaling, HOMEODOMAIN-LEUCINE ZIPPER (HD-ZIP) ATHB6 [55]. In contrast to DREB2A, ATHB6 is a negative regulator of ABA responses, including drought and heat tolerance [78,79,84]. BPM silencing, as well as ATHB6 overexpression, leads to an increase in the level of the ATHB6 protein together with the decrease of its ubiquitination status. However, overexpression of BPM6 accelerated the turnover of ATHB6, indicating that BPMs influence ATHB6 stability [55]. Furthermore, BPM silencing results in ABA insensitivity, and reduced stomatal closure, leading to increased transpiration and water loss, indicating that BPMs play a positive role in ABA signaling [55,85]. Under non-stress conditions, BPMs promote the proteasome-dependent degradation of ATHB6 [55]. This turnover of ATHB6 can be inhibited by ABA and may serve to attenuate ABA responses. Furthermore, all six BPM proteins can interact with three different HD-ZIP transcription factors (ATHB5, ATHB6, and ATHB16), implying that this family of E3 ligases may regulate other processes as well [55]. Subsequently, it was reported that BPM1 can interact with another transcription factor, RAV1 (related to ABI3/VP1), which negatively regulates the transcription of ABI3/4/5, unless it is phosphorylated by SnRK2s [59,60].

The RING-type E3 ligase MYB30-INTERACTING E3 LIGASE 1 (MIEL1) was first described to control the abundance of R2R3-type transcription factor MYB30 that negatively regulates ABA signaling and activates defense and HR responses [58,86]. MIEL1 expression is rapidly downregulated after bacterial infection to allow MYB30 accumulation, thus promoting defense and the hypersensitive response (HR), and the *miel1* mutant displays enhanced resistance responses after inoculation with bacteria [58]. In addition to its role in the control of biotic stress, MIEL1 regulates abiotic stress in an ABA-dependent manner by targeting the transcription factor MYB96, which normally promotes the expression of ABI4 [56]. In the absence of ABA, MIEL1 accumulates and promotes the degradation of MYB96 to silence ABA signal transduction, whereas in the presence of the hormone, MIEL1 is degraded by UPS, promoting MYB96 transcriptional activity. According to these findings, MIEL1 may facilitate the ABA-mediated interplay between biotic and abiotic stress [87]. As an added complication, in addition to MIEL1, another RING E3 ligase, RING-H2 FINGER PROTEIN 2B (RHA2B), is also able to bind and promote the degradation of MYB30 in an ABA-dependent manner during drought response [57].

In contrast to the E3 ligases described above and defined as negative regulators of ABA signaling, E3 ligase SALT- AND DROUGHT-INDUCED RING FINGER 1 (SDIR1) is genetically characterized as a positive regulator. The *sdir1* mutant has reduced sensitivity to ABA in seed germination and early seedling growth assays and reduced stomatal closure in response to ABA, while overexpression of SDIR1 leads to quite opposite phenotypes [64,88]. Furthermore, rice OsSDIR1 is also a functional E3 ligase, and its overexpression leads to markedly increased drought tolerance [89]. SDIR1 interacts with its substrate SDIR1-INTERACTING PROTEIN 1 (SDIRIP1) at the endoplasmic reticulum membrane to promote its degradation via the 26S proteasome. SDIRIP1 negatively regulates ABI5 expression to impact ABA-mediated seed germination and responses to

salt [64]. SDIRIP1 RNAi lines are ABA hypersensitive, suggesting that SDIRIP1 functions as a negative regulator of ABA signaling. Another RING-type E3 ligase has been described to target SDIRIP1 in the cytosol. AtAIRP2 regulated the turnover of SDIRIP1 in cell-free degradation and protoplast co-transfection assays [63]. AtAIRP2 and SDIR1 have combinatory roles in seed germination, as evidenced by reciprocal complementation of the ABA- and salt-insensitive germination phenotypes of *sdir1* and *atairp2* mutants, respectively. Recently, another link that could promote SDIRIP1 degradation was described [90]. RESPONSE TO LOW SULFUR 1–4 (LSU1–4), proteins, belonging to a small plant-specific protein family, were identified as important stress-related hubs that bind to SDIRIP1 [90]. LSUs are also able to bind to NBR1, a selective autophagy receptor; therefore, it is tempting to speculate that LSUs might serve as a bridge to facilitate autophagic degradation of SDIRIP1.

In Arabidopsis, ABA plays a major role in the drought stress response with a well-described cis-acting ABA-RESPONSIVE ELEMENT (ABRE) involved in transcriptional activation [91]. Under drought stress conditions, transcription factor ABF2 bind to the ABRE sequence of drought-responsive genes. ABF2 recruits the ADA2B-GCN5 protein complex, in which ADA2B is a transcriptional co-activator protein, while GCN5 acts as a histone acetyltransferase [92,93]. A recent study revealed that ADA2B is targeted for proteasomal degradation by the At1g08710 protein [61]. That F-box protein is localized in the nucleus and forms an SCF complex by interacting with SKP1 and cullin1 proteins. At1g08710 mutant plants present a better survival rate under water deficit conditions and a high accumulation of the ADA2B transcript following ABA treatment. Additionally, transcripts of drought-responsive genes RD22, RD29A, and ABI3 were upregulated under drought stress. These results suggest that F-box protein At1g08710 imparts drought stress tolerance through ADA2B degradation. This very recent report proves that ABF2 is an *in vitro* ubiquitination substrate of KEG [52].

UPS selectivity is accomplished not only at the ubiquitination level but also by the proteasome itself [94]. The 19S regulatory particle (RP) of the 26S proteasome confers ATP dependence, as well as substrate specificity to the holoenzyme, especially for substrates bearing a poly-Ub tag. One of the subunits of the RP in Arabidopsis is RPN10. It is essential for the degradation of ABI5 by the proteasome, as evidenced by the increased ABA sensitivity and ABI5 accumulation in the *rpn10-1* mutant. This suggests that ABI5 turnover is accomplished through the interaction between ABI5 and RPN10. It is not known whether this interaction is direct or indirect, dependent or independent of the ubiquitination status of ABI5, but it promotes its association with proteasomal proteases. This association may be prevented by the ABA signal which stabilizes ABI5 [95].

The regulation of the stability of ABA-related transcription factors is complex and multilayered. A single E3 ligase like KEG or BPM can target multiple proteins. Alternatively, a single stress-related protein can be targeted for degradation by multiple E3 ligases. As described, ABI5 abundance is regulated by at least five different E3 ligases, which are used to inhibit its activity under different circumstances [96]. Such a mechanism may rapidly integrate different inputs to control the level of a key transcription factor, leading to the appropriate response. Furthermore, a single Ub ligase can conjugate Ub to substrates in a variety of ways (e.g., Lys63- vs. Lys48-linked Ub chains), depending on the signal obtained or the identity of the substrate, leading to different outcomes. An interesting example of such regulation is rice RING E3 ligase IPA1 INTERACTING PROTEIN 1 (IPI1), which ubiquitinates IDEAL PLANT ARCHITECTURE 1 (IPA1) that is involved in plant development [97]. By adding Lys48-linked poly-Ub chains, IPI1 promotes the degradation of IPA1 in panicles, while it stabilizes IPA1 in shoot apices by adding Lys63-linked poly-Ub chains.

3.3. Other E3 ligases known to modify ABA-mediated responses

Numerous E3 ligases have been linked to ABA-regulated responses; however, in many cases, their targets have not yet been characterized. These enzymes have been categorized based on their functions as positive and negative modulators of ABA signaling (Table 2 and references therein). In many cases, it is also unclear whether they are present in the same growth conditions, cell types, organs, or developmental stages.

Table 2. E3 ligases with unidentified targets that are known to regulate ABA responses. The original references are provided next to the E3 ligase; however, the table is a compilation of the tables from several earlier reviews [6,7,15,98,99], as well as new data that were not previously reviewed.

E3 family	E3 ligase [Reference]	Influence on ABA response/ Remarks
RING-type	ATL61 [100]	Positive regulator in the ABA-mediated drought stress response
SCF	ARKP1 [101]	Positive role in ABA signaling network; mutants displaying reduced ABA-mediated inhibition of seed germination, root elongation, and water loss rate of detached leaves
RING-type	CHYR1/RZP34 [102]	Promotes ABA-induced stomatal closure, reactive oxygen species production, and plant drought tolerance; activity is regulated by SnRK2.6
RING-type	SINA2 [103]	Regulates plant responses to ABA and osmotic stress; activity is regulated by CDKG1 (cyclin dependent kinase G1)
RING-type	AtPPRT1 [104,105]	Negative role in ABA and drought stress responses
RING-type	AtARRE/ATL27 [106]	Negative regulation in ABA signaling
BTB-type	BPH1 [107]	Negatively involved in ABA-mediated cellular events; mutation caused delayed seed germination in response to ABA and resulted in hyper-induction of a large portion of ABA-inducible genes in response to ABA; mutants exhibited enhanced stomatal closure under ABA application and reduced water loss
RING-type	JUL1 [108]	ABA-mediated microtubule disorganization; regulates stomatal closure, and tolerance to drought stress
RING-type	COP1 [109]	ABA-mediated microtubule disorganization, stomatal closure
RING-type	RDUF1/2 [110]	Negatively regulates ABA signaling
RING-type	RHA2a/2b [111,112]	Positively regulate ABA signaling
RING-type	AIRP4 [113]	Positively regulate ABA signaling
SCF	TLP3/9 [114]	Positively regulate ABA signaling
RING-type	ATL43 [115]	Positively regulate ABA signaling
RING-type	CER9 [116]	Negatively regulates ABA signaling
U-box	PUB9 [117]	Negatively regulates ABA signaling
U-box	PUB18/19 [118,119]	Negatively regulates plant drought response and ABA signaling
SCF	DOR [120]	Negatively regulates ABA signaling
SCF	EDL3 [121]	Positive regulator in seed germination and root growth; positively regulates ABA signaling
SCF	MAX2 [122]	Negatively regulates plant drought stress through mediating ABA signaling; negatively regulates ABA signaling
DDB	DWA3 [123]	Negatively regulates ABA signaling; mutants exhibited ABA-hypersensitivity
U-box	PUB44/SAUL1 [124]	Negatively regulates ABA signaling
RING-type	XERICO [125]	Positively regulates ABA-dependent drought response; overexpression leads to ABA over-accumulation
SCF	FOF2 [126]	Plays an important negative role in ABA-mediated seed germination and early seedling development, as well as a positive role in ABA-mediated drought tolerance

4. Autophagy as a cellular degradation pathway and a part of intracellular trafficking

A multistep ubiquitination process resulting in the attachment of Lys48- and Lys63-linked poly-Ub tags is used to label proteins degraded by the 26S proteasome (a part of UPS) and via autophagy, respectively. UPS is the key regulator of proteostasis; however, the proteasome and autophagy complement each other in the maintenance of

appropriate protein balance in plant development and stress response (for review [5]). Several types of autophagy exist; however, macroautophagy is better described, and it will be called autophagy below. It is characterized by the presence of autophagosomes, double-membrane vesicles surrounding cargos designed for degradation and delivering it to the vacuole, after fusion of the external autophagosomal membrane with the tonoplast, (for recent reviews [3,127]). The selectivity of autophagy is achieved due to the involvement of selective autophagy cargo receptors. They recognize cargo and dock to the autophagosomes [128,129]. Autophagy, in contrast to the 26S proteasome, can degrade not only proteins and protein aggregates but also entire organelles or their parts. Therefore, it must be well controlled to avoid excessive degradation of the cellular content. It is negatively regulated by the Target of Rapamycin (TOR) kinase, which along with its effectors is active under nutrient-rich conditions when it upregulates cell growth and translation, but it is inactive during nutrient deficiency. The TOR kinase complex is modulated by diverse upstream inputs to phosphorylate different proteins in the nucleus, nucleolus, and cytosol, thus controlling transcription, the cell cycle, rRNA transcription, ribosome biogenesis, translation, and metabolism, which are pivotal to cell proliferation and growth (for review [130,131]).


Autophagy is a part of the intracellular vesicular system; therefore, extensive interlinking exists between autophagy and trafficking pathways, and functional endocytic and exocytic pathways are essential for efficient autophagy flux in all eukaryotes (for reviews [132–135]). Autophagy controls degradation, and exocytosis (secretion) is responsible for the transport of cargo to the plasma membrane or extracellular space, while endocytosis transports cargo from the plasma membrane or extracellular space to intracellular organelles (for review [136–138]). Plasma membrane receptors and transporters must be tightly regulated, and part of this control is mediated by their ubiquitination and recruitment to endosomal sorting complexes (ESCORTs) by proteins interacting with ESCORTs, of which some are capable of Ub-binding (for review [138,139]). Actin and microtubule cytoskeletons and their reorganization play an important role in the control of all three processes (for review [140,141]).

5. Role of TOR kinase, autophagy, and intracellular trafficking in the control of ABA signaling

Various elements of ABA signaling are ubiquitinated and degraded via the 26S proteasome (for review [6,7,99]). The RPN10 subunit of the 26S proteasome controls ABA signaling, as demonstrated by *rpn10-1* mutants that were more sensitive to ABA [95]. Interestingly, the same research group determined that RPN10 functions as a selective cargo receptor in proteaphagy (autophagic degradation of 26S proteasome) [142]. Data on the interplay of ABA with autophagy are still limited, but growing evidence indicates that they can co-regulate each other. For example, ABA activates autophagy by reducing the ATG4 persulfidation level. The enzymatic activity of ATG4 is reversibly regulated by sulfide, and this regulation effectively controls autophagy flux. ATG4 is necessary for processing newly synthesized ATG8 proteins that can be further lipidated to progress autophagy [143]. However, some data suggest that elevated autophagy might induce ABA responses. For example, the overexpression of banana ATG8f modulated drought stress resistance in *Arabidopsis* [144], and overexpression of NBR1 in *Arabidopsis* resulted in increased ABA signaling [145].

TOR kinase might be involved in the synthesis and/or distribution of ABA [146,147]. Additionally, the reciprocal regulation of ABA signaling and TOR kinase has been reported [148]. In growth-promoting conditions, the phosphorylation of ABA receptors by TOR disrupts their interaction with PP2C phosphatases, which can now interact with SnRK2s. This leads to the inactivation of SnRK2 kinases, thus keeping the ABA response silent. Under stress, ABA-activated SnRK2s induce the ABA response but also inactivate TOR and, in turn, activate autophagy [148]. There are additional indications for TOR-ABA connections. For instance, protein phosphatase 2A (PP2A) associated proteins

Furthermore, there is an interplay of ABA signaling with endocytosis, exocytosis, the cytoskeleton, and selective targeting of ABA-responsive transcription factors (Figure 4B). The activity of soluble and membrane-located ABA receptors is regulated by proteins linked to endocytosis and autophagy. The connection of some RCARs with the proteins interacting with ESCORT complexes, such as VACUOLAR PROTEIN SORTING23A (VPS23A) [156], FYVE domain protein required for endosomal sorting 1 (FREE1) [157,158], or ALG-2 INTERACTING PROTEIN-X (ALIX) [159], have been revealed. The turnover of ABA receptors is controlled by the targeted ubiquitination of their partners. The XBAT35 E3 ligase is a positive regulator of ABA responses since it accelerates the turnover of VPS23A through the 26S proteasome, thereby reducing vacuolar degradation of RCAR10/PYL4 [160]. In the absence of ABA, the ubiquitinated RCAR10/PYL4 and RCAR11/PYR1 are recognized by VPS23A and FREE1 and targeted for vacuolar degradation. In the presence of ABA, VPS23A and FREE1 are targeted by the SINAT E3 ligase and directed to proteasomal and vacuolar degradation [161]. The termination of ABA responses leads to autophagic clearance of the excess of SINAT-VPS23A and SINAT-FREE1 complexes and the reset of cellular homeostasis.



ABA-mediated stomatal movement is also regulated by the exocyst subunit Exo70B1 [162]. Moreover, ABA mediates microtubule disorganization, which can influence all types of vesicular trafficking [108,109]. The links via Rho GTPases/Rho Of Plants (ROP), which not only regulate numerous cytoskeleton and vesicular trafficking-based processes but also suppress ABA-induced responses, have also been revealed [163–165].

Interestingly, ABA also induces Arabidopsis multi stress regulator tryptophan-rich sensory protein-related (TSPO), which acts as a heme scavenger and binds the excessive or deleterious heme and then is targeted for degradation through autophagy [163,166].

The ABA-autophagy links also take place at the level of ABA-responsive transcription factors. Recently, an interesting possibility for the regulatory circuit between selective autophagy and ABA-responsive transcription factors was suggested by the observation that ABI3, ABI4, and ABI5 interact with selective autophagy cargo receptor NBR1 [145]. Moreover, ABA-activated SnRK2s phosphorylate FREE1, allowing for its nuclear import. In the nucleus, FREE1 interacts with the basic leucine zipper transcription factors ABA-RESPONSIVE ELEMENTS BINDING FACTOR4 (ABF4) and ABA-INSENSITIVE5 (ABI5) to reduce binding to the cis-regulatory sequences of downstream genes and diminish ABA responses [167].

Both ABA signaling and autophagy can also be involved in crosstalk related to ROS (reactive oxygen species) homeostasis. For example, autophagy affects stomatal opening by controlling ROS production in guard cells [168] and regulates glucose-mediated root meristem activity by affecting ROS homeostasis [169]. Conversely, ABI5 regulates ROS during seedling germination by CAT1 catalase activation [170], and ABI4 (acting antagonistically to ETHYLENE-INSENSITIVE3 (EIN3)) regulates ROS accumulation by modulating ascorbic acid biosynthesis [171]. TOR also controls the ROS level by regulating mitochondrial activity [172].

6. ABA crosstalk with other hormones and their possible control by autophagy

ABA is involved in extensive crosstalk with other phytohormones. Cytokinins (CKs) and brassinosteroids (BRs) regulate many aspects of plant growth antagonistically to ABA. In addition, modulation of ABA interplay with CKs and BRs is balanced by selective autophagy.

ABA-CKs crosstalk is particularly apparent in the regulatory loop consisting of type-A and -B response regulators (ARRs) and SnRK2s (for review [173]). Unlike type-B ARRs, which lack a DNA binding domain, type-A ARRs act as negative regulators of cytokinin signaling. Phosphorylation in response to cytokinin stabilizes a subset of the type-A ARRs. Plants modulate responses to CKs, at least in part, via the autophagic regulation of type-A ARRs mediated by members of the EXO70D, a subclade of the EXO70 gene family, which act as receptors to recruit type-A ARRs to the autophagosome for subsequent degradation [174]. Additionally, type-A ARRs interact with ABA-responsive transcription factor ABI5 and downregulate ABI5 transcription [175]. The other ABA-responsive transcription factor ABI4 activates ABI4 transcription. Moreover, the interaction of ABI4 and ABI5 with selective autophagy cargo receptor NBR1 with ABI4 and ABI5 suggest that both transcription factors might be degraded via autophagy in an NBR1-dependent manner [145]. A fragment of the CK signaling pathway with ABA-CKs-autophagy interplay is illustrated in Figure 5A.




Figure 5. Interplay of autophagy with ABA-cytokinin crosstalk (a) and ABA-brassinosteroid crosstalk (b). Yellow arrows mark ABA, CK, and BR signals. Elements of the CK signaling pathway (AHK, histidine kinase receptor(s); AHP, histidine phosphotransfer protein(s)) and BR signaling pathway (BRI1, leucine-rich repeat receptor-like kinase family; BSU1/BAK1, protein kinases; BIN2, GSK3-like kinase; BES1, transcription factor) are marked in panel A and B, respectively. See text for further details.

BR signaling has recently attracted attention (for reviews [17,176–179]). The cross-regulation of ABA- and BR-signaling pathways is marked in Figure 5B. BRASSINOSTEROID-INSENSITIVE 2 (BIN2) interacts with central components of the ABA signaling pathway, such as bZIP transcription factor ABI5 [180] and protein phosphatase 2C ABI1 [181], and phosphorylates SNF1-related protein kinases SnRK2.2 and SnRK2.3 [182]. Moreover, ABA promotes BR signaling via ABI1/2, which, upon binding to BIN2, leads to reduced phosphorylation of BRI1-EMS-SUPPRESSOR 1 (BES1) [183]. The links to autophagy are apparent on several levels. TOR, a negative regulator of autophagy positively regulates BR signaling via RPS6 kinase 2 (S6K2), which, in turn, phosphorylates BIN2 [184]. In addition, transcription factor BES1, an important element of BR signaling, is selectively degraded via autophagy after being targeted to autophagosomes by selective autophagy cargo receptor DSK2 [185]. Interesting crosslinks have been observed in tomato, where autophagy-related genes ATG2, ATG6, and NBR1 are induced by BES1, which results in the induction of the autophagy pathway under chilling stress [186].

7. Closing remarks

Perception and signaling of stress-triggering conditions are crucial for plant growth and development. Similarly, the control of ending these signaling events when they are no longer needed and reestablishing homeostasis is important. Ubiquitination, precisely provided by dedicated E3 ligases, is a universal mark for protein degradation. This mark is recognized by two major proteolytic pathways: UPS and autophagy. There are many unanswered questions on how plants coordinate and choose between these different proteolysis pathways to target the same or similar regulatory components. The clue might be in poly-Ub marks that have different architecture. However, what determines how particular E3 ligases structure the Ub chain is unknown.

During plant growth and development, as well as under stress, different phytohormones are mutually antagonistic and synergistic. Those phytohormones regulate autophagy, and vice-versa, autophagy impacts the balance of phytohormones. Though

the field is actively investigated, there is still a lot to discover. The future challenge is to identify phytohormone-related proteins that are specific targets of autophagy that are in line with their receptors. This knowledge would extend our understanding about stress regulation and, consequently, provide some new approaches for crop improvement, eventually leading to an increase in stress tolerance and yield.

Funding: Work in the lab is funded by the National Science Center, Poland, grant No. 2018/31/F/NZ1/02234 (SignalOAS).

Conflicts of Interest: The authors declare no conflict of interest.

References

- Orosa B, Ustun S, Calderon Villalobos LIA, Genschik P, Gibbs D, Holdsworth MJ, Isono E, Lois M, Trujillo M, Sadanandom A. Plant proteostasis - shaping the proteome: a research community aiming to understand molecular mechanisms that control protein abundance. *New Phytol* **2020**, 227, 1028-1033, doi:10.1111/nph.16664.
- Bence NF, Sampat RM, Kopito RR. Impairment of the ubiquitin-proteasome system by protein aggregation. *Science* **2001**, 292, 1552-1555, doi:10.1126/science.292.5521.1552.
- Marshall RS, Vierstra RD. Autophagy: The Master of Bulk and Selective Recycling. *Annu Rev Plant Biol* **2018**, 69, 173-208, doi:10.1146/annurev-arplant-042817-040606.
- Signorelli S, Tarkowski LP, Van den Ende W, Bassham DC. Linking Autophagy to Abiotic and Biotic Stress Responses. *Trends Plant Sci* **2019**, 24, 413-430, doi:10.1016/j.tplants.2019.02.001.
- Su T, Yang M, Wang P, Zhao Y, Ma C. Interplay between the Ubiquitin Proteasome System and Ubiquitin-Mediated Autophagy in Plants. *Cells* **2020**, 9, doi:10.3390/cells9102219.
- Doroodian P, Hua Z. The Ubiquitin Switch in Plant Stress Response. *Plants* **2021**, 10, 246.
- Stone SL. Role of the Ubiquitin Proteasome System in Plant Response to Abiotic Stress. *Int Rev Cell Mol Biol* **2019**, 343, 65-110, doi:10.1016/bs.ircmb.2018.05.012.
- Xu FQ, Xue HW. The ubiquitin-proteasome system in plant responses to environments. *Plant Cell Environ* **2019**, 42, 2931-2944, doi:10.1111/pce.13633.
- Li Y, Lin Y, Li X, Guo S, Huang Y, Xie Q. Autophagy Dances with Phytohormones upon Multiple Stresses. *Plants (Basel)* **2020**, 9, doi:10.3390/plants9081038.
- Finkelstein R. Absciscic Acid synthesis and response. *Arabidopsis Book* **2013**, 11, e0166, doi:10.1199/tab.0166.
- Yoshida T, Mogami J, Yamaguchi-Shinozaki K. Omics Approaches Toward Defining the Comprehensive Absciscic Acid Signaling Network in Plants. *Plant Cell Physiol* **2015**, 56, 1043-1052, doi:10.1093/pcp/pcv060.
- Tal L, Gil MXA, Guercio AM, Shabek N. Structural Aspects of Plant Hormone Signal Perception and Regulation by Ubiquitin Ligases. *Plant Physiol* **2020**, 182, 1537-1544, doi:10.1104/pp.19.01282.
- Yu F, Xie Q. Non-26S Proteasome Endomembrane Trafficking Pathways in ABA Signaling. *Trends Plant Sci* **2017**, 22, 976-985, doi:10.1016/j.tplants.2017.08.009.
- Yu FF, Xie Q. Ubiquitination modification precisely modulates the ABA signaling pathway in plants. *Yi Chuan* **2017**, 39, 692-706, doi:10.16288/j.ycz.17-043.
- Ali A, Pardo JM, Yun DJ. Desensitization of ABA-Signaling: The Swing From Activation to Degradation. *Front Plant Sci* **2020**, 11, 379, doi:10.3389/fpls.2020.00379.
- Chen K, Li GJ, Bressan RA, Song CP, Zhu JK, Zhao Y. Absciscic acid dynamics, signaling, and functions in plants. *J Integr Plant Biol* **2020**, 62, 25-54, doi:10.1111/jipb.12899.
- Bulgakov VP, Avramenko TV. Linking Brassinosteroid and ABA Signaling in the Context of Stress Acclimation. *International journal of molecular sciences* **2020**, 21, doi:10.3390/ijms21145108.
- Hai NN, Chuong NN, Tu NHC, Kisiala A, Hoang XLT, Thao NP. Role and Regulation of Cytokinins in Plant Response to Drought Stress. *Plants (Basel)* **2020**, 9, doi:10.3390/plants9040422.
- Verma V, Ravindran P, Kumar PP. Plant hormone-mediated regulation of stress responses. *BMC Plant Biol* **2016**, 16, 86, doi:10.1186/s12870-016-0771-y.
- Wang Q, Yu F, Xie Q. Balancing growth and adaptation to stress: Crosstalk between brassinosteroid and absciscic acid signaling. *Plant Cell Environ* **2020**, 43, 2325-2335, doi:10.1111/pce.13846.
- Fujii H, Zhu JK. Arabidopsis mutant deficient in 3 absciscic acid-activated protein kinases reveals critical roles in growth, reproduction, and stress. *Proc Natl Acad Sci U S A* **2009**, 106, 8380-8385, doi:10.1073/pnas.0903144106.
- Soon FF, Ng LM, Zhou XE, West GM, Kovach A, Tan MH, Suino-Powell KM, He Y, Xu Y, Chalmers MJ et al. Molecular mimicry regulates ABA signaling by SnRK2 kinases and PP2C phosphatases. *Science* **2012**, 335, 85-88, doi:10.1126/science.1215106.
- Dejonghe W, Okamoto M, Cutler SR. Small Molecule Probes of ABA Biosynthesis and Signaling. *Plant Cell Physiol* **2018**, 59, 1490-1499, doi:10.1093/pcp/pcy126.
- Buetow L, Huang DT. Structural insights into the catalysis and regulation of E3 ubiquitin ligases. *Nat Rev Mol Cell Biol* **2016**, 17, 626-642, doi:10.1038/nrm.2016.91.

25. French ME, Koehler CF, Hunter T. Emerging functions of branched ubiquitin chains. *Cell Discov* **2021**, 7, 6, doi:10.1038/s41421-020-00237-y.
26. Zientara-Rytter K, Subramani S. The Roles of Ubiquitin-Binding Protein Shuttles in the Degradative Fate of Ubiquitinated Proteins in the Ubiquitin-Proteasome System and Autophagy. *Cells* **2019**, 8, doi:10.3390/cells8010040.
27. Shu K, Yang W. E3 Ubiquitin Ligases: Ubiquitous Actors in Plant Development and Abiotic Stress Responses. *Plant Cell Physiol* **2017**, 58, 1461-1476, doi:10.1093/pcp/pcx071.
28. Ahn MY, Oh TR, Seo DH, Kim JH, Cho NH, Kim WT. Arabidopsis group XIV ubiquitin-conjugating enzymes AtUBC32, AtUBC33, and AtUBC34 play negative roles in drought stress response. *J Plant Physiol* **2018**, 230, 73-79, doi:10.1016/j.jplph.2018.08.010.
29. Irigoyen ML, Iniesto E, Rodriguez L, Puga MI, Yanagawa Y, Pick E, Strickland E, Paz-Ares J, Wei N, De Jaeger G et al. Targeted degradation of abscisic acid receptors is mediated by the ubiquitin ligase substrate adaptor DDA1 in Arabidopsis. *Plant Cell* **2014**, 26, 712-728, doi:10.1105/tpc.113.122234.
30. Li D, Zhang L, Li X, Kong X, Wang X, Li Y, Liu Z, Wang J, Li X, Yang Y. AtRAE1 is involved in degradation of ABA receptor RCAR1 and negatively regulates ABA signalling in Arabidopsis. *Plant Cell Environ* **2018**, 41, 231-244, doi:10.1111/pce.13086.
31. Zhao J, Zhao L, Zhang M, Zafar SA, Fang J, Li M, Zhang W, Li X. Arabidopsis E3 Ubiquitin Ligases PUB22 and PUB23 Negatively Regulate Drought Tolerance by Targeting ABA Receptor PYL9 for Degradation. *International journal of molecular sciences* **2017**, 18, doi:10.3390/ijms18091841.
32. Li Y, Zhang L, Li D, Liu Z, Wang J, Li X, Yang Y. The Arabidopsis F-box E3 ligase RIFP1 plays a negative role in abscisic acid signalling by facilitating ABA receptor RCAR3 degradation. *Plant Cell Environ* **2016**, 39, 571-582, doi:10.1111/pce.12639.
33. Bueso E, Rodriguez L, Lorenzo-Orts L, Gonzalez-Guzman M, Sayas E, Munoz-Bertomeu J, Ibanez C, Serrano R, Rodriguez PL. The single-subunit RING-type E3 ubiquitin ligase RSL1 targets PYL4 and PYR1 ABA receptors in plasma membrane to modulate abscisic acid signaling. *The Plant journal : for cell and molecular biology* **2014**, 80, 1057-1071, doi:10.1111/tjp.12708.
34. Fernandez MA, Belda-Palazon B, Julian J, Coego A, Lozano-Juste J, Inigo S, Rodriguez L, Bueso E, Goossens A, Rodriguez PL. RBR-Type E3 Ligases and the Ubiquitin-Conjugating Enzyme UBC26 Regulate Abscisic Acid Receptor Levels and Signaling. *Plant Physiol* **2020**, 182, 1723-1742, doi:10.1104/pp.19.00898.
35. Kong L, Cheng J, Zhu Y, Ding Y, Meng J, Chen Z, Xie Q, Guo Y, Li J, Yang S et al. Degradation of the ABA co-receptor ABI1 by PUB12/13 U-box E3 ligases. *Nat Commun* **2015**, 6, 8630, doi:10.1038/ncomms9630.
36. Julian J, Coego A, Lozano-Juste J, Lechner E, Wu Q, Zhang X, Merilo E, Belda-Palazon B, Park SY, Cutler SR et al. The MATH-BTB BPM3 and BPM5 subunits of Cullin3-RING E3 ubiquitin ligases target PP2CA and other clade A PP2Cs for degradation. *Proc Natl Acad Sci U S A* **2019**, 116, 15725-15734, doi:10.1073/pnas.1908677116.
37. Pan W, Lin B, Yang X, Liu L, Xia R, Li J, Wu Y, Xie Q. The UBC27-AIRP3 ubiquitination complex modulates ABA signaling by promoting the degradation of ABI1 in Arabidopsis. *Proc Natl Acad Sci U S A* **2020**, 117, 27694-27702, doi:10.1073/pnas.2007366117.
38. Chen Q, Bai L, Wang W, Shi H, Ramon Botella J, Zhan Q, Liu K, Yang HQ, Song CP. COP1 promotes ABA-induced stomatal closure by modulating the abundance of ABI/HAB and AHG3 phosphatases. *New Phytol* **2021**, 229, 2035-2049, doi:10.1111/nph.17001.
39. Wu Q, Zhang X, Peirats-Llobet M, Belda-Palazon B, Wang X, Cui S, Yu X, Rodriguez PL, An C. Ubiquitin Ligases RGLG1 and RGLG5 Regulate Abscisic Acid Signaling by Controlling the Turnover of Phosphatase PP2CA. *Plant Cell* **2016**, 28, 2178-2196, doi:10.1105/tpc.16.00364.
40. Baek W, Lim CW, Luan S, Lee SC. The RING finger E3 ligases PIR1 and PIR2 mediate PP2CA degradation to enhance abscisic acid response in Arabidopsis. *The Plant journal : for cell and molecular biology* **2019**, 100, 473-486, doi:10.1111/tjp.14507.
41. Cheng C, Wang Z, Ren Z, Zhi L, Yao B, Su C, Liu L, Li X. SCFAtPP2-B11 modulates ABA signaling by facilitating SnRK2.3 degradation in Arabidopsis thaliana. *PLoS Genet* **2017**, 13, e1006947, doi:10.1371/journal.pgen.1006947.
42. Ali A, Kim JK, Jan M, Khan HA, Khan IU, Shen M, Park J, Lim CJ, Hussain S, Baek D et al. Rheostatic Control of ABA Signaling through HOS15-Mediated OST1 Degradation. *Mol Plant* **2019**, 12, 1447-1462, doi:10.1016/j.molp.2019.08.005.
43. Lee JH, Yoon HJ, Terzaghi W, Martinez C, Dai M, Li J, Byun MO, Deng XW. DWA1 and DWA2, two Arabidopsis DWD protein components of CUL4-based E3 ligases, act together as negative regulators in ABA signal transduction. *Plant Cell* **2010**, 22, 1716-1732, doi:10.1105/tpc.109.073783.
44. Stone SL, Williams LA, Farmer LM, Vierstra RD, Callis J. KEEP ON GOING, a RING E3 ligase essential for Arabidopsis growth and development, is involved in abscisic acid signaling. *Plant Cell* **2006**, 18, 3415-3428, doi:10.1105/tpc.106.046532.
45. Seo KI, Lee JH, Nezames CD, Zhong S, Song E, Byun MO, Deng XW. ABD1 is an Arabidopsis DCAF substrate receptor for CUL4-DBB1-based E3 ligases that acts as a negative regulator of abscisic acid signaling. *Plant Cell* **2014**, 26, 695-711, doi:10.1105/tpc.113.119974.
46. Jin D, Wu M, Li B, Bucker B, Keil P, Zhang S, Li J, Kang D, Liu J, Dong J et al. The COP9 Signalosome regulates seed germination by facilitating protein degradation of RGL2 and ABI5. *PLoS Genet* **2018**, 14, e1007237, doi:10.1371/journal.pgen.1007237.
47. Kurup S, Jones HD, Holdsworth MJ. Interactions of the developmental regulator ABI3 with proteins identified from developing Arabidopsis seeds. *The Plant journal : for cell and molecular biology* **2000**, 21, 143-155, doi:10.1046/j.1365-313x.2000.00663.x.
48. Zhang X, Garreton V, Chua NH. The AIP2 E3 ligase acts as a novel negative regulator of ABA signaling by promoting ABI3 degradation. *Genes Dev* **2005**, 19, 1532-1543, doi:10.1101/gad.1318705.

49. Xu X, Chi W, Sun X, Feng P, Guo H, Li J, Lin R, Lu C, Wang H, Leister D et al. Convergence of light and chloroplast signals for de-etiolation through ABI4-HY5 and COP1. *Nat Plants* **2016**, 2, 16066, doi:10.1038/nplants.2016.66.
50. Chen YT, Liu H, Stone S, Callis J. ABA and the ubiquitin E3 ligase KEEP ON GOING affect proteolysis of the Arabidopsis thaliana transcription factors ABF1 and ABF3. *The Plant journal : for cell and molecular biology* **2013**, 75, 965-976, doi:10.1111/tpj.12259.
51. Liu H, Stone SL. Cytoplasmic degradation of the Arabidopsis transcription factor abscisic acid insensitive 5 is mediated by the RING-type E3 ligase KEEP ON GOING. *J Biol Chem* **2013**, 288, 20267-20279, doi:10.1074/jbc.M113.465369.
52. Linden KJ, Chen Y-T, Kyaw K, Schultz B, Callis J. Factors that affect protein abundance of the bZIP transcription factor ABRE-BINDING FACTOR 2 (ABF2), a positive regulator of abscisic acid signaling. *bioRxiv* **2021**, 10.1101/2021.01.25.428157, 2021.2001.2025.428157, doi:10.1101/2021.01.25.428157.
53. Qin F, Sakuma Y, Tran LS, Maruyama K, Kidokoro S, Fujita Y, Fujita M, Umezawa T, Sawano Y, Miyazono K et al. Arabidopsis DREB2A-interacting proteins function as RING E3 ligases and negatively regulate plant drought stress-responsive gene expression. *Plant Cell* **2008**, 20, 1693-1707, doi:10.1105/tpc.107.057380.
54. Morimoto K, Ohama N, Kidokoro S, Mizoi J, Takahashi F, Todaka D, Mogami J, Sato H, Qin F, Kim JS et al. BPM-CUL3 E3 ligase modulates thermotolerance by facilitating negative regulatory domain-mediated degradation of DREB2A in Arabidopsis. *Proc Natl Acad Sci U S A* **2017**, 114, E8528-E8536, doi:10.1073/pnas.1704189114.
55. Lechner E, Leonhardt N, Eisler H, Parmentier Y, Alioua M, Jacquet H, Leung J, Genschik P. MATH/BTB CRL3 receptors target the homeodomain-leucine zipper ATHB6 to modulate abscisic acid signaling. *Dev Cell* **2011**, 21, 1116-1128, doi:10.1016/j.devcel.2011.10.018.
56. Lee HG, Seo PJ. The Arabidopsis MIEL1 E3 ligase negatively regulates ABA signalling by promoting protein turnover of MYB96. *Nat Commun* **2016**, 7, 12525, doi:10.1038/ncomms12525.
57. Zheng Y, Chen Z, Ma L, Liao C. The Ubiquitin E3 Ligase RHA2b Promotes Degradation of MYB30 in Abscisic Acid Signaling. *Plant Physiol* **2018**, 178, 428-440, doi:10.1104/pp.18.00683.
58. Marino D, Froidure S, Canonne J, Ben Khaled S, Khafif M, Pouzet C, Jauneau A, Roby D, Rivas S. Arabidopsis ubiquitin ligase MIEL1 mediates degradation of the transcription factor MYB30 weakening plant defence. *Nat Commun* **2013**, 4, 1476, doi:10.1038/ncomms2479.
59. Chen L, Lee JH, Weber H, Tohge T, Witt S, Roje S, Fernie AR, Hellmann H. Arabidopsis BPM proteins function as substrate adaptors to a cullin3-based E3 ligase to affect fatty acid metabolism in plants. *Plant Cell* **2013**, 25, 2253-2264, doi:10.1105/tpc.112.107292.
60. Feng CZ, Chen Y, Wang C, Kong YH, Wu WH, Chen YF. Arabidopsis RAV1 transcription factor, phosphorylated by SnRK2 kinases, regulates the expression of ABI3, ABI4, and ABI5 during seed germination and early seedling development. *The Plant journal : for cell and molecular biology* **2014**, 80, 654-668, doi:10.1111/tpj.12670.
61. Rao V, Virupapuram V. Arabidopsis F-box protein At1g08710 interacts with transcriptional protein ADA2b and imparts drought stress tolerance by negatively regulating seedling growth. *Biochem Biophys Res Commun* **2021**, 536, 45-51, doi:10.1016/j.bbrc.2020.12.054.
62. Cho SK, Ryu MY, Seo DH, Kang BG, Kim WT. The Arabidopsis RING E3 ubiquitin ligase AtAIRP2 plays combinatory roles with AtAIRP1 in abscisic acid-mediated drought stress responses. *Plant Physiol* **2011**, 157, 2240-2257, doi:10.1104/pp.111.185595.
63. Oh TR, Kim JH, Cho SK, Ryu MY, Yang SW, Kim WT. AtAIRP2 E3 Ligase Affects ABA and High-Salinity Responses by Stimulating Its ATP1/SDIRIP1 Substrate Turnover. *Plant Physiol* **2017**, 174, 2515-2531, doi:10.1104/pp.17.00467.
64. Zhang H, Cui F, Wu Y, Lou L, Liu L, Tian M, Ning Y, Shu K, Tang S, Xie Q. The RING finger ubiquitin E3 ligase SDIR1 targets SDIR1-INTERACTING PROTEIN1 for degradation to modulate the salt stress response and ABA signaling in Arabidopsis. *Plant Cell* **2015**, 27, 214-227, doi:10.1105/tpc.114.134163.
65. Li Q, Serio RJ, Schofield A, Liu H, Rasmussen SR, Hofius D, Stone SL. Arabidopsis RING-type E3 ubiquitin ligase XBAT35.2 promotes proteasome-dependent degradation of ACD11 to attenuate abiotic stress tolerance. *The Plant journal : for cell and molecular biology* **2020**, 104, 1712-1723, doi:10.1111/tpj.15032.
66. Kim JH, Kim WT. The Arabidopsis RING E3 ubiquitin ligase AtAIRP3/LOG2 participates in positive regulation of high-salt and drought stress responses. *Plant Physiol* **2013**, 162, 1733-1749, doi:10.1104/pp.113.220103.
67. Lyzenga WJ, Liu H, Schofield A, Muike-Hennessey A, Stone SL. Arabidopsis CIPK26 interacts with KEG, components of the ABA signalling network and is degraded by the ubiquitin-proteasome system. *J Exp Bot* **2013**, 64, 2779-2791, doi:10.1093/jxb/ert123.
68. Lyzenga WJ, Sullivan V, Liu H, Stone SL. The Kinase Activity of Calcineurin B-like Interacting Protein Kinase 26 (CIPK26) Influences Its Own Stability and that of the ABA-regulated Ubiquitin Ligase, Keep on Going (KEG). *Front Plant Sci* **2017**, 8, 502, doi:10.3389/fpls.2017.00502.
69. Luo J, Shen G, Yan J, He C, Zhang H. AtCHIP functions as an E3 ubiquitin ligase of protein phosphatase 2A subunits and alters plant response to abscisic acid treatment. *The Plant journal : for cell and molecular biology* **2006**, 46, 649-657, doi:10.1111/j.1365-3113X.2006.02730.x.
70. Park GG, Park JJ, Yoon J, Yu SN, An G. A RING finger E3 ligase gene, Oryza sativa Delayed Seed Germination 1 (OsDSG1), controls seed germination and stress responses in rice. *Plant Mol Biol* **2010**, 74, 467-478, doi:10.1007/s11103-010-9687-3.
71. Chandrasekaran U, Luo X, Zhou W, Shu K. Multifaceted Signaling Networks Mediated by Abscisic Acid Insensitive 4. *Plant Commun* **2020**, 1, 100040, doi:10.1016/j.xplc.2020.100040.

72. Finkelstein R, Lynch T, Reeves W, Petitfils M, Mostachetti M. Accumulation of the transcription factor ABA-insensitive (ABI)4 is tightly regulated post-transcriptionally. *J Exp Bot* **2011**, 62, 3971-3979, doi:10.1093/jxb/err093.
73. Hua Z, Vierstra RD. The cullin-RING ubiquitin-protein ligases. *Annu Rev Plant Biol* **2011**, 62, 299-334, doi:10.1146/annurev-arplant-042809-112256.
74. Lopez-Molina L, Mongrand S, Kinoshita N, Chua NH. AFP is a novel negative regulator of ABA signaling that promotes ABI5 protein degradation. *Genes Dev* **2003**, 17, 410-418, doi:10.1101/gad.1055803.
75. Tang N, Ma S, Zong W, Yang N, Lv Y, Yan C, Guo Z, Li J, Li X, Xiang Y et al. MODD Mediates Deactivation and Degradation of OsZIP46 to Negatively Regulate ABA Signaling and Drought Resistance in Rice. *Plant Cell* **2016**, 28, 2161-2177, doi:10.1105/tpc.16.00171.
76. Lopez-Molina L, Mongrand S, Chua NH. A postgermination developmental arrest checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in Arabidopsis. *Proc Natl Acad Sci U S A* **2001**, 98, 4782-4787, doi:10.1073/pnas.081594298.
77. Miura K, Lee J, Jin JB, Yoo CY, Miura T, Hasegawa PM. Sumoylation of ABI5 by the Arabidopsis SUMO E3 ligase SIZ1 negatively regulates abscisic acid signaling. *Proc Natl Acad Sci U S A* **2009**, 106, 5418-5423, doi:10.1073/pnas.0811088106.
78. Sakuma Y, Maruyama K, Osakabe Y, Qin F, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. Functional analysis of an Arabidopsis transcription factor, DREB2A, involved in drought-responsive gene expression. *Plant Cell* **2006**, 18, 1292-1309, doi:10.1105/tpc.105.035881.
79. Sakuma Y, Maruyama K, Qin F, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K. Dual function of an Arabidopsis transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression. *Proc Natl Acad Sci U S A* **2006**, 103, 18822-18827, doi:10.1073/pnas.0605639103.
80. Morimoto K, Mizoi J, Qin F, Kim JS, Sato H, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K. Stabilization of Arabidopsis DREB2A is required but not sufficient for the induction of target genes under conditions of stress. *PLoS One* **2013**, 8, e80457, doi:10.1371/journal.pone.0080457.
81. Zhang N, Yin Y, Liu X, Tong S, Xing J, Zhang Y, Pudake RN, Izquierdo EM, Peng H, Xin M et al. The E3 Ligase TaSAP5 Alters Drought Stress Responses by Promoting the Degradation of DRIP Proteins. *Plant Physiol* **2017**, 175, 1878-1892, doi:10.1104/pp.17.01319.
82. Choi H, Han S, Shin D, Lee S. Polyubiquitin recognition by AtSAP5, an A20-type zinc finger containing protein from Arabidopsis thaliana. *Biochem Biophys Res Commun* **2012**, 419, 436-440, doi:10.1016/j.bbrc.2012.02.044.
83. Wang F, Liu Y, Shi Y, Han D, Wu Y, Ye W, Yang H, Li G, Cui F, Wan S et al. SUMOylation Stabilizes the Transcription Factor DREB2A to Improve Plant Thermotolerance. *Plant Physiol* **2020**, 183, 41-50, doi:10.1104/pp.20.00080.
84. Himmelbach A, Hoffmann T, Leube M, Hohener B, Grill E. Homeodomain protein ATHB6 is a target of the protein phosphatase ABI1 and regulates hormone responses in Arabidopsis. *EMBO J* **2002**, 21, 3029-3038, doi:10.1093/emboj/cdf316.
85. Johannesson H, Wang Y, Hanson J, Engstrom P. The Arabidopsis thaliana homeobox gene ATHB5 is a potential regulator of abscisic acid responsiveness in developing seedlings. *Plant Mol Biol* **2003**, 51, 719-729, doi:10.1023/a:1022567625228.
86. Zheng Y, Schumaker KS, Guo Y. Sumoylation of transcription factor MYB30 by the small ubiquitin-like modifier E3 ligase SIZ1 mediates abscisic acid response in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* **2012**, 109, 12822-12827, doi:10.1073/pnas.1202630109.
87. Lee SC, Luan S. ABA signal transduction at the crossroad of biotic and abiotic stress responses. *Plant Cell Environ* **2012**, 35, 53-60, doi:10.1111/j.1365-3040.2011.02426.x.
88. Zhang Y, Yang C, Li Y, Zheng N, Chen H, Zhao Q, Gao T, Guo H, Xie Q. SDIR1 is a RING finger E3 ligase that positively regulates stress-responsive abscisic acid signaling in Arabidopsis. *Plant Cell* **2007**, 19, 1912-1929, doi:10.1105/tpc.106.048488.
89. Gao T, Wu Y, Zhang Y, Liu L, Ning Y, Wang D, Tong H, Chen S, Chu C, Xie Q. OsSDIR1 overexpression greatly improves drought tolerance in transgenic rice. *Plant Mol Biol* **2011**, 76, 145-156, doi:10.1007/s11103-011-9775-z.
90. Niemiro A, Cysewski D, Brzywczy J, Wawrzynska A, Sienko M, Poznanski J, Sirko A. Similar but Not Identical-Binding Properties of LSU (Response to Low Sulfur) Proteins From Arabidopsis thaliana. *Front Plant Sci* **2020**, 11, 1246, doi:10.3389/fpls.2020.01246.
91. Gomez-Porras JL, Riano-Pachon DM, Dreyer I, Mayer JE, Mueller-Roeber B. Genome-wide analysis of ABA-responsive elements ABRE and CE3 reveals divergent patterns in Arabidopsis and rice. *BMC Genomics* **2007**, 8, 260, doi:10.1186/1471-2164-8-260.
92. Stockinger EJ, Mao Y, Regier MK, Triezenberg SJ, Thomashow MF. Transcriptional adaptor and histone acetyltransferase proteins in Arabidopsis and their interactions with CBF1, a transcriptional activator involved in cold-regulated gene expression. *Nucleic Acids Res* **2001**, 29, 1524-1533, doi:10.1093/nar/29.7.1524.
93. Vlachonassios KE, Thomashow MF, Triezenberg SJ. Disruption mutations of ADA2b and GCN5 transcriptional adaptor genes dramatically affect Arabidopsis growth, development, and gene expression. *Plant Cell* **2003**, 15, 626-638, doi:10.1105/tpc.007922.
94. Smalle J, Vierstra RD. The ubiquitin 26S proteasome proteolytic pathway. *Annu Rev Plant Biol* **2004**, 55, 555-590, doi:10.1146/annurev-arplant.55.031903.141801.
95. Smalle J, Kurepa J, Yang P, Emborg TJ, Babiychuk E, Kushnir S, Vierstra RD. The pleiotropic role of the 26S proteasome subunit RPN10 in Arabidopsis growth and development supports a substrate-specific function in abscisic acid signaling. *Plant Cell* **2003**, 15, 965-980, doi:10.1105/tpc.009217.

96. Liu H, Stone SL. Regulation of ABI5 turnover by reversible post-translational modifications. *Plant Signal Behav* **2014**, 9, e27577, doi:10.4161/psb.27577.
97. Wang J, Yu H, Xiong G, Lu Z, Jiao Y, Meng X, Liu G, Chen X, Wang Y, Li J. Tissue-Specific Ubiquitination by IPA1 INTERACTING PROTEIN1 Modulates IPA1 Protein Levels to Regulate Plant Architecture in Rice. *Plant Cell* **2017**, 29, 697-707, doi:10.1105/tpc.16.00879.
98. Yang W, Zhang W, Wang X. Post-translational control of ABA signalling: the roles of protein phosphorylation and ubiquitination. *Plant Biotechnol J* **2017**, 15, 4-14, doi:10.1111/pbi.12652.
99. Yu F, Wu Y, Xie Q. Ubiquitin-Proteasome System in ABA Signaling: From Perception to Action. *Mol Plant* **2016**, 9, 21-33, doi:10.1016/j.molp.2015.09.015.
100. Yang R, Wang T, Shi W, Li S, Liu Z, Wang J, Yang Y. E3 ubiquitin ligase ATL61 acts as a positive regulator in abscisic acid mediated drought response in Arabidopsis. *Biochem Biophys Res Commun* **2020**, 528, 292-298, doi:10.1016/j.bbrc.2020.05.067.
101. Li Y, Liu Z, Wang J, Li X, Yang Y. The Arabidopsis Kelch Repeat F-box E3 Ligase ARKP1 Plays a Positive Role for the Regulation of Abscisic Acid Signaling. *Plant Mol Biol Rep* **2016**, 34, 582-591.
102. Ding S, Zhang B, Qin F. Arabidopsis RZFP34/CHYR1, a Ubiquitin E3 Ligase, Regulates Stomatal Movement and Drought Tolerance via SnRK2.6-Mediated Phosphorylation. *Plant Cell* **2015**, 27, 3228-3244, doi:10.1105/tpc.15.00321.
103. Chen Y, Fokar M, Kang M, Chen N, Allen RD, Chen Y. Phosphorylation of Arabidopsis SINA2 by CDKG1 affects its ubiquitin ligase activity. *BMC Plant Biol* **2018**, 18, 147, doi:10.1186/s12870-018-1364-8.
104. Liu Y, Pei L, Xiao S, Peng L, Liu Z, Li X, Yang Y, Wang J. AtPPRT1 negatively regulates salt stress response in Arabidopsis seedlings. *Plant Signal Behav* **2020**, 15, 1732103, doi:10.1080/15592324.2020.1732103.
105. Pei L, Peng L, Wan X, Xiong J, Liu Z, Li X, Yang Y, Wang J. Expression Pattern and Function Analysis of AtPPRT1, a Novel Negative Regulator in ABA and Drought Stress Responses in Arabidopsis. *International journal of molecular sciences* **2019**, 20, doi:10.3390/ijms20020394.
106. Wang B, Li C, Kong X, Li Y, Liu Z, Wang J, Li X, Yang Y. AtARRE, an E3 ubiquitin ligase, negatively regulates ABA signaling in Arabidopsis thaliana. *Plant Cell Rep* **2018**, 37, 1269-1278, doi:10.1007/s00299-018-2311-8.
107. Woo OG, Kim SH, Cho SK, Kim SH, Lee HN, Chung T, Yang SW, Lee JH. BPH1, a novel substrate receptor of CRL3, plays a repressive role in ABA signal transduction. *Plant Mol Biol* **2018**, 96, 593-606, doi:10.1007/s11103-018-0717-x.
108. Yu SG, Kim JH, Cho NH, Oh TR, Kim WT. Arabidopsis RING E3 ubiquitin ligase JUL1 participates in ABA-mediated microtubule depolymerization, stomatal closure, and tolerance response to drought stress. *The Plant journal : for cell and molecular biology* **2020**, 103, 824-842, doi:10.1111/tpj.14775.
109. Khanna R, Li J, Tseng TS, Schroeder JI, Ehrhardt DW, Briggs WR. COP1 jointly modulates cytoskeletal processes and electrophysiological responses required for stomatal closure. *Mol Plant* **2014**, 7, 1441-1454, doi:10.1093/mp/ssu065.
110. Kim SJ, Ryu MY, Kim WT. Suppression of Arabidopsis RING-DUF1117 E3 ubiquitin ligases, AtRDUF1 and AtRDUF2, reduces tolerance to ABA-mediated drought stress. *Biochem Biophys Res Commun* **2012**, 420, 141-147, doi:10.1016/j.bbrc.2012.02.131.
111. Bu Q, Li H, Zhao Q, Jiang H, Zhai Q, Zhang J, Wu X, Sun J, Xie Q, Wang D et al. The Arabidopsis RING finger E3 ligase RHA2a is a novel positive regulator of abscisic acid signaling during seed germination and early seedling development. *Plant Physiol* **2009**, 150, 463-481, doi:10.1104/pp.109.135269.
112. Li H, Jiang H, Bu Q, Zhao Q, Sun J, Xie Q, Li C. The Arabidopsis RING finger E3 ligase RHA2b acts additively with RHA2a in regulating abscisic acid signaling and drought response. *Plant Physiol* **2011**, 156, 550-563, doi:10.1104/pp.111.176214.
113. Yang L, Liu Q, Liu Z, Yang H, Wang J, Li X, Yang Y. Arabidopsis C3HC4-RING finger E3 ubiquitin ligase AtAIRP4 positively regulates stress-responsive abscisic acid signaling. *J Integr Plant Biol* **2016**, 58, 67-80, doi:10.1111/jipb.12364.
114. Bao Y, Song WM, Jin YL, Jiang CM, Yang Y, Li B, Huang WJ, Liu H, Zhang HX. Characterization of Arabidopsis Tubby-like proteins and redundant function of AtTLP3 and AtTLP9 in plant response to ABA and osmotic stress. *Plant Mol Biol* **2014**, 86, 471-483, doi:10.1007/s11103-014-0241-6.
115. Serrano M, Parra S, Alcaraz LD, Guzman P. The ATL gene family from Arabidopsis thaliana and Oryza sativa comprises a large number of putative ubiquitin ligases of the RING-H2 type. *J Mol Evol* **2006**, 62, 434-445, doi:10.1007/s00239-005-0038-y.
116. Zhao H, Zhang H, Cui P, Ding F, Wang G, Li R, Jenks MA, Lu S, Xiong L. The Putative E3 Ubiquitin Ligase ECERIFERUM9 Regulates Abscisic Acid Biosynthesis and Response during Seed Germination and Postgermination Growth in Arabidopsis. *Plant Physiol* **2014**, 165, 1255-1268, doi:10.1104/pp.114.239699.
117. Samuel MA, Mudgil Y, Salt JN, Delmas F, Ramachandran S, Chilelli A, Goring DR. Interactions between the S-domain receptor kinases and AtPUB-ARM E3 ubiquitin ligases suggest a conserved signaling pathway in Arabidopsis. *Plant Physiol* **2008**, 147, 2084-2095, doi:10.1104/pp.108.123380.
118. Bergler J, Hoth S. Plant U-box armadillo repeat proteins AtPUB18 and AtPUB19 are involved in salt inhibition of germination in Arabidopsis. *Plant Biol (Stuttg)* **2011**, 13, 725-730, doi:10.1111/j.1438-8677.2010.00431.x.
119. Liu YC, Wu YR, Huang XH, Sun J, Xie Q. AtPUB19, a U-box E3 ubiquitin ligase, negatively regulates abscisic acid and drought responses in Arabidopsis thaliana. *Mol Plant* **2011**, 4, 938-946, doi:10.1093/mp/ssr030.
120. Zhang Y, Xu W, Li Z, Deng XW, Wu W, Xue Y. F-box protein DOR functions as a novel inhibitory factor for abscisic acid-induced stomatal closure under drought stress in Arabidopsis. *Plant Physiol* **2008**, 148, 2121-2133, doi:10.1104/pp.108.126912.
121. Koops P, Pelser S, Ignatz M, Klose C, Marrocco-Selden K, Kretsch T. EDL3 is an F-box protein involved in the regulation of abscisic acid signalling in Arabidopsis thaliana. *J Exp Bot* **2011**, 62, 5547-5560, doi:10.1093/jxb/err236.

122. Bu Q, Lv T, Shen H, Luong P, Wang J, Wang Z, Huang Z, Xiao L, Engineer C, Kim TH et al. Regulation of drought tolerance by the F-box protein MAX2 in Arabidopsis. *Plant Physiol* **2014**, *164*, 424-439, doi:10.1104/pp.113.226837.
123. Lee JH, Terzaghi W, Deng XW. DWA3, an Arabidopsis DWD protein, acts as a negative regulator in ABA signal transduction. *Plant Sci* **2011**, *180*, 352-357, doi:10.1016/j.plantsci.2010.10.008.
124. Salt JN, Yoshioka K, Moeder W, Goring DR. Altered germination and subcellular localization patterns for PUB44/SAUL1 in response to stress and phytohormone treatments. *PLoS One* **2011**, *6*, e21321, doi:10.1371/journal.pone.0021321.
125. Ko JH, Yang SH, Han KH. Upregulation of an Arabidopsis RING-H2 gene, XERICO, confers drought tolerance through increased abscisic acid biosynthesis. *The Plant journal : for cell and molecular biology* **2006**, *47*, 343-355, doi:10.1111/j.1365-3113X.2006.02782.x.
126. Qu L, Sun M, Li X, He R, Zhong M, Luo D, Liu X, Zhao X. The Arabidopsis F-box protein FOF2 regulates ABA-mediated seed germination and drought tolerance. *Plant Sci* **2020**, *301*, 110643, doi:10.1016/j.plantsci.2020.110643.
127. Su T, Li X, Yang M, Shao Q, Zhao Y, Ma C, Wang P. Autophagy: An Intracellular Degradation Pathway Regulating Plant Survival and Stress Response. *Front Plant Sci* **2020**, *11*, 164, doi:10.3389/fpls.2020.00164.
128. Lin Y, Guo R, Ji C, Zhou J, Jiang L. New insights into AtNBR1 as a selective autophagy cargo receptor in Arabidopsis. *Plant Signal Behav* **2021**, *16*, 1839226, doi:10.1080/15592324.2020.1839226.
129. Luo S, Li X, Zhang Y, Fu Y, Fan B, Zhu C, Chen Z. Cargo Recognition and Function of Selective Autophagy Receptors in Plants. *International journal of molecular sciences* **2021**, *22*, doi:10.3390/ijms22031013.
130. Ingargiola C, Turqueto Duarte G, Robaglia C, Leprince AS, Meyer C. The Plant Target of Rapamycin: A Conduc TOR of Nutrition and Metabolism in Photosynthetic Organisms. *Genes (Basel)* **2020**, *11*, doi:10.3390/genes11111285.
131. Xiong Y, Sheen J. Novel links in the plant TOR kinase signaling network. *Curr Opin Plant Biol* **2015**, *28*, 83-91, doi:10.1016/j.pbi.2015.09.006.
132. Papandreou ME, Tavernarakis N. Crosstalk between Endo/Exocytosis and Autophagy in Health and Disease. *Biotechnol J* **2020**, *15*, e1900267, doi:10.1002/biot.201900267.
133. Pecenkova T, Markovic V, Sabol P, Kulich I, Zarsky V. Exocyst and autophagy-related membrane trafficking in plants. *J Exp Bot* **2017**, *69*, 47-57, doi:10.1093/jxb/erx363.
134. Saeed B, Brillada C, Trujillo M. Dissecting the plant exocyst. *Curr Opin Plant Biol* **2019**, *52*, 69-76, doi:10.1016/j.pbi.2019.08.004.
135. Zientara-Rytter K, Sirko A. To deliver or to degrade - an interplay of the ubiquitin-proteasome system, autophagy and vesicular transport in plants. *FEBS J* **2016**, *283*, 3534-3555, doi:10.1111/febs.13712.
136. Elliott L, Moore I, Kirchhelle C. Spatio-temporal control of post-Golgi exocytic trafficking in plants. *J Cell Sci* **2020**, *133*, doi:10.1242/jcs.237065.
137. Kanazawa T, Ueda T. Exocytic trafficking pathways in plants: why and how they are redirected. *New Phytol* **2017**, *215*, 952-957, doi:10.1111/nph.14613.
138. Mosesso N, Nagel MK, Isono E. Ubiquitin recognition in endocytic trafficking - with or without ESCRT-0. *J Cell Sci* **2019**, *132*, doi:10.1242/jcs.232868.
139. Gao C, Zhuang X, Shen J, Jiang L. Plant ESCRT Complexes: Moving Beyond Endosomal Sorting. *Trends Plant Sci* **2017**, *22*, 986-998, doi:10.1016/j.tplants.2017.08.003.
140. Geitmann A, Nebenfuhr A. Navigating the plant cell: intracellular transport logistics in the green kingdom. *Mol Biol Cell* **2015**, *26*, 3373-3378, doi:10.1091/mbc.E14-10-1482.
141. Koppers M, Ozkan N, Farias GG. Complex Interactions Between Membrane-Bound Organelles, Biomolecular Condensates and the Cytoskeleton. *Front Cell Dev Biol* **2020**, *8*, 618733, doi:10.3389/fcell.2020.618733.
142. Marshall RS, Li F, Gemperline DC, Book AJ, Vierstra RD. Autophagic Degradation of the 26S Proteasome Is Mediated by the Dual ATG8/Ubiquitin Receptor RPN10 in Arabidopsis. *Mol Cell* **2015**, *58*, 1053-1066, doi:10.1016/j.molcel.2015.04.023.
143. Laureano-Marin AM, Aroca A, Perez-Perez ME, Yruela I, Jurado-Flores A, Moreno I, Crespo JL, Romero LC, Gotor C. Abscissic Acid-Triggered Persulfidation of the Cys Protease ATG4 Mediates Regulation of Autophagy by Sulfide. *Plant Cell* **2020**, *32*, 3902-3920, doi:10.1105/tpc.20.00766.
144. Li B, Liu G, Wang Y, Wei Y, Shi H. Overexpression of Banana ATG8f Modulates Drought Stress Resistance in Arabidopsis. *Biomolecules* **2019**, *9*, doi:10.3390/biom9120814.
145. Tarnowski L, Rodriguez MC, Brzywczy J, Piecho-Kabacik M, Krckova Z, Martinec J, Wawrzynska A, Sirko A. A selective autophagy cargo receptor NBR1 modulates abscisic acid signalling in Arabidopsis thaliana. *Sci Rep* **2020**, *10*, 7778, doi:10.1038/s41598-020-64765-z.
146. Kravchenko A, Citerne S, Jehanno I, Bersimbaev RI, Veit B, Meyer C, Leprince AS. Mutations in the Arabidopsis Lst8 and Raptor genes encoding partners of the TOR complex, or inhibition of TOR activity decrease abscisic acid (ABA) synthesis. *Biochem Biophys Res Commun* **2015**, *467*, 992-997, doi:10.1016/j.bbrc.2015.10.028.
147. Salem MA, Li Y, Wiszniewski A, Giavalisco P. Regulatory-associated protein of TOR (RAPTOR) alters the hormonal and metabolic composition of Arabidopsis seeds, controlling seed morphology, viability and germination potential. *The Plant journal : for cell and molecular biology* **2017**, *92*, 525-545, doi:10.1111/tpj.13667.
148. Wang P, Zhao Y, Li Z, Hsu CC, Liu X, Fu L, Hou YJ, Du Y, Xie S, Zhang C et al. Reciprocal Regulation of the TOR Kinase and ABA Receptor Balances Plant Growth and Stress Response. *Mol Cell* **2018**, *69*, 100-112 e106, doi:10.1016/j.molcel.2017.12.002.

149. Hu R, Zhu Y, Shen G, Zhang H. TAP46 plays a positive role in the ABSCISIC ACID INSENSITIVE5-regulated gene expression in Arabidopsis. *Plant Physiol* **2014**, *164*, 721-734, doi:10.1104/pp.113.233684.
150. Punzo P, Ruggiero A, Grillo S, Batelli G. TIP41 network analysis and mutant phenotypes predict interactions between the TOR and ABA pathways. *Plant Signal Behav* **2018**, *13*, e1537698, doi:10.1080/15592324.2018.1537698.
151. Punzo P, Ruggiero A, Possenti M, Nurcato R, Costa A, Morelli G, Grillo S, Batelli G. The PP2A-interactor TIP41 modulates ABA responses in Arabidopsis thaliana. *The Plant journal : for cell and molecular biology* **2018**, *94*, 991-1009, doi:10.1111/tbj.13913.
152. Waadt R, Manalansan B, Rauniyar N, Munemasa S, Booker MA, Brandt B, Waadt C, Nusinow DA, Kay SA, Kunz HH et al. Identification of Open Stomata1-Interacting Proteins Reveals Interactions with Sucrose Non-fermenting1-Related Protein Kinases2 and with Type 2A Protein Phosphatases That Function in Absciscic Acid Responses. *Plant Physiol* **2015**, *169*, 760-779, doi:10.1104/pp.15.00575.
153. Li L, Song Y, Wang K, Dong P, Zhang X, Li F, Li Z, Ren M. TOR-inhibitor insensitive-1 (TRIN1) regulates cotyledons greening in Arabidopsis. *Front Plant Sci* **2015**, *6*, 861, doi:10.3389/fpls.2015.00861.
154. Barrada A, Djendli M, Desnos T, Mercier R, Robaglia C, Montane MH, Menand B. A TOR-YAK1 signaling axis controls cell cycle, meristem activity and plant growth in Arabidopsis. *Development* **2019**, *146*, doi:10.1242/dev.171298.
155. Kim D, Ntui VO, Xiong L. Arabidopsis YAK1 regulates absciscic acid response and drought resistance. *FEBS Lett* **2016**, *590*, 2201-2209, doi:10.1002/1873-3468.12234.
156. Yu F, Lou L, Tian M, Li Q, Ding Y, Cao X, Wu Y, Belda-Palazon B, Rodriguez PL, Yang S et al. ESCRT-I Component VPS23A Affects ABA Signaling by Recognizing ABA Receptors for Endosomal Degradation. *Mol Plant* **2016**, *9*, 1570-1582, doi:10.1016/j.molp.2016.11.002.
157. Belda-Palazon B, Rodriguez L, Fernandez MA, Castillo MC, Anderson EM, Gao C, Gonzalez-Guzman M, Peirats-Llobet M, Zhao Q, De Winne N et al. FYVE1/FREE1 Interacts with the PYL4 ABA Receptor and Mediates Its Delivery to the Vacuolar Degradation Pathway. *Plant Cell* **2016**, *28*, 2291-2311, doi:10.1105/tpc.16.00178.
158. Gao C, Zhuang X, Cui Y, Fu X, He Y, Zhao Q, Zeng Y, Shen J, Luo M, Jiang L. Dual roles of an Arabidopsis ESCRT component FREE1 in regulating vacuolar protein transport and autophagic degradation. *Proc Natl Acad Sci U S A* **2015**, *112*, 1886-1891, doi:10.1073/pnas.1421271112.
159. Garcia-Leon M, Cuyas L, El-Moneim DA, Rodriguez L, Belda-Palazon B, Sanchez-Quant E, Fernandez Y, Roux B, Zamarrano AM, Garcia-Mina JM et al. Arabidopsis ALIX Regulates Stomatal Aperture and Turnover of Absciscic Acid Receptors. *Plant Cell* **2019**, *31*, 2411-2429, doi:10.1105/tpc.19.00399.
160. Yu F, Cao X, Liu G, Wang Q, Xia R, Zhang X, Xie Q. ESCRT-I Component VPS23A Is Targeted by E3 Ubiquitin Ligase XBAT35 for Proteasome-Mediated Degradation in Modulating ABA Signaling. *Mol Plant* **2020**, *13*, 1556-1569, doi:10.1016/j.molp.2020.09.008.
161. Xia FN, Zeng B, Liu HS, Qi H, Xie LJ, Yu LJ, Chen QF, Li JF, Chen YQ, Jiang L et al. SINAT E3 Ubiquitin Ligases Mediate FREE1 and VPS23A Degradation to Modulate Absciscic Acid Signaling. *Plant Cell* **2020**, *32*, 3290-3310, doi:10.1105/tpc.20.00267.
162. Seo DH, Ahn MY, Park KY, Kim EY, Kim WT. The N-Terminal UND Motif of the Arabidopsis U-Box E3 Ligase PUB18 Is Critical for the Negative Regulation of ABA-Mediated Stomatal Movement and Determines Its Ubiquitination Specificity for Exocyst Subunit Exo70B1. *Plant Cell* **2016**, *28*, 2952-2973, doi:10.1105/tpc.16.00347.
163. Jurkiewicz P, Batoko H. Protein degradation mechanisms modulate absciscic acid signaling and responses during abiotic stress. *Plant Sci* **2018**, *267*, 48-54, doi:10.1016/j.plantsci.2017.10.017.
164. Mucha E, Fricke I, Schaefer A, Wittinghofer A, Berken A. Rho proteins of plants--functional cycle and regulation of cytoskeletal dynamics. *Eur J Cell Biol* **2011**, *90*, 934-943, doi:10.1016/j.ejcb.2010.11.009.
165. Nagawa S, Xu T, Yang Z. RHO GTPase in plants: Conservation and invention of regulators and effectors. *Small GTPases* **2010**, *1*, 78-88, doi:10.4161/sgtp.1.2.14544.
166. Vanhee C, Batoko H. Autophagy involvement in responses to absciscic acid by plant cells. *Autophagy* **2011**, *7*, 655-656, doi:10.4161/auto.7.6.15307.
167. Li H, Li Y, Zhao Q, Li T, Wei J, Li B, Shen W, Yang C, Zeng Y, Rodriguez PL et al. The plant ESCRT component FREE1 shuttles to the nucleus to attenuate absciscic acid signalling. *Nat Plants* **2019**, *5*, 512-524, doi:10.1038/s41477-019-0400-5.
168. Yamauchi S, Mano S, Oikawa K, Hikino K, Teshima KM, Kimori Y, Nishimura M, Shimazaki KI, Takemiya A. Autophagy controls reactive oxygen species homeostasis in guard cells that is essential for stomatal opening. *Proc Natl Acad Sci U S A* **2019**, *116*, 19187-19192, doi:10.1073/pnas.1910886116.
169. Huang L, Yu LJ, Zhang X, Fan B, Wang FZ, Dai YS, Qi H, Zhou Y, Xie LJ, Xiao S. Autophagy regulates glucose-mediated root meristem activity by modulating ROS production in Arabidopsis. *Autophagy* **2019**, *15*, 407-422, doi:10.1080/15548627.2018.1520547.
170. Bi C, Ma Y, Wu Z, Yu YT, Liang S, Lu K, Wang XF. Arabidopsis ABI5 plays a role in regulating ROS homeostasis by activating CATALASE 1 transcription in seed germination. *Plant Mol Biol* **2017**, *94*, 197-213, doi:10.1007/s11103-017-0603-y.
171. Yu Y, Wang J, Li S, Kakan X, Zhou Y, Miao Y, Wang F, Qin H, Huang R. Ascorbic Acid Integrates the Antagonistic Modulation of Ethylene and Absciscic Acid in the Accumulation of Reactive Oxygen Species. *Plant Physiol* **2019**, *179*, 1861-1875, doi:10.1104/pp.18.01250.
172. Schieke SM, Finkel T. Mitochondrial signaling, TOR, and life span. *Biol Chem* **2006**, *387*, 1357-1361, doi:10.1515/BC.2006.170.

173. Zubo YO, Schaller GE. Role of the Cytokinin-Activated Type-B Response Regulators in Hormone Crosstalk. *Plants (Basel)* **2020**, 9, doi:10.3390/plants9020166.
174. Acheampong AK, Shanks C, Cheng CY, Schaller GE, Dagdas Y, Kieber JJ. EXO70D isoforms mediate selective autophagic degradation of type-A ARR proteins to regulate cytokinin sensitivity. *Proc Natl Acad Sci U S A* **2020**, 117, 27034-27043, doi:10.1073/pnas.2013161117.
175. Wang Y, Li L, Ye T, Zhao S, Liu Z, Feng YQ, Wu Y. Cytokinin antagonizes ABA suppression to seed germination of Arabidopsis by downregulating ABI5 expression. *The Plant journal : for cell and molecular biology* **2011**, 68, 249-261, doi:10.1111/j.1365-3113X.2011.04683.x.
176. Lv M, Li J. Molecular Mechanisms of Brassinosteroid-Mediated Responses to Changing Environments in Arabidopsis. *International journal of molecular sciences* **2020**, 21, doi:10.3390/ijms21082737.
177. Nolan TM, Vukasinovic N, Liu D, Russinova E, Yin Y. Brassinosteroids: Multidimensional Regulators of Plant Growth, Development, and Stress Responses. *Plant Cell* **2020**, 32, 295-318, doi:10.1105/tpc.19.00335.
178. Planas-Riverola A, Gupta A, Betegon-Putze I, Bosch N, Ibanes M, Cano-Delgado AI. Brassinosteroid signaling in plant development and adaptation to stress. *Development* **2019**, 146, doi:10.1242/dev.151894.
179. Wei Z, Li J. Regulation of Brassinosteroid Homeostasis in Higher Plants. *Front Plant Sci* **2020**, 11, 583622, doi:10.3389/fpls.2020.583622.
180. Hu Y, Yu D. BRASSINOSTEROID INSENSITIVE2 interacts with ABSCISIC ACID INSENSITIVE5 to mediate the antagonism of brassinosteroids to abscisic acid during seed germination in Arabidopsis. *Plant Cell* **2014**, 26, 4394-4408, doi:10.1105/tpc.114.130849.
181. Jiang H, Tang B, Xie Z, Nolan T, Ye H, Song GY, Walley J, Yin Y. GSK3-like kinase BIN2 phosphorylates RD26 to potentiate drought signaling in Arabidopsis. *The Plant journal : for cell and molecular biology* **2019**, 100, 923-937, doi:10.1111/tpj.14484.
182. Cai Z, Liu J, Wang H, Yang C, Chen Y, Li Y, Pan S, Dong R, Tang G, Barajas-Lopez Jde D et al. GSK3-like kinases positively modulate abscisic acid signaling through phosphorylating subgroup III SnRK2s in Arabidopsis. *Proc Natl Acad Sci U S A* **2014**, 111, 9651-9656, doi:10.1073/pnas.1316717111.
183. Wang H, Tang J, Liu J, Hu J, Liu J, Chen Y, Cai Z, Wang X. Absciscic Acid Signaling Inhibits Brassinosteroid Signaling through Dampening the Dephosphorylation of BIN2 by ABI1 and ABI2. *Mol Plant* **2018**, 11, 315-325, doi:10.1016/j.molp.2017.12.013.
184. Xiong F, Zhang R, Meng Z, Deng K, Que Y, Zhuo F, Feng L, Guo S, Datla R, Ren M. Brassinosteroid Insensitive 2 (BIN2) acts as a downstream effector of the Target of Rapamycin (TOR) signaling pathway to regulate photoautotrophic growth in Arabidopsis. *New Phytol* **2017**, 213, 233-249, doi:10.1111/nph.14118.
185. Nolan TM, Brennan B, Yang M, Chen J, Zhang M, Li Z, Wang X, Bassham DC, Walley J, Yin Y. Selective Autophagy of BES1 Mediated by DSK2 Balances Plant Growth and Survival. *Dev Cell* **2017**, 41, 33-46 e37, doi:10.1016/j.devcel.2017.03.013.
186. Chi C, Li X, Fang P, Xia X, Shi K, Zhou Y, Zhou J, Yu J. Brassinosteroids act as a positive regulator of NBR1-dependent selective autophagy in response to chilling stress in tomato. *J Exp Bot* **2020**, 71, 1092-1106, doi:10.1093/jxb/erz466.