

Modulation of endocannabinoids by caloric restriction is conserved in mice but is not required for the protection from acute kidney injury

Karla Johanna Ruth Hoyer-Allo^{1,2*}, Martin Richard Späth^{1,2*}, Ruth Hanßen^{3,4}, Marc Johnsen^{1,2}, Susanne Brodesser², Kathrin Kaufmann², Katharina Kiefer², Felix Carlo Koehler^{1,2}, Heike Goebel⁵, Torsten Kubacki^{1,2}, Franziska Grundmann¹, Bernhard Schermer^{1,2}, Jens Brüning^{3,4}, Thomas Benzing^{1,2}, Volker Burst^{1,#} and Roman-Ulrich Müller^{1,2,#}

¹ Department II of Internal Medicine and Center for Molecular Medicine Cologne, University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany

² CECAD, University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany

³ Max-Planck-Institute for Metabolism Research, University of Cologne, Faculty of Medicine and University Hospital Cologne Cologne, Germany and,

⁴ Polyclinic for Endocrinology, Diabetes and Preventive Medicine (PEPD), University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany

⁵ Institute of Pathology, University Hospital of Cologne, Cologne, Germany

*** authors contributed equally**

#Address correspondence to:

Roman-ulrich.mueller@uk-koeln.de

Volker.burst@uk-koeln.de

Abstract (max. 200 words)

Acute kidney injury (AKI) is a frequent and critical complication in the clinical setting. In rodents AKI can be prevented effectively through caloric restriction (CR), which has also been shown to increase lifespan in many species. In *Caenorhabditis elegans* (*C. elegans*) longevity studies revealed that a marked CR-induced reduction of endocannabinoids may be a key mechanism. Thus, we hypothesized that regulation of endocannabinoids, in particular arachidonoyl ethanolamide (AEA), might also play a role in CR-mediated protection from renal ischemia-reperfusion injury (IRI) in mammals including humans. In male C57Bl6J mice, CR significantly reduced renal IRI and led to a significant decrease of AEA. Supplementation of AEA to near-normal serum concentrations by repetitive intraperitoneal administration in CR mice, however, did not abrogate the protective effect of CR. We also analyzed serum samples taken before and after CR from patients of three different pilot trials of dietary interventions. In contrast to mice and *C. elegans*, we detected an increase of AEA. We conclude that endocannabinoid levels in mice are modulated by CR, but CR-mediated renal protection does not depend on this effect. Moreover, our results indicate that modulation of endocannabinoids by CR in humans may differ fundamentally from the effects in animal models.

Keywords

Acute kidney injury, ischemia-reperfusion injury, preconditioning, caloric restriction, stress resistance, endocannabinoid, anandamide

Introduction

Acute kidney injury (AKI) is one of the most frequent complications in hospitalized patients. In particular in the elderly, AKI shows a rapidly rising incidence and is strongly associated with increased mortality and morbidity [1]. Although, kidney function tends to recover at least partially, AKI is associated with a deterioration of clinical outcomes including an increased risk of chronic kidney disease (CKD) and progression to end-stage renal disease (ESRD) [2-7]. In general, progression to ESRD can be the result of repetitive, minor events of AKI during lifetime [8, 9]. While pre- and postrenal causes of AKI can be corrected rapidly in most of the cases, there is no causal therapy for intrarenal AKI – e.g. due to ischemia-reperfusion injury (IRI) or drug toxicity - in the clinical setting [10-16].

Despite knowledge on risk factors and decades of research on AKI, targeted approaches towards prevention are still missing. In contrast, extremely effective approaches have been developed that are capable of preventing or attenuating AKI in rodent models [17-20]. These preconditioning strategies enhance cellular stress resistance before exposure to the injurious stimulus. Caloric restriction (CR) is one of the most potent preconditioning protocols and has been shown to increase stress resistance and increase lifespan in a multitude of species ranging from the nematode *Caenorhabditis elegans* (*C. elegans*) to non-human primates [18-22].

The molecular mechanisms underlying the impact of CR on resilience are still incompletely understood. However, this knowledge would be crucial to develop targeted approaches for organ protection in the patient setting. In *C. elegans*, CR has been shown to result in a significant reduction of N-acyl ethanolamines (NAEs). This decrease of NAEs was required and sufficient for the CR-mediated extension of lifespan [23]. NAEs are endocannabinoids and as such lipid-derived signaling molecules that are – among other organs - synthesized in the kidney [24-26]. The aim of this study was to evaluate whether (1) CR-induced reduction of NAE levels is conserved in mammals and (2) whether this reduction is involved in CR-induced renal organ protection. To reach this goal, we used a murine model

of renal IRI and supplemented anandamide (AEA), one of the two most prominent NAEs in mammals [27], to calorically restricted male mice.

Results

Anandamide is reduced by CR and can be increased by intraperitoneal administration in mice

To examine if the CR-induced reduction of endocannabinoids previously observed in *C. elegans* was conserved in mice, we analyzed plasma samples of male C57Bl6J wildtype mice after CR and without intervention. In *C. elegans*, among a row of different NAEs reduced by CR, the supplementation of two C20-fatty-acid-containing NAEs – eicosapentaenoyl ethanolamide (EPEA) and arachidonoyl ethanolamide (AEA) – was able to rescue phenotypes associated with NAE depletion [23]. Whilst EPEA is more abundant in worms, this NAE was not detectable in mouse serum (Fig. S1). However, AEA was the C20-fatty-acid containing endocannabinoid most strongly reduced by CR in rodents (Fig. 1A).




Fig. 1

Fig. 1: Anandamide is reduced by CR and dose finding reveals a linear increase.

A. AEA levels normalized to the mean of ad libitum fed male mice (CTRL) in comparison to calorically restricted male mice (CR). **B.** Schematic illustration of the dose-finding experiment. **C.** AEA levels after intraperitoneal injections of 0.00 (vehicle), 0.02, 0.05, 0.075 and 0.1 mg/g bodyweight resulted in a linear rise of AEA. AEA concentrations were normalized to the mean of the vehicle-treated mice. Abb.: AEA: Anandamide; CR: caloric restriction; CTRL: no intervention; d: day; veh: vehicle control; bars indicate mean \pm standard deviation; ***: p-value < 0.001; dashed line: chosen target dose of 0.01 mg/g bodyweight.

Consequently, we decided to examine whether the reduction in AEA was required for CR-mediated renal organ protection using a mouse model of unilateral nephrectomy and contralateral IRI. As a first step, we performed a dose-finding experiment for AEA

supplementation. Since the protective effects of CR can be shown already after a few days, we deemed a period of 8 days of supplementation sufficient for this purpose (Fig. 1B). Intraperitoneal injections of 0.00 (vehicle), 0.02, 0.05, 0.075 and 0.1 mg/g bodyweight resulted in a linear rise ($r^2= 0.96$, $p=0.38$) of AEA. Since already the lowest concentration resulted in a strong increase of AEA serum levels (Fig. 1C), we continued the experiment using a concentration of 0.01 mg/g bodyweight.

The CR-mediated protection from renal IRI does not depend on the reduction of AEA

To examine, if CR-mediated protection from AKI depends on the loss of AEA, we injected AEA (0.01 mg/g bodyweight) or vehicle to calorically restricted mice (CR) or *ad libitum* fed controls (CTRL). All groups underwent renal IRI or sham-IRI (sham) simultaneously (Fig. 2A).




Fig. 2

Fig. 2: Intrapерitoneal AEA administration does not abrogate the CR-mediated kidney protection in mice.

A. Schematic illustration of the experimental setup **B.** Plasma creatinine values 24 h after ischemia-reperfusion injury (IRI). **C.** Plasma blood urea nitrogen values 24 h after IRI.

Abb.: CR: caloric restriction; CR-AEA: calorically restricted mice treated with AEA-injections on the last eight days prior to ischemia-reperfusion injury (IRI); CR-veh: calorically restricted mice treated with vehicle-injections on the last eight days prior to ischemia-reperfusion injury (IRI); CTRL: no preconditioning; CTRL-AEA: *ad libitum* fed mice

with AEA-injections on the last eight days prior to ischemia-reperfusion injury (IRI); CTRL-veh: *ad libitum* fed mice with vehicle-injections on the last eight days prior to ischemia-reperfusion injury; d: day; IRI: ischemia-reperfusion injury; sham: right nephrectomy followed by no-clamping of the left renal pedicle. Bars indicate mean \pm standard deviation; ***: p-value<0.001; n.s.: p-value>0.05.

Intraperitoneal supplementation of AEA or vehicle did not influence weight loss upon CR (Fig. S2). As expected, 24 h after IRI, creatinine and blood urea nitrogen (BUN) were significantly increased in *ad libitum* fed mice (CTRL) compared to sham operated animals (sham) (Fig. 2B, C). This almost 15-fold increase was entirely prevented by CR (Fig. 2B, C). The protective effect of CR was not affected by AEA supplementation (Fig. 2B, C). Likewise, AEA or vehicle supplementation in *ad libitum* fed mice did not affect the outcome after IRI (CTRL_AEA/CTRL_veh, Fig. S3 A, B).

Intraperitoneal AEA supplementation does not abrogate the CR-mediated protection from cell death and tubular damage

For histological examination of tubular injury Periodic Acid-Schiff-Staining (PAS) was performed. As expected, kidneys of *ad libitum* fed mice (CTRL) showed increased pyknosis, stronger brush border loss and stronger epithelial flattening after IRI in comparison to sham-operated (sham) and to CR-mice (Fig. 3).




Fig. 3

Fig. 3: **Histological confirmation: intraperitoneal AEA administration does not abrogate the protective effect of CR.**

Histological analyses reveal higher tubular damage in ad libitum fed animals in comparison to calorically restricted male mice. AEA-injection does not abrogate the protection by CR in mice.

Abb.: Asterisk: brush border loss, arrowhead: epithelial flattening; CR: caloric restriction; CR-AEA: calorically restricted mice treated with AEA-injections on the last eight days prior to ischemia-reperfusion injury (IRI); CR-veh: calorically restricted mice treated with vehicle-injections on the last eight days (d) prior to ischemia-reperfusion injury (IRI); CTRL: no preconditioning; d: day; IRI: ischemia-reperfusion injury; sham: right nephrectomy followed by no-clamping of the left renal pedicle; square: pyknosis; ; PAS: Periodic acid-Schiff reaction; caspase3: immunohistochemical staining against cleaved-caspase3; TUNEL: TdT-mediated dUTP-biotin nick end labeling.

Kidneys of CR-mice injected with AEA or vehicle (CR_AEA/CR_veh) were indistinguishable from CR animals without intraperitoneal injections (CR) after IRI on the level of histology. Likewise, supplementation of AEA or vehicle to *ad libitum* fed mice did not affect the histological outcome after IRI (CTRL_AEA/CTRL_veh). This was confirmed in a semi-

quantitative analysis of the histological damage using a composite damage scoring system (Fig. S4).

In addition, we assessed cell death in kidneys 24 h after IRI using cleaved-caspase 3 stainings and terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL). Whilst kidneys of *ad libitum*-fed mice (CTRL) showed a strong increase in cell death, this increase was entirely prevented by CR. Again, AEA supplementation or vehicle administration did not have any impact on this protection (CR-AEA/CR-veh, Fig. 3) and did not change the induction of cell death after IRI in AL-fed mice (CTRL-AEA/CTRL-veh) (Fig. S5).

CR is associated with an increase of AEA in humans

To answer the question if CR-induced reduction of endocannabinoids and specifically AEA is even conserved in humans, we assembled a set of longitudinally acquired human serum samples before and after CR derived from different investigator initiated clinical trials at our center. The first trial investigated the protective effect of CR in patients undergoing elective cardiac surgery on cardiopulmonary bypass (CR_KCH, NCT01534364, [28]). After randomization, subjects received a formula diet that supplied 60% of daily energy expenditure (DEE) for the last 7 days prior to surgery, control subjects were not restricted in their food intake. In contrast to the findings in mice and worms, analysis of serum samples of 19 patients (> 60 years of age) revealed significantly higher AEA levels after CR (Fig. 4A).




Fig. 4

Fig. 4: **AEA-increase after short-term CR in humans.**

A. Normalized AEA levels in human serum samples withdrawn from elderly subjects (> 60 years) before (pre-diet) and after dietary intervention (post-diet) with a formula diet consisting of the amount of 60 % of daily energy expenditure (DEE) for 7 days prior to a coronary-arteria bypass graft. **B.** Normalized AEA levels in human serum samples withdrawn from middle-aged (42 – 64 years, n=8) healthy living kidney donors before (pre-diet) and after dietary intervention (post-diet) with formula diet with the amount of 50 % of the DEE for the last week prior to kidney donation. **C.** Normalized AEA levels in human serum samples withdrawn from middle-aged (29 – 65 years, n=8) obese (BMI > 30 kg/m²) subjects before (pre-diet) and after dietary intervention (post-diet) with the fixed amount of 800 kcal/d for 3 months. Abb.: AEA (Anandamide) concentrations were normalized to the mean of the pre-diet concentrations. Depicted are means \pm standard deviation; *: p-value<0.05, n.s.: p-value>0.05.

Consequently, we decided to check AEA in serum samples from a second study examining middle-aged (42 – 64 years, n=6) healthy living kidney donors that were provided a formula diet containing 50% of the DEE for the last week prior to kidney donation (CR_LSP, NCT02745444) (Fig. 4B). Again, CR led to a significant increase of AEA (Fig. 4B).

Finally, samples from a third cohort (optifast) of obese individuals (BMI > 34 kg/m², 29 – 65 years, n=8) undergoing a 3-months formula diet with the fixed amount of 800 kcal/d to reach weight-loss showed a trend towards higher AEA levels after CR (p-value: 0.1296) (Fig. 4C). Detailed baseline characteristics of the three cohorts are shown in supplemental table S1.

Discussion

CR-mediated preconditioning has been shown to be extremely effective in prevention of AKI in rodents [16, 18, 21, 29]. In *C. elegans* CR is a classical longevity intervention increasing lifespan and metabolic health. In line with the effect on organ protection in mammals, longevity pathways also increase stress resistance in nematodes. Reduced N-acyl-ethanolamine (NAE) levels upon CR have been shown to be both sufficient and required for CR-mediated lifespan extension [23].

NAEs comprise a heterogeneous group of lipid-derived molecules that can activate the G-protein-coupled receptors CB-1 and CB-2 in the central nervous system and peripheral organs, e.g. the kidney [27]. NAE signaling is involved in the regulation of food intake [30], energy metabolism [31], inflammation [32] and vascular tone [24]. Nevertheless, the exact underlying mechanisms remain elusive [24].

CR led to a reduction of 6 different NAEs in the nematode, whilst only supplementation of the C20- fatty acid containing NAEs – EPEA and AEA – were able to prevent dauer arrest and promote reproductive growth of *daf-2* mutants (Lucanic, Held et al. 2011). In the present study we were able to show that CR significantly reduced NAEs, including AEA, in mice.

Notably, CR consisting of 80 % of the daily intake for two weeks did not influence AEA levels of C57Bl6J mice in one previous study [33]. We assume this difference to be the result of a more stringent CR protocol, i.e. a longer period of lower calorie intake in the study at hand.

EPEA is known to be more abundant in worms than in rodents [34], explaining why this NAE was not detected in our mouse serum. Instead, AEA was the C20-fatty-acid containing NAE most strongly reduced by CR in our experiment. Therefore, we decided to evaluate the impact of AEA-reduction on CR-induced renal organ protection in mice.

AEA is the first NAE identified [27, 35]. Whilst there is ample evidence that AEA is generated by enzymatic cleavage of N-arachidonoyl-phosphatidylethanolamine (NAPE) through NAPE-phospholipase D [36-39], the underlying mechanism of NAPE-generation in kidneys remains elusive [27]. AEA shows high abundance in renal tissue [35, 40] and was detected in endothelial and mesangial cells [26]. AEA showed higher concentrations in the

renal medulla than in the cortex [27, 41]. Regarding its function, AEA can activate CB-1/2 receptors, but also transient receptor potential vanilloid type I channels (TRPV1) localized in the vasculature [27, 42]. Whilst CB-1 is expressed nearly ubiquitously in the nephron [43], CB-2 was only detected in podocytes [44], proximal tubular [45] and mesangial cells [26]. AEA gets inactivated enzymatically by fatty acid amide hydrolase (FAAH) and monoacylglycerol lipase (MGL) [46]. Under physiologic conditions AEA contributes to the regulation of glomerular filtration, the vascular tonus and sodium-transport [24].

Current mechanistic concepts suggest, that CR-mediated cellular resistance is based on reduced IGF-1/Insulin signaling [18], activation of AMPK [47] and furthermore an inhibition of the mammalian target of rapamycin (mTOR) [48]. In the nematode, Lucanic *et al.* concluded, that the TOR-pathway might control NAE levels in response to nutrient intake and that reduced NAE signaling is a component of CR-mediated lifespan extension [23]. On the other hand, there is evidence that NAEs activate mTOR [49, 50] and AMPK signaling in renal proximal tubular cells [51] as well as hepatocytes [52].

In the context of AKI, the role of NAEs seems to be ambivalent. Whereas CB-1-inhibition reduced the decline of kidney function after cisplatin administration [53], CB-1 and CB-2 activation was demonstrated to mediate protective effects in renal IRI by increasing the renal blood flow through vasodilatation [54-56]. Furthermore, inhibition of the degradation of another NAE - of 2-Arachidonylglycerol (2-AG) - was shown to improve the outcome after bilateral renal IRI [57].

Here, we provide the first detailed study of CR-dependent NAE-modulation in mice and humans combined with pharmacological testing of the contribution of AEA to CR-mediated renal organ protection. In line with previous investigations, we found AKI induced by renal IRI to be completely prevented by CR in C57Bl6J mice. This protection was accompanied by the reduction in AEA. However repetitive intraperitoneal supplementation of AEA did not show any impact on the outcome 24 h after renal IRI (Fig. 2 & S3). It is important to emphasize a number of differences in study design compared to previously published work on NAEs in renal IRI. First, different models of renal IRI (uni-/ bilateral without nephrectomy) were used [54-57].

Second, this is the first study employing repetitive intraperitoneal supplementation of AEA. Finally, differences between mouse strains may also play a role, since only Moradi *et al.* used C57Bl6 mice in a model of bilateral IRI [57].

Since the lack of effect of AEA-supplementation could have been a consequence of the experimental protocol used, we decided to – before moving on to additional phenotypic mouse studies - check whether AEA reduction upon CR was indeed also conserved in humans. This knowledge would also be important as a potential basis to human trials modulating NAE levels by CR.

Interestingly, the literature concerning dietary influences on AEA in humans is much more controversial than in rodents [33] or *C. elegans* [23]. Therefore, we analyzed human serum samples acquired before and after short-term CR in three independent cohorts. Of note, results from two of these cohorts revealed a significant increase in AEA after CR. The third study showed a trend towards this direction ($p=0.1296$) not reaching statistical significance, possibly as a consequence of comparably high heterogeneity of this cohort. Notably, recent publications provide evidence for either no influence on AEA [58, 59] or a reduction of AEA after longer-term interventions (e.g. gastric-bypass, life style interventions programs or prolonged CR) [60-62]. Considering that fasting raises and semi-starvation reduces endocannabinoid levels [63, 64] it seems possible that the observed differences are related to the duration of the particular interventions. The mechanisms underlying the impact of different CR protocols on NAE levels will be an interesting question for future studies.

Taken together, this study clearly shows, that NAEs are reduced by CR in mice. Nevertheless, this effect does not seem to be involved in CR-mediated protection from murine renal IRI. Furthermore, our analyses indicate that the CR-induced modulation of NAEs in humans may differ fundamentally from the effects in animal models.

Methods

Animal procedures

All animal experiments were carried out in accordance with the LANUV NRW (Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, State Agency for Nature, Environment and Consumer Protection North Rhine-Westphalia; VSG 84-02.04.2013.A158). Male wildtype C57/Bl6J mice were provided by Charles River and were used for all experiments. All mice were kept under similar specific pathogen free (SPF) conditions in group-cages with up to 5 mice at the same time and a daily 12 hour light and 12 hour dark rhythm. The mice received *ad libitum* access to drinking water and food with one exception for the caloric restriction. The standard food chow was purchased from sniff GmbH (Soest, Germany).

As previously described [16, 21, 29] calorically restricted mice were provided with 66 % (3 g per day per mouse) daily in the morning on the last 28 days prior to surgery. Before initiation of dietary preconditioning CR-mice were set individually in new cages. Plasma samples of calorically restricted animals were obtained in the morning before feeding.

Anandamide was purchased by Cayman Chemical (Ann Arbor, Michigan, USA) and administrated (0.01 mg/g bodyweight) by daily intraperitoneal injections in a volume of 10 μ L/g bodyweight for 1 week prior to IRI (11 to 12 weeks of age). Standard sodium-chloride solution (0.9 %) with 0.05% Ethanol in an amount of 10 μ L/g bodyweight was used for the vehicle control group.

Ischemia-Reperfusion Injury (IRI)

Renal IRI surgery was performed in 12-13-week-old male C57Bl6J mice under anesthesia (ketamine (Zoetis (Berlin, Germany))/xylazine (Bayer (Leverkusen, Germany))). To maintain the body temperature surgery was performed on a temperature-controlled heating pad (Havard apparatus, Holliston, Massachusetts, USA). As described before [21, 29], following midline abdominal incision the vessels of the right kidney were ligated with sutures

and the kidney was removed afterwards. Following this, the left renal artery and vein underwent unilateral clamping for 40 minutes with an atraumatic micro-vascular clamp and subsequent removal of the clamp for reperfusion, which was controlled visually. The abdominal incision was closed in two layers by silk suture in the body wall and clips in the skin.

Sham-operated animals underwent the same approach with the exception that the left renal pedicles were not clamped. After surgery the mice obtained normal saline containing 0.2 mg Buprenorphine subcutaneously and access to drinking water supplemented with metamizole. To prevent mutilation of the wound by fellow mice, all operated mice were transferred to single cages.

24 h after reperfusion the mice received intraperitoneal injections of ketamine/xylazine for anesthesia. First blood was taken in heparin coated 1 mL syringes by cardiac puncture of the right ventricle and centrifuged at 4000 g for 10 minutes at 4°C. As done 24 h before to the right kidneys, the left kidneys were cut into two similar sections, whereby one section was fixed in 4% formaldehyde and subsequently embedded in paraffin and the other section was snap frozen in liquid nitrogen for further analyses.

Functional data and statistics

Measurements of plasma urea and creatinine were implemented by the central laboratory of the University Hospital of Cologne using a Cobas C 702 (Roche Diagnostics, Mannheim, Penzberg, Germany). For statistical analysis GraphPad Prism 8 (GraphPad, San Diego, CA, USA) was used. For analyses of mouse studies with three or more treatment groups one-way analyses of variances (ANOVA) followed by Tukey posthoc-tests were used. For analyses of the longitudinally acquired human blood samples paired Student's t-tests were performed. In all cases two-tailed p-values <0.05 were defined as significant and all data are represented as means \pm standard deviations (SD).

Histopathology

Following the fixation in 4% formaldehyde all kidney tissues were embedded in paraffin and cut into 2 μm slices. By Periodic acid-Schiff (PAS) reaction staining the slices were used for morphological evaluations. One experienced nephropathologist analyzed the tubular damage of each kidney in a blinded fashion by using a composite score as previously described [16, 65]. The score reflects the percentage of tubules which presented epithelial flattening, loss of nucleus, epithelial vacuolization, debris and loss of brush boarder (0: no change, 1: 1-24%; 2: 25-50%; 3: >50%). The mean sum for each treatment group was plotted.

Immunohistochemical analysis was carried out by staining Anti-cleaved Caspase-3 antibody (Asp175, Cell Signaling Technology, Danvers, Massachusetts, USA) in a dilution of 1:400 according to the manufacturer's protocol. Images were acquired in a 20x magnification using a slidescanner SCN4000 (Leica, Wetzlar, Germany). Beyond that, terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) staining was done using the DeadEnd Fluorometric TUNEL System (Promega, Madison, Wisconsin, USA) according to the manufacturer's protocol.

Image acquisition

Images were taken by an Axiovert 200M microscope provided with ZEN software (version Zen 2.6, Zeiss, Oberkochen, Germany).

Quantification of ethanolamides

Levels of ethanolamides in human and murine serum samples were determined by Liquid Chromatography coupled to Electrospray Ionization Tandem Mass Spectrometry (LC-ESI-MS/MS):

Extraction of serum samples was performed as previously described (Zoerner et al., 2012) with several modifications: After thawing on ice, 200 μL of serum were diluted with 800 μL of ice-cold PBS. To all samples 10 μL of an internal standard mixture, containing 100 $\text{pg}/\mu\text{L}$ of each of the following deuterated ethanolamides were added: palmitoyl(16:0) ethanolamide-

d₄, oleoyl(18:1) ethanolamide-d₄, and arachidonoyl(20:4) ethanolamide-d₄ (Cayman Chemical). After incubation on ice for 15 minutes, 1 ml of toluene was added. Extraction was performed by shaking the samples in a ThermoMixer (Eppendorf) for 1 min at 4 °C and 2000 rpm. After centrifugation (16,100 RCF, 5 min, 4 °C), the upper (organic) phase was transferred to a new tube and dried under a stream of nitrogen. The dried extract was resolved in 300 µl of methanol/water 3:1 (v/v). The samples were again centrifuged, and 100 µl of supernatant were transferred to autoinjector vials.

LC-MS/MS analysis was performed by injection of 35 µl of sample onto a Core-Shell Kinetex C18 column (150 mm × 2.1 mm ID, 2.6 µm particle size, 100 Å pore size, Phenomenex) and detection using a QTRAP 6500 triple quadrupole/linear ion trap mass spectrometer (SCIEX). The LC (1260 Infinity Binary LC System, Agilent) was operated at 40 °C and at a flow rate of 0.5 ml/min with a mobile phase of 5 mM ammonium acetate and 0.1 % acetic acid in water, pH 3.5 (solvent A) and acetonitrile (solvent B). Ethanolamides were eluted with the following gradient: initial, 10 % B; 3 min, 100 % B; 7 min, 100 % B; 8 min, 90 % B; 11 min, 90 % B.

Ethanolamides were monitored in the positive ion mode with their specific Multiple Reaction Monitoring (MRM) transitions (Williams *et al.*, 2007). The instrument settings for nebulizer gas (Gas 1), turbogas (Gas 2), curtain gas, and collision gas were 70 psi, 55 psi, 35 psi, and medium, respectively. The interface heater was on, the Turbo V ESI source temperature was 550 °C, and the ionspray voltage was 5 kV. For all MRM transitions the values for the entrance potential and the cell exit potential were 10 V and 8 V, respectively. The collision energies ranged from 17 to 25 V, the declustering potentials from 70 to 110 V.

The LC chromatogram peaks of endogenous ethanolamides and deuterated internal standards were integrated using the Analyst 1.6.3 software (SCIEX, Darmstadt, Germany). Endogenous ethanolamides were quantified on the basis of external calibration curves which were calculated from LC-MS/MS measurements of serially diluted synthetic palmitoyl ethanolamide, oleoyl ethanolamide, and arachidonoyl ethanolamide standard solutions within

the range of 0.0 to 40.8 pmol on column. To each dilution fixed amounts of the deuterated internal standards were added. The standard calibration curves were plotted based on molar concentration versus peak area ratio of ethanolamide standards to deuterated internal standards. Linearity and correlation coefficients (R^2) of the calibration curves were obtained via linear regression analysis. R^2 of the calibration curves were >0.99 . Linoleoyl(18:2) ethanolamide was quantified on the basis of the calibration curve of oleoyl ethanolamide, for the quantification of docosahexaenoyl (22:6) ethanolamide the calibration curve of arachidonoyl ethanolamide was used. The calculated amounts of endogenous ethanolamides were normalized to the volume of the serum sample.

Human sample acquisition

Human samples were obtained as part of three different clinical studies at our center. The first cohort consisted of elderly subjects (> 60 years) and was provided with formula diet with the amount of 60 % of daily energy expenditure (DEE) for 7 days prior to a coronary-arteria bypass graft ([28], study name: CR_KCH, clinicaltrials.gov identifier: NCT01534364). The second cohort (CR_LSP, clinicaltrials.gov identifier: NCT02745444) consisted of middle-aged (42 – 64 years, $n=6$) healthy living kidney donors provided with formula diet with the amount of 50 % of the DEE for the last week prior to kidney donation. The subjects of the third cohort (OptiFast, application number of the local ethics committee: 10-226) were middle-aged (29 – 65 years, $n=8$) and obese ($BMI > 30 \text{ kg/m}^2$). They were kept on a formula with the fixed amount of 800 kcal/d for 3 months. Blood samples we included in our analyses were taken at study inclusion and after the diet after at least 5 h of fasting. All studies were operated in accordance with the Declaration of Helsinki and the good clinical practice guidelines by the International Conference on Harmonization. All patients provided informed consent and approval of each study protocol was obtained from the local institutional review board (Ethics committee of the University of Cologne, Cologne, Germany).

Supplemental Material

Supplemental table 1: **Detailed patient characteristics.**

	CR_KCH	CR_LSP	Optifast
diet	CR	CR	CR
number of subjects	19	6	8
mean age (min/max)	68.2 (55/82)	52.4 (42.72/63.14)	49.88 (29/59)
male (%)	73.7	66.66	37.5
mean bodyweight at screening/kg (min/max)	79.95 (55.9/98.1)	90.93 (80.6/112.2)	129.91 (106.75/157.6)
mean BMI (min/max)	27.38 (21/34.7)	28.98 (25.2/33.5)	42.21 (34.8/47.2)
mean Δ weight (min/max)	-2.78 (-5.6/0.6)	-2.18 (-4/-0.6)	-22.48 (-33.4/-13.9)

Supplemental table legend

Supplemental table 1: **Detailed patient characteristics.** BMI: body mass index; min.: minimum; max.: maximum

Supplemental figure legends

Fig. S1: **CR leads to a significant reduction of several endocannabinoids.**

Levels of the measured set of endocannabinoids on ad libitum diet in comparison to CR. Concentrations were normalized to the mean AEA levels in ad libitum fed mice (Fig. 1). Abb.: Bars indicate means \pm standard deviation; ***: p-value<0.001. ad lib: ad libitum; CR: caloric restriction; 16:0 N-palmitoyl ethanolamide; 18:1: N-oleoyl ethanolamide; 18:2: N-linoleoyl ethanolamide; 20:1: N-eicosenoyl ethanolamide; 20:3: N-eicosatrienoyl ethanolamide; 20:4: N-arachidonoyl ethanolamide/Anandamide; 22:6: N-Docosahexaenoyl ethanolamide.

Fig. S2: Mean bodyweight of the different experimental groups.

Schematic illustration of the mean bodyweight of 4, 3, 2, 1 weeks and directly prior to and 24 h after IRI. Abb.: AEA: ad libitum fed mice treated with anandamide-injections on the last eight days (d) prior to ischemia-reperfusion injury (IRI); CR: caloric restriction; CR-AEA: calorically restricted mice treated with AEA-injections on the last eight days (d) prior to ischemia-reperfusion injury (IRI); CR-veh: calorically restricted mice treated with vehicle-injections on the last eight days (d) prior to ischemia-reperfusion injury (IRI); CTRL: no preconditioning; d: day; IRI: ischemia-reperfusion injury; sham: right nephrectomy followed by no-clamping of the left renal pedicle; veh: *ad libitum* fed mice treated with vehicle-injections on the last eight days (d) prior to ischemia-reperfusion injury (IRI).

Fig. S3: Plasma analyses reveal no difference between AEA- and veh-treated *ad libitum* fed mice.

A. Plasma creatinine values of *ad libitum* fed mice with vehicle-injections (veh) in comparison to AEA-injections on the last eight days prior to ischemia-reperfusion injury **B.** Plasma blood urea nitrogen (BUN) values of male *ad libitum* fed mice with vehicle-injections (veh) in comparison to AEA-injections on the last eight days prior to ischemia-reperfusion injury.

Fig. S4: Quantification of histological damage.

Semiquantitative analysis of histological damage by a score composed of tubular epithelial flattening, tubular loss of nucleus, epithelial vacuolization, debris and loss of brush border. N=3 kidneys of each experimental group in this study were scored by a pathologist in a blinded manner and the mean for each treatment group was plotted. Values indicate means \pm standard deviation. Abb.: CR: caloric restriction; CR-AEA: calorically restricted mice treated with AEA-injections on the last eight days prior to ischemia-reperfusion injury; CR-sham: caloric restriction prior to right nephrectomy followed by no-clamping of the left renal pedicle; CR-veh: calorically restricted mice treated with vehicle-injections on the last eight days prior to ischemia-reperfusion injury; CTRL: no preconditioning; sham: right nephrectomy followed by

no-clamping of the left renal pedicle; CTRL-AEA: ad libitum fed mice treated with anandamide-injections on the last eight days prior to ischemia-reperfusion injury; CTRL-veh: *ad libitum* fed mice treated with vehicle-injections on the last eight days prior to ischemia-reperfusion injury.

Fig. S5: AEA-supplementation does not change the histological outcome in ad libitum fed mice.

Histological analyses do not show differences between ad libitum fed mice after intraperitoneal supplementation of anandamide (AEA) or vehicle (veh).

Abb.: CTRL-AEA: *ad libitum* fed mice after AEA supplementation on the last eight days prior to ischemia-reperfusion injury; CTRL-veh: ad libitum fed mice after vehicle supplementation on the last eight days prior to ischemia-reperfusion injury; PAS: Periodic acid-Schiff reaction; caspase3: immunohistochemical staining against cleaved-caspase3; TUNEL: TdT-mediated dUTP-biotin nick end labeling.

Funding

M.R.S. is supported by the Koeln Fortune program and by the Cologne Clinician Scientist Program (CCSP) / Faculty of Medicine / University of Cologne. Funded by the German Research Foundation (DFG, FI 773/15-1)/ Faculty of Medicine, University of Cologne. F.C.K. is supported by the Koeln Fortune program / Faculty of Medicine, University of Cologne, and has received grants from the German Federal Ministry of Research and Education outside the submitted work. R.-U. M. is supported by the Nachwuchsgruppen. NRW program of the Ministry of Science North Rhine Westfalia (MIWF, to R.-U.M.) and received further funding from the German Research Foundation (CRU 329, MU 3629/3-1). R.-U.M. is further funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – CECAD, EXC 2030 – 390661388. F.G is supported by the Koeln Fortune program, / Faculty of Medicine / University of Cologne and Funded by the German Research Foundation

Acknowledgments

The authors acknowledge support from the CECAD imaging facility (head: Astrid Schauss) and the proficient technical support by Martyna Brütting (CECAD) and Serena Greco-Torres (CECAD).

Author Contributions:

Conceptualization, K.J.R.H.-A., M.R.S., B.S., J.B., T.B., V.B and R.-U.M.; Methodology, K.J.R.H.-A., M.R.S., M.J., T.K. and S.B., Katharina K., Kathrin K., F.C.K. and H.G.; Validation, K.J.R.H.-A. and M.R.S.; Formal Analysis, K.J.R.H.-A., M.R.S., M.J. and S.B ; providing of human resources, R.H., F. G. and J.B.; Writing – Original Draft Preparation, K.J.R.H.-A. and M.R.S.; Writing – Review & Editing, K.J.R.H.-A., M.R.S., F.C.K.,T.K.; Visualization, K.J.R.H.-A. and M.R.S.; Supervision, V.B and R.-U.M.; Project Administration, V.B and R.-U.M.; Funding Acquisition, M.R.S., F.C.K., F.G., R.-U.M..

Institutional Review Board Statement

All animal experiments were carried out in accordance with the LANUV NRW (Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, State Agency for Nature, Environment and Consumer Protection North Rhine-Westphalia; VSG 84-02.04.2013.A158).

The human studies were conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Ethics committee of the University of Cologne, Cologne, Germany (CR_KCH: protocol code 11-296 and date of approval: 2011-12-09; CR_LSP: protocol code: 15-031, date of approval: 2015-04-27; optifast: protocol code: 10-226, date of approval: 2012-05-10).

Informed consent statement

Informed consent was obtained from all subjects involved in the study.

Data Availability Statement

All data in this study can be obtained from the authors.

Conflicts of interests

The Dept. II of Internal Medicine received research funding from Fresenius Kabi. The authors declare that there are no personal conflicts of interest regarding the publication of this paper.

References

1. Rewa, O.; Bagshaw, S. M., Acute kidney injury—epidemiology, outcomes and economics. *Nature Reviews Nephrology* **2014**, 10, 193.
2. Wald, R.; Quinn, R. R.; Luo, J.; Li, P.; Scales, D. C.; Mamdani, M. M.; Ray, J. G., Chronic dialysis and death among survivors of acute kidney injury requiring dialysis. *JAMA* **2009**, 302, (11), 1179-85.
3. Thakar, C. V.; Christianson, A.; Himmelfarb, J.; Leonard, A. C., Acute kidney injury episodes and chronic kidney disease risk in diabetes mellitus. *Clin. J. Am. Soc. Nephrol.* **2011**, 6, (11), 2567-72.
4. Lafrance, J. P.; Miller, D. R., Acute kidney injury associates with increased long-term mortality. *J. Am. Soc. Nephrol.* **2010**, 21, (2), 345-52.
5. Prendecki, M.; Blacker, E.; Sadeghi-Alavijeh, O.; Edwards, R.; Montgomery, H.; Gillis, S.; Harber, M., Improving outcomes in patients with Acute Kidney Injury: the impact of hospital based automated AKI alerts. *Postgrad. Med. J.* **2016**, 92, (1083), 9-13.
6. Chertow, G. M.; Burdick, E.; Honour, M.; Bonventre, J. V.; Bates, D. W., Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *J. Am. Soc. Nephrol.* **2005**, 16, (11), 3365-70.
7. Coca, S. G.; Yusuf, B.; Shlipak, M. G.; Garg, A. X.; Parikh, C. R., Long-term risk of mortality and other adverse outcomes after acute kidney injury: a systematic review and meta-analysis. *Am. J. Kidney Dis.* **2009**, 53, (6), 961-73.
8. Rodríguez, E.; Arias-Cabrales, C.; Bermejo, S.; Sierra, A.; Burballa, C.; Soler, M. J.; Barrios, C.; Pascual, J., Impact of Recurrent Acute Kidney Injury on Patient Outcomes. *Kidney Blood Press. Res.* **2018**, 43, (1), 34-44.
9. Dear, J. W.; Yuen, P. S., Setting the stage for acute-on-chronic kidney injury. *Kidney Int.* **2008**, 74, (1), 7-9.

10. Mao, H.; Katz, N.; Ariyanon, W.; Blanca-Martos, L.; Adýbelli, Z.; Giuliani, A.; Danesi, T. H.; Kim, J. C.; Nayak, A.; Neri, M.; Virzi, G. M.; Brocca, A.; Scalzotto, E.; Salvador, L.; Ronco, C., Cardiac surgery-associated acute kidney injury. *Cardiorenal Med.* **2013**, *3*, (3), 178-199.
11. Kinsey, G. R.; Okusa, M. D., Pathogenesis of acute kidney injury: foundation for clinical practice. *Am. J. Kidney Dis.* **2011**, *58*, (2), 291-301.
12. Krüger, B.; Krick, S.; Dhillon, N.; Lerner, S. M.; Ames, S.; Bromberg, J. S.; Lin, M.; Walsh, L.; Vella, J.; Fischereeder, M.; Krämer, B. K.; Colvin, R. B.; Heeger, P. S.; Murphy, B. T.; Schröppel, B., Donor Toll-like receptor 4 contributes to ischemia and reperfusion injury following human kidney transplantation. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, (9), 3390-5.
13. Salvadori, M.; Rosso, G.; Bertoni, E., Update on ischemia-reperfusion injury in kidney transplantation: Pathogenesis and treatment. *World J Transplant* **2015**, *5*, (2), 52-67.
14. Peerapornratana, S.; Manrique-Caballero, C. L.; Gómez, H.; Kellum, J. A., Acute kidney injury from sepsis: current concepts, epidemiology, pathophysiology, prevention and treatment. *Kidney Int.* **2019**, *96*, (5), 1083-1099.
15. Peres, L. A.; da Cunha, A. D., Jr., Acute nephrotoxicity of cisplatin: molecular mechanisms. *J Bras Nefrol* **2013**, *35*, (4), 332-40.
16. Späth, M. R.; Bartram, M. P.; Palacio-Escat, N.; Hoyer, K. J. R.; Debes, C.; Demir, F.; Schroeter, C. B.; Mandel, A. M.; Grundmann, F.; Ciarimboli, G.; Beyer, A.; Kizhakkedathu, J. N.; Brodesser, S.; Göbel, H.; Becker, J. U.; Benzing, T.; Schermer, B.; Höhne, M.; Burst, V.; Saez-Rodriguez, J.; Huesgen, P. F.; Müller, R.-U.; Rinschen, M. M., The proteome microenvironment determines the protective effect of preconditioning in cisplatin-induced acute kidney injury. *Kidney Int.* **2019**, *95*, (2), 333-349.
17. Cochrane, J.; Williams, B. T.; Banerjee, A.; Harken, A. H.; Burke, T. J.; Cairns, C. B.; Shapiro, J. I., Ischemic preconditioning attenuates functional, metabolic, and morphologic injury from ischemic acute renal failure in the rat. *Ren. Fail.* **1999**, *21*, (2), 135-45.
18. Mitchell, J. R.; Verweij, M.; Brand, K.; van de Ven, M.; Goemaere, N.; van den Engel, S.; Chu, T.; Forrer, F.; Muller, C.; de Jong, M.; van, I. W.; JN, I. J.; Hoeijmakers, J. H.; de Bruin, R. W., Short-term dietary restriction and fasting precondition against ischemia reperfusion injury in mice. *Aging Cell* **2010**, *9*, (1), 40-53.
19. Bernhardt, W. M.; Campean, V.; Kany, S.; Jurgensen, J. S.; Weidemann, A.; Warnecke, C.; Arend, M.; Klaus, S.; Gunzler, V.; Amann, K.; Willam, C.; Wiesener, M. S.; Eckardt, K. U., Preconditional activation of hypoxia-inducible factors ameliorates ischemic acute renal failure. *J. Am. Soc. Nephrol.* **2006**, *17*, (7), 1970-8.
20. Yang, C. C.; Lin, L. C.; Wu, M. S.; Chien, C. T.; Lai, M. K., Repetitive hypoxic preconditioning attenuates renal ischemia/reperfusion induced oxidative injury via upregulating HIF-1 alpha-dependent bcl-2 signaling. *Transplantation* **2009**, *88*, (11), 1251-60.
21. Johnsen, M.; Kubacki, T.; Yeroslaviz, A.; Späth, M. R.; Mörsdorf, J.; Göbel, H.; Bohl, K.; Ignarski, M.; Meharg, C.; Habermann, B.; Altmüller, J.; Beyer, A.; Benzing, T.; Schermer, B.; Burst, V.; Müller, R.-U., The Integrated RNA Landscape of Renal Preconditioning against Ischemia-Reperfusion Injury. *J. Am. Soc. Nephrol.* **2020**, ASN.2019050534.
22. Mattison, J. A.; Roth, G. S.; Beasley, T. M.; Tilmont, E. M.; Handy, A. M.; Herbert, R. L.; Longo, D. L.; Allison, D. B.; Young, J. E.; Bryant, M.; Barnard, D.; Ward, W. F.; Qi, W.; Ingram, D. K.; de Cabo, R., Impact of caloric restriction on health and survival in rhesus monkeys from the NIA study. *Nature* **2012**, *489*, (7415), 318-21.

23. Lucanic, M.; Held, J. M.; Vantipalli, M. C.; Klang, I. M.; Graham, J. B.; Gibson, B. W.; Lithgow, G. J.; Gill, M. S., N-acylethanolamine signalling mediates the effect of diet on lifespan in *Caenorhabditis elegans*. *Nature* **2011**, 473, (7346), 226-9.
24. Chua, J. T.; Argueta, D. A.; DiPatrizio, N. V.; Kovesdy, C. P.; Vaziri, N. D.; Kalantar-Zadeh, K.; Moradi, H., Endocannabinoid System and the Kidneys: From Renal Physiology to Injury and Disease. *Cannabis and cannabinoid research* **2019**, 4, (1), 10-20.
25. Silva, G. B.; Atchison, D. K.; Juncos, L. I.; Garcia, N. H., Anandamide inhibits transport-related oxygen consumption in the loop of Henle by activating CB1 receptors. *Am. J. Physiol. Renal Physiol.* **2013**, 304, (4), F376-81.
26. Deutsch, D. G.; Goligorsky, M. S.; Schmid, P. C.; Krebsbach, R. J.; Schmid, H. H.; Das, S. K.; Dey, S. K.; Arreaza, G.; Thorup, C.; Stefano, G.; Moore, L. C., Production and physiological actions of anandamide in the vasculature of the rat kidney. *J. Clin. Invest.* **1997**, 100, (6), 1538-46.
27. Ritter, J. K.; Li, G.; Xia, M.; Boini, K., Anandamide and its metabolites: what are their roles in the kidney? *Front Biosci (Schol Ed)* **2016**, 8, 264-77.
28. Grundmann, F.; Muller, R. U.; Reppenhorst, A.; Hulswitt, L.; Spath, M. R.; Kubacki, T.; Scherner, M.; Faust, M.; Becker, I.; Wahlers, T.; Schermer, B.; Benzing, T.; Burst, V., Preoperative Short-Term Calorie Restriction for Prevention of Acute Kidney Injury After Cardiac Surgery: A Randomized, Controlled, Open-Label, Pilot Trial. *J Am Heart Assoc* **2018**, 7, (6).
29. Johnsen, M.; Späth, M. R.; Denzel, M. S.; Göbel, H.; Kubacki, T.; Hoyer, K. J. R.; Hinze, Y.; Benzing, T.; Schermer, B.; Antebi, A.; Burst, V.; Müller, R.-U., Oral Supplementation of Glucosamine Fails to Alleviate Acute Kidney Injury in Renal Ischemia-Reperfusion Damage. *PLoS One* **2016**, 11, (8), e0161315.
30. Di Marzo, V.; Matias, I., Endocannabinoid control of food intake and energy balance. *Nat. Neurosci.* **2005**, 8, (5), 585-9.
31. Bermudez-Silva, F. J.; Viveros, M. P.; McPartland, J. M.; Rodriguez de Fonseca, F., The endocannabinoid system, eating behavior and energy homeostasis: the end or a new beginning? *Pharmacol. Biochem. Behav.* **2010**, 95, (4), 375-82.
32. Turcotte, C.; Chouinard, F.; Lefebvre, J. S.; Flamand, N., Regulation of inflammation by cannabinoids, the endocannabinoids 2-arachidonoyl-glycerol and arachidonoyl-ethanolamide, and their metabolites. *J. Leukoc. Biol.* **2015**, 97, (6), 1049-70.
33. King-Himmelreich, T. S.; Möser, C. V.; Wolters, M. C.; Schmetzer, J.; Möller, M.; Schreiber, Y.; Ferreirós, N.; Geisslinger, G.; Niederberger, E., AMP-activated kinase and the endogenous endocannabinoid system might contribute to antinociceptive effects of prolonged moderate caloric restriction in mice. *Mol. Pain* **2017**, 13, 1744806917703111.
34. McDougle, D. R.; Watson, J. E.; Abdeen, A. A.; Adili, R.; Caputo, M. P.; Krapf, J. E.; Johnson, R. W.; Kilian, K. A.; Holinstat, M.; Das, A., Anti-inflammatory ω -3 endocannabinoid epoxides. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, 114, (30), E6034-e6043.
35. Devane, W. A.; Hanus, L.; Breuer, A.; Pertwee, R. G.; Stevenson, L. A.; Griffin, G.; Gibson, D.; Mandelbaum, A.; Etinger, A.; Mechoulam, R., Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* **1992**, 258, (5090), 1946-9.
36. Schmid, H. H.; Schmid, P. C.; Natarajan, V., N-acylated glycerophospholipids and their derivatives. *Prog. Lipid Res.* **1990**, 29, (1), 1-43.

37. Di Marzo, V.; Fontana, A.; Cadas, H.; Schinelli, S.; Cimino, G.; Schwartz, J. C.; Piomelli, D., Formation and inactivation of endogenous cannabinoid anandamide in central neurons. *Nature* **1994**, 372, (6507), 686-91.
38. Piomelli, D., The molecular logic of endocannabinoid signalling. *Nat. Rev. Neurosci.* **2003**, 4, (11), 873-84.
39. Elphick, M. R.; Egertová, M., The phylogenetic distribution and evolutionary origins of endocannabinoid signalling. *Handb. Exp. Pharmacol.* **2005**, (168), 283-97.
40. Long, J. Z.; LaCava, M.; Jin, X.; Cravatt, B. F., An anatomical and temporal portrait of physiological substrates for fatty acid amide hydrolase. *J. Lipid Res.* **2011**, 52, (2), 337-44.
41. Ritter, J. K.; Li, C.; Xia, M.; Poklis, J. L.; Lichtman, A. H.; Abdullah, R. A.; Dewey, W. L.; Li, P. L., Production and actions of the anandamide metabolite prostamide E2 in the renal medulla. *J. Pharmacol. Exp. Ther.* **2012**, 342, (3), 770-9.
42. Zygmunt, P. M.; Petersson, J.; Andersson, D. A.; Chuang, H.; Sørsgård, M.; Di Marzo, V.; Julius, D.; Högestätt, E. D., Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide. *Nature* **1999**, 400, (6743), 452-7.
43. Larrinaga, G.; Varona, A.; Perez, I.; Sanz, B.; Ugalde, A.; Candenás, M. L.; Pinto, F. M.; Gil, J.; Lopez, J. I., Expression of cannabinoid receptors in human kidney. *Histol. Histopathol.* **2010**, 25, (9), 1133-8.
44. Barutta, F.; Piscitelli, F.; Pinach, S.; Bruno, G.; Gambino, R.; Rastaldi, M. P.; Salvidio, G.; Di Marzo, V.; Cavallo Perin, P.; Gruden, G., Protective role of cannabinoid receptor type 2 in a mouse model of diabetic nephropathy. *Diabetes* **2011**, 60, (9), 2386-96.
45. Jenkin, K. A.; McAinch, A. J.; Grinfeld, E.; Hryciw, D. H., Role for cannabinoid receptors in human proximal tubular hypertrophy. *Cell. Physiol. Biochem.* **2010**, 26, (6), 879-86.
46. Howlett, A. C., The cannabinoid receptors. *Prostaglandins Other Lipid Mediat.* **2002**, 68-69, 619-31.
47. Afsar, B.; Afsar, R. E.; Copur, S.; Sag, A. A.; Ortiz, A.; Kanbay, M., The effect of energy restriction on development and progression of chronic kidney disease: review of the current evidence. *Br. J. Nutr.* **2020**, 1-14.
48. Rojas-Morales, P.; Tapia, E.; Leon-Contreras, J. C.; Gonzalez-Reyes, S.; Jimenez-Osorio, A. S.; Trujillo, J.; Pavon, N.; Granados-Pineda, J.; Hernandez-Pando, R.; Sanchez-Lozada, L. G.; Osorio-Alonso, H.; Pedraza-Chaverri, J., Mechanisms of Fasting-Mediated Protection against Renal Injury and Fibrosis Development after Ischemic Acute Kidney Injury. *Biomolecules* **2019**, 9, (9).
49. Di Marzo, V.; Stella, N.; Zimmer, A., Endocannabinoid signalling and the deteriorating brain. *Nature reviews. Neuroscience* **2015**, 16, (1), 30-42.
50. Garza-Lombó, C.; Gonsebatt, M. E., Mammalian Target of Rapamycin: Its Role in Early Neural Development and in Adult and Aged Brain Function. *Front. Cell. Neurosci.* **2016**, 10, (157).
51. Udi, S.; Hinden, L.; Earley, B.; Drori, A.; Reuveni, N.; Hadar, R.; Cinar, R.; Nemirovski, A.; Tam, J., Proximal Tubular Cannabinoid-1 Receptor Regulates Obesity-Induced CKD. *J. Am. Soc. Nephrol.* **2017**, 28, (12), 3518-3532.
52. Liu, J.; Godlewski, G.; Jourdan, T.; Liu, Z.; Cinar, R.; Xiong, K.; Kunos, G., Cannabinoid-1 Receptor Antagonism Improves Glycemic Control and Increases Energy Expenditure Through Sirtuin-1/Mechanistic Target of Rapamycin Complex 2 and 5'Adenosine Monophosphate-Activated Protein Kinase Signaling. *Hepatology* **2019**, 69, (4), 1535-1548.

53. Mukhopadhyay, P.; Pan, H.; Rajesh, M.; Batkai, S.; Patel, V.; Harvey-White, J.; Mukhopadhyay, B.; Hasko, G.; Gao, B.; Mackie, K.; Pacher, P., CB1 cannabinoid receptors promote oxidative/nitrosative stress, inflammation and cell death in a murine nephropathy model. *Br. J. Pharmacol.* **2010**, 160, (3), 657-68.
54. Cakir, M.; Tekin, S.; Doganyigit, Z.; Cakan, P.; Kaymak, E., The protective effect of cannabinoid type 2 receptor activation on renal ischemia-reperfusion injury. *Mol. Cell. Biochem.* **2019**.
55. Feizi, A.; Jafari, M. R.; Hamedivafa, F.; Tabrizian, P.; Djahanguiri, B., The preventive effect of cannabinoids on reperfusion-induced ischemia of mouse kidney. *Exp. Toxicol. Pathol.* **2008**, 60, (4-5), 405-10.
56. Fouad, A. A.; Al-Mulhim, A. S.; Jresat, I., Cannabidiol treatment ameliorates ischemia/reperfusion renal injury in rats. *Life Sci.* **2012**, 91, (7-8), 284-92.
57. Moradi, H.; Oveisi, F.; Khanifar, E.; Moreno-Sanz, G.; Vaziri, N. D.; Piomelli, D., Increased Renal 2-Arachidonoylglycerol Level Is Associated with Improved Renal Function in a Mouse Model of Acute Kidney Injury. *Cannabis Cannabinoid Res* **2016**, 1, (1), 218-228.
58. Engeli, S.; Böhnke, J.; Feldpausch, M.; Gorzelniak, K.; Janke, J.; Bátkai, S.; Pacher, P.; Harvey-White, J.; Luft, F. C.; Sharma, A. M.; Jordan, J., Activation of the peripheral endocannabinoid system in human obesity. *Diabetes* **2005**, 54, (10), 2838-43.
59. Engeli, S.; Heusser, K.; Janke, J.; Gorzelniak, K.; Bátkai, S.; Pacher, P.; Harvey-White, J.; Luft, F. C.; Jordan, J., Peripheral endocannabinoid system activity in patients treated with sibutramine. *Obesity (Silver Spring)* **2008**, 16, (5), 1135-7.
60. van Eyk, H. J.; van Schinkel, L. D.; Kantae, V.; Dronkers, C. E. A.; Westenberg, J. J. M.; de Roos, A.; Lamb, H. J.; Jukema, J. W.; Harms, A. C.; Hankemeier, T.; van der Stelt, M.; Jazet, I. M.; Rensen, P. C. N.; Smit, J. W. A., Caloric restriction lowers endocannabinoid tonus and improves cardiac function in type 2 diabetes. *Nutr. Diabetes* **2018**, 8, (1), 6.
61. Di Marzo, V.; Côté, M.; Matias, I.; Lemieux, I.; Arsenault, B. J.; Cartier, A.; Piscitelli, F.; Petrosino, S.; Alméras, N.; Després, J. P., Changes in plasma endocannabinoid levels in viscerally obese men following a 1 year lifestyle modification programme and waist circumference reduction: associations with changes in metabolic risk factors. *Diabetologia* **2009**, 52, (2), 213-7.
62. Quercioli, A.; Montecucco, F.; Pataky, Z.; Thomas, A.; Ambrosio, G.; Staub, C.; Di Marzo, V.; Ratib, O.; Mach, F.; Golay, A.; Schindler, T. H., Improvement in coronary circulatory function in morbidly obese individuals after gastric bypass-induced weight loss: relation to alterations in endocannabinoids and adipocytokines. *Eur. Heart J.* **2013**, 34, (27), 2063-73.
63. Hanus, L.; Avraham, Y.; Ben-Shushan, D.; Zolotarev, O.; Berry, E. M.; Mechoulam, R., Short-term fasting and prolonged semistarvation have opposite effects on 2-AG levels in mouse brain. *Brain Res.* **2003**, 983, (1-2), 144-51.
64. Kirkham, T. C.; Williams, C. M.; Fezza, F.; Di Marzo, V., Endocannabinoid levels in rat limbic forebrain and hypothalamus in relation to fasting, feeding and satiation: stimulation of eating by 2-arachidonoyl glycerol. *Br. J. Pharmacol.* **2002**, 136, (4), 550-7.
65. Patschan, D.; Hildebrandt, A.; Rinneburger, J.; Wessels, J. T.; Patschan, S.; Becker, J. U.; Henze, E.; Krüger, A.; Müller, G. A., The hormone melatonin stimulates renoprotective effects of "early outgrowth" endothelial progenitor cells in acute ischemic kidney injury. *American Journal of Physiology-Renal Physiology* **2012**, 302, (10), F1305-F1312.

