

# Physical activity and inhibitory control: The mediating role of sleep quality and sleep efficiency

Lin Li <sup>1,2,#</sup>, Qian Yu <sup>3,#</sup>, Wen-rui Zhao <sup>1,2</sup>, Fabian Herold <sup>4,5</sup>, Boris Cheval <sup>6,7</sup>, Zhaowei Kong <sup>8</sup>, Jinming Li <sup>3</sup>, Notger Mueller <sup>4,5</sup>, Arthur F. Kramer <sup>9,10</sup>, Jie Cui <sup>1,2</sup>, Hua-wei Pan <sup>1,2</sup>, Zhu-xuan Zhan <sup>1,2</sup>, Min-qiang Hui <sup>1,2</sup> and Liye Zou <sup>3\*</sup>

1. Key Laboratory Ministry of education of adolescent health assessment and exercise intervention, East China Normal University, Shang Hai 200241, China; lilin.xtt@163.com (LL); zhaowenrui108@163.com (WRZ); 1375515922@qq.com (JC); 13758114178@163.com (HWP); zzx951105@163.com (ZXZ); 893291859@qq.com (MQH);
2. School of physical education and health, East China Normal University, Shang Hai 200241, China;
3. Exercise & Mental Health Laboratory; Institute of KEEP Collaborative Innovation, School of Psychology, Shenzhen University, Shenzhen, 518060, China; yuqianmiss@163.com (QY); jinmingli1999@gmail.com (LJM); liyezou123@gmail.com (LZ);
4. Department of Neurology, Medical Faculty, Otto von Guericke University, Leipziger Str. 44, 39120 Magdeburg, Germany; fabian.herold@st.ovgu.de (FH); notger.mueller@dzne.de (NM);
5. Research Group Neuroprotection, German Center for Neurodegenerative Diseases (DZNE), Leipziger Str. 44, 39120 Magdeburg, Germany;
6. Swiss Center for Affective Sciences, University of Geneva, Geneva, Switzerland; Boris.Cheval@unige.ch (BC);
7. Laboratory for the Study of Emotion Elicitation and Expression (E3Lab), Department of Psychology, FPSE, University of Geneva, Geneva, Switzerland;
8. Faculty of Education, University of Macau, Macao, China; [zwkong@um.edu.mo](mailto:zwkong@um.edu.mo) (ZWK);
9. Center for Cognitive and Brain Health, Department of Psychology, Northeastern University, Boston, MA 02115, USA. Email: [a.kramer@northeastern.edu](mailto:a.kramer@northeastern.edu) (KA)
10. Beckman Institute, University of Illinois at Urbana-Champaign, Champaign, IL 61801, USA.

# Equally contributed to this study.

\* Co-correspondence: [liyezou123@gmail.com](mailto:liyezou123@gmail.com) (LYZ)

**Abstract: Objectives.** The current study aimed to investigate the relationship between physical activity (PA) level and inhibitory control performance and then determine whether this association was mediated by multiple sleep parameters (i.e., subjective sleep quality, sleep duration, sleep efficiency, sleep disturbance). **Methods.** 180 healthy university students (age:  $20.15 \pm 1.92$  years) from the East China Normal University were recruited in the present study. PA level, sleep parameters, and inhibitory control performance were assessed using the International Physical Activity Questionnaire (IPAQ), the Pittsburgh Sleep Quality Index Scale (PSQI), and a Stroop test, respectively. Data were analyzed using structural equation modeling. **Results.** A higher level of PA was linked to better cognitive performance. Furthermore, higher subjective sleep quality and sleep efficiency were associated with better inhibitory control performance. The mediation analysis revealed that subjective sleep quality and sleep efficiency mediated the relationship between PA level and inhibitory control performance. **Conclusion.** Our results are in accordance with the literature and buttress the idea that a healthy lifestyle that involves a relatively high level of regular PA and adequate sleep patterns is beneficial for cognition (e.g., inhibitory control performance). Furthermore, our study adds to the literature that sleep quality and sleep efficiency mediates the relationship of PA and inhibitory control performance expanding our knowledge in the field of exercise-cognition.

**Keywords:** physical activity; sleep; inhibitory performance; mediating effects

## Introduction

Brain health and cognitive enhancement being important factors for a successful study, better career opportunity and a higher quality of life and thus has received widespread attention from different disciplines including medicine and education [1]. Emerging evidence demonstrates that regular physical activity (PA) can enhance cognitive performance, especially in the domain of executive functioning (EF) [2-5]. Among the diverse subcomponents of EF, controlled inhibition (the capacity to suppress irrelevant information or prepotent responses in monitoring and updating information) has been reported to benefit more from PA than the other aspects of EF (e.g., shifting or updating) [6]. Considering that brain develops until the age of 30 years [7-10] and the important role of inhibitory control to master life successfully [11], there is an urgency to investigate how regular PA influences the inhibitory control performance and how to maximize PA-induced cognitive benefits among the young ages. Of note, existing researches mostly focused on children and older individuals, while there is a lack of studies on young to middle-aged adults [1]. Additionally, although PA has been proven to maintain and improve neurocognitive function, much less is known about the pathway by which it exerts salutary effects on inhibitory control performance. Among the suggested potential mediators, sleep has been proposed as an ideal candidate to explain the PA-cognition links, although empirical evidence in younger adults is currently lacking [12].

The improved cognitive performance among individuals who engaged in regular PA may be explained, at least partly, by sleep health. Indeed, young adults regularly experience restricted sleep (70% people sleeping less than 6 hours at least one night a week) due to educational, vocational and social responsibilities [13]. Mounting evidence shows that the increased PA contributes to sleep health (i.e., sleep duration, efficiency, and quality). For example [14, 15], a randomized controlled trials conducted in 1997 first indicated the PA (i.e., moderate-intensity exercise, daily activity)-induced benefits in self-reported sleep quality. A review by Vanderlinden et al. indicated that moderate intensity exercise intervention (lasted 12 weeks to 6 months), with a frequency of three times per week, showed the highest improvements in sleep outcomes in older adults [16]. Even among patients with sleep disruptions, exercise intervention was also found to positively influence the sleep outcomes [17].

The provided evidence support the idea that better sleep was linked to improved cognitive performance [20-22]. For example, sufficient sleep quality has been shown to be essential for an optimal cognitive performance, especially in the domain of EF (i.e., inhibitory control) [23]. However, although the exact neurobiological mechanisms of the association of sleep and cognitive performance are not fully understood. Sleep might work through complex ultradian, circadian and homeostatic regulations to positively influence the prefrontal circuits which are known to be important neural correlates for EF [24-26]. The improved prefrontal circuits could be well reflected by higher voltage and slower brain waves during non-rapid eye movement (NREM) coinciding with the deactivation of dorsolateral prefrontal cortex during rapid eye movement (REM) [27].

To the best of our knowledge, there are only two studies examining the potential mediating role of sleep health on the association between PA and cognitive function [28-29]. Wilckens et al. found that sleep efficiency may be one pathway by which PA benefited executive control, but this study only investigated two sleep health indicators (sleep efficiency and total sleep time as measured by accelerometer-based sleep assessment) [28]. Contrary to outcomes of Wilckens et al, Falck et al. reported that PA and sleep quality interacted with cognitive function via independent mechanisms without mediation effects [29] among the elderly (above 55 years). To address the gap and inconsistency, the current study attempted to explore how PA influences young adults' inhibitory control via wider range of sleep parameters [30]. We aimed to: (i) investigate the relationships among regular PA level, inhibitory control and multiple sleep parameters (subjective sleep quality, sleep duration, sleep efficiency, sleep disturbance) among the university students; (ii) examine how sleep parameters mediate the relationship between PA level and inhibitory control performance.

## Methods

### *Participants*

In total, 180 healthy university students (age:  $20.15 \pm 1.92$  years ; 109 females, 71 males) from the East China Normal University were recruited in the present study. PA level of participants, sleep status, inhibitory control performance was assessed via the International Physical Activity Questionnaire (IPAQ), the Pittsburgh Sleep Quality Index Scale (PSQI), and Stroop test, respectively. Moreover, all participants were screened for depression and anxiety by the Self-rating depression scale (SDS) and the Self-Rating Anxiety Scale (SAS). Exclusion criteria included: (1) SDS score  $\geq 53$ ; (2) SAS score  $\geq 50$ ; (3) failed to complete all the assessment; (4) left-handedness. The whole procedure was approved by the ethic committee of the East China Normal University (No. HR 085-2018) and all participants signed the informed consent before the onset of this experiment.

### *Demographic Information*

A self-developed scale on demographic information was used to acquire participants' information like gender, age, height and weight. Height and weight were assessed by Height Weight Meter. Body Mass Index (BMI) was calculated as weight (in kilograms) divided by the square of height (in metres).

### *International Physical Activity Questionnaire*

The IPAQ was used to assess the PA level of each individual (content validity = 0.98, calibration validity = 0.41, test-retest reliability = 0.78). The IPAQ totally consists of five parts: (1) job-related PA; (2) transportation PA; (3) housework, house maintenance, and caring for family; (4) recreation, sport, and leisure-time PA; and (5) time spent sitting. MET (Metabolic Equivalent of Task) minutes per week, as a continuous variable, represents the amount of energy expended carrying out physical activity.

### *Pittsburgh Sleep Quality Index Scale*


The subjective level of sleep quality was assessed using the PSQI which queries possible factors related to sleep disturbance over the past four weeks[31]. It includes 19 items within seven sleep-related domains: Subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleep medication, and daytime dysfunction[31]. The score of each domain is rated from 0 to 3 and seven sub-scores can be added to produce a total sleep quality score ranging from 0 to 21[31], with higher total score indicating lower sleep quality[31].

### *Measurement of Inhibitory Control Performance*

Inhibitory control performance was tested by a Stroop task in which participants were asked to differentiate the stimulation in the mixed congruent and incongruent conditions[32]. The stimuli used in this task were composed of four-color words (red, yellow, blue, and green in Chinese). In the congruent condition, the word was shown in matching color; in the incongruent condition, the word did not match with the color. The whole task lasted for 240 s, including two blocks with 32 trials (8 congruent and 24 incongruent trials) in total. In each trial, a fixed cross was presented first to attract participants' attention and then the color words were shown randomly, with each one lasting for 2000 ms. The inter-stimulus interval were also random (2000, 4000, 6000, 8000, or 10000 ms). The participants were asked to report the correct color of word by pressing the corresponding button in the keyboard as quickly and accurately as possible, ignoring the interference brought by the word's semantic meaning. Subjects performed 20 practice trials before performing the experimental trial blocks. Accuracy and reaction time of each condition were also recorded. According to the operational definition of inhibitory performance in Stroop test, the efficiency of inhibitory performance was characterized by the difference of reaction time. The smaller the Stroop effect ( $\Delta RT = RT_{\text{incongruent}} - RT_{\text{congruent}}$ ), the stronger the inhibitory performance.

### Statistical Analysis

Statistical analyses were conducted using SPSS (Statistical Product and Service Solutions) software, version 23. Mean and standard deviation for all variables were calculated. Unless noted to the contrary, for all effects described as significant,  $p < .05$ . Partial correlation analyses, with sex, age and BMI as covariates, were used to examine the relationships between PA level (MET score), inhibitory control performance, and seven sleep parameters (subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleep medication, and daytime dysfunction). Moreover, a structural equation modeling (SEM) with the help of AMOS (Analysis of Moment Structure) was conducted to assess whether the sleep parameters mediated the association between PA and inhibitory control performance. The indices used to evaluate the overall fit of model included Chi-square ( $\chi^2$ ) test, normed Chi-square ( $\chi^2/df$ ), normed fit index (NFI), incremental fit index (IFI), and comparative fit index (CFI). Usually, a well-fitted model should meet the following requirements: (1) a non-significant Chi-square; (2) 0.90 or lesser NFI, IFI, CFI value; (3) 0.06 or lesser normed Chi-square value [33]. The paths from the mediation model were examined using direct and indirect effects of study variables (PA level, sleep parameters, and stroop effect); bootstrapping approach was used to test statistical significance of mediation.



**Figure 1.** Stroop Test.

## Results

The results of demographic and anthropometric parameters, the IPAQ, the PSQI, and performance in Stroop test are shown in **Table 1**.

**Table 1.** Demographic Information on Variables.

	Males (n=71)	Females (n=109)	Total (N=180)
<i>Demographic and Anthropometric Parameters</i>			
age (years)	20.31±1.45	19.68±1.64	19.93±1.59
BMI (kg/m <sup>2</sup> )	21.66±2.05	21.20±2.55	21.37±2.37
<i>International Physical Activity Questionnaire</i>			
MET-min/w	4967.05±1447.06	5171.22±1556.39	5090.69±1513.41
<i>Pittsburgh Sleep Quality Index</i>			
Subjective sleep quality	0.68±0.69	0.76±0.79	0.73±0.75
Sleep time	0.62±0.70	0.54±0.69	0.57±0.69
Sleep efficiency	0.73±0.72	0.68±0.76	0.70±0.74
Sleep disorders	0.80±0.71	0.97±0.55	0.91±0.62
<i>Stroop Test</i>			
Congruent accuracy (%)	95.00±9.35	97.00±5.80	96.00±7.46
Incongruent accuracy (%)	95.00±8.65	97.00±4.70	96.00±6.66
Congruent reaction time (ms)	899.98±107.20	867.53±135.39	875.47±141.42
Incongruent reaction time(ms)	1000.18±107.79	964.34±148.10	973.07±152.56
Stroop effect (ms)	100.21±45.64	96.81±47.13	98.15±46.45

**Note.** BMI: body mass index; MET: metabolic equivalent of task.

### Correlation analysis

As shown in **Table 2**, results indicate negative correlations of regular PA level with (1) subjective sleep quality ( $r = -0.42$ ,  $p < 0.001$ ), (2) habitual sleep efficiency ( $r = -0.52$ ,  $p < 0.001$ ), and (3) the Stroop effect ( $r = -0.39$ ,  $p < 0.001$ ) (higher score in the PSQI indicating worse sleep status; smaller Stroop effect indicating better inhibitory control performance). Additionally, the scores of subjective sleep quality and sleep efficiency were found to positively correlate with the Stroop effect ( $r = 0.83$ ,  $p < 0.001$ ;  $r = 0.61$ ,  $p < 0.001$ ).

Table 2. Correlation Analyses Among Outcome Measures.

	MET-min/w	S1	S2	S3	S4	S5	S6	S7	Acc.1	Acc.2	RT1	RT2	Stroop effect
MET-min/w	-												
S1	-0.425***	-											
S2	-0.033	0.068	-										
S3	-0.006	0.082	0.107	-									
S4	-0.519***	0.652***	0.033	0.126	-								
S5	0.180**	0.100	0.186**	-0.106	-0.008	-							
S6	0.153**	0.060	0.125	-0.210**	0.081	0.331***	-						
S7	0.058	0.047	0.289***	-0.039	0.002	0.198**	0.412***	-					
Acc.1	-0.018	0.057	0.064	0.074	-0.023	0.103	-0.019	0.136	-				
Acc.2	-0.056	-0.032	-0.016	0.116	-0.062	-0.002	-0.106	0.017	0.486***	-			
RT1	0.036	0.139	0.021	-0.053	0.043	0.047	0.116	0.033	-0.076	-0.107	-		
RT2	-0.101	0.419***	0.036	-0.006	0.252***	0.060	0.142	0.032	-0.027	-0.109	0.938***	-	
Stroop effect	-0.386***	0.831***	0.045	0.125	0.611***	0.045	0.098	0.003	0.125	-0.026	0.008	0.355***	-

**Note.** MET = metabolic equivalent of task; S1 = Subjective sleep quality; S2 = sleep latency; S3 = sleep duration; S4 = habitual sleep efficiency; S5 = sleep disturbances; S6 = use of sleep medication; S7 = daytime dysfunction. Acc.1 = congruent accuracy; Acc.2 = incongruent accuracy; RT1 = congruent reaction time; RT2 = incongruent reaction time.

### Multiple Mediation Model

Based on the outcomes of correlation analyses, only two sleep parameters (subjective sleep quality and sleep efficiency) were included in the mediation model. The structured equation model was used to determine the mediating role of sleep parameters on the relationship between PA and inhibitory control (Figure 2). In the initial model, RMSEA (0.07) and SRMR (0.06) were higher than 0.05, which meant that the initial model failed to present a good fit. Therefore, the model was modified as specified by modification indices ( $\chi^2/df = 64.43$ , NFI = 0.83, IFI = 0.83, and CFI = 0.82). After removing the non-significant path between physical activity and inhibitory control performance, the model was re-analyzed and all indices within the new model show a good fit ( $\chi^2/df = 0.06$ , NFI = 0.92, IFI = 0.92, and CFI = 0.93) (Table 3). In the final model, the mediating effect explained 76% of variance variation of dependent variable, subjective sleep quality and sleep efficiency explained 53.6% and 22.4% of variance variation respectively. In order to reduce the interaction between the mediating variables, residual correlation analysis between the two mediating variables was conducted (path coefficient = 0.55).


The paths from the mediation model were examined using direct and indirect effect of study variables (Table 4; Figure 2). Results showed the PA-induced indirect effect on inhibitory control performance while adjusting for sleep parameters among the university students. Thus, sleep parameters (subjective sleep quality, sleep efficiency) played a mediating role in the indirect relationships. Additionally, the non-significant direct effect of physical activity on inhibitory control performance is 0.001.

**Table 3.** Modification indices in Multiple mediation model 2

Model	$\chi^2$	<i>p</i>	<i>f</i>	$\chi^2 / DF$	NFI	IFI	CFI	RMSEA	SRMR
Initial model	64.43	0.001	1	64.43	0.83	0.83	0.82	0.07	0.06
Final model	0.06	0.06	1	0.06	0.92	0.92	0.93	0.05	0.04
$\Delta\chi^2$	64.37								

**Table 4.** The Bootstrap Analysis of the Multiple Mediation Model 2.

Route	Indirect Effects of standardization	Average Indirect Effect	95% Confidence Interval		<i>p</i>
			Lower Limit	Upper Limit	
Subjective Sleep Quality	(0.42) * (0.77) = 0.32	0.41	0.27	0.55	0.01
Sleep Efficiency	(0.53) * (0.13) = 0.07	0.51	0.39	0.63	0.01



**Figure 2.** The Multiple Mediation Model 2.



## Discussion

Based on previous literature, the mediating role of sleep parameters on the relationship between PA and EF (i.e., inhibitory control) is expected, but this test has not been formally assessed in a young sample and with a wider range of sleep health indicators [12, 34]. In the present study, we observed in our cohort of healthy younger adults (1) a negative correlation between PA and inhibitory control performance, (2) a negative correlation between PA and subjective sleep quality as well as sleep efficiency ( $[\text{number of sleep hours} / \text{number of hours spent in bed}] \times 100$ ), and (3) positive correlations between inhibitory control and sleep parameters (subjective sleep quality, sleep efficiency). Moreover, the mediation analysis reveals that subjective sleep quality and sleep efficiency play mediating roles in the PA-EF (inhibitory control).

In this study, positive correlations were observed between PA level and sleep indicators (subjective sleep quality and habitual sleep efficiency), which is in line with the findings of existing studies [28, 35]. Over the last decades, it has been shown that among younger adults, higher PA level is positively associated with sleep health [36, 37]. Notably, recent studies proposed that PA-sleep relationship may be influenced by PA slots [38]. To be specific, some researchers emphasized that light PA in the daytime was correlated with better sleep status, whereas PA at night may reduce the sleep duration and sleep time window [39]. However, as we used the IPAQ to assess PA, the time of day in PA engagement was not in the present study but such an assessment should be considered in future studies to deepen our understanding of the relationship between PA, sleep, and cognitive performance.

Results from partial correlation analysis indicate that EF (inhibitory control) was positively correlated with both subjective sleep quality and sleep efficiency but not linked to sleep duration. Such results are consistent with previous researches in which sleep parameters were measured with actigraphic device and polysomnography [40-42]. PA-sleep relationships may be partially explained by following neurobiological mechanisms: (1) Higher sleep efficiency and sleep quality enhanced the efficacy of slow-wave sleep in restoration of prefrontal cortex function (which is responsible for inhibitory control performance) [43]; (2) Prior sleep deprivation, sleep disorders, sleep fragmentation and medication effects usually occur along with the longer sleep duration which does not reflect real sleep health [44-46]. Non-restorative sleep caused by recurrent awakenings and stage shifts can even be observed in the population with normal sleep duration [47]. Due to this, sleep duration cannot be regarded as a reliable indicator for sleep health.

Only an indirect relationship (mediating by sleep parameters) rather than a direct relationship between PA and cognitive performance was observed among university students in the present study. This finding is partially supported Wilckens et al., who found that PA level was indirectly correlated with inhibitory control in both adolescent (aged 21-30) and older adults (aged 55-80) and the sleep efficiency, but not total sleep time or sleep duration, played the mediating role. In the present study, we observed that both sleep efficiency and subjective sleep quality played an important role in the PA-EF relationship. Based on these outcomes, it could be inferred that ease of falling asleep and staying asleep rather than any aspect of sleep work on the PA-cognition connections. Moreover, it is essential to highlight that the direct relationship between PA and inhibitory control performance reflects the overall relationship with cognition, which consists of sleep parameters and other mediators not tested here. And some of the potential mediators may suppress the association between PA and inhibitory control [48, 49].

Besides the primary outcome carried out from the present study, some novel findings were summarized based on comparison with previous researches [28, 29, 50]. It is suggested that the mediating role of sleep efficiency in the PA-EF (inhibitory control) relationship is observed across different age groups [28]. PA can even attenuate the negative impact of low sleep efficiency on executive function, with the clearest effects observed using direct measurements of sleep and PA [50]. Even though sleep quality and sleep duration were also found to positively associate with PA and cognition among the young adults [35], apparent functional weakening in such relationships appear in the aging population [29]. It seems that sleep measures like sleep quality and sleep duration are more sensitive to aging-related neurobiology (e.g., neural atrophy, nocturnal hypoxia,



neuroendocrine changes, and altered neuromodulation), which may reduce the potential to impact cognitive performance through strategies such as PA[51].

Finally, this study has certain theoretical and practical significance, but some limitations must be acknowledged when interpreting our findings. First, the cross-sectional design does not allow to make strong conclusions regarding the causality of the observed correlations. Secondly, this study did not account for the time of day when participants engaged in PA, which is known to influence sleep patterns and thus might influence cognitive performance, too. Thirdly, our sleep pattern assessment relies on the PSQI which is an international recognized questionnaire with good psychometric properties but relies on subjective ratings rather than objective ratings. Given difference in subjective and objective ratings of sleep [52, 53], this might have influenced our analysis. Thus, caution should be taken when comparing our results with findings of other studies quantifying sleep parameters by objective tools (e.g., actigraphy) although those study points in the same direction (better sleep patterns are associated with better cognitive performance) [54]. In this regard, we recommend that further studies investigating the relationship between regular PA level, sleep patterns, and cognitive performance, should consider to assess both subjective ratings and objective ratings of sleep. For studies with more stern requirements, polysomnography, the gold standard for sleep research is recommended. Lastly, we focused only on one cognitive parameter (inhibitory control), preventing to generalize our results to other EF components and cognitive measures.

## Conclusion

Sleep efficiency and subjective sleep quality were found to statistically mediate the significant relationship between physical activity and inhibitory control in the population of university students. Joint measurements including both objective and subjective assessment and participants in wider age range should be considered in the further studies to test the potential neurobehavioral mechanisms and moderators.

## Reference

1. Erickson KI, Hillman C, Stillman CM, Ballard RM, Bloodgood B, Conroy DE, et al.(2019). Physical activity, cognition, and brain outcomes: a review of the 2018 physical activity guidelines. 51,1242.
2. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee I-M, et al.(2011). American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. 43,1334-1359.
3. Verburgh L, Königs M, Scherder EJ, Oosterlaan JJ *Bjosc*m.(2014). Physical exercise and executive functions in preadolescent children, adolescents and young adults: a meta-analysis. 48,973-979.
4. Li JW, O'Connor H, O'Dwyer N, Orr RJ *Jos, sport mi*.(2017). The effect of acute and chronic exercise on cognitive function and academic performance in adolescents: A systematic review. 20,841-848.
5. Xue Y, Yang Y, Huang TJ *Bjosc*m.(2019). Effects of chronic exercise interventions on executive function among children and adolescents: a systematic review with meta-analysis. 53,1397-1404.
6. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD *JCp*.(2000). The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: A latent variable analysis. 41,49-100.
7. Lenroot RK, Giedd JN *JN*, reviews b.(2006). Brain development in children and adolescents: insights from anatomical magnetic resonance imaging. 30,718-729.
8. Whitford TJ, Rennie CJ, Grieve SM, Clark CR, Gordon E, Williams LM *JHbm*.(2007). Brain maturation in adolescence: concurrent changes in neuroanatomy and neurophysiology. 28,228-237.
9. Lebel C, Walker L, Leemans A, Phillips L, Beaulieu CJN.(2008). Microstructural maturation of the human brain from childhood to adulthood. 40,1044-1055.
10. Tamnes CK, Herting MM, Goddings A-L, Meuwese R, Blakemore S-J, Dahl RE, et al.(2017). Development of the cerebral cortex across adolescence: a multisample study of inter-related longitudinal changes in cortical volume, surface area, and thickness. 37,3402-3412.

11. Diamond A, Lee KJS.(2011). Interventions shown to aid executive function development in children 4 to 12 years old. 333,959-964.
12. Stillman CM, Cohen J, Lehman ME, Erickson KIJFihh.(2016). Mediators of physical activity on neurocognitive function: a review at multiple levels of analysis. 10,626.
13. Wong ML, Lau EYY, Wan JHY, Cheung SF, Hui CH, Mok DSYJJopr.(2013). The interplay between sleep and mood in predicting academic functioning, physical health and psychological health: A longitudinal study. 74,271-277.
14. King AC, Oman RF, Brassington GS, Bliwise DL, Haskell WLJJ.(1997). Moderate-intensity exercise and self-rated quality of sleep in older adults: a randomized controlled trial. 277,32-37.
15. Singh NJS.(1997). Sleep, sleep deprivation, and daytime activities. 20,95-101.
16. Vanderlinden J, Boen F, van Uffelen JIJJoBN, Activity P.(2020). Effects of physical activity programs on sleep outcomes in older adults: a systematic review. 17,11.
17. Lins-Filho OL, Pedrosa RP, Gomes JM, Moraes SLD, Vasconcelos BCE, Lemos CAA, et al.(2020). Effect of exercise training on subjective parameters in patients with obstructive sleep apnea: a systematic review and meta-analysis. 69,1-7.
18. Lang C, Kalak N, Brand S, Holsboer-Trachsler E, Pühse U, Gerber MJSmr.(2016). The relationship between physical activity and sleep from mid adolescence to early adulthood. A systematic review of methodological approaches and meta-analysis. 28,32-45.
19. Atoui S, Chevance G, Romain A-J, Kingsbury C, Lachance J-P, Bernard PJSmr.(2021). Daily associations between sleep and physical activity: a systematic review and meta-analysis.101426.
20. Lo JC, Groeger JA, Cheng GH, Dijk D-J, Chee MWJSm.(2016). Self-reported sleep duration and cognitive performance in older adults: a systematic review and meta-analysis. 17,87-98.
21. Naismith SL, Mowszowski LJ Coip.(2018). Sleep disturbance in mild cognitive impairment: a systematic review of recent findings. 31,153-159.
22. Smithies TD, Toth AJ, Dunican IC, Caldwell JA, Kowal M, Campbell MJJS.(2021). The Effect of Sleep Restriction on Cognitive Performance in Elite Cognitive Performers: A Systematic Review.
23. Benitez A, Gunstad JJTCN.(2012). Poor sleep quality diminishes cognitive functioning independent of depression and anxiety in healthy young adults. 26,214-223.
24. Horn JBJP.(1993). Human sleep, sleep loss and behavior. Implications for the prefrontal cortex and psychiatric behavior. 162,413-419.
25. Harrison Y, Horne JJJosr.(1998). Sleep loss impairs short and novel language tasks having a prefrontal focus. 7,95-100.
26. Harrison Y, Horne JA, Rothwell AJS.(2000). Prefrontal neuropsychological effects of sleep deprivation in young adults--a model for healthy aging? 23,1067-1073.
27. Muzur A, Pace-Schott EF, Hobson JAJTics.(2002). The prefrontal cortex in sleep. 6,475-481.
28. Wilckens KA, Erickson KI, Wheeler MEJBsm.(2018). Physical activity and cognition: a mediating role of efficient sleep. 16,569-586.
29. Falck RS, Best JR, Davis JC, Liu-Ambrose TJJoAsD.(2018). The independent associations of physical activity and sleep with cognitive function in older adults. 63,1469-1484.
30. Cheval B, Maltagliati S, Sieber S, Beran D, Chalabaev A, Sander D, et al.(2021). Why are individuals with diabetes less active? The mediating role of physical, emotional, and cognitive factors.
31. Buysse DJ, Reynolds III CF, Monk TH, Berman SR, Kupfer DJJPr.(1989). The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. 28,193-213.
32. Cui J, Zou L, Herold F, Yu Q, Jiao C, Zhang Y, et al.(2020). Does Cardiorespiratory Fitness Influence the Effect of Acute Aerobic Exercise on Executive Function? 14.
33. Hu Lt, Bentler PM.(1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6,1-55. 10.1080/10705519909540118.
34. Stillman CM, Esteban-Cornejo I, Brown B, Bender CM, Erickson KI.(2020). Effects of Exercise on Brain and Cognition Across Age Groups and Health States. *Trends in neurosciences*, 43,533-543. 10.1016/j.tins.2020.04.010.
35. Youngstedt SDJCism.(2005). Effects of exercise on sleep. 24,355-365.

36. Kalak N, Gerber M, Kirov R, Mikoteit T, Yordanova J, Pühse U, et al.(2012). Daily morning running for 3 weeks improved sleep and psychological functioning in healthy adolescents compared with controls. 51,615-622.
37. Lang C, Brand S, Feldmeth AK, Holsboer-Trachsler E, Pühse U, Gerber MJP, et al.(2013). Increased self-reported and objectively assessed physical activity predict sleep quality among adolescents. 120,46-53.
38. Kline CEJAjlm.(2014). The bidirectional relationship between exercise and sleep: implications for exercise adherence and sleep improvement. 8,375-379.
39. Wendt A, da Silva ICM, Gonçalves H, Menezes A, Barros F, Wehrmeister FCJJoS, et al.(2020). Short-term effect of physical activity on sleep health: A population-based study using accelerometry.
40. Bastien CH, Fortier-Brochu E, Rioux I, LeBlanc M, Daley M, Morin CMJJopr.(2003). Cognitive performance and sleep quality in the elderly suffering from chronic insomnia: relationship between objective and subjective measures. 54,39-49.
41. Foley DJ, Masaki K, White L, Larkin EK, Monjan A, Redline SJS.(2003). Sleep-disordered breathing and cognitive impairment in elderly Japanese-American men. 26,596-599.
42. Blackwell T, Yaffe K, Ancoli-Israel S, Schneider JL, Cauley JA, Hillier TA, et al.(2006). Poor sleep is associated with impaired cognitive function in older women: the study of osteoporotic fractures. 61,405-410.
43. Wilckens KA, Erickson KI, Wheeler MEJNp.(2012). Age-related decline in controlled retrieval: the role of the PFC and sleep. 2012.
44. Harrison Y, Horne JJP.(1996). Long-term extension to sleep-Are we really chronically sleep deprived? 33,22-30.
45. Youngstedt SD, Kripke DFJSmr.(2004). Long sleep and mortality: rationale for sleep restriction. 8,159-174.
46. Monk TH, Buysse DJ, Begley AE, Billy BD, Fletcher MEJCi.(2009). Effects of a two-hour change in bedtime on the sleep of healthy seniors. 26,526-543.
47. Shrivastava D, Jung S, Saadat M, Sirohi R, Crewson KJJochimp.(2014). How to interpret the results of a sleep study. 4,24983.
48. Preacher KJ, Hayes AFJBrm.(2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. 40,879-891.
49. Zhao X, Lynch Jr JG, Chen QJJocr.(2010). Reconsidering Baron and Kenny: Myths and truths about mediation analysis. 37,197-206.
50. Lambiase MJ, Gabriel KP, Kuller LH, Matthews KAJJoGSABS, Sciences M.(2014). Sleep and executive function in older women: the moderating effect of physical activity. 69,1170-1176.
51. Scullin MK, Bliwise DLJPoPS.(2015). Sleep, cognition, and normal aging: integrating a half century of multidisciplinary research. 10,97-137.
52. Landry GJ, Best JR, Liu-Ambrose TJFian.(2015). Measuring sleep quality in older adults: a comparison using subjective and objective methods. 7,166.
53. Hughes JM, Song Y, Fung CH, Dzierzewski JM, Mitchell MN, Jouldjian S, et al.(2018). Measuring sleep in vulnerable older adults: a comparison of subjective and objective sleep measures. 41,145-157.
54. Okano K, Kaczmarzyk JR, Dave N, Gabrieli JD, Grossman JCJNsol.(2019). Sleep quality, duration, and consistency are associated with better academic performance in college students. 4,1-5.