

Article

Not peer-reviewed version

Space Theory

[S. Chen](#) *

Posted Date: 6 November 2024

doi: 10.20944/preprints202105.0197.v12

Keywords: space; gravitation



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Space Theory

S. Chen

Rheast LLC, 1331 Lamar St, Houston, TX 77010, USA; chen@rheast.com

Abstract: The author introduces a novel theoretical framework that suggests matter and energy are both converted from curved space. In Space Theory, gravitation is generated by the flow of space, instead of being transmitted by the graviton as in String Theory. This theory also suggests that Newton's gravitational constant, denoted as G , may not be truly constant but could vary over time. The equivalent equation of space is $S = Ec^2 = mc^4$, and the gravitational force formula is $S_{\mu\nu} = 4\pi Gm = (4/3)\pi((r + a)^3 - r^3)$. The Space Theory also predicts that the surface gravitational acceleration of the neutron star Crab Pulsar (PSR B0531+21) is approximately $8.21924883 \times 10^6 \text{ m/s}^2$.

Keywords: space; gravitation

Introduction

Newton's law of universal gravitation [1] provides the equation for the gravitational force, which states that:

$$\mathbf{F} = G \frac{Mm}{r^2},$$

When a star dies, it undergoes a contraction and blows off its outer envelope, forming a planetary nebula. [3,4] If the star collapses to within its Schwarzschild radius, it forms a black hole. [5]



Figure 1. A star forms a black hole.

Before the star collapses, the gravitational force between the planet and the star is represented by $\mathbf{F}_1$. After the collapse, the gravitational force between the planet and the black hole is represented by $\mathbf{F}_2$. The gravitational force of a black hole is extremely strong and nothing, not even light, can escape it. [6] Therefore, $\mathbf{F}_2$ is greater than $\mathbf{F}_1$. In this equation, M_1 represents the mass of the star, M_2 represents the mass of the planet, m represents the mass of the black hole, r represents the distance between their centers of mass, and G is the gravitational constant.

$$\mathbf{F}_1 < \mathbf{F}_2,$$
$$G \frac{M_1 m}{r^2} < G \frac{M_2 m}{r^2}.$$

As the star collapses into a black hole, it blows off its outer envelope and loses mass. [7] Assuming the star loses 0.2% of its mass during this process, the mass of the black hole can be represented as 99.8% of m_1 . This can be rewritten as:

$$G \frac{M_1 m}{r^2} < G \frac{0.998 M_1 m}{r^2}.$$

When the common parameters are removed, the equation can be simplified to:

$$1 < 0.998.$$

How is it even possible that 1 is less than 0.998? As the mass decreases from m_1 to m_2 and the distance r between the objects remains unchanged, it suggests that the gravitational constant G has increased. In Einstein’s theory of relativity [2], matter curves spacetime, and the Einstein field equations can be expressed in the following form:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}.$$

In this equation, $G_{\mu\nu}$ is the Einstein tensor and G is the gravitational constant. However, if the gravitational constant G is constantly changing, it raises the question of whether Einstein’s theory is still accurate.

Model

In Einstein’s theory of relativity [2], matter curves spacetime. What if the situation were reversed, and both matter and energy were converted from curved space?

Matter can release energy through annihilation, fission, and fusion [2], as matter and energy are different forms of the same thing. Therefore, matter curves spacetime because it is converted from curved space, and the flow of space released by matter creates gravitational force. The greater the amount of space released, the stronger the gravitational force generated. If one object releases much more space than another object, the flow of space will narrow the distance between the two objects. It is the space that moves, not the objects.



Figure 2. Gravitation.

When two galaxies are very far apart, the space they release accumulates in the middle, and this expands their distance. Therefore, the expansion of the universe and the phenomenon of redshift [8] are caused by the increase in space.




Figure 3. Expansion of the universe.

Thus, space curves in one dimension, converting matter and energy.




Figure 4. Curved space.

Since an electron is matter and there are only two directions in one dimension, matter curves in two different directions, creating two types of electric charges: positive and negative. Like charges repel each other because they occupy the same position in one dimension, while unlike charges attract each other because they occupy opposite positions in one dimension. Since they are all matter, they can only attract or repel each other, not annihilate.




Figure 5. Electric charges.

Quarks have three color charges [9]: red, green, and blue, which are converted by matter curving in three different dimensions. When a gluon is transferred between quarks, a color change occurs in both. For example, if a red quark emits a red–antigreen gluon, it becomes green; if a green quark absorbs a red–antigreen gluon, it becomes red. Quantum chromodynamics [10] proves that these three values of quark color can be converted into each other: they are different forms of the same thing.




Figure 6. Quark color.

Since there are only three values of quark color, space has only three dimensions. Therefore, space curves in six different directions of three dimensions, creating six types of matter: ordinary matter, antimatter, two types of dark matter, and two types of anti-dark matter.

Matter and antimatter in opposite directions attract and annihilate each other, while matter and dark matter perpendicular to each other in another dimension have no electromagnetic force. Both matter, antimatter, dark matter, and anti-dark matter are affected by gravitation.

During the Big Bang, matter and antimatter moved in opposite directions, so antimatter ended up on the other side of the universe. Currently, there is only dark matter and anti-dark matter in the observable universe. Particle collision experiments [11] prove that ordinary matter can be converted into antimatter: they are different forms of the same thing.




Figure 7. Six types of matter.

Matter is converted from curved space, and it also curves time.

In the double-slit experiment, an interference pattern emerges as the particles build up one by one. [12] This occurs because matter curves time, causing the particle from the past to interfere with the particle from the future.

In which-way experiment, if particle detectors are positioned at the slits, showing through which slit a photon goes, the interference pattern will disappear. [13] The mass of the observer is much greater than the particle, and when the observation occurs, the time of the observer engulfs the time of the particle, similar to how a black hole swallows a star, resulting in wave function collapse.

The Wheeler's delayed choice experiment demonstrates that extracting "which path" information after a particle passes through the slits can appear to retroactively alter its previous behavior at the slits. [14] Because matter curves time, the particle from the future can interfere with the particle from the past, meaning that the present behavior can have an impact on the past.

The quantum eraser experiment further shows that wave behavior can be restored by erasing or making permanently unavailable the "which path" information. [15] Since the time of the particle has no connection with the time of the observer, no wave function collapse occurs.




Figure 8. Matter curves time.

Every elementary particle is matter, and they are all converted from curved space, which gives them rest mass. However, the current commonly accepted physical theories imply or assume that the photon is strictly massless [16]. The Lorentz factor γ is defined as [17]:

$$\gamma = \frac{1}{\sqrt{1 - (v/c)^2}}.$$

In Newton's law of universal gravitation [1], the gravitational acceleration is:

$$\ddot{r} = \frac{Gm_0/r^2}{\sqrt{1 - (v/c)^2}}.$$

In Einstein's theory of relativity [2], the stress-energy tensor $T^{\alpha\beta}$ for a non-interacting particle with rest mass m_0 and trajectory $\mathbf{x}_p(t)$ is given by:

$$T^{\alpha\beta}(\mathbf{x}, t) = \frac{m_0 v^\alpha(t) v^\beta(t)}{\sqrt{1 - (v/c)^2}} \delta(\mathbf{x} - \mathbf{x}_p(t)).$$

If the photon has no rest mass, it would not be subject to gravitation and could escape from a black hole, contradicting observations. [6]

Of course, there is no doubt that the Standard Model is 100% accurate, so the blame is on the photon. The mistake, therefore, lies with the photon, which should behave precisely as predicted by scientists. The photon is both a wave and a particle, both matter and antimatter, and both has and does not have rest mass.

According to the standard model in particle physics [18], neutrino has zero rest mass and is a spin-half particle. Unfortunately, experimental observations by the Super-Kamiokande Observatory and the Sudbury Neutrino Observatories have shown that the neutrino actually has a non-zero rest mass, revealing the limitations of the Standard Model. [19–21]

Furthermore, as is widely known, dark matter possesses mass. Is this mass conferred by the Higgs field? If the answer is yes, does the Higgs boson composed of both matter, antimatter, and dark matter? If the answer is no, where does the mass of dark matter originate from?

Gravitation

Since there are only three values for the color charge of quarks, so the space has only three dimensions. Therefore, the space released by matter takes the form of a three-dimensional sphere. As this space flows outward, it forms a hollow sphere, and its volume can be written in the following form:

$$V_3 = \frac{4}{3} \pi r^3,$$

$$S_{\mu\nu} = \frac{4}{3} \pi ((r + \dot{r})^3 - r^3).$$

Where $S_{\mu\nu}$ is the space released by the matter, r is the radius of the sphere, and $\ddot{r}$ is the degree of space curvature. The greater the degree of space curvature, the greater the gravitational acceleration of the object. When the outward flow of space occurs, its volume remains constant, allowing for the direct calculation of the gravitational acceleration $\ddot{r}_2$ at a different distance using the gravitational acceleration $\ddot{r}_1$ at a known distance. The equation can be rewritten to:

$$S_{\mu\nu} = \frac{4}{3}\pi((r_1 + \ddot{r}_1)^3 - r_1^3) = \frac{4}{3}\pi((r_2 + \ddot{r}_2)^3 - r_2^3),$$
$$\ddot{r}_2 = \sqrt[3]{(r_1 + \ddot{r}_1)^3 - r_1^3 + r_2^3} - r_2.$$

The mass of the Earth is approximately 5.97237×10^{24} kg, and the average distance from its center to its surface is about 6.37123×10^6 m. [22,23] According to Newton’s law of universal gravitation, the value of the gravitational constant is approximately 6.67408×10^{-11} m³/kg/s². [24] The gravitational acceleration $\ddot{r}_1$ at the Earth’s surface can be calculated using this law and is given by:

$$\ddot{r}_1 = \frac{Gm}{r_1^2} \approx 9.81954911 \text{ m/s}^2.$$

If the distance up to 10^7 m, the gravitational acceleration $\ddot{r}_2$ is:

$$\ddot{r}_2 = \frac{Gm}{r_2^2} \approx 3.98600751 \text{ m/s}^2.$$

When the gravitational acceleration $\ddot{r}_1$ is equal to the Newtonian gravitational acceleration $\ddot{r}_1$ at r_1 , the gravitational acceleration $\ddot{r}_2$ at r_2 is:

$$\ddot{r}_2 = \sqrt[3]{(r_1 + \ddot{r}_1)^3 - r_1^3 + r_2^3} - r_2 \approx 3.98601207 \text{ m/s}^2.$$

As you can see, the value of the gravitational acceleration $\ddot{r}_2$ calculated using the Space Theory is extremely close to the value calculated using Newton’s law of universal gravitation. This confirms that the Space Theory can be used to accurately calculate the gravitational acceleration. The gravitational acceleration of the Earth at different distances in both models is shown in the following table:

Table 1. Gravitational acceleration of Earth.

Distance of Earth	0 m	1.15×10^3 m	10^4 m	10^5 m	10^6 m	10^7 m
Newton’s law	∞	$3 \times 10^8 \text{ m/s}^2$	3986007 m/s^2	39860 m/s^2	398.600 m/s^2	3.986007 m/s^2
Space Theory	106141 m/s^2	104991 m/s^2	96171 m/s^2	29976 m/s^2	398.442 m/s^2	3.986012 m/s^2

The mass of the Moon is approximately 7.342×10^{22} kg, its mean radius is about 1.737×10^6 m, and the time-averaged distance between the centers of the Earth and Moon is about 3.844×10^8 m. [25–27] When considering different distances, the gravitational acceleration of the Moon in two different models is shown in the following table:

Table 2. Gravitational acceleration of Moon.

Distance of Moon	0 m	1.27×10^2 m	10^5 m	10^6 m	10^7 m	3.844×10^8 m
Newton’s law	∞	$3 \times 10^8 \text{ m/s}^2$	490.01 m/s^2	4.900109 m/s^2	0.04900 m/s^2	0.00003316 m/s^2
Space Theory	24496 m/s^2	24369 m/s^2	487.60 m/s^2	4.899863 m/s^2	0.04899 m/s^2	0.00003316 m/s^2

The mass of the Sun is approximately 1.9885×10^{30} kg, with a mean radius of about 6.96342×10^8 m, and the mean distance between the centers of the Earth and the Sun is about 1.496×10^{11} m. [28,29] The table below shows the gravitational acceleration of the Sun in two different models at varying distances:

Table 3. Gravitational acceleration of Sun.

Distance of Sun	0 m	$6.65 \times 10^5 \text{ m}$	10^7 m	10^9 m	10^{10} m	$1.496 \times 10^{11} \text{ m}$
Newton's law	∞	$3 \times 10^8 \text{ m/s}^2$	1327140 m/s^2	132.71 m/s^2	1.3271 m/s^2	0.005929 m/s^2
Space Theory	7356638 m/s^2	6693449 m/s^2	1181938 m/s^2	132.71 m/s^2	1.3271 m/s^2	0.005929 m/s^2

When the distance is zero, Newton's gravitational acceleration becomes infinite, which is obviously incorrect. In contrast, the gravitation of the Space Theory is more accurate and does not require the use of Newton's constant of gravitation. The Schwarzschild radius [5] is a physical parameter that appears in the Schwarzschild solution to Einstein's field equations. It corresponds to the radius defining the event horizon of a black hole and can be expressed as:

$$r_s = \frac{2Gm}{c^2}.$$

The Schwarzschild radius of Earth is approximately 0.00887 m . However, when the distance is 1150 m , the Newton's gravitational acceleration exceeds approximately $3 \times 10^8 \text{ m/s}^2$, it is greater than the speed of light in vacuum, approximately $2.9979 \times 10^8 \text{ m/s}$, and this leads to the formation of a black hole, which is obviously wrong.

Expanding the formula, the gravitational acceleration of the Space Theory can be derived as:

$$S_{\mu\nu} = 4\pi r^2 \ddot{r} \left(1 + \frac{\dot{r}}{r} + \frac{\dot{r}^2}{3r^2} \right).$$

Then introduce a new variable β to represent $1 + \dot{r}/r + \dot{r}^2/(3r^2)$. The equation can be simplified to:

$$S_{\mu\nu} = 4\pi r^2 \ddot{r} \beta.$$

The expression implies that β can approach 1 as closely as desired by increasing the distance r to infinity.

$$\lim_{r \rightarrow \infty} \beta = 1.$$

At the surface of the Earth, the ratio of the gravitational acceleration $\ddot{r}$ to the distance r is approximately 0.00000153 , which is negligible and can be omitted. The formula can be rewritten as:

$$S_{\mu\nu} = 4\pi r^2 \ddot{r}.$$

Since the accelerations of the two formulas are equal at long distances, the formula can be simplified to:

$$\ddot{r} = \frac{S_{\mu\nu}}{4\pi r^2} = \frac{Gm}{r^2}.$$

After removing the same parameters, the gravitation of the Space Theory $S_{\mu\nu}$ thus takes the form:

$$S_{\mu\nu} = 4\pi Gm.$$

The result demonstrates that the space released by matter per kilogram is precisely equal to $4\pi G$, providing evidence that matter is converted from curved space, indicating that they are different forms of the same thing. The further the distance between the objects, the closer the values of the formulas.




Figure 9. Gravitational acceleration of Earth.

In Einstein's theory of relativity [2], matter curves spacetime, and the Einstein field equations can be expressed in the following form:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu},$$

$$\kappa = \frac{8\pi G}{c^4}.$$

Where $G_{\mu\nu}$ is the Einstein tensor, $T_{\mu\nu}$ is the stress-energy tensor, c is the speed of light in vacuum, and κ is the Einstein constant of gravitation.

In the geometrized unit system, the value of $4\pi G$ is set equal to unity. Therefore, it is possible to remove the Newton's constant of gravitation from the equations to avoid mistakes, especially since the Newton's law of universal gravitation does not apply to black holes. The equation can be rewritten as:

$$4\pi G = 1,$$

$$\kappa = \frac{2}{c^4}.$$

Since the theory is that matter and energy are both converted from curved space, the form should also be reversed. As space S is proportional to mass m and the fourth power of the speed of light in vacuum c^4 , the equivalent equation for space S can be expressed as:

$$S = Ec^2 = mc^4,$$

$$S_{\mu\nu} = \frac{4\pi G}{c^4} S.$$

Of course, under ordinary circumstances, if the Newton's "constant" of gravitation doesn't change too much, it can still be used to calculate the gravitational acceleration, the form can be rewritten to:

$$S_{\mu\nu} = 4\pi Gm = \frac{4}{3}\pi((r + \ddot{r})^3 - r^3),$$

$$\ddot{r} = \sqrt[3]{3Gm + r^3} - r.$$

On the surface of Earth, the gravitational acceleration is:

$$\ddot{r} = \frac{Gm}{r^2} \approx 9.81954911 \text{ m/s}^2.$$

$$\ddot{r} = \sqrt[3]{3Gm + r^3} - r \approx 9.81953396 \text{ m/s}^2.$$

And the relative error is:

$$\eta \approx 0.00000154284.$$

As you can see, this value is still extremely close to the original. What's more, this formula does not introduce any new variables.

In 1665, Newton extended the binomial theorem [1] to include real exponents, expanding the finite sum into an infinite series. To achieve this, he needed to give binomial coefficients a definition with an arbitrary upper index, which could not be accomplished through the traditional factorial formula. Nonetheless, for any given number n , it is possible to define the coefficients as:

$$\binom{n}{k} = \frac{n(n-1)(n-2)\cdots(n-k+1)}{k!} = \frac{(n)_k}{k!}.$$

The Pochhammer symbol $(\cdot)_k$ is used to represent a falling factorial, which is defined as a polynomial:

$$(n)_k = n^{\underline{k}} = n(n-1)(n-2)\cdots(n-k+1) = \prod_{k=1}^n (n-k+1).$$

This formula holds true for the usual definitions when n is a nonnegative integer. For any complex number n and real numbers x and y with $|x| > |y|$, the following equation holds:

$$(x + y)^n = \sum_{k=0}^{\infty} \binom{n}{k} x^{n-k} y^k = x^n + nx^{n-1}y + \frac{n(n-1)}{2!}x^{n-2}y^2 + \frac{n(n-1)(n-2)}{3!}x^{n-3}y^3 + \dots$$

The series for the cube root can be obtained by setting $n = 1/3$, which gives:

$$(x + y)^{\frac{1}{3}} = x^{\frac{1}{3}} + \frac{1}{3}x^{-\frac{2}{3}}y - \frac{1}{9}x^{-\frac{5}{3}}y^2 + \frac{5}{81}x^{-\frac{8}{3}}y^3 + \dots$$

At long distances where $|r| > |\sqrt[3]{3Gm}|$, the equation becomes:

$$\ddot{r} = \sqrt[3]{r^3 + 3Gm} - r = r + \frac{1}{3}r^{-2}3Gm - \frac{1}{9}r^{-5}(3Gm)^2 + \frac{5}{81}r^{-8}(3Gm)^3 + \dots - r,$$

$$\ddot{r} = \frac{Gm}{r^2} - \frac{1}{r}\left(\frac{Gm}{r^2}\right)^2 + \frac{5}{3r^2}\left(\frac{Gm}{r^2}\right)^3 + \dots$$

Then introduce a new variable β , the equation can be simplified to:

$$\ddot{r} = \frac{Gm}{r^2} + \beta,$$

$$\beta = -\frac{1}{r}\left(\frac{Gm}{r^2}\right)^2 + \frac{5}{3r^2}\left(\frac{Gm}{r^2}\right)^3 + \dots$$

At the surface of the Earth, the variable β is approximately -0.00001513417 , which is negligible and can be omitted. This is the reason why it is extremely close to Newton's value at long distances.

$$\lim_{r \rightarrow \infty} \beta = 0.$$

At short distances where $|\sqrt[3]{3Gm}| > |r|$, the form of the equation is different:

$$\ddot{r} = \sqrt[3]{3Gm + r^3} - r = (3Gm)^{\frac{1}{3}} + \frac{1}{3}(3Gm)^{-\frac{2}{3}}r^3 - \frac{1}{9}(3Gm)^{-\frac{5}{3}}r^6 + \frac{5}{81}(3Gm)^{-\frac{8}{3}}r^9 + \dots - r,$$

$$\ddot{r} = \sqrt[3]{3Gm} - r \left(1 - \frac{1}{3}\left(\frac{r}{\sqrt[3]{3Gm}}\right)^2 + \frac{1}{9}\left(\frac{r}{\sqrt[3]{3Gm}}\right)^5 - \frac{5}{81}\left(\frac{r}{\sqrt[3]{3Gm}}\right)^8 + \dots \right).$$

In summary, formulas inversely proportional to the distance r raised to the power of n can all be replaced with the hollow sphere model:

$$(x^n + ny)^{\frac{1}{n}} = x^{\frac{n-1}{n}} + \frac{1}{n}x^{\frac{1}{n}}ny + \dots = x + \frac{y}{x^{\frac{n-1}{n}}} + \dots,$$

$$\lim_{x \rightarrow \infty} \frac{y}{x^n} = \lim_{x \rightarrow \infty} \sqrt[n+1]{(n+1)y + x^{n+1}} - x.$$

In Chapter 7 of the book "Why String Theory?", it is mentioned that there is no direct experimental evidence for the String Theory. [30] To avoid encountering the same issue of unverifiability as String Theory, the author makes a prediction about the surface gravity of neutron star.

The Crab Pulsar (PSR B0531+21) has a mass of 1.4 solar masses ($M_{\odot}$) and a radius of approximately 10500 m. [31] The solar mass ($M_{\odot}$) is a standard unit of mass in astronomy, equal to approximately 1.98847×10^{30} kg. [28] On the surface of Crab Pulsar, the gravitational acceleration is:

$$\ddot{r} = \frac{Gm}{r^2} \approx 1.68523274 \times 10^{12} \text{ m/s}^2,$$

$$\ddot{r} = \sqrt[3]{3Gm + r^3} - r \approx 8.21924883 \times 10^6 \text{ m/s}^2.$$

Newton's prediction of $1.68523274 \times 10^{12} \text{ m/s}^2$, which is significantly greater than the speed of light in vacuum, $2.9979 \times 10^8 \text{ m/s}$, and this leads to the formation of a black hole, clearly deviates from what is expected. In contrast, the predicted value from the Space Theory, $8.21924883 \times 10^6 \text{ m/s}^2$, aligns more closely with reality.

In addition, the distance can also be determined by the gravitational acceleration, it can be rewritten as:

$$r = \sqrt{\frac{Gm}{\ddot{r}}},$$

$$r = \sqrt{\frac{Gm}{\ddot{r}} - \frac{1}{12}\ddot{r}^2} - \frac{1}{2}\ddot{r}.$$

And the product of variables, G and m , becomes:

$$Gm = \ddot{r}r^2,$$

$$Gm = \ddot{r}r^2 + \ddot{r}^2r + \frac{1}{3}\ddot{r}^3.$$

As you can see, when the acceleration is very small, the results between the two formulas will be very close. Due to the modification of the Newton's law, the orbital velocity also requires a corresponding adjustment, and it can be expressed in the following form:

$$\lim_{r \rightarrow \infty} \frac{x}{r} = \lim_{r \rightarrow \infty} \sqrt{2x + r^2} - r,$$

$$\dot{r} = \sqrt{\sqrt{2Gm + r^2} - r}.$$

The mass of the Earth is approximately 5.97237×10^{24} kg, and the mass of the Moon is approximately 7.342×10^{22} kg. [23,25] The time-averaged distance between the centers of the Earth and Moon is about 3.844×10^8 m. [27] The orbital velocity is:

$$\dot{r} = \sqrt{\frac{G(m_1 + m_2)}{r}} \approx 1024.54383 \text{ m/s},$$

$$\dot{r} = \sqrt{\sqrt{2G(m_1 + m_2) + r^2} - r} \approx 1023.84606 \text{ m/s}.$$

The modified formula resolves the issue of approaching infinity as the distance nears zero. Similarly, the distance of the orbit can also be determined by velocity, it can be rewritten as:

$$r = \frac{Gm}{\dot{r}^2} - \frac{1}{2}\dot{r}^2.$$

When the orbital velocity is 1024 m/s , the distance is:

$$r = \frac{G(m_1 + m_2)}{\dot{r}^2} \approx 3.84808 \times 10^8 \text{ m},$$

$$r = \frac{G(m_1 + m_2)}{\dot{r}^2} - \frac{1}{2}\dot{r}^2 \approx 3.84284 \times 10^8 \text{ m}.$$

To ensure the accuracy of calculations, the author recommends using the Decimal Module in Python, and suggests setting precision to 100 digits.

Electromagnetism

The same formula can also be used to calculate the electrostatic force. Interpret the formula as describing the propagation of photons in three-dimensional space, extending as a hollow sphere, with the number of particles remaining constant, and the density decreasing with distance, causing changes in the electric field. Therefore, the electric field can also be represented using the formula:

$$\lim_{r \rightarrow \infty} \frac{x}{r^2} = \lim_{r \rightarrow \infty} \sqrt[3]{3x + r^3} - r,$$

$$\mathbf{E} = \sqrt[3]{3k_e|q| + r^3} - r.$$

where **E** is the electric field of photons, k_e is the Coulomb constant, and q is the quantity of each charge. The Coulomb constant is about $8.98755179 \times 10^9 \text{ Nm}^2/\text{C}^2$, and the electric charge of a single electron is approximately $1.602176634 \times 10^{-19} \text{ C}$ [32]. If the distance of an electron is 1 m , the electric field is:

$$\mathbf{E} = k_e \frac{|q|}{r^2} \approx 1.43996426069 \times 10^{-9} \text{ N/C},$$

$$\mathbf{E} = \sqrt[3]{3k_e|q| + r^3} - r \approx 1.43996425861 \times 10^{-9} \text{ N/C}.$$

The further the distance between the objects, the closer the values of the formulas.




Figure 10. The electric field of an electron.

In the geometrized unit system, where $\hbar = e = G = k_e = 1$ and $\epsilon_0 = 1/(4\pi)$. For the convenience of calculation, here Z represents the atomic number of the atom, and the quantity of charge is set to 1. The Bohr radius, symbolized a_0 , is approximately 52.9 pm . When the distance is equal to the Bohr radius, the electric field of the hydrogen atom $Z = 1$ is:

$$\mathbf{E} = \frac{1}{(r/a_0)^2} = 1,$$

$$\mathbf{E} = \sqrt[3]{3 + (r/a_0)^3} - r/a_0 \approx 0.58740105.$$

The electric field is shown in the following table:

Table 4. The electric field of the Hydrogen atom.

r/a_0	0	0.25	0.5	1	2	4
Coulomb's law	∞	16	4	1	0.25	0.0625
Space Theory	1.4422	1.1947	0.962	0.5874	0.2239	0.0615

The variation is illustrated in the diagram:




Figure 11. The electric field of a Hydrogen atom.

And the orbital velocity is:

$$\dot{r} = \sqrt{\frac{1}{r/a_0}} = 1,$$

$$\dot{r} = \sqrt{2 + (r/a_0)^2} - r/a_0 \approx 0.85559967.$$

The velocity is shown in the following table:

Table 5. The orbital velocity of a Hydrogen atom.

r/a_0	0	0.25	0.5	1	2	4
Newton's law	∞	2	1.4142	1	0.7071	0.5
Space Theory	1.1892	1.0891	1	0.8555	0.6704	0.4925

The revised formula shows that the orbital velocity of an electron near the atomic nucleus increases by only 39% compared to when it's at the Bohr radius, which is completely different with the value in classical physics. After the correction, the electron will no longer fall into the nucleus when the distance is zero.




Figure 12. The orbital velocity of a Hydrogen atom.

Carbon, hydrogen, nitrogen, oxygen, phosphorus, and sulfur are the six most important chemical elements whose covalent combinations make up most biological molecules on Earth. [33] The following table shows atomic radius, Van der Waals radius and covalent bond from theoretical models. [34–39]

Table 6. Atomic radius from theoretical models.

Number	Symbol	Name	Radius			Bond		
			Calculated	Empirically	Van der Waals	Single	Double	Triple
1	H	Hydrogen	53 pm	25 pm	120 pm	32 pm		
2	He	Helium	31 pm	120 pm	140 pm			
6	C	Carbon	67 pm	70 pm	170 pm	75 pm	67 pm	60 pm
7	N	Nitrogen	56 pm	65 pm	155 pm	71 pm	60 pm	54 pm
8	O	Oxygen	48 pm	60 pm	152 pm	63 pm	57 pm	53 pm
15	P	Phosphorus	98 pm	100 pm	180 pm	111 pm	102 pm	94 pm
16	S	Sulfur	88 pm	100 pm	180 pm	103 pm	94 pm	95 pm

Table 7. Typical bond lengths and bond energies.

Bond	Length	Energy	Bond	Length	Energy	Bond	Length	Energy
C–O	143 pm	358 kJ/mo	N–C	147 pm	305 kJ/mo	N–O	144 pm	201 kJ/mo
C=O	123 pm	745 kJ/mo	N=C	127 pm	615 kJ/mo	N=O	120 pm	607 kJ/mo
C≡O	113 pm	1072 kJ/mo	N≡C	115 pm	891 kJ/mo	O–O	148 pm	146 kJ/mo
C–C	154 pm	347 kJ/mo	N–N	147 pm	160 kJ/mo	O=O	121 pm	495 kJ/mo

C=C	133 pm	614 kJ/mo	N=N	124 pm	418 kJ/mo	O-H	96 pm	467 kJ/mo
C≡C	120 pm	839 kJ/mo	N≡N	110 pm	941 kJ/mo	H-H	74 pm	436 kJ/mo
C-H	109 pm	413 kJ/mo	N-H	101 pm	391 kJ/mo			

The Lennard-Jones potential characterizes intermolecular interactions within pairs. This potential model accounts for both soft repulsive and attractive (van der Waals) forces and is applicable to electronically neutral atoms or molecules. The commonly employed expression for the Lennard-Jones potential is: [40–42]

$$V(r) = 4\varepsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right],$$

where r represents the distance between two interacting particles, ε represents the depth of the potential well, commonly known as the ‘dispersion energy’, and σ represents the distance at which the particle-particle potential energy $V(r)$ equals zero, often termed the ‘size of the particle’. The formula is rewritten using the hollow sphere model:

$$\lim_{r \rightarrow \infty} \frac{x}{r^6} = \lim_{r \rightarrow \infty} \sqrt[7]{7x + r^7} - r,$$

$$V(r) = 4\varepsilon \left[\left(\sqrt[7]{7\sigma^6 + r^7} - r - \frac{1}{2} \right)^2 - \frac{1}{4} \right].$$

The bond length of the O-H molecule is 96 pm, so the σ is:

$$\sigma = r \sqrt[6]{\frac{1}{2}} \approx 85.52627 \text{ pm},$$

$$\sigma = \sqrt[6]{\frac{1}{7} \left(\left(\frac{1}{2} + r \right)^7 - r^7 \right)} \approx 85.74948 \text{ pm}.$$

When r is 96 pm and σ is 85.52 pm, the result is:

$$\frac{V(r)}{\varepsilon} = 4 \left[\left(\left(\frac{\sigma}{r} \right)^6 - \frac{1}{2} \right)^2 - \frac{1}{4} \right] \approx -0.99999,$$

$$\frac{V(r)}{\varepsilon} = 4 \left[\left(\sqrt[7]{7\sigma^6 + r^7} - r - \frac{1}{2} \right)^2 - \frac{1}{4} \right] \approx -0.99975.$$

The variation is illustrated in the diagram:




Figure 13. Lennard-Jones potential of a hydrogen bond.

The formula can also be further modified:

$$\lim_{r \rightarrow \infty} \frac{x}{r^{12}} = \lim_{r \rightarrow \infty} \sqrt[13]{13x + r^{13}} - r,$$

$$V(r) = 4\varepsilon \left[\sqrt[13]{13\sigma^{12} + r^{13}} - \sqrt[7]{7\sigma^6 + r^7} \right].$$

When r is 96 pm and σ is 85.52 pm, the result is:




Figure 14. Lennard-Jones potential of a hydrogen bond.

Table 8. Lennard-Jones potential of a hydrogen bond.

Radius	0	25 pm	50 pm	100 pm	200 pm	400 pm
Lennard-Jones	∞	10264104	2407	-0.9526	-0.0243	-0.000382
Space Theory	56.6872	56.6111	48.0927	-0.9401	-0.0242	-0.000382

The Schrödinger equation for a hydrogen atom can be solved by separation of variables. [43] In this case, spherical polar coordinates are the most convenient.

$$\Psi(r,\theta,\varphi) = R(r)Y_l^m(\theta,\varphi) = R(r)\Theta(\theta)\Phi(\varphi).$$

The Spherical Harmonic functions indicate the electron’s position around the proton, and the radial function shows the electron’s distance from the proton. The solutions to the radial differential equation are products of the associated Laguerre polynomials, an exponential factor, and a normalization factor.

$$R(r) = N_{n,l} \left(\frac{r}{a_0} \right)^l \times L_{n,l}(r) \times e^{-\frac{r}{na_0}}.$$

The first six radial functions are provided in Table below.

Table 9. Radial wave functions for wavefunctions of the first three shells.

n	l		$\Phi(\varphi)$	$\Theta(\theta)$	$R(r)$
1	0	1s	$\frac{1}{\sqrt{2\pi}}$	$\frac{1}{\sqrt{2}}$	$2 \left(\frac{Z}{a_0} \right)^{\frac{3}{2}} e^{-\frac{Zr}{a_0}}$

2	0	2s	$\frac{1}{\sqrt{2\pi}}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2\sqrt{2}}\left(\frac{Z}{a_0}\right)^{\frac{3}{2}}\left[2 - \frac{Zr}{a_0}\right]e^{-\frac{Zr}{2a_0}}$
2	1	2p	$\frac{1}{\sqrt{2\pi}}$	$\frac{\sqrt{6}}{2}\cos\theta$	$\frac{1}{2\sqrt{6}}\left(\frac{Z}{a_0}\right)^{\frac{3}{2}}\left[\frac{Zr}{a_0}\right]e^{-\frac{Zr}{2a_0}}$
3	0	3s	$\frac{1}{\sqrt{2\pi}}$	$\frac{1}{\sqrt{2}}$	$\frac{2}{81\sqrt{3}}\left(\frac{Z}{a_0}\right)^{\frac{3}{2}}\left[27 - 18\frac{Zr}{a_0} + 2\left(\frac{Zr}{a_0}\right)^2\right]e^{-\frac{Zr}{3a_0}}$
3	1	3p	$\frac{1}{\sqrt{2\pi}}$	$\frac{\sqrt{6}}{2}\cos\theta$	$\frac{4}{81\sqrt{6}}\left(\frac{Z}{a_0}\right)^{\frac{3}{2}}\left[6\frac{Zr}{a_0} - \left(\frac{Zr}{a_0}\right)^2\right]e^{-\frac{Zr}{3a_0}}$
3	2	3d	$\frac{1}{\sqrt{2\pi}}$	$\frac{\sqrt{10}}{4}(3\cos^2\theta - 1)$	$\frac{4}{81\sqrt{30}}\left(\frac{Z}{a_0}\right)^{\frac{3}{2}}\left(\frac{Zr}{a_0}\right)^2e^{-\frac{Zr}{3a_0}}$

The radial probability density function is:

$$P(r) = |\Psi|^2 4\pi r^2.$$

For the 1s orbital, the radial wave function is given as:

$$P(r) = \left[\frac{1}{\sqrt{\pi}} \left(\frac{Z}{a_0} \right)^{\frac{3}{2}} e^{-\frac{Zr}{a_0}} \right]^2 4\pi r^2 = 4 \left(\frac{Z}{a_0} \right)^3 \left(r e^{-\frac{Zr}{a_0}} \right)^2.$$

The uv differentiation formula is:

$$(uv)' = u'v + uv' = 0.$$

Remove the constant terms, take the derivative with respect to r , and set it to zero to obtain the radius of maximum probability.

$$\begin{aligned} (r)' &= 1, \\ \left(e^{-r\frac{Z}{a_0}} \right)' &= -\frac{Z}{a_0} e^{-r\frac{Z}{a_0}}. \end{aligned}$$

Which gives:

$$\begin{aligned} \left[1 - r \frac{Z}{a_0} \right] e^{-r\frac{Z}{a_0}} &= 0, \\ r &= \frac{a_0}{Z}. \end{aligned}$$

Slater's rules [44] provide a value for the screening constant, denoted by s , which relates the effective and actual nuclear charges as:

$$Z_{\text{eff}} = Z - s.$$

Stars obtained their energy from nuclear fusion of hydrogen to form helium. [45] In this case, the radius for maximum probability will change from 52.9 pm to 31.1 pm. [34] This means that as the distance between the two atoms changes, the radius for maximum probability will also change. Which gives:

$$Z_{\text{eff}} = 1 + (1 - s)\alpha,$$

Set the distance between the two atoms as x , and $0 \leq \alpha \leq 1$, the formula can be modified to:

$$\alpha = \sqrt[3]{1 + (x/a_0)^3} - x/a_0.$$

The shielding constant s of a 1s electron is 0.3 [44], and the bond length in a H₂ molecule is 74 pm [39], so the radius for maximum probability of a H₂ molecule is:

$$Z_{\text{eff}} \approx 1.1071,$$

$$r = \frac{a_0}{Z_{\text{eff}}} \approx 47.7825.$$

The variation is illustrated in the diagram:




Figure 15. The radius for maximum probability (nuclear fusion).

And the radial probability density function $P(r)$ is:




Figure 16. The radial probability density (1s).

Since the revised orbital formula prevents electrons from falling into the nucleus, the model needs to be recalculated.



Figure 17. Hydrogen and Helium.

And the orbital velocity is:

$$\dot{r} = \sqrt{\sqrt{2 + (r/a_0)^2} - r/a_0},$$

$$\lim_{r \rightarrow 0} \dot{r} = \sqrt[4]{2} \approx 1.189207115.$$

The sum of $\mathbf{E}$ pointing toward the nucleus is:

$$\mathbf{E} = Z\mathbf{E}_p - (Z - 1)\mathbf{E}_e,$$

$$\mathbf{E}_p = \sqrt[3]{3 + \left(\frac{r}{a_0}\right)^3} - \frac{r}{a_0},$$

$$\mathbf{E}_e = \frac{4}{\pi^2} \int_0^{\pi/2} f(\theta) d\theta.$$

where $f(\theta)$ is:

$$f(\theta) = \sin\left(\frac{\pi}{2} - \theta\right) \left[\sqrt[3]{3 + \left(\cos(\theta) \frac{2r}{a_0}\right)^3} - \cos(\theta) \frac{2r}{a_0} \right].$$

The author used the quad function from the SciPy library to perform numerical integration, this function is part of the scipy.integrate module and is widely used for calculating definite integrals efficiently. [46] The maximum radial probability in the 1s orbital of helium occurs at a distance of 31 pm, which gives:

$$E_p \approx 0.88779184,$$

$$E_e \approx 0.26614705,$$

$$E = 2E_p - E_e \approx 1.50943663.$$

And the effective nuclear charge is:

$$s = 0.3,$$

$$Z_{\text{eff}} = Z - s = 1.7.$$

When considering E_p as 1 unit of charge, the result is:

$$s = \frac{E_e}{E_p} \approx 0.29978542 \approx 0.3,$$

$$Z_{\text{eff}} = \frac{E}{E_p} \approx 1.70021457 \approx 1.7.$$

Discussion

This paper is not yet complete. As of this writing, graviton has not been found yet.

Funding: None.

Acknowledgments: I would like to thank the scientists who study String Theory and make outstanding contributions to physics. The thickening speed of physics journals has surpassed the speed of light, but this doesn't violate the theory of relativity, because those journals aren't transmitting any information.

Conflict of Interest: None.

References

1. I. Newton. (1687). "Philosophiae Naturalis Principia Mathematica." The Mathematical Principles of Natural Philosophy.
2. A. Einstein. (1916). "Die Grundlage der allgemeinen Relativitätstheorie." Annalen der Physik, Vierte Folge, Band 49, 769-822.
3. R. Penrose. (1965). "Gravitational Collapse and Space-Time Singularities." Physical Review Letters. American Physical Society, 14, 57-59. doi:10.1103/physrevlett.14.57
4. A. Frankowski, N. Soker. (2009). "Very late thermal pulses influenced by accretion in planetary nebulae." New Astronomy, 14, 8, 654-658. doi:10.1016/j.newast.2009.03.006
5. K. Schwarzschild. (1916). "Über das Gravitationsfeld eines Massenpunktes nach der Einsteinschen Theorie." Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften, Seite 189-196.
6. D. Clery. (2020). "Black holes caught in the act of swallowing stars." Science, 367, 6477, 495. doi:10.1126/science.367.6477.495
7. S.W. Hawking. (1974). "Black hole explosions." Nature 248, 30-31. doi:10.1038/248030a0
8. E. Hubble. (1929). "A relation between distance and radial velocity among extra-galactic nebulae." PNAS 15, 3, 168-173. doi:10.1073/pnas.15.3.168
9. O.W. Greenberg. (1964). "Spin and Unitary Spin Independence in a Paraquark Model of Baryons and Mesons." Physical Review Letters, 13, 598-602.
10. R.P. Feynman. (2006). "QED: The strange theory of light and matter." Princeton University Press.
11. V. Hatton. (1991). "Operational history of the SPS collider 1981-1990." Conference Record of the 1991 IEEE Particle Accelerator Conference, 2952-2954. doi:10.1109/PAC.1991.165151
12. R. Pfleegor, L. Mandel. (1967). "Interference of Independent Photon Beams." Physical Review, 159, 5, 1084-1088. doi:10.1103/PhysRev.159.1084

13. W. Wootters, W. Zurek. (1979). "Complementarity in the double-slit experiment: Quantum nonseparability and a quantitative statement of Bohr's principle." *Physical Review D*, 19, 2, 473–484. doi:10.1103/PhysRevD.19.473
14. A. Peruzzo, P. Shadbolt, et al. (2012). "A Quantum Delayed-Choice Experiment." *Science*, 338, 6107, 634–637. doi:10.1126/science.1226719
15. X. Ma, J. Kofler, et al. (2012). "Quantum erasure with causally disconnected choice." *Proceedings of the National Academy of Sciences*, 110, 4, 110–1226. doi:10.1073/pnas.1213201110
16. M. Alonso, E. Finn. (1968). "Fundamental University Physics Volume III: Quantum and Statistical Physics." Addison Wesley, ISBN 978-0-201-00262-1.
17. J. Forshaw, G. Smith. (2014). "Dynamics and Relativity." John Wiley & Sons, ISBN 978-1-118-93329-9.
18. R. Oerter. (2006). "The Theory of Almost Everything: The Standard Model, the Unsung Triumph of Modern Physics." Penguin Group, ISBN 978-0-13-236678-6.
19. B. Pontecorvo. (1958). "Mesonium and Antimesonium." *Soviet Journal of Experimental and Theoretical Physics*, 6, 429–431.
20. E. Kearns, T. Kajita, Y. Totsuka. (1999). "Detecting Massive Neutrinos." *Scientific American*, 281, 64–71. doi:10.1038/scientificamerican0899-64
21. A. McDonald, J. Klein, D. Wark. (2006). "Solving the Solar Neutrino Problem." *Scientific American*, 288, 40–49. doi:10.1038/scientificamerican0403-40
22. B. Luzum, N. Capitaine, A. Fienga, et al. (2011). "The IAU 2009 system of astronomical constants: The report of the IAU working group on numerical standards for Fundamental Astronomy." *Celestial Mechanics and Dynamical Astronomy*, 110, 293–304. doi:10.1007/s10569-011-9352-4
23. F. Chambat, B. Valette. (2001). "Mean radius, mass, and inertia for reference Earth models." *Physics of the Earth and Planetary Interiors*, 124, 3–4, 234–253. doi:10.1016/S0031-9201(01)00200-X
24. P. Mohr, D. Newell, B. Taylor. (2016). "CODATA Recommended Values of the Fundamental Physical Constants: 2014." *Journal of Physical and Chemical Reference Data*, 45, 4, 1527–1605. doi:10.1063/1.4954402
25. M. Wieczorek, B. Jolliff, A. Khan, et al. (2006). "The Constitution and Structure of the Lunar Interior." *Reviews in Mineralogy and Geochemistry*, 60, 1, 221–364. doi:10.2138/rmg.2006.60.3
26. C. Hirt, W.E. Featherstone. (2012). "A 1.5 km-resolution field model of the Moon." *Earth and Planetary Science Letters*, 329–330, 22–30. doi:10.1016/j.epsl.2012.02.012
27. J. Battat, T. Murphy, E. Adelberger, et al. (2009). "The Apache Point Observatory Lunar Laser-ranging Operation (APOLLO): Two Years of Millimeter-Precision Measurements of the Earth-Moon Range." *Astronomical Society of the Pacific*, 121, 875, 29–40. doi:10.1086/596748
28. G. Poole. (2019). "Cosmic Power Generation and Gravity." *Journal of High Energy Physics, Gravitation and Cosmology*, 5, 920-927. doi:10.4236/jhepgc.2019.53047
29. E. Pitjeva, E. Standish. (2009). "Proposals for the masses of the three largest asteroids, the Moon-Earth mass ratio and the Astronomical Unit." *Celestial Mechanics and Dynamical Astronomy*, 103, 365–372. doi:10.1007/s10569-009-9203-8
30. J. Conlon. (2016). "Why string theory?" CRC Press.
31. D. Jeffery. (n.d.). "Neutron Stars and the Crab Nebula." University of Nevada, Las Vegas. https://www.physics.unlv.edu/~jeffery/astro/neutron_star/crab_nebula.html
32. NIST. (2019). "2018 CODATA Value: vacuum electric permittivity." The NIST Reference on Constants, Units, and Uncertainty.
33. E. Frieden. (1972). "The chemical elements of life." *Scientific American*, 227(1), 52-64.
34. E. Clementi, D. L. Raimondi, W. P. Reinhardt. (1967). "Atomic screening constants from SCF functions. II. Atoms with 37 to 86 electrons." *The Journal of chemical physics*, 47(4), 1300-1307.
35. A. V. Bondi. (1964). "van der Waals volumes and radii." *The Journal of physical chemistry*, 68(3), 441-451.
36. P. Pyykkö, M. Atsumi. (2009). "Molecular single-bond covalent radii for elements 1–118." *Chemistry–A European Journal*, 15(1), 186-197.
37. P. Pyykkö, M. Atsumi. (2009). "Molecular double-bond covalent radii for elements Li–E112." *Chemistry–A European Journal*, 15(46), 12770-12779.
38. P. Pyykkö, S. Riedel, M. Patzschke. (2005). "Triple-bond covalent radii." *Chemistry–A European Journal*, 11(12), 3511-3520.
39. G. Carruth, E. Ehrlich. "Bond Energies." Volume Library. Ed. Carruth, Gorton. Vol 1. Tennessee: Southwestern, 2002.
40. JE Jones. "On the determination of molecular fields.—I. From the variation of the viscosity of a gas with temperature." *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 106.738 (1924): 441-462.
41. JE Jones. "On the determination of molecular fields.—II. From the equation of state of a gas." *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 106.738 (1924): 463-477.
42. JE Lennard-Jones. "Cohesion." *Proceedings of the Physical Society* 43.5 (1931): 461.

43. E Schrödinger. "An undulatory theory of the mechanics of atoms and molecules." Physical review 28.6 (1926): 1049.
44. JC Slater. "Atomic shielding constants." Physical review 36.1 (1930): 57.
45. DD Clayton. Principles of stellar evolution and nucleosynthesis. University of Chicago press, 1983.
46. Virtanen, P., et al. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. Nature Methods, 17, 261–272.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.