

Review

Then there were plenty –ring meristems giving rise to many stamen whorls

Doudou Kong¹ and Annette Becker^{1,*}

¹ Justus-Liebig-University, Institute of Botany, Heinrich-Buff-Ring 38, 35392 Giessen, Germany; Doudou.Kong@bot1.bio.uni-giessen.de

* Correspondence: Annette.becker@bot1.bio.uni-giessen.de; Tel.: +49 641 9935200

Abstract: Floral meristems are dynamic systems that generate floral organ primordia at their flanks and, in most species, terminate while giving rise to the gynoeceum primordia. However, we find species with floral meristems that generate additional ring meristems repeatedly throughout angiosperm history. Ring meristems produce only stamen primordia, resulting in polystemous flowers (having stamen numbers more than double that of petals or sepals), and act independently of the floral meristem activity. Most of our knowledge on floral meristem regulation is derived from molecular genetic studies of *Arabidopsis thaliana*, a species with a fixed number of floral organs and as such of only limited value to understand ring meristem function, regulation, and ecological value. This review provides an overview of the main molecular players regulating floral meristem activity in *A. thaliana*, and summarizes our knowledge of ring primordia morphology and occurrence in dicots. Our work provides a first step towards understanding the significance and molecular genetics of ring meristem regulation and evolution.

Keywords: floral meristem, polystemony, numerous stamens, evo-devo, ring meristem

1. Introduction

Flowers are among the most beautiful examples for mutually beneficial relationships between animals and plants. Flowers are the major innovation of angiosperms and their astonishing diversification is thought to be the major driver of the enormous success of the flowering plants. Even though most flowers are composed of only four different types of organs, their variability in structure, color, and scent is amazing and fine-tuned for attracting insect or bird pollinators, or to allow effective wind pollination [1–3].

In this review, we will first outline the general morphogenesis of the floral meristem and describe the functions of the key genes involved in regulating maintenance and termination of the floral meristem. We will then explain the concept of polystemony and identify lineages with independent origin of polystemony.

In many species, flower primordia arise from the flanks of the inflorescence meristem in a precise pattern (phyllotaxy), and in few cases only, the shoot apical meristem turns into a floral meristem. For several species, the positioning of the flower primordia follows a localized auxin maximum via PIN1 mediated auxin transport in the epidermal L1 layer [4]. After initiation of the flower primordium, a meristem-to-organ boundary forms separating the floral primordium from the rest of the plant body and the floral meristem enters its growth phase [4,5]. Auxin transport is now reversed, away from the primordium, to create an auxin sink in the central tissues of the shoot or inflorescences mediating the connection of the floral primordium to the vasculature [6].

In *Arabidopsis thaliana*, all floral organs arise from the floral meristem which, unlike the shoot or root meristem, undergoes a genetically defined succession of initiation, maintenance, and termination such that stem cell activity ceases once all floral organs are formed [5]. The floral meristem releases floral organ primordia at its flanks, and the number and position of stem cells is tightly regulated ensuring meristem stability as well as

allowing cells to accumulate before organogenesis. Thus, the meristem is a defined structure even though its constituent cells are constantly changing. Cells exiting the meristem adopt their future cell fate according to their position within the meristem, while remaining in their original cell layers. Floral organs arise as primordia, composed of undifferentiated cells, which adopt a floral organ identity, and subsequently grow and differentiate into the floral organs. In most flowers, floral organs are arranged in whorls, with sepals being the outermost organs, the second whorl are the petals, then the pollen grain producing stamens, and in the center of the flower the gynoecium is formed, harboring the ovules. After fertilization, the ovules develop into seeds and the gynoecium is transformed into the fruit [7]. The development of floral organs follows a genetically determined blueprint, which was elucidated by molecular genetics studies in model species like *Arabidopsis thaliana*, *Antirrhinum majus* and grasses. Several recent reviews summarize our knowledge on floral organ identity, its variation and origin, and remaining open questions [8–11] which will not be covered here.

Floral meristems are thus dynamic systems, balancing proliferation and termination in a species-specific manner, leading to a species-specific size and number of the four different floral organ types. The shoot and inflorescence meristems are often, and to at least some extent indeterminate, but stem cell maintenance in the floral meristem ends after the initiation of the gynoecium rendering the floral meristem determinate [12]. *Arabidopsis thaliana* has a fixed number of floral organs and is typical for the canalized floral ground plan of many core eudicots. However, deviations from this ground plan are numerous and little is known about the conservation of the gene regulatory network (GRN) balancing floral meristem action. While it seems reasonable to assume that some degree of conservation exists in floral meristem GRNs, this genetic system also allows extreme examples of variation of floral organ number (merosity): the floral meristem of the Ranunculaceae *Laccopetalum giganteum* generates up to 10.000 small carpels [13] while the bisexual Hydatellaceae (sister to all other Nymphaeales) flowers consist of either a single stamen or a single carpel [14].

2. Regulation of stem cell activity in the floral meristem

The genetic networks regulating flower development have been characterized to the largest extent in the genetic model organism *Arabidopsis thaliana*. Molecular data from other species are largely missing, thus we are showing the *A. thaliana* way of modulating floral meristem fate keeping in mind that this floral meristem generates a fixed number of whorls and organs per whorl.

The regulation of floral meristem termination involves a highly complex regulatory interplay of transcription factors, micro RNAs, receptor-like kinases and their ligands, MAP kinases, and chromatin remodeling: the homeodomain-containing transcription factors WUSCHEL (*WUS*) and SHOOT MERISTEMLESS (*STM*) function synergistically during shoot, inflorescence, and floral meristem development and maintenance. In *wus* and *stm* mutants, the floral meristem ceases prematurely because a smaller number of stem cells in the central zone is formed which cannot keep up with the production of a larger number of cells in the periphery required for floral organ formation [15–18]. Conversely, overexpression of *WUS* leads to a conversion of cells in organ primordia into cells with stem cell characteristics [19]. *WUS* activation is limited by the CLAVATA (*CLV*) signaling loop where the small apoplastic signaling peptide *CLV3*, activated by *WUS* in L1 and L2, moves toward L3 where it binds to the *CLV1* and *CLV2* receptors and other receptor-like kinases to indirectly repress *WUS* expression [20]. When any of the *CLV* genes are non-functional, the meristem's central zone increases due to an overexpression of *WUS* [21,22]. *WUS* also activates the floral homeotic gene *AGAMOUS* (*AG*) by direct binding to its second intron, and thus, *WUS* regulates not only stem cell maintenance but is also required for floral organ initiation [23]. Once the floral meristem has produced all floral organ primordia, it is consumed in the process of carpel development and stem cell activity ceases. For this process, the floral homeotic C function protein *AG* is required to repress *WUS* by

directly binding to its genomic locus and by the recruitment of Polycomb group proteins that methylate chromatin to silence the *WUS* genomic locus [18,24]. In addition, *AG* activates *KNUCKLES (KNU)* by replacing repressing histone marks with activating ones in a time-dependent manner, and *KNU* then represses *WUS* expression [25]. Also, the floral homeotic A function gene *APETALA2 (AP2)*, which also promotes stem cell fate in the floral meristem by directly antagonizing *AG*, is translationally repressed by miR172d to achieve floral determinacy [26,27]. Further, *AP2* not only negatively regulates the expression of *AG*, it also independently regulates the activity of floral meristem [28]. At the same time, there is evidence that *AP2* negatively regulates the *CLV* signaling pathway [29].

Moreover, additional transcription factors, such as *SUPERMAN (SUP)*, *CRABS CLAW (CRC)*, *PERIANTHA (PAN)*, and *ULTRAPETALA (ULT)*, also regulate floral meristem maintenance and termination, but to a lesser extent and in a partially redundant way, suggesting a high level of genetic robustness in the floral meristem termination network [30–32].

More recently, auxin signaling was reported to play a major role in floral meristem termination on several levels. For example, *SUP* recruits chromatin remodelers to suppress auxin biosynthesis genes at early stages of floral meristem development. If this function is lacking, the floral meristem is enlarged and active for a prolonged time [33]. The *AG* target *CRC* connects floral meristem termination with gynoecium development by activating the putative auxin transporter *TORNADO2* to promote gynoecium formation while the floral meristem terminates [34]. The auxin response factor *ARF3* is regulated by auxin levels, but also by *AG* and *AP2* and is required to terminate *WUS* activity in the floral meristem [35]. However, it remains unclear how auxin biosynthesis and transport contributes to floral meristem termination on the molecular level.

While several meristem regulators in other species such as *Solanum lycopersicum* (tomato), *Oryza sativa* (rice), and *Zea mays* (corn) were already identified and functionally characterized, they all turned out to be orthologs of the genes analyzed in *A. thaliana* [36] except for *BEARDED-EAR* and *MOSAIC FLORAL ORGANS1* from corn and tomato [37,38], respectively. Interestingly, many components of the floral organ identity and floral meristem termination regulatory network are conserved among seed and even non-seed plants. For example, *WUS* homologs are required for the initiation of cell growth during stem cell formation in the moss *Physcomitrella patens* [39] and several *WUS* homologs were also identified in gymnosperms [40].

3. How to generate multiple stamen primordia

Considering variations in the floral ground plan, the number of stamens is especially labile and we will concentrate on this trait in the following section. The flower of the most recent common ancestor (MRCA) of all angiosperms had most likely more than ten stamens arranged in three stamens per whorl suggesting at least four whorls generating stamen primordia [41]. In recent species, the most common stamen arrangement in monocots and core eudicots is in two whorls with the outer stamens in alternate position to the petals [42]. In the ANITA grade, whose members are sister to all other angiosperms, stamen numbers are especially labile and range from one in Hydatellaceae and Chloranthaceae up to 300 in Schisandraceae [42]. In species with an increased number of stamens, those are often not arranged spirally, but rather an androecial ring primordium forms generating whorls of stamens. The direction of stamen primordia initiation can be towards the center of the flower (centripetal), the periphery (centrifugal), or bidirectionally (Figure 1). The ring primordium sometimes subdivides into sectors and is thus fragmented [42]. This additional ring meristem provides a mean to uncouple stamen initiation from carpel initiation, such that the last stamens initiate after the carpels are already far advanced in their development (Figure 1).




Figure 1. Flowers with ring meristems. (A) closed ring meristem and (B) fragmented ring meristem, the ring meristem is orange. Black arrows indicate the direction of stamen primordia initiation. (C–D) Ring meristem of *Eschscholzia californica* forming stamen primordia. (C) Bud of a late stage 4 flower in which the floral dome separates from the ring meristem (indicated by a blurred orange line). (D) Stage 5 bud showing the continuous formation of stamen primordia while the gynoecium has already initiated. (E) Stage 6 bud showing the formation of new stamen primordia (white arrow) at the time when the early stamens already form a flat surface for microsporangia initiation. The sepals were removed in all images, and staging was done according to Becker et al. (2005) [43]. Abbreviations: b, bidirectional; cp, centripetal; cf, centrifugal; FD, floral dome; fm, floral meristem; G, gynoecium; p, petal; RM, ring meristem; SE, sepal; SP, stamen primordium; S, stamen. Scale bar in A = 86 μm , in B = 100 μm , in C = 120 μm (A and B modified from Endress, 2011, C – E from Becker, 2016 with permission) [7,42].

4. Ring meristems in eudicots

Eudicots comprise 44 orders, and ring meristems producing extra stamen whorls are found in 13 of them (AGP, 2016 and Figure 2) [44]. Interestingly, the fraction of families with ring meristems per order is unequally distributed across the phylogeny. In Ranunculales, all families have members with ring meristems and almost all combinations of closed/fragmented architecture and directions of stamen primordia initiation occurs. Asterales consist of 11 families and a staggering number of almost 27,000 species, but include only the genus *Taraxacum* within the Asteraceae that has a ring meristem. In most orders, only a fraction of members develops ring meristems. For example, in Proteales, Fabales, Rosales, and Myrtales, only a single family includes members with ring meristems. Interestingly, most ring meristems are of the closed type and the number of orders with centripetal and centrifugal initiation of primordia is similar. However, bidirectional initiation is rarely found (Figure 2). Interestingly, centripetal primordia initiation is predominant in the non-core eudicot orders of Ranunculales and Proteales, but in the core eudicots, centrifugal initiation is found more often. In several groups, a closed ring meristem is associated with centripetal primordia initiation, for example in the Ranunculales, Proteales, Fabales, Hamamelidaceae, and Portulacaceae. In contrast, all combinations of primordia initiation direction and closed/fragmented ring meristem occur within the Malvales. Based on our data, there seems no preference in co-occurrence between closed and fragmented ring meristems and the direction of stamen primordia initiation. However, data compilation on the species level are lacking so far.




Figure 2. Distribution and types of ring meristems across eudicots. On the left, a simplified phylogeny shows only orders and families with ring meristem occurrence. Numbers in brackets next to the orders indicate the number of families in the respective order. On the right, the type of ring meristem is indicated. Numbers below summarize occurrences of morphological traits. Based on [42,44–55].

5. Are ring meristems in dicots of independent origin?

In the Proteales and Ranunculales, which are sister to all other eudicots, ring meristems are found in all (Ranunculales) or at least some (Proteales) families, but in Buxales and Trochodendrales, also non-core eudicots, ring meristems are missing. It thus remains unclear if the MRCA of eudicots had a floral ring meristem and if this was the case, we need to propose losses of ring meristems in the majority of eudicot orders. Generating more stamen and, in consequence, more costly pollen grains, requires a selective advantage for trait maintenance. If a high stamen number is not advantageous, independent losses may be rather frequent. In contrast, it seems also reasonable to propose that the eudicot MRCA had no ring meristem, but according to this hypothesis, at least 13 independent gains of ring meristem activity need to be postulated. However, the diversity of ring meristem architectures in dicots may be a hint towards several independent gains, as not even in the Ranunculales, an evolutionary trend from an ancestral to a more derived condition in ring meristem structure can be safely derived.

6. Conclusion and outlook

Here, we have compiled recent findings on transcription factors, signaling pathways and phytohormone action, all involved in floral meristem termination of *Arabidopsis*. We further demonstrated that at least some components of the floral meristem regulatory systems are conserved between eudicots and mosses. However, the molecular regulation of floral meristem termination is mainly analyzed in *Arabidopsis* and even knowledge on this process in the monocot model species rice is lagging behind. Meristematic activity

regulation in species with ring meristems has, to our knowledge, not been studied so far, leaving many questions open. This raises the question how the meristem activity in the central floral meristem and the ring meristem is differentiated and correlated in space and time, when presumably at least partially similar developmental mechanisms govern this activity. Unlike the ring meristem, which gives rise to only stamen primordia, the floral meristem releases lateral organs whose order and identity are precisely regulated. In most species, termination of the floral meristem activity is coupled to gynoecium initiation. However, ring meristems give rise to several whorls of stamen and it remains unclear how these meristems are initiated, separated from the floral meristem, and then terminated. Further, besides open questions related to their developmental regulation remaining unanswered so far, their ecological significance is unclear. For example, it would be interesting to test whether ring meristem occurrence and polystemony is correlated to specific pollination modes and pollinator preferences. Or how fast polystemony disappears when selfing becomes more prominent. Further, ring meristems producing varying numbers of stamen may shift the balance between male and female contributions towards sexual reproduction. Thus, the regulation of ring meristem activity termination, at least in some species [43], may be more flexible than that of the central floral meristem. For example, it may depend on environmental cues or on the accumulation of sufficient reserves, or other metabolic changes during a plant's maturation process. Thus, our review provides only a first glance into a morphological structure that is enigmatic in terms of molecular mechanism of its regulation and with respect to its origin, evolution, and ecological significance.

Author Contributions: Writing—original draft preparation, D.-D.K, A.B.; writing—review and editing, D.-D.K.; visualization, D.-D.K.; acquisition, A.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by German Research Foundation (DFG), grant number BE 2547/24-1

Acknowledgments: We thank Dr. Matthias Lange for interesting discussion on the subject already years ago, and Katrin Ehlers, Dominik Lotz, and Clemens Roessner.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Chanderbali, A.S.; Berger, B.A.; Howarth, D.G.; Soltis, D.E.; Soltis, P.S. Evolution of floral diversity: genomics, genes and gamma. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **2017**, *372*, doi:10.1098/rstb.2015.0509.
2. Soltis, P.S.; Brockington, S.F.; Yoo, M.-J.; Piedrahita, A.; Latvis, M.; Moore, M.J.; Chanderbali, A.S.; Soltis, D.E. Floral variation and floral genetics in basal angiosperms. *Am. J. Bot.* **2009**, *96*, 110–128, doi:10.3732/ajb.0800182.
3. Endress, P.K. The early evolution of the angiosperm flower. *Trends in Ecology & Evolution* **1987**, *2*, 300–304, doi:10.1016/0169-5347(87)90082-6.
4. Denay, G.; Chahtane, H.; Tichtinsky, G.; Parcy, F. A flower is born: an update on Arabidopsis floral meristem formation. *Curr. Opin. Plant Biol.* **2017**, *35*, 15–22, doi:10.1016/j.pbi.2016.09.003.
5. Chang, W.; Guo, Y.; Zhang, H.; Liu, X.; Guo, L. Same Actor in Different Stages: Genes in Shoot Apical Meristem Maintenance and Floral Meristem Determinacy in Arabidopsis. *Front. Ecol. Evol.* **2020**, *8*, doi:10.3389/fevo.2020.00089.
6. Furutani, M.; Nakano, Y.; Tasaka, M. MAB4-induced auxin sink generates local auxin gradients in Arabidopsis organ formation. *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 1198–1203, doi:10.1073/pnas.1316109111.
7. Becker, A.; Ehlers, K. Arabidopsis flower development—of protein complexes, targets, and transport. *Protoplasma* **2016**, *253*, 219–230, doi:10.1007/s00709-015-0812-7.
8. Schragel-Lavelle, A.; Klein, H.; Fisher, A.; Bartlett, M. Grass flowers: An untapped resource for floral evo-devo. *Jnl of Sytematics Evolution* **2017**, *55*, 525–541, doi:10.1111/jse.12251.
9. Theißen, G.; Melzer, R.; Rümpler, F. MADS-domain transcription factors and the floral quartet model of flower development: linking plant development and evolution. *Development* **2016**, *143*, 3259–3271, doi:10.1242/dev.134080.
10. Kramer, E.M. Plus ça change, plus c'est la même chose: The developmental evolution of flowers. *Curr. Top. Dev. Biol.* **2019**, *131*, 211–238, doi:10.1016/bs.ctdb.2018.11.015.
11. Smyth, D.R. Evolution and genetic control of the floral ground plan. *New Phytol.* **2018**, *220*, 70–86, doi:10.1111/nph.15282.
12. Sun, B.; Xu, Y.; Ng, K.-H.; Ito, T. A timing mechanism for stem cell maintenance and differentiation in the Arabidopsis floral meristem. *Genes Dev.* **2009**, *23*, 1791–1804, doi:10.1101/gad.1800409.

13. Emadzade, K.; Lehnebach, C.; Lockhart, P.; Hörandl, E. A molecular phylogeny, morphology and classification of genera of Ranunculaceae (Ranunculaceae). *Taxon* **2010**, *59*, 809–828, doi:10.1002/tax.593011.
14. Rudall, P.J.; Sokoloff, D.D.; Remizowa, M.V.; Conran, J.G.; Davis, J.I.; Macfarlane, T.D.; Stevenson, D.W. Morphology of Hydatellaceae, an anomalous aquatic family recently recognized as an early-divergent angiosperm lineage. *Am. J. Bot.* **2007**, *94*, 1073–1092, doi:10.3732/ajb.94.7.1073.
15. Long, J.A.; Moan, E.L.; Medford, J.I.; Barton, M.K. A member of the KNOTTED class of homeodomain proteins encoded by the *STM* gene of *Arabidopsis*. *Nature* **1996**, *379*, 66–69, doi:10.1038/379066a0.
16. Laux, T.; Mayer, K.F.; Berger, J.; Jürgens, G. The *WUSCHEL* gene is required for shoot and floral meristem integrity in *Arabidopsis*. *Development* **1996**, *122*, 87–96, doi:10.1242/dev.122.1.87.
17. Clark, S.E.; Jacobsen, S.E.; Levin, J.Z.; Meyerowitz, E.M. The *CLAVATA* and *SHOOT MERISTEMLESS* loci competitively regulate meristem activity in *Arabidopsis*. *Development* **1996**, *122*, 1567–1575, doi:10.1242/dev.122.5.1567.
18. Lenhard, M.; Bohnert, A.; Jürgens, G.; Laux, T. Termination of Stem Cell Maintenance in *Arabidopsis* Floral Meristems by Interactions between *WUSCHEL* and *AGAMOUS*. *Cell* **2001**, *105*, 805–814, doi:10.1016/S0092-8674(01)00390-7.
19. Gallois, J.-L.; Nora, F.R.; Mizukami, Y.; Sablowski, R. *WUSCHEL* induces shoot stem cell activity and developmental plasticity in the root meristem. *Genes Dev.* **2004**, *18*, 375–380, doi:10.1101/gad.291204.
20. Kinoshita, A.; Betsuyaku, S.; Osakabe, Y.; Mizuno, S.; Nagawa, S.; Stahl, Y.; Simon, R.; Yamaguchi-Shinozaki, K.; Fukuda, H.; Sawa, S. RPK2 is an essential receptor-like kinase that transmits the CLV3 signal in *Arabidopsis*. *Development* **2010**, *137*, 3911–3920, doi:10.1242/dev.048199.
21. Brand, U.; Fletcher, J.C.; Hobe, M.; Meyerowitz, E.M.; Simon, R. Dependence of stem cell fate in *Arabidopsis* on a feedback loop regulated by *CLV3* activity. *Science* **2000**, *289*, 617–619, doi:10.1126/science.289.5479.617.
22. Fletcher, J.C.; Brand, U.; Running, M.P.; Simon, R.; Meyerowitz, E.M. Signaling of cell fate decisions by *CLAVATA3* in *Arabidopsis* shoot meristems. *Science* **1999**, *283*, 1911–1914, doi:10.1126/science.283.5409.1911.
23. Lohmann, J.U.; Hong, R.L.; Hobe, M.; Busch, M.A.; Parcy, F.; Simon, R.; Weigel, D. A Molecular Link between Stem Cell Regulation and Floral Patterning in *Arabidopsis*. *Cell* **2001**, *105*, 793–803, doi:10.1016/s0092-8674(01)00384-1.
24. Ji, L.; Liu, X.; Yan, J.; Wang, W.; Yumul, R.E.; Kim, Y.J.; Dinh, T.T.; Liu, J.; Cui, X.; Zheng, B.; et al. *ARGONAUTE10* and *ARGONAUTE1* regulate the termination of floral stem cells through two microRNAs in *Arabidopsis*. *PLoS Genet.* **2011**, *7*, e1001358, doi:10.1371/journal.pgen.1001358.
25. Sun, B.; Looi, L.-S.; Guo, S.; He, Z.; Gan, E.-S.; Huang, J.; Xu, Y.; Wee, W.-Y.; Ito, T. Timing mechanism dependent on cell division is invoked by Polycomb eviction in plant stem cells. *Science* **2014**, *343*, 1248559, doi:10.1126/science.1248559.
26. Huang, Z.; Shi, T.; Zheng, B.; Yumul, R.E.; Liu, X.; You, C.; Gao, Z.; Xiao, L.; Chen, X. *APETALA2* antagonizes the transcriptional activity of *AGAMOUS* in regulating floral stem cells in *Arabidopsis thaliana*. *New Phytol.* **2017**, *215*, 1197–1209, doi:10.1111/nph.14151.
27. Chen, X. A microRNA as a translational repressor of *APETALA2* in *Arabidopsis* flower development. *Science* **2004**, *303*, 2022–2025, doi:10.1126/science.1088060.
28. Zhao, L.; Kim, Y.; Dinh, T.T.; Chen, X. miR172 regulates stem cell fate and defines the inner boundary of *APETALA3* and *PIS-TILLATA* expression domain in *Arabidopsis* floral meristems. *The Plant Journal* **2007**, *51*, 840–849. doi:10.1111/j.1365-3113.2007.03181.x.
29. Würschum, T.; Groß-Hardt, R.; Laux, T. *APETALA2* regulates the stem cell niche in the *Arabidopsis* shoot meristem. *The Plant Cell* **2006**, *18*, 295–307. doi:10.1105/tpc.105.038398.
30. Prunet, N.; Morel, P.; Thierry, A.-M.; Eshed, Y.; Bowman, J.L.; Negrutiu, I.; Trehin, C. *REBELOTE*, *SQUINT*, and *ULTRAPETALA1* function redundantly in the temporal regulation of floral meristem termination in *Arabidopsis thaliana*. *The Plant Cell* **2008**, *20*, 901–919, doi:10.1105/tpc.107.053306.
31. Maier, A.T.; Stehling-Sun, S.; Offenburger, S.-L.; Lohmann, J.U. The bZIP Transcription Factor *PERIANTHIA*: A Multifunctional Hub for Meristem Control. *Front. Plant Sci.* **2011**, *2*, 79, doi:10.3389/fpls.2011.00079.
32. Carles, C.C.; Choffnes-Inada, D.; Reville, K.; Lertpiriyapong, K.; Fletcher, J.C. *ULTRAPETALA1* encodes a SAND domain putative transcriptional regulator that controls shoot and floral meristem activity in *Arabidopsis*. *Development* **2005**, *132*, 897–911, doi:10.1242/dev.01642.
33. Xu, Y.; Prunet, N.; Gan, E.-S.; Wang, Y.; Stewart, D.; Wellmer, F.; Huang, J.; Yamaguchi, N.; Tatsumi, Y.; Kojima, M.; et al. *SUPERMAN* regulates floral whorl boundaries through control of auxin biosynthesis. *EMBO J.* **2018**, *37*, doi:10.15252/embj.201797499.
34. Yamaguchi, N.; Huang, J.; Xu, Y.; Tanoi, K.; Ito, T. Fine-tuning of auxin homeostasis governs the transition from floral stem cell maintenance to gynoecium formation. *Nat. Commun.* **2017**, *8*, 1125, doi:10.1038/s41467-017-01252-6.
35. Zhang, K.; Wang, R.; Zi, H.; Li, Y.; Cao, X.; Li, D.; Guo, L.; Tong, J.; Pan, Y.; Jiao, Y.; et al. *AUXIN RESPONSE FACTOR3* Regulates Floral Meristem Determinacy by Repressing Cytokinin Biosynthesis and Signaling. *The Plant Cell* **2018**, *30*, 324–346, doi:10.1105/tpc.17.00705.
36. Galli, M.; Gallavotti, A. Expanding the Regulatory Network for Meristem Size in Plants. *Trends Genet.* **2016**, *32*, 372–383, doi:10.1016/j.tig.2016.04.001.
37. Thompson, B.E.; Bartling, L.; Whipple, C.; Hall, D.H.; Sakai, H.; Schmidt, R.; Hake, S. *bearded-ear* encodes a MADS box transcription factor critical for maize floral development. *The Plant Cell* **2009**, *21*, 2578–2590, doi:10.1105/tpc.109.067751.

38. Ohmori, S.; Kimizu, M.; Sugita, M.; Miyao, A.; Hirochika, H.; Uchida, E.; Nagato, Y.; Yoshida, H. *MOSAIC FLORAL ORGANS1*, an AGL6-like MADS box gene, regulates floral organ identity and meristem fate in rice. *The Plant Cell* **2009**, *21*, 3008–3025, doi:10.1105/tpc.109.068742.
39. Sakakibara, K.; Reisewitz, P.; Aoyama, T.; Friedrich, T.; Ando, S.; Sato, Y.; Tamada, Y.; Nishiyama, T.; Hiwatashi, Y.; Kurata, T.; et al. *WOX13-like* genes are required for reprogramming of leaf and protoplast cells into stem cells in the moss *Physcomitrella patens*. *Development* **2014**, *141*, 1660–1670, doi:10.1242/dev.097444.
40. Nardmann, J.; Werr, W. Symplesiomorphies in the *WUSCHEL* clade suggest that the last common ancestor of seed plants contained at least four independent stem cell niches. *New Phytol.* **2013**, *199*, 1081–1092, doi:10.1111/nph.12343.
41. Sauquet, H.; Balthazar, M. von; Magallón, S.; Doyle, J.A.; Endress, P.K.; Bailes, E.J.; Barroso de Morais, E.; Bull-Hereñu, K.; Carrive, L.; Chartier, M.; et al. The ancestral flower of angiosperms and its early diversification. *Nat. Commun.* **2017**, *8*, 16047, doi:10.1038/ncomms16047.
42. Endress, P.K. Evolutionary diversification of the flowers in angiosperms. *Am. J. Bot.* **2011**, *98*, 370–396, doi:10.3732/ajb.1000299.
43. Becker, A.; Gleissberg, S.; Smyth, D.R. Floral and Vegetative Morphogenesis in California Poppy (*Eschscholzia californica* Cham.). *International Journal of Plant Sciences* **2005**, *166*, 537–555, doi:10.1086/429866.
44. The Angiosperm Phylogeny Group; Chase, M.W.; Christenhusz, M.J.M.; Fay, M.F.; Byng, J.W.; Judd, W.S.; Soltis, D.E.; Mabberty, D.J.; Sennikov, A.N.; Soltis, P.S.; et al. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot. J. Linn. Soc.* **2016**, *181*, 1–20, doi:10.1111/boj.12385.
45. Zhao, L.; Bachelier, J.B.; ZHANG, X.-H.; Ren, Y. Floral organogenesis in *Dysosma versipellis* (Berberidaceae) and its systematic implications. *Botany* **2016**, *94*, 359–368, doi:10.1139/cjb-2016-0019.
46. Zhang, X.H.; Ren, Y.I.; Tian, X.H. Floral morphogenesis in *Sinofranchetia* (Lardizabalaceae) and its systematic significance. *Bot. J. Linn. Soc.* **2009**, *160*, 82–92, doi:10.1111/j.1095-8339.2009.00835.x.
47. Wang, X.; Gong, J.; Zhao, L.; Che, X.; Li, H.; Ren, Y. Flower morphology and development of the monotypic Chinese genus *Anemoclema* (Ranunculaceae). *Plant Syst Evol* **2016**, *302*, 683–690, doi:10.1007/s00606-016-1298-2.
48. Tucker, S.C. Floral ontogeny in *Swartzia* (Leguminosae: Papilionoideae: Swartzieae): distribution and role of the ring meristem. *Am. J. Bot.* **2003**, *90*, 1271–1292, doi:10.3732/ajb.90.9.1271.
49. Tian, X.; Zhang, L.; Ren, Y. Development of flowers and inflorescences of *Circaeaster* (Circaeasteraceae, Ranunculales). *Plant Syst Evol* **2005**, *256*, 89–96, doi:10.1007/s00606-005-0377-6.
50. Schuchovski, C.; Meulia, T.; Sant'Anna-Santos, B.F.; Fresnedo-Ramírez, J. Inflorescence Development and Floral Organogenesis in *Taraxacum kok-saghyz*. *Plants (Basel)* **2020**, *9*, 1258, doi:10.3390/plants9101258.
51. Ren, Y.; Li, H.-F.; Zhao, L.; Endress, P.K. Floral morphogenesis in *Euptelea* (Eupteleaceae, Ranunculales). *Ann Bot* **2007**, *100*, 185–193, doi:10.1093/aob/mcm106.
52. Endress, P.K.; Matthews, M.L. Progress and problems in the assessment of flower morphology in higher-level systematics. *Plant Syst Evol* **2012**, *298*, 257–276, doi:10.1007/s00606-011-0576-2.
53. Endress, P.K. Flower Structure and Trends of Evolution in Eudicots and Their Major Subclades 1. *Annals of the Missouri Botanical Garden* **2010**, *97*, 541–583, doi:10.3417/2009139.
54. Endress, P.K. Angiosperm Floral Evolution: Morphological Developmental Framework. In *Advances in botanical research: Incorporating advances in plant pathology*; Callow, J.A., Ed.; Elsevier Academic Press: Amsterdam, Netherlands, Boston, MA, 2006; pp 1–61, ISBN 9780120059447.
55. Endress, P.K. The families and Genera of Vascular Plants II. *Flowering Plants* **1993**, 599–602.