

Article

Impact of Rowing Training on Quality of Life and Physical Activity Levels in Female Breast Cancer Survivors

Juan Gavala-González ^{1,2}, Amanda Torres-Pérez ^{2,3*} and José Carlos Fernández-García ^{2,3}

¹ Department of Physical Education and Sports, University of Seville, Seville, Spain; jgavala@us.es

² Researching in Sport Science: Research group (CTS-563) of the Andalusian Research Plan.

³ Department of Didactics of Languages, Arts and Sport, University of Malaga, Andalucía-Tech, IBIMA, Málaga, Spain; jcfcg@uma.es

* Correspondence: amandatorres@uma.es

Abstract: The aim of this study was to determine whether a rowing training program improved the quality of life and the physical activity levels in female breast cancer survivors ($n = 28$) (stage I - 4.54%; II - 36.36%; III - 54.54%; and IV - 4.54%) diagnosed 4.68 ± 3.00 years previously, who had undergone a subsequent intervention (preservation 56.53% and total mastectomy 43.47%) and had a current mean age of 52.30 ± 3.78 years. The participants ($n = 28$) engaged in a 12-week training program, each week comprising three sessions and each session lasting 60–90 min. The short form of the International Physical Activity Questionnaire (IPAQ-SF) and the Short Form 36 Health Survey (SF-36) were also administered. The results showed statistically significant improvements in levels of physical activity and in the dimensions of quality of life. We can conclude that a 12-week rowing training program tailored to women who have had breast cancer increases physical activity levels, leading to improved health status and quality of life.

Keywords: breast cancer; rowing; exercise; quality of life; perceived health; IPAQ-SF; SF-36.

1. Introduction

Cancer is the second leading cause of death worldwide, representing about 9.6 million deaths in 2018, which means that one in six deaths globally is due to this disease [1,2]. In women, breast cancer is the most common cancer, affecting around 2.1 million women in 2018, i.e., one in four cancers diagnosed is breast cancer [2–4].

The rise in the number of cancer cases diagnosed in recent years has been associated with population growth, closely linked to increased life expectancy and therefore with aging, considering age as a fundamental risk factor for developing cancer. It has also been related to the increase in early detection, improvements in primary care and in early diagnosis programs which, although they lead to higher numbers of cases, are in turn related to a decrease in mortality [4–6].

One third of the diagnosed cancer cases could be prevented if exposure to various lifestyle-related risk factors were eliminated or reduced, such as the consumption of harmful substances such as smoking or alcohol, an unhealthy, high-calorie diet with a high intake of saturated animal fats and sugars, or a sedentary lifestyle [1,4,5,7]. A large body of scientific evidence shows that physical activity has positive effects on the general population, improving health status, mood, body composition, quality of life [8–12], and preventing the onset of numerous diseases, including various types of cancer, such as breast cancer [12,15,16].

Physical activity has been associated with a lower risk of developing breast cancer [13–15], with a decrease in the probability of relapse and with a higher survival rate [14–16]. In individuals with cancer, physical activity has benefits for their health: reduced fatigue, improved strength levels, and improved quality of life and physical function [14,17–20].

However, despite the evidence supporting physical activity, two out of three cancer patients do not perform the minimum levels of exercise recommended by the American College of Sports Medicine (ACSM), which considers it essential to perform 150 min of moderate aerobic activity or 75 min of vigorous aerobic activity per week and at least 2 days of resistance training [19,21,22].

The relationship between physical activity and breast cancer has been demonstrated in several studies that analyzed and compared the effect of different exercise programs in breast cancer survivors, finding significant improvements in quality of life [21,23], physical function and muscle strength [21,22]. More specifically, the study by Wiskemann et al. (2016) based on a 12-week resistance training program showed gains in muscle strength [24], and in several studies where the training programs combined endurance with aerobic exercise for 12 weeks, the improvements were significant in muscle strength, level of physical activity and quality of life [21,25–29]. In addition, studies on mixed programs combining aerobic and strength exercises [30,31], showed important improvements in aerobic capacity, maximum oxygen consumption, reduction in the percentage of fat mass, muscle strength and, above all, in quality of life. Improvements were also found in physical, psychological, social and quality of life parameters in training programs based on dragon boat rowing in women with breast cancer [19,32–34].

Women breast cancer survivors have found rowing to be an activity that improves the sequels of the disease [35], such as reducing pain, increasing the range of movement in the upper limbs, improving muscle activation and increasing strength and muscle function [36,37].

In this sense, rowing is considered one of the most complete water sports, involving the work of the musculature of both the upper and lower limbs [38] and almost all the body's musculature [39]. It is a sport in which symmetrical movements are performed that do not require forced position and which combines the work of strength and aerobic endurance [19]. Several studies have shown that this type of activity improves the quality of life of cancer patients, including psychological, physical, social and emotional aspects, favoring their rehabilitation, self-esteem and normalizing their daily life [37,40,41].

In addition, the practice of rowing has psychosocial benefits for cancer survivors [37,42], such as the fact that it is a team sport promotes the development of social relationships, they find the support they need in other women who have gone through or are going through the same situation or that it is an outdoor activity, which provides them with extra motivation to adhere to physical activity and improves their quality of life [41,43,44].

Finally, in studies using the short form of the International Physical Activity Questionnaire (IPAQ-SF) to examine changes in physical activity [28,29] and the Short Form 36 Health Survey (SF-36) for changes in quality of life [27,30] following physical activity interventions in breast cancer patients, it was found that activity levels increased and the different domains of quality of life improved after completion of the training programs.

The purpose of the present study is to determine the influence of a 12-week rowing training program on quality of life and physical activity levels in women who have survived breast cancer.

2. Materials and Methods

2.1. Design and Participants

Participants ($n = 28$) aged 52.3 ± 3.8 years were recruited under the condition of having overcome breast cancer. The women were diagnosed 4.7 ± 3.0 years earlier, had different stages of disease and had undergone surgery as shown in Table 1.

Breast Cancer Survivors (BCS) is the name given to women who have been diagnosed with breast cancer and have had to undergo surgery and chemotherapy and/or radiotherapy treatments.

To take part in this research, we searched among various breast cancer associations in Malaga for women who wanted to do sport (rowing) and who met the following

characteristics: having overcome breast cancer, having completed chemotherapy and radiotherapy treatments, and having the oncologist's approval to do physical activity. No more than 10 years had passed since the cancer diagnosis. All of them were still taking tamoxifen.

The sample was selected based on compliance with these inclusion criteria.

Table 1. Characteristics of the breast cancer survivor sample.

Age (Years)	52.30 ± 3,78	
Years from Diagnosis	4.68 ± 3.00	
Stage (%)	I	4,54
	II	36,36
	III	54,54
	IV	4,54
Surgery (%)	Preservation	56,53
	Total Mastectomy	43,47

The study was carried out at the RC Mediterráneo in Málaga involving women from the Sport Association Málaga D.B. 48 participants were invited to take part: 10 initially withdrew due to compatibility/work/family and transport problems, and 10 were excluded because they did not attend 90% of the sessions.

After the initial selection, the nature of the study was explained to the participants, indicating that their anonymity would be maintained at all times, following the ethical considerations of the Sport and Exercise Science Research [45], as well as the principles included in the Declaration of Helsinki [46], which define the ethical guidelines for research in human subjects. The University of Malaga assigned the identification number 65-2020-H, which is registered with the Ethics Committee. The participants provided written informed consent and throughout the intervention and afterwards, we acted under the provisions of the Organic Law 3/2018, of December 5, on the Protection of Personal Data and Guarantee of Digital Rights, regarding the protection of personal data under Spanish legislation. After signing the informed consent, the physical activity (IPAQ-SF) and the health-related quality of life (SF-36) questionnaires were administered.

The intervention lasted 12 uninterrupted weeks in which the women carried out two weekly sessions as described above. Both at the beginning and at the end of the program, the participants were asked to complete the questionnaires.

2.2. Instruments

The participants also completed the short version of the International Physical Activity (IPAQ-SF) questionnaire to assess physical activity levels over the last 7 days. This questionnaire consists of seven questions which have acceptable measurement properties to monitor physical activity levels for adults aged 18 to 65 years in various settings and also reports the number of metabolic equivalents (METs) over the last 7 days [47]. Several studies have demonstrated the reliability of the IPAQ-SF for measuring the level of physical activity or the number of METs achieved during the last 7 days, obtaining similar results to other types of tests such as accelerometry or podometry [48–50].

The SF-36 Health Survey was used to assess health-related quality of life. This questionnaire consists of 36 items that report both positive and negative health status covering eight dimensions: physical function, social function, physical role, emotional role, mental health, vitality, bodily pain, and general health.

2.3. Intervention

Before starting the training program, participants were asked to complete the questionnaires: IPAQ-SF and SF-36 Health Survey and, in addition, they filled in the informed consent document to participate in the study.

The 12-week rowing training program was carried out at the RC Mediterráneo in Málaga, and it was divided into three parts of 4 weeks each. These stages progressively increased in intensity and were regulated through the participants' subjective perception of effort, using the Börg scale [51].

- Initial phase: mobility, proprioceptive and postural control exercises. Main phase with rowing training. Final phase with stretching. Börg scale 5-6.
- Intermediate phase: mobility, proprioceptive and postural control exercises. Main phase with rowing training. Final phase with stretching. Börg scale 6-7.
- Final phase: mobility, proprioceptive and postural control exercises. Main phase with rowing training. Final phase with stretching. Börg scale 7-8.

Throughout the program, a weekly schedule was established consisting of three training days lasting 60–90 min per session. These sessions were supervised by a trainer who ensured attendance, correct execution of the tasks and intensity of the sessions, in addition to excluding from the study those subjects who did not comply with at least 90% participation. All the exercises in these sessions were performed in group. Exercises were generic and adjusted for people who have never rowed before. Each of the training sessions had the same structure:

1. Initial phase: performed with warm-up, mobility, proprioceptive and postural control exercises, all exercises carried out in a multipurpose room (10–15 min).
2. Intermediate phase: performed in the Mediterranean Sea near the port of Malaga (Cruise Terminal/Malagueta Beach) using fixed-bench boats, typical of the Spanish Mediterranean, called *Llauts*, which are propelled by eight rowers and a coxswain or skipper [52], with each rower holding a single oar (40–60 min).
3. Final phase: flexibility exercises to relax the musculature and bring the body back to its initial state after exercise (10–15 min).

Both at the beginning and at the end of the program, sessions were held to discuss issues, and the participants were asked to complete the questionnaires and to visit the Human Motricity Laboratory at the University of Malaga for anthropometric (weight and height) and body composition (DEXA) assessments.

2.4. Data Analysis

All analyses were performed with IBM SPSS, version 25. The significance level was defined as $p < 0.05$. The fit of the different variables to the normal distribution was assessed using both graphic procedures and the Shapiro–Wilk test.

To examine the differences resulting from the rowing training performed by the participants, the medians of each variable pre- and post-intervention were analyzed using the Wilcoxon test for related samples (paired data). In addition, the graphic analysis of the different variables was carried out using box and whisker plots.

3. Results

The descriptive analysis of the different study variables (Table 2), level of physical activity and quality of life are shown below, differentiating between pre-intervention and post-intervention values.

Table 2. Descriptive analysis of the variables pretest and posttest.

Variables	Mean		Percentiles						Minimum		Maximum	
			25		50 (Mdn)		75					
	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
IPAQ Walking (METS)	587,8 ± 432.1	1522,1 ± 1353.1	305,3	693,0	495,0	1287,0	693,0	1683,0	0	231	1485	6930

IPAQ Moderate (METs)	185,0 ± 260,0	1250,0 ± 1664,2	0,0	480,0	0,0	720,0	240,0	1140,0	0	240	960	8400
IPAQ Vigorous (METs)	416,7 ± 513,1	1620,0 ± 1608,0	0,0	780,0	80,0	1440,0	960,0	1920,0	0	0	1440	7680
IPAQ Total (METs)	1189,5 ± 835,0	4392,1 ± 3341,7	495,0	2388,0	1075,5	3483,0	1611,0	5442,7	0	1413	3108	16290
IPAQ Sitting (METs)	267,8 ± 155,4	197,8 ± 135,2	120,0	104,3	240,0	150,0	435,0	285,0	30	0	540	520
Physical Function (Points)	75,4 ± 18,8	85,2 ± 13,3	70,0	77,5	82,5	90,0	90,0	93,8	30	50	95	100
Physical Role (Points)	62,5 ± 26,2	76,6 ± 25,4	37,5	62,5	71,9	84,4	85,9	100,0	6,3	12,5	100	100
Bodily Pain (Points)	49,7 ± 18,6	59,3 ± 22,2	41,0	43,5	51,5	61,0	62,0	72,0	10	22	84	100
General Health (Points)	60,1 ± 17,6	67,1 ± 19,9	46,3	50,5	61,0	72,0	72,0	85,8	20	30	92	97
Vitality (Points)	50,0 ± 12,9	59,8 ± 13,4	45,0	50,0	50,0	60,0	60,0	70,0	20	25	70	80
Social Function (Points)	72,4 ± 22,7	87,5 ± 18,4	62,5	75,0	75,0	100,0	87,5	100,0	12,5	37,5	100	100
Emotional Role (Points)	76,0 ± 15,8	89,9 ± 14,7	62,5	77,1	75,0	100,0	89,6	100,0	50	50	100	100
Mental Health (Points)	58,8 ± 13,9	67,0 ± 11,9	48,0	56,0	58,0	68,0	71,0	76,0	32	36	84	84

Pre = Pretest; Post = Posttest; Mdn = Median IPAQ = International Physical Activity Questionnaire; METs = metabolic equivalents; 1 MET = 3,5 mL O₂ * kg⁻¹ * min⁻¹.

For the variables associated with engaging in physical activity obtained from the IPAQ-SF, Table 3 displays the medians of each, as well as the results of the Wilcoxon Test to determine whether significant differences exist between the pretest and posttest data.

Table 3. Statistical analysis of the Physical Activity variables (IPAQ-SF).

Variables	Mdn (50)		Wilcoxon Test		
	Pre	Post	T	z	p
IPAQ Walking (METs)	495,0	1287,0	276,00	- 4,201	0,000**
IPAQ Moderate (METs)	0,0	720,0	300,00	-4,314	0,000**
IPAQ Vigorous (METs)	80,0	1440,0	231,00	- 4,043	0,000**
IPAQ Total (METs)	1075,5	3483,0	300,00	- 4,286	0,000**
IPAQ Sitting (METs)	240,0	150,0	23,50	-3,075	0,002*

* = $p < 0,05$; ** = $p < 0,001$; Pre = Pretest; Post = Posttest; Mdn = Median; T = t test statistic; z = z-score; IPAQ = International Physical Activity Questionnaire; METs = metabolic equivalents; 1 MET = 3,5 mL O₂ * kg⁻¹ * min⁻¹.

Figure 1 depicts all the physical activity variables, showing improvements after the intervention in levels of Walking, Moderate, Vigorous and Total Physical Activity, all of which are significant. The lower value obtained for the IPAQ Sitting variable after the intervention compared to before the intervention indicates the participants were more active and spent less time sitting throughout the week.




Figure 1. Physical Activity levels (IPAQ-SF); PRE = Pretest; POST = Posttest; METS = metabolic equivalents

Regarding the quality of life of the participants, the differences between the results obtained in the various measurements through the SF-36 questionnaire can be seen in Table 4.

Table 4. Statistical analysis of the Quality-of-Life variables (SF-36).

Variables	Mdn (50)		Wilcoxon Test		
	Pre	Post	T	z	p
Physical Function (Pts)	82,5	90,0	117	- 3,256	0,001*
Physical Role (Pts)	71,9	84,4	226	- 3,866	0,000**
Bodily Pain (Pts)	51,5	61,0	186,5	- 2,472	0,013*
General Health (Pts)	61,0	72,0	188	-2,006	0,045*
Vitality (Pts)	50,0	60,0	232	- 2,879	0,004*
Social Function (Pts)	75,0	100,0	130	- 3,247	0,001*
Emotional Role (Pts)	75,0	100,0	160	- 3,268	0,001*
Mental Health (Pts)	58,0	68,0	180,5	- 2,836	0,005*

Mdn = Median; Pre = Pretest; Post = Posttest; Pts = Points; * = $p < 0,05$; ** = $p < 0,001$,

Figure 2 and Figure 3 displays the results for the variables associated with quality of life, mental and physical dimensions, before and after the intervention.




Figure 2. Quality of Life (SF-36) – Mental Dimensions; PRE = Pretest; POST = Posttest.

Focusing on some of the physical dimensions, Figure 3 shows the variation in bodily pain. That this value is higher after the intervention may indicate that the participants report less pain in general. In other words, engaging in rowing decreases bodily pain in women who are breast cancer survivors or increases the pain threshold in the women studied. Also, illustrates the improvement after the intervention in the physical role, which indicates that the participants have fewer limitations when performing any physical activity compared to prior to the intervention. This suggests that a 12-week rowing training program, adapted to women who have had breast cancer, helps to improve the perceived ability to perform other activities.




Figure 3. Quality of Life (SF-36) – Physical Dimensions; PRE = Pretest; POST = Posttest.

Finally, in terms of overall perceived health of the participants, a significant improvement was detected after the physical activity intervention (Figure 4), demonstrating that after completing the rowing program, a tendency towards an improved perception of health emerged: The number of women who reported having poor or fair health decreased and those who claimed having “very good health” increased, with 75% of the participants reporting their health status to be good or very good.




Figure 4. Variation in Perceived Health; Pts = Points.

In summary, after an intervention based on a 12-week rowing training program tailored to women who have had breast cancer, all the variables included, both those concerning level of physical activity and quality of life, showed significant improvements. That is, the participants were more physically active and less sedentary throughout the week and, in addition, their perceived physical, emotional, and mental health status improved, which indicates fewer limitations in terms of physical activity, social life and vitality.

4. Discussion

To date, studies relating rowing to improvements in quality of life or in the level of physical activity in breast cancer survivors are practically non-existent. In this sense, some research can be found in which other types of sports practice, such as dragon boating [32,33], yoga and pilates [53,54] or endurance, strength or aerobic exercise training programs [22,24,31], report improvements at a physical, psychological and social level in women breast cancer survivors.

More specifically, upon examining studies that used different types of physical activity programs, the 12-week resistance training program proposed by Wiskemann et al. (2016) reported improvements only in muscle strength [24]. Other programs that combined aerobic and strength exercises for 24 weeks led to improvements in muscle strength, aerobic capacity and only some of the quality of life dimensions [31], while those that combined resistance and aerobic exercises for 16 weeks showed improvements in the quality of life and physical fitness of the participants [26]. Harris (2012) and McDonough et al. (2018) reported that dragon boat rowing [32,33] led to physical, psychological and social improvements in women breast cancer survivors.

In our study, based on a rowing training program lasting only 12 weeks, improvements were observed in all levels of physical activity as measured by the IPAQ-SF, increasing the levels of walking, moderate and vigorous physical activity, as well as total physical activity and decreasing the sedentary activity of the participants, and improvements were shown in all dimensions of quality of life. The results obtained in the present research are superior to those reported in previous studies and, in addition, a shorter intervention time was required.

These results may be due to the fact that rowing is a sport that involves the muscles of both the lower and upper limbs and almost every muscles of the body[38]. In addition, rowing is a cyclic and symmetrical sport that combines overall strength with aerobic endurance [19] and it is based on cyclic and alternating movements of flexion-extension of the limbs and stabilization of the trunk and back muscles [55]. In this sense, the

involvement of the whole body in physical activity, such as rowing, leads to improvements in quality of life, physical function or reduction of body fat in women breast cancer survivors [33,56,57].

In the study by Park et al. (2019), through a training program combining aerobic and resistance exercises, physical activity levels increased after the 12-week intervention, rising in 63,4% of the participants. At baseline, 33% of participants were inactive, 49.6% minimally active and 17.4% health-enhancing physical activity (HEPA) active, and after intervention the data improved, being the percentages 15.3%, 50.4% and 34.2% respectively [28]. It is of interest to compare these results with those of our study, which uses an intervention of the same duration. In our study, overall physical activity levels increased in all the participants after the rowing program with respect to the initial measurement. It should also be noted that prior to the intervention, about 50% of the participants reported no moderate or vigorous physical activity, whereas afterwards, all the participants reported engaging in moderate physical activity and less than 10% reported no vigorous activity.

Regarding quality of life parameters, the study by Mascherini et al. (2020) shows improvements in Physical Function, Social Function, General Health and Mental Health after 6 months of training [58] and Di Blasio et al. (2017) after 12 weeks of intervention only found improvements in Physical Function, Physical Role, Bodily Pain and General Health [22]. Dolan et al. (2018) also showed that after a 22-week training program all SF-36 quality of life dimensions improved with the exception of Bodily Pain [59].

Some of the limitations of this study reside in the scarcity of the sample which, although it presents homogeneity in the variables evaluated for this article, it would be possible to contemplate the possibility of knowing what the effects of this rowing program are from different times since the cancer was diagnosed or whether mastectomy is present or not.

The results of the present study demonstrate that rowing training has a greater influence on quality of life in breast cancer survivors than previous studies, as significant improvements were found in all the quality-of-life parameters after an intervention of shorter duration. This implies that both perceived physical, emotional, and mental health status and the perceived ability to perform other activities improved to a greater extent after our rowing-based training program. Indeed, 75% of the participants reported their health status as good or very good after the intervention, with a considerable increase compared to the initial data.

5. Conclusions

While studies have reported improved quality of life and reduced physical inactivity in female breast cancer survivors through training programs such as aerobic, strength, resistance, and dragon boat exercises. In the present study, we have shown that an intervention of just twelve weeks in length using rowing training tailored to women who have had breast cancer produced improvements in all levels of physical activity and reduced physical inactivity with respect to the initial measurement. In addition, after the intervention, the physical, emotional, and mental health status of the participants improved, leading to fewer limitations in their daily routines, physical activity, social life, and vitality. Our rowing program showed greater benefits in health and quality of life than other studies of various durations. We can therefore conclude that rowing training contributes to increasing daily physical activity and to improving health status, as well as the different dimensions of quality of life in women who have survived breast cancer.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided, The following statements should be used “Conceptualization, J,G,G, and A,T,P;; methodology, J,C,F,G and J,G,G;; software, A,T,P;; validation, J,C,F,G, and J,G,G;; formal analysis, J,C,F,G, and A,T,P;; investigation, J,C,F,G and J,G,G;; resources, J,C,F,G and J,G,G,; data curation, A,T,P;; writing—original draft preparation, J,G,G, and A,T,P;; writing—review and editing, J,G,G;; A,T,P, and J,C,F,G; visualization, J,G,G;; A,T,P, and

J,C,F,G,; supervision, J,C,F,G,; project administration, J,C,F,G and J,G,G,; funding acquisition, J,C,F,G. All authors have read and agreed to the published version of the manuscript,”

Funding: This research was funded by the “Researching in Sport Sciences” research group (CTS-563) of the Andalusian Research Plan,

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by Ethics Committee of University of Málaga (protocol code 65-2020-H and date of approval 30/09/2020),

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study,

Acknowledgments: We would like to thank the Real Club Mediterraneo of Malaga, its rowing captain: Juan Carlos Marfil Rodríguez and the team of the Malaga Dragon Boat BCS for opening their doors to us, giving us all the facilities in the world and for giving us the most important thing: their time and inspiring us with their impetus and energy, thanks to which we can present the first results of this study. To Maria Repice, for her help with the English version of this manuscript. Finally, we would also like to thank Alejandro Sánchez for his participation in the data collection for this study. All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. World Health Organization Fact Sheets: Cancer Available online: <https://www.who.int/news-room/fact-sheets/detail/cancer> (accessed on Aug 10, 2020).
2. International Agency for Research on Cancer (IARC) Cancer Today Available online: <https://gco.iarc.fr/today/home> (accessed on Aug 10, 2020).
3. World Health Organization Breast Cancer: prevention and control. Available online: <https://www.who.int/cancer/detection/breastcancer/en/> (accessed on Aug 12, 2020).
4. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA. Cancer J. Clin.* **2018**, *68*, 394–424, doi:10.3322/caac.21492.
5. Sociedad Española de Oncología Médica Las cifras del cáncer en España 2020. *Soc. Española Oncol. Médica* **2020**, *36*.
6. World Health Organization Early Cancer Diagnosis saves lives cuts treatments costs Available online: <https://www.who.int/es/news/item/03-02-2017-early-cancer-diagnosis-saves-lives-cuts-treatment-costs> (accessed on Dec 5, 2020).
7. World Health Organization Prevención del cáncer Available online: <https://www.who.int/cancer/prevention/es/> (accessed on Dec 5, 2020).
8. Loria Calderon, T.M.; Carmona Gómez, C.D. Efecto agudo del baile como ejercicio aeróbico sobre el balance estático en personas mayores de 50 años. *Rev. Iberoam. Ciencias la Act. Física y el Deporte.* **2020**, *9*, 61–74, doi:10.24310/riccafd.2020.v9i2.8022.
9. Gómez-Cabello, A.; Pardos-Mainer, E.; González-Gálvez, N.; Sagarra-Romero, L. Actividad Física y Calidad de Vida en las Personas Mayores: Estudio Piloto PQS. *Rev. Iberoam. Ciencias La Act. Fis. Y El Deporte.* **2018**, *8*, 96–109.
10. Warburton, D.E.R.; Bredin, S.S.D. Health benefits of physical activity: a systematic review of current systematic reviews. *Curr. Opin. Cardiol.* **2017**, *32*, 541–556, doi:10.1097/HCO.0000000000000437.
11. Gálvez Fernández, I. Pérdida de Peso y Masa Grasa con Auto-Cargas en Mujeres. *Rev. Iberoam. Ciencias la Act. Física y el Deporte.* **2017**, *6*, 30–37, doi:10.24310/riccafd.2017.v6i2.3780.
12. Díaz, J.; Muñoz, D.; Cordero, J.C.; Robles, M.C.; Courel-Ibañez, J.; Sánchez-Alcaraz, B.J. Estado de Ánimo y Calidad de Vida en Mujeres Adultas practicantes de Pádel. *Rev. Iberoam. Ciencias La Act. Fis. Y El Deporte.* **2018**, *8*, 34–43.
13. World Cancer Research Fund (WCRF); American Institute for Cancer Research (AICR) *Diet, Nutrition, Physical Activity and Cancer: a Global Perspective*; 2018; ISBN 9781912259465.
14. Johnsson, A.; Broberg, P.; Krüger, U.; Johnsson, A.; Tornberg, Å.B.; Olsson, H. Physical activity and survival following breast cancer. *Eur. J. Cancer Care (Engl).* **2019**, *28*, 1–8, doi:10.1111/ecc.13037.

15. Pollán, M.; Casla-Barrio, S.; Alfaro, J.; Esteban, C.; Seguí-Palmer, M.A.; Lucia, A.; Martín, M. Exercise and cancer: a position statement from the Spanish Society of Medical Oncology. *Clin. Transl. Oncol.* **2020**, *22*, 1710–1729, doi:10.1007/s12094-020-02312-y.
16. Schmidt, M.E.; Chang-Claude, J.; Vrieling, A.; Seibold, P.; Heinz, J.; Obi, N.; Flesch-Janys, D.; Steindorf, K. Association of pre-diagnosis physical activity with recurrence and mortality among women with breast cancer. *Int. J. Cancer* **2013**, *133*, 1431–1440, doi:10.1002/ijc.28130.
17. Pennington, K.P.; McTiernan, A. The role of physical activity in breast and gynecologic cancer survivorship. *Gynecol. Oncol.* **2018**, *149*, 198–204, doi:10.1016/j.ygyno.2018.01.020.
18. Mascherini, G.; Tosi, B.; Giannelli, C.; Grifoni, E.; Degl'Innocenti, S.; Galanti, G. Breast cancer: effectiveness of a one-year unsupervised exercise program. *J. Sports Med. Phys. Fitness* **2018**, *59*, 283–289, doi:10.23736/S0022-4707.18.08131-8.
19. Gavala-González, J.; Gálvez-Fernández, I.; Mercadé-Melé, P.; Fernández-García, J.C. Rowing training in breast cancer survivors: A longitudinal study of physical fitness. *Int. J. Environ. Res. Public Health* **2020**, *17*, 1–12, doi:10.3390/ijerph17144938.
20. Furmaniak, A.; Menig, M.; Markes, M. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst. Rev.* **2016**, doi:10.1002/14651858.CD005001.pub3.www.cochranelibrary.com.
21. Courneya, K.S.; McKenzie, D.C.; Mackey, J.R.; Gelmon, K.; Friedenreich, C.M.; Yasui, Y.; Reid, R.D.; Cook, D.; Jespersen, D.; Proulx, C.; et al. Effects of Exercise Dose and Type During Breast Cancer Chemotherapy: Multicenter Randomized Trial. *J. Natl. Cancer Inst.* **2013**, *105*, 1821–1832, doi:10.1093/jnci/djt297.
22. Di Blasio, A.; Morano, T.; Cianchetti, E.; Gallina, S.; Bucci, I.; Di Santo, S.; Tinari, C.; Di Donato, F.; Izzicupo, P.; Di Baldassarre, A.; et al. Psychophysical health status of breast cancer survivors and effects of 12 weeks of aerobic training. *Complement. Ther. Clin. Pract.* **2017**, *27*, 19–26, doi:10.1016/j.ctcp.2017.01.004.
23. Mishra, S.I.; Scherer, R.W.; Snyder, C.; Geigle, P.; Gotay, C. Are Exercise Programs Effective for Improving Health-Related Quality of Life Among Cancer Survivors? A Systematic Review and Meta-Analysis. *Oncol. Nurs. Forum* **2014**, *41*, E326–E342, doi:10.1188/14.ONF.E326-E342.
24. Wiskemann, J.; Schmidt, M.E.; Klassen, O.; Debus, J.; Ulrich, C.M.; Potthoff, K.; Steindorf, K. Effects of 12-week resistance training during radiotherapy in breast cancer patients. *Scand. J. Med. Sci. Sport.* **2016**, *27*, 1500–1510, doi:10.1111/sms.12777.
25. Soriano-Maldonado, A.; Carrera-Ruiz, Á.; Díez-Fernández, D.M.; Esteban-Simón, A.; Maldonado-Quesada, M.; Moreno-Poza, N.; García-Martínez, M.D.M.; Alcaraz-García, C.; Vázquez-Sousa, R.; Moreno-Martos, H.; et al. Effects of a 12-week resistance and aerobic exercise program on muscular strength and quality of life in breast cancer survivors: Study protocol for the EFICAN randomized controlled trial. *Medicine (Baltimore)*. **2019**, *98*, 1–9, doi:10.1097/MD.00000000000017625.
26. Dieli-Conwright, C.M.; Courneya, K.S.; Demark-Wahnefried, W.; Sami, N.; Lee, K.; Sweeney, F.C.; Stewart, C.; Buchanan, T.A.; Spicer, D.; Tripathy, D.; et al. Aerobic and resistance exercise improves physical fitness, bone health, and quality of life in overweight and obese breast cancer survivors: A randomized controlled trial. *Breast Cancer Res.* **2018**, *20*, 1–10, doi:10.1186/s13058-018-1051-6.
27. Thomas, G.A.; Cartmel, B.; Harrigan, M.; Fiellin, M.; Capozza, S.; Zhou, Y.; Ercolano, E.; Gross, C.P.; Hershman, D.; Ligibel, J.; et al. The effect of exercise on body composition and bone mineral density in breast cancer survivors taking aromatase inhibitors. *Obesity* **2017**, *25*, 346–351, doi:10.1002/oby.21729.
28. Park, S.W.; Lee, I.; Kim, J. Il; Park, H.; Lee, J.D.; Uhm, K.E.; Hwang, J.H.; Lee, E.S.; Jung, S.Y.; Park, Y.H.; et al. Factors associated with physical activity of breast cancer patients participating in exercise intervention. *Support. Care Cancer* **2019**, *27*, 1747–1754, doi:10.1007/s00520-018-4427-3.
29. Uhm, K.E.; Yoo, J.S.; Chung, S.H.; Lee, J.D.; Lee, I.; Kim, J. Il; Lee, S.K.; Nam, S.J.; Park, Y.H.; Lee, J.Y.; et al. Effects of exercise intervention in breast cancer patients: is mobile health (mHealth) with pedometer more effective than conventional program using brochure? *Breast Cancer Res. Treat.* **2017**, *161*, 443–452, doi:10.1007/s10549-016-4065-8.
30. Smith, T.M.; Broomhall, C.N.; Crecelius, A.R. Physical and Psychological Effects of a 12-Session Cancer Rehabilitation Exercise Program. *Clin. J. Oncol. Nurs.* **2016**, *20*, 653–659, doi:10.1188/16.CJON.653-659.
31. De Luca, V.; Minganti, C.; Borriore, P.; Grazioli, E.; Cerulli, C.; Guerra, E.; Bonifacino, A.; Parisi, A. Effects of concurrent aerobic

- and strength training on breast cancer survivors: a pilot study. *Public Health* **2016**, *136*, 126–132, doi:10.1016/j.puhe.2016.03.028.
32. Harris, S.R. Were all in the same boat: A review of the benefits of dragon boat racing for women living with breast cancer. *Evidence-based Complement. Altern. Med.* **2012**, *2012*, 1–8, doi:10.1155/2012/167651.
 33. McDonough, M.H.; Patterson, M.C.; Weisenbach, B.B.; Ullrich-French, S.; Sabiston, C.M. The difference is more than floating: factors affecting breast cancer survivors' decisions to join and maintain participation in dragon boat teams and support groups. *Disabil. Rehabil.* **2018**, *41*, 1788–1796, doi:10.1080/09638288.2018.1449259.
 34. Fong, A.J.; Saxton, H.R.; Kauffeldt, K.D.; Sabiston, C.M.; Tomasone, J.R. “We’re all in the same boat together”: exploring quality participation strategies in dragon boat teams for breast cancer survivors. *Disabil. Rehabil.* **2020**, *0*, 1–12, doi:10.1080/09638288.2020.1733676.
 35. McKenzie, D.C. Abreast in a boat - A race against breast cancer. *CMAJ* **1998**, *159*, 376–378.
 36. Asensio-García, M. del R.; Tomás-Rodríguez, M.I.; Palazón-Bru, A.; Hernández-Sánchez, S.; Nouni-García, R.; Romero-Aledo, A.L.; Gil-Guillén, V.F. Effect of rowing on mobility, functionality, and quality of life in women with and without breast cancer: a 4-month intervention. *Support. Care Cancer* **2020**, doi:10.1007/s00520-020-05757-7.
 37. Iacrossi, L.; Gambalunga, F.; Molinaro, S.; De Domenico, R.; Giannarelli, D.; Fabi, A. The effectiveness of the sport “dragon boat racing” in reducing the risk of lymphedema incidence: An observational study. *Cancer Nurs.* **2019**, *42*, 323–331, doi:10.1097/NCC.0000000000000615.
 38. Das, A.; Mandal, M.; Syamal, A.K.; Majumdar, P. *Monitoring Changes of Cardio-Respiratory Parameters During 2000m Rowing Performance*; 2019; Vol. 12;.
 39. Yoshiga, C.C.; Higuchi, M. Rowing performance of female and male rowers. *Scand. J. Med. Sci. Sports* **2003**, *13*, 317–321, doi:10.1034/j.1600-0838.2003.00321.x.
 40. McNeely, M.L.; Campbell, K.L.; Courneya, K.S.; Mackey, J.R. Effect of acute exercise on upper-limb volume in breast cancer survivors: A pilot study. *Physiother. Canada* **2009**, doi:10.3138/physio.61.4.244.
 41. Ray, H.; Jakubec, S.L. Nature-based experiences and health of cancer survivors. *Complement. Ther. Clin. Pract.* **2014**, *20*, 188–192, doi:10.1016/j.ctcp.2014.07.005.
 42. Sablston, C.M.; McDonough, M.H.; Crocker, P.R.E. Psychosocial experiences of breast cancer survivors involved in a dragon boat program: Exploring links to positive psychological growth. *J. Sport Exerc. Psychol.* **2007**, *29*, 419–438, doi:10.1123/jsep.29.4.419.
 43. Hadd, V.; Sabiston, C.M.; McDonough, M.H.; Crocker, P.R.E. Sources of stress for breast cancer survivors involved in dragon boating: Examining associations with treatment characteristics and self-esteem. *J. Women's Heal.* **2010**, *19*, 1345–1353, doi:10.1089/jwh.2009.1440.
 44. McDonough, M.H.; Sabiston, C.M.; Ullrich-French, S. The development of social relationships, social support, and posttraumatic growth in a dragon boating team for breast cancer survivors. *J. Sport Exerc. Psychol.* **2011**, *33*, 627–648, doi:10.1123/jsep.33.5.627.
 45. Harriss, D.; Macsween, A.; Atkinson, G. Standards for Ethics in Sport and Exercise Science Research. *Int. J. Sports Med.* **2017**, *38*, 1126–1131.
 46. Ebihara, A. World medical association declaration of Helsinki. *Japanese Pharmacol. Ther.* **2000**, *28*, 983–986.
 47. Barrera, R. Cuestionario Internacional de actividad física (IPAQ). *Rev. Enfermería del Trab.* **2017**, *7*, 49–54.
 48. Kathleen, Y.; Wolin, D.; Heil, S.; Charles, E.; Gary, G. Validation of the International Physical Activity Questionnaire-Short Among Blacks. *J Phys Act Heal.* **2008**, *5*, 746–760.
 49. Cleland, C.; Ferguson, C.; Ellis, C.; Hunter, R. Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol.* **2018**, *18*, 176.
 50. Kurtze, N.; Rangul, V.; Hustvedt, B.E. Reliability and validity of the international physical activity questionnaire in the Nord-Trøndelag health study (HUNT) population of men. *BMC Med. Res. Methodol.* **2008**, *8*.
 51. Börg, G. Psychophysical bases of perceived exertion. *Med. Sci. Sports Exerc.* **1982**, *14*, 377–381, doi:10.1249/00005768-198205000-

00012.

52. Gavala-González, J. Las modalidades del remo: el remo en banco fijo. Available online: <http://tv.us.es/las-modalidades-del-remo-el-remo-en-banco-fijo/> (accessed on Jan 12, 2021).
53. Patsou, E.D.; Alexias, G.D.; Anagnostopoulos, F.G.; Karamouzis, M. V. Effects of physical activity on depressive symptoms during breast cancer survivorship: A meta-analysis of randomised control trials. *ESMO Open* **2017**, doi:10.1136/esmoopen-2017-000271.
54. Panchik, D.; Masco, S.; Zinnikas, P.; Hillriegel, B.; Lauder, T.; Suttman, E.; Chinchilli, V.; McBeth, M.; Hermann, W. Effect of Exercise on Breast Cancer-Related Lymphedema: What the Lymphatic Surgeon Needs to Know. *J. Reconstr. Microsurg.* **2019**, doi:10.1055/s-0038-1660832.
55. José González Aramendi, R.M. *Remo olímpico y remo tradicional: aspectos biomecánicos, fisiológicos y nutricionales Olympic rowing and traditional rowing: biomechanical, physiological and nutritional aspects*; 2014;
56. Lane, K.; Jespersen, D.; McKenzie, D.C. The effect of a whole body exercise programme and dragon boat training on arm volume and arm circumference in women treated for breast cancer. *Eur. J. Cancer Care (Engl)*. **2005**, *14*, 353–358, doi:10.1111/j.1365-2354.2005.00595.x.
57. Quintana, V.A.; Díaz, K.; Caire, G. Intervenciones para promover estilos de vida saludables y su efecto en las variables psicológicas en sobrevivientes de cáncer de mama: revisión sistemática. *Nutr. Hosp.* **2018**, *35*, 979–992.
58. Mascherini, G.; Tosi, B.; Giannelli, C.; Ermini, E.; Osti, L.; Galanti, G. Adjuvant Therapy Reduces Fat Mass Loss during Exercise Prescription in Breast Cancer Survivors. *J. Funct. Morphol. Kinesiol.* **2020**, *5*, 1–10, doi:10.3390/JFMK5030049.
59. Dolan, L.B.; Barry, D.; Petrella, T.; Davey, L.; Minnes, A.; Yantzi, A.; Marzolini, S.; Oh, P. The Cardiac Rehabilitation Model Improves Fitness, Quality of Life, and Depression in Breast Cancer Survivors. *J. Cardiopulm. Rehabil. Prev.* **2018**, *38*, 246–252, doi:10.1097/HCR.0000000000000256.