Article:

Title: Persistence of anti-SARS-CoV-2 antibodies six months after infection in an outbreak with five hundred COVID-19 cases in Borriana (Spain): A prospective cohort study.

Authors:

Salvador Domènech-Montoliu¹, Joan Puig-Barberà², Maria Rosario Pac-Sa³, Paula Vidal-Utrillas⁴, Marta Latorre-Poveda¹; Alba Del Rio-González⁴, Sara Ferrando-Rubert⁴, Gema Ferrer-Abad⁴, Manuel Sánchez-Urbano¹, Laura Aparisi-Esteve⁵, Gema Badenes-Marques¹, Belén Cervera-Ferrer¹, Ursula Clerig-Arnau¹, Claudia Dols-Bernad⁶, Maria Fontal-Carcel⁷, Lorna Gomez-Lanas¹, David Jovani-Sales⁵, Maria Carmen León-Domingo⁸, Maria Dolores Llopico-Vilanova¹, Mercedes Moros-Blasco⁵, Cristina Notari-Rodríguez¹, Raquel Ruíz-Puig¹, Sonia Valls-López¹ and Alberto Arnedo-Pena^{3,9,10}.

 1. Emergency Service Hospital de la Plana, Vila-real, Castellon, Spain. ; pttcarmen@hotmail.com (SDM) martalapo@hotmail.com (MRP) martalapo@hotmail.com (MSU) gemabamar@hotmail.com (GBM) belencerveraferrer@hotmail.com (BCF) ursuclerig@gmail.com (UCA) lornagl78@gmail.com (LGL) lopivila@hotmail.com (MDLLV) ; notari-cri@gva.es (CNR) raquelruizpuig@gmail.com (RRP) Sonia.valls.lopez@gmail.com (SVL).

2. Vaccines Research Area FISABIO, Valencia, Spain. jpuigb55@gmail.com

3. Public Health Center, Castelló de la Plana, Castellon, Spain. charopac@gmail.com (MRPS) albertoarnedopena@gmail.com (AAP)

4. Health Centers I and II, Borriana, Castellon, Spain. <u>vidalutrillaspaula@gmail.com</u> (PVU) <u>delrio_alb@gva.es</u> (ADRG) <u>sfr1812@gmail.com</u> (SFR) <u>jovasal1987@gamail.com</u> (GFA).

5. Carinyena Health Center, Vila-real, Castellon, Spain.<u>lauraaparisiesteve@gmail.com</u> (LAE) jovasal1987@gamail.com (DJS) mercedesmb1094@hotmail.com (MMB).

- 6. Health Center, Onda, Castellon, Spain. claudiadb1294@hotmail.com
- 7. Health Center, La Vall d'Uixó, Castellon, Spain. fontalcarcel93@hotmail.com
- 8. Villa Fátima School. Borriana. Castellon. Spain.; carmendole@hotmail.com
- 9. Department of Health Science, Public University Navarra. Pamplona, Spain.
- 10. Epidemiology and Public Health (CIBERESP), Madrid, Spain

Correpondence: Alberto Arnedo-Pena: albertoarnedopena@gmail.com. +35-622-573979

Abstract:

(cc)

In March 2020, several mass gathering events were related to the *Falles* festival in Borriana (Spain), resulting in a 536 laboratory-confirmed COVID-19 cases outbreak among participants. Our objective was to estimate anti-SARS-CoV-2 antibodies persistence six months after and factors associated with antibody response. A prospective population-based cohort study was carried out by the Public Health Center of Castellon and the Emergency and Clinical Analysis and Microbiology Services of Hospital de la Plana in Vila-real. In October 2020, sero-epidemiologic study to estimate the persistence of anti-SARS-CoV-2

antibodies by a electrochemiluminescence immunoassay (ECLIA) was implemented. We enrolled 484 (90.2%) of the 536 members of the initial outbreak cohort and detected persistent antibodies in 479 (99%) without re-infection episodes. Five participants had a negative antibody test. Factors associated with a negative result were a lower body mass index (BMI), and less contact with other COVID-19 cases. Among the 469 participants with two ECLIA tests, 96 (20.5%) had an increase of antibodies and 373 (79.5%) a decline. Increased antibodies were associated with older age, higher BMI, more severe illness, and low current smokers. After a COVID-19 infection, a high proportion of cases maintained detectable anti-SARS-CoV-2 antibodies.

Keywords: SARS-CoV-2, COVID-19, antibodies, cohort, population-based, body mass index, ECLIA.

1. Introduction

At the time of writing, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused 160,457,476 confirmed cases and 3,331,604 deaths globally [1]. One of the main unknowns is the duration of immunity elicited after coronavirus disease (COVID-19) in patients who have recovered. Other aspects of interest are the protection against SARS-CoV-2 re-infection, and factors related to the development of persistent SARS-Cov-2 immunity. Serologic surveys are needed to gather evidence on the persistence and protection of the immune response to SARS-CoV-2 infection [2-5].

In the first days of March 2020, several mass gathering events (MGEs) for the *Falles* festival took place in Borriana, a city of 35,000 inhabitants in Castellon (Spain), and a COVID-19 outbreak ensued. During March-June 2020, an epidemiologic study whit a serologic survey of this outbreak found 570 COVID-19 cases (536 laboratory-confirmed and 34 with clinical and epidemiologic criteria), 13 admissions and one death, among 1338 participants in the MGEs; the attack rate (AR) was 42.6 %.

The follow-up of a representative sample of the COVID-19 cases through a serologic survey to determine SARS-CoV-2 antibodies would offer the opportunity to know the duration of immunity after COVID19. Accordingly, our objective was to conduct a sero-epidemiological study, six months after the first study, to estimate the persistence of anti-SARS-CoV-2 antibodies among MGEs participants who suffered a laboratory-confirmed COVID-19 infection and the potential factors associated with the persistence and intensity of the immune response.

2. Materials and Methods

The study, a prospective population-base cohort study, was carried out by the Public Health Center of Castellon and the Emergency and Clinical Analysis and Microbiology Services of Hospital de la Plana in Vila-real. We invited all the subjects who tested positive for SARS-CoV-2 in the outbreak of the MGEs of *Falles* festival to participate in a second seroepidemiological study in October 2020.

The tests used to ascertain SARS-CoV-2 infection in the first study were: qualitative Electrochemiluminescence immunoassay (ECLIA) (Elecsys® Anti-SARS-CoV-2, Roche Diagnostics) [6], in 514 subjects. Lateral flow immunochromatographic assay (LFIC) (Healgen Scientific LLC for COVID-19 IgG/IgM rapid test cassette [7], in 15 subjects. Reverse-transcriptase polymerase chain reaction (RT-PCR) including LightMix® Modular Sarbecovirus E-gene with the LightCycler® 480 II system [8], in 39 subjects. In 32 subjects, we obtained both ECLIA and PCR results.

The outbreak cohort members with a laboratory-confirmed COVID-19 test were 536 cases from the first serologic survey between March and June 2020. The second serologic survey to determine anti-SARS-CoV-2 antibodies by ECLIA was implemented during October 2020, and 484 of the 536 members of the cohort took part (90.3%). Overall, 469 members had two ECLIA determinations after the second serologic survey (Figure 1).

Figure 1.Flow chart of Borriana COVID-19 outbreak cohort



Figure 1. Flow chart of Borriana COVID-19 outbreak cohort.

The ECLIA test, already described above, uses a modified double-antigen sandwich immunoassay using recombinant SARS-CoV-2 nucleocapsid protein (N) [6]. The test results are reported as numeric values, considering a reactive cutoff index (COI)>1.0 as a positive result and a COI <0.1 as a negative result.

We explore factors associated with the immune response, comparing the geometric mean of total anti-SARS-CoV-2 antibodies of participants in the first and second serologic surveys and the change in magnitude between tests as an increase or a decline. We also explored the potential associations [9-12] with COVID19 of ABO blood groups, ascertained by the gel test (ID-Card ABO/RhD, DiaMed GmbH, Bio-Rad Laboratories Switzerland) [13] and 25-hydroxy vitamin D [25(OH)D] levels, determined by electrochemiluminisence-based assay (Elecsys vitamin D total II Roche Diagnostic, Germany) [14].

Health staff of the Hospital de la Plana, and the health centers of Borriana, Vila-real, Onda, and La Vall d'Uixo conducted a telephone questionnaire survey in October 2020, asking the participants about their health situation, the occurrence of disease in the last six months, medical assistance, evolution of COVID-19 disease, sequelae, SARS-CoV-2 re-infections, and subsequent exposures. We obtained information about other potential factors associated with the immune response such as age, sex, weight, height, body mass index (BMI) (kg/m²), occupation, level of physical exercise, smoking habits, consumption of alcohol, chronic illness, and COVID-19 exposures, either in the household or in other settings from the questionnaire survey during May-June 2020. **Statistical analysis**

We used the Chi2 test and Fisher exact test for qualitative variables, Kruskal-Wallis test for quantitative variables, and Wilcoxon matched-pairs signed-rank test and equality of median for matched pairs test to compare the changes in anti-SARS-CoV-2 antibodies between the two serologic surveys. Positive or negative detection of anti-SARS-CoV-2 antibodies in the second respect to the first survey was the dependent variable, and we estimated the associations between this variable and several factors by relative risks (RR) through univariate and multivariate (adjusted RR) Poisson regression, robust or exact according to models conditions, with 95% confidence intervals (CI). Besides, we analysed the increase and decline in anti-SARS-CoV-2 antibody arbitrary units following the same statistical approach. As Vitamin D was only measured in the second serologic survey, we did not include it in multivariate models. After a review of SARS-CoV-2 medical literature, we studied the potential confounding

factors by Directed Acyclic Graphs (DAGs)(15) using the DAGitti 3.0 program [16]. With a sample size of 454 cohort members, the study had a power of 80% and alpha error of 5% to detect at least a 2% change in the fraction of positives between the first and second serologic surveys. We used Stata ® version 14 statistical program for all calculations and estimates.

The following of the study's cohort was a part of the public health surveillance as a prolongation of the COVID-19 outbreak control measures and the response of the COVID-19 pandemic. It was exempt from the Ethics Review Board approval's protocol according to the Spanish legislation. The study was approved by the director of Public Health Center of Castelló and the management of the Health Department of la Plana. All participants or the parents of minors provided the informed written consent to be included in the study.

3. Results

A total of 484 (90.2%) of 536 subjects accepted to participate (Figure 1) (Table1). We did not observe significant differences between participants and no-participants, except for MGEs assistance, which was lower in non-participants (p<0.001).

Table 1. Participants and non-participants characteristics. COVID-19 Borriana cohort second SARS-CoV-2 serologic survey, October 2020.

	Participant	No-participants	1
Variables	N=484 (%)	N=52 (%)	p-value
Female	301(62.2)	32(61.5)	1.000
Age mean ± Standard Deviation	37.2±17.1	33.5±16.7	0.064
0-24 years	143(29.5)	19 (36.5)	
25-44	157(32.4)	17 (32.7)	0.492
45-64	166(34.3)	16 (30.8)	0.405
65 and over	18 (3.7)	0 (0.0)	
Body mass index ¹ Kg/m ²			
<18.0	41 (8.5)	6 (11.5)	
18.0-24.9	210 (43.7)	24 (46.2)	0 747
25.0-29.9	148 (30.8)	13 (25.0)	0.747
≥30.0	85 (17.7)	9 (17.3)	
Occupation I-II ^{2 3}	145 (30.1)	16 (30.7)	1.000
Current smoker ⁴	65 (13.9)	12 (23.1)	0.221
Physical exercise	389 (80.4)	24 (46.2)	0.075
Alcohol beverages ⁵	108 (23.0)	10 (19.2)	0.604
Chronic illness ⁶	166 (34.6)	14 (26.9)	0.285
COVID-19			
Family with COVD-19 case ⁷	303 (62.7)	31 (59.6)	0.554
Probable contact COVID-19 case ⁸	390 (81.8)	39 (75.0)	0.349
Assistance Mass Gathering events ≥2 and	205 (61 0)	16 (20.8)	0.000
over ⁹	295 (61.0)	10 (50.8)	0.000
Hospitalisations	9 (1.9)	3 (5.8)	0.101
PCR positive	26 (5.4)	13 (25.0)	NC^{10}
Asymptomatic	54 (11.2)	10 (19.2)	0.111
Medical consultation	208 (43.0)	25 (48.1)	0.556

 Missing information 3 participants. 2. Occupation groups I-II: professional, managerial and technical occupations. Groups III-VI: skilled, non-manual or manual, partly-skilled, unskilled occupations. 3. Missing information 3 participants. 4. Missing information 16 participants. 5 Missing information 14 participants. 6. Missing information 4 participants. 7. Missing information 1 participant. 8. Missing information 7 participants.9. Excluding participants with COVID-19 symptoms before 6 March or after 31 March 2020.10. No calculable only positive PCR cases recorded.

From 484 participants in the outbreak cohort persistent antibodies were detected in 479 (99%) and re-infection episodes were detected. Only five of the 484 participants (1.0%) had a negative anti-SARS-CoV-2 antibody in the second determination.

A non-adjusted comparison of subjects with a negative and a positive ECLIA is presented in Table 2. Negative ECLIA subjects were younger with a lower BMI and reported lower frequency of probably contacts with COVID-19 cases and MGEs attendance; shorter duration, no medical consultations, and full recovery COVID-19 disease. In contrast, one-third of the ECLIA positive subjects suffered some sequelae, not recovered completely and had a worse health level than before the COVID-19 disease. The ECLIA results did not differ by the O positive, ABO blood group, or vitamin D mean serum concentration.

 Table 2.
 Subjects characteristic distribution by SARS-CoV-2 ECLIA result, and unadjusted relative risk of comparing negative versus persistent antibody (positive ECLIA) results.
 COVID-19 Borriana cohort, second seroepidemiological survey, October 2020.

	Negative	Persistent			
Variables	Antibodies	Antibodies	RR^1	95% CI ²	p-value
	n=5(%)	n=479 (%)			1
Female	4 (80.0)	297 (62.0)	0.41	0.05-3.66	0.426
Age mean ± standard deviation	24.0±16.7	37.4±17.1	0.95	0.90-1.01	0.106
0-24 years	4 (80.0)	139 (29.0)			
25-44	0	157 (32.8)			0.100
45-64	1 (20.0)	165 (34.4)			0.123
65 and over	0	18 (3.8)			
O ABO ³	4(80)	195 (40.8)	5.71	0.64-50.8	0.116
Vitamin D ng/ml	28.2±8.9	29.8±9.2	0.98	0.87-1.09	0.687
Body mass index ⁴ Kg/m ²	21.3±2.7	25.0±5.0	0.84	0.76-0.94	0.002
<18.0	1(20.0)	40 (8.4)			
18.0-24.9	4 (80.0)	206 (43.3)			0.020
25.0-29.9	0	148 (31.1)			0.038
≥30.0	0	85 (17.9)			
Occupation I-II ⁵	1(20.0)	144 (30.3)	0.57	0.07-5.15	0.624
Current smoker ⁶	1(20.0)	64 (13.8)	1.19	0.37-3.82	0.776
Physical exercise	4 (80.0)	285 (59.5)	2.70	0.30-24.08	0.653
Alcohol consumption ⁷	1 (20.0)	107 (23.0)	0.83	0.29-7.44	0.874
Chronic illness ⁸	0	166 (34.9)	0.28	0.00-2.06	0.240
COVID-19 disease					
Family with COVD-1 9 case ⁹	2 (40.0)	301(63.0)	0.39	0.07-2.35	0.308
Probable contact COVID-19 case ¹⁰	2 (40.0)	388(82.2)	0.15	0.03-0.88	0.044
Assistance Mass Gathering Events ≥2	1 (359/)	255 ((0, ()	0.22	0.02.2.10	0.190
and over ¹¹	1 (25%)	255 (60.6)	0.22	0.02-2.10	0.189
Hospitalisations	0	9 (1.9)	7.85	0.00-19.23	1.000
PCR positive	0	26 (5.4)	2.61	0.00-57.60	1.000
Asymptomatic	1 (20.0)	53 (11.1)	1.99	0.23-17.62	0.448
Medical consultation	0	208 (43.4)	0.19	0.00-1.45	0.121
Illness duration	4.3±6.7	10.4±17.5	0.93	0.76-1.13	0.457
Post-COVID-19					
Sequelae	0	159 (33.2)	0.30	0.00-2.23	0.273
Health as before the disease ¹²	5 (100.0)	397 (83.1)	1.36	0.18-∞	0.732
Recover health ¹³	5 (100.0)	390 (81.6)	1.50	0.20-∞	0.800
Exposure post-COVID-19					
Social contact	4 (80.0)	103 (78.5)	1.09	0.12-9.79	0.935
Gathering people	0	25 (5.2)	2.73	0.00-20.04	1.000
Trip out of Borriana ¹⁴	1(20.0)	109 (22.9)	0.85	0.10-7.50	0.880
Restaurant assistance	0	142 (29.6)	0.36	0.00-2.63	0.352
Terrace assistance ¹⁵	3 (60.0)	286 (59.8)	2.19	0.37-13.04	0.400

RR = Relative risk. 2. CI = Interval confidence. 3. Missing information 1 participant. 4 Missing information 3 participants. 5. Missing information 3 participants. 6. Missing information 16 participants. 7. Missing information 14 participants. 8. Missing information 4 participants. 7. 9. Missing information 1 participants. 10. Missing information 7 participants. 11. Excluding participants with COVID-19 symptoms before 6 March or after 31 March 2020.12. Missing information 1 participants. 13. Missing information 1 participants. 14. Missing information 2 participants. 15. Missing information 1 participants.

We show in Table 3 the multivariate Poisson regression adjusted relative risk (aRR) of the association between the measured factors and an ECLIA negative result. The negative anti-SARS-COV-2 antibody group had a lower BMI (aRR=0.83 95% CI 0.72-0.99), less probable contact with COVID-19 cases (aRR=0.13 95% CI 0.03-0.51), and was younger (aRR=0.96 95% CI 0.91-1.00).

Table 3. Adjusted relative risks (aRR) of factors associated with negative anti-SARS-CoV-2 antibody group versus persistent(positive ECLIA result), by Poisson regression. COVID-19 Borriana cohort 2020.

Factors	aRR	95% CI ¹	p-value	Pearson goodness of fit
Body mass index (Kg/m ²) ²	0.87	0.77-0.99	0.037	0.817
Age (years) ³	0.96	0.91-1.00	0.076	0.971
Sex: Female⁴	0.59	0.09-3.96	0.590	0.971
O ABO blood group⁵	5.52	0.61-49.57	0.127	0.971
Occupation I-II ⁶	0.66	0.08-5.45	0.695	0.587
Current smoker ⁷	2.43	0.14-41.87	0.541	0.072
Physical exercise ⁷	3.05	0.36-25.71	0.305	0.999
Alcohol consumption ⁷	1.07	0.10-11.77	0.954	0.398
Chronic illness ⁸	0.42	0.00-3.35	0.456	NC ⁹
COVID-19 disease				
Family with COVD-19 case ¹⁰	0.49	0.08-2.85	0.428	0.540
Probable contact COVID-19 case ¹¹	0.18	0.04-0.82	0.027	1.000
Assistance Mass Gathering Events ≥2 and over ¹²¹³	0.20	0.02-1.78	0.150	0.458
Medical consultation ¹⁴	0.26	0.00-2.05	0.227	NC ⁹
Illness duration ¹⁵	0.94	0.77-1.15	0.570	1.000

1. CI = Confidence interval. 2. Adjusted for age sex ABO, occupation current smoker physical exercise alcohol consumption. 3. Adjusted for sex ABO. 4. Adjusted for ABO age. 5. Adjusted for sex age. 6. Adjusted for age ABO sex. 7. Adjusted for age sex ABO occupation. 8. Adjusted for age sex ABO current smoker physical exercise alcohol beverage. 9. NC = No computable 10. Adjusted for age sex ABO chronic illness current smoker physical exercise alcohol consumption assistance mass gathering events \geq 2 and over.11. Adjusted for age, sex ABO occupation current smoker physical exercise alcohol consumption assistance mass gathering events \geq 2 and over. 12 Excluding participants with symptoms before 6 March or after 31 March 2020.13. Adjusted for age, sex, ABO, occupation, current smoker physical exercise alcohol consumption assistance mass gathering events \geq 2 and over. 14. Adjusted for age sex, ABO, chronic illness occupation. 15. Adjusted by age sex, body mass index, ABO, chronic illness.

Among the 484 participants, 469 (96.7%) had two anti-SARS-CoV-2 antibody determinations by ECLIA, one in the first study (June 2020) and the second in the current survey. We observed an overall decline in the values of the arbitrary units of the ECLIA test from June 2020, geometric mean 46.41 (95% CI 41.87-51.44) to October 2020, geometric mean 26.55 (95% CI 23.30-30.25) (p<0.001). However, ninety-six participants (20.5%) had an increase of antibodies, whereas 373 participants (79.5%) had a decline. The comparison of the two groups is in Table 4. Participants with an increase in antibodies were older, not current smokers, had a higher BMI, a higher frequency of probable contact with COVID-19 cases, more medical consultation, and a longer duration of the illness. Vitamin D status and O ABO blood levels were similar in both groups.

Table 4. Comparison between participants with increase and decline anti-SARS-CoV-2 antibodies, considering the first and seco	nd
SARS-CoV-2 serologic surveys. COVID-19 Borriana cohort 2020.	

	Increase	Decline	
Variables	SARS-CoV-2	SARS-CoV-2	Pvalue
	N=96 (%)	n=373 (%)	
Female	60(62.5)	236 (63.5)	0.906
Age mean ± standard deviation	45.3±18.0	34.9±16.3	0.001
0-24 years	15 (15.6)	126 (33.8)	
25-44	24 (25.0)	131 (35.1)	0.000
45-64	47 (49.0)	108 (29.0)	0.000
65 and over	10 (10.4)	8 (2.1)	
O ABO ¹	45 (46.9)	149 (40.1)	0.246
Vitamin D ng/ml	29.4±9.0	30.0±9.3	0.621
Body mass index (Kg/m²)	26.9±5.0	24.5±4.9	0.001
<18.0	6 (6.3)	38 (10.2)	
18.0-24.9	27 (28.1)	176 (47.2)	0.001
25.0-29.9	35 (36.5)	107 (28.7)	0.001
≥30.0	38 (40.0)	55 (14.7)	
Occupation I-II ²	29 (30.2)	107 (28.9)	0.802
Current smoker ³	7 (7.5)	56 (15.6)	0.003
Physical exercise	57 (59.4)	222 (59.5)	1.000
Alcohol beverages ⁴	24 (25.8)	82 (22.7)	0.583
Chronic illness ⁵	41 (43.2)	123 (45.6)	0.092
COVID-19			
Family with COVD-19 case ⁶	63 (65.6)	230 (61.8)	0.555
Probable contact COVID-19 case ⁷	83 (89.2)	294 (79.7)	0.036
Assistance Mass Gathering Events ≥2 and over ⁸	61 (63.5)	227 (60.9)	0.724
Hospitalisations	3 (3.1)	6 (1.6)	0.398
PCR positive	8 (8.3)	17 (4.6)	0.198
Asymptomatic	7 (7.3)	46 (12.3)	0.206
Medical consultation	56 (58.3)	144 (38.6)	0.001
Illness duration	12.0±15.0	10±18.2	0.034

 Missing information 1 participant. 2. Missing information 3 participants. 4. Missing information 11 participants. 5. Missing information 3 participants. 6 Missing information 1 participants. 7. Missing information 7 participants. 8. Excluding participants with COVID-19 symptoms before 6 March or after 31 March 2020.

After an adjusted Poisson regression analysis (Table 5), high BMI, older age, a medical consultation and more contact with COVID-19 cases were associated with an increase of anti-SARS-CoV-2 antibodies, being a current smoker was associated with a decline in anti-SARS-CoV-2 antibodies (Figure 2).

Table 5. Adjusted relative risks (aRR) of factors with associated with an increase of anti-SARS-CoV-2 antibodies, considering the first and second SARS-CoV-2 serologic surveys, by Poisson regression. COVID-19 Borriana cohort 2020.

Variables	Variables	n value	Pearson	
d		95% CI ¹	p-value	goodness of fit
Body mass index (Kg/m ²) ²	1.06	1.02-1.10	0.001	0.996
Age (years) ³	1.03	1.02-1.04	0.000	0.991
Current smoker ⁴	0.48	0.24-0.96	0.037	0.997
Probable contact COVID-19 case ⁵	2.04	1.12-3.74	0.022	0.994
Medical consultation ⁶	1.62	1.12-2.34	0.010	0.991
Illness duration ⁷	0.99	0.99-1.01	0.543	0.966

1. CI = Confidence interval. 2. Adjusted for age sex ABO, occupation current smoker physical exercise alcohol consumption. 3.

Adjusted for sex ABO. 4. Adjusted for age sex ABO occupation. 4. Adjusted for age, sex ABO occupation current smoker physical

exercise alcohol consumption assistance mass gathering events \geq 2 and over. 6. Adjusted for age sex, ABO, chronic illness occupation. 7. Adjusted by age sex, body mass index, ABO, chronic illness.



Figure 2. Absolute difference in ECLIA arbitrary units between second and first serosuvey. ECLIA titers evolution COVID-19 Borriana cohort 2020.

4. Discussion

Our results show that 99 % (479/484) of COVID19 patients in the cohort maintained anti-SARS-CoV-2 antibodies in the first six months after the COVID-19 episode. We did not find evidence of any SARS-CoV-2 re-infection episode in this group in the period between the first and second serosurvey.

The factors associated with a negative result in the second serosurvey were a lower BMI, lower frequency of probable contact with COVID-19 cases, a younger age, and a milder COVID19 episode when compared to the group with antibody persistence. Besides, we observed a significant overall decline in anti-SARS-CoV-2 antibodies. However, a fifth of participants showed up an increase in anti-SARS-CoV-2 antibodies, which was probably related to the infection course or the frequency of contacts with other cases.

The persistence of anti-SARS-CoV-2 antibodies after the infection, measured by different serologic techniques, including ECLIA [17-18] or other methods [19-22], has been reported to last between 4 and 8 months. However, significant differences in antibodies persistence associated with the detection techniques have been found, and total antibodies against S1and the S1 receptor-binding domain (RBD) were those detected for longer [23].

Several studies have described a similar decline of neutralising SARS-CoV-2 antibodies. This finding supports the need for follow-up studies to gather evidence on the duration of the antibody response and the protection against re-infections [24-25]. Alt-

hough we did not observe re-infection episodes in our cohort, others have reported re-infection episodes. A 96% protective effectiveness in prior infected patients was estimated in care homes outbreaks [26-27].

In the United States, and from data generated from commercial laboratory analysis of SARS-COV-2 antibody tests of 3.2 million patients, Harvey and co-authors found 0.30% re-infections [28]. In the United Kingdom, two cohort studies of health care workers comparing positive anti-SARS-COV-2 antibody group versus negative group during six months of follow-up (29), have found re-infection rate ratios of 0.11 and 0.16, respectively [30]. In a vessel outbreak, three crew members with previous positive anti-SARS-COV-2 antibodies had not suffered infection versus 104 of 122 (85.2%) crew members with negative antibodies, who suffered a SARS-CoV-2 infection (p=0.002) [31].

A more severe COVID-19 disease is usually associated with higher anti-SARS-CoV-19 antibodies [32-34], and antibodies correlate with duration of the infection, older age and hospitalisation [35-36]. Still, in other studies, age and symptoms were not associated with antibody levels [37], suggesting high variability. Also, high levels of anti-SARS-CoV IgG and neutralising antibodies were observed in COVID-19 patients with high BMI and patients with metabolic syndrome [33, 38-40]. These results are in line with our finding, where low BMI was associated with negative antibodies. However, in obese COVID-19 patients, a reverse association has been observed between high BMI and lower SARS-CoV-2 antibodies [41]. In addition, the increase of SARS-CoV-2 antibodies in the second survey was associated with age older, higher BMI, more severe disease (42) and low current smoke. A hypothesis was suggested that a tobacco mosaic RNA virus, presented in the respiratory tract of smokers, could have some protection against SARS-COV-2 virus (43).

Factors indicating less exposure to COVID-19 cases in the group with negative antibodies suggest that the intensity, type, and duration of exposure could play a role in the development of the disease and its course [37, 44-45]. Then, high viral load exposure with high duration may be decisive for the development and course of COVID-19 infection [46].

The O ABO blood group was not associated with negative antibody groups, in line with the finding of Wendel and co-author (38). Still, the small sample of the negative antibody group prevents a definitive conclusion. Also, Vitamin D status was not related to negative antibody group, but the measure as prevalence exclude a potential role [47]. On the other hand, negative antibodies were not associated with sex in contrast with Markmann and co-authors article [33], where male had more neutralising antibody levels than female.

Serologic surveys of SARS-CoV-2 infection are a useful approach in the control and prevention of COVID-19 disease, including the surveillance of the disease, to characterise the effectiveness of vaccinations against the disease, the antibody response relation to disease course, and the factors associated with the duration of immune response [3].

Many serologic tests of SARS-CoV-2 measure IgA, IgG, and IgM, with a high heterogeneity in sensitivity and specificity [48]. The ECLIA technique has a good accuracy [3, 49-50] and has been recommended for population screening [51]. Anti-SARS-CoV-2 antibodies obtained by ECLIA positively correlate with neutralising antibodies, but their sensitivity should be improved [52-53].

Our study has some strengths; first, a prospective cohort design; second, the elevated participation rate; third, the use of a serologic test with reasonable sensitivity and specificity; four, the use of multivariate analysis to control potential confounding; fifth, a follow-up of up to 6 months; and, finally, a population-based approach.

As limitations, the quantitative ECLIA anti-SARS-CoV-2 antibodies test that we used has a significant but feeble correlation (Pearson's correlation coefficient of 0.37 p<0.001 and poor linear relationship with the enzyme-linked immunosorbent assay (ELISA AESKULISA^R) [54]. The ECLIA test does not measure the response to S antigen of the SARS-CoV-2, but detects antibodies against the nucleocapsid N. Finally, the small number of participants with negative anti-SARS-CoV-2 antibodies decreases the power to

detect associated factors; the younger age and, in general, milder COVID-19 disease of the cohort does not permit a generalisation of results, and we cannot discard the existence of unknown factors that we did not consider in this new disease.

In subsequent studies, a quantitative determination of neutralising antibodies against RDB antigen [55-56] could be helpful to study the immunity against the SARS-CoV-2 virus considering the importance to define correlate levels of protective immunity [57-58] and more precise estimation of potential factors associated with the clinical course [59-60] As we can assume the future existence of vaccinated participants against SARS-CoV-2 in the cohort, the follow-up of the cohort would provide information on the response after vaccination against SARS-CoV-2 in patients who have suffered the infection [61] and inform the need of additional vaccine doses.

5. Conclusion

In the first six months after a COVID-19 infection, a high proportion of participants maintained detectable anti-SARS-CoV-2 antibodies, and we did not observe new COVID-19 episodes in the follow-up period.

6. Patens

Author contributions: Conceptualization SDM, AAP, MRPS, LGL, DJS, LAE, UCA, SFR, MSU.

Methodology AAP,SDM, JPB, PVU, MLP, ADRG, SFR, MSU, GFA, LAE, GBM, UCA, CDB, MMB; software AAP, MRPS, JPB,MCLD, ; validation JPB, CNR MRPS ; formal analysis MRPS, AAP, JPB, SDM ; investigation SDM, MRPS, PVU, MLP, ADRG, SFR, GFA, MSU, LAE, GBM, BCF, UCA, CDB, MFC, LGL, DJS, MCLD, MDLLV, MMB, CNR, RRP, SVL ; resources SDM, GFA, GBM, BCF, MFC, LGL, MDLLV, MLP, CNR, RRP, SVL ; data curation AAP, JPB, PVU, MLP, ADRG, MRPS ; writing original draft preparation AAP, JPB, MRPS, MCLD, SDM, SFR ; writing-review and editing JPB, AAP, SDM, UCA, MFC, MCDL ; visualization JPB, DJS, LAE, LGL ; supervisión MRPS, MCLD; project admistration SDM, AAP, MCLD, GFA ; funding acquision SDM, MCLD, AAP . All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, The study of this cohort was a part of the public health surveillance as a prolongation of the COVID-19 outbreak in the Falles festival of Borriana control measures (1) and was exempt from the Ethics Review Board approval's protocol according to the Spanish legislation, including the General Low including the General Law of Health (2), the Law of Cohesion and Quality of the National System of Health (3), and the Law General of Public Health (4). In addition, the cohort was followed to respond to a new disease, the COVID-19 pandemic (5).

1. Domènech-Montoliu S, Pac-Sa MR, Vidal-Utrillas P, Latorre-Poveda M, Del Rio-González A, Ferrando-Rubert S, et al. "Mass gathering events and COVID-19 transmission in Borriana (Spain): A retrospective cohort study". PLoS One. 2021 Aug 26;16(8):e0256747. doi: 10.1371/journal.pone.0256747.

2. Ley 14/1986, de 25 de abril, General de Sanidad (Law General of Health): <u>https://www.boe.es/eli/es/l/1986/04/25/14/con</u> Access: 2 August 2021

3. Ley 16/2003, de 28 de mayo, de cohesión y calidad del Sistema Nacional de Salud. (Law of Cohesion and Quality of the National System of Health). <u>https://www.boe.es/eli/es/l/2003/05/28/16</u> Access: 2 August 2021.

33/2011, 4. Lev de 4 de octubre, General de Salud Pública (Law General of Public Health). https://www.boe.es/eli/es/l/2011/10/04/33/con Access: 2 August 2021.

5. Ministerio de Sanidad Acordado en Consejo Interterritorial del Sistema Nacional de Salud el 16 de julio de 2020. Plan de respuesta temprana en un escenario de control de la pandemia por el COVID-19.pdf (Ministry of Health Agreed in the Interterritorial Council of the National Health System on July 16, 2020". Early response plan in a COVID-19 pandemic control scenario. pdf).

Informed Consent Statement: Informated consent was obtained from all subject involved in the study.

Data Availability Statement: Data of the this study can be consulted if the authors are requested. Dataset: borrianacohort.dta.

Acknowledgments: We thank the participants of the cohort and the Borriana's *Falles* organisation for the support that made it possible to perform this study. In addition, we appreciate the assistance and support of Roser Blasco-Gari, Helena Buj-Jorda, Israel Borras-Acosta, Lucia Castell-Agusti, Mercedes De Francia-Valero, Maria Domènech-Molinos, Marc Garcia, Maria Gil-Fortuño, Elena Grañana-Toran, Noelia Hernández-Perez, Laura Lopez-Diago, Salvador Martinez-Parra, Sara Moner-Marin, Silvia Pesudo-Calatayud, Lara Sabater-Hernández, Maria Luisa Salve-Martinez, Irene Suarez-Linares, Juan José Ventura-Buchardo, and Alberto Yagüe-Muñoz to carry out the study.

Conflits of Interest: The authors declare no conflict of interest.

References

1. Johns Hopkins University. COVID-19 Data in Motion: 12 May, 2021. https://coronavirus.jhu.edu/

2. Clapham H, Hay J, Routledge I, Takahashi S, Choisy M, Cummings D, Grenfell B, Metcalf CJE, Mina M, Barraquer IR, Salje H, Tam CC. Seroepidemiologic study designs for determining SARS-COV-2 transmission and immunity. Emerg Infect Dis. 2020 ;26:1978-1986. doi: 10.3201/eid2609.201840.

3.Coste AT, Jaton K, Papadimitriou-Olivgeris M, Greub G, Croxatto A. Comparison of SARS-CoV-2 serological tests with different antigen targets. J Clin Virol. 2021 ;134:104690. doi: 10.1016/j.jcv.2020.104690.

4. Lee CY, Lin RTP, Renia L, Ng LFP. Serological approaches for COVID-19: Epidemiologic perspective on surveillance and control. Front Immunol. 2020;11:879. doi: 10.3389/fimmu.2020.00879.

5. Liu G, Rusling JF. COVID-19 antibody tests and their limitations. ACS Sens. 2021;6:593-612. doi: 10.1021/acssensors.0c02621.

6.Egger M, Bundschuh C, Wiesinger K, Gabriel C, Clodi M, Mueller T, Dieplinger B. Comparison of the Elecsys® Anti-SARS-CoV-2 immunoassay with the EDI[™] enzyme linked immunosorbent assays for the detection of SARS-CoV-2 antibodies in human plasma. Clin Chim Acta. 2020;509:18-21. doi: 10.1016/j.cca.2020.05.049.

7.Dellière S, Salmona M, Minier M, Gabassi A, Alanio A, Le Goff J, Delaugerre C, Chaix ML; Saint-Louis CORE (COvid REsearch) group. Evaluation of the COVID-19 IgG/IgM rapid test from Orient Gene Biotech. J Clin Microbiol. 2020;58:e01233-20. doi: 10.1128/JCM.01233-20.

8.Yip CC, Sridhar S, Cheng AK, Leung KH, Choi GK, Chen JH, Poon RW, Chan KH, Wu AK, Chan HS, Chau SK, Chung TW, To KK, Tsang OT, Hung IF, Cheng VC, Yuen KY, Chan JF. Evaluation of the commercially available LightMix® Modular E-gene kit using clinical and proficiency testing specimens for SARS-CoV-2 detection. J Clin Virol. 2020;129:104476. doi: 10.1016/j.jcv.2020.104476.

9. Fan Q, Zhang W, Li B, Li DJ, Zhang J, Zhao F. Association between ABO Blood Group System and COVID-19 susceptibility in Wuhan. Front Cell Infect Microbiol. 2020;10:404. doi: 10.3389/fcimb.2020.00404.

10.Wu Y, Feng Z, Li P, Yu Q. Relationship between ABO blood group distribution and clinical characteristics in patients with COVID-19. Clin Chim Acta. 2020 ;509:220-223. doi: 10.1016/j.cca.2020.06.026.

11.Merzon E, Tworowski D, Gorohovski A, Vinker S, Golan Cohen A, Green I, Frenkel-Morgenstern M. Low plasma 25(OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study. FEBS J. 2020 ;287:3693-3702. doi: 10.1111/febs.15495.

12. Kaufman HW, Niles JK, Kroll MH, Bi C, Holick MF. SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. PLoS One. 2020;15:e0239252. doi: 10.1371/journal.pone.0239252

13.Lapierre Y, Rigal D, Adam J, Josef D, Meyer F, Greber S, Drot C. The gel test: a new way to detect red cell antigen-antibody reactions. Transfusion. 1990;30:109-13. doi: 10.1046/j.1537-2995.1990.30290162894.x.

14.Asif M, Groboske SE, Leung EKY, Yeo KJ, van Wijk XMR. Evaluation of a new generation automated assay for 25-hydroxy vitamin D based on competitive protein binding. J Appl Lab Med. 2019;4:247-253. . doi: 10.1373/jalm.2018.028415.

15. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. Epidemiology. 1999;10:37-48.

16. Textor J, van der Zander B, Gilthorpe MS, Liskiewicz M, Ellison GT. Robust causal inference using directed acyclic graphs: the R package 'dagitty'. Int J Epidemiol. 2016;45:1887-1894. doi: 10.1093/ije/dyw341.

17. Fejt V, Krátká Z, Zelená H, Fürst T. Age is not a disease: Evolution of protective antibodies against SARS-CoV-2 in seniors from the Břevnice nursing home. Cas Lek Cesk. 2020 Winter;159(7-8):303-311.

18. Choe PG, Kim KH, Kang CK, Suh HJ, Kang E, Lee SY, Kim NJ, Yi J, Park WB, Oh MD. Antibody responses 8 months after asymptomatic or mild SARS-CoV-2 infection. Emerg Infect Dis. 2021;27:928-931. doi: 10.3201/eid2703.204543.

19. Wajnberg A, Amanat F, Firpo A, Altman DR, Bailey MJ, Mansour M, McMahon M, Meade P, Mendu DR, Muellers K, Stadlbauer D, Stone K, Strohmeier S, Simon V, Aberg J, Reich DL, Krammer F, Cordon-Cardo C. Robust neutralizing antibodies to SARS-CoV-2 infection persist for months. Science. 2020 ;370:1227-1230. doi: 10.1126/science.abd7728.

20.Gudbjartsson DF, Norddahl GL, Melsted P, Gunnarsdottir K, Holm H, Eythorsson E, Arnthorsson AO, Helgason D, Bjarnadottir K, Ingvarsson RF, Thorsteinsdottir B, Kristjansdottir S, Birgisdottir K, Kristinsdottir AM, Sigurdsson MI, Arnadottir GA, Ivarsdottir EV, Andresdottir M, Jonsson F, Agustsdottir AB, Berglund J, Eiriksdottir B, Fridriksdottir R, Gardarsdottir EE, Gottfredsson M, Gretarsdottir OS, Gudmundsdottir S, Gudmundsson KR, Gunnarsdottir TR, Gylfason A, Helgason A, Jensson BO, Jonasdottir A, Jonsson H, Kristjansson T, Kristinsson KG, Magnusdottir DN, Magnusson OT, Olafsdottir LB, Rognvaldsson S, le Roux L, Sigmundsdottir G, Sigurdsson A, Sveinbjornsson G, Sveinsdottir KE, Sveinsdottir M, Thorarensen EA, Thorbjornsson B, Thordardottir M, Saemundsdottir J, Kristjansson SH, Josefsdottir KS, Masson G, Georgsson G, Kristjansson M, Moller A, Palsson R, Gudnason T, Thorsteinsdottir U, Jonsdottir I, Sulem P, Stefansson K. Humoral immune response to SARS-CoV-2 in Iceland. N Engl J Med. 2020 ;383:1724-1734. doi: 10.1056/NEJMoa2026116.

21.Ripperger TJ, Uhrlaub JL, Watanabe M, Wong R, Castaneda Y, Pizzato HA, Thompson MR, Bradshaw C, Weinkauf CC, Bime C, Erickson HL, Knox K, Bixby B, Parthasarathy S, Chaudhary S, Natt B, Cristan E, El Aini T, Rischard F, Campion J, Chopra M, Insel M, Sam A, Knepler JL, Capaldi AP, Spier CM, Dake MD, Edwards T, Kaplan ME, Scott SJ, Hypes C, Mosier J, Harris DT, LaFleur BJ, Sprissler R, Nikolich-Žugich J, Bhattacharya D. Orthogonal SARS-CoV-2 serological assays enable surveillance of low-prevalence communities and reveal durable humoral immunity. Immunity. 2020;53:925-933.e4. doi: 10.1016/j.immuni.2020.10.004.

22. Dan JM, Mateus J, Kato Y, Hastie KM, Yu ED, Faliti CE, Grifoni A, Ramirez SI, Haupt S, Frazier A, Nakao C, Rayaprolu V, Rawlings SA, Peters B, Krammer F, Simon V, Saphire EO, Smith DM, Weiskopf D, Sette A, Crotty S. Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. Science. 2021;371:eabf4063. doi: 10.1126/science.abf4063.

23. Favresse J, Eucher C, Elsen M, Gillot C, Van Eeckhoudt S, Dogné JM, Douxfils J. Persistence of anti-SARS-CoV-2 antibodies depends on the analytical kit: A report for up to 10 months after infection. Microorganisms. 2021;9:556. doi: 10.3390/microorganisms9030556.

24.Nguyen-Contant P, Embong AK, Kanagaiah P, Chaves FA, Yang H, Branche AR, Topham DJ, Sangster MY. S Protein-Reactive IgG and memory B cell production after human SARS-CoV-2 infection includes broad reactivity to the S2 subunit. mBio. 2020 ;11:e01991-20. doi: 10.1128/mBio.01991-20.

25. Seow J, Graham C, Merrick B, Acors S, Pickering S, Steel KJA, Hemmings O, O'Byrne A, Kouphou N, Galao RP, Betancor G, Wilson HD, Signell AW, Winstone H, Kerridge C, Huettner I, Jimenez-Guardeño JM, Lista MJ, Temperton N, Snell LB, Bisnauthsing K, Moore A, Green A, Martinez L, Stokes B, Honey J, Izquierdo-Barras A, Arbane G, Patel A, Tan MKI, O'Connell L, O'Hara G, MacMahon E, Douthwaite S, Nebbia G, Batra R, Martinez-Nunez R, Shankar-Hari M, Edgeworth JD, Neil SJD, Malim MH, Doores KJ. Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. Nat Microbiol. 2020;5:1598-1607. doi: 10.1038/s41564-020-00813-8.

26. West J, Everden S, Nikitas N. A case of COVID-19 re-infection in the UK. Clin Med (Lond). 2021;21:e52-e53. doi: 10.7861/clinmed.2020-0912.

27. Jeffery-Smith A, Iyanger N, Williams SV, Chow JY, Aiano F, Hoschler K, Lackenby A, Ellis J, Platt S, Miah S, Brown K, Amirthalingam G, Patel M, Ramsay ME, Gopal R, Charlett A, Ladhani SN, Zambon M. Antibodies to SARS-CoV-2 protect against re-infection during outbreaks in care homes, September and October 2020. Euro Surveill. 2021;26:2100092. doi: 10.2807/1560-7917.ES.2021.26.5.2100092.

28. Harvey RA, Rassen JA, Kabelac CA, Turenne W, Leonard S, Klesh R, Meyer WA 3rd, Kaufman HW, Anderson S, Cohen O, Petkov VI, Cronin KA, Van Dyke AL, Lowy DR, Sharpless NE, Penberthy LT. Association of SARS-CoV-2 seropositive antibody test with risk of future infection. JAMA Intern Med. 2021 ;181:672-679. doi: 10.1001/jamainternmed.2021.0366.

29. Lumley SF, O'Donnell D, Stoesser NE, Matthews PC, Howarth A, Hatch SB, Marsden BD, Cox S, James T, Warren F, Peck LJ, Ritter TG, de Toledo Z, Warren L, Axten D, Cornall RJ, Jones EY, Stuart DI, Screaton G, Ebner D, Hoosdally S, Chand M, Crook DW, O'Donnell AM, Conlon CP, Pouwels KB, Walker AS, Peto TEA, Hopkins S, Walker TM, Jeffery K, Eyre DW; Oxford University Hospitals Staff Testing Group. Antibody status and incidence of SARS-CoV-2 infection in health care workers. N Engl J Med. 2021;384:533-540. doi: 10.1056/NEJMoa2034545.

30. Hall VJ, Foulkes S, Charlett A, Atti A, Monk EJM, Simmons R, Wellington E, Cole MJ, Saei A, Oguti B, Munro K, Wallace S, Kirwan PD, Shrotri M, Vusirikala A, Rokadiya S, Kall M, Zambon M, Ramsay M, Brooks T, Brown CS, Chand MA, Hopkins S; SIREN Study Group. SARS-CoV-2 infection rates of antibody-positive compared with antibody-negative health-care workers in England: a large, multicentre, prospective cohort study (SIREN). Lancet. 2021;397:1459-1469. doi: 10.1016/S0140-6736(21)00675-9.

31. Addetia A, Crawford KHD, Dingens A, Zhu H, Roychoudhury P, Huang ML, Jerome KR, Bloom JD, Greninger AL. Neutralizing antibodies correlate with protection from SARS-CoV-2 in humans during a fishery vessel outbreak with a high attack rate. J Clin Microbiol. 2020;58:e02107-20. doi: 10.1128/JCM.02107-20.

32. Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, Wang X, Yuan J, Li T, Li J, Qian S, Hong C, Wang F, Liu Y, Wang Z, He Q, Li Z, He B, Zhang T, Fu Y, Ge S, Liu L, Zhang J, Xia N, Zhang Z. Antibody responses to SARS-CoV-2 in patients with novel coronavirus disease 2019. Clin Infect Dis. 2020;71:2027-2034. doi: 10.1093/cid/ciaa344..

33. Markmann AJ, Giallourou N, Bhowmik DR, Hou YJ, Lerner A, Martinez DR, Premkumar L, Root H, van Duin D, Napravnik S, Graham SD, Guerra Q, Raut R, Petropoulos CJ, Wrin T, Cornaby C, Schmitz J, Kuruc J, Weiss S, Park Y, Baric R, de Silva AM, Mar-

golis DM, Bartelt LA. Sex disparities and neutralizing antibody durability to SARS-CoV-2 infection in convalescent individuals. medRxiv [Preprint]. 2021 Feb 3:2021.02.01.21250493. doi: 10.1101/2021.02.01.21250493.

34. Zhang X, Lu S, Li H, Wang Y, Lu Z, Liu Z, Lai Q, Ji Y, Huang X, Li Y, Sun J, Wu Y, Xu X, Hou J. Viral and antibody kinetics of COVID-19 patients with different disease severities in acute and convalescent phases: A 6-month follow-up study. Virol Sin. 2020;35:820-829. doi: 10.1007/s12250-020-00329-9.

35. Benner SE, Patel EU, Laeyendecker O, Pekosz A, Littlefield K, Eby Y, Fernandez RE, Miller J, Kirby CS, Keruly M, Klock E, Baker OR, Schmidt HA, Shrestha R, Burgess I, Bonny TS, Clarke W, Caturegli P, Sullivan D, Shoham S, Quinn TC, Bloch EM, Casadevall A, Tobian AAR, Redd AD. SARS-CoV-2 antibody avidity responses in COVID-19 patients and convalescent plasma donors. J Infect Dis. 2020;222:1974-1984. doi: 10.1093/infdis/jiaa581.

36. Shields AM, Faustini SE, Perez-Toledo M, Jossi S, Allen JD, Al-Taei S, Backhouse C, Dunbar L, Ebanks D, Emmanuel B, Faniyi AA, Garvey MI, Grinbergs A, McGinnell G, O'Neill J, Watanabe Y, Crispin M, Wraith DC, Cunningham AF, Drayson MT, Richter AG. Serological responses to SARS-CoV-2 following non-hospitalised infection: clinical and ethnodemographic features associated with the magnitude of the antibody response. medRxiv [Preprint]. 2020 Nov 16:2020.11.12.20230763. doi: 10.1101/2020.11.12.20230763.

37.Ladhani SN, Jeffery-Smith A, Patel M, Janarthanan R, Fok J, Crawley-Boevey E, Vusirikala A, Fernandez Ruiz De Olano E, Perez MS, Tang S, Dun-Campbell K, Evans EW, Bell A, Patel B, Amin-Chowdhury Z, Aiano F, Paranthaman K, Ma T, Saavedra-Campos M, Ellis J, Chand M, Brown K, Ramsay ME, Hopkins S, Shetty N, Chow JY, Gopal R, Zambon M. High prevalence of SARS-CoV-2 antibodies in care homes affected by COVID-19: Prospective cohort study, England. EClinicalMedicine. 2020;28:100597. doi: 10.1016/j.eclinm.2020.100597.

38.Wendel S, Fontão-Wendel R, Fachini R, Candelaria G, Scuracchio P, Achkar R, Brito M, Reis LF, Camargo A, Amano M, Machado R, Araujo D, Soares C, Durigon E. A longitudinal study of convalescent plasma (CCP) donors and correlation of ABO group, initial neutralizing antibodies (nAb), and body mass index (BMI) with nAb and anti-nucleocapsid (NP) SARS-CoV-2 antibody kinetics: Proposals for better quality of CCP collections. Transfusion. 2021;61:1447-1460. doi: 10.1111/trf.16323.

39. Racine-Brzostek SE, Yang HS, Jack GA, Chen Z, Chadburn A, Ketas TJ, Francomano E, Klasse PJ, Moore JP, McDonough KA, Girardin RC, Dupuis AP, Payne AF, Ma LX, Sweeney J, Zhong E, Yee J, Cushing MM, Zhao Z. Postconvalescent SARS-CoV-2 IgG and neutralizing antibodies are elevated in individuals with poor metabolic health. J Clin Endocrinol Metab. 2021;106:e2025-e2034. doi: 10.1210/clinem/dgab004.

40. Gerhards C, Thiaucourt M, Kittel M, Becker C, Ast V, Hetjens M, Neumaier M, Haselmann V. Longitudinal assessment of anti-SARS-CoV-2 antibody dynamics and clinical features following convalescent from COVID-19 infection. Int J Infect Dis. 2021 Apr 28:S1201-9712(21)00392-1. doi: 10.1016/j.ijid.2021.04.080.

41. Frasca D, Reidy L, Cray C, Diaz A, Romero M, Kahl K, Blomberg BB. Influence of obesity on serum levels of SARS-CoV-2-specific antibodies in COVID-19 patients. PLoS One. 2021;16:e0245424. doi: 10.1371/journal.pone.0245424.

42.De Giorgi V, West KA, Henning AN, Chen L, Holbrook MR, Gross R, Liang J, Postnikova E, Trenbeath J, Pogue S, Scinto T, Alter HJ, Cantilena CC. Anti-SARS-CoV-2 Serology persistence over time in COVID-19 Convalescent Plasma Donors. medRxiv [Pre-print]. 2021 Mar 10:2021.03.08.21253093. doi: 10.1101/2021.03.08.21253093.

43. de Bernardis E, Busà L. A putative role for the tobacco mosaic virus in smokers' resistance to COVID-19. Med Hypotheses. 2020 Oct;143:110153. doi: 10.1016/j.mehy.2020.110153.

44. Matricardi PM, Dal Negro RW, Nisini R. The first, holistic immunological model of COVID-19: Implications for prevention, diagnosis, and public health measures. Pediatr Allergy Immunol. 2020;31:454-470. doi: 10.1111/pai.13271.

45.Iwuji K, Islam E, Berdine G, Nugent K, Test V, Tijerina A. Prevalence of coronavirus antibody among first responders in Lubbock, Texas. J Prim Care Community Health. 2020;11:2150132720971390. doi: 10.1177/2150132720971390.

46. Calisti R. SARS-CoV-2: exposure to high external doses as determinants of higher viral loads and of increased risk for COVID-19. A systematic review of the literature. Epidemiol Prev. 2020;44(5-6 Suppl 2):152-159. doi: 10.19191/EP20.5-6.S2.114.

47. Barassi A, Pezzilli R, Mondoni M, Rinaldo RF, DavÌ M, Cozzolino M, Melzi D'Eril GV, Centanni S. Vitamin D in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) patients with non-invasive ventilation support. Panminerva Med. 2021 Jan 25. doi: 10.23736/S0031-0808.21.04277-4.

48.Deeks JJ, Dinnes J, Takwoingi Y, Davenport C, Spijker R, Taylor-Phillips S, Adriano A, Beese S, Dretzke J, Ferrante di Ruffano L, Harris IM, Price MJ, Dittrich S, Emperador D, Hooft L, Leeflang MM, Van den Bruel A; Cochrane COVID-19 Diagnostic Test Accuracy Group. Antibody tests for identification of current and past infection with SARS-CoV-2. Cochrane Database Syst Rev. 2020;6:CD013652. doi: 10.1002/14651858.CD013652.

49.Afzal N, Tariq N, Raza S, Shakeel D. Diagnostic accuracy of electro-chemiluminescence immunoassay anti-SARS-CoV-2 sero-logical test. Cureus. 2021;13:e12588. doi: 10.7759/cureus.12588.

50. Kittel M, Findeisen P, Muth MC, Thiaucourt M, Gerhards C, Neumaier M, Haselmann V. Specificity testing by point prevalence as a simple assessment strategy using the Roche Elecsys® anti-SARS-CoV-2 immunoassay. Int J Infect Dis. 202;105:632-638. doi: 10.1016/j.ijid.2021.02.024.

51. Speletas M, Kyritsi MA, Vontas A, Theodoridou A, Chrysanthidis T, Hatzianastasiou S, Petinaki E, Hadjichristodoulou C. Evaluation of two chemiluminescent and three ELISA immunoassays for the detection of SARS-CoV-2 IgG antibodies: Implications for disease diagnosis and patients' management. Front Immunol. 2020;11:609242. doi: 10.3389/fimmu.2020.609242.

52. Kohmer N, Westhaus S, Rühl C, Ciesek S, Rabenau HF. Brief clinical evaluation of six high-throughput SARS-CoV-2 IgG antibody assays. J Clin Virol. 2020;129:104480. doi: 10.1016/j.jcv.2020.104480.

53.Müller L, Ostermann PN, Walker A, Wienemann T, Mertens A, Adams O, Andree M, Hauka S, Lübke N, Keitel V, Drexler I, Di Cristanziano V, Hermsen DF, Kaiser R, Boege F, Klein F, Schaal H, Timm J, Senff T. Sensitivity of anti-SARS-CoV-2 serological assays in a high-prevalence setting. Eur J Clin Microbiol Infect Dis. 2021;40:1063-1071. doi: 10.1007/s10096-021-04169-7.

54.Khawaja S, Asif M, Naz Mukry S, Sultan Shamsi T. Possible correlation of electrochemiluminescence based numerical cut off index value with concentration of anti-SARS-CoV-2 antibody: Is it worth reporting? J Public Health Res. 2021;10:2079. doi: 10.4081/jphr.2021.2079.

55.Peterhoff D, Glück V, Vogel M, Schuster P, Schütz A, Neubert P, Albert V, Frisch S, Kiessling M, Pervan P, Werner M, Ritter N, Babl L, Deichner M, Hanses F, Lubnow M, Müller T, Lunz D, Hitzenbichler F, Audebert F, Hähnel V, Offner R, Müller M, Schmid S, Burkhardt R, Glück T, Koller M, Niller HH, Graf B, Salzberger B, Wenzel JJ, Jantsch J, Gessner A, Schmidt B, Wagner R. A highly specific and sensitive serological assay detects SARS-CoV-2 antibody levels in COVID-19 patients that correlate with neutralization. Infection. 2021;49:75-82. doi: 10.1007/s15010-020-01503-7.

56.Figueiredo-Campos P, Blankenhaus B, Mota C, Gomes A, Serrano M, Ariotti S, Costa C, Nunes-Cabaço H, Mendes AM, Gaspar P, Pereira-Santos MC, Rodrigues F, Condeço J, Escoval MA, Santos M, Ramirez M, Melo-Cristino J, Simas JP, Vasconcelos E, Afonso Â, Veldhoen M. Seroprevalence of anti-SARS-CoV-2 antibodies in COVID-19 patients and healthy volunteers up to 6 months post disease onset. Eur J Immunol. 2020;50:2025-2040. doi: 10.1002/eji.202048970.

57. Krammer F. Correlates of protection from SARS-CoV-2 infection. Lancet. 2021;397:1421-1423. doi: 10.1016/S0140-6736(21)00782-0.

58. Furukawa K, Arii J, Nishimura M, Tjan LH, Lystia Poetranto A, Ren Z, Aktar S, Huang JR, Sutandhio S, Kurahashi Y, Nishino A, Shigekuni S, Takeda Y, Uto K, Matsui K, Sato I, Inui Y, Endo K, Kosaka Y, Oota T, Saegusa J, Mori Y. Seroepidemiological survey of the antibody for severe acute respiratory syndrome coronavirus 2 with neutralizing activity at hospitals: A cross-sectional study in Hyogo Prefecture, Japan. JMA J. 2021;4:41-49. doi: 10.31662/jmaj.2020-0094.

59. Schaffner A, Risch L, Aeschbacher S, Risch C, Weber MC, Thiel SL, Jüngert K, Pichler M, Grossmann K, Wohlwend N, Lung T, Hillmann D, Bigler S, Bodmer T, Imperiali M, Renz H, Kohler P, Vernazza P, Kahlert CR, Twerenbold R, Paprotny M, Conen D, Risch M. Characterization of a pan-immunoglobulin assay quantifying antibodies directed against the receptor binding domain of the SARS-CoV-2 S1-Subunit of the spike protein: A population-based study. J Clin Med. 2020;9:3989. doi: 10.3390/jcm9123989.

60. Anichini G, Terrosi C, Gandolfo C, Gori Savellini G, Fabrizi S, Miceli GB, Cusi MG. SARS-CoV-2 Antibody Response in Persons with Past Natural Infection. N Engl J Med. 2021:NEJMc2103825. doi: 10.1056/NEJMc2103825.

61. Saadat S, Rikhtegaran Tehrani Z, Logue J, Newman M, Frieman MB, Harris AD, Sajadi MM. Binding and Neutralization Antibody Titers After a Single Vaccine Dose in Health Care Workers Previously Infected With SARS-CoV-2. JAMA. 2021;325:1467-1469. doi: 10.1001/jama.2021.3341.