

Table A-1 MV Model factors

IMV Model factor	18-30 (N = 161)	31+ (N=176)	Significance against Core
Defeat	47 (33 %)	82 (52 %)	.01*
<i>At discharge</i>	9 (6 %)	14 (8 %)	.90
Hopelessness	16 (11 %)	15 (10 %)	.98
<i>At discharge</i>	1 (1 %)	0	.96
Humiliation	73 (51 %)	105 (67 %)	.64
<i>At discharge</i>	12 (8 %)	28 (18 %)	.76
Entrapment	65 (46 %)	97 (62 %)	.02*
<i>At discharge</i>	19 (14 %)	26 (17 %)	.01*
Social problemsolving	31 (22 %)	43 (27 %)	.61
<i>At discharge</i>	27 (19 %)	48 (31 %)	.71
Coping	27 (19 %)	38 (24 %)	.86
<i>At discharge</i>	70 (49 %)	80 (51 %)	.48
Memory biases	54 (38 %)	79 (50 %)	.07
Rumination	110 (77 %)	123 (78 %)	.82
<i>At discharge</i>	45 (32 %)	63 (40 %)	.44
Thwarted Belongingness	101 (71 %)	111 (71 %)	.40
<i>At discharge</i>	40 (28 %)	47 (30 %)	.16
Burdensomeness	67 (47 %)	77 (49 %)	.74
<i>At discharge</i>	0	0	.06
Absence of positive future thinking	66 (47 %)	87 (44 %)	.41
<i>At discharge</i>	11 (8 %)	23 (15 %)	.03*
Unrealistic goals	20 (14 %)	21 (14 %)	.79
<i>At discharge</i>	12 (9 %)	9 (6 %)	.12
Not engaging in new goals	53 (38 %)	73 (47 %)	.02*
<i>At discharge</i>	19 (14 %)	27 (18 %)	.33
Social norms	7 (5 %)	14 (9 %)	.73
Resilience	20 (14 %)	38 (25 %)	.82
<i>At discharge</i>	43 (31 %)	64 (42 %)	.21
Social support	66 (47 %)	82 (53 %)	.12
<i>At discharge</i>	47 (33 %)	69 (45 %)	.02*
Social isolation	12 (9 %)	6 (4 %)	.69
Positive attitudes towards suicide	19 (14 %)	28 (18 %)	.001*
Suicide plan	18 (13 %)	14 (9 %)	.45
<i>At discharge</i>	1 (1 %)	4 (3 %)	.09
Exposure to suicidality	42 (30 %)	60 (39 %)	.91
Impulsivity	62 (44 %)	79 (51 %)	.95
Pain tolerance	12 (9 %)	23 (15 %)	.47
Fearlessness of death	18 (13 %)	30 (20 %)	.07
Imagery of death/suicide	50 (36 %)	53 (34 %)	.73
<i>At discharge</i>	12 (9 %)	16 (10 %)	.71
Past suicide attempt or self-harm	113 (75 %)	121 (74 %)	.35

*p<.05