

Article

The healthcare professionals' perspectives on antimicrobial resistance (AMR) associated factors and their consequences: A cross sectional study in Eastern province of Saudi Arabia.

Mohamed A. Baraka^{1,2*}, Amany Alboghdadly³, Samar Alshawwa^{4*}, Asim Ahmed Elnour⁵, Hassan Alsultan⁶, Taha Als Salman⁶, Hussain Alaithan⁶, Md. Ashraf Islam⁶, Kareem Ahmed El-Fass⁷, Yehia Mohamed^{8,9}, Abdulsalam A. Alasser¹⁰ and Khairi Mustafa Salem¹¹.

- 1 Assistant professor, Clinical Pharmacy program, College of Pharmacy, Al Ain University, Al Ain campus, United Arab Emirates. Mohamed.baraka@au.ac.ae
- 2 Assistant professor, Clinical Pharmacy department, College of Pharmacy, Al-Azhar University, Cairo, Egypt. Mohamed.baraka2020@gmail.com
- 3 Department of Clinical Pharmacy, School of Pharmaceutical Sciences, Universiti Sains Malaysia, Penang, Malaysia. amanyalboghdadly@gmail.com
- 4 Samar Z. Alshawwa, Assistant Professor, Pharmaceutical Sciences Department, College of Pharmacy, Princess Nourah bint Abdulrahman University; Riyadh, Saudi Arabia; samarzuhair@yahoo.com
- 5 Program of Clinical Pharmacy, College of Pharmacy, Al Ain University, Abu Dhabi Campus-UAE. Email: assahura2021@gmail.com
- 6 Pharmacy Practice department, College of Clinical Pharmacy, Imam Abdulrahman Bin Faisal University (University of Dammam), Saudi Arabia. hassanalsultan3@gmail.com, maislam@iau.edu.sa
- 7 Department of Pharmacy practice, King Fahd University, Riyadh, Saudi Arabia; kelfass@kfu.edu.sa
- 8 College of Medicine, Ajman University, Ajman, UAE. y.mohamed@ajman.ac.ae
- 9 Faculty of Pharmacy Boys, Al-Azhar University, Cairo, Egypt
- 10 Director of pharmacy Services department, King Fahd Hospital of the University (KFHU), Imam Abdulrahman Bin Faisal University (University of Dammam), Saudi Arabia. aaasseri@iau.edu.sa
- 11 Pharmaceutical Sciences, College of Pharmacy, Al Ain University, Al Ain, Abu Dhabi, United Arab Emirates. Pharmacy@au.ac.ae

* Correspondence: Mohamed.baraka2020@gmail.com, Mohamed Baraka
samarzuhair@yahoo.com, Samar Z. Alshawwa.

Abstract: (1) Background: Factors reported in literature associated with inappropriate prescribing of antimicrobials, such as physicians with less experience, uncertain diagnosis, and patient caregiver influences on physicians' decisions. Monitoring antimicrobial resistance is critical for identifying emerging resistance patterns, developing, and assessing the effectiveness of mitigation strategies. Improvement in prescribing 'antimicrobials would minimize the risk of resistance and, consequently, improve patients' clinical and health outcomes. The purpose of the study is to delineate factors associated with antimicrobial resistance, describe the factors influencing prescriber's choice during prescribing of antimicrobial, and examine factors related to consequences of inappropriate prescribing of antimicrobial. (2) Methods: Cross-sectional study conducted among healthcare providers (190) in six tertiary hospitals in the Eastern province of Saudi Arabia. The research panel has developed validated and piloted survey specific with closed-ended questions. A value of $P < 0.05$ was considered for statistical significance. All data analysis was performed using Statistical Package for Social Sciences (IBM SPSS version 23.0). (3) Results: 72.7% of the respondents have agreed that poor skills and knowledge are key factors that contribute to the inappropriate prescribing of antimicrobials. All the respondents acknowledged effectiveness and previous experience with the antimicrobial, and reading scientific materials (such as books, articles, and the internet) were key factors influencing physicians' choice during antimicrobial prescribing. (4) Conclusion: The current study has identified comprehensive education and training needs for healthcare providers about antimicrobial resistance. Using antimicrobials unnecessarily, insufficient duration of antimicrobial use, and using broad spectrum antimicrobials, were reported to be common practices. Further, poor skills and knowledge were a key factor that contributed to the inappropriate use and overuse of antimicrobials and using antimicrobials without physician prescription (self-medication) were the

key factors which contribute to AMR from participants' perspectives. Furthermore, internal policy and guidelines are needed to ensure that the antimicrobials are prescribed in accordance with standard protocols and clinical guidelines.

Keywords: antimicrobial resistance; AMR; Infections; antibiotics; inappropriate prescribing; healthcare professionals; education; training; antimicrobial stewardship programs; continuous professional development.

1. Introduction

Infectious diseases are caused by the microorganism's penetration into the epidermis, dermis, soft tissues of the skin, and other body tissues, leading to infectious diseases and comorbidities that can ultimately lead to sepsis and death. Antimicrobial agents have the advantage of treating infectious diseases, and they have the potential not only to retain the quality of life of the patient but have also proved to be lifesaving in a variety of severe infectious conditions [1]. Improper use of antimicrobials has led to the development of antimicrobial resistant (AMR), which has been associated with failure in managing infections, increased in length of hospital stay, increased mortality, morbidity [2,3], and health costs [4,5]. Several factors have been described in the literature to be correlated with inappropriate prescribing of antimicrobials, such as physicians with less experience, level of knowledge, uncertain diagnosis, and patient caregiver influences on physician's decision [6]. Over two million infections and 23,000 deaths annually in the United States were due to AMR [7]. Studies from diverse settings estimate that between 25% and 50% of antibiotic use in hospitals was suboptimal or unnecessary [8,9]. Gulf Cooperation Council countries face challenges of emerging AMR with no clear, regional guidelines for antimicrobial use or specific policies for restricting and monitoring antimicrobial prescribing [10]. Monitoring AMR is critical for identifying emerging resistance and developing, monitoring, and evaluating the effectiveness of mitigation strategies [11]. Improvement in antimicrobial prescribing would minimize the risk of AMR, reduce healthcare costs, and improve patient clinical outcomes. In attempt from world health organization (WHO) to reduce AMR, several initiatives and programs; have been established to educate healthcare providers about appropriate antimicrobial prescribing for evidence-based indications [12-13]. A study conducted in Saudi Arabia has revealed that medications most commonly used were antibiotics [14]. In another study, it has been confirmed that certain types of medications, such as chemotherapeutic agents and antipsychotics, are exclusively prescribed by the specialists involved, while antibiotics are vigorously prescribed by all physicians, as well as by allied health practitioners, irrespective of their knowledge or experience about antibiotics prescribed [15]. According to, general practitioners are the most common prescribers of antibiotics in Saudi Arabia relative to specialists and residents. The objectives of the recent study were to delineate the factors associated with antimicrobial resistant, describe the factors influencing prescriber's choice during prescribing of antimicrobial, and examine the factors related to consequences of inappropriate prescribing of antimicrobial.

2. Results

A total of 184 healthcare providers have responded to the survey with 96.8% response rate. Results have demonstrated that more than half of the respondents (53.8%) were pharmacists, (35.8%) were physicians, and (10.9%) were nurses (56% were males). Nearly 56% of respondents' age was less than 30 years; around one-third (30.4%) were in the age strata 31 to 40 years, and only 13.6% were aged ≥ 41 years. Regarding years of practice, 37% of the respondents had less than 3 years of clinical experience, whereas only 16% reported that their experience exceeded ten years. The majority (72.8%) have worked more than 8 hours per day. Regarding the average of continuous professional development (CPD), (71%) of the respondents had more than 20 CPD hours per year; Table 1.

Table 1. Descriptive statistics for socio-demographic characteristics of respondents (n=184)

Characteristics	Group	Frequency (n)	Percentage (%)
Gender	Male	103	56.0
	Female	81	44.0
Age (year)	Less than 30	103	56.0
	31-40	56	30.4
	≥ 41	25	13.6
Nationality	Saudi	154	83.7
	Non-Saudi	30	16.3
Region	Eastern of Saudi Arabia	178	96.7
	Others	6	3.3
Profession	Nurses	20	10.9
	Physicians	65	35.3
	Pharmacists	99	53.8
Country of last professional degree	Saudi Arabia	149	81.0
	Others	35	19.0
Working hours (per day)	Less than 8 hours	50	27.2
	8 hours and more	134	72.8
Years of Practice	Less than 3 years	68	37.0
	3-6 years	50	27.2
	6-10 years	36	19.6
	More than 10 years	30	16.3
Average of CME (continuous medical education) hours/year	Less than 20	53	28.8
	20 and above	131	71.2

In the current study, six factors with various questions (including yes/no response) on antimicrobial dispensing and prescribing practices were analyzed. At first, we have assessed the factors influencing physicians' choice during antimicrobial prescribing,

which was reflected on eight questions. Results revealed that for the first question, "Parent/patient's demand" 32.6% have responded positively. For the second question, "Reading scientific materials (such as books, articles, and internet)" 56.0% have responded positively as well. Responses retrieved from the third question, "Attending courses and lectures" were positively reported in 36.4%. Nearly half of the respondents (46.2%) agreed with the fourth question, "Cost of antibiotic" while (59.8%) agreed with the fifth question, "Effectiveness and previous experience with the prescribed antimicrobial. Only 23.4% responded yes for the sixth question, "Recommendation by other colleagues, while (39.7%) responded yes to the seventh question, "The knowledge gained during undergraduate or postgraduate training". Nearly half (49.5%) of the respondents have agreed with the eighth question, "Availability of the antibiotic in the hospital formulary"; Table 2.

Table 2. Factors influencing physicians' choice during antimicrobial prescribing

Items	Short item name	Responses	
		Yes (n & %)	No (n & %)
Parent/patient's demand	PD	60 (32.6)	124 (67.4)
Reading scientific materials (such as, books, articles, and internet)	RSM	103 (56.0)	81 (44.0)
Attending courses and lectures.	ACL	67 (36.4)	117 (63.6)
Cost of antibiotic	CA	85 (46.2)	99 (53.8)
Effectiveness and previous experience with the drug antimicrobial	EEDA	110 (59.8)	74 (40.2)
Recommendation by other colleagues	RC	43 (23.4)	141 (76.6)
The knowledge gained during undergraduate or postgraduate training	KT	73 (39.7)	111 (60.3)
Availability of the antibiotic in the hospital formulary	AAHF	91 (49.5)	93 (50.5)

Findings revealed that maximum 72.7% of the respondents have agreed that poor skills and knowledge was a key factor which contributes to the inappropriate prescribing of antimicrobials. However, nearly half of respondents (47.5%) have concurred that unrestricted availability of antimicrobials, inadequate supervision (45.4%), overworked/busy health care providers (32.2%), lack of physicians' interest in the subject of antimicrobial prescribing (41.5%) and lack of microbiology lab facilities (43.7%) were the major causes of inappropriate use of antimicrobial; Table 3.

Table 3. Factors related to important causes of inappropriate prescribing of antimicrobial

Items	Short item name	Responses	
		Yes (n & %)	No (n & %)
Poor skills and knowledge	PSK	133 (72.7)	50 (27.3)
Unrestricted availability of antimicrobials	UAA	87 (47.5)	96 (52.5)
Inadequate supervision	IS	83 (45.4)	100 (54.6)
Lack of physician interest in the subject of antimicrobial prescribing and infection management	LPI	76 (41.5)	107 (58.5)
Lack of effective hospital policies	LHP	85 (46.4)	98 (53.6)
Overworked/busy health care providers	OHP	59 (32.2)	124 (67.8)
Lack of microbiology lab facilities (lack of antimicrobial susceptibility test results)	LML	80 (43.7)	103 (56.3)

Concerning the causes of AMR, the highest number of respondents (80.4%) has concurred that using antimicrobial when they were not necessary, whereas not completing the full course of antimicrobial (59.8%), using antimicrobials without physician prescription as self-medication (66.3%) were reported as the key factors which contribute to AMR; Table 4.

Table 4. Factor associated with the causes of antimicrobial resistant

Items	Short item name	Responses	
		Yes (n & %)	No (n & %)
Using antimicrobial when they are not necessary	UANN	148 (80.4)	36 (19.6)
Not completing the full course of antimicrobial	NCFA	110 (59.8)	74 (40.2)
Using the same antimicrobial with a different brand	USADB	47 (25.5)	137 (74.5)
Using antimicrobials without physician prescription (self-medication)	UAWPP	122 (66.3)	62 (33.7)
Using broad spectrum antimicrobial, as a common practice	UBSA	94 (51.1)	90 (48.9)
Patient pressure for antibiotics as part of treatment	PPAT	77 (41.8)	107 (58.2)
Patients are able to buy antibiotics over-the-counter	PABA	80 (43.5)	104 (56.5)

Regarding the factors detecting the important consequences of antimicrobial overuse, the results have shown that the majority (78.8%) of respondents agreed that it will lead to AMR. The inverse results found for the question 'reduce the duration of hospitalization' have shown that 79.9% of respondents disagreed with this statement. Further, 85.9% of respondents didn't believe that better patient outcomes can be a consequence of antimicrobial overuse. (54.9%) of respondents have denied that waste of resources and recurrence of infections (56.5%) can be the consequence of antimicrobials overuse; Table 5.

Table 5. Factor related to important consequences of antimicrobial overuse

Items	Factor short name	Responses	
		Yes (n & %)	No (n & %)
Antimicrobial resistant	AMR	145 (78.8)	39 (21.2)
Adverse drug reactions and medication errors	ADRME	78 (42.4)	106 (57.6)
Reduce duration of hospitalization	RDH	37 (20.1)	147 (79.9)
Better patient outcome	BPO	26 (14.1)	158 (85.9)
Waste of resources	WR	83 (45.1)	101 (54.9)
Recurrence of infections	RI	80 (43.5)	104 (56.5)

In order to assess the factors that may help to control AMR, nine questions were used. Results demonstrated that the question about 'Physician education' reported a higher proportion (66.8%) of respondent's agreement. Whereas the question of 'consulting with infectious diseases experts' 55.4% of respondents have agreed, and 58.2% have believed that providing local antimicrobial guidelines will contribute to the control of AMR. However, only 56% have agreed with the statement that providing knowledge of pathogens and antimicrobial susceptibility test results will contribute to control AMR ¹⁶, ¹⁷. The respondents have rejected the assumptions that obtaining local AMR profiles, practicing antimicrobial restriction, removing catheters when not essential, and targeting antimicrobial therapy to likely pathogens could contribute to controlling AMR (51.6%), (59.2%), (66.3%), and (70.1%), respectively; Table 6.

Table 6. Factors that may help to control antimicrobial resistant

Items	Factor short name	Responses	
		Yes (n & %)	No (n & %)
Treating infection, not contamination or colonization	TINCC	74 (40.2)	110 (59.8)
Physician education on appropriate antimicrobial therapy	PEAAT	123 (66.8)	61 (33.2)
Consulting with infectious diseases experts	CIDE	102 (55.4)	82 (44.6)
Providing local antimicrobial guidelines	PLAG	107 (58.2)	77 (41.8)
Knowledge of pathogens and antimicrobial susceptibility test results	KPASR	103 (56.0)	81 (44.0)
Obtaining local antibiotic resistant profiles	OLARP	89 (48.4)	95 (51.6)
Practicing antimicrobial restriction	PAR	75 (40.8)	109 (59.2)
Removing catheters when not essential	RCNE	62 (33.7)	122 (66.3)
Targeting antimicrobial therapy to likely pathogens	TALP	55 (29.9)	129 (70.1)

The survey's final theme was designed to know about the most common resources used by healthcare providers for knowledge towards antimicrobials. Slightly above half (57.6%) of respondents have agreed that the Lexicomp/Sanford guide/up-to-date or other online resources were common resources. Most respondents, on the other hand, did not

agree with other common resources, such as national/Saudi guidance, hospital/internal guidelines, smartphone medical applications, Red Book, Quick reference guides or booklets or medical journals, and other guidelines by professional organizations, infectious diseases specialists/service in the hospital, colleagues or seniors from your own team or specialty. The clinical pharmacists and medical representatives have not been considered as a source of information for antimicrobials Table 7.

Table 7. Factors associated with most common resources used for knowledge toward antimicrobials

Items	Factor short name	Responses	
		Yes (n & %)	No (n & %)
National/Saudi guidelines	NG	83 (45.1)	101 (54.9)
Hospital/internal guidelines	HG	78 (42.4)	106 (57.6)
Lexicomp/Sanford guide/UPTODATE or other Online resources	LG	106 (57.6)	78 (42.4)
Smart phone medical applications	SMA	80 (43.5)	104 (56.5)
Red Book, Quick reference guides or booklets or Medical journals	RBRJ	41 (22.3)	143 (77.7)
Other guidelines by professional organizations (IDSA,CDC,...etc)	OG	62 (33.7)	122 (66.3)
Infectious diseases specialists/service in the hospital	IDSH	66 (35.9)	118 (64.1)
Colleagues or seniors from your own team or specialty	CTS	36 (19.6)	148 (80.4)
Hospital/Clinical pharmacists or Drug information centres	HPGIC	62 (33.7)	122 (66.3)
Medical/Pharmaceutical representatives	MR	31 (16.8)	153 (83.2)
Others	Others	9 (4.9)	175 (95.1)

3. Discussion

Our study documented the perceptions of healthcare professionals working in six tertiary hospitals in the eastern province of Saudi Arabian tertiary hospitals. The current study has delineated the factors associated with antimicrobial resistance (AMR), described the factors influencing prescriber’s choice during prescribing antimicrobials, and examined the factors related to the consequences of inappropriate prescribing of antimicrobials.

This study has revealed that all the respondents acknowledged the importance of previous experience with antimicrobials, and reading scientific materials as the key factors influencing physicians’ choice during antimicrobial prescribing. Conforming with a study conducted in Jeddah city in Saudi Arabia, our results confirmed their study findings [1]. Interestingly, attending courses and lectures was not considered a factor that may help control AMR. This might explain the poor knowledge among our respondents about the importance of attending courses and lectures to improve their antimicrobial practices.

Similar to healthcare professionals from other parts of the world, 72.7% of the respondents agreed that poor skills and knowledge were a key factor that contributes to the inappropriate use of antimicrobials [1,18]. However, most of the respondents have agreed that unrestricted availability of antimicrobials, overworked/busy health care providers and lack of physicians' interest in the subject of antimicrobial prescribing and infection management were not causes of inappropriate prescribing of antimicrobials.

Many healthcare providers in our study have agreed that using antimicrobials when they were not necessary, not completing the full course of antimicrobial, and using broad spectrum antimicrobial, were common practices. Further, using antimicrobials without physician prescription (self-medication) were the key factors which contribute to AMR [16,19]. Our findings lend support to previously published studies [1,14,18]. In response to this and based on the current study results, this can be controlled via increasing the sessions of CPD for healthcare providers on appropriate antimicrobial treatment, providing local antimicrobial guidelines and consulting with experts on infectious diseases. The majority of healthcare professionals have believed that AMR was a consequence of antimicrobial overuse. Whereas they have denied that the waste of resources and recurrence of infections can be the consequences of antimicrobials overuse. We highly encourage each hospital to generate their internal policy or guidelines to guide and restrict antimicrobials' overuse. In response to this and based on healthcare professionals' responses here, this can be rectified via CPD awareness programs and education on appropriate antimicrobial therapy, providing local antimicrobial guidelines, consulting with infectious diseases experts, and above all implementing antimicrobial stewardship program. Yusef ,D. et al. concluded that antimicrobial stewardship programs (ASPs) helped to reduce the use of antifungals in hospitalized patients and highlighted the importance of promoting antifungal stewardship among antibiotic stewardship [20]. Also, Rodolfo E. Quirós et al. showed significant improvement in antimicrobial utilization after multiple and intensive ASPs [3]. Interestingly, targeting antimicrobial therapy to likely pathogens was not mostly considered a factor that may help control AMR. This may be due to the poor of knowledge in our respondents on the importance of targeting pathogens. Some recent studies conducted in Saudi Arabia by Al-Harthi and colleague were in line with our results, where they identified the importance of healthcare professional education on appropriate antimicrobial therapy, providing local antimicrobial guidelines and consulting with infectious diseases experts as crucial factors that can control AMR and its consequence. [5]

In fact, further studies are needed to target the reasons behind the gap in knowledge, experience, and skills on antimicrobials, as revealed by the current study respondents. This will help in controlling AMR and its consequences in our region.

Limitations of the study:

There are some limitations to be considered when interpreting the study findings. First, although the research was conducted at six hospitals in the Eastern province that were well-resourced, the findings of the study may not be generalizable to other regions in the Kingdom. Second, participants were selected randomly and the characteristics of those who did not participate in the survey are unknown. Further, purposive sampling

was used to target doctors, nurses, and pharmacists from departments with high antibiotic use; therefore, there are departments such as psychiatry for example that are underrepresented. Therefore, it is possible that the sampling approach and recruitment limit the generalizability of the study results to other departments or hospitals. Finally, the sample included in the study was small, with a moderate response rate. We believe the results from the current study will be valuable in guiding educational efforts in order to prevent antimicrobial resistance and well received by clinicians in their respective institutions. However, our study strength is that, for the best of our knowledge, it is the first study that investigates factors affecting healthcare providers' perceptions regarding AMR consequences.

4. Materials and Methods

The study was a cross-sectional study among health care providers (190) in six tertiary hospitals (4 governmental and 2 private hospitals with more than 500 beds) in the Eastern province of Saudi Arabia.

Sample size

A published statistical formula was used to calculate the sample size for the unknown nature of population size. Finally, the sample was adjusted for potential missing or non-response error using the formula:

$$n = \frac{(Z_{1-\beta})^2 [p(1-p)]}{d^2}$$

Where, n = required sample size; $Z_{1-\beta}$ = Z value at power $1-\beta$ (at power 90% this value is 1.24); p = preferred population proportion (0.5); d = margin of error (ideal value is 0.05). And. $n_1 = n/(1 - e)$ Where 'n' is the required sample size as per formula, 'n1' is adjusted sample size and 'd' is the potential missing or non-response rate. A minimum number of 190 samples was required to achieve 90% power when considering a 5% margin of error and a 10% missing/non-response error rate.

Data collection instrument and quality assurance

Survey development

The research panel has developed validated and piloted survey (divided into five parts) with closed ended questions. The initial version of the structured survey was developed after in-depth literature review to assess AMR associated factors and their consequences. The expert advisory panel (three pharmacists, statistician, physician, and the director of clinical pharmacy services) finalized the survey and assured the quality after addressing the comments and corrections.

The survey was subdivided into five parts; first (part A), including respondents' characteristics and experiences. The second part (part B) has described the factors related to antimicrobial prescribing and dispensing practices. This has contained a total of six factors with different options (allowed to choose multiple options). In addition to evaluating the viability of the survey, the survey has been piloted to ensure face validity and

has resolved both improvements in terminology and ease of use. However, the data collected from the pilot study were not involved in the final analysis. In terms of reliability, a variety of factors have been taken into account when planning this analysis to reduce the risk to the reliability. This has included: the data collection process was clearly documented, and research procedures were followed as per the data collection protocol during the research process. Further, the results were obtained on one occasion to simplify the capture of answers from participants, and to help mitigate any unintended bias in the analysis of responses, closed-ended questions were purposely selected for this survey design.

Administration of the survey

A standard set of the pre-tested validated survey was used to collect data among health care providers (physicians, pharmacists, and nurses) by self-administrative procedure. In addition, questions for online surveys (QuestionPro) have been distributed to the hospital administration, pharmacy directors, and nursing supervisors via e-mails and related social media accounts to boost data collection. The data collection procedure was administrated by the investigators via several necessary visits, reminders by phone calls as well as emails.

Statistical Analysis

The online survey (QuestionPro) has enabled the view of some missing cases among the data set. Therefore, we have used the 'last-observation-carried-forward method', some of these cases were treated, and the rest were cleaned from the final analysis. Additionally, informal technique was used to check and cleaned the whole data set. Variables were coded and recoded as per the requirements of study objectives. Descriptive statistics were utilized for data analysis. Frequencies and their corresponding percentages were reported to describe the factors and their respective items. A value of $p < 0.05$ was considered for statistical significance. The whole data analysis was conducted using Statistical Package for Social Sciences (IBM SPSS version 23.0).

5. Conclusions

The current study has identified comprehensive education and training needs for healthcare providers about the AMR and how to prevent such problem. Using antimicrobials unnecessarily, insufficient duration of antimicrobial use, and using broad spectrum antimicrobials, were reported to be common practices. Further, poor skills and knowledge were a key factor that contributed to the inappropriate use and overuse of antimicrobials and using antimicrobials without physician prescription (self-medication) were the key factors which contribute to AMR from participants' perspectives. This problem can be controlled via increasing the sessions of CPD for healthcare providers on appropriate antimicrobial treatment, providing local antimicrobial guidelines and consulting with experts on infectious diseases. More studies about local AMR patterns are deemed, and more efforts are warranted to educate clinicians about antimicrobial resistance. We highly encourage each hospital to generate their internal policy or guidelines to guide and restrict antimicrobials' overuse to combat AMR.

Author Contributions: “Conceptualization, Mohamed A. Baraka, Hassan Al Sultan, Taha Al Salman, and Hussain Alaithan, Md. Ashraf Islam, Amany Alboghdadly, Samar Alshawwa, Asim Ahmed. and Abdulsalam A. Alasser; methodology, Mohamed A. Baraka, Amany Alboghdadly, Samar Alshawwa, Asim Ahmed Elnour, Hassan Alsultan, Md. Ashraf Islam, Kareem Ahmed El-Fass, Yehia Mohamed, Abdulsalam A. Alasser and Khairi Mustafa Salem; software, Mohamed A. Baraka, Md. Ashraf Islam and Kareem Ahmed El-Fass.; validation, Mohamed A. Baraka, Amany Alboghdadly, Md. Ashraf Islam, Kareem Ahmed El-Fass, Yehia Mohamed.; formal analysis, Mohamed A. Baraka, Md. Ashraf Islam and Kareem Ahmed El-Fass.; investigation, Mohamed A. Baraka, Samar Alshawwa, Hassan Alsultan, Taha Al Salman, Hussain Alaithan, Abdulsalam A. Alasser.; resources, Mohamed A. Baraka, Amany Alboghdadly, Samar Alshawwa and Khairi Mustafa Salem.; data curation, Mohamed A. Baraka, Hassan Alsultan, Taha Al Salman, Hussain Alaithan, Md. Ashraf Islam.; writing—original draft preparation, Mohamed A. Baraka, Amany Alboghdadly, Samar Alshawwa, Asim Ahmed Elnour, Hassan Alsultan, Taha Al Salman, Hussain Alaithan, Md. Ashraf Islam.; writing—review and editing, Mohamed A. Baraka, Amany Alboghdadly, Samar Alshawwa, Asim Ahmed Elnour, Md. Ashraf Islam, Kareem Ahmed El-Fass, Yehia Mohamed and Khairi Mustafa Salem.; visualization, Mohamed A. Baraka, Amany Alboghdadly, Samar Alshawwa, Asim Ahmed Elnour.; supervision, Mohamed A. Baraka, Amany Alboghdadly, Samar Alshawwa and Khairi Mustafa Salem.; project administration, Mohamed A. Baraka, Samar Alshawwa and Khairi Mustafa Salem; funding acquisition, Mohamed A. Baraka, Samar Alshawwa and Alasser. All authors have read and agreed to the published version of the manuscript

Funding: This research was funded by the Deanship of Scientific Research at Princess Nourah bint Abdulrahman University through the Fast-track Research Funding Program.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of Imam Abdulrahman Bin Faisal University (IRB – UGS – 2019 – 5 – 088).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data will be available upon request.

Acknowledgments: In this section, you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- (1) Al-Harthi, S. E.; Khan, L. M.; Abed, H. H.; Alkreathy, H. M.; Ali, A. S. Appraisal of Antimicrobial Prescribing Practices of Governmental and Non-Governmental Dentists for Hospitals in the Western Region of Saudi Arabia. *Saudi Med J* **2013**, *34*, 1262–1269.
- (2) Alshammari, T. M. ; Larrat, E. P. ; Caffrey, A. R. ; Quilliam, B. J. ; LaPlante, K. L. The Risk of Hepatotoxicity with Fluoroquinolones: A National Case-Control Safety Study. *Am J Heal. Syst Pharm* **2014**, *71*, 37–43. <https://doi.org/10.2146/ajhp130165>.
- (3) Quirós, R. E. ; Bardossy, A. C. ; Angeleri, P. ; Zurita, J. ; Aleman Espinoza, W. R. ; Carneiro, M. ; Guerra, S. ; Medina, J. ; Castañeda Luquerna, X. ; Guerra, A. ; Vega, S. ; Cuellar Ponce de Leon, L. E. ; Munita, J. ; Escobar, E. D. ; Maki, G. ; Prentiss, T. ; Zervos, M. Antimicrobial Stewardship Programs in Adult Intensive Care Units in Latin America: Implementation, Assessments, and Impact on Outcomes. *Infect. Control Hosp. Epidemiol.* **2021**, 1–10. <https://doi.org/10.1017/ice.2021.80>.
- (4) Nasr, Z. ; Babiker, A. ; Elbasheer, M. ; Osman, A. ; Elazzazy, S. ; Wilby, K., J. Practice Implications of an Antimicrobial Stewardship Intervention in a Tertiary Care Teaching Hospital, Qatar. *EMHJ* **2019**, *25*, 172–180. <https://doi.org/10.26719/emhj.18.026>.

- (5) Al-Harthi, S. E. ; Khan, L. M. ; Osman, A. M. ; Alim, M. A. ; Saadah, O. I. ; Almohammadi, A. A. ; Khan, F. M. ; Kamel, F. O. Perceptions and Knowledge Regarding Antimicrobial Stewardship among Clinicians in Jeddah, Saudi Arabia. *Saudi Med J* **2015**, *36*, 813–820. <https://doi.org/10.15537/smj.2015.7.11833>.
- (6) Suaifan, G. ; Shehadeh, M. ; Darwish, D. ; Al-Ijel, H. ; Yousef, A. ; Darwish, R. A Cross-Sectional Study on Knowledge, Attitude and Behavior Related to \ Antibiotic Use and Resistance among Medical and Non-Medical University Students in Jordan. *AJPP* **2012**, *6*, 763–770. <https://doi.org/10.5897/ajpp12.080>.
- (7) Centers for Disease Control and Prevention. *Antibiotic Resistance Threats in the United States, 2013* . US Department of Health and Human Services, Centers for Disease Control and Prevention.; 2013.
- (8) Willemsen, I. ; Groenhuijzen, A. ; Bogaers, D. ; Stuurman, A. ; Van Keulen, P. ; Kluytmans, J. Appropriateness of Antimicrobial Therapy Measured by Repeated Prevalence Surveys. *Antimicrob. Agents Chemother.* **2007**, *51*, 864–867. <https://doi.org/10.1128/AAC.00994-06>.
- (9) Kapadia, S. N. ; Abramson, E. L. ; Carter, E. J. ; Loo, A. S. ; Kaushal, R. ; Calfee, D. P. ; Simon, M. S. The Expanding Role of Antimicrobial Stewardship Programs in Hospitals in the United States: Lessons Learned from a Multisite Qualitative Study. *Jt. Comm. J. Qual. Patient Saf.* **2018**, *44*, 68–74. <https://doi.org/10.1016/j.jcjq.2017.07.007>.
- (10) Aly, M. ; Balkhy, H. H. The Prevalence of Antimicrobial Resistance in Clinical Isolates from Gulf Corporation Council Countries. *Antimicrob. Resist. Infect. Control* **2012**, *1*. <https://doi.org/10.1186/2047-2994-1-26>.
- (11) Karp, B. E. ; Tate, H. ; Plumblee, J. R. ; Dessai, U. ; Whichard, J. M. ; Thacker, E. L. ; Hale, K. R. ; Wilson, W.; Friedman, C. R.; Griffin, P. M.; McDermott, P. F. National Antimicrobial Resistance Monitoring System: Two Decades of Advancing Public Health through Integrated Surveillance of Antimicrobial Resistance. *Foodborne Patho Dis.* **2017**, *14*, 545–557. <https://doi.org/10.1089/fpd.2017.2283>.
- (12) Vickers, H. International Antibiotic Resistance Crisis. *BMJ* **2011**, *342*, d3207. <https://doi.org/10.1136/sbmj.d3207>.
- (13) Tonkin-Crine, S. ; Yardley, L. ; Little, P. Antibiotic Prescribing for Acute Respiratory Tract Infections in Primary Care: An Updated and Expanded Meta-Ethnography. *J Antimicrob Chemother* **2011**, *66*, 2215–2223. <https://doi.org/10.3399/bjgp18X697889>.
- (14) AlKhamees, O. A. ; AlNemer, K. A. ; Bin Maneea, M. W. ; AlSugair, F. A. ; AlEnizi, B. H. ; Alharf, A. A. Top 10 Most Used Drugs in the Kingdom of Saudi Arabia 2010–2015. *SPJ* **2018**, *26* (2), 211–216. <https://doi.org/10.1016/j.jsps.2017.12.009>.
- (15) Edgar, T. ; Boyd, S. D. ; Palamé, M. J. Sustainability for Behaviour Change in the Fight against Antibiotic Resistance: A Social Marketing Framework. *JAC* **2009**, *63*, 230–237. <https://doi.org/10.1093/jac/dkn508>.
- (16) Hayat, K. ; Rosenthal, M. ; Gillani, A. H. ; Chang, J. ; Ji, W.; Yang, C. ; Jiang, M. ; Zhao, M. ; Fung, Y. Perspective of Key Healthcare Professionals on Antimicrobial Resistance and Stewardship Programs: A Multicenter Cross-Sectional Study from Pakistan. *Front. Pharmacol.* **2020**, *10*. <https://doi.org/10.3389/fphar.2019.01520>.
- (17) Labi, A.-K. ; Obeng-Nkrumah, N. ; Bjerrum, S. ; Aryee, N. A. A. ; Ofori-Adjei, Y. A. ; Yawson, A. E. ; Newman, M. J. Physicians' Knowledge, Attitudes, and Perceptions Concerning Antibiotic Resistance: A Survey in a Ghanaian Tertiary Care Hospital. *BMC Health Serv. Res.* **2018**, *18*, 126. <https://doi.org/10.1186/s12913-018-2899-y>.
- (18) Baadani, A. M. ; Baig, K. ; Alfahad, W. A. ; Aldalbahi, S. ; Omrani, A. S. Physicians' Knowledge, Perceptions, and Attitudes toward Antimicrobial Prescribing in Riyadh, Saudi Arabia. *Saudi Med. J.* **2015**, *36*, 613–619. <https://doi.org/10.15537/smj.2015.5.11726>.
- (19) Gillani, A. H. ; Ji, W. ; Hussain, W. ; Imran, A. ; Chang, J. ; Yang, C. ; Fang, Y. Antibiotic Self-Medication among Non-Medical University Students in Punjab, Pakistan: A Cross-Sectional Survey. *Int.J Environ Res. Public Health.* **2017**. <https://doi.org/10.3390/ijerph14101152>.
- (20) Yusef, D. ; Hayajneh, W. A. ; Al-Azzam, S. ; Jaradat, S. K. ; Gould, I. ; Lattyak, E. A. ; Lattyak, W. J. ; Conway, B. R. ; Bond, S.; Conlon-Bingham, G.; Aldeyab, M. A. Impact of Antimicrobial Stewardship Interventions on Reducing Antifungal Use in

Hospitals in Jordan. *Infect. Control Hosp. Epidemiol.* **2021**, 1–2. <https://doi.org/10.1017/ice.2021.105>.