

Article

Malnutrition in infants aged under 6 months attending community health centres: A cross sectional survey

Carlos S. Grijalva-Eternod ^{1,2,*}, Emma Beaumont ¹, Ritu Rana ^{3,4}, Nahom Abate ⁵, Hatty Barthorp ⁴, Marie McGrath ^{1,6}, Ayenew Negesse ⁷, Mubarek Abera ⁸, Alemseged Abdissa ^{8,9}, Tsinuel Girma ^{8,10}, Elizabeth Allen ¹, Marko Kerac ¹, Melkamu Berhane ⁸

¹ London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK; Carlos.Grijalva-Eternod@lshtm.ac.uk (C.S.G.-E.); Emma.Beaumont@lshtm.ac.uk (E.M.); Elizabeth.Allen@lshtm.ac.uk (E.A.); Marko.Kerac@lshtm.ac.uk (M.K.)

² UCL Institute for Global Health, 30 Guilford Street, London, WC1N 1EH, UK

³ Indian Institute of Public Health, Gandhinagar 382042, Gujarat, India; rrana@iiphg.org (R.R.)

⁴ GOAL Global, Carnegie House, Library Road, Dun Laoghaire, A96 C7W7, Dublin, Ireland; hbarthorp@goal.ie (H.B.)

⁵ GOAL Ethiopia, Yeka Subcity, Woreda 9, Addis Ababa, P.O. Box 5504, Ethiopia; nahoma@et.goal.ie (N.A.)

⁶ Emergency Nutrition Network, 69 High Street, Marlborough House, Kidlington, Oxfordshire, OX5 2DN, UK; marie@ennonline.net (M.M.)

⁷ Debre Markos University, Debre Markos, Ethiopia; ayenewnegesse@gmail.com (A.N.)

⁸ Jimma University, Jimma, Ethiopia; abmubarek@gmail.com (M.A.); alemseged.abdissa@gmail.com (A.A.); tsinuel@yahoo.com (T.G.); melkamuarefayine@gmail.com (M.B.)

⁹ Armauer Hansen Research Institute, Addis Ababa, P.O. Box 1005, Ethiopia

¹⁰ Harvard Chan School of Public Health, Addis Ababa, Ethiopia

* Correspondence: Carlos.Grijalva-Eternod@lshtm.ac.uk

Abstract: Poor understanding of malnutrition burden is a common reason for not prioritizing the care of small and nutritionally at-risk infants aged under-six months (infants u6m). We aimed to estimate the anthropometric deficit prevalence in infants u6m attending health centres, using the Composite Index of Anthropometric Failure (CIAF); and to assess the overlap of different individual indicators. We undertook a two-week survey of all infants u6m visiting each of 18 health centres in two zones of the Oromia region, Ethiopia. We measured weight, length, and MUAC (Mid upper arm circumference); and calculated weight-for-length (WLZ), length-for-age (LAZ), and weight-for-age z-scores (WAZ). Overall, 21.7% (95% CI: 19.2; 24.3) of infants u6m presented CIAF and of these, 10.7% (95% CI: 8.9; 12.7) had multiple anthropometric deficits. Low MUAC overlapped with 47.5% (95% CI: 38.0; 57.3), 43.8% (95% CI: 34.9; 53.1), and 42.6% (95% CI: 36.3; 49.2) of the stunted, wasted and CIAF prevalence, respectively. Underweight overlapped with 63.4% (95% CI: 53.6; 72.2), 52.7% (95% CI: 43.4; 61.7), and 59.6% (95% CI: 53.1; 65.9) of the stunted, wasted and CIAF prevalence, respectively. Anthropometric deficits, single and multiple, are prevalent in infants attending health centres. WAZ overlaps more with other forms of anthropometric deficits than MUAC.

Keywords: Anthropometric deficit; infants under 6 months; malnutrition; MAMI, Ethiopia; weight-for-age; the Composite Index of Anthropometric Failure

1. Introduction

Globally, millions of infants aged under-six months (henceforth infants u6m) are small and nutritionally at-risk, with different forms of anthropometric deficits (e.g. wasted, underweight, or stunted) or low birthweight (LBW) [1]. These deficits are associated with increased risk of mortality, morbidity, subsequent malnutrition, and impaired development [2]. For instance, an estimated 8.5 million infants u6m are wasted, of whom 3.8 million are severely wasted [3]. In addition, an estimated 20.5 million livebirths have

a LBW [4]. These first six months of life are recognised as a crucial developmental period, often with unique nutritional challenges, but with life-long adverse consequences if malnutrition develops [5,6].

Over the last 20 years, Community-based Management of Acute Malnutrition (CMAM) has positively transformed the care of malnourished children aged 6-59 months (henceforth children) by relocating the focus of care from inpatient to outpatient settings [7]. This relocation expanded care accessibility and promoted early identification of at-risk children; improving in turn programme coverage and impact [5,8]. Despite these successes, malnourished infants u6m have been left behind [5]. In many settings, they are neither identified nor supported to survive and thrive as per the Every Woman Every Child Global strategy [5,9]. This is despite evidence of their markedly greater risk of mortality compared with malnourished children [10].

The 2013 update to the WHO guideline for management of severe malnutrition included for the first time a section on infants u6m, which incorporated recommendations for outpatient care for clinically stable malnourished infants u6m, rather than inpatient care for all, irrespective of clinical status, as was conventionally practiced [11]. However, most low- and middle-income countries (LMICs), have not adopted these updated recommendations into national protocols. Reasons for this include (i) a poor understanding of the malnutrition burden in infants u6m [5]; (ii) a problematic admission criteria based on weight-for-length z-scores (WLZ) [11] that has low reliability [12] and it is challenging to obtain in infants u6m, as measuring their length is difficult (more than in children), and WLZ cannot be calculated for lengths <45 cm [13]; (iii) infants u6m treatment needs are complex [5]; and (iv) the low quality evidence on the best strategies to meet those needs [11].

To address the first two reasons, weight-for-age z-score (WAZ) and the mid-upper arm circumference (MUAC) have been suggested as better alternatives for identifying small and nutritionally at-risk infants u6m, but there is sparse evidence on their prognostic value in this age group. In addition, optimal cut-off values still need to be determined [5,14-16]. The growing appreciation of the frequent co-existence of different types of anthropometric deficit further complicates our understanding on the best ways to define malnutrition programme admission criteria [17]. Towards this, the Composite Index of Anthropometric Failure (CIAF) is an aggregated indicator for malnutrition combining the conventional indicators wasted, stunted and underweight [18,19], but to our knowledge it has not been used to characterise malnutrition in infants u6m.

As in many other LMICs, infant malnutrition is a public health concern in Ethiopia [20]. Current Ministry of Health Guidelines recommend inpatient-only treatment for infants u6m, following identification using a WLZ <-3 [21]. To inform future programme planning and research (including sample size and logistics consideration for a randomised controlled trial (RCT) we are planning in Ethiopia), we aimed to generate caseload-evidence relevant to infants u6m malnutrition. The primary objectives of this study were to assess the prevalence of different indicators of anthropometric deficit in infants u6m attending health centres (this is where small and nutritionally at-risk infants u6m would be screened and recruited into future programmes) and to describe the patterns of overlap between different forms of anthropometric deficit in infants u6m. Our secondary objective was to assess the infant feeding practices of the surveyed infants u6m.

2. Materials and Methods

2.1. Study site

The study sites were in Deder woreda, East Hararge zone and in Jimma zone, Ethiopia. Though geographically separate, these are both located in the Oromia Region. They were chosen because they are the sites of our future RCT. The study was implemented in 18 health centres; ten in Jimma zone and eight in Deder woreda.

Jimma zone is one of the most populous areas of the Oromia Regional State, with a population of over 3 million people. Deder woreda has a population of some 315,000

people. Both sites have a high burden of malnutrition. Their main livelihood in the area is agriculture, petty trade of cash crops such as khat and coffee, fattening of oxen, and local casual labour. Jimma zone and Deder woreda have about 124 and 8 health centres, respectively; each serving an average population between 15,000 to 30,000.	100
	101
	102
	103
<i>2.2. Study design and population</i>	104
We undertook a health centre-based cross-sectional survey, surveying all infants u6m who attended the selected health centres for any reason over an average period of two weeks in each centre. Reasons for attendance included: born at the health centre; immunization clinics; growth monitoring clinics; under-5 clinics (where children present with a variety of acute illnesses). We collected data from 1,060 infants u6m between October 12 th , 2020 and January 29 th , 2021.	105
	106
	107
	108
	109
	110
<i>2.3. Sample size estimation</i>	111
We lacked prior information on how many infants u6m attending the health centres would have anthropometric deficits. Consequently, to estimate a sufficiently robust sample, we assumed a 50% prevalence of anthropometric deficit in infants u6m, and a 3% precision. Using these assumptions, we estimated to need a sample of 1067 infants u6m, an average of 60 infants u6m per health centre. To plan for field logistics, we assumed that each health centre would have an average attendance of 30 infants u6m per week and set the average duration of data collection for each health centre to a two-week period.	112
	113
	114
	115
	116
	117
	118
	119
<i>2.4. Health centre selection</i>	120
In Deder woreda, we included all eight available health centres. In Jimma zone, we selected ten out of the 124 available health centres as follows: First, we undertook a register review in all 124 health centres to collect eligibility information on ease of access and patient load. We excluded 60 health centres from which we were unable to gather complete eligibility data. We further excluded seven health centres that were difficult to access. We ranked the remaining 57 health centres according to patient load and randomly selected ten centres from the top 50%.	121
	122
	123
	124
	125
	126
	127
<i>2.5. Training of data collectors</i>	128
We undertook a training for the teams of enumerators and supervisors, one team in Jimma zone and one in Deder woreda, to ensure consistency and high quality of data collection. Our training included learning how to obtain anthropometric measurements (e.g. weight, length, MUAC), assess infant feeding practices, obtain economic and demographic data, use digital data gathering devices (DDGs), take informed consent, and obtain clinical history data. Our training also included the piloting of data collection, prior to initiating the actual survey data collection, to ensure collection of high-quality data and to identify and correct any sources of data collection errors. During the pilot, we also assessed field-challenges of the survey tool for final editing.	129
	130
	131
	132
	133
	134
	135
	136
	137
<i>2.6. Data and measurements</i>	138
All data was collected using an electronic questionnaire designed using the REDCap (Research Electronic Data Capture) project system (https://redcap.am.lshtm.ac.uk/redcap/).	139
	140
	141
At the household level, we obtained information about the sex and formal education of the household head, household size, number of dependent children aged <18 years. From mothers or primary caregivers, we obtained information on age, formal education, and religion.	142
	143
	144
	145
From all infants u6m we obtained sex and date of birth data. We asked mothers/caregivers to recall their infants' age in weeks. We asked whether the infant was born singleton or was a twin or a triplet; the infant's birth order; how many siblings aged <18 years they have, and whether any of them died recently. We collected data on	146
	147
	148
	149

infants' feeding practices in the past 24 hours. We asked about current and past breast- 150
feeding and whether they received any liquids such as water, milk, juice, broth, runny 151
porridge, yogurt, or other liquids apart from those mentioned. We asked whether they 152
were fed using a bottle and whether they received any solid, semi-solid or soft foods 153
[22]. 154

We measured weight with the infant undressed using a digital weight scale (Seca 155
354) to the nearest 5 g if weighing <10 kg, or to the nearest 10 g if weighing ≥10 kg. We 156
measured length using a UNICEF length/height board (infant/child/adult) to the nearest 157
completed 0.1 cm. We measured knee-heel length, from the heel to the upper part of the 158
distal femur (knee), using a digital Vernier calliper (Sealey AK9623EV) to the nearest 159
completed 0.1 cm. We measured MUAC using UNICEF MUAC tapes to the nearest completed 160
0.1 cm. We measured head circumference using standard anthropometric tapes to the 161
nearest completed 0.1 cm. We obtained subscapular and triceps skinfolds using a 162
Harpden skinfold calliper to the nearest completed 0.2 mm. 163

We collected all anthropometric measurements in pairs, as per WHO Child Growth 164
Standards protocols [23]. In brief, for each anthropometric measurement two enumerators 165
independently took one measurement. If the difference between these two measurements 166
exceeded the pre-set maximum allowed difference (see **Appendix Table A1**), each 167
enumerator took a second measurement and we excluded the first pair of measurements. 168
If the difference for their second pair of measurements also exceeded the allowed 169
difference, they each took a third measurement, and we excluded the first two pairs of 170
measurements. If the difference in their third pair of measurements exceeded the allowed 171
difference, no further measurements were taken, and we excluded all three pairs of 172
measurements. 173

2.7. Data handling and analysis

We handled and analysed data using Stata (StataCorp. 2019. Stata Statistical Software: 174
Release 16. College Station, TX: StataCorp LLC). 175

We estimated infant's age in weeks as the difference between the survey's date and 176
the infant's date of birth divided by seven. For infants whose date of birth was not 177
available, we used maternal recalled age. We grouped infants by age, in 5-weeks brackets. 178

We identified infants as being exclusively breastfed if their mothers or primary 179
caregivers reported in their 24 hours recall to having breastfed the infants but answer no 180
to all questions regarding any liquid intake, bottle feeding or feeding the infant with 181
solid, semi-solid or soft foods. 182

We calculated the mean value for each of the anthropometric measurements using 183
the pair of measurements that did not exceed the allowed difference. We used weight, 184
length and age data to estimate the anthropometric indices weight-for-age, length-for- 185
age, and weight-for-length z-scores (WAZ, LAZ, and WLZ, respectively) based on the 186
2006 WHO Child Growth Standards [13], using the zanthro Stata command [24]. We 187
marked anthropometric indices as outliers using the 2006 WHO recommendations [25], 188
i.e.: if they were >5 or <-6, >6 or <-6, and >5 or <-5 for WAZ, LAZ, and WLZ values, 189
respectively. 190

We defined underweight, stunted, and wasted as WAZ, LAZ, and WLZ <-2, respectively. 191
We used CIAF to assess overall malnutrition prevalence in infants u6m [18]. We 192
defined CIAF as all infants u6m that were either underweight, stunted or wasted and we 193
generated the following subcategories: wasted only, wasted and underweight, wasted, 194
stunted and underweight, stunted and underweight, stunted only, and underweight 195
only [18]. We defined the Composite Index of Severe Anthropometric Failure (CISAF) as 196
all infants u6m that were severely underweight, stunted or wasted, as defined by a 197
WAZ, LAZ, or WLZ <-3, respectively [26]. 198

To explore the overlap between MUAC and CIAF in infants, we explored the use of 199
different thresholds to define low MUAC: MUAC <11.5cm; MUAC <11.0cm if aged <6 200
weeks and <11.5 thereafter; MUAC <11.0cm if aged <7 weeks and <11.5 thereafter; 201
MUAC <11.0cm if aged <13 weeks and <11.5 thereafter; MUAC <11.0cm if aged <17 202
203

weeks and <11.5 thereafter; MUAC <11.0cm; and MUAC <10.5cm. These thresholds were chosen to match those in past/present use in older children; and the age thresholds to match timings of immunization clinic visits when future programmes would use MUAC for identification of at-risk infants u6m. 204
205
206
207

For the analysis, we excluded infants u6m for which we could not estimate all anthropometric indices as they had weight or length missing, or their length were <45cm, or if one or more of their anthropometric indices were marked as outliers. In addition, we excluded infants from the assessment of overlap between different anthropometric indicators if they presented with oedema. 208
209
210
211
212

We estimated means or proportions, along with the 95% confidence intervals (95% CI), for all variables. We compared basic characteristics between Jimma and Deder using t-test and z-test with the lincom Stata command. To determine the prevalence of different anthropometric deficits in infants, we estimated summary statistics for all anthropometric variables. 213
214
215
216
217

3. Results

3.1. Participants flow

Figure 1 shows the survey's participants flow. Of the infants u6m whose mothers or primary caregivers agreed to participate, a total of 2.17% (95% CI: 1.45; 3.25) had either missing weight or length data, their WLZ could not be estimated because their length were <45 cm, or had any of the anthropometric indices WLZ, WAZ, or LAZ identified as outliers. 220
221
222
223
224




Figure 1. Study participants' flow chart. 225
226

3.2. Sample characteristics

Key household and caregiver characteristics of 1,037 surveyed infants u6m with complete anthropometric information are presented in **Table 1**. Overall, most households were male-headed and comprised of an average of five members, three of whom 227
228
229
230

were children or adolescents aged <18 years. The infants' primary caregiver was most often the mother, who was on average in her mid-twenties. Levels of formal education were low: 34% of household heads and 41% of mothers/primary caregivers had no formal education. The infants' male:female sex ratio was 1.24:1; they were on average 13 weeks of age; most were singleton births. Mean WAZ, LAZ and WLZ were all below the 2016 WHO Child Growth Standards median. We found seven infants u6m who presented with oedema: these were excluded from subsequent anthropometric deficits assessment.

We found differences in some of the above mentioned characteristics between the Jimma and Deder sites: these are presented in **Supplementary Materials Table S1**. The main difference was that infants u6m in Deder presented with significantly worse nutritional status, as evidenced by greater underweight, wasted, and stunted prevalence.

3.3. Infant feeding practices

A summary of key infant feeding practices is also presented in **Table 1**. Reported breastfeeding was almost universal with over 98% of mothers reporting that infants u6m had ever been breastfed, 96% receiving breastmilk as the first food after birth, and 95% breastfed in the past 24 hours. Almost half of the infants u6m were reported to be exclusively breastfed at time of assessment and about 15% were bottle-fed. Only 2% were reported to have initiated consumption of any solid, semi-solid or soft foods.

Table 1. Characteristics of the 1,037 surveyed infants under six months of age.

	mean or %	95% CI
Household Characteristics		
Household members	5.4	5.3; 5.6
Household children aged <18 years	3.2	3.1; 3.3
Household head is male (%)	98.9	98.1; 99.4
Household head formal education		
No education (%)	33.9	31.0; 36.9
Grade 1-8 (%)	45.5	42.4; 48.6
Grade 9-12 (%)	14.0	11.9; 16.3
Technical and Vocational Education (%)	0.60	0.30; 1.40
College/University (%)	6.10	4.70; 7.70
Mother/Caregiver Characteristics		
Mother as main caregiver (%)	99.2	98.5; 99.6
Grandparent help in infant care (%)	35.8	32.9; 38.7
Mother/Caregiver age (years)	25.9	25.6; 26.2
Mother/Caregiver is married (%)	98.8	98.0; 99.3
Mother/Caregiver age at marriage (years)	17.5	17.3; 17.6
Time married (years)	8.45	8.10; 8.79
Mother/Caregiver formal education		
No education (%)	40.9	37.9; 43.9
Grade 1-8 (%)	45.8	42.8; 48.9
Grade 9-12 (%)	9.16	7.55; 11.1
Technical and Vocational Education (%)	1.25	0.73; 2.15
College/University (%)	2.89	2.03; 4.11
Mother/Caregiver religion		

	mean or %	95% CI
Muslim (%)	94.2	92.6; 95.5
Orthodox Christian (%)	4.60	3.50; 6.10
Protestant (%)	1.10	0.60; 1.90
Refused to answer (%)	0.10	0.00; 0.70
Infant Characteristics		
Infant's age (weeks)	13.4	13.0; 13.8
Infant is male (%)	55.4	52.3; 58.4
Infant is singleton (%)	98.6	97.6; 99.1
Siblings aged <18 years	2.2	2.0; 2.3
Infant's birth order		
1st (%)	25.0	22.4; 27.7
2nd (%)	20.6	18.3; 23.2
3rd (%)	15.0	13.0; 17.4
4th (%)	12.3	10.4; 14.4
5th (%)	11.3	9.49; 13.4
6th+ (%)	15.8	13.7; 18.2
Recent death of sibling (%)	14.9	12.8; 17.2
Infant feeding		
Infant ever breastfed (%)	98.9	98.1; 99.4
Infant received breastmilk as first food (%)	96.1	94.7; 97.1
Infant was breastfed in the past 24 hrs (%)	94.1	92.5; 95.4
Breastfeeding frequency in past 24 hrs (times)	10.4	10.3; 10.6
Infant exclusively breastfed (%)	48.9	45.9; 51.9
Infant was bottle-fed (%)	15.2	13.2; 17.6
Infant fed any solid, semi-solid or soft foods (%)	2.03	1.32; 3.09
Infant anthropometry		
Bilateral pitting oedema (%)	0.68	0.32; 1.41
Weight (kg)	5.62	5.55; 5.70
Length (cm)	59.6	59.4; 59.9
Mid-upper arm circumference (cm)	12.4	12.4; 12.5
Triceps skinfold (mm)	7.7	7.6; 7.8
Subscapular skinfold (mm)	6.8	6.7; 6.9
Head circumference (cm)	40.4	40.2; 40.5
Knee-to-heel length (mm)	148	147; 149
Weight-for-age z-score	-0.65	-0.72; -0.57
Length-for-age z-score	-0.35	-0.43; -0.26
Weight-for-length z-score	-0.47	-0.55; -0.40


In Figure 2 we showed the age-smoothed proportion of infants u6m that were exclusively breastfed, were fed different liquids, or were bottle-fed. Median duration of exclusive breastfeeding was 10.6 weeks; and by the age of 16.7 weeks, half of the infants u6m were being given water.

251

252

253

254



255

Figure 2. Age-smoothed proportion of exclusively breastfed infants aged under six months. The figure also presents age-smoothed proportions of bottle feeding and the consumption of water, milk and other liquids (other than water, milk, juice, broth, runny porridge, or yogurt).

256

257

258

3.4. Prevalence of anthropometric deficit

259

Table 2 presents the prevalence of anthropometric deficits in infants u6m, overall and by age categories. We observed that over one in five infants u6m had some form of anthropometric deficit, as indexed by CIAF, and over 4% had a severe anthropometric deficit, as indexed by CISAF. Multiple anthropometric deficits (i.e. wasted and underweight; stunted and underweight; and wasted, stunted and underweight) affected 10.7% of infants u6m (95% CI: 8.93; 12.7). We did not observe any marked age-dependent variability in the prevalence of infants u6m wasted, stunted, underweight, CIAF or CISAF.

260

261

262

263

264

265

266

3.5. Prevalence of low MUAC

267

Table 3 shows the prevalence of low MUAC using different thresholds. According to the threshold used, we found that overall low MUAC prevalence ranged from 6.71% to 19.1%. We also observed large age-dependent variations: low MUAC was very common in the youngest age groups and decreased sharply in older infants.

268

269

270

271

Table 2. Proportion of infants aged under six months with different indicators of anthropometric deficit.

272

Anthropometric indicator	All		0-5 weeks		6-10 weeks		11-15 weeks		16-20 weeks		20-25 weeks	
	n = 1,030	%	n = 93	%	n = 268	%	n = 285	%	n = 243	%	n = 141	%
Wasted	10.9	9.11; 12.9	12.9	7.47; 21.4	8.21	5.46; 12.2	10.2	7.16; 14.3	12.8	9.11; 17.6	12.8	8.19; 19.4
Severely wasted	2.43	1.64; 3.57	--	--	--	--	--	--	--	--	--	--
Stunted	9.81	8.13; 11.8	7.53	3.63; 15.0	9.70	6.69; 13.9	9.12	6.28; 13.1	10.7	7.38; 15.3	11.4	7.07; 17.7
Severely stunted	3.47	2.51; 4.78	--	--	--	--	--	--	--	--	--	--
Underweight	12.9	11.0; 15.1	7.53	3.63; 15.0	13.8	10.2; 18.5	10.5	7.45; 14.7	16.5	12.3; 21.7	13.5	8.76; 20.2
Severely underweight	4.56	3.44; 6.02	--	--	--	--	--	--	--	--	--	--
CIAF	21.7	19.2; 24.3	19.4	12.5; 28.7	21.3	16.8; 26.6	20.4	16.1; 25.4	24.3	19.3; 30.1	22.0	15.9; 29.6
No CIAF	78.4	75.7; 80.8	80.7	71.3; 87.5	78.7	73.4; 83.2	79.7	74.6; 83.9	75.7	69.9; 80.7	78.0	70.4; 84.1
<i>CIAF categories</i>												
Wasted only	5.15	3.95; 6.68	--	--	--	--	--	--	--	--	--	--
Wasted and Underweight	4.47	3.36; 5.91	--	--	--	--	--	--	--	--	--	--
Wasted, Stunted and Underweight	1.26	0.73; 2.16	--	--	--	--	--	--	--	--	--	--
Stunted and Underweight	4.95	3.78; 6.46	--	--	--	--	--	--	--	--	--	--
Stunted only	3.59	2.61; 4.92	--	--	--	--	--	--	--	--	--	--
Underweight only	2.23	1.49; 3.34	--	--	--	--	--	--	--	--	--	--
CISAF	4.27	3.19; 5.69	1.08	0.15; 7.25	4.85	2.84; 8.18	5.61	3.47; 8.97	5.35	3.13; 9.00	0.71	0.10; 4.87

CIAF: Composite Index of Anthropometric Failure. CISAF: Composite Index of Severe Anthropometric Failure.

Underweight, stunted and wasted was defined as weight-for-age (WAZ), length-for-age (LAZ) and weight-for-length (WLZ) z-scores <-2, respectively. Severe underweight, stunted and wasted was defined as WAZ, LAZ and WLZ <-3, respectively. CIAF are all infants u6m that were either underweight, stunted or wasted. CISAF are all infants u6m that were either severely underweight, stunted or wasted.

273

274

Table 3. Proportion of infants aged under six months with low MUAC as defined by different thresholds.

275

MUAC indicator	All		0-5 weeks		6-10 weeks		11-15 weeks		16-20 weeks		20-25 weeks	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI
MUAC <10.5cm	6.71	5.33; 8.42	44.1	34.4; 54.3	4.12	2.29; 7.29	3.16	1.65; 5.96	2.47	1.11; 5.39	1.43	0.36; 5.54
MUAC <11.0cm	12.0	10.1; 14.1	65.6	55.4; 74.5	11.6	8.28; 16.0	5.26	3.20; 8.55	4.53	2.52; 7.99	3.57	1.49; 8.30
MUAC <11.0cm if <17 weeks, <11.5 thereafter	12.7	10.8; 14.9	65.6	55.4; 74.5	11.6	8.28; 16.0	5.26	3.20; 8.55	5.76	3.44; 9.50	7.14	3.88; 12.8
MUAC <11.0cm if <13 weeks, <11.5 thereafter	13.4	11.5; 15.7	65.6	55.4; 74.5	11.6	8.28; 16.0	7.72	5.13; 11.5	5.76	3.44; 9.50	7.14	3.88; 12.8
MUAC <11.0cm if <7 weeks, <11.5 thereafter	16.5	14.4; 18.9	65.6	55.4; 74.5	20.6	16.2; 25.9	10.5	7.45; 14.7	5.76	3.44; 9.50	7.14	3.88; 12.8
MUAC <11.0cm if <6 weeks, <11.5 thereafter	17.6	15.4; 20.1	65.6	55.4; 74.5	24.7	19.9; 30.3	10.5	7.45; 14.7	5.76	3.44; 9.50	7.14	3.88; 12.8
MUAC <11.5cm	19.1	16.8; 21.6	81.7	72.5; 88.3	24.7	19.9; 30.3	10.5	7.45; 14.7	5.76	3.44; 9.50	7.14	3.88; 12.8

MUAC: Mid-upper arm circumference.

276

277

3.6. Overlap of low MUAC and underweight with different anthropometric deficits 278

In Table 4 we show the proportion of underweight, wasted, stunted, CIAF and CISAF infants that would be identified by different MUAC thresholds or by underweight if that were used as a sole criterion for identification of small and nutritionally at-risk infants u6m. Higher MUAC thresholds identify greater proportions of the different definitions of anthropometric deficit. However, they also identified greater proportions of infants u6m that had no anthropometric deficits, as indexed by CIAF. On comparing different MUAC thresholds and underweight for identifying CIAF and CISAF, we observed that underweight identified larger proportions of CIAF and CISAF infants u6m than any MUAC threshold. A visual representation of the overlap, in two different age groups, between the wasted, stunted and underweight and low MUAC indicators, using a MUAC threshold <11.0cm if aged <6 weeks and <11.5 thereafter, is shown in Figure 3.

Supplementary Materials Table S2 and S3 also cross-tabulate MUAC data with CIAF and CISAF categories. The overlap between different MUAC thresholds and CIAF is again observed to be strongly influenced by the infant's age and how inclusive the MUAC threshold is. For the youngest infants (aged 0-5weeks) the least inclusive MUAC threshold we examined (<10.5cm) overlaps with 61% of other forms of CIAF, but also with 40% of infants with no other anthropometric deficits. The most inclusive MUAC threshold (<11.5cm) overlaps with most (89%) but not all of other forms of CIAF but also with 80% of infants with no other anthropometric deficits. As the age of the infant increases, this proportion of MUAC-identified infants u6m with no CIAF is reduced (to 0% at ages 11-15 weeks) but overlap with CIAF also decreases. Underweight has a more consistent pattern of overlapping with 39% of all forms of CIAF at age 0-5weeks; with 65% at ages 6-10weeks; with 52% at ages 11-15weeks; with 68% at 16-20weeks and 61% at ages 21-25 weeks.

305

Table 4. Overlap of different anthropometric deficits with different low MUAC thresholds and Underweight

306

Indicator	Underweight			Stunted			Wasted			CIAF			CISAF			No CIAF	
	%	95% CI	n = 133	%	95% CI	n = 101	%	95% CI	n = 112	%	95% CI	n = 223	%	95% CI	n = 44	%	95% CI
MUAC <10.5cm	24.1	17.6; 32.1	22.8	15.6; 32.0	17.0	11.1; 25.1	17.0	12.7; 22.6	36.4	23.6; 51.4	3.85	2.72; 5.43					
MUAC <11.0cm	44.4	36.2; 52.9	38.6	29.6; 48.4	27.7	20.2; 36.7	30.5	24.8; 36.9	43.2	29.5; 58.0	6.83	5.28; 8.80					
MUAC <11.0cm if <17 weeks, <11.5 thereafter	50.4	41.9; 58.8	42.6	33.3; 52.4	33.9	25.8; 43.2	34.1	28.2; 40.6	43.2	29.5; 58.0	6.83	5.28; 8.80					
MUAC <11.0cm if <13 weeks, <11.5 thereafter	54.1	45.6; 62.4	43.6	34.2; 53.4	36.6	28.2; 45.9	36.3	30.3; 42.9	45.5	31.5; 60.2	7.08	5.50; 9.07					
MUAC <11.0cm if <7 weeks, <11.5 thereafter	60.9	52.4; 68.8	47.5	38.0; 57.3	43.8	34.9; 53.1	42.6	36.3; 49.2	45.5	31.5; 60.2	9.32	7.49; 11.5					
MUAC <11.0cm if <6 weeks, <11.5 thereafter	60.9	52.4; 68.8	47.5	38.0; 57.3	43.8	34.9; 53.1	42.6	36.3; 49.2	45.5	31.5; 60.2	10.7	8.73; 13.0					
MUAC <11.5cm	61.7	53.1; 69.5	47.5	38.0; 57.3	46.4	37.4; 55.7	44.0	37.6; 50.5	45.5	31.5; 60.2	12.2	10.1; 14.6					
Underweight	--	--	63.4	53.6; 72.2	52.7	43.4; 61.7	59.6	53.1; 65.9	65.9	50.9; 78.3	--	--					

CIAF: Composite Index of Anthropometric Failure. CISAF: Composite Index of Severe Anthropometric Failure.

Underweight, stunted and wasted was defined as weight-for-age (WAZ), length-for-age (LAZ) and weight-for-length (WLZ) z-scores <-2, respectively. Severe underweight, stunted and wasted was defined as WAZ, LAZ and WLZ <-3, respectively. CIAF are all infants u6m that were either underweight, stunted or wasted. CISAF are all infants u6m that were either severely underweight, stunted or wasted.

307



Figure 3. A Venn-diagram showing the overlap of different malnutrition indicators in two age groups.

Low MUAC was defined as MUAC <11.0cm if <6 weeks, <11.5 thereafter. Underweight, stunted and wasted was defined as weight-for-age, length-for-age and weight-for-length z-scores <-2, respectively. MUAC: Mid-upper arm circumference.

308

309

310

311

312

313

314

315

316

4. Discussion	317
<i>4.1. Summary of results</i>	318
To our knowledge, ours is the first study on infants u6m – not just in Ethiopia, but in any setting, to estimate the malnutrition burden as indexed by CIAF and its subcategories. The major finding was of a common problem: over 20% of infants u6m attending clinics had some form of anthropometric deficit (CIAF), of which a fifth (4%) were severe (CISAF) and over half (11%) were multiple anthropometric deficits with combinations of wasted, stunted or underweight.	319 320 321 322 323 324
Estimating malnutrition burden using the simplest index – MUAC – resulted in a wide prevalence range depending on the infants' age and the threshold used to define low MUAC. Whilst MUAC-defined prevalence overlapped with CIAF-defined prevalence, the extent of overlap varied markedly and was lowest in the youngest age group where many infants with low MUAC had no other CIAF-defined anthropometric deficit. In contrast to marked age-related differences for MUAC-based estimates, wasted-, stunted-, underweight-, CIAF-, and CISAF-based estimates were similar across the different age categories. There were no consistent upwards or downwards prevalence trends across the age categories. Underweight as the sole case definition criterion overlapped with at least half of the infants u6m with other forms of anthropometric deficit and identified larger proportions of CIAF and CISAF infants than any MUAC threshold.	325 326 327 328 329 330 331 332 333 334 335
<i>4.2. Burden of malnutrition and programme implications</i>	336
Even though our study used a health centre-based rather than population-based sample, our observed wasted and underweight prevalence are comparable with recent national estimates from the 2019 Ethiopian Mini DHS. However, our infants u6m sample were markedly less stunted than the national prevalence (9.8% vs 17.1%) [20]. Our data on the overlap between different types of anthropometric deficit was also consistent with that from older children where this has been observed; and where there is current focus on the overlap between wasting and stunting since this can greatly increase mortality risk [27]. In older children, there is also increased focus on underweight as a simple single measure and way of capturing this overlap without having to assess height-based indices [27].	337 338 339 340 341 342 343 344 345 346
Current Ethiopian guidelines (and most other national guidelines which are also based on the 2013 update to the WHO guideline for management of severe malnutrition) for admitting infants u6m to malnutrition treatment programmes focus only on those who are severely wasted, as indicated by a WLZ <-3 [21]. Our data show that this focus misses many infants u6m with other forms of anthropometric deficit. Widening admission criteria would certainly have programme caseload and capacity implications. Our results showed that the number of admissions would almost double (from a 2.4% to 4.3%) if CISAF was to become the criterion for admission, or it would increase by ten-fold if the criteria was any form of anthropometric deficit as denoted by CIAF. Key questions for future research are what impact there would be on programme outcomes, notably mortality and morbidity, whether the caseload expansion would be justified by the benefits, and whether it would be cost-effective. A closely related question is what the benefits of a shift to outpatient-based care for at-risk infants u6m are. This alone would markedly improve programme capacity over current inpatient-only models of care.	347 348 349 350 351 352 353 354 355 356 357 358 359 360
<i>4.3. The utility of simple measurements: MUAC and WAZ</i>	361
A small but growing number of studies have explored the utility of different anthropometric criteria to identify malnutrition in infants u6m [28–36]. The best of these look at the association between anthropometric measures and subsequent mortality. This conceptual approach recognises that what matters is clinical outcome rather than body size; that there is no one anthropometric measure considered the 'gold standard' for identifying malnutrition; that all anthropometric measures are only proxy measures of nutritional status, each with advantages and disadvantages and differing in sensitivity, specificity, and positive and negative predictive value when identifying infants u6m	362 363 364 365 366 367 368 369

at-risk of adverse outcomes, notably mortality and morbidity [17]. Despite differing designs and contexts, most of these studies on infants u6m agree that WLZ is poor at identifying high-risk infants and that WAZ or MUAC are better and more practical. WAZ is already widely measured in growth monitoring programmes so would be particularly easy to adopt. Whilst our cross-sectional study design could not assess the prognostic value of these two indices in our population, we have shown the potential caseload implications.

The extent to which low MUAC, especially in the first six weeks of life, overlaps with LBW is a major unknown. This overlap is plausible and matters because LBW is a well-established risk factor for both short- and long-term mortality and morbidity [37], even when other anthropometric indicators are within normal range [29]. If a large overlap between low MUAC and LBW in the first weeks of life is present, it would strongly support the use of low MUAC to identify infants u6m for enrolment to treatment/support programmes at this age, even when the overlap with other anthropometric deficits is poor. A recent study of newborns in Ethiopia found a MUAC ≤ 9.8 cm, measured within 24 hours of life, to be a useful diagnostic tool for LBW [38]. Conversely, if there is a poor overlap between low MUAC and LBW then there might be less value in using low MUAC, as these infants might have smaller arms because they are young and hence small.

Either way, future research is needed to assess the independent role of MUAC in identifying infants u6m at high risk of mortality/morbidity. MUAC-for-age tables are available and might be used to improve the predictive value of low MUAC – but this might work against the practical and programmatic advantage of MUAC, namely simplicity and speed of assessment.

Much current research on anthropometric deficits as a risk factor has focused on how well it identifies individuals at-risk of short-term morbidity or mortality [39]. However, less attention has been given to assess how well it predicts a longer-term risk of impaired physical or cognitive development. Evidence from a recent systematic review shows that adverse nutrition in infancy and childhood is associated with long-term adult non-communicable disease [6]. Future research should explore how both MUAC and the various subcategories of CIAF predict these longer-term risk and possible underlying mechanisms and pathways. The role of body composition and body proportions, both indicators of development, could help explain what lies behind these simple anthropometric measurements. Past research has shown that MUAC and WLZ have different associations with body composition, where WLZ appears to have similar associations with lean and fat mass, whilst it also has a small but at times negligible association with length. MUAC on the other hand, appears to have a stronger association with fat mass than with lean mass, but also has a strong association with length; that is each index MUAC and WLZ appears to select infants u6m for different phenotypes [40].

4.4. Implications for research and programming

Our data has immediate implications for programming and research using a new “Management of small and nutritionally At-risk Mothers and Infants (MAMI)” Care Pathway, just launched in May 2021 [1]. Building on past experiences [28], the MAMI Care Pathway emphasises the potential of MUAC as an independent programme enrolment criterion. Our results highlight that more data is needed to understand the prognostic utility of MUAC, especially in the youngest infants u6m where overlap with other forms of anthropometric deficits are the least. Balancing the needs for simplicity and recognising the rapid increase in MUAC, especially in the first month of life, our own planned research in Ethiopia will now include both single indicators, a WAZ <-2 and a MUAC <11.0 cm if aged <6 weeks and <11.5 thereafter; where of the total sample of included infants u6m, about 26% of them will be identified by both indicators.

4.5. Strengths and limitations

We recognise the limitations of our work. Most importantly, our data from health centres might not be representative of the wider infants u6m population. However, it is still important since recruitment into future programmes is likely to occur at health centre level and hence our data matters for planning of such programmes. Existing programmes such as immunization have very high coverage and thus our strategy of screening at such visits would be both efficient and would identify many high risk infants u6m early. Related to this, our health centres selection was not randomised and did not seek to be representative of all the health centres in the Oromia region. We selected based on ease of access and on the number of infants u6m who receive care, informed by register data. Whilst this limits our ability to generalise the absolute prevalence results to the wider region/country, there is no reason to believe that the overlaps between indicators and the age-related variations should be very different elsewhere. Nonetheless, this should be examined in future work.

As already noted, missing birthweight data was an important limitation. Whilst we tried to collect birthweight data, this was unavailable for most infants u6m. As a result, we were unable to disentangle how much of the anthropometric deficit we observed could be explained by poor foetal growth, and which of the different indicators of anthropometric deficits are best at identifying small infants that were LBW.

In this analysis, we also did not analyse the reasons underlying or associated with anthropometric deficit. This should be the focus for future work. A particularly important future question is how much anthropometric deficit / growth failure is reversible with nutritional and other interventions and what are the clinical and functional benefits (e.g. on mortality, morbidity and child development) of these interventions. These questions we seek to address in our own planned RCT, and we hope this paper will inspire others will do similar in other settings.

Finally, we did not stratify our analysis by the infant's sex and as such, we are not able to understand whether our findings differ by sex. Data from children has shown that boys are often more underweight, wasted and stunted whereas there is a sex bias towards girls when identifying malnutrition using MUAC [41,42]. Nonetheless, none of these limitations are likely to invalidate the findings of our study, rather it forms the basis of future work.

Our study has also strengths. In our study, we sampled infants u6m attending health centres for any reason. Most studies evaluating malnutrition burden in infants u6m focused on community representative samples [28,31–33], on infants hospitalised and receiving in-patient care [30,34–36], or they have followed-up a birth cohort [29]. Our study was a facility-based survey collecting data on infants attending health centres for any reason, which provides better understanding of the potential for care provision to the population covered by these health centres. Future enrolment into programmes for small and nutritionally at-risk infants u6m is likely to occur at health centres so it is vital to know numbers at these facilities.

We took on the recent call for using aggregate measures to quantify malnutrition and quantify the burden of malnutrition as any manifestation of anthropometric deficit as indexed by CIAF [17]. Published research on malnutrition in infants u6m has focused strongly on severe wasting as denoted by a WLZ <-3 [29,30,33,36], but our data adds to growing calls for more inclusive case definitions for identifying infants at risk of malnutrition, mortality or impaired development. As mentioned above, we are currently developing an intervention trial that will test the benefit of a more inclusive approach and we hope that our data will encourage others to do likewise. The analysis in the present study provides the first step towards quantifying the burden of CIAF and of MUAC in infants u6m and the degree of convergence between indicators and across age sub-groups.

5. Conclusions

422
423
424
425
426
427
428
429
430
431
432
433
434

435
436
437
438
439

440
441
442
443
444
445
446

447
448
449
450
451
452

453
454
455
456
457
458
459
460
461

462
463
464
465
466
467
468
469
470
471
472

473

Single and multiple anthropometric deficits are prevalent in Ethiopian infants u6m attending health centres. To identify any form of anthropometric deficits, as classified by wasted, stunted, underweight, CIAF or CISAF, WAZ appears to perform better than MUAC, whilst both are good in identifying infants u6m with multiple anthropometric deficits. Further research is needed to understand which criteria or combination of criteria are best for future programmes managing small and nutritionally at-risk infants u6m; and to understand the associated functional and clinical outcomes, notably short term-risks of mortality and morbidity. 474
 475
 476
 477
 478
 479
 480
 481

Supplementary Materials: The following are available online at www.mdpi.com/xxx/xxx, Table S1: Characteristics of the 1,037 surveyed infants under six months of age, by study site, Table S2: Proportion of infants aged under six months with and without CIAF overlapping with low MUAC or underweight, Table S3: Proportion of CIAF categories overlapping with low MUAC. 482
 483
 484
 485

Author Contributions: Conceptualization, C.S.G.-E., E.A., E.B., H.B., M.A., M.B., M.K. and M.M.; methodology, C.S.G.-E., E.A., E.B., M.A., M.B. and M.K.; formal analysis, C.S.G.-E.; investigation, C.S.G.-E., E.B., M.A., M.B. and R.R.; data curation, C.S.G.-E. and E.B.; writing—original draft preparation, C.S.G.-E.; writing—review and editing, All authors; supervision, M.B. and N.A.; funding acquisition, A.A., H.B., M.A., M.B., M.K., M.M. and T.G. All authors have read and agreed to the published version of the manuscript. 486
 487
 488
 489
 490
 491

Funding: This study was generously supported by the Eleanor Crook Foundation grant to the LSHTM (grant reference: EPPHZR37). 492
 493

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki, and all procedures involving research study participants were approved by the Institutional Review Board of Jimma University, Ethiopia (reference: IHRPGD/478/2020) and the Research Ethics Committee at the London School of Hygiene and Tropical Medicine, United Kingdom (reference: 18022). 494
 495
 496
 497
 498

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study. 499
 500

Data Availability Statement: The data presented in this study are available on request from the corresponding author. 501
 502

Acknowledgments: We are grateful to all study participants for kindly agreeing to take part in this study. We are thankful to Belay Kebede and Abdulhakim Ali for field coordination during data collection. 503
 504
 505

Conflicts of Interest: The authors declare no conflict of interest. The funder had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results. 506
 507
 508

Appendix A 509

Table A1. Pre-set maximum allowable differences of various anthropometric measurements. 510

Anthropometric measurement (units)	Maximum allowable difference
Weight (g)	100
Length (cm)	0.7
MUAC (mm)	5
Head circumference (cm)	0.5
Knee-heel length (mm)	6
Subscapular skinfold thickness (mm)	2
Triceps skinfold thickness (mm)	2

511

1. MAMI (Management of small & nutritionally At-risk Infants under six months & their Mothers): Project summary. Available online: <http://www.ennonline.net/ourwork/research/mami> (accessed on May 14, 2021). 514
515
2. Christian, P.; Lee, S.E.; Donahue Angel, M.; Adair, L.S.; Arifeen, S.E.; Ashorn, P.; Barros, F.C.; Fall, C.H.D.; Fawzi, W.W.; Hao, W.; et al. Risk of childhood undernutrition related to small-for-gestational age and preterm birth in low- and middle-income countries. *Int. J. Epidemiol.* 2013, 42, 1340–1355, doi:10.1093/ije/dyt109. 516
517
518
3. Kerac, M.; Blencowe, H.; Grijalva-Eternod, C.; McGrath, M.; Shoham, J.; Cole, T.J.; Seal, A. Prevalence of wasting among under 6-month-old infants in developing countries and implications of new case definitions using WHO growth standards: a secondary data analysis. *Arch. Dis. Child.* 2011, 96, 1008–1013, doi:10.1136/adc.2010.191882. 519
520
521
4. Blencowe, H.; Krasevec, J.; de Onis, M.; Black, R.E.; An, X.; Stevens, G.A.; Borghi, E.; Hayashi, C.; Estevez, D.; Cegolon, L.; et al. National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *Lancet Glob. Heal.* 2019, 7, e849–e860, doi:10.1016/S2214-109X(18)30565-5. 522
523
524
5. Kerac, M.; Mwangome, M.; McGrath, M.; Haider, R.; Berkley, J.A. Management of Acute Malnutrition in Infants Aged under 6 Months (MAMI): Current Issues and Future Directions in Policy and Research. *Food Nutr. Bull.* 2015, 36, S30–S34, doi:10.1177/15648265150361S105. 525
526
527
6. Grey, K.; Gonzales, G.B.; Abera, M.; Lelijveld, N.; Thompson, D.; Berhane, M.; Abdissa, A.; Girma, T.; Kerac, M. Severe malnutrition or famine exposure in childhood and cardiometabolic non-communicable disease later in life: a systematic review. *BMJ Glob. Heal.* 2021, 6, e003161, doi:10.1136/bmjgh-2020-003161. 528
529
530
7. Collins, S.; Dent, N.; Binns, P.; Bahwere, P.; Sadler, K.; Hallam, A. Management of severe acute malnutrition in children. *Lancet* 2006, 368, 1992–2000, doi:10.1016/S0140-6736(06)69443-9. 531
532
8. Bhutta, Z.A.; Berkley, J.A.; Bandsma, R.H.J.; Kerac, M.; Trehan, I.; Briend, A. Severe childhood malnutrition. *Nat. Rev. Dis. Prim.* 2017, 3, 17067, doi:10.1038/nrdp.2017.67. 533
534
9. Every Woman Every Child Global Strategy for Women's, Children's and Adolescents Health 2016-2030 Available online: <https://www.everywomaneverychild.org/global-strategy/> (accessed on May 14, 2021). 535
536
10. Grijalva-Eternod, C.S.; Kerac, M.; McGrath, M.; Wilkinson, C.; Hirsch, J.C.; Delchevalerie, P.; Seal, A.J. Admission profile and discharge outcomes for infants aged less than 6 months admitted to inpatient therapeutic care in 10 countries. A secondary data analysis. *Matern. Child Nutr.* 2017, 13, e12345, doi:10.1111/mcn.12345. 537
538
539
11. WHO Guideline: Updates on the management of severe acute malnutrition in infants and children. Available online: <https://www.who.int/publications/i/item/9789241506328> (accessed on May 14, 2021). 540
541
12. Mwangome, M.K.; Berkley, J.A. The reliability of weight-for-length/height Z scores in children. *Matern. Child Nutr.* 2014, 10, 474–480, doi:10.1111/mcn.12124. 542
543
13. WHO Multicentre Growth Reference Study Group WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length, weight- for-height and body mass index-for-age: methods and development.; Geneva, 2006; 544
545
14. Kerac, M.; Frison, S.; Connell, N.; Page, B.; McGrath, M. Informing the management of acute malnutrition in infants aged under 6 months (MAMI): risk factor analysis using nationally-representative demographic & health survey secondary data. *PeerJ* 2019, 6, e5848, doi:10.7717/peerj.5848. 546
547
548
15. Angood, C.; McGrath, M.; Mehta, S.; Mwangome, M.; Lung'aho, M.; Roberfroid, D.; Perry, A.; Wilkinson, C.; Israel, A.-D.; Bizouerne, C.; et al. Research Priorities to Improve the Management of Acute Malnutrition in Infants Aged Less Than Six Months (MAMI). *PLOS Med.* 2015, 12, e1001812, doi:10.1371/journal.pmed.1001812. 549
550
551
16. Lelijveld, N.; Kerac, M.; McGrath, M.; Mwangome, M.; Berkley, J.A. A review of methods to detect cases of severely malnourished infants less than 6 months for their admission into therapeutic care. Available online: <https://www.ennonline.net/mamicasedetectionreview> (accessed on May 14, 2021). 552
553
554
17. Kerac, M.; McGrath, M.; Connell, N.; Kompara, C.; Moore, W.H.; Bailey, J.; Bandsma, R.; Berkley, J.A.; Briend, A.; Collins, S.; et al. 'Severe malnutrition': thinking deeply, communicating simply. *BMJ Glob. Heal.* 2020, 5, e003023, doi:10.1136/bmjgh-2020-003023. 555
556
557
18. Svedberg, P. How Many People Are Malnourished? *Annu. Rev. Nutr.* 2011, 31, 263–283, doi:10.1146/annurev-nutr-081810-160805. 558
559
19. Kuiti, B.K.; Bose, K. The Concept of Composite Index of Anthropometric Failure (CIAF): Revisited and Revised. *Anthropol. - Open J.* 2018, 3, 32–35, doi:10.17140/ANTPOJ-3-118. 560
561
20. Ethiopian Public Health Institute (EPHI) [Ethiopia]; ICF International. Ethiopia Mini Demographic and Health Survey 2019: Key Indicators; 2019; 562
563
21. Federal Ministry of Health National Guideline for the Management of Acute Malnutrition in Ethiopia. 2019. 564
22. WHO Indicators for assessing infant and young child feeding practices part 2: measurement; Geneva, 2010; 565
23. de Onis, M.; Onyango, A.W.; Van den Broeck, J.; Chumlea, W.C.; Martorell, R. Measurement and Standardization Protocols for Anthropometry Used in the Construction of a New International Growth Reference. *Food Nutr. Bull.* 2004, 25, S27–S36, doi:10.1177/15648265040251S105. 566
567
568
24. Vidmar, S.I.; Cole, T.J.; Pan, H. Standardizing anthropometric measures in children and adolescents with functions for egen: Update. *Stata J.* 2013, 13, 366–378. 569
570
25. Crowe, S.; Seal, A.; Grijalva-Eternod, C.; Kerac, M. Effect of nutrition survey 'cleaning criteria' on estimates of malnutrition prevalence and disease burden: secondary data analysis. *PeerJ* 2014, 2, e380, doi:10.7717/peerj.380. 571
572

26. Vollmer, S.; Harttgen, K.; Kupka, R.; Subramanian, S. V. Levels and trends of childhood undernutrition by wealth and education according to a Composite Index of Anthropometric Failure: evidence from 146 Demographic and Health Surveys from 39 countries. *BMJ Glob. Heal.* 2017, 2, e000206, doi:10.1136/bmigh-2016-000206. 573

27. Myatt, M.; Khara, T.; Schoenbuchner, S.; Pietzsch, S.; Dolan, C.; Lelijveld, N.; Briand, A. Children who are both wasted and stunted are also underweight and have a high risk of death: a descriptive epidemiology of multiple anthropometric deficits using data from 51 countries. *Arch. Public Heal.* 2018, 76, 28, doi:10.1186/s13690-018-0277-1. 575

28. Moore, L.; O'Mahony, S.; Shevlin, M.; Hyland, P.; Barthorp, H.; Vallières, F. Towards identifying malnutrition among infants under 6 months: a mixed-methods study of South-Sudanese refugees in Ethiopia. *Public Health Nutr.* 2020, 1–10, doi:10.1017/S1368980020004048. 578

29. Mwangome, M.; Ngari, M.; Bwahere, P.; Kabore, P.; McGrath, M.; Kerac, M.; Berkley, J.A. Anthropometry at birth and at age of routine vaccination to predict mortality in the first year of life: A birth cohort study in BurkinaFaso. *PLoS One* 2019, 14, e0213523, doi:10.1371/journal.pone.0213523. 582

30. Mwangome, M.; Ngari, M.; Fegan, G.; Mturi, N.; Shebe, M.; Bauni, E.; Berkley, J.A. Diagnostic criteria for severe acute malnutrition among infants aged under 6 mo. *Am. J. Clin. Nutr.* 2017, 105, ajcn149815, doi:10.3945/ajcn.116.149815. 584

31. Mwangome, M.K.; Fegan, G.; Fulford, T.; Prentice, A.M.; Berkley, J.A. Mid-upper arm circumference at age of routine infant vaccination to identify infants at elevated risk of death: a retrospective cohort study in the Gambia. *Bull. World Health Organ.* 2012, 90, 887–894, doi:10.2471/BLT.12.109009. 587

32. Rasmussen, J.; Andersen, A.; Fisker, A.B.; Ravn, H.; Sodemann, M.; Rodrigues, A.; Benn, C.S.; Aaby, P. Mid-upper-arm-circumference and mid-upper-arm circumference z-score: the best predictor of mortality? *Eur. J. Clin. Nutr.* 2012, 66, 998–1003, doi:10.1038/ejcn.2012.95. 590

33. Chowdhury, R.; Nitika; Choudhary, T.S.; Taneja, S.; Martines, J.; Bhandari, N.; Bahl, R. Diagnostic measures for severe acute malnutrition in Indian infants under 6 months of age: a secondary data analysis. *BMC Pediatr.* 2021, 21, 158, doi:10.1186/s12887-021-02629-9. 594

34. Ezeofor, I.O.; Garcia, A.L.; Wright, C.M. Criteria for undernutrition screening in hospitalised infants under 6 months: a diagnostic accuracy study in a resource-poor setting. *Arch. Dis. Child.* 2020, 105, 524–529, doi:10.1136/archdischild-2019-318313. 597

35. Kumar, P.; Meiyappan, Y.; Rogers, E.; Daniel, A.; Sinha, R.; Basu, S.; Kumar, V.; De Wagt, A. Outcomes of Hospitalized Infants Aged One to Six Months in Relation to Different Anthropometric Indices – An Observational Cohort Study. *Indian J. Pediatr.* 2020, 87, 699–705, doi:10.1007/s12098-020-03236-9. 599

36. Jimma, B.R.; Hassen, H.Y.; Getnet, Y.; Bahwere, P.; Gebreyesus, S.H. Diagnostic performance of midupper arm circumference for detecting severe wasting among infants aged 1–6 months in Ethiopia. *Am. J. Clin. Nutr.* 2021, 113, 55–62, doi:10.1093/ajcn/nqaa294. 602

37. Khan, A.M.; Carducci, B.; Bhutta, Z.A. Low Birth Weight and Small for Gestational Age in the Context of 1,000 Days. In *The Biology of the First 1,000 Days*; Karakochuk, C.D., Whitfield, K.C., Green, T.J., Kraemer, K., Eds.; CRC Press, 2017; pp. 171–187 ISBN 9781315152950. 605

38. Gidi, N.W.; Berhane, M.; Girma, T.; Abdissa, A.; Lim, R.; Lee, K.; Nguyen, C.; Russell, F. Anthropometric measures that identify premature and low birth weight newborns in Ethiopia: a cross-sectional study with community follow-up. *Arch. Dis. Child.* 2019, archdischild-2019-317490, doi:10.1136/archdischild-2019-317490. 608

39. Randev, S. Malnutrition in Infants under 6 months: Is it Time to Change Recommendations? *Indian J. Pediatr.* 2020, 87, 684–685, doi:10.1007/s12098-020-03439-0. 611

40. Grijalva-Eternod, C.S.; Wells, J.C.K.; Girma, T.; Kæstel, P.; Admassu, B.; Friis, H.; Andersen, G.S. Midupper arm circumference and weight-for-length z scores have different associations with body composition: evidence from a cohort of Ethiopian infants. *Am. J. Clin. Nutr.* 2015, 102, 593–599, doi:10.3945/ajcn.114.106419. 613

41. Berkley, J.; Mwangi, I.; Griffiths, K.; Ahmed, I.; Mithwani, S.; English, M.; Newton, C.; Maitland, K. Assessment of Severe Malnutrition Among Hospitalized Children in Rural Kenya. *JAMA* 2005, 294, 591, doi:10.1001/jama.294.5.591. 616

42. Thurstans, S.; Opondo, C.; Seal, A.; Wells, J.; Khara, T.; Dolan, C.; Briand, A.; Myatt, M.; Garenne, M.; Sear, R.; et al. Boys are more likely to be undernourished than girls: a systematic review and meta-analysis of sex differences in undernutrition. *BMJ Glob. Heal.* 2020, 5, e004030, doi:10.1136/bmigh-2020-004030. 618