

Review

From Jekyll to Hyde: The yeast-hyphal transition of *Candida albicans*

Eve Wai Ling Chow ¹, Li Mei Pang ² and Yue Wang ^{1,3*}

¹ Institute of Molecular and Cell Biology (IMCB), Agency for Science, Technology and Research (A*STAR), 61 Biopolis Drive, Proteos, Singapore 138673; wlechow@imcb.a-star.edu.sg

² National Dental Research Institute Singapore (NDRIS), National Dental Centre Singapore, 5 Second Hospital Ave, Singapore 168938; pang.li.mei@ndcs.com.sg

³ Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, 10 Medical Drive, Singapore 117597

* Correspondence: mcbwangy@imcb.a-star.edu.sg

Abstract: *Candida albicans* is a major fungal pathogen of humans, accounting for 15% of nosocomial infections with an estimated attributable mortality of 47%. *C. albicans* is usually a benign member of the human microbiome in healthy people. Under constant exposure to highly dynamic environmental cues in diverse host niches, *C. albicans* has successfully evolved to adapt to both commensal and pathogenic lifestyles. The ability of *C. albicans* to undergo a reversible morphological transition from yeast to filamentous forms is a well-established virulent trait. Over the past few decades, a significant amount of research has been carried out to understand the underlying regulatory mechanisms, signaling pathways, and transcription factors that govern the *C. albicans* yeast-to-hyphal transition. This review will summarize our current understanding of well-elucidated signal transduction pathways that activate *C. albicans* hyphal morphogenesis in response to various environmental cues and the cell cycle machinery involved in the subsequent regulation and maintenance of hyphal morphogenesis.

Keywords: polymorphism; hyphal morphogenesis; hyphal activation; signal transduction pathways; cell cycle regulation

1. Introduction

Candida albicans is a commensal fungus that is usually a benign member of the microflora in the gastrointestinal tract, genitourinary tract, mouth, and skin of most healthy individuals [1-4]. *C. albicans* is also an opportunistic fungal pathogen responsible for infections ranging from mild superficial infections to life-threatening candidemia [5]. The use of modern medical therapies such as broad-spectrum antibiotics, cancer chemotherapy, and solid organ transplant has led to an increase in the population vulnerable to invasive candidiasis [6,7]. *C. albicans* is a leading cause of hospital-acquired infections; in the intensive care unit (ICU), candidemia may represent up to 15% of nosocomial infections with an estimated attributable mortality of 47% [7-11].

C. albicans displays a wide range of virulence factors and fitness attributes, including its capacity for rapid evolution of resistance to commonly used antifungals (e.g., azoles, polyenes, and echinocandins) and its ability to form biofilms on medical devices, contributing to its success as a pathogen. One striking feature that allows *C. albicans* to cross the commensal-to-pathogen boundary is its ability to switch reversibly between two morphological forms, namely unicellular budding yeast, or filamentous form (hyphae and pseudohyphae), in response to various environmental cues that reflect the host environment [12-19].

Yeast, hyphal, and pseudohyphal forms of *C. albicans* are all present in tissues of human patients and animals with systemic invasive candidiasis [20,21]. Yeast cells exhibit a round-to-oval cell morphology that arises from budding and nuclear division [22]. In contrast, hyphae consist of tubular cells that remain firmly attached following cytokinesis without a constriction at the site of separation. Pseudohyphae share features resembling both yeasts and hyphae, which are branched chains of elongated yeast cells with

constrictions at the septum. Both yeast and hyphal forms have crucial and complementary roles important for infection [23]. For instance, the yeast form is required for adhesion to endothelial cells and dissemination into the bloodstream, while the hyphal form is required for tissue penetration during the early stages of infection and yielding resistance towards phagocytosis [24-29]. Hyphae-specific virulence factors such as adhesins (Hwp1, Als3, Als10, Fav2, and Pga55), host tissue degrading proteases (Sap4, Sap5, and Sap6), and cytolytic peptide toxin (Ece1), aggrandize the host cell damage during infection [22,30].

Although *C. albicans* can undergo an array of morphological transitions such as the formation of chlamydospores, gray cells, GUT (gastrointestinally induced transition) phenotype, and white/opaque cells, the yeast-to-hyphae transition appears to be the most critical virulence trait [12,15]. Mutants locked in either the hyphal or yeast form have shown diminished virulence, suggesting that the ability to switch between the two morphological forms is essential for virulence [31,32]. Recent advances in mechanistic studies have provided insights into the morphological regulation, coordination, and interplay between environmental factors and genes associated with pathogenesis. This review provides an update on the signal transduction pathways involved in activating *C. albicans* hyphal morphogenesis and how the cell cycle progression and its machinery further aid the regulation and maintenance of sustained hyphal growth.

2. Environmental cues inducing the yeast-to-hyphae transition

C. albicans has adapted to growth in the human host and can transit from yeast to hyphae under a diverse range of environmental cues, as shown in Figure 1 [16,17,19]. Depending upon the cues encountered, morphogenesis can be triggered via several pathways which activate different regulatory circuits.




Figure 1. External hyphal-inducing signals. The yeast-to-hyphae transition in *C. albicans* can be triggered by various environmental cues such as high temperature (37°C), high CO₂ concentration (~5%), pH 7, nutrition deprivation, serum, peptidoglycan, *N*-acetylglucosamine, and quorum sensing molecules from endogenous and exogenous sources.

2.1. Host niches

The human host presents one of the most favorable environments for *C. albicans* morphogenesis due to the presence of multiple inducing factors such as elevated (body) temperature (37°C), the presence of serum, elevated carbon dioxide (CO₂) levels (~5%) and low glucose content (0.1%) [33-36]. Hyphal initiation requires an increase in temperature to 37°C and the release of quorum sensing molecules (e.g., farnesol) for the temporary clearance of the major transcription repressor of hyphal morphogenesis Nrg1 [37]. Elevated temperature is also known to promote filamentation through the molecular chaperone Hsp90 and the transcription factor Sfl2 [38,39]. In combination with elevated

temperature, the serum is one of the strongest inducers of *C. albicans* hyphal morphogenesis [40]. Our group has previously demonstrated that bacterial peptidoglycans present in the host serum trigger the hyphal growth of *C. albicans* by directly activating the cyclic AMP (cAMP)-protein kinase A (PKA) signaling cascades through adenylyl cyclase Cyr1 [35,41]. Similarly, CO₂, another potent inducer of filamentous growth, is also known to activate the cAMP-PKA pathway by binding to Cyr1 [42]. In *C. albicans*, a carbonic anhydrase Nce3 is involved in CO₂ signaling and conversion of CO₂ to bicarbonate (HCO₃⁻) [33]. Especially in host niches with limited CO₂ (e.g., on the skin), the CO₂/HCO₃⁻ equilibration controlled by Nce3 is crucial for the pathogenesis of *C. albicans*. The G protein-coupled receptor Grp1 and the Gα protein Gpa2 act as the glucose-sensing network for *C. albicans* morphogenesis [43]. Low glucose concentration present in the bloodstream results in the maximal hyphal formation, while high glucose concentrations repress it [36]. The factors mentioned above have been shown to activate the fungal cAMP-PKA signaling pathway [33,44-46].

2.2. Hypoxia (low oxygen)

Hypoxia is a clinical characteristic of inflammatory conditions, representing zones of intense immune activity [47,48]. *C. albicans* can modulate the host response under hypoxia and anoxia (absence of oxygen) to evade immune responses [47]. As a commensal, *C. albicans* adapts to hypoxia condition by repressing the transcription factor of filamentous growth Efg1. Interestingly, Efg1 has a dual role in hyphal morphogenesis. Under hypoxia, it acts as a repressor at temperatures ≤35°C, while under normoxia (normal oxygen level), Efg1 is a strong inducer of hyphal formation [48]. *efg1Δ/Δ* mutants displayed hyperfilamentous growth at temperatures ≤35°C during hypoxic growth on agar surface or during embedment in agar but not during growth in liquid media [48,49]. In contrast, Ace2 is essential for hyphal morphogenesis under hypoxia while being dispensable under normoxia [50,51]. Efg1 and Ace2 share functional overlap; chromatin immunoprecipitation on microchips (ChIP) analyses revealed that hypoxic repressors (Efg1 and Bcr1) and hypoxic activators (Ace2 and Brg1) are connected in regulatory circuits in controlling hyphal morphogenesis under hypoxia conditions [48]. Additionally, Efg1 was implicated in the Cek1-mediated pathway under hypoxia at ≤35°C; low Efg1 phosphorylation levels inhibit Cek1 and Cph1, preventing hyphal morphogenesis. The low Efg1 phosphorylation levels also inhibited hyphal morphogenesis through the cAMP-PKA pathway.

2.3. pH conditions

C. albicans is constantly exposed to fluctuations in pH ranging from acidic to slightly alkaline in different human body niches such as the digestive tract, vagina, oral cavity, blood, and tissues [52]. pH sensing is mediated through Rim101, an important regulator of the yeast-to-hyphae morphological transition [53-55]. Upon activation, the transcription factor Rim101 enters the nucleus and mediates pH-dependent responses [56]. Remarkably, *C. albicans* is not only capable of sensing and adapting to environmental pH but can also modulate extracellular pH by alkalinizing its surrounding environment and auto-induces hyphal formation [57]. Furthermore, alkalinization has been shown to counter the macrophage acidification during engulfment, promoting its survival in the macrophage [58].


2.4. N-acetylglucosamine (GlcNAc)

GlcNAc is commonly found as a structural component of the mucosa of the gastrointestinal tract, bacterial cell wall peptidoglycan, and fungal cell wall chitin [59,60]. Given the ubiquitous nature of GlcNAc in host niches and microbial cells, it could potentially serve as a critical signaling molecule that regulates the switch between commensalism and pathogenicity of *C. albicans* [61]. Figure 2 depicts an update on GlcNAc

signaling pathways and their involvement in hyphal morphogenesis. Ngt1 was identified as a membrane transporter specific for GlcNAc, indicating the importance of GlcNAc in intracellular signaling [62]. However, metabolism or breakdown of GlcNAc intracellularly is not required in the *C. albicans* hyphal morphogenesis as triple deletion mutants that lack all three catabolic genes (*HXK1*, *NAG1*, and *DAC1*) can exhibit filamentous growth with the addition of exogenous GlcNAc [63]. Interestingly, genetic screens have revealed two novel transcription factors *NGS1* and *RON1*, which play essential roles in both the GlcNAc catabolism and GlcNAc-induced filamentous growth [64]. *NGS1* encodes a protein similar to the GNAT family of histone acetyltransferase Gcn5, while *RON1* encodes a protein similar to the Ndt80-like DNA-binding domain [65]. Ngs1 was discovered as a novel GlcNAc signal sensor and transducer for GlcNAc-induced transcription in *C. albicans* [65]. Ngs1 targets the promoters of GlcNAc-inducible genes constitutively via the transcription factor Rep1 [65]. Ron1 was initially thought to act as both an activator and a repressor of hyphal morphogenesis. However, *ron1Δ/Δ* mutants constructed using the CRISPR/Cas9 method did not display observable GlcNAc-induced filamentous growth [64,66]. Noteworthy, upon the addition of GlcNAc, *ndt80 ron1* double deletion mutants could overcome the hyphal defects observed in *ndt80* mutants. Collectively, it suggests that Ron1 functions as a repressor of filamentous growth in the absence of Ndt80 [66]. The GlcNAc signaling pathway was initially believed to be related to the cAMP-PKA pathway as *cyr1Δ/Δ* mutants cannot form hyphae under a broad range of conditions, including GlcNAc [67]. However, it was later discovered that the fast-growing *cyr1* pseudo revertant strains could undergo filamentous growth in GlcNAc containing medium [68]. This indicates that GlcNAc can stimulate a signaling pathway independent of the cAMP-PKA pathway that has yet to be fully elucidated. An alternative pathway involved in the GlcNAc signaling is the pH-sensing Rim101 pathway. Production of excess ammonia during GlcNAc catabolism results in an increase in extracellular pH (>5), which indirectly stimulates the hyphal induction in *C. albicans* via the Rim101 pathway [60,69].

2.5. Amino acids sensing

C. albicans can utilize amino acids as alternative carbon sources during growth in glucose-poor, amino acid-rich conditions [57]. Amino acids that can be catabolized to arginine and proline are potent inducers of hyphal morphogenesis [70-72]. *C. albicans* detects extracellular amino acids via the plasma membrane-localized SPS (Ssy1-Ptr3-Ssy5) complex, which regulates two paralogous transcription factors, Stp1 and Stp2 (Figure 2) [73-75]. In the presence of extracellular amino acids, the amino acid sensor Ssy1 (Cys1) activates amino acid permease (AAP) genes [76], while the peripherally membrane-associated Ptr3 recruits casein kinase I (CKI), which activates the endoproteolytic activity of the endoprotease Ssy5 [76,77]. Ssy5 endoproteolytically cleaves the nuclear exclusion domain of Stp1 and Stp2, facilitating their translocation to the nucleus [74]. Processed Stp1 regulates the expression of *SAP2*, which encodes the major secreted aspartyl proteinase, and *OPT1*, which encodes an oligopeptide transporter. The active Stp2 activates the expression of a subset of AAP genes [74,75]. The endoplasmic reticulum (ER) chaperone protein Csh3 is required for the proper expression and plasma membrane localization of Ssy1 and AAPs [78]. *ssy1Δ/Δ*, *ptr3Δ/Δ*, *ssy5Δ/Δ*, *csh3Δ/Δ*, and *stp2Δ/Δ* mutants fail to respond to the presence of extracellular amino acids and display impaired filamentous growth [73-76,78]. Amino acid-induced morphogenesis has recently been shown to be dependent on proline catabolism, with a strict requirement for Ras1 activity [79]. Proline catabolism in the mitochondria leads to elevated cellular ATP levels, which exceed the critical threshold of ATP needed to induce cAMP synthesis, leading to hyphal morphogenesis [71,79,80].



In addition to host environmental cues, *C. albicans* morphogenesis is also regulated by several endogenous and exogenous quorum-sensing molecules (QSM) [81-85]. Tyrosol and farnesol are well-known QSM produced by *C. albicans*, which accelerate and inhibit the yeast-to-hyphae transition, respectively [81,86,87]. *C. albicans* also produces aromatic

alcohols such as ethyl alcohol, isoamyl alcohol, 2-phenylethyl alcohol, tryptophol, 1-dodecanol, 2-dodecanol, and nerolidol that inhibit filamentation and subsequent biofilm formation [88,89]. Farnesol and 1-dodecanol were implicated in the Ras1-cAMP signaling pathway, and hyphal defects can be restored upon the addition of dibutyryl-cAMP [89]. Farnesol also inhibits filamentous growth through the negative regulators Tup1 and Nrg1 [90]. The hyphal morphogenesis of *C. albicans* can also be regulated by interaction with other microorganisms found in the host environment [91,92]. For instance, the coexistence of *C. albicans* and Gram-negative bacteria, such as *Pseudomonas aeruginosa*, *Stenotrophomonas maltophilia*, and *Burkholderia cepacia*, is commonly found as mixed infections in the lungs of cystic fibrosis (CF) patients [93]. Exogenous QSM, namely, 3-oxo-C12-homoserine lactone and phenazines (pyocyanin, phenazine methosulfate, and phenazine-1-carboxylate) secreted by *P. aeruginosa*, were found to inhibit the hyphal development of *C. albicans* [82,94]. Diffusible signal factor (DSF), representing a new class of widely conserved quorum-sensing signals from Gram-negative bacteria, has been implicated in inter-kingdom signaling between *C. albicans* and bacteria [95]. DSF (*cis*-11-methyl-2-dodecenoic acid) produced by *S. maltophilia* and *Burkholderia cenocepacia* DSF or BDSF (*cis*-2-dodecenoic acid) produced by *B. cenocepacia* plays a role in the yeast-to-hyphae transition [95]. DSF released by *S. maltophilia* has been reported to interfere with two key virulence factors of *C. albicans*: the yeast-to-hyphae transition and biofilm formation [96]. Recent microarray studies revealed the involvement of repressors (Ubi4 and Sfl1) and the activator (Sfl2) of filamentous growth in BDSF regulation of hyphal morphogenesis [97]. With the addition of BDSF, elevated levels of Ubi4 and Sfl1 and degradation of Sfl2 block the yeast-to-hyphae transition.

2.7. *In vitro* conditions

Hyphal growth can also be induced using synthetic growth media such as Lee's medium (pH 7), spider medium, and mammalian tissue culture M199 under laboratory conditions [98-100]. Nitrogen starvation-induced filamentation occurs in the low nitrogen SLAD medium via ammonium permease Mep2 sensing [101,102]. Methionine, an amino acid in Lee's medium, has been reported as the main inducer of yeast-to-hyphae transition via G-protein-coupled receptor Gpr1 sensing [36]. Recently, the methionine permease Mup1 and the S-adenosylmethionine decarboxylase Spe2 were discovered to be crucial for cAMP production in response to methionine [103]. Both nitrogen and amino-acid catabolism activate hyphal morphogenesis via the cAMP-PKA pathway.

3. The cAMP-PKA pathway

The cyclic adenosine monophosphate (cAMP)-protein kinase A (PKA) pathway is highly conserved in eukaryotes and regulates many cellular processes in *C. albicans* [44,104]. This pathway plays a critical role in morphogenesis, positively regulating filamentation [105-107]. One of the well-studied regulatory targets of the cAMP-PKA pathway is the transcription factor Efg1, which stimulates the expression of numerous hyphal-specific genes through the activation of the transcription factor Ume6 (Figure 3).




Figure 3. Signal transduction pathways that govern hyphal growth in *C. albicans*. Activation of filamentous growth in *C. albicans* by various environmental cues and signal transduction pathways; the cAMP-PKA pathway, the Cek1-mediated pathway, the PKC pathway, and the embedded matrix.

The cAMP-PKA pathway is triggered either directly through the adenylyl cyclase Cyr1 or via the small GTPase Ras1, which activates Cyr1, depending upon the stimuli encountered (Figure 3) [108]. Cyr1, in direct association with Cap1 (cyclase-associated protein), drives the conversion of ATP to cAMP [46]. PKA holoenzyme is activated upon

cAMP binding to the homodimer regulatory subunit Bcy1, inducing a conformational change releasing the two catalytic subunits, Tpk1 and Tpk2, which then activate downstream target proteins or genes through phosphorylation or the binding of promoter regions to induce transcription (Figure 3) [104,109-111]. Tpk1 and Tpk2 are partially redundant. Tpk1 is required for hyphal formation on solid media, whereas Tpk2 is needed for hyphal formation in liquid media and invasive growth into solid media [110]. While the loss of either subunit does not block filamentation, loss of both Tpk1 and Tpk2 completely blocks filamentation [39,110]. cAMP levels are negatively regulated by Pde1 (low-affinity phosphodiesterase) and Pde2 (high-affinity cAMP phosphodiesterase), which increase the rate of cAMP degradation [112,113]. Loss-of-function mutations or deletion of *PDE2* increase cAMP levels, leading to constitutive activation of the pathway and hyper-filamentation [113,114]. The *pde2Δ/Δ* mutants exhibit reduced virulence due to reduced adhesion capability [115]. On the contrary, the *pde1Δ/Δ* mutants can still undergo filamentation [116]. Interestingly, *pde1 pde2* double mutants exhibit attenuated virulence as compared to *pde2* mutants [115].

The adenylyl cyclase Cyr1 is required for hyphal development and virulence but is not essential for basal growth in *C. albicans* [67]. Deletion of *CYR1* has a global impact on gene expression, resulting in many alterations in response to environmental cues [67,117]. Cyr1 contains several highly conserved functional domains, which include a Gα domain, a Ras-association (RA) domain, a leucine-rich repeat (LRR) domain, a cyclase catalytic (CYCc) domain, and a Cap1 (cyclase-associated protein 1) binding domain (CBD) [118,119].

The small GTPase Ras1, an upstream activator of Cyr1, transduces extracellular signals (serum in combination with elevated temperature or nitrogen starvation) to Cyr1 [44,101,108]. Ras1 usually exists in the cell in an inactive (GDP-bound) form, and its switch to the active form (GTP-bound) is regulated by the GTPase-activating protein (GAP) Ira2; the guanine nucleotide exchange factor (GEF) Cdc25 drive the GTP-Ras1-to-GDP-Ras1 switch (Figure 3) [120]. Active Ras1 directly interacts with Cyr1 through the RA domain, stimulating cAMP production [44,108]. Cyr1 activity depends upon the binding of Cap1 at the CBD domain and the binding of G-actin to Cap1 to form a tripartite complex, which serves to maintain the activation of the pathway [46,121]. Deletion of *CAP1* results in lowered cAMP levels and blocks in morphogenesis.

The presence of serum drives morphogenesis via Ras1 activation of the cAMP-PKA pathway. Deletion of *RAS1* impairs serum-induced filamentous growth, which can be overcome by supplementation with cAMP, and overexpression of cAMP signaling components rescued its defects [122,123]. The serum contains various active factors, such as glucose and bacterial peptidoglycan fragments, that can stimulate the pathway. The Gα domain of Cyr1 is the binding site for a heterotrimeric G-protein α subunit Gpa2, which is activated by the G-protein-coupled receptor Gpr1 in response to glucose and amino acids [34,124]. However, neither Gpr1 nor Gpa2 is required for serum-induced hyphal formation in liquid media [43]. Glucose-induced activation of cAMP synthesis appears to be mediated by Cdc25-Ras1 interaction and not Gpr1 binding of Cyr1 [36,43,45]. The LRR domain of Cyr1 recognizes and binds muramyl dipeptides (MDP), subunits of bacterial peptidoglycan present in serum [35,39,41]. Deletion or mutation of the LRR domain abolishes cAMP-PKA activation in the presence of MDPs [35]. CO₂ or HCO₃⁻ bind to the CYCc domain, stimulating the production of cAMP required for hyphal growth [33]. Both endogenous and exogenous QSM farnesol and 3-oxo-C12-homoserine lactone (HSL) block the hyphal growth by binding to the CYCc domain and inhibiting the activity of Cyr1 [82].

Temperature-dependent morphogenesis via the cAMP-PKA pathway is governed by the heat shock chaperone protein, Hsp90, whose expression is regulated by the heat shock transcription factor, Hsf1 [125]. Under basal conditions, Hsp90 and its cochaperone Sgt2 interact with Cyr1 and repress it [39,126]. Temperature elevation results in cellular stress leading to an increase in competing Hsp90 client proteins, thereby relieving Hsp90 repression of Cyr1. Inhibition of Hsp90 leads to filamentous growth under non-inducing conditions.

Cell cycle perturbation also induces morphogenesis via the Ras-cAMP-PKA signaling pathway. Disrupting cell cycle progression by treating with the DNA synthesis inhibitor hydroxyurea (HU) arrests cells in the S phase, while prolonged depletion of Cln3 arrests cells in the G1 phase [127,128]. Arrested cells switch to filamentous growth and re-enter the cell cycle via Ras1-activation of the cAMP-PKA pathway [127,128]. However, while cell cycle arrest in G1 and S phase induces morphogenesis, different mechanisms are involved. HU-induced filamentation does not require the downstream transcription factor Efg1, and the hyphal-specific transcription factor Ume6 and the G1 cyclin Hgc1 but involves other PKA target genes [127,129]. In contrast, filamentation due to Cln3 depletion requires Efg1, Ume6 and Hgc1 [127-129].

4. The MAPK pathways

The mitogen-activated protein kinase (MAPK) signal transduction pathway consists of three components: the MAP kinase kinase kinase (MAPKKK), the MAP kinase kinase (MAPKK), and the MAP kinase (MAPK) (Figure 3). MAPK signaling is dependent on three phosphotransfer steps. Upon activation, MAPKKK becomes phosphorylated and triggers the phosphorylation of the MAPKK, which in turn phosphorylates the MAPK [130,131]. In *C. albicans*, the Cek1-mediated MAPK pathway and the PKC MAPK pathway are activated by different stimuli. They serve as patterns of cascades that are essential for its morphogenesis and virulence, as shown in Figure 3 [131,132]. Apart from the cAMP-PKA signaling pathway, Ras1 also signals through the MAPK signaling cascade (Cek1-mediated) to coordinate filamentation in response to nitrogen starvation conditions via the Mep2 sensor [101].

4.1. The Cek1-mediated MAPK pathway

C. albicans extracellular signal-regulated kinase (ERK)-like 1 (Cek1)-mediated MAPK pathway is involved in cell wall biogenesis and virulence [133,134]. The Cek1-mediated MAPK pathway also plays an important role in hyphal development through the activation of downstream transcription factor Cph1, a positive regulator of filamentation [100,135]. This pathway can be induced by several factors such as low nitrogen, cell wall damage, osmotic stress, and embedding matrix. Under nitrogen starvation conditions, this pathway is activated by the ammonium permease Mep2 via a Ras1-dependent manner [101]. Cdc42, an essential GTPase, and its GEF Cdc24 are required for filamentous growth and virulence [136,137]. Upon interactions with Cdc24 and Ras1, activated Cdc42 turns on downstream effectors, including p21-activated kinase (PAK) Cst20 and Cla4, which then triggers concerted phosphorylation of the Ste11 (MAPKKK), Hst7 (MAPKK), and Cek1 (MAPK) (Figure 3) [132]. Mutations in the Cek1-mediated cascade cause defects in hyphal development to a different degree under certain conditions and result in attenuated virulence in animal models [138,139]. The Cek1-mediated MAPK pathway can also be activated through its upstream transmembrane proteins via cell wall damage or osmotic stress. Sho1, Opy2, and Msb2 form a complex that interacts with Cdc42 and Cst20, triggering Cek1 phosphorylation [138]. *sho1Δ/Δ*, *opy2Δ/Δ*, and *msb2Δ/Δ* mutants displayed altered sensitivity to cell wall damaging agents such as Congo Red, zymoylase, and tunicamycin, suggesting their roles in cell wall biogenesis [140,141]. *sho1Δ/Δ* mutants are sensitive to osmotic stress (i.e., 1 M sodium chloride), suggesting its additional role in osmotic stress signaling. The Cek1-mediated MAPK pathway responds to embedded matrix conditions by initiating a signaling cascade that ultimately activates Cph1 via Cek1 [22]. Rac1, Lmo1, and its exchange factor Dck1 are essential for invasive filamentous growth in the embedding matrix [142,143]. In contrast to Cdc42, Rac1 is not required for serum-induced hyphal growth [144]. *rac1Δ/Δ*, *lmo1Δ/Δ*, and *dck1Δ/Δ* mutants were observed to exhibit filamentous growth defects on solid agar and increased sensitivity to cell wall damaging agents, such as Calcofluor White and Congo Red [142]. Intriguingly, the overexpression of the Cek1 MAP kinase in *rac1Δ/Δ*, *lmo1Δ/Δ*, and *dck1Δ/Δ* deletion mutants, restores invasive filamentous growth on solid media, suggesting that Rac1,

Lmo1, and Dck1 function together upstream of the Cek1-mediated MAPK pathway during invasive filamentous growth [142]. The downstream transcription factor, Cph1, is essential for hyphal growth on solid agar but not in liquid media [100]. It was found that *efg1 cph1* double deletion mutants cannot form filaments under hypha-inducing conditions and are avirulent in animal models [31]. However, *efg1 cph1* double deletion mutants can form filamentation when embedded in the matrix, suggesting the involvement of other transcription factors for hyphal development under this condition [145].

4.2. The PKC MAPK pathway

The Protein kinase C (PKC) MAPK pathway is commonly known as the cell wall integrity pathway [131]. Pkc1 activation leads to a MAPK cascade activation of Bck1 (MAPKKK), Mkk1 (MAPKK), and Mkc1 (MAPK). Cellular morphogenesis in *C. albicans* is a highly dynamic process controlled by a master regulator, Rho1, in response to various stressors (Figure 3) [146]. Rho1, the master regulator of the cell wall integrity signaling cascade, is activated by the GEF Rom2 and inactivated by the GAP Lrg1 [147]. Recently, the PKC-MAPK pathway was discovered to regulate *C. albicans* morphogenesis through the co-regulation of cAMP signaling [148]. Interestingly, Rho1 plays an important role in filamentation through Pkc1. Pkc1 was found to be a global regulator of *C. albicans* morphogenesis through the regulation of adenylyl cyclase Cyr1. A reduction of Cyr1 activity was observed in *pkc1Δ/Δ* mutants. Lrg1 deactivates Rho1 by locking it in its inactive form, which suppresses the yeast-to-hyphae transition. *C. albicans* morphogenesis is independent of its canonical MAPK cascade. Mutants deleted of *BCK1* or *MKC1* do not impair the filamentous growth in response to the Hsp90 inhibitor geldanamycin or serum. Although the downstream transcription factors of Mkc1 have previously been proposed to be Efg1, Czf1, and Bcr1, to date, *C. albicans* morphogenesis through distinct effector(s) remains elusive [148,149].

5. Negative regulators of hyphal morphogenesis

C. albicans morphogenesis is negatively regulated by the transcriptional repressors Tup1, Nrg1, and Rfg1 [150-152]. Tup1 is a global transcriptional repressor, and its inactivation leads to constitutive filamentous growth and derepression of hyphal-specific genes [124,150,153]. Nrg1 and Rfg1 are well characterized DNA-binding proteins, which regulate different subsets of hyphal-specific genes by recruiting co-repressor Tup1. A DNA microarray analysis revealed significant upregulation of 61 genes in response to serum and 37°C [154]. Approximately half of these genes are found to be repressed by the transcription factors Tup1, Nrg1, and Rfg1, suggesting their importance in repressing hyphal morphogenesis. *C. albicans* cells that lack these repressors develop into pseudohyphae with the expression of hyphal-specific genes [155]. Surprisingly, only *nrg1Δ/Δ* mutants form hyphae in response to serum. In addition, *nrg1Δ/Δ* mutants appear to display stronger hyphal phenotypes than *rfg1Δ/Δ* mutants, suggesting its predominant role in the negative regulation of hyphal growth [156].

5.1. The farnesol-mediated inhibition pathway

Though Tup1 is found to act independently of the cAMP-PKA and MAPK pathways to regulate morphogenesis, it seems to play a crucial role in the farnesol response pathway [90,153]. Farnesol, an endogenous QSM, is produced when the cell densities of *C. albicans* are high. While farnesol can block the yeast-to-hyphae transition, it cannot block the elongation of preexisting filaments [157-159]. Morphological and transcriptional studies, which investigated the possible functional overlap between farnesol and hyphal transcriptional repressors, have demonstrated the direct involvement of Tup1 in the farnesol-mediated inhibition of filamentous growth [90]. *tup1Δ/Δ* and *nrg1Δ/Δ* mutants display elevated levels of farnesol and are constitutively filamentous even in the presence of exogenous farnesol. In the presence of farnesol, *TUP1* levels increase, but *NRG1* and *RFG1* levels are unaffected [90]. Further targeted studies on the farnesol-mediated

inhibition pathway have unraveled its dedicated mechanistic control of filamentous growth [160]. Upon inoculation of cells, where farnesol inhibition is relieved, the transcriptional repressor Cup9 is constantly degraded by the N-end rule E3 ubiquitin ligase Ubr1, allowing the expression of kinase Sok1 and subsequent degradation of Nrg1. In contrast, the presence of farnesol inhibits the degradation of Cup9, thereby repressing Sok1 expression, which in turn blocks the degradation of Nrg1 and hyphal development [160].

5.2. The roles of negative regulators Tup1 and Nrg1 in hyphal elongation

Critical regulators of hyphal initiation and the activation of hypha-associated genes, such as Efg1, Cph1, Czf1, and Flo8, are shown in Figure 4. Thereafter, a second regulatory network is required for the hyphal elongation process and long-term maintenance of hyphal growth through Hgc1, Eed1, and Ume6, which are negatively regulated by Tup1 and Nrg1 [138,161,162]. Eed1 was first identified in oral tissue infections from patients suffering from oral disease, and its associated regulatory network was explored through comprehensive transcriptomics analysis [161]. Eed1 is positively regulated by Efg1 as the overexpression of *EED1* partially rescues the hyphal defects in *efg1Δ/Δ* mutants. *EED1* expression is significantly up-regulated in the continuously filamentous *nrg1Δ/Δ* and *tup1Δ/Δ* mutants under non-hyphae inducing conditions [161]. In contrast, under hyphae-inducing conditions, *EED1* levels were slightly decreased in *nrg1Δ/Δ* and *tup1Δ/Δ* mutants, but elevated 10-fold in wild-type cells. Collectively, this suggests that *EED1* is repressed by both Nrg1 and Tup1 in wild-type *C. albicans*. Ume6 acts downstream of Eed1 as the overexpression of *UME6* restored the hyphal elongation defect observed in *eed1Δ/Δ* mutants [161]. *UME6* expression levels were significantly down-regulation in *eed1Δ/Δ* mutants [161]. The *HGC1* expression is detected within 5 minutes of hyphal induction, whereas the *UME6* expression is only detected after 15 min upon induction [163]. This suggests that a Ume6-independent mechanism initially induces *HGC1*. Nrg1 and Tup1 negatively regulate both Ume6 and Hgc1 [155,164]. Ume6 could also be induced as a result of relief of transcriptional repression by the Nrg1-Tup1 complex.




Figure 4. Regulation of hyphal elongation requires mechanisms for initiation and long-term maintenance. Initiation of hypha growth requires transcription factors such as Efg1, Cph1, Czf1 and Flo8. Subsequent elongation process and maintenance require the involvement of Hgc1, Eed1 and Ume6. Both Eed1 and Ume6 are negatively regulated by Tup1 and Nrg1.

5.3. O₂ and CO₂ signaling pathways for sustained hyphal development

The stability of hyphae-specific transcription factor Ume6 is governed by two parallel pathways in response to O₂ and CO₂ concentrations [165,166]. Ofd1 negatively regulates the stability of Ume6 by E3 ubiquitin ligase Ubr1 under hypoxia conditions. *ofd1Δ/Δ* and *ubr1Δ/Δ* mutants can maintain hyphal elongation in atmospheric O₂ and 5% CO₂ [165,166]. However, the deletion of *UBR1* does not block Ume6 degradation in atmospheric CO₂, suggesting the involvement of additional E3 ubiquitin ligase in response to CO₂ [166]. Recently, it was discovered that CO₂, an inducer of filamentous growth, also plays a critical role in the sustenance of hyphal growth in response to high CO₂ (5%) [166]. In the CO₂ signaling of sustained hyphal growth, a type 2C protein phosphatase (PP2C)

Ptc2 and a cyclin-dependent kinase Ssn3 were identified to be the major positive and negative regulators, respectively [166]. High CO₂ induces Ptc2-mediated dephosphorylation of Ssn3. Consequently, the hypophosphorylated Ssn3 fails to phosphorylate Ume6 at the S437 residue. This prevents Ume6 from the E3 ubiquitin ligase SCF^{Grr1} for subsequent ubiquitination, resulting in stabilization of Ume6 for the sustenance of hyphal growth.

5.4. Negative regulators as potential drug targets

Recent discoveries have introduced novel compounds that inhibit *C. albicans* hyphal and biofilm formation through the up-regulation of negative regulators Tup1 and Nrg1 [167,168]. Treatment of *C. albicans* with novel synthetic SR analogues, 5-[3-substituted-4-(4-substituted benzyloxy)-benzylidene]-2-thioxo-thiazolidin-4-one derivatives, resulted in a 3 to 4-fold increase in the expression of *TUP1* and a 2-fold increase in the expression of *NRG1*, which effectively inhibits the hyphal morphogenesis [167]. Copper oxide nanoparticle (Cu₂O-NP) was found to inhibit the yeast-to-hyphae transition through the down-regulation of *RAS1* and up-regulation of *NRG1* and *TUP1* [168]. Exploiting the negative regulators as drug targets holds excellent potential for future clinical applications. There is a growing interest in applying nanoparticles on medical devices, prosthetic devices, and catheters to combat polymicrobial biofilms in clinical settings.

6. Mechanisms of hyphal morphogenesis

6.1. Septin ring formation

Although the septin subunits are static in budding yeast cells, upon hyphal induction, Cdc3, Cdc12, and Sep7 form a stable core, while the Cdc10 subunit becomes dynamic, shuttling between the septin ring and the cytoplasm [169]. Cdc3 and Cdc12 are essential, whereas Cdc10 and Cdc11 are not. However, the deletion of *CDC10* and *CDC11* leads to defects in cytokinesis. During hyphal growth, Cdc11 is phosphorylated by the cyclin-CDK (cyclin-dependent kinase) complex Ccn1-Cdc28, and another cyclin-CDK complex Hgc1-Cdc28 maintains its phosphorylated state; and mutations to the phosphorylation sites in Cdc11 impairs maintenance of polarized growth [170]. Cdc11 phosphorylation by the septin ring-associated kinase Gin4 primes it for further phosphorylation by Ccn1-Cdc28 [170]. Both *cdc10* and *cdc11* mutants have abnormalities in septum formation during hyphal growth and form curved hyphae [171,172]. Cdc10 dynamics is dependent on Sep7 and its phosphorylation status [169]. *sep7Δ/Δ* mutants can form hyphae, but the hyphal compartments separate after cytokinesis. Ccn1-Cdc28 and Hgc1-Cdc28 phosphorylate Gin4, which in turn phosphorylates Sep7 [173,174]. Deletion of *GIN4* disrupts the formation of septin rings in germ tubes resulting in a severe cytokinesis defect; *gin4* mutants form pseudohyphae constitutively and cannot form true hyphae upon serum induction [175]. Depletion of Gin4 in G1 cells blocks septin ring formation [174]. Sep7 is dephosphorylated by the protein phosphatase 2A (PP2A), mediated by the structural subunit Tpd3 and the catalytic subunit Pph21 [176]. Deletion of *PPH21* or *TPD3* or its regulators, *CDC55* or *RTS1*, leads to the hyperphosphorylation of Sep7 and the disruption of septin organization [176,177]. *cdc55Δ/Δ* mutants grow as pseudohyphae under yeast growth conditions, while *rts1Δ/Δ* mutants grow as round, enlarged multinucleated cells. Both *cdc55Δ/Δ* and *rts1Δ/Δ* mutants display hyphal defects.

The nucleus migrates out from the mother cell to the septin band within the developing hyphae, and the first nuclear division occurs in this subapical compartment [178]. One daughter nucleus migrates back to the mother cell, while the other nucleus migrates to the apical compartment. After mitosis, the protein phosphatase Cdc14, which regulates mitotic exit, localizes to the septum in yeast cells and dephosphorylates the Mob2-Cbk1 complex, allowing the transcription factor Ace2 to translocate to the nucleus and activate the transcription of genes involved in cell separation [179]. However, in hyphal cells, Cdc14 does not localize to the septum, and Mob2-Cbk1 remains at the hyphal tip [179]; thus, cytokinesis does not result in cell separation or the formation of a

constriction between cells as observed in yeast or pseudohyphae, respectively. The septin ring splits into two rings with the formation of the primary septum dividing the hyphal compartments. Both rings are maintained in hyphal cells, unlike in yeast and pseudohyphal cells, where the septin rings are disassembled after cytokinesis. However, in *sep7Δ/Δ* mutants, Cdc14 can localize to the hyphal septum, activating the Ace2-dependent cell separation program, resulting in hyphal cell separation [180]. The subapical compartment of the hyphae is vacuolated and remains in the G1 phase.

The nucleosome assembly protein, Nap1, plays a role in septin ring formation and dynamics [181]. Deletion of *NAP1* results in constitutively pseudohyphal cells that can transit to true hyphae under hyphal-inducing conditions. Phosphorylation of Nap1 occurs in a cell-cycle-dependent manner, which involves Gin4 and Cla4, a second septin ring-associated kinase. Phosphorylated Nap1 translocates from the cytoplasm to the emerging bud neck. In *cdc10Δ/Δ* and *cdc11Δ/Δ* mutants, Nap1 remains in the cytoplasm even though it is hyperphosphorylated. After mitosis, Nap1 is dephosphorylated in a manner that is dependent upon PP2A and Cdc14.

6.2. Polarization of the actin cytoskeleton

Actin cytoskeleton polarization is required for the morphogenesis of *C. albicans*, regardless of cell type. The actin cytoskeleton, made up of actin patches and cables, maintains directional growth by directing vesicular flow for tip expansion. In yeast and pseudohyphae, polarized growth is driven by the polarisome, a complex that includes the polarisome scaffold protein Spa2, the formin Bni1 that serves as the actin cable nucleator, and the formin-actin-binding protein Bud6 [182]. Actin cables, comprised of long bundles of actin filaments, converge at the apical site. During polarized growth, post-Golgi membrane-bound secretory vesicles are continuously delivered to the apical site, supplying material required to expand the plasma membrane and synthesize new cell walls. The vesicles are tethered to the actin cables by the Rab-type GTPase Sec4, activated by the GEF Sec2 [183,184], while the class V myosin, Myo2, complexed to the regulatory light chain Mlc1, provides the motive force for vesicle transport [185]. Upon arrival at the plasma membrane, the secretory vesicles dock with the exocyst before fusing with the plasma membrane. The exocyst is a complex that comprises Sec3, Sec5, Sec6, Sec8, Sec10, Sec15, Exo70, and Exo84 [186]. Sec4 mediates vesicle tethering with the exocyst through its interaction with Sec15 [183,184].

Although the polarisome and exocyst complexes also localize to the hyphal tip, polarized growth in hyphae is driven by a Spitzenkörper, a vesicle-rich structure responsible for hyphal growth directionality, which is present during all stages of the cell cycle, including septation [187]. Spa2, Bni1, and Bud6 coordinate the functions of the Spitzenkörper and the polarisome complex at the hyphal tip [185,187]. During hyphal growth, the post-Golgi secretory vesicles travel along actin cables to the Spitzenkörper, which acts as a vesicle supply center and is maintained at a fixed distance from the hyphal tip (Figure 5). The vesicle-associated proteins Sec4, Sec2, and Mlc1 are localized to the Spitzenkörper during hyphal growth [185,188]. At the Spitzenkörper, the secretory vesicles are loaded onto actin cables nucleated by the polarisome and transported to the plasma membrane, where they dock with the exocyst. Actin cables are essential in hyphal growth, as their disruption inhibits hyphal formation [187]. Loss of *BNI1* does not affect bud emergence, as germ tube formation can be initiated in *bni1Δ/Δ* mutants. However, the germ tubes are wider in diameter, and *bni1Δ/Δ* mutants cannot maintain polarized cell growth [189]. Deletion of *SPA2* leads to polarity and hyphal growth defects [190]. *spa2Δ/Δ* mutants displayed random budding with multiple surface protrusions. Similar to the *bni1Δ/Δ* mutants, *spa2Δ/Δ* mutants can form germ tubes. However, unlike in *bni1Δ/Δ* mutants, hyphal growth can be maintained in the *spa2Δ/Δ* mutants, albeit in the form of severely swollen and curvy hyphae. Actin depolymerizing drugs, cytochalasin A and latrunculin A, disrupt the actin cytoskeleton, thus inhibiting hyphal growth and also suppressing the

at the hyphal tip throughout hyphal growth [197]. Endocytosis in *C. albicans* mainly occurs via clathrin-mediated endocytosis, and various genes involved in the process have been studied. Sla1 and Sla2 are actin cytoskeletal proteins involved in actin patch organization and dynamics, as well as actin cable polarization, and necessary for normal endocytosis [199-204]. Cortical actin patches formed in *sla1Δ/Δ* mutants are depolarized and less dynamic and formed short filaments [201,205]. *sla2Δ/Δ* mutants cannot undergo hyphal and pseudohyphal growth as the localization and orientation of actin patches and cables are defective [200,206]. *sla2Δ/Δ* mutants grow slower and form enlarged cells, as Swe1, the morphogenesis checkpoint kinase, delays cell cycle progression. Swe1 phosphorylates the Clb2-Cdc28 complex in response to perturbations to the actin cytoskeleton, thus delaying the normal transition from polarized growth to isotropic bud growth and delaying nuclear division. Pan1 is a clathrin-mediated endocytosis scaffold protein that is essential for endocytosis [207]. Depletion of Pan1 leads to the formation of thick and swollen cells that have abnormal filamentation. The inhibitory protein kinase Akl1 interacts with Pan1 to repress endocytosis, suppressing hyphal elongation [208]. Deletion of *AKL1* results in faster hyphal elongation rates and longer hyphae, while *AKL1* overexpression reduced hyphal elongation rates. However, overexpression of *PAN1* counteracts the effects of *AKL1* overexpression.

The myosin type I protein Myo5, the Wiskott-Aldrich Syndrome protein (WASP) homolog *WAL1*, and the WASP-interacting protein Vpr1 form a complex similar to that in *Saccharomyces cerevisiae* [209]. The Vpr1-Wal1-Myo5 complex is required for the polarized distribution of cortical actin patches. The deletion of *MYO5* leads to mislocalization of cortical actin patches, with the patches dispersed throughout the bud and the mother cell, resulting in excessive isotropic growth [210]. *myo5Δ/Δ* mutants are unable to endocytose and cannot form hyphae [210]. Deletion of *WAL1* and *VRP1* leads to defects in polarized growth [209,211]. *wal1Δ/Δ* mutants can initiate but cannot maintain hyphal growth. Instead, *wal1Δ/Δ* mutants form elongated, pseudohyphal cells under hyphae-inducing conditions. *vpr1Δ/Δ* mutants have a defect in hyphal formation that is slightly less severe than in *wal1Δ/Δ* mutants. Cortical actin patches are depolarized in both the mother cells and buds of *vpr1Δ/Δ* mutants. Myo5 and Wal1 activate the actin module Arp2/3 complex to initiate actin polymerization. Deleting *ARP2* or *ARP3* leads to an inability to form hyphae, although endocytosis is not abolished [204,212]. Deleting *RVS161* and *RVS167*, which encode Bin-Amphiphysin-Rvs (BAR) domain proteins, results in defective actin patch polarization, with the *rvs161Δ/Δ* mutant displaying a more severe defect in endocytosis and morphogenesis than the *rvs167Δ/Δ* mutant [202].

6.3. The role of Ras- and Rho-family GTPase

The small Ras- and Rho-family GTPases play essential roles in hyphal maintenance. The small Rho GTPase Cdc42 is the master regulator of polarized growth. Cdc42 affects hyphal growth and maintenance in at least two ways. Firstly, Cdc42 affects morphogenesis at the transcriptional level. Reduced expression levels of Cdc42 lead to decreased expression of hyphal-specific genes [213]. Secondly, decreasing cellular levels of active Cdc42 results in yeast and hyphae larger and rounder in shape, indicative of polarized growth defect [213,214]. Cdc42 cycles between GDP- and GTP- bound states. The GEF Cdc24 mediates the formation of GTP-bound Cdc42 [137,213]. The GAPs Rga2 and Bem3 mediate the return of Cdc42 from the GTP- to the GDP-bound form. Cdc42 and Cdc24, both required for viability, localize to the hyphal tip during hyphal growth [137,213]. Bem3 is localized to the apical zone of polarized growth, while Rga2 is localized to the septum and is phosphorylated in a hyphal-specific manner [215]. Loss of Rga2 and Bem3 results in forming a Spitzenkörper-like structure under pseudohyphal-promoting conditions, and the mutants have a morphology resembling true hyphae. Bem1 is a polarity establishment scaffolding protein that binds GTP-bound Cdc42, keeping it localized to the apical site [216].

The Ras-like GTPase Rsr1, a landmark protein that is the master regulator of the bud site selection system, regulates the amount and distribution of Cdc42 activity at the hyphal tip [217,218]. Rsr1 activity is regulated by the GEF Bud5 and the GAP Bud2. Bud5 is localized to the apical site, while Bud2 is localized to the subapical region and septin ring. *rsr1Δ/Δ* mutants have defects in polarized growth; yeast cells are larger and rounder, while the hyphae are wider than wild-type cells. Rsr1 is involved in regulating the recruitment and spatial distribution of vesicles at the hyphal tip [217,218]. Loss of Rsr1 affects the size of the fixed region to which vesicles are delivered and also affects the localization of exocyst subunits [219]. Rsr1 may play a role in limiting the competition for Cdc42 between the septum and the hyphal tip.

7. CDKs, cyclins, and their roles in hyphal morphogenesis

Maintenance of cell signalling is important for cell cycle progression and cell growth. The cell-cycle-associated cyclins and CDKs tightly regulate the small GTPases and other components of polarized growth. *C. albicans* has three G1 cyclins (Ccn1, Cln3, and Hgc1), two B-type mitotic cyclins (Clb2, Clb4), of which only Cln3 and Clb2 are essential. The essential CDK Cdc28 serves as the master regulator that controls cell cycle progression at G1/S and G2/M phases via specific cyclin interactions that dictate the timing of the phases. Levels of the G1 and B-type mitotic cyclins oscillate during the cell cycle, and a single cyclin-Cdc28 complex can regulate multiple events within each phase of the cell cycle. Cdc28 usually is stable and present at constant levels throughout the cell cycle; however, its depletion leads to filamentous growth [220]. Ccn1 and Cln3 levels in yeast cells are high in the G1 phase, coinciding with bud emergence and apical growth, and decline in the early G2 phase. Clb2 levels peak in the early G2/M phase, while Clb4 levels reach the peak in the mid-G2/M phase [221]. Levels of both B-cyclins start to decline in the M phase and disappear during exit from mitosis [179,221].

In hyphal cells, polarized growth continues at the apical site throughout the cell cycle, indicating the decoupling of cell elongation from the cell cycle. Ccn1 and Cln3 levels are accumulated earlier and persist for a longer time during hyphal growth [221,222], extending the G1 phase in the early germ tube. Accumulation of the mitotic cyclins, Clb2 and Clb4, is delayed in hyphal cells. Although it is not required for the initiation of hyphal growth, high levels of Ccn1 are required for maintenance of hyphal growth, along with Cln3. The forkhead family transcription factor, Fkh2, usually undergoes cell-cycle dependent phosphorylation to induce the expression of genes that regulate cell-cycle progression [223,224]. However, upon hyphal induction, Fkh2 is phosphorylated by Ccn1/Cln3-Cdc28 and Mob2-Cbk1 in a cell cycle-independent manner, redirecting it to enhance the expression of hyphal-specific genes such as the hyphal-specific G1 cyclin *HGC1* (Figure 6) [223,224]. *fkh2Δ/Δ* mutants grow constitutively as pseudohyphae under both yeast and hyphal-inducing conditions [223,224]. During hyphal growth, Ccn1-Cdc28 and Cln3-Cdc28 complexes phosphorylate Mob2, the activator of Cbk1, the cell wall integrity kinase, inhibiting the activation of Ace2 (Figure 6) [225]. Cln3-Cdc28 complex regulates cortical actin patches via phosphorylation of Sla1 [201].




Figure 6. *C. albicans* morphogenesis is tightly regulated by the cell-cycle-associated cyclins and cyclin-dependent kinase (CDK). The essential CDK Cdc28 serves as the master regulator that controls the cell cycle progression at G1/S and G2/M phases by forming CDK complexes with specific cyclins. Levels of Cdc28 are relatively stable throughout the cell cycle and depletes during hyphal growth. In contrast, levels of the G1 and B-type mitotic cyclins oscillate during the cell cycle. G1 cyclins Cln3 and Ccn1 peaks in the G1 phase and decline in the early G2 phase, while B-type mitotic cyclin Clb2 peaks at the early G2/M phase and declines in the M phase. Upon hyphal induction, Fkh2 is phosphorylated by Cln3-Cdc28 and Ccn1-Cdc28 complexes in a cell-cycle-dependent manner to enhance the expression of hyphal-specific genes. The Hgc1-Cdc28 complex is essential for the maintenance of hyphal growth. The exocyst subunit Exo84 is phosphorylated by the Hgc1-Cdc28 complex for the regulation of polarized secretion. Phosphorylation of the septin subunit Cdc11 (by Ccn1-Cdc28 and Hgc1-Cdc28 complexes), GTPase-activating protein (GAP) Rga2 (by the Hgc1-Cdc28 complex), and the polarisome protein Spa2 (by Hgc1-Cdc28 and Clb2-Cdc28 complexes) promote polarized growth. Rga2 is phosphorylated and inactivated by Hgc1-Cdc28, which relieves the repression of the GTPase Cdc42. Phosphorylation of the transcription factor Efg1 and protein kinase Gin4 inhibit cell separation. Phosphorylated Efg1 binds to promoters of Ace2 target genes, inhibiting their transcription. Phosphorylated Gin4 modifies the dynamics of the septin ring by subsequent phosphorylation of Sep7 and deactivating the cell separation program via inhibition of the protein phosphatase Cdc14.

The hyphal-specific G1 cyclin Hgc1 does not regulate the cell cycle but plays a critical role in hyphal morphogenesis (Figure 6). Besides suppression by Tup1 and Nrg1, the expression of *HGC1* is positively regulated by the transcription factor Ume6, which ensures that Hgc1 is expressed throughout the cell cycle as long as the inducing conditions remain. Hgc1 interacts with Cdc28, forming a complex regulating by phosphorylation regulators and components of cell polarity, membrane trafficking, and cell separation, which is required to maintain hyphal growth (Figure 6). The Hgc1-Cdc28 complex phosphorylates and inactivates Rga2, sequestering it from the hyphal tip to allow Cdc42 localization at the hyphal tip to persist during polarized growth [214,226]. Hgc1-Cdc28, together with Clb2-Cdc28, phosphorylates Spa2, localizing the polarisome to the hyphal tip [227]. Hgc1-Cdc28 complex phosphorylates the exocyst subunits Exo84 and Sec2, allowing them to be recycled at the growing hyphal tip [188,228]. The Hgc1-Cdc28 complex phosphorylates Efg1, leading to Efg1 competitively binding to promoters of Ace2 target genes, thereby repressing the expression of cell separation activators to prevent cell

separation after cytokinesis [229]. The hgc1-Cdc28 complex also plays a role in regulating the septin ring dynamics during hyphal growth via Sep7.

The cyclins Pcl1 and Pcl5 and the CDK Pho85, although not essential for cell cycle progression, contribute to morphogenesis in response to environmental cues. The Pcl1-Pho85 complex is required for temperature-dependent filamentation induced by Hsp90 inhibition [230]. The transcription factor Hms1 is required for filamentation induced by high temperatures. The Pcl1-Pho85 complex phosphorylates Hms1, which then binds to hyphal-specific genes. It also regulates the degradation of the transcription factor Gcn4, which is indirectly involved in filamentous growth in response to amino acid starvation [231]. Gcn4 induces *PCL5* expression, and the Pcl5-Pho85 complex phosphorylates Gcn4, leading to its degradation [232,233].

8. Cell cycle perturbation leads to morphogenesis

Although hyphal growth is not directly controlled by the cell cycle, perturbing the cell cycle can cause significant pseudohyphal growth under non-hyphal-inducing conditions or block hyphal growth under hyphal-inducing conditions [128,129,234]. Loss of *Ccn1* does not induce morphogenesis but causes a filamentation defect under serum induction [222]. Depletion of *Clb3*, *Clb2*, or *Clb4* results in filamentous growth in the absence of hyphal-inducing stimuli [128,234]. Cells depleted of *Clb3* undergo cell cycle arrest in the G1 phase and form filaments before the resumption of the cell cycle [129]. In the absence of mitotic cyclins, polarized growth promoted by G1 cyclins is not entirely suppressed, and filamentation occurs. *Clb2*-depleted cells form elongated projections during cell cycle arrest and are inviable, whereas *Clb4*-depleted cells grow constitutively as pseudohyphae and remain viable [221]. Depletion of *Cdc28* also promotes filamentous growth [220].

Genotoxic stresses that disrupt cell cycle progression and activate DNA damage/replication checkpoints lead to filamentation [235]. Pharmacological inhibition of cell cycle progression by the DNA replication inhibitors hydroxyurea (HU) and aphidicolin (AC) or DNA damage induced by UV radiation or the alkylating agent methylmethane sulfonate (MMS) result in S phase arrest, inducing filamentous growth [127,236,237]. Checkpoint proteins play a crucial role in response to DNA replication and DNA damage stresses. The protein kinase Rad53 plays a central role in the DNA replication and DNA damage checkpoint. *rad53Δ/Δ* mutants cannot switch to filamentous growth in response to DNA replication inhibitors and DNA damage; mutations to the Rad53 FHA domains inhibit filamentation in response to DNA damage, but not cell cycle arrest [236,237]. Deletion of *RAD9*, which encodes a checkpoint protein upstream of Rad53, blocks DNA damage-induced filamentation [236]. Depleting the DNA repair protein Rad52 or deleting its gene results in the accumulation of spontaneous DNA damages that trigger the DNA damage checkpoint, resulting in filamentous growth [238]. After the stress is relieved, deactivation of the cell cycle checkpoint is necessary for the cell cycle to progress. Rad53 is dephosphorylated by the protein phosphatase 2A-like complex Pph3-Psy2. Deletion of *PPH3* that encodes a protein phosphatase, *PSY2* that encodes the regulatory subunit of Psy2, or *TIP41* that encodes the regulator of the Pph3-Psy2 complex enhances MMS-induced filamentous growth and delays the filament-to-yeast transition following DNA damage stress relief [239,240]. The histone acetyltransferases Hat1 and Hat2 are required for the repair of DNA damages caused by endogenous and exogenous agents; *hat1Δ/Δ* mutants accumulate DNA damages rapidly and switch to filamentous growth [241].

Perturbations to mitosis can also lead to the switch to filamentous growth under non-hyphal-inducing conditions. The cell cycle regulatory polo-like kinase Cdc5 is required for the early stages of nuclear division and chromatin separation and mediates spindle formation during the S phase [242]. Depletion of *Cdc5* leads to mitotic inhibition and blocks the cell cycle in the G2 phase, leading to hyphal-like growth; however, the cells eventually lose viability [242]. The cytoplasmic dynein, Dyn1, mediates nuclear movement

during mitosis. Deletion of *DYN1* or depletion of Dyn1 results in filamentous growth, which requires the spindle position checkpoint protein Bub2 [243,244]. Pharmacological perturbation of mitosis by the microtubule inhibitor nocodazole and deletion of *MAD2* that encodes a spindle assembly checkpoint protein leads to pseudohyphal growth [245].

Degradation of cyclins and CDK inhibitors is regulated mainly by ubiquitin-proteasome-dependent proteolysis and is required for orderly cell cycle progression. Degradation of these cell cycle regulatory proteins is mediated by two multiprotein ubiquitin ligase E3 complexes, the Skp1-Cullin/Cdc53-F-box (SCF) complex and the anaphase-promoting complex/cyclosome (APC/C) complex. The multiprotein SCF complex, consisting of the linker protein Skp1, the scaffold protein Cullin/Cdc53, and a substrate recognition F-box protein, plays a central role in regulating the temporal and spatial degradation of cell cycle regulatory proteins. Depleting the essential *CDC53* or the deletion of the F-box protein genes *CDC4* and *GRR1* lead to filamentous growth. SCF^{Cdc4} is required for the degradation of the CDK inhibitor Sol1 and the transcription factors Ume6 and Gcn4 [232,233,246]. Deleting *CDC4* leads to constitutively filamentous growth with a mix of hyphal and pseudohyphal cells [247]. As Sol1 represses the Clb2-Cdc28 complex, deletion of *CDC4* stabilizes Sol1, inhibiting the switch from apical to isotropic growth, resulting in a pseudohyphal phenotype. However, deleting *SOL1* in the *cdc4Δ/Δ* mutant background still gave rise to constitutively filamentous growth [246]. SCF^{Grr1} is required for the degradation of Ccn1 and Cln3. The deletion of *GRR1* stabilizes Ccn1 and Cln3 levels, and the *grr1Δ/Δ* mutants grow constitutively as pseudohyphae under yeast conditions [248].

The APC/C complex mediates protein degradation during mitotic progression [249]. While little is known about the APC/C complex in *C. albicans*, the two coactivators Cdc20 and Cdh1, have been identified recently. Cdc20 is essential and mediates the degradation of Clb2 and Cdc5. Depletion of Cdc20 results in the accumulation of Clb2 and Cdc5, leading to a delay in metaphase and telophase; and Cdc20-depleted cells grow as long filaments over time [249]. Cdh1 likely plays a role in regulating mitotic exit by influencing Clb2 and Cdc5 degradation; deletion of *CDH1* results in a delay in Clb2 degradation and elevated levels of Cdc5 [249]. *cdh1Δ/Δ* mutants display pleiotropic phenotypes, with a mix of yeast, elongated buds, and pseudohyphae.

9. Conclusion

In summary, the extensive research findings over the years have provided us with illuminating insights into the activation and regulation of hyphal morphogenesis in *C. albicans*. The factors involved are often crucial in controlling the balance between commensalism and invasive infection by *C. albicans*. The yeast-to-hyphal transition in *C. albicans* is highly dependent on the complex interplay between internal signal transduction signalling pathways and external environmental cues that reflect the host niches. This transition is governed by a complex network of signalling pathways, including the cAMP-PKA pathway, the MAP kinase pathways, and the Cek1-mediated and PKC pathways. Activation of these pathways in response to various cues triggers the activation of specific transcription factors such as Efg1, Flo8, Ume6, Tec1, and Cph1. Crosstalks between the cAMP-PKA and MAP kinase pathways add a further layer of complexity to the existing signalling network as multiple signalling pathways can converge to the same set of transcription factors. Following hyphal initiation, subsequent hyphal development requires delicate mechanisms to maintain hyphal elongation. Polarized growth requires continuous delivery of membrane-bound secretory vesicles, along the actin cables, to its site of polarized growth. The vesicles accumulate as Spitzenkörper in the subapical region before docking with the exocyst and polarisome components. This exocytosis process is counterbalanced by the endocytosis process. Cell morphology of *C. albicans* is known to be tightly linked to cell-cycle progression through cyclin-CDK complexes. One of the most important CDK complexes is the Hgc1-Cdc28 complex, which governs multiple cellular processes required for hyphal development. The Hgc1-Cdc28 complex plays important

roles in polarized growth, polarized secretion, and inhibition of cell separation, which ensures the formation of long tubular cells without constriction at the septal junction. Lastly, perturbation of the cell cycle can either induce or impair the highly polarized growth in *C. albicans* under different conditions.

10. Future Directions

However, much remains to be explored and unraveled in *C. albicans* morphogenesis. While the mechanisms behind hyphal induction during cell cycle arrest have been uncovered, there are still missing gaps. Future work could uncover the genes that regulate filamentation in response to the cell cycle perturbation and elucidate how the signals are transduced to the cAMP-PKA pathway and the activated downstream transcriptional regulators. The link between nutrient and osmotic stress and filamentous growth via the PKC pathway has been uncovered, but the downstream transcriptional regulators remain elusive. Applying evolutionary tools of systems biology in combination with animal-based studies will propel discoveries in this field. The recent development of the transposon-mediated mutagenesis systems in haploid *C. albicans* strains would allow genome-wide screening for novel genes with functions that influence morphogenesis [250,251]. Future work towards identifying additional downstream transcriptional regulatory genes could open up new avenues towards antifungal therapy development.

Author Contributions: All authors contributed to the manuscript preparation. E.W.L.C. conceived the original idea and drafted the manuscript; L.M.P. assisted in the initial draft preparation; Y.W. supervised the project and made substantial scientific contributions. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the funding from National Research Foundation (NRF2019-NRF-ISF003-3039) and National Medical Research Council (NMRC/OFIRG/0072/2018 and NMRC/OFIRG/0055/2019) of Singapore to Y.W..

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Scully, C.; El-Kabir, M.; Samaranayake, L.P. *Candida* and Oral Candidosis: A Review. *Crit Rev Oral Biol Med* **1994**, *5*, 125-157, doi:10.1177/10454411940050020101.
2. Achkar, J.M.; Fries, B.C. *Candida* Infections of the Genitourinary Tract. *Clin Microbiol Rev* **2010**, *23*, 253-273, doi:10.1128/cmr.00076-09.
3. Rosenbach, A.; Dignard, D.; Pierce, J.V.; Whiteway, M.; Kumamoto, C.A. Adaptations of *Candida albicans* for Growth in the Mammalian Intestinal Tract. *Eukaryot cell* **2010**, *9*, 1075-1086, doi:10.1128/ec.00034-10.
4. Kühbacher, A.; Burger-Kentischer, A.; Rupp, S. Interaction of *Candida* Species with the Skin. *Microorganisms* **2017**, *5*, 32, doi:10.3390/microorganisms5020032.
5. Pappas, P.G.; Lionakis, M.S.; Arendrup, M.C.; Ostrosky-Zeichner, L.; Kullberg, B.J. Invasive candidiasis. *Nat Rev Dis Primers* **2018**, *4*, 18026, doi:10.1038/nrdp.2018.26.
6. Bongomin, F.; Gago, S.; Oladele, R.O.; Denning, D.W. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J Fungi (Basel)* **2017**, *3*, 57, doi:10.3390/jof3040057.
7. Bassetti, M.; Mikulska, M.; Viscoli, C. Bench-to-bedside review: therapeutic management of invasive candidiasis in the intensive care unit. *Crit Care* **2010**, *14*, 244-244, doi:10.1186/cc9239.

8. Gudlaugsson, O.; Gillespie, S.; Lee, K.; Vande Berg, J.; Hu, J.; Messer, S.; Herwaldt, L.; Pfaller, M.; Diekema, D. Attributable mortality of nosocomial candidemia, revisited. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **2003**, *37*, 1172-1177, doi:10.1086/378745.
9. Macphail, G.L.; Taylor, G.D.; Buchanan-Chell, M.; Ross, C.; Wilson, S.; Kureishi, A. Epidemiology, treatment and outcome of candidemia: a five-year review at three Canadian hospitals. *Mycoses* **2002**, *45*, 141-145, doi:10.1046/j.1439-0507.2002.00741.x.
10. Zaoutis, T.E.; Argon, J.; Chu, J.; Berlin, J.A.; Walsh, T.J.; Feudtner, C. The epidemiology and attributable outcomes of candidemia in adults and children hospitalized in the United States: a propensity analysis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **2005**, *41*, 1232-1239, doi:10.1086/496922.
11. Morgan, J.; Meltzer, M.I.; Plikaytis, B.D.; Sofair, A.N.; Huie-White, S.; Wilcox, S.; Harrison, L.H.; Seaberg, E.C.; Hajjeh, R.A.; Teutsch, S.M. Excess mortality, hospital stay, and cost due to candidemia: a case-control study using data from population-based candidemia surveillance. *Infection control and hospital epidemiology* **2005**, *26*, 540-547, doi:10.1086/502581.
12. Brown, A.J.; Gow, N.A. Regulatory networks controlling *Candida albicans* morphogenesis. *Trends in microbiology* **1999**, *7*, 333-338, doi:10.1016/s0966-842x(99)01556-5.
13. Gow, N.A. Germ tube growth of *Candida albicans*. *Curr Top Med Mycol* **1997**, *8*, 43-55.
14. Mukaremera, L.; Lee, K.K.; Mora-Montes, H.M.; Gow, N.A.R. *Candida albicans* Yeast, Pseudohyphal, and Hyphal Morphogenesis Differentially Affects Immune Recognition. *Frontiers in immunology* **2017**, *8*, 629, doi:10.3389/fimmu.2017.00629.
15. Nemecek, J.C.; Wüthrich, M.; Klein, B.S. Global control of dimorphism and virulence in fungi. *Science (New York, N.Y.)* **2006**, *312*, 583-588, doi:10.1126/science.1124105.
16. Trevijano-Contador, N.; Rueda, C.; Zaragoza, O. Fungal morphogenetic changes inside the mammalian host. *Seminars in cell & developmental biology* **2016**, *57*, 100-109, doi:10.1016/j.semcdb.2016.04.008.
17. Slutsky, B.; Buffo, J.; Soll, D.R. High-frequency switching of colony morphology in *Candida albicans*. *Science (New York, N.Y.)* **1985**, *230*, 666-669, doi:10.1126/science.3901258.
18. Sudbery, P.; Gow, N.; Berman, J. The distinct morphogenic states of *Candida albicans*. *Trends in microbiology* **2004**, *12*, 317-324, doi:10.1016/j.tim.2004.05.008.
19. Braunsdorf, C.; Mailänder-Sánchez, D.; Schaller, M. Fungal sensing of host environment. *Cellular microbiology* **2016**, *18*, 1188-1200, doi:10.1111/cmi.12610.
20. Chin, V.K.; Foong, K.J.; Maha, A.; Rusliza, B.; Norhafizah, M.; Chong, P.P. Multi-step pathogenesis and induction of local immune response by systemic *Candida albicans* infection in an intravenous challenge mouse model. *Int J Mol Sci* **2014**, *15*, 14848-14867, doi:10.3390/ijms150814848.
21. Di Carlo, P.; Di Vita, G.; Guadagnino, G.; Cocorullo, G.; D'Arpa, F.; Salamone, G.; Salvatore, B.; Gulotta, G.; Cabibi, D. Surgical pathology and the diagnosis of invasive visceral yeast infection: two case reports and literature review. *World J Emerg Surg* **2013**, *8*, 38, doi:10.1186/1749-7922-8-38.
22. Noble, S.M.; Gianetti, B.A.; Witchley, J.N. *Candida albicans* cell-type switching and functional plasticity in the mammalian host. *Nat Rev Microbiol* **2017**, *15*, 96.
23. van der Meer, J.W.M.; van de Veerdonk, F.L.; Joosten, L.A.B.; Kullberg, B.-J.; Netea, M.G. Severe *Candida* spp. infections: new insights into natural immunity. *Int J Antimicrob Agents* **2010**, *36*, S58-S62, doi:10.1016/j.ijantimicag.2010.11.013.
24. Erwig, L.P.; Gow, N.A.R. Interactions of fungal pathogens with phagocytes. *Nat Rev Microbiol* **2016**, *14*, 163-176, doi:10.1038/nrmicro.2015.21.
25. Fradin, C.; De Groot, P.; MacCallum, D.; Schaller, M.; Klis, F.; Odds, F.C.; Hube, B. Granulocytes govern the transcriptional response, morphology and proliferation of *Candida albicans* in human blood. *Molecular microbiology* **2005**, *56*, 397-415, doi:10.1111/j.1365-2958.2005.04557.x.

26. Lorenz, M.C.; Bender, J.A.; Fink, G.R. Transcriptional response of *Candida albicans* upon internalization by macrophages. *Eukaryot cell* **2004**, *3*, 1076-1087, doi:10.1128/ec.3.5.1076-1087.2004.
27. Naglik, J.R.; Moyes, D.L.; Wächter, B.; Hube, B. *Candida albicans* interactions with epithelial cells and mucosal immunity. *Microbes Infect* **2011**, *13*, 963-976, doi:10.1016/j.micinf.2011.06.009.
28. Grubb, S.E.W.; Murdoch, C.; Sudbery, P.E.; Saville, S.P.; Lopez-Ribot, J.L.; Thornhill, M.H. Adhesion of *Candida albicans* to endothelial cells under physiological conditions of flow. *Infect Immun* **2009**, *77*, 3872, doi:10.1128/IAI.00518-09.
29. Yang, W.; Yan, L.; Wu, C.; Zhao, X.; Tang, J. Fungal invasion of epithelial cells. *Microbiological research* **2014**, *169*, 803-810, doi:10.1016/j.micres.2014.02.013.
30. Rogiers, O.; Frising, U.C.; Kucharíková, S.; Jabra-Rizk, M.A.; van Loo, G.; Van Dijck, P.; Wullaert, A. Candidalysin Crucially Contributes to Nlrp3 Inflammasome Activation by *Candida albicans* Hyphae. *mBio* **2019**, *10*, e02221-02218, doi:10.1128/mBio.02221-18.
31. Lo, H.J.; Köhler, J.R.; DiDomenico, B.; Loebenberg, D.; Cacciapuoti, A.; Fink, G.R. Nonfilamentous *C. albicans* mutants are avirulent. *Cell* **1997**, *90*, 939-949, doi:10.1016/s0092-8674(00)80358-x.
32. Schweizer, A.; Rupp, S.; Taylor, B.N.; Rölinghoff, M.; Schröppel, K. The TEA/ATTS transcription factor CaTec1p regulates hyphal development and virulence in *Candida albicans*. *Molecular microbiology* **2000**, *38*, 435-445, doi:<https://doi.org/10.1046/j.1365-2958.2000.02132.x>.
33. Klengel, T.; Liang, W.-J.; Chaloupka, J.; Ruoff, C.; Schröppel, K.; Naglik, J.R.; Eckert, S.E.; Mogensen, E.G.; Haynes, K.; Tuite, M.F.; et al. Fungal adenylyl cyclase integrates CO₂ sensing with cAMP signaling and virulence. *Current Biology* **2005**, *15*, 2021-2026, doi:10.1016/j.cub.2005.10.040.
34. Brock, M. Fungal metabolism in host niches. *Current opinion in microbiology* **2009**, *12*, 371-376, doi:<https://doi.org/10.1016/j.mib.2009.05.004>.
35. Xu, X.-L.; Lee, R.T.H.; Fang, H.-M.; Wang, Y.-M.; Li, R.; Zou, H.; Zhu, Y.; Wang, Y. Bacterial Peptidoglycan Triggers *Candida albicans* Hyphal Growth by Directly Activating the Adenylyl Cyclase Cyr1p. *Cell Host & Microbe* **2008**, *4*, 28-39, doi:<https://doi.org/10.1016/j.chom.2008.05.014>.
36. Maidan, M.M.; Thevelein, J.M.; Van Dijck, P. Carbon source induced yeast-to-hypha transition in *Candida albicans* is dependent on the presence of amino acids and on the G-protein-coupled receptor Gpr1. *Biochem Soc Trans* **2005**, *33*, 291-293, doi:10.1042/bst0330291.
37. Lu, Y.; Su, C.; Wang, A.; Liu, H. Hyphal development in *Candida albicans* requires two temporally linked changes in promoter chromatin for initiation and maintenance. *PLoS biology* **2011**, *9*, e1001105, doi:10.1371/journal.pbio.1001105.
38. Song, W.; Wang, H.; Chen, J. *Candida albicans* Sfl2, a temperature-induced transcriptional regulator, is required for virulence in a murine gastrointestinal infection model. *FEMS yeast research* **2011**, *11*, 209-222, doi:10.1111/j.1567-1364.2010.00710.x.
39. Shapiro, R.S.; Uppuluri, P.; Zaas, A.K.; Collins, C.; Senn, H.; Perfect, J.R.; Heitman, J.; Cowen, L.E. Hsp90 orchestrates temperature-dependent *Candida albicans* morphogenesis via Ras1-PKA signaling. *Curr Biol* **2009**, *19*, 621-629, doi:10.1016/j.cub.2009.03.017.
40. Taschdjian, C.L.; Burchall, J.J.; Kozinn, P.J. Rapid identification of *Candida albicans* by filamentation on serum and serum substitutes. *A.M.A. journal of diseases of children* **1960**, *99*, 212-215, doi:10.1001/archpedi.1960.02070030214011.
41. Tan, C.T.; Xu, X.; Qiao, Y.; Wang, Y. A peptidoglycan storm caused by beta-lactam antibiotic's action on host microbiota drives *Candida albicans* infection. *Nat Commun* **2021**, *12*, 2560, doi:10.1038/s41467-021-22845-2.
42. Hall, R.A.; De Sordi, L.; Maccallum, D.M.; Topal, H.; Eaton, R.; Bloor, J.W.; Robinson, G.K.; Levin, L.R.; Buck, J.; Wang, Y.; et al. CO₂ acts as a signalling molecule in populations of the fungal pathogen *Candida albicans*. *PLoS Pathog* **2010**, *6*, e1001193-e1001193, doi:10.1371/journal.ppat.1001193.

43. Miwa, T.; Takagi, Y.; Shinozaki, M.; Yun, C.W.; Schell, W.A.; Perfect, J.R.; Kumagai, H.; Tamaki, H. Gpr1, a putative G-protein-coupled receptor, regulates morphogenesis and hypha formation in the pathogenic fungus *Candida albicans*. *Eukaryot cell* **2004**, *3*, 919-931, doi:10.1128/EC.3.4.919-931.2004.
44. D'Souza, C.A.; Heitman, J. Conserved cAMP signaling cascades regulate fungal development and virulence. *FEMS Microbiology Reviews* **2001**, *25*, 349-364, doi:10.1111/j.1574-6976.2001.tb00582.x.
45. Maidan, M.M.; De Rop, L.; Serneels, J.; Exler, S.; Rupp, S.; Tournu, H.; Thevelein, J.M.; Van Dijck, P. The G protein-coupled receptor Gpr1 and the Galpha protein Gpa2 act through the cAMP-protein kinase A pathway to induce morphogenesis in *Candida albicans*. *Mol Biol Cell* **2005**, *16*, 1971-1986, doi:10.1091/mbc.e04-09-0780.
46. Zou, H.; Fang, H.M.; Zhu, Y.; Wang, Y. *Candida albicans* Cyr1, Cap1 and G-actin form a sensor/effector apparatus for activating cAMP synthesis in hyphal growth. *Mol Microbiol* **2010**, *75*, 579-591, doi:10.1111/j.1365-2958.2009.06980.x.
47. Lopes, J.P.; Stylianou, M.; Backman, E.; Holmberg, S.; Jass, J.; Claesson, R.; Urban, C.F. Evasion of Immune Surveillance in Low Oxygen Environments Enhances *Candida albicans* Virulence. *mBio* **2018**, *9*, e02120-02118, doi:10.1128/mBio.02120-18.
48. Desai, P.R.; van Wijlick, L.; Kurtz, D.; Juchimiuk, M.; Ernst, J.F. Hypoxia and Temperature Regulated Morphogenesis in *Candida albicans*. *PLOS Genetics* **2015**, *11*, e1005447, doi:10.1371/journal.pgen.1005447.
49. Giusani, A.D.; Vines, M.; Kumamoto, C.A. Invasive filamentous growth of *Candida albicans* is promoted by Czf1p-dependent relief of Efg1p-mediated repression. *Genetics* **2002**, *160*, 1749-1753.
50. Mulhern, S.M.; Logue, M.E.; Butler, G. *Candida albicans* transcription factor Ace2 regulates metabolism and is required for filamentation in hypoxic conditions. *Eukaryot cell* **2006**, *5*, 2001-2013, doi:10.1128/ec.00155-06.
51. Saputo, S.; Kumar, A.; Krysan, D.J. Efg1 directly regulates ACE2 expression to mediate cross talk between the cAMP/PKA and RAM pathways during *Candida albicans* morphogenesis. *Eukaryot cell* **2014**, *13*, 1169-1180, doi:10.1128/ec.00148-14.
52. Mayer, F.L.; Wilson, D.; Hube, B. *Candida albicans* pathogenicity mechanisms. *Virulence* **2013**, *4*, 119-128, doi:10.4161/viru.22913.
53. Davis, D.; Edwards, J.E., Jr.; Mitchell, A.P.; Ibrahim, A.S. *Candida albicans* RIM101 pH response pathway is required for host-pathogen interactions. *Infect Immun* **2000**, *68*, 5953-5959, doi:10.1128/iai.68.10.5953-5959.2000.
54. Davis, D.; Wilson, R.B.; Mitchell, A.P. RIM101-dependent and-independent pathways govern pH responses in *Candida albicans*. *Molecular and cellular biology* **2000**, *20*, 971-978, doi:10.1128/mcb.20.3.971-978.2000.
55. El Barkani, A.; Kurzai, O.; Fonzi, W.A.; Ramon, A.; Porta, A.; Frosch, M.; Mühlischlegel, F.A. Dominant Active Alleles of RIM101 (PRR2) Bypass the pH Restriction on Filamentation of *Candida albicans*. *Molecular and cellular biology* **2000**, *20*, 4635-4647, doi:10.1128/mcb.20.13.4635-4647.2000.
56. Davis, D.A. How human pathogenic fungi sense and adapt to pH: the link to virulence. *Curr Opin Microbiol* **2009**, *12*, 365-370, doi:10.1016/j.mib.2009.05.006.
57. Vylkova, S.; Carman, A.J.; Danhof, H.A.; Collette, J.R.; Zhou, H.; Lorenz, M.C. The fungal pathogen *Candida albicans* autoinduces hyphal morphogenesis by raising extracellular pH. *mBio* **2011**, *2*, e00055-00011, doi:10.1128/mBio.00055-11.
58. Vesely, E.M.; Williams, R.B.; Konopka, J.B.; Lorenz, M.C. N-Acetylglucosamine Metabolism Promotes Survival of *Candida albicans* in the Phagosome. *mSphere* **2017**, *2*, doi:10.1128/mSphere.00357-17.
59. Pande, K.; Chen, C.; Noble, S.M. Passage through the mammalian gut triggers a phenotypic switch that promotes *Candida albicans* commensalism. *Nat Genet* **2013**, *45*, 1088-1091, doi:10.1038/ng.2710.
60. Naseem, S.; Konopka, J.B. N-acetylglucosamine Regulates Virulence Properties in Microbial Pathogens. *PLoS Pathog* **2015**, *11*, e1004947, doi:10.1371/journal.ppat.1004947.
61. Sudbery, P.E. Growth of *Candida albicans* hyphae. *Nat Rev Microbiol* **2011**, *9*, 737-748, doi:10.1038/nrmicro2636.
62. Alvarez, F.J.; Konopka, J.B. Identification of an N-Acetylglucosamine Transporter That Mediates Hyphal Induction in *Candida albicans*. *Molecular biology of the cell* **2007**, *18*, 965-975, doi:10.1091/mbc.e06-10-0931.

63. Naseem, S.; Gunasekera, A.; Araya, E.; Konopka, J.B. N-acetylglucosamine (GlcNAc) induction of hyphal morphogenesis and transcriptional responses in *Candida albicans* are not dependent on its metabolism. *J Biol Chem* **2011**, *286*, 28671-28680, doi:10.1074/jbc.M111.249854.
64. Naseem, S.; Min, K.; Spitzer, D.; Gardin, J.; Konopka, J.B. Regulation of Hyphal Growth and N-Acetylglucosamine Catabolism by Two Transcription Factors in *Candida albicans*. *Genetics* **2017**, *206*, 299-314, doi:10.1534/genetics.117.201491.
65. Su, C.; Lu, Y.; Liu, H. N-acetylglucosamine sensing by a GCN5-related N-acetyltransferase induces transcription via chromatin histone acetylation in fungi. *Nat Commun* **2016**, *7*, 12916, doi:10.1038/ncomms12916.
66. Min, K.; Biermann, A.; Hogan, D.A.; Konopka, J.B. Genetic Analysis of NDT80 Family Transcription Factors in *Candida albicans* Using New CRISPR-Cas9 Approaches. *mSphere* **2018**, *3*, e00545-00518, doi:10.1128/mSphere.00545-18.
67. Rocha, C.R.; Schröppel, K.; Harcus, D.; Marcil, A.; Dignard, D.; Taylor, B.N.; Thomas, D.Y.; Whiteway, M.; Leberer, E. Signaling through adenylyl cyclase is essential for hyphal growth and virulence in the pathogenic fungus *Candida albicans*. *Mol Biol Cell* **2001**, *12*, 3631-3643, doi:10.1091/mbc.12.11.3631.
68. Parrino, S.M.; Si, H.; Naseem, S.; Groudan, K.; Gardin, J.; Konopka, J.B. cAMP-independent signal pathways stimulate hyphal morphogenesis in *Candida albicans*. *Molecular microbiology* **2017**, *103*, 764-779, doi:<https://doi.org/10.1111/mmi.13588>.
69. Min, K.; Naseem, S.; Konopka, J.B. N-Acetylglucosamine Regulates Morphogenesis and Virulence Pathways in Fungi. *J Fungi (Basel)* **2020**, *6*, 8.
70. Dabrowa, N.; Taxer, S.S.; Howard, D.H. Germination of *Candida albicans* induced by proline. *Infect Immun* **1976**, *13*, 830-835, doi:10.1128/iai.13.3.830-835.1976.
71. Land, G.A.; McDonald, W.C.; Stjernholm, R.L.; Friedman, T.L. Factors affecting filamentation in *Candida albicans*: relationship of the uptake and distribution of proline to morphogenesis. *Infect Immun* **1975**, *11*, 1014-1023, doi:10.1128/iai.11.5.1014-1023.1975.
72. Ghosh, S.; Navarathna, D.H.; Roberts, D.D.; Cooper, J.T.; Atkin, A.L.; Petro, T.M.; Nickerson, K.W. Arginine-induced germ tube formation in *Candida albicans* is essential for escape from murine macrophage line RAW 264.7. *Infect Immun* **2009**, *77*, 1596-1605, doi:10.1128/IAI.01452-08.
73. Miramon, P.; Lorenz, M.C. The SPS amino acid sensor mediates nutrient acquisition and immune evasion in *Candida albicans*. *Cellular microbiology* **2016**, *18*, 1611-1624, doi:10.1111/cmi.12600.
74. Martinez, P.; Ljungdahl, P.O. Divergence of Stp1 and Stp2 transcription factors in *Candida albicans* places virulence factors required for proper nutrient acquisition under amino acid control. *Molecular and cellular biology* **2005**, *25*, 9435-9446, doi:10.1128/MCB.25.21.9435-9446.2005.
75. Miramon, P.; Pountain, A.W.; van Hoof, A.; Lorenz, M.C. The Paralogous Transcription Factors Stp1 and Stp2 of *Candida albicans* Have Distinct Functions in Nutrient Acquisition and Host Interaction. *Infection and immunity* **2020**, *88*, doi:10.1128/IAI.00763-19.
76. Brega, E.; Zufferey, R.; Mamoun, C.B. *Candida albicans* Csy1p is a nutrient sensor important for activation of amino acid uptake and hyphal morphogenesis. *Eukaryot Cell* **2004**, *3*, 135-143, doi:10.1128/EC.3.1.135-143.2004.
77. Abdel-Sater, F.; Jean, C.; Merhi, A.; Vissers, S.; Andre, B. Amino acid signaling in yeast: activation of Ssy5 protease is associated with its phosphorylation-induced ubiquitylation. *J Biol Chem* **2011**, *286*, 12006-12015, doi:10.1074/jbc.M110.200592.
78. Martinez, P.; Ljungdahl, P.O. An ER packaging chaperone determines the amino acid uptake capacity and virulence of *Candida albicans*. *Mol Microbiol* **2004**, *51*, 371-384, doi:10.1046/j.1365-2958.2003.03845.x.
79. Silao, F.G.S.; Ward, M.; Ryman, K.; Wallström, A.; Brindefalk, B.; Udekwu, K.; Ljungdahl, P.O. Mitochondrial proline catabolism activates Ras1/cAMP/PKA-induced filamentation in *Candida albicans*. *PLOS Genetics* **2019**, *15*, e1007976, doi:10.1371/journal.pgen.1007976.

80. Grahl, N.; Demers, E.G.; Lindsay, A.K.; Harty, C.E.; Willger, S.D.; Piispanen, A.E.; Hogan, D.A. Mitochondrial Activity and Cyr1 Are Key Regulators of Ras1 Activation of *C. albicans* Virulence Pathways. *PLoS Pathog* **2015**, *11*, e1005133, doi:10.1371/journal.ppat.1005133.
81. Davis-Hanna, A.; Piispanen, A.E.; Stateva, L.I.; Hogan, D.A. Farnesol and dodecanol effects on the *Candida albicans* Ras1-cAMP signalling pathway and the regulation of morphogenesis. *Mol Microbiol* **2008**, *67*, 47-62, doi:10.1111/j.1365-2958.2007.06013.x.
82. Hall, R.A.; Turner, K.J.; Chaloupka, J.; Cottier, F.; De Sordi, L.; Sanglard, D.; Levin, L.R.; Buck, J.; Mühlshlegel, F.A. The quorum-sensing molecules farnesol/homoserine lactone and dodecanol operate via distinct modes of action in *Candida albicans*. *Eukaryot cell* **2011**, *10*, 1034-1042, doi:10.1128/EC.05060-11.
83. Chauhan, N.M.; Raut, J.S.; Karuppayil, S.M. A morphogenetic regulatory role for ethyl alcohol in *Candida albicans*. *Mycoses* **2011**, *54*, e697-703, doi:10.1111/j.1439-0507.2010.02002.x.
84. Martins, M.; Henriques, M.; Azeredo, J.; Rocha, S.M.; Coimbra, M.A.; Oliveira, R. Morphogenesis control in *Candida albicans* and *Candida dubliniensis* through signaling molecules produced by planktonic and biofilm cells. *Eukaryot cell* **2007**, *6*, 2429-2436, doi:10.1128/EC.00252-07.
85. Lindsay, A.K.; Morales, D.K.; Liu, Z.; Grahl, N.; Zhang, A.; Willger, S.D.; Myers, L.C.; Hogan, D.A. Analysis of *Candida albicans* mutants defective in the Cdk8 module of mediator reveal links between metabolism and biofilm formation. *PLoS Genetics* **2014**, *10*, e1004567-e1004567, doi:10.1371/journal.pgen.1004567.
86. Chen, H.; Fujita, M.; Feng, Q.; Clardy, J.; Fink, G.R. Tyrosol is a quorum-sensing molecule in *Candida albicans*. *Proceedings of the National Academy of Sciences of the United States of America* **2004**, *101*, 5048-5052, doi:10.1073/pnas.0401416101.
87. Alem, M.A.S.; Oteef, M.D.Y.; Flowers, T.H.; Douglas, L.J. Production of tyrosol by *Candida albicans* biofilms and its role in quorum sensing and biofilm development. *Eukaryot cell* **2006**, *5*, 1770-1779, doi:10.1128/EC.00219-06.
88. Ghosh, S.; Kebaara, B.W.; Atkin, A.L.; Nickerson, K.W. Regulation of aromatic alcohol production in *Candida albicans*. *Applied and environmental microbiology* **2008**, *74*, 7211-7218, doi:10.1128/aem.01614-08.
89. Chauhan, N.M.; Mohan Karuppayil, S. Dual identities for various alcohols in two different yeasts. *Mycology* **2020**, *12*, 25-38, doi:10.1080/21501203.2020.1837976.
90. Kebaara, B.W.; Langford, M.L.; Navarathna, D.H.M.L.P.; Dumitru, R.; Nickerson, K.W.; Atkin, A.L. *Candida albicans* Tup1 Is Involved in Farnesol-Mediated Inhibition of Filamentous-Growth Induction. *Eukaryot cell* **2008**, *7*, 980-987, doi:10.1128/ec.00357-07.
91. Lindsay, A.K.; Hogan, D.A. *Candida albicans*: Molecular interactions with *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *Fungal Biol Rev* **2014**, *28*, 85-96, doi:<https://doi.org/10.1016/j.fbr.2014.10.002>.
92. Hogan, D.A.; Vik, Å.; Kolter, R. A *Pseudomonas aeruginosa* quorum-sensing molecule influences *Candida albicans* morphology. *Molecular microbiology* **2004**, *54*, 1212-1223, doi:<https://doi.org/10.1111/j.1365-2958.2004.04349.x>.
93. Haiko, J.; Saeedi, B.; Bagger, G.; Karpati, F.; Özenci, V. Coexistence of *Candida* species and bacteria in patients with cystic fibrosis. *Eur J Clin Microbiol Infect Dis* **2019**, *38*, 1071-1077, doi:10.1007/s10096-019-03493-3.
94. Morales, D.K.; Grahl, N.; Okegbe, C.; Dietrich, L.E.P.; Jacobs, N.J.; Hogan, D.A. Control of *Candida albicans* Metabolism and Biofilm Formation by *Pseudomonas aeruginosa* Phenazines. *mBio* **2013**, *4*, e00526-00512, doi:10.1128/mBio.00526-12.
95. Ryan, R.P.; An, S.-q.; Allan, J.H.; McCarthy, Y.; Dow, J.M. The DSF Family of Cell-Cell Signals: An Expanding Class of Bacterial Virulence Regulators. *PLoS Pathog* **2015**, *11*, e1004986-e1004986, doi:10.1371/journal.ppat.1004986.
96. de Rossi, B.P.; García, C.; Alcaraz, E.; Franco, M. *Stenotrophomonas maltophilia* interferes via the DSF-mediated quorum sensing system with *Candida albicans* filamentation and its planktonic and biofilm modes of growth. *Revista Argentina de microbiologia* **2014**, *46*, 288-297, doi:10.1016/s0325-7541(14)70084-7.

97. Yang, D.; Hu, Y.; Yin, Z.; Gao, Q.; Zhang, Y.; Chan, F.Y.; Zeng, G.; Weng, L.; Wang, L.; Wang, Y. *Candida albicans* Ubiquitin and Heat Shock Factor-Type Transcriptional Factors Are Involved in 2-Dodecenoic Acid-Mediated Inhibition of Hyphal Growth. *Microorganisms* **2020**, *8*, 75.
98. Calera, J.A.; Zhao, X.J.; Calderone, R. Defective hyphal development and avirulence caused by a deletion of the SSK1 response regulator gene in *Candida albicans*. *Infect Immun* **2000**, *68*, 518-525, doi:10.1128/iai.68.2.518-525.2000.
99. Lee, K.L.; Buckley, H.R.; Campbell, C.C. An amino acid liquid synthetic medium for the development of mycelial and yeast forms of *Candida albicans*. *Sabouraudia* **1975**, *13*, 148-153, doi:10.1080/00362177585190271.
100. Liu, H.; Kohler, J.; Fink, G. Suppression of hyphal formation in *Candida albicans* by mutation of a STE12 homolog. *Science* **1994**, *266*, 1723-1726, doi:10.1126/science.7992058.
101. Biswas, K.; Morschhäuser, J. The Mep2p ammonium permease controls nitrogen starvation-induced filamentous growth in *Candida albicans*. *Molecular microbiology* **2005**, *56*, 649-669, doi:<https://doi.org/10.1111/j.1365-2958.2005.04576.x>.
102. Dabas, N.; Schneider, S.; Morschhäuser, J. Mutational Analysis of the *Candida albicans* Ammonium Permease Mep2p Reveals Residues Required for Ammonium Transport and Signaling. *Eukaryot cell* **2009**, *8*, 147-160, doi:10.1128/ec.00229-08.
103. Schrevers, S.; Van Zeebroeck, G.; Riedelberger, M.; Tournu, H.; Kuchler, K.; Van Dijck, P. Methionine is required for cAMP-PKA-mediated morphogenesis and virulence of *Candida albicans*. *Molecular microbiology* **2018**, *108*, 258-275, doi:<https://doi.org/10.1111/mmi.13933>.
104. Cao, C.; Wu, M.; Bing, J.; Tao, L.; Ding, X.; Liu, X.; Huang, G. Global regulatory roles of the cAMP/PKA pathway revealed by phenotypic, transcriptomic and phosphoproteomic analyses in a null mutant of the PKA catalytic subunit in *Candida albicans*. *Mol Microbiol* **2017**, *105*, 46-64, doi:10.1111/mmi.13681.
105. Hogan, D.A.; Sundstrom, P. The Ras/cAMP/PKA signaling pathway and virulence in *Candida albicans*. *Future microbiology* **2009**, *4*, 1263-1270, doi:10.2217/fmb.09.106.
106. Bockmühl, D.P.; Ernst, J.F. A potential phosphorylation site for an A-type kinase in the Efg1 regulator protein contributes to hyphal morphogenesis of *Candida albicans*. *Genetics* **2001**, *157*, 1523-1530.
107. Cao, F.; Lane, S.; Raniga, P.P.; Lu, Y.; Zhou, Z.; Ramon, K.; Chen, J.; Liu, H. The Flo8 transcription factor is essential for hyphal development and virulence in *Candida albicans*. *Molecular biology of the cell* **2006**, *17*, 295-307, doi:10.1091/mbc.e05-06-0502.
108. Fang, H.M.; Wang, Y. RA domain-mediated interaction of Cdc35 with Ras1 is essential for increasing cellular cAMP level for *Candida albicans* hyphal development. *Mol Microbiol* **2006**, *61*, 484-496, doi:10.1111/j.1365-2958.2006.05248.x.
109. Taylor, S.S.; Buechler, J.A.; Yonemoto, W. cAMP-dependent protein kinase: framework for a diverse family of regulatory enzymes. *Annual review of biochemistry* **1990**, *59*, 971-1005, doi:10.1146/annurev.bi.59.070190.004543.
110. Bockmühl, D.P.; Krishnamurthy, S.; Gerads, M.; Sonneborn, A.; Ernst, J.F. Distinct and redundant roles of the two protein kinase A isoforms Tpk1p and Tpk2p in morphogenesis and growth of *Candida albicans*. *Molecular microbiology* **2001**, *42*, 1243-1257, doi:10.1046/j.1365-2958.2001.02688.x.
111. Cloutier, M.; Castilla, R.; Bolduc, N.; Zelada, A.; Martineau, P.; Bouillon, M.; Magee, B.B.; Passeron, S.; Giasson, L.; Cantore, M.L. The two isoforms of the cAMP-dependent protein kinase catalytic subunit are involved in the control of dimorphism in the human fungal pathogen *Candida albicans*. *Fungal genetics and biology : FG & B* **2003**, *38*, 133-141, doi:10.1016/s1087-1845(02)00520-0.
112. Hoyer, L.L.; Cieslinski, L.B.; McLaughlin, M.M.; Torphy, T.J.; Shatzman, A.R.; Livi, G.P. A *Candida albicans* cyclic nucleotide phosphodiesterase: cloning and expression in *Saccharomyces cerevisiae* and biochemical characterization of the recombinant enzyme. *Microbiology (Reading, England)* **1994**, *140* (Pt 7), 1533-1542, doi:10.1099/13500872-140-7-1533.
113. Jung, W.H.; Stateva, L.I. The cAMP phosphodiesterase encoded by CaPDE2 is required for hyphal development in *Candida albicans*. *Microbiology (Reading, England)* **2003**, *149*, 2961-2976, doi:10.1099/mic.0.26517-0.

114. Bahn, Y.S.; Staab, J.; Sundstrom, P. Increased high-affinity phosphodiesterase PDE2 gene expression in germ tubes counteracts CAP1-dependent synthesis of cyclic AMP, limits hypha production and promotes virulence of *Candida albicans*. *Molecular microbiology* **2003**, *50*, 391-409, doi:10.1046/j.1365-2958.2003.03692.x.
115. Wilson, D.; Tutulan-Cunita, A.; Jung, W.; Hauser, N.C.; Hernandez, R.; Williamson, T.; Piekarska, K.; Rupp, S.; Young, T.; Stateva, L. Deletion of the high-affinity cAMP phosphodiesterase encoded by PDE2 affects stress responses and virulence in *Candida albicans*. *Molecular microbiology* **2007**, *65*, 841-856, doi:10.1111/j.1365-2958.2007.05788.x.
116. Wilson, D.; Fiori, A.; Brucker, K.D.; Dijck, P.V.; Stateva, L. *Candida albicans* Pde1p and Gpa2p comprise a regulatory module mediating agonist-induced cAMP signalling and environmental adaptation. *Fungal genetics and biology : FG & B* **2010**, *47*, 742-752, doi:10.1016/j.fgb.2010.06.006.
117. Harcus, D.; Nantel, A.; Marcil, A.; Rigby, T.; Whiteway, M. Transcription profiling of cyclic AMP signaling in *Candida albicans*. *Molecular biology of the cell* **2004**, *15*, 4490-4499, doi:10.1091/mbc.e04-02-0144.
118. Wang, Y. Fungal Adenylyl Cyclase Acts As a Signal Sensor and Integrator and Plays a Central Role in Interaction with Bacteria. *PLoS Pathog* **2013**, *9*, e1003612, doi:10.1371/journal.ppat.1003612.
119. Huang, G.; Huang, Q.; Wei, Y.; Wang, Y.; Du, H. Multiple roles and diverse regulation of the Ras/cAMP/protein kinase A pathway in *Candida albicans*. *Molecular microbiology* **2019**, *111*, 6-16, doi:10.1111/mmi.14148.
120. Boguski, M.S.; McCormick, F. Proteins regulating Ras and its relatives. *Nature* **1993**, *366*, 643-654, doi:10.1038/366643a0.
121. Bahn, Y.S.; Sundstrom, P. CAP1, an adenylate cyclase-associated protein gene, regulates bud-hypha transitions, filamentous growth, and cyclic AMP levels and is required for virulence of *Candida albicans*. *J Bacteriol* **2001**, *183*, 3211-3223, doi:10.1128/JB.183.10.3211-3223.2001.
122. Feng, Q.; Summers, E.; Guo, B.; Fink, G. Ras signaling is required for serum-induced hyphal differentiation in *Candida albicans*. *J Bacteriol* **1999**, *181*, 6339-6346, doi:10.1128/jb.181.20.6339-6346.1999.
123. Leberer, E.; Harcus, D.; Dignard, D.; Johnson, L.; Ushinsky, S.; Thomas, D.Y.; Schröppel, K. Ras links cellular morphogenesis to virulence by regulation of the MAP kinase and cAMP signalling pathways in the pathogenic fungus *Candida albicans*. *Molecular microbiology* **2001**, *42*, 673-687, doi:10.1046/j.1365-2958.2001.02672.x.
124. Brown, A.J.; Barelle, C.J.; Budge, S.; Duncan, J.; Harris, S.; Lee, P.R.; Leng, P.; Macaskill, S.; Abdul Murad, A.M.; Ramsdale, M.; et al. Gene regulation during morphogenesis in *Candida albicans*. *Contributions to microbiology* **2000**, *5*, 112-125, doi:10.1159/000060347.
125. Leach, M.D.; Farrer, R.A.; Tan, K.; Miao, Z.; Walker, L.A.; Cuomo, C.A.; Wheeler, R.T.; Brown, A.J.; Wong, K.H.; Cowen, L.E. Hsf1 and Hsp90 orchestrate temperature-dependent global transcriptional remodelling and chromatin architecture in *Candida albicans*. *Nat Commun* **2016**, *7*, 11704, doi:10.1038/ncomms11704.
126. Shapiro, R.S.; Zaas, A.K.; Betancourt-Quiroz, M.; Perfect, J.R.; Cowen, L.E. The Hsp90 co-chaperone Sgt1 governs *Candida albicans* morphogenesis and drug resistance. *PLoS One* **2012**, *7*, e44734, doi:10.1371/journal.pone.0044734.
127. Bachewich, C.; Nantel, A.; Whiteway, M. Cell cycle arrest during S or M phase generates polarized growth via distinct signals in *Candida albicans*. *Mol Microbiol* **2005**, *57*, 942-959, doi:10.1111/j.1365-2958.2005.04727.x.
128. Bachewich, C.; Whiteway, M. Cyclin Cln3p links G1 progression to hyphal and pseudohyphal development in *Candida albicans*. *Eukaryot cell* **2005**, *4*, 95-102, doi:10.1128/EC.4.1.95-102.2005.
129. Chen, C.; Zeng, G.; Wang, Y. G1 and S phase arrest in *Candida albicans* induces filamentous growth via distinct mechanisms. *Molecular microbiology* **2018**, *110*, 191-203, doi:10.1111/mmi.14097.
130. Chen, R.E.; Thorner, J. Function and regulation in MAPK signaling pathways: lessons learned from the yeast *Saccharomyces cerevisiae*. *Biochim Biophys Acta* **2007**, *1773*, 1311-1340, doi:10.1016/j.bbamcr.2007.05.003.
131. Monge, R.A.; Román, E.; Nombela, C.; Pla, J. The MAP kinase signal transduction network in *Candida albicans*. *Microbiology (Reading, England)* **2006**, *152*, 905-912, doi:<https://doi.org/10.1099/mic.0.28616-0>.

132. Román, E.; Arana, D.M.; Nombela, C.; Alonso-Monge, R.; Pla, J. MAP kinase pathways as regulators of fungal virulence. *Trends in microbiology* **2007**, *15*, 181-190, doi:<https://doi.org/10.1016/j.tim.2007.02.001>.
133. Eisman, B.; Alonso-Monge, R.; Román, E.; Arana, D.; Nombela, C.; Pla, J. The Cek1 and Hog1 Mitogen-Activated Protein Kinases Play Complementary Roles in Cell Wall Biogenesis and Chlamydospore Formation in the Fungal Pathogen *Candida albicans*. *Eukaryot cell* **2006**, *5*, 347-358, doi:10.1128/ec.5.2.347-358.2006.
134. Galán-Díez, M.; Arana, D.M.; Serrano-Gómez, D.; Kremer, L.; Casasnovas, J.M.; Ortega, M.; Cuesta-Domínguez, A.; Corbí, A.L.; Pla, J.; Fernández-Ruiz, E. *Candida albicans* beta-glucan exposure is controlled by the fungal CEK1-mediated mitogen-activated protein kinase pathway that modulates immune responses triggered through dectin-1. *Infect Immun* **2010**, *78*, 1426-1436, doi:10.1128/iai.00989-09.
135. Köhler, J.R.; Fink, G.R. *Candida albicans* strains heterozygous and homozygous for mutations in mitogen-activated protein kinase signaling components have defects in hyphal development. *Proceedings of the National Academy of Sciences of the United States of America* **1996**, *93*, 13223-13228, doi:10.1073/pnas.93.23.13223.
136. Hazan, I.; Liu, H. Hyphal tip-associated localization of Cdc42 is F-actin dependent in *Candida albicans*. *Eukaryot cell* **2002**, *1*, 856-864, doi:10.1128/ec.1.6.856-864.2002.
137. Bassilana, M.; Blyth, J.; Arkowitz, R.A. Cdc24, the GDP-GTP exchange factor for Cdc42, is required for invasive hyphal growth of *Candida albicans*. *Eukaryot cell* **2003**, *2*, 9-18, doi:10.1128/ec.2.1.9-18.2003.
138. Chen, H.; Zhou, X.; Ren, B.; Cheng, L. The regulation of hyphae growth in *Candida albicans*. *Virulence* **2020**, *11*, 337-348, doi:10.1080/21505594.2020.1748930.
139. Huang, G. Regulation of phenotypic transitions in the fungal pathogen *Candida albicans*. *Virulence* **2012**, *3*, 251-261, doi:10.4161/viru.20010.
140. Herrero de Dios, C.; Román, E.; Díez, C.; Alonso-Monge, R.; Pla, J. The transmembrane protein Opy2 mediates activation of the Cek1 MAP kinase in *Candida albicans*. *Fungal Genet and Biol* **2013**, *50*, 21-32, doi:<https://doi.org/10.1016/j.fgb.2012.11.001>.
141. Román, E.; Cottier, F.; Ernst, J.F.; Pla, J. Msb2 signaling mucin controls activation of Cek1 mitogen-activated protein kinase in *Candida albicans*. *Eukaryot cell* **2009**, *8*, 1235-1249, doi:10.1128/ec.00081-09.
142. Hope, H.; Schmauch, C.; Arkowitz, R.A.; Bassilana, M. The *Candida albicans* ELMO homologue functions together with Rac1 and Dck1, upstream of the MAP Kinase Cek1, in invasive filamentous growth. *Mol Microbiol* **2010**, *76*, 1572-1590, doi:<https://doi.org/10.1111/j.1365-2958.2010.07186.x>.
143. Hope, H.; Bogliolo, S.; Arkowitz, R.A.; Bassilana, M. Activation of Rac1 by the Guanine Nucleotide Exchange Factor Dck1 Is Required for Invasive Filamentous Growth in the Pathogen *Candida albicans*. *Molecular biology of the cell* **2008**, *19*, 3638-3651, doi:10.1091/mbc.e07-12-1272.
144. Bassilana, M.; Arkowitz, R.A. Rac1 and Cdc42 Have Different Roles in *Candida albicans* Development. *Eukaryot cell* **2006**, *5*, 321-329, doi:10.1128/ec.5.2.321-329.2006.
145. Riggle, P.J.; Andrutis, K.A.; Chen, X.; Tzipori, S.R.; Kumamoto, C.A. Invasive Lesions Containing Filamentous Forms Produced by a *Candida albicans* Mutant That Is Defective in Filamentous Growth in Culture. *Infect Immun* **1999**, *67*, 3649-3652, doi:10.1128/iai.67.7.3649-3652.1999.
146. Levin, D.E. Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the cell wall integrity signaling pathway. *Genetics* **2011**, *189*, 1145-1175, doi:10.1534/genetics.111.128264.
147. Perez, P.; Rincón, S.A. Rho GTPases: regulation of cell polarity and growth in yeasts. *The Biochemical journal* **2010**, *426*, 243-253, doi:10.1042/bj20091823.
148. Xie, J.L.; Grahl, N.; Sless, T.; Leach, M.D.; Kim, S.H.; Hogan, D.A.; Robbins, N.; Cowen, L.E. Signaling through Lrg1, Rho1 and Pkc1 Governs *Candida albicans* Morphogenesis in Response to Diverse Cues. *PLOS Genetics* **2016**, *12*, e1006405, doi:10.1371/journal.pgen.1006405.

149. Biswas, S.; Van Dijk, P.; Datta, A. Environmental Sensing and Signal Transduction Pathways Regulating Morphopathogenic Determinants of *Candida albicans*. *Microbiol Mol Biol Rev* **2007**, *71*, 348-376, doi:10.1128/mmbr.00009-06.
150. Braun, B.R.; Johnson, A.D. Control of Filament Formation in *Candida albicans* by the Transcriptional Repressor *TUP1*. *Science (New York, N.Y.)* **1997**, *277*, 105-109, doi:10.1126/science.277.5322.105.
151. Murad, A.M.; Leng, P.; Straffon, M.; Wishart, J.; Macaskill, S.; MacCallum, D.; Schnell, N.; Talibi, D.; Marechal, D.; Tekai, F.; et al. NRG1 represses yeast-hypha morphogenesis and hypha-specific gene expression in *Candida albicans*. *The EMBO journal* **2001**, *20*, 4742-4752, doi:10.1093/emboj/20.17.4742.
152. Kadosh, D.; Johnson, A.D. Rfg1, a protein related to the *Saccharomyces cerevisiae* hypoxic regulator Rox1, controls filamentous growth and virulence in *Candida albicans*. *Molecular and cellular biology* **2001**, *21*, 2496-2505, doi:10.1128/mcb.21.7.2496-2505.2001.
153. Braun, B.R.; Johnson, A.D. TUP1, CPH1 and EFG1 make independent contributions to filamentation in *Candida albicans*. *Genetics* **2000**, *155*, 57-67.
154. Kadosh, D.; Johnson, A.D. Induction of the *Candida albicans* filamentous growth program by relief of transcriptional repression: a genome-wide analysis. *Molecular biology of the cell* **2005**, *16*, 2903-2912, doi:10.1091/mbc.e05-01-0073.
155. Banerjee, M.; Thompson, D.S.; Lazzell, A.; Carlisle, P.L.; Pierce, C.; Monteagudo, C.; López-Ribot, J.L.; Kadosh, D. UME6, a novel filament-specific regulator of *Candida albicans* hyphal extension and virulence. *Molecular biology of the cell* **2008**, *19*, 1354-1365, doi:10.1091/mbc.e07-11-1110.
156. Braun, B.R.; Kadosh, D.; Johnson, A.D. NRG1, a repressor of filamentous growth in *C. albicans*, is down-regulated during filament induction. *The EMBO journal* **2001**, *20*, 4753-4761, doi:10.1093/emboj/20.17.4753.
157. Jensen, E.C.; Hornby, J.M.; Pagliaccetti, N.E.; Wolter, C.M.; Nickerson, K.W.; Atkin, A.L. Farnesol restores wild-type colony morphology to 96% of *Candida albicans* colony morphology variants recovered following treatment with mutagens. *Genome* **2006**, *49*, 346-353, doi:10.1139/g05-117.
158. Mosel, D.D.; Dumitru, R.; Hornby, J.M.; Atkin, A.L.; Nickerson, K.W. Farnesol concentrations required to block germ tube formation in *Candida albicans* in the presence and absence of serum. *Applied and environmental microbiology* **2005**, *71*, 4938-4940, doi:10.1128/AEM.71.8.4938-4940.2005.
159. Ramage, G.; Saville, S.P.; Wickes, B.L.; López-Ribot, J.L. Inhibition of *Candida albicans* biofilm formation by farnesol, a quorum-sensing molecule. *Applied and environmental microbiology* **2002**, *68*, 5459-5463, doi:10.1128/aem.68.11.5459-5463.2002.
160. Lu, Y.; Su, C.; Unoje, O.; Liu, H. Quorum sensing controls hyphal initiation in *Candida albicans* through Ubr1-mediated protein degradation. *Proceedings of the National Academy of Sciences of the United States of America* **2014**, *111*, 1975-1980, doi:10.1073/pnas.1318690111.
161. Martin, R.; Moran, G.P.; Jacobsen, I.D.; Heyken, A.; Domey, J.; Sullivan, D.J.; Kurzai, O.; Hube, B. The *Candida albicans*-Specific Gene EED1 Encodes a Key Regulator of Hyphal Extension. *PLOS ONE* **2011**, *6*, e18394, doi:10.1371/journal.pone.0018394.
162. Mendelsohn, S.; Pinsky, M.; Weissman, Z.; Kornitzer, D. Regulation of the *Candida albicans* Hypha-Inducing Transcription Factor Ume6 by the CDK1 Cyclins Cln3 and Hgc1. *mSphere* **2017**, *2*, e00248-00216, doi:10.1128/mSphere.00248-16.
163. Carlisle, P.L.; Kadosh, D. *Candida albicans* Ume6, a filament-specific transcriptional regulator, directs hyphal growth via a pathway involving Hgc1 cyclin-related protein. *Eukaryot cell* **2010**, *9*, 1320-1328, doi:10.1128/EC.00046-10.
164. Zheng, X.; Wang, Y.; Wang, Y. Hgc1, a novel hypha-specific G1 cyclin-related protein regulates *Candida albicans* hyphal morphogenesis. *The EMBO journal* **2004**, *23*, 1845-1856, doi:10.1038/sj.emboj.7600195.
165. Lu, Y.; Su, C.; Solis, Norma V.; Filler, Scott G.; Liu, H. Synergistic Regulation of Hyphal Elongation by Hypoxia, CO₂, and Nutrient Conditions Controls the Virulence of *Candida albicans*. *Cell Host & Microbe* **2013**, *14*, 499-509, doi:10.1016/j.chom.2013.10.008.

166. Lu, Y.; Su, C.; Ray, S.; Yuan, Y.; Liu, H. CO₂ Signaling through the Ptc2-Ssn3 Axis Governs Sustained Hyphal Development of *Candida albicans* by Reducing Ume6 Phosphorylation and Degradation. *mBio* **2019**, *10*, doi:10.1128/mBio.02320-18.
167. Hamdy, R.; Soliman, S.S.M.; Alsaadi, A.I.; Fayed, B.; Hamoda, A.M.; Elseginy, S.A.; Hussein, M.I.; Ibrahim, A.S. Design and synthesis of new drugs inhibitors of *Candida albicans* hyphae and biofilm formation by upregulating the expression of *TUP1* transcription repressor gene. *Eur J Pharm Sci* **2020**, *148*, 105327, doi:<https://doi.org/10.1016/j.ejps.2020.105327>.
168. Padmavathi, A.R.; P, S.M.; Das, A.; Priya, A.; Sushmitha, T.J.; Pandian, S.K.; Toleti, S.R. Impediment to growth and yeast-to-hyphae transition in *Candida albicans* by copper oxide nanoparticles. *Biofouling* **2020**, *36*, 56-72, doi:10.1080/08927014.2020.1715371.
169. González-Novo, A.; Correa-Bordes, J.; Labrador, L.; Sánchez, M.; Vázquez de Aldana, C.R.; Jiménez, J. Sep7 is essential to modify septin ring dynamics and inhibit cell separation during *Candida albicans* hyphal growth. *Mol Biol Cell* **2008**, *19*, 1509-1518, doi:10.1091/mbc.e07-09-0876.
170. Sinha, I.; Wang, Y.M.; Philp, R.; Li, C.R.; Yap, W.H.; Wang, Y. Cyclin-dependent kinases control septin phosphorylation in *Candida albicans* hyphal development. *Developmental cell* **2007**, *13*, 421-432, doi:10.1016/j.devcel.2007.06.011.
171. Sudbery, P.E. The germ tubes of *Candida albicans* hyphae and pseudohyphae show different patterns of septin ring localization. *Molecular microbiology* **2001**, *41*, 19-31, doi:<https://doi.org/10.1046/j.1365-2958.2001.02459.x>.
172. Warena, A.J.; Konopka, J.B. Septin function in *Candida albicans* morphogenesis. *Molecular biology of the cell* **2002**, *13*, 2732-2746, doi:10.1091/mbc.e02-01-0013.
173. Au Yong, J.Y.; Wang, Y.M.; Wang, Y. The Nim1 kinase Gin4 has distinct domains crucial for septin assembly, phospholipid binding and mitotic exit. *J Cell Sci* **2016**, *129*, 2744-2756, doi:10.1242/jcs.183160.
174. Li, C.R.; Au Yong, J.Y.; Wang, Y.M.; Wang, Y. CDK regulates septin organization through cell-cycle-dependent phosphorylation of the Nim1-related kinase Gin4. *J Cell Sci* **2012**, *125*, 2533-2543, doi:10.1242/jcs.104497.
175. Wightman, R.; Bates, S.; Amornrattanapan, P.; Sudbery, P. In *Candida albicans*, the Nim1 kinases Gin4 and Hsl1 negatively regulate pseudohypha formation and Gin4 also controls septin organization. *J Cell Biol* **2004**, *164*, 581-591, doi:10.1083/jcb.200307176.
176. Liu, Q.; Han, Q.; Wang, N.; Yao, G.; Zeng, G.; Wang, Y.; Huang, Z.; Sang, J.; Wang, Y. Tpd3-Pph21 phosphatase plays a direct role in Sep7 dephosphorylation in *Candida albicans*. *Molecular microbiology* **2016**, *101*, 109-121, doi:10.1111/mmi.13376.
177. Han, Q.; Pan, C.; Wang, Y.; Wang, N.; Wang, Y.; Sang, J. The PP2A regulatory subunits, Cdc55 and Rts1, play distinct roles in *Candida albicans*' growth, morphogenesis, and virulence. *Fungal genetics and biology : FG & B* **2019**, *131*, 103240, doi:10.1016/j.fgb.2019.103240.
178. Finley, K.R.; Berman, J. Microtubules in *Candida albicans* hyphae drive nuclear dynamics and connect cell cycle progression to morphogenesis. *Eukaryot cell* **2005**, *4*, 1697-1711, doi:10.1128/EC.4.10.1697-1711.2005.
179. Clemente-Blanco, A.; Gonzalez-Novo, A.; Machin, F.; Caballero-Lima, D.; Aragon, L.; Sanchez, M.; de Aldana, C.R.; Jimenez, J.; Correa-Bordes, J. The Cdc14p phosphatase affects late cell-cycle events and morphogenesis in *Candida albicans*. *J Cell Sci* **2006**, *119*, 1130-1143, doi:10.1242/jcs.02820.
180. González-Novo, A.; Vázquez de Aldana, C.R.; Jiménez, J. Fungal septins: one ring to rule it all? *Cent Eur J Biol* **2009**, *4*, 274-289, doi:10.2478/s11535-009-0032-2.
181. Huang, Z.X.; Zhao, P.; Zeng, G.S.; Wang, Y.M.; Sudbery, I.; Wang, Y. Phosphoregulation of Nap1 plays a role in septin ring dynamics and morphogenesis in *Candida albicans*. *mBio* **2014**, *5*, e00915-00913, doi:10.1128/mBio.00915-13.
182. Xie, Y.; Loh, Z.Y.; Xue, J.; Zhou, F.; Sun, J.; Qiao, Z.; Jin, S.; Deng, Y.; Li, H.; Wang, Y.; et al. Orchestrated actin nucleation by the *Candida albicans* polarisome complex enables filamentous growth. *J Biol Chem* **2020**, *295*, 14840-14854, doi:10.1074/jbc.RA120.013890.
183. Caballero-Lima, D.; Hautbergue, G.M.; Wilson, S.A.; Sudbery, P.E. In *Candida albicans* hyphae, Sec2p is physically associated with SEC2 mRNA on secretory vesicles. *Mol Microbiol* **2014**, *94*, 828-842, doi:10.1111/mmi.12799.

184. Caballero-Lima, D.; Kaneva, I.N.; Watton, S.P.; Sudbery, P.E.; Craven, C.J. The spatial distribution of the exocyst and actin cortical patches is sufficient to organize hyphal tip growth. *Eukaryot cell* **2013**, *12*, 998-1008, doi:10.1128/EC.00085-13.
185. Jones, L.A.; Sudbery, P.E. Spitzenkorper, exocyst, and polarisome components in *Candida albicans* hyphae show different patterns of localization and have distinct dynamic properties. *Eukaryot cell* **2010**, *9*, 1455-1465, doi:10.1128/EC.00109-10.
186. TerBush, D.R.; Maurice, T.; Roth, D.; Novick, P. The Exocyst is a multiprotein complex required for exocytosis in *Saccharomyces cerevisiae*. *The EMBO journal* **1996**, *15*, 6483-6494.
187. Crampin, H.; Finley, K.; Gerami-Nejad, M.; Court, H.; Gale, C.; Berman, J.; Sudbery, P. *Candida albicans* hyphae have a Spitzenkorper that is distinct from the polarisome found in yeast and pseudohyphae. *J Cell Sci* **2005**, *118*, 2935-2947, doi:10.1242/jcs.02414.
188. Bishop, A.; Lane, R.; Beniston, R.; Chapa-y-Lazo, B.; Smythe, C.; Sudbery, P. Hyphal growth in *Candida albicans* requires the phosphorylation of Sec2 by the Cdc28-Ccn1/Hgc1 kinase. *The EMBO journal* **2010**, *29*, 2930-2942, doi:10.1038/emboj.2010.158.
189. Martin, R.; Walther, A.; Wendland, J. Ras1-induced hyphal development in *Candida albicans* requires the formin Bni1. *Eukaryot cell* **2005**, *4*, 1712-1724, doi:10.1128/EC.4.10.1712-1724.2005.
190. Zheng, X.D.; Wang, Y.M.; Wang, Y. CaSPA2 is important for polarity establishment and maintenance in *Candida albicans*. *Molecular microbiology* **2003**, *49*, 1391-1405, doi:10.1046/j.1365-2958.2003.03646.x.
191. Akashi, T.; Kanbe, T.; Tanaka, K. The role of the cytoskeleton in the polarized growth of the germ tube in *Candida albicans*. *Microbiology (Reading, England)* **1994**, *140* (Pt 2), 271-280, doi:10.1099/13500872-140-2-271.
192. Hazan, I.; Liu, H. Hyphal tip-associated localization of Cdc42 is F-actin dependent in *Candida albicans*. *Eukaryot Cell* **2002**, *1*, 856-864, doi:10.1128/ec.1.6.856-864.2002.
193. Wolyniak, M.J.; Sundstrom, P. Role of actin cytoskeletal dynamics in activation of the cyclic AMP pathway and HWP1 gene expression in *Candida albicans*. *Eukaryot cell* **2007**, *6*, 1824-1840, doi:10.1128/EC.00188-07.
194. Yokoyama, K.; Kaji, H.; Nishimura, K.; Miyaji, M. The role of microfilaments and microtubules in apical growth and dimorphism of *Candida albicans*. *J Gen Microbiol* **1990**, *136*, 1067-1075, doi:10.1099/00221287-136-6-1067.
195. Pruyne, D.; Bretscher, A. Polarization of cell growth in yeast. *J Cell Sci* **2000**, *113* (Pt 4), 571-585.
196. Kaksonen, M.; Sun, Y.; Drubin, D.G. A pathway for association of receptors, adaptors, and actin during endocytic internalization. *Cell* **2003**, *115*, 475-487, doi:10.1016/s0092-8674(03)00883-3.
197. Anderson, J.M.; Soll, D.R. Differences in actin localization during bud and hypha formation in the yeast *Candida albicans*. *J Gen Microbiol* **1986**, *132*, 2035-2047, doi:10.1099/00221287-132-7-2035.
198. Smith, M.G.; Swamy, S.R.; Pon, L.A. The life cycle of actin patches in mating yeast. *J Cell Sci* **2001**, *114*, 1505-1513.
199. Asleson, C.M.; Bensen, E.S.; Gale, C.A.; Melms, A.S.; Kurischko, C.; Berman, J. *Candida albicans* INT1-induced filamentation in *Saccharomyces cerevisiae* depends on Sla2p. *Molecular biology of the cell* **2001**, *21*, 1272-1284, doi:10.1128/MCB.21.4.1272-1284.2001.
200. Gale, C.A.; Leonard, M.D.; Finley, K.R.; Christensen, L.; McClellan, M.; Abbey, D.; Kurischko, C.; Bensen, E.; Tzafrir, I.; Kauffman, S.; et al. SLA2 mutations cause SWE1-mediated cell cycle phenotypes in *Candida albicans* and *Saccharomyces cerevisiae*. *Microbiology (Reading, England)* **2009**, *155*, 3847-3859, doi:10.1099/mic.0.033233-0.
201. Zeng, G.; Wang, Y.M.; Wang, Y. Cdc28-Cln3 phosphorylation of Sla1 regulates actin patch dynamics in different modes of fungal growth. *Mol Biol Cell* **2012**, *23*, 3485-3497, doi:10.1091/mbc.E12-03-0231.
202. Douglas, L.M.; Martin, S.W.; Konopka, J.B. BAR domain proteins Rvs161 and Rvs167 contribute to *Candida albicans* endocytosis, morphogenesis, and virulence. *Infect Immun* **2009**, *77*, 4150-4160, doi:10.1128/iai.00683-09.
203. Epp, E.; Nazarova, E.; Regan, H.; Douglas, L.M.; Konopka, J.B.; Vogel, J.; Whiteway, M. Clathrin- and Arp2/3-independent endocytosis in the fungal pathogen *Candida albicans*. *mBio* **2013**, *4*, e00476, doi:10.1128/mBio.00476-13.

204. Epp, E.; Walther, A.; Lépine, G.; Leon, Z.; Mullick, A.; Raymond, M.; Wendland, J.; Whiteway, M. Forward genetics in *Candida albicans* that reveals the Arp2/3 complex is required for hyphal formation, but not endocytosis. *Molecular microbiology* **2010**, *75*, 1182-1198, doi:10.1111/j.1365-2958.2009.07038.x.
205. Reijnt, P.; Jorde, S.; Wendland, J. *Candida albicans* SH3-domain proteins involved in hyphal growth, cytokinesis, and vacuolar morphology. *Curr Genet* **2010**, *56*, 309-319, doi:10.1007/s00294-010-0301-7.
206. Asleson, C.M.; Bensen, E.S.; Gale, C.A.; Melms, A.S.; Kurischko, C.; Berman, J. *Candida albicans* INT1-induced filamentation in *Saccharomyces cerevisiae* depends on Sla2p. *Mol Cell Biol* **2001**, *21*, 1272-1284, doi:10.1128/MCB.21.4.1272-1284.2001.
207. Martin, R.; Hellwig, D.; Schaub, Y.; Bauer, J.; Walther, A.; Wendland, J. Functional analysis of *Candida albicans* genes whose *Saccharomyces cerevisiae* homologues are involved in endocytosis. *Yeast* **2007**, *24*, 511-522, doi:10.1002/yea.1489.
208. Bar-Yosef, H.; Gildor, T.; Ramirez-Zavala, B.; Schmauch, C.; Weissman, Z.; Pinsky, M.; Naddaf, R.; Morschhauser, J.; Arkowitz, R.A.; Kornitzer, D. A Global Analysis of Kinase Function in *Candida albicans* Hyphal Morphogenesis Reveals a Role for the Endocytosis Regulator Akl1. *Front Cell Infect Microbiol* **2018**, *8*, 17, doi:10.3389/fcimb.2018.00017.
209. Borth, N.; Walther, A.; Reijnt, P.; Jorde, S.; Schaub, Y.; Wendland, J. *Candida albicans* Vrp1 is required for polarized morphogenesis and interacts with Wal1 and Myo5. *Microbiology (Reading, England)* **2010**, *156*, 2962-2969, doi:10.1099/mic.0.041707-0.
210. Oberholzer, U.; Marcil, A.; Leberer, E.; Thomas, D.Y.; Whiteway, M. Myosin I is required for hypha formation in *Candida albicans*. *Eukaryot cell* **2002**, *1*, 213-228, doi:10.1128/ec.1.2.213-228.2002.
211. Walther, A.; Wendland, J. Polarized hyphal growth in *Candida albicans* requires the Wiskott-Aldrich Syndrome protein homolog Wal1p. *Eukaryot cell* **2004**, *3*, 471-482, doi:10.1128/ec.3.2.471-482.2004.
212. Epp, E.; Nazarova, E.; Regan, H.; Douglas, L.M.; Konopka, J.B.; Vogel, J.; Whiteway, M. Clathrin- and Arp2/3-independent endocytosis in the fungal pathogen *Candida albicans*. *mBio* **2013**, *4*, e00476-00413, doi:10.1128/mBio.00476-13.
213. Bassilana, M.; Hopkins, J.; Arkowitz, R.A. Regulation of the Cdc42/Cdc24 GTPase module during *Candida albicans* hyphal growth. *Eukaryot cell* **2005**, *4*, 588-603, doi:10.1128/EC.4.3.588-603.2005.
214. Ushinsky, S.C.; H Marcus, D.; Ash, J.; Dignard, D.; Marcil, A.; Morschhauser, J.; Thomas, D.Y.; Whiteway, M.; Leberer, E. CDC42 is required for polarized growth in human pathogen *Candida albicans*. *Eukaryot cell* **2002**, *1*, 95-104, doi:10.1128/ec.1.1.95-104.2002.
215. Court, H.; Sudbery, P. Regulation of Cdc42 GTPase activity in the formation of hyphae in *Candida albicans*. *Molecular biology of the cell* **2007**, *18*, 265-281, doi:10.1091/mbc.e06-05-0411.
216. Kozubowski, L.; Saito, K.; Johnson, J.M.; Howell, A.S.; Zyla, T.R.; Lew, D.J. Symmetry-Breaking Polarization Driven by a Cdc42p GEF-PAK Complex. *Curr Biol* **2008**, *18*, 1719-1726, doi:<https://doi.org/10.1016/j.cub.2008.09.060>.
217. Hausauer, D.L.; Gerami-Nejad, M.; Kistler-Anderson, C.; Gale, C.A. Hyphal guidance and invasive growth in *Candida albicans* require the Ras-like GTPase Rsr1p and its GTPase-activating protein Bud2p. *Eukaryot cell* **2005**, *4*, 1273-1286, doi:10.1128/EC.4.7.1273-1286.2005.
218. Pulver, R.; Heisel, T.; Gonias, S.; Robins, R.; Norton, J.; Haynes, P.; Gale, C.A. Rsr1 focuses Cdc42 activity at hyphal tips and promotes maintenance of hyphal development in *Candida albicans*. *Eukaryot cell* **2013**, *12*, 482-495, doi:10.1128/EC.00294-12.
219. Guo, P.P.; Yong, J.Y.A.; Wang, Y.M.; Li, C.R. Sec15 links bud site selection to polarised cell growth and exocytosis in *Candida albicans*. *Scientific Reports* **2016**, *6*, 26464, doi:10.1038/srep26464.
220. Umeyama, T.; Kaneko, A.; Niimi, M.; Uehara, Y. Repression of CDC28 reduces the expression of the morphology-related transcription factors, Efg1p, Nrg1p, Rbf1p, Rim101p, Fkh2p and Tec1p and induces cell elongation in *Candida albicans*. *Yeast* **2006**, *23*, 537-552, doi:10.1002/yea.1373.
221. Bensen, E.S.; Clemente-Blanco, A.; Finley, K.R.; Correa-Bordes, J.; Berman, J. The mitotic cyclins Clb2p and Clb4p affect morphogenesis in *Candida albicans*. *Mol Biol Cell* **2005**, *16*, 3387-3400, doi:10.1091/mbc.e04-12-1081.

222. Loeb, J.D.; Sepulveda-Becerra, M.; Hazan, I.; Liu, H. A G1 cyclin is necessary for maintenance of filamentous growth in *Candida albicans*. *Molecular and cellular biology* **1999**, *19*, 4019-4027, doi:10.1128/mcb.19.6.4019.
223. Bensen, E.S.; Filler, S.G.; Berman, J. A forkhead transcription factor is important for true hyphal as well as yeast morphogenesis in *Candida albicans*. *Eukaryotic cell* **2002**, *1*, 787-798, doi:10.1128/EC.1.5.787-798.2002.
224. Greig, J.A.; Sudbery, I.M.; Richardson, J.P.; Naglik, J.R.; Wang, Y.; Sudbery, P.E. Cell Cycle-Independent Phospho-Regulation of Fkh2 during Hyphal Growth Regulates *Candida albicans* Pathogenesis. *PLoS Pathog* **2015**, *11*, e1004630, doi:10.1371/journal.ppat.1004630.
225. Gutiérrez-Escribano, P.; González-Novo, A.; Suárez, M.B.; Li, C.R.; Wang, Y.; de Aldana, C.R.; Correa-Bordes, J. CDK-dependent phosphorylation of Mob2 is essential for hyphal development in *Candida albicans*. *Molecular biology of the cell* **2011**, *22*, 2458-2469, doi:10.1091/mbc.E11-03-0205.
226. Zheng, X.-D.; Lee, R.T.H.; Wang, Y.-M.; Lin, Q.-S.; Wang, Y. Phosphorylation of Rga2, a Cdc42 GAP, by CDK/Hgc1 is crucial for *Candida albicans* hyphal growth. *The EMBO journal* **2007**, *26*, 3760-3769, doi:10.1038/sj.emboj.7601814.
227. Wang, H.; Huang, Z.X.; Au Yong, J.Y.; Zou, H.; Zeng, G.; Gao, J.; Wang, Y.; Wong, A.H.; Wang, Y. CDK phosphorylates the polarisome scaffold Spa2 to maintain its localization at the site of cell growth. *Molecular microbiology* **2016**, *101*, 250-264, doi:10.1111/mmi.13386.
228. Caballero-Lima, D.; Sudbery, P.E. In *Candida albicans*, phosphorylation of Exo84 by Cdk1-Hgc1 is necessary for efficient hyphal extension. *Molecular biology of the cell* **2014**, *25*, 1097-1110, doi:10.1091/mbc.E13-11-0688.
229. Wang, A.; Raniga, P.P.; Lane, S.; Lu, Y.; Liu, H. Hyphal chain formation in *Candida albicans*: Cdc28-Hgc1 phosphorylation of Efg1 represses cell separation genes. *Molecular and cellular biology* **2009**, *29*, 4406-4416, doi:10.1128/mcb.01502-08.
230. Shapiro, R.S.; Sellam, A.; Tebbji, F.; Whiteway, M.; Nantel, A.; Cowen, L.E. Pho85, Pcl1, and Hms1 signaling governs *Candida albicans* morphogenesis induced by high temperature or Hsp90 compromise. *Curr Biol* **2012**, *22*, 461-470, doi:10.1016/j.cub.2012.01.062.
231. Tripathi, G.; Wiltshire, C.; Macaskill, S.; Tournu, H.; Budge, S.; Brown, A.J. Gcn4 co-ordinates morphogenetic and metabolic responses to amino acid starvation in *Candida albicans*. *The EMBO journal* **2002**, *21*, 5448-5456, doi:10.1093/emboj/cdf507.
232. Gildor, T.; Shemer, R.; Atir-Lande, A.; Kornitzer, D. Coevolution of cyclin Pcl5 and its substrate Gcn4. *Eukaryot Cell* **2005**, *4*, 310-318, doi:10.1128/EC.4.2.310-318.2005.
233. Shemer, R.; Meimoun, A.; Holtzman, T.; Kornitzer, D. Regulation of the transcription factor Gcn4 by Pho85 cyclin PCL5. *Molecular and cellular biology* **2002**, *22*, 5395-5404, doi:10.1128/MCB.22.15.5395-5404.2002.
234. Chapa y Lazo, B.; Bates, S.; Sudbery, P. The G1 cyclin Cln3 regulates morphogenesis in *Candida albicans*. *Eukaryot Cell* **2005**, *4*, 90-94, doi:10.1128/EC.4.1.90-94.2005.
235. Berman, J. Morphogenesis and cell cycle progression in *Candida albicans*. *Curr Opin Microbiol* **2006**, *9*, 595-601, doi:10.1016/j.mib.2006.10.007.
236. Shi, Q.M.; Wang, Y.M.; Zheng, X.D.; Lee, R.T.; Wang, Y. Critical role of DNA checkpoints in mediating genotoxic-stress-induced filamentous growth in *Candida albicans*. *Molecular biology of the cell* **2007**, *18*, 815-826, doi:10.1091/mbc.e06-05-0442.
237. Loll-Kripplleber, R.; d'Enfert, C.; Feri, A.; Diogo, D.; Perin, A.; Marcet-Houben, M.; Bougnoux, M.E.; Legrand, M. A study of the DNA damage checkpoint in *Candida albicans*: uncoupling of the functions of Rad53 in DNA repair, cell cycle regulation and genotoxic stress-induced polarized growth. *Molecular microbiology* **2014**, *91*, 452-471, doi:10.1111/mmi.12471.
238. Andaluz, E.; Ciudad, T.; Gomez-Raja, J.; Calderone, R.; Larriba, G. Rad52 depletion in *Candida albicans* triggers both the DNA-damage checkpoint and filamentation accompanied by but independent of expression of hypha-specific genes. *Molecular microbiology* **2006**, *59*, 1452-1472, doi:10.1111/j.1365-2958.2005.05038.x.
239. Wang, H.; Gao, J.; Li, W.; Wong, A.H.; Hu, K.; Chen, K.; Wang, Y.; Sang, J. Pph3 dephosphorylation of Rad53 is required for cell recovery from MMS-induced DNA damage in *Candida albicans*. *PLoS One* **2012**, *7*, e37246, doi:10.1371/journal.pone.0037246.

240. Feng, J.; Duan, Y.; Sun, W.; Qin, Y.; Zhuang, Z.; Zhu, D.; Sun, X.; Jiang, L. CaTip41 regulates protein phosphatase 2A activity, CaRad53 deactivation and the recovery of DNA damage-induced filamentation to yeast form in *Candida albicans*. *FEMS yeast research* **2016**, *16*, fow009, doi:10.1093/femsyr/fow009.
241. Tscherner, M.; Stappler, E.; Hnisz, D.; Kuchler, K. The histone acetyltransferase Hat1 facilitates DNA damage repair and morphogenesis in *Candida albicans*. *Molecular microbiology* **2012**, *86*, 1197-1214, doi:10.1111/mmi.12051.
242. Bachewich, C.; Thomas, D.Y.; Whiteway, M. Depletion of a polo-like kinase in *Candida albicans* activates cyclase-dependent hyphal-like growth. *Mol Biol Cell* **2003**, *14*, 2163-2180, doi:10.1091/mbc.02-05-0076.
243. Finley, K.R.; Bouchonville, K.J.; Quick, A.; Berman, J. Dynein-dependent nuclear dynamics affect morphogenesis in *Candida albicans* by means of the Bub2p spindle checkpoint. *J Cell Sci* **2008**, *121*, 466-476, doi:10.1242/jcs.015172.
244. Martin, R.; Walther, A.; Wendland, J. Deletion of the dynein heavy-chain gene DYN1 leads to aberrant nuclear positioning and defective hyphal development in *Candida albicans*. *Eukaryot Cell* **2004**, *3*, 1574-1588, doi:10.1128/EC.3.6.1574-1588.2004.
245. Bai, C.; Ramanan, N.; Wang, Y.M.; Wang, Y. Spindle assembly checkpoint component CaMad2p is indispensable for *Candida albicans* survival and virulence in mice. *Molecular microbiology* **2002**, *45*, 31-44, doi:10.1046/j.1365-2958.2002.02995.x.
246. Atir-Lande, A.; Gildor, T.; Kornitzer, D. Role for the SCFCDC4 ubiquitin ligase in *Candida albicans* morphogenesis. *Mol Biol Cell* **2005**, *16*, 2772-2785, doi:10.1091/mbc.e05-01-0079.
247. Shieh, J.C.; White, A.; Cheng, Y.C.; Rosamond, J. Identification and functional characterization of *Candida albicans* CDC4. *J Biomed Sci* **2005**, *12*, 913-924, doi:10.1007/s11373-005-9027-9.
248. Li, W.J.; Wang, Y.M.; Zheng, X.D.; Shi, Q.M.; Zhang, T.T.; Bai, C.; Li, D.; Sang, J.L.; Wang, Y. The F-box protein Grr1 regulates the stability of Ccn1, Cln3 and Hof1 and cell morphogenesis in *Candida albicans*. *Molecular microbiology* **2006**, *62*, 212-226, doi:10.1111/j.1365-2958.2006.05361.x.
249. Chou, H.; Glory, A.; Bachewich, C. Orthologues of the anaphase-promoting complex/cyclosome coactivators Cdc20p and Cdh1p are important for mitotic progression and morphogenesis in *Candida albicans*. *Eukaryot Cell* **2011**, *10*, 696-709, doi:10.1128/EC.00263-10.
250. Gao, J.; Wang, H.; Li, Z.; Wong, A.H.; Wang, Y.Z.; Guo, Y.; Lin, X.; Zeng, G.; Liu, H.; Wang, Y.; et al. *Candida albicans* gains azole resistance by altering sphingolipid composition. *Nat Commun* **2018**, *9*, 4495, doi:10.1038/s41467-018-06944-1.
251. Segal, E.S.; Gritsenko, V.; Levitan, A.; Yadav, B.; Dror, N.; Steenwyk, J.L.; Silberberg, Y.; Mielich, K.; Rokas, A.; Gow, N.A.R.; et al. Gene Essentiality Analyzed by In Vivo Transposon Mutagenesis and Machine Learning in a Stable Haploid Isolate of *Candida albicans*. *mBio* **2018**, *9*, doi:10.1128/mBio.02048-18.