

Review

Tackling dysfunction of mitochondrial bioenergetics in the brain by *omics* technologies

Paola Zanfardino¹, Stefano Doccini², Filippo M. Santorelli^{2,*} and Vittoria Petruzzella^{1,*}

¹University of Bari Aldo Moro, Department of Medical Basic Sciences, Neurosciences and Sense Organs, Bari, 70124 Italy;

²IRCCS Fondazione Stella Maris, Calambrone, 56128 Pisa, Italy;

*Correspondence: filippo3364@gmail.com and vittoria.petruzzella@uniba.it

Abstract: Oxidative phosphorylation (OxPhos) is the basic function of mitochondria although the landscape of mitochondrial functions is continuously growing to include more aspects of cellular homeostasis. Thanks to the application of *-omics* technologies to the study of the OxPhos system, novel features emerge from the cataloging of novel proteins as mitochondrial thus adding details to the mitochondrial proteome and defining novel metabolic cellular interrelations, especially in the human brain. We focussed on the diversity of bioenergetics demand and different aspects of mitochondrial structure, functions, and dysfunction in the brain. Definition as ‘mitoexome’, ‘mitoproteome’ and ‘mitointeractome’ have entered the field of ‘mitochondrial medicine’. In this context, we reviewed several genetic defects that hamper the last step of aerobic metabolism mostly involving the nervous tissue as one of the most prominent energy-dependent tissues and, as consequence, as a primary target of mitochondrial dysfunction. The dual genetic determination of the OxPhos complexes is one of the reasons for the complexity of the genotype-phenotype correlation when facing human diseases associated with mitochondria defects; clinically, are characterized by extremely heterogeneous symptoms, ranging from organ-specific to multisystemic dysfunction with different clinical courses. Finally, we briefly discuss the future directions of the multi-omics study of human brain disorders.

Keywords: mitochondria; mitochondrial DNA; nervous tissue; OxPhos complexes; bioenergetics; genomics; proteomics; mitochondrial diseases

1. Introduction

The *panoply* of mitochondrial functions reflects on highly heterogeneous clinical presentations when an error in a mitochondrial protein or function occurs. Mitochondria are dynamic and mobile organelles representing a hub where exchange of information among the nucleus and other cellular compartments takes place to modulate energy production and metabolites provision to the cell’s specific needs and nutrient availability [1]. The basic function of mitochondria is the generation of more than 90% of cellular energy via the oxidative phosphorylation (OxPhos) system [2] but, in addition, they play many fold roles in the different types of cells: compartmentalize metabolites for the maintenance of redox homeostasis; function as centers for metabolic waste management [3]; surveil calcium homeostasis [4]; initiate caspase-dependent apoptosis and other intermediate cellular stress response [5]; provide sulfur metabolism and iron-sulfur cluster biogenesis [6,7]; house the synthesis of cardiolipin, steroids, quinone, and heme [8,9]; breakdown fatty acids through β -oxidation; and serve as a metabolic platform for the TCA and urea cycles [10]. All these functions include homeostatic regulation of organelle morphology and dynamics [11], quality control [12], and participation in the immune response [13,14]. Alteration of each of the above functions and activities can affect differently according to

the specificity of the organ and cell type, but alteration of mitochondrial energy production can impact tissues with the highest energy requirements such as the nervous either central (CNS) and peripheral (PNS) [15,16].

The term ‘mitochondrial medicine’ categorizes the ample array of clinical presentations associated with all types of mitochondrial defects having directly or secondarily defect of one or several mitochondrial functions although ‘mitochondrial diseases’ traditionally indicate dysfunction of the OxPhos system [6,17]. The direct link between human disease and the genetic alteration of a mitochondrial function has taken a breakthrough with the application of *-omics* technologies (*i.e.*, genomics, transcriptomics, proteomics, metabolomics, and epigenomics, etc.). Rapidly, high-throughput *omics* techniques — that is detection of massive differences in a multitude of molecular constituents in organisms supported by sophisticated bioinformatics tools — have allowed progress in cataloging the predicted human mitochondrial proteins thus revealing new details and providing clues to elucidating still unknown basic aspects of mitochondrial structure and function. These novel high-throughput techniques have enhanced the final diagnosis of several mitochondrial disorders. This is a very relevant aspect, especially considering that mitochondrial diseases individually are rare but are probably the most frequent genetic disorder in adults (incidence of 1 in 5000 live births) [18]. More recently, genome editing technology applied to neural cultures and cerebral organoids generated from patients-derived iPSCs is revolutionizing the landscape and offering new opportunities for understanding the pathogenetic effects of mutations in nervous tissue.

This review aims to focus on the dysfunction of OxPhos defects mostly in the nervous system to highlighting the contributions of powerful omics technologies to mitochondrial medicine to land from the laboratory to the clinic.

2. Mitoxome, mitochondrial proteome, and mitointeractome

Before Next-Generation Sequencing (NGS) improved our understanding of how mutations cause diseases, first attempts to identify the mitochondrial proteome were based on ‘cyberscreening’ of available genome databases. This allowed the discovery of few human mitochondrial genes presenting orthologs in lower eukaryotes. An example of the cyberscreening strategy used *Saccharomyces cerevisiae* proteins as ‘probes’ to identify *BCS1*, *PET112*, *SCO1*, *COX15*, and *COX11*, five yeast genes that present orthologs (respectively *BCS1L*, *GATB*, *SCO1*, *COX15*, and *COX11*) in human [19]. Except for *COX11*, a COX-assembly, all genes have been implicated in mitochondrial diseases [OMIM 603647.0001-603647.0013; OMIM 603645.0001-603645.0002; OMIM 603644.0001-603644.0002; OMIM 603646.0001-603646.0004], see paragraphs 4.3 and 4.4. To date, whole-exome (WES) and whole-genome (WGS) resequencing have dramatically enhanced the ability to identify the underlying gene mutations in patients with isolated or multiple mitochondrial respiratory chain complex defects [20,21]. The collection of mt genes and coding exons of the 1,034 nuclear genes encoding the human mitochondrial proteome is defined as “*MitoExome*” [22,23]. This multigene panel is useful in performing targeted resequencing of the OxPhos nuclear genes because it includes not only the 77 nuclear structural OxPhos subunits and the 37 mtDNA genes [24] but also genes for mitochondrial proteins either already known or not to be associated with a specific mitochondrial disease, including assembly factors and electron carriers’ genes which represent a large fraction of the overall mitochondrial genes that can cause mitochondrial dysfunction [21]. Application of *MitoExome* resequencing provides novel mutation candidates, enables the discovery of unusual clinical variants [25,26] and new clinical phenotypes [26] (Figure 1). Furthermore, the integration of *MitoExome* sequencing with the study of mitochondrial proteome potentiates the detection of variants causing protein destabilization and/or aberrantly low expression [27].




Figure 1. Omics strategies advance in understanding mitochondrial function and dysfunction in brain disorders related to OxPhos gene mutations. Mitochondrial bioenergetics involves activities whose function and structure have been deeply elucidated by *omics* technologies; **(a)** The introduction of high-resolution technologies has been resolute to deepen the structure of the respiratory chain complexes and supercomplexes; **(b)** Quantitative proteomics *e.g.* LC-MS/MS enable the identification and quantification of mitoproteins and provide large amounts of data. Through Network-based approaches analyzing protein-protein interactions, the huge amount of information allows the discovery of novel accessory subunits and assembly factors of the five multi-subunit enzyme complexes; **(c)** *MitoExome* resequencing enhances the ability to identify gene mutations in patients and provides novel gene mutation candidates for diagnostic screening; **(d)** Novel multi-omics analysis, based on single-cell *omics*, is applied to two-dimensional (2D) neural cultures and three-dimensional (3D) cerebral organoids generated from patients-derived iPSCs that can be engineered by CRISPR/Cas9. **Abbreviations:** LC-MS/MS: Liquid Chromatography with tandem mass spectrometry; PPIs: Protein-protein Interactions; nDNA: nuclear DNA; mtDNA: mitochondrial DNA; iPSCs: Induced Pluripotent Stem cells.

Biochemical and ultrastructural characterizations have enhanced the heterogeneity of mitochondria in their function, trafficking patterns, lifespan, and morphology across cell types and different cellular compartments. Different tissues, cell types, and cellular states have unique signatures of protein localization to mitochondria. In the proteomic comparison of the mitochondrial proteins, almost half are found as *core* components in virtually all tissues, whereas the remaining are tissue-specific [28,29]. The study of mitochondrial proteome starts with the isolation of mt compartment from cells and tissues and stands behind the availability of methodologies to isolate pure mitochondria from different sources to define exactly the function of each protein in each cell type of the human body [30]. The performance of proteomics analysis is driven by the reduction of sample complexity, enhancement of mass spectrometry (MS) power of resolution, and the possibility to reduce the contamination of the sample with non-mitochondrial proteins owed to chemical and physical similarities between mitochondria and other cellular components (*e.g.* lysosomes). Since the initial rough estimates, it has been suggested that the mammalian mitochondrial proteome encompassed about 1000 -1500 distinct proteins – including the 13 mtDNA-encoded proteins [24] – that represents an important subset of the ~20,000 distinct mammalian proteins [31,32] (**Figure 2**).




Figure 2. Functional diversity of mitochondrial proteins and bioenergetics consequences of OxPhos system dysfunction. The mammalian mitochondrial proteome includes both mitochondrial and nuclear DNA encoded proteins. Most of the proteins required for the various activities in which mitochondria are involved are encoded by the nuclear genome, whereas the mitochondrial energy-producing system, i.e., the OxPhos complexes, has either mitochondrial DNA (mtDNA) and nuclear DNA (nDNA) encoded components. The enlarged mitochondrion shows most of the bioenergetics consequences (indicated by the red bolt lightning) of genetic defects involving the OxPhos complexes. **Abbreviations:** IP3: Inositol Trisphosphate; IP3R: Inositol Trisphosphate Receptor; VDAC: Voltage-dependent anion channel; ETC: Electron Transport Chain; TCA: Tricarboxylic Acid Cycle; ATP: Adenosine triphosphate.

Quantitative two-dimensional (2D) gels of highly purified mitochondria estimated ~1500 distinct spots [33], a number higher than the ~1000 distinct protein products encoded by the genomes of alpha-proteobacteria, which are the closest living relatives of modern-day mitochondria [34]. Several databases have been used to integrate the experimental data with bioinformatic predictions based on mitochondrial localization or interaction. For example, the *MitoProteome* is an object-related database developed at the UCSD Supercomputer Center, which contains information on mitochondria-localized proteins [35,36]. Each entry in the MitoProteome corresponds to a gene encoding a protein that is localized within mitochondria and its basic information, along with annotations of isoforms, splice variants, and functions of the corresponding protein. To date, the most comprehensive study elucidating the mitochondrial proteome of different mammalian tissues is represented by the *MitoCarta inventory* [29,37]. This catalog combines multiple experimental and computational approaches *i.e.*, mass spectrometry (MS) analysis of mitochondria isolated from 14 mouse tissues, large-scale GFP-fusion microscopy analysis, and bioinformatics using data mining, prediction, evolutionary conservation, and a Bayesian integration of seven additional data sources. The first release was represented by MitoCarta1.0 (<http://www.broadinstitute.org/pubs/MitoCarta/>) which contained about 1000 distinct gene *loci* (Pagliarini et al., 2008). Updated in 2016, MitoCarta 2.0 listed about

1200 genes [37]. Another dedicated database that collected, curated, and annotated information on mitochondrial proteins is the *MitoMiner database* (<http://mitominer.mrc-mbu.cam.ac.uk/>) [38] (version 4.0, 2018). It is based on literature and proteomics data based on both LC-MS and 2D gel studies, antibody staining, other subcellular localization data, and provides a collective score for each protein's probability to have the mitochondrial association. MitoMiner records mitochondrial proteins from 12 organisms [38]. Using the data contained within MitoMiner, the *Integrated Mitochondrial Protein Index* (IMPI) was also developed (<http://www.mrc-mbu.cam.ac.uk/impi>). IMPI version Q2 (2018) contains 1626 human genes that encode mitochondrially localized proteins, 1184 known to be mitochondrial and 442 predicted to be mitochondrial. The large amount of information provided by mito-databases as MitoMiner 4.0 v2018 JUN (<http://mitominer.mrc-mbu.cam.ac.uk>), makes it possible to define different score systems for mitochondrial confidence combining data from various mitochondrial and functional annotation databases. These strategies allow increasing the stringency of protein accepted as inherently mitochondrial [39]. An exhaustive list of the major data sources loaded with the latest version and links to the relevant resources is reported in the Data Sources section of the Mitominer (<https://mitominer.mrc-mbu.cam.ac.uk/release-4.0/dataCategories.do>).

More recent advances in the experimental proteomic approaches, specifically in labeling and MS methods, have further expanded and defined the known mitochondrial proteome and have simultaneously revealed the sub-mitochondrial localization of many of them [40,41]. A novel spatial proteomics pipeline demonstrated that many proteins cannot be classified to a single localization as they either transit between compartments or carry out their functional role(s) in multiple locations [41]. The redundant functions, or functions affecting multiple cellular processes, rendered difficult the study and it was estimated that about ~20% of mitochondrial proteins remained uncharacterized [42].

Along with technological progress that has enabled the discovery of approximately 196,111 human proteins [based on Swiss-Prot and TrEMBL entries in Universal Protein Resource (The UniProt Consortium), as of April 15, 2021], derives the challenge of identifying a large amount of potential protein-protein interactions (PPIs). An example of the network-based approaches analyzing protein-protein interaction is represented by *MitoInteractome*, a web-based portal containing 6,549 protein sequences extracted from SwissProt (<http://www.expasy.ch/sprot/>), MitoP (<http://www.mitop.de:8080/mitop2/>), MitoProteome (<http://www.mitoproteome.org/>), HPRD (<http://www.hprd.org>) and Gene Ontology database (<http://www.geneontology.org>). This enables the elucidation of integrative mitochondrial functions and can expedite the discovery of novel interactions which otherwise may have been missed using traditional experimental techniques. *MEGADOCK* [43,44], a structure-based PPI prediction method, was first developed and then followed the MEGADOCK-Web-Mito database which is a PPI prediction data archive, that includes prediction results for protein pairs of 654 mitochondria-related human proteins [45]. All these approaches have been key in the study of PPI as a means to infer functions for uncharacterized proteins and to enable the discovery of novel protein, *e.g.* several complex I assembly factors [46,47] (**Figure 1**).

3. Diversity of bioenergetics demand in the brain

The brain relies on glucose metabolism for ATP generation and many other activities and an inappropriate supply of either glucose or oxygen degrades brain function. The high energy request of the brain is owed to neuronal signaling that accounts for ~70% of the calculated energy expense, whereas non-signaling activities (metabolism of protein and phospholipids, oligonucleotide turnover, axonal transport, mitochondrial proton leak, actin cytoskeleton remodeling, etc.) expend an estimated 30%. Excitatory neurons consume ~80–85% of calculated ATP use, whereas inhibitory neurons and glia account for 15–20% [48]. Gray matter has regionally heterogeneous metabolic rates that are considerably higher than white matter [49] that have a higher fraction of non-signaling energy demands than gray matter. In addition to ATP generation, glucose is also the precursor

for the synthesis of many compounds within the brain, including neurotransmitters and neuromodulators. For these reasons, mitochondria are quite heterogeneous as anatomical localization, activity, and metabolism at regional, cellular, subcellular levels and during differentiation, when the upregulation of mitochondrial metabolism is the basis of cell proliferation in neuronal stem cells and progenitor cells. Although different brain regions contain about half mitochondria to the heart, brain mitochondria are qualitatively different to sustain the high metabolic demand that requires, for instance, tight cooperation between neurons and astrocytes [50]. Astrocytes provide crucial metabolic and structural support [50-52] and are key players in neurotransmission [53,54] and behavior [55,56]. The ATP used by neurons is supplied by the OxPhos process, whereas most energy needs of astrocytes are satisfied by glycolysis [57,58]. The mitochondrial ATP production per molecule of glucose oxidized is ~16 times more than glycolysis. The survival of neurons requires OxPhos [50] and in mature neurons, the local ATP supply provided by mitochondria is used to regulate axonal and dendritic development, axonal regeneration, as well as contributing to synaptic transmission and plasticity. The different energy metabolisms of the two cell types are closely coupled, with astrocytes releasing the glycolytic end-product, lactate, which is used by neighboring neurons to drive OxPhos [59,60].

An example of heterogeneity of mitochondria in metabolic enzyme diversity has been provided by a study comparing the mitochondrial proteome of the three major cerebellar cell types: Granule cells (GC), the most abundant excitatory neuron; Purkinje cells (PC), the major inhibitory neuron of the cerebellum and astrocytes. [61]. In the adult cerebellum, ~15% of the annotated mitochondrial proteome was shown to be differentially regulated among the three cell types. Fatty acids were more efficiently metabolized by astrocytic than neuronal mitochondria due to the enrichment of two beta-oxidation enzymes, *i.e.*, short-chain-specific acyl-coenzyme A dehydrogenase and carnitine palmitoyl-transferase 1a, a rate-limiting enzyme in long-chain fatty acids oxidation [61]. Notably, the astrocytic mitochondrial proteome contained a substantial enrichment of peroxisomal proteins, some of which are known to have dual-targeting (*i.e.* catalase) [62] or tether to mitochondria (*i.e.* Eci2 and Pex11b). In the same work, the mitochondrial calcium uniporter (*MCU*) [4,61,63] and its regulators were detected mostly in GC [61]. Recent studies suggest that the markedly different modes of ATP production in the neurons and astrocytes reside also in the supra-organization of the mitochondrial respiratory chain in supercomplexes (*see* 4.6 paragraph) able to regulate different rates of respiration and mitochondrial ROS production [64].

The brain mitochondrial proteome is not a *unicum* also when considering synaptic and non-synaptic mitochondria (sMito and nsMito). Proteomic profiling of sMito *vs* nsMito revealed mitochondrial complex I as an upstream regulator of degenerative processes associated with a high range of age-related neuropathologies characterized by synaptic dysfunction [65]. An accurate analysis of quantitative proteomics was performed to differentiate sMito and nsMito in mouse brain mitochondria by *stable isotope labeling with amino acids in cell culture (SILAC)*. In SILAC, cells are differentially labeled by growing them in a 'light' medium, containing normal amino acids, or a 'heavy' medium, containing a stable isotope [66]. Significant differential expression was shown for 522 proteins involved in several pathways including the OxPhos system, mitochondrial fission/fusion, calcium transport, and mtDNA replication and maintenance. Lower levels of Pyruvate dehydrogenase (PDH) subunits in the synapse to other parts of the cell and reduced expression of complex I, II, and IV (except for COX4I2) suggested decreased bioenergetic function of sMito compared to nsMito [66]. Consistent with this finding, sMito exhibited increased age-associated mtDNA deletions and reduced levels of TFAM and mtSod2, suggesting a reduced ability of sMito to withstand ROS, thus providing insights into synaptic mitochondrial susceptibility to damage [66] (**Figure 3**).




Figure 3. Diversity of mitochondrial proteome in synaptic and non-synaptic mitochondria. Synaptic mitochondria show defects in energy metabolism due to low levels of Pyruvate dehydrogenase (PDH) subunits (A), reduced expression of complex I, II, and IV (B), and increased susceptibility to damage (increased mtDNA deletions) (C), compared to non-synaptic mitochondria. **Abbreviations:** ADP: Adenosine diphosphate; ATP: Adenosine triphosphate.

The extreme heterogeneity of mitochondria activities and functioning has been recently shown by a novel and fine imaging approach that specifically allows to label and monitor mitochondrial translation products for microscopic fluorescent imaging. In neuronal cultures, mitochondrial translation was monitored in axonal and dendritic mitochondria as well as in pre- and post-synaptic regions of neurites by specifically labeling the peptides newly synthesized by mitochondrial ribosomes revealing that not all mitochondria translate to the same extent in different cell types [67]. Finally, the fundamental role of mitochondria during neurogenesis has been recapitulated in the cellular and organoid model of Leigh syndrome (LS), a severe manifestation of mitochondrial disease in children [68]. Mutations in *SURF1*, a complex IV assembly gene, cause neuronal impairment because of defective metabolic programming of neural progenitor cells (NPCs) that prevents the establishment of neuronal morphogenesis. Using CRISPR/Cas9 engineered *SURF1* patient-derived iPSCs, a human model of LS was developed. Single-cell RNA-sequencing and multi-omics analysis revealed compromised neuronal morphogenesis in mutant 2D neural cultures and 3D brain organoids (Figure 2). The defects already emerged at the level of NPCs, which were unable to shift toward OxPhos and retained a proliferative glycolytic state that fails to instruct neuronal morphogenesis. Interestingly, gene augmentation and *PGC1A* induction via Bezafibrate treatment inducing mitochondria biogenesis supported the metabolic programming of LS NPCs, leading to restored neuronal morphogenesis [68].

4. Structure, assembly, and disorders of bioenergetics complexes

The development of mito-omics-based approaches has been crucial in understanding the functional and bioenergetic consequences of mutations responsible for the onset of primary mitochondrial diseases. The OxPhos is the process by which mitochondria produce the ATP needed by the cells. The reactions are performed by five multimeric enzyme complexes (EC): Complex I (EC 1.6.5.3) or NADH: ubiquinone reductase, CI, 45 subunits; Complex II (EC 1.3.5.1) or succinate dehydrogenase, CII, 4 subunits; Complex III (EC 1.10.2.2) or quinol-cytochrome c (cyt c) reductase, CIII, 10 subunits; Complex IV (EC 1.9.3.1) or cyt c oxidase (COX), CIV, 13 subunits; Complex V (EC 3.6.14) or FoF1-ATPase, CV, 16 subunits; and two-electron transport carriers, namely, ubiquinone (coenzyme Q, CoQ) and cyt c [69]. Reactions catalyzed by CI, CIII, and CIV result in the release of protons in the inner membrane space, thereby creating the proton gradient needed for ATP synthase activity. The correct function of the OxPhos system depends on the concerted action of several chaperones and other assembly factors that play essential roles in the formation, regulation, and stability of the five complexes and the mobile electron carriers, and nucleotide transporters [70]. *Early*-stage assembly factors, that have critical roles in the structural assembly of individual subunits and subcomplexes, and *late*-stage accessory factors, also known as the LYRM (leucine-tyrosine-arginine motif) proteins, that regulate the incorporation and/or activation of final subunits and/or co-factors (*i.e.* Fe-S clusters) provide optimal assembly of CI, CII, CIII, and CV. The human mitoproteome contains at least 12 LYRM proteins [71].

The OxPhos system is under a dual genetic control: 13 subunits are of mtDNA origin [24] and the remaining are encoded by the nuclear DNA (nDNA) [72]. MtDNA is a small circular genome [24] that encodes only 13 mitochondrial proteins, 22 mt-tRNAs, and 2 mt-rRNAs. Hence, the nuclear-encoded mitochondrial proteome requires sophisticated machinery for the transport into mitochondria [73-75]. Over the last years, a growing number of human proteins involved in mtDNA replication, and expression have been identified owing to the study of primary mitochondrial diseases. The coordination between the two genomes is crucial for mtDNA integrity, copy number regulation, and mitochondrial protein synthesis because mutations in nuclear genes encoding proteins for mtDNA replication and maintenance may affect its integrity and properties [76]. Dedicated reviews on these topics, including also the specific mechanisms regulating mtDNA replication [77], transcription [78], and translation [79,80] are available elsewhere.

Genetically, the mitochondrial diseases associated with the OxPhos system are split into two broad genetic categories: disorders due to mutations in the mtDNA, observing the rules of mitochondrial genetics; disorders due to mutations in the nDNA, transmitted as a Mendelian trait [6,81]. To date, mutations in both mitochondrial and nuclear genomes have been reported to cause mitochondrial disease manifesting with characteristic leukoencephalopathy and other clinical phenotypes either multisystemic or with single tissue involvement [82-84].

Since the first descriptions of mtDNA mutations [85-87], the number of mutations has been growing more and more until it counts over 1000 heteroplasmic rearrangements (large deletions/duplications) (<http://mitobreak.portugene.com>), and over 350-point mutations possibly pathogenic among the 700 variants reported, which affect all mtDNA genes (<https://www.mitomap.org>). A few major clinical phenotypes in adults have been recently reviewed [88]: LHON [87,89]; Neuropathy, ataxia, retinitis pigmentosa (NARP)/maternally inherited Leigh syndrome (MILS) [90,91]; Maternally inherited non-syndromic deafness, associated or not with aminoglycosides use [92]; Myoclonus, epilepsy, ragged-red-fibers syndrome (MERRF) [93,94]; Mitochondrial encephalopathy, lactic acidosis stroke-like syndrome (MELAS) [95,96]; Chronic progressive external ophthalmoplegia (CPEO) spectrum [85]; Kearns-Sayre syndrome (KSS) [97,98] and Pearson's syndrome [99,100]. LHON and NARP/MILS are disorders that affect single OxPhos complex, complex I in LHON [101], and complex V in NARP/MILS [102], respectively. All these phenotypes are maternally inherited, displaying the hallmarks of mitochondrial diseases

including variability of the phenotype, incomplete penetrance, and overlapping clinical features. The exception is represented by CPEO/KSS/Pearson associated with single mtDNA deletions, which are mostly sporadic [103,104].

Herein, we will provide some rapid information on structure, assembly, and disorders related to each of the OxPhos complexes.

4.1. NADH–Ubiquinone Oxidoreductase - Complex I

NADH–Ubiquinone Oxidoreductase (Complex I, CI) couples the electron transfer of the two electrons derived from NADH oxidation to the ubiquinone with the translocation of four protons into the intermembrane space (IMS) [105-107]. Most of the molecular studies of mitochondrial diseases have focused on Complex I, which is the largest and most complicated among the respiratory complexes. Of 45 subunits, seven are encoded by the mtDNA (MT-ND1-6 and MT-ND4L), and the remaining, including the dual copy of the acyl-carrier protein NDUFAB1 [108], are encoded by nDNA [108,109]. Structurally, CI is an L-shaped complex that is composed of two domains: the hydrophilic head protruding into the matrix and the hydrophobic part within the inner mitochondrial membrane (IMM) [110]. Fourteen core subunits, conserved from bacteria to humans, perform catalytic activities [108,111,112]. Seven core subunits in the hydrophilic arm contain the redox-active centers: a non-covalently bound FMN and seven Fe–S clusters [113]. All the seven mtDNA-encoded CI subunits are in the hydrophobic arm and form the proton channels [109]. The remaining 30 subunits are ‘supernumerary’ but important for assembly and stability [114]. Most accessory subunits are only found in eukaryotic complex I. A notable exception is represented by subunits NDUFS4, NDUFS6, and NDUFA12 that are already present in complex I from α -proteobacteria [115].

The complete mammalian CI structure has been elucidated [105,116] and determined by X-ray crystallography [111,117] and cryo-EM [112,118-123]. It is organized in six independent modules, N, Q, ND1/PP-a, ND2/PP-b, ND4/PD, and ND5/PD-b, that, assisted by specific assembly factors, are incorporated in a specific order [124]. The overall L-shaped CI structure derives from the assembly of the N- and Q modules in the peripheral arm, and ND1, ND2, ND4, and ND5 modules in the P part of the membrane arm forming, at the hinge between the two arms, the channel of the CoQ binding site (Q-module) [114,113]. The *N module*, situated at the head of the hydrophilic part, contains the NADH-binding site and a flavomononucleotide (FMN) cofactor which oxidizes NADH to release 2 electrons [125]; the *Q module* for Q reduction, situated in the hydrophilic arm, contains eight Fe–S clusters where electrons flow to reach ubiquinone [125]. The N and Q modules form the peripheral arm containing the 7 “core” subunits (NDUFV1, NDUFV2, NDUFS1, NDUFS2, NDUFS3, NDUFS7, and NDUFS8) whereas the 30 accessory subunits are necessary to stabilize the enzyme [126]. The *P-module* constitutes the membrane arm and is composed of the seven mtDNA-encoded proteins: ND1- ND4, ND4L, ND5, and ND6, involved in proton translocation [127]. Specific factors assisting the preassembly of the modules and the role of protein import machinery are summarized in **Table 1**.

Table 1. Complex I assembly factors with interacting module/function, associated clinical phenotypes, and references. Adapted from [70], [47] and [162].

Assembly factors	CI interacting module/function	Associated clinical phenotypes	References
ACAD9	ND2/PP-b module Component of MCIA complex, necessary for insertion of ND2	Cardiorespiratory depression, hypertrophic cardiomyopathy, encephalopathy, and severe lactic acidosis	[128,129]
ECSIT	ND2/PP-b module	-	[130]

	Component of MCIA complex, necessary for insertion of ND2		
FOXRED1	ND4/PD module	Leigh syndrome, congenital lactic acidosis, athetoid movements of the limbs in early childhood, hypotonia and cerebellar atrophy, mitochondrial respiratory chain deficiency associated with Leigh syndrome, encephalomyopathy, or ataxia	[125,131,132]
ATP5SL/DMAC2	ND4/PD module	-	[133]
TMEM70	ND4/PD module	Mitochondrial complex V deficiency, nuclear type 2	[134,135]
NDUFAF1	N module, ND1 Component of MCIA complex, necessary for insertion of ND2	Hypertrophic cardiomyopathy, developmental delay, lactic acidosis, hypotonia, and Wolff-Parkinson-White syndrome	[136,137]
NDUFAF2	N module. Stabilization of pre-CI or 830 kDa subcomplex	Ataxia, lethargy, nystagmus, hypotonia, optic atrophy, and episodic respiratory insufficiency, generic encephalopathic syndromes, or Leigh syndrome	[138]
NDUFAF3 (C3ORF60)	Q module	Macrocephaly, weak cry, no eye contact, wide anterior fontanel and axial hypotonia	[139]
NDUFAF4 (C6ORF66)	Q module	Severe encephalopathy and antenatal cardiomyopathy	[140]
NDUFAF5 (C20ORF7)	Not known Catalyze hydroxylation of NDUF57 and dimethylation of NDUF52 of the Q module	Facial dysmorphism, progressive lactic acidosis and neurological defects, severe early-onset encephalopathy	[141,142]
NDUFAF6	Not known. Maintain a normal level of mt-ND1 subunit	Fatal infantile lactic acidosis, neonatal myopathy, encephalopathy, and lactic acidosis	[29,143]
NDUFAF7	Not known. Catalyze hydroxylation of NDUF57 and dimethylation of NDUF52 of the Q module	-	[144,145]
NDUFAF8/C17ORF89	Not known. Stabilization of NDUFAF5	Leigh syndrome	[146]
NUBPL	Supposed to interact with the developing N module and possibly Q module. Insertion of iron-sulfur clusters in N and Q module subunits	Infantile onset hepatopathy, renal tubular acidosis, developmental delay, short stature, leukoencephalopathy, myopathy, nystagmus, and ataxia	[147-149]
TIMMDC1/C10RF1	ND1/PP-a Insertion of ND1	Infantile onset hypotonia, failure to thrive, delayed or minimal psychomotor development, sensorineural deafness, dysmetria, dyskinetic movements, peripheral neuropathy, nystagmus, and Leigh syndrome	[133,150,151]
TMEM126A	ND4 module Component of MCIA complex, necessary for building the intermediate ND2 module	Autosomal recessive optic atrophy	[152-156]
TMEM126B	ND2/PP-b module Component of MCIA complex, necessary for building the intermediate ND2 module	Exercise intolerance, muscle weakness, myalgia, early-onset renal tubular acidosis, and hypertrophic cardiomyopathy	[157-159]

TMEM186	ND2/PP-b module- Interact strongly with newly synthesized ND3	-	[160]
DMAC1/TMEM261	ND5/PD-b	-	[114]
COA1/MITRAC15	ND2/PP-b module	-	[160]
COA7	-	Autosomal recessive spinocerebellar ataxia with axonal neuropathy type 3 and mitochondrial myo- pathy	[161]
LYRM-2	NADH-Dehydrogenase module Maturation of N-module	-	-

A wide range of pathological phenotypes of the nervous system has been found to affect CI stability/activity both involving mitochondrial- and nuclear-encoded subunits [6]. Many pathological variants in the seven mtDNA encoded subunits, *MT-ND1-6* and *ND4L* have been associated with a wide *spectrum* of syndromes with the age of onset occurring mostly during late childhood or early adulthood [163-166]. Mutations in three *MT-ND* genes are the main cause of Leber’s hereditary optic neuropathy (LHON) [OMIM 535 000], the most common mtDNA inherited disease [167]. LHON is one cause of bilateral acute or subacute, painless loss of central vision in young men (more than 80% of LHON patients are male, because of degeneration of retinal ganglion cell layers [168,169]. Important clues to understanding the pathogenesis of LHON, which is characterized by yet undisclosed genetic and environmental factors affecting the incomplete penetrance, have been obtained by analysis of mtDNA copy number and by proteomics approaches [170-173]. Mitochondrial DNA copy number is a key factor in differentiating LHON affected individuals from the unaffected mutation carriers [170-173]. A mitochondrial proteomic profile of 11778G>A mutant fibroblasts using 2-Dimensional Polyacrylamide Gel Electrophoresis (2-DE) and MS [174] disclosed that most of the mitochondrial proteins - including those involved in intermediary metabolic processes, nucleoid-related proteins, chaperones, *cristae* remodeling ones, and an antioxidant enzyme - were down-regulated, and some OxPhos subunits were altered [174]. The major bioenergetics consequences, particularly of *MT-ND4* and *MT-ND1* mutations, resulted in CI-dependent reduction of ATP synthesis and redox balance leading to increased ROS levels and decreased antioxidant enzyme activities [175-177].

The main pathological mutations found in structural CI subunits are summarized in **Table 2**.

Table 2. Complex II subunits with location, associated clinical phenotypes, and references. Adapted from [70], [47] and [162].

Subunits	Location	Associated clinical phenotypes	References
MTND1	ND1-module	Leber optic atrophy, MELAS syndrome, dystonia, spasticity, and myopathy	[178-180]
MTND2	ND2-module	Leber optic atrophy	[181]
MTND3	ND2-module	Infantile encephalopathy and Leigh syndrome	[182]
MTND4	ND4-module	Leber optic atrophy and MELAS syndrome	[183,184]
MTND4L	ND2-module	Leber optic atrophy	[185]
MTND5	ND5-module	Leber optic atrophy and MELAS syndrome	[186,187]
MTND6	ND2-module	Leber optic atrophy and MELAS syndrome	[186,188]
NDUFV1	N-module	Severe encephalopathy and neurologic abnormalities	[189,190]
NDUFV2	N-module	Hypertrophic cardiomyopathy, truncal hypotonia, and encephalopathy	[191]
NDUFV3	N-module	Complex I deficiency	

NDUFS1	N-module	Growth retardation, axial hypotonia, hepatomegaly, dys-tonia, and persistently hyperlactatemia	[190]
NDUFS2	Q-module	Neonatal lactic acidosis and hypertrophic cardiomyopa- thy	[192]
NDUFS3	Q-module	Leigh syndrome, severe axial dystonia with oral and pharyngeal motor dysfunction, dysphagia and a tetraparetic syndrome	[193]
NDUFS4	Q-module	Muscular hypotonia, absence of visual and auditive at- tention, and cardiac defects	[194]
NDUFS6	Q-module	Fatal infantile lactic acidosis, neonatal myopathy, en- cephalopathy, and lactic acidosis	[195,196]
NDUFS7	Q-module	Leigh syndrome, feeding problems, dysarthria, and ataxia	[197]
NDUFS8	Q-module	Leigh syndrome, poor feeding, and episodes of apnea and cyanosis	[198]
NDUFA11	ND2-module	Fatal infantile metabolic acidosis, brain atrophy, no mo- tor development and hypertrophic cardiomyopathy	[199]
NDUFA1	ND1-module	Leigh syndrome, hypotonia, nystagmus, generalized choreoathetosis, and decreased reflexes	[200]
NDUFA2	N-module	Leigh syndrome, hypertrophic cardiomyopathy, and de- velopmental delay	[201]
NDUFA3	ND1-module	-	
NDUFA5	Q-module	-	
NDUFA6 (LYRM-6)	LYR protein	Auditory and optic neuropathy, mitochondrial-related infantile death, brain disorder, leukoencephalopathy	[202]
NDUFA7	N-module		
NDUFA8	IMS protein (ND1-module)	Intrauterine growth retardation, respiratory insuffi- ciency, lactic acidosis and hypoglycemia	[163]
NDUFA9	Q-module	Severe neonatal hypotonia, dysmorphic features, epi- lepsy, and signs of brainstem involvement	[203]
NDUFA10	ND2-module	Leigh syndrome	
NDUFA11	ND2-module	Encephalocardiomyopathy and fatal infantile lactic acidemia, neuromuscular disorder	
NDUFA12	N-module	Respiratory and metabolic acidosis, hearing loss, apneas, and retinitis pigmentosa	[204]
NDUFA13	ND1-module	Leigh syndrome, progressive loss of motor abilities, sco- liosis, and dystonia	[205]
NDUFB1	ND4-module		
NDUFB2	ND5-module		
NDUFB3	ND5-module	Delayed development, hypotonia, poor eye contact, ab- normal eye movements, poor feeding, encephalopathy, and hearing loss	[206]
NDUFB4	ND4-module		
NDUFB5	ND4-module		
NDUFB6	ND5-module		
NDUFB7	ND5-module		
NDUFB8	ND5-module	Encephalopathy, myopathy, hypotonia, developmental delay, and lactic acidosis, mitochondrial Complex I Defi- ciency in Individuals with Leigh-like Encephalomyopa- thy	[207]
NDUFB9 (LYRM-3)	LYR protein	Leigh syndrome, respiratory failure, seizures, hypotonia, cardiac hypertrophy, failure to thrive and severely delayed psychomotor develop- ment	[206]
NDUFB10	IMS protein (ND4 module)	Progressive hypotonia associated with increased serum lactate	[208]
NDUFB11	ND4-module	Lethal complex I deficiency, X-linked microphthalmia with linear skin defects (MLS) syndrome	[209-211]

NDUFC1	ND2-module	
NDUFC2	ND2-module	X-linked microphthalmia with linear skin defects (MLS) syndrome, cardiomyopathy and other congenital anomalies [212]
NDUFS5	IMS protein (ND2 module)	

Quantitative proteomics has revealed the importance of the 30 non-catalytically active supernumerary subunits of CI. Pathological variants causing CI deficiency have been described in NDUFAF1 [CIA30], ACAD9, and TMEM126B that together with ECSIT, COA1 and TMEM186, form the Mitochondrial Complex I Intermediate Assembly (*MCIA*) [157] important for the biogenesis of the ND2-module. NDUFAF3 (C3ORF60) and NDUFAF4 (C6ORF66) working together in the assembly of the *Q-module*, have been found mutated in different cases of infantile mitochondrial disease [139,140,213- 216].

The gene *NDUFS4* (NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, NM_002495.2), is a hotspot for pathogenic mutations. Inactivation of the *NDUFS4* gene is known to cause mostly, Leigh or Leigh-like syndrome [217-225], a rare disease with a prevalence of roughly 1:40.000 live births [226,227]. Unfortunately, the prognosis of *NDUFS4*-linked LS is poor. Loss of *NDUFS4* affects complex I assembly and causes detrimental structural changes in assembled complex I [217,228]. Several pieces of evidence have suggested that *NDUFS4* plays a role in the late stage of complex I assembly [218,220,229]. *NDUFS4* knock out mouse models [230,231], human and murine cell lines, and more recently induced pluripotent stem cells (iPSCs) from LS patients carrying mtDNA mutations in the *NDUFS4* [68] have been set up to explore strategies to counteract pathophysiological consequences of complex I deficiency. LS patient-derived neural cells have shown defective bioenergetics [232,233], decreased protein synthesis [234], impaired mitochondrial calcium homeostasis [233,235], and abnormal corticogenesis [236]. The presence of defective neurite outgrowth has been confirmed also in neural progenitor cells (NPCs) carrying mutations in the *NDUFS4* as well as in the *SURF1* (Surfeit locus protein 1, NM_003172.2) genes, another well-known cause of LS [237-239].

Structural subunits and specific factors assisting the assembly associated with human diseases are summarized in **Tables 1** and **2**.

4.2 Succinate–Ubiquinone Oxidoreductase - Complex II

Succinate dehydrogenase (SDH, complex II, CII), a ~120 kDa integral membrane complex, participates in both the TCA cycle and the respiratory chain. CII transfers the electrons to CoQ and does not contribute to proton pumping across the mitochondrial membrane. All four subunits are encoded by the nuclear genome. The largest hydrophilic domain is a heterodimer composed of SDHA and SDHB that protrude toward the matrix and contain the redox-active groups' flavin adenine dinucleotide (FAD(H₂)) and three Fe–S clusters, respectively. The smaller hydrophobic domain is composed of SDHC and SDHD and contains two CoQ binding sites [240] providing reduction of ubiquinone to ubiquinol, the mobile electron carrier that links to CIII. Four specific chaperones [SDH assembly factor 1–4 (SDHAF1–4)] participate in the stabilization and incorporation of the prosthetic groups into each of the structural subunits SDHA, SDHB, and SDHC + SDHD [124,241]. In the late stage of assembly of CII, ACN9, similarly to LYRM-8 (also known as SDHAF1), is important for the formation and stabilization of CII throughout the insertion or retention of the Fe-S centers within the protein backbones and FMC1 (Formation of mitochondrial complex V assembly factor 1) [242].

CII defects are quite rare and represent less than 10% of OxPhos deficiency cases [243]. Different forms of encephalopathy and rare neuroendocrine tumors are the two main pathological manifestations that can originate from mutations in CII subunits or assembly factors. Mutations in SDHA, encoding the 70 kDa Flavoprotein subunit, have also been found in rare cases of Leigh syndrome [244-249]. Ultrarare association of bi-genomic

variants in the SDHB and mitochondrial MT-CYB genes has been described in a patient with clinical and metabolic features of a ME-LAS-like syndrome [250].

The main pathological mutations found in CII subunits or assembly factors are summarized in **Tables 3**.

Table 3. Complex II subunits and assembly factors with function, associated clinical phenotypes, and references. Adapted from [264] and [70].

Subunits	Function	Associated clinical phenotypes	References
SDHA	CII subunit	Leigh syndrome, neonatal dilated cardiomyopathy, catecholamine-secreting extra-adrenal paraganglioma	[244-252]
SDHB	CII subunit	Paraganglioma, pheochromocytoma, gastrointestinal stromal tumors	[253,254]
SDHC	CII subunit	Paraganglioma, gastric stromal sarcoma	[255,256]
SDHD	CII subunit	Paraganglioma, pheochromocytoma, gastric stromal sarcoma	[256,257]
Assembly factors			
SDHAF1/LYRM-8	Insert Fe/S clusters into mature SDHB	Leukoencephalopathy, spastic quadriplegia, psychomotor regression	[242]
SDHAF2	Insert FAD cofactor into apo-protein SDHA	Paraganglioma and pheochromocytomas, astrointestina cell sarcomas	[255,257-261]
SDHAF3/NDUFV1/LYRM-10	Maintain SHDB stability	Familial and sporadic pheochromocytomas and paraganglioma	[262]
SDHAF4	Protect the subunit from auto-oxidation and facilitates the assembly with SDHB	Vagal paragangliomas	[263]

4.3 Ubiquinol: cytochrome c oxidoreductase - Complex III

The ubiquinol: cytochrome c oxidoreductase (cytochrome bc1, complex III, CIII) constitutes the central part of the respiratory chain. CIII accepts two electrons from reduced CoQ (CoQH2) and donates them, one by one, to cytochrome c, using catalytic proteins: cytochrome b (MT-CYB - human nomenclature), containing two CoQ binding sites and two heme b groups; UQCRFS1, the Rieske Fe-S protein; and CYC1, containing heme c as the prosthetic group. Each of the two ‘monomers’ is composed of 10 different subunits and associate as a symmetric dimer [265]. The complex assembly starts with the synthesis, membrane insertion, and hemylation of cytochrome b, mediated by UQCC1–3 in humans [266-268], followed by the sequential incorporation of the remaining subunits into a dimeric pre-CIII2 [268]. MZM1L (LYRM7), BCS1L, and tetratricopeptide repeat domain-containing protein 19 (TTC19) are the three assembly factors, known to be involved in the stabilization, incorporation, and metabolism of UQCRFS1 [270-277]. LYRM7 chaperone binds the Rieske protein before its incorporation as the last step of the biogenesis of the nascent CIII dimer (CIII2), acted by BCS1L [271,273,278].

The first mutations found in CIII were identified in MT-CYB, the only subunit encoded by mtDNA [279-282]. Most of these pathological variants were found in heteroplasmy and mainly associated with late-onset sporadic myopathy and exercise intolerance [279-285]. Other MT-CYB mutations were associated with histiocytoid cardiomyopathy [286], parkinsonism and MELAS overlap syndrome [281], or multisystem disorders [287-290].

Among the cases of CIII deficiency of nuclear origin are mutations in assembly factors [291] and the most common are nonsense and missense mutations in TTC19 [292], LYRM7 [293], and BCS1L [294], which cause defective CIII assembly/stability and decreased ubiquinol:cyt *c* oxidoreductase activity. Interestingly, a shuttle of electrons from NADH and/or ubiquinol to CIII, Pyocianine, has been used to efficiently recover mitochondrial function thus ameliorating bioenergetic efficiency in fibroblasts derived from patients' dysfunction due to TTC19, BCS1L, and LYRM7 [278].

The main pathological mutations found in CIII subunits or assembly factors are summarized in **Table 4**.

Table 4. Complex III subunits and assembly factors with function, associated clinical phenotypes, and references. Adapted from [263] and [70].

Subunits	Function	Associated clinical phenotypes	References
UQCRC1	CIII subunit	Parkinsonism with polyneuropathy	[295]
UQCRC2	CIII subunit	Hypoglycemia, lactic acidosis, ketosis, and hyperammonemia	[296]
MT-CYB	CIII subunit	Leber optic atrophy, exercise intolerance, encephalomyopathy, cardiomyopathy, and multisystemic disorder, histiocytosis cardiomyopathy, parkinsonism, and MELAS overlap syndrome	[280,281,286,287,297,298]
CYC1	CIII subunit	Neurologic deterioration, insulin-responsive hyperglycemia, ketoacidosis with increased serum lactate, liver failure, and hyperammonemia	[299]
UQCRCF1	CIII subunit	Cardiomyopathy and alopecia totalis	[300]
UQCRH	CIII subunit	-	-
UQCRB	CIII subunit	Gastroenteritis, liver enlargement, hypoglycemia, and metabolic acidosis but normal psychomotor development at age 4, hepatopathy	[301]
UQCRQ	CIII subunit	Severe neurologic phenotype, early-onset severe encephalopathy	[302]
UQCR10	CIII subunit	-	-
UQCR11	CIII subunit	-	-
UQCRCF1	CIII subunit	Decreased CIII activity, lactic acidosis, cardiomyopathy, and alopecia totalis	[300]
Assembly factors			
UQCC1	Cytochrome <i>b</i> assembly factor		
UQCC2	Cytochrome <i>b</i> assembly factor	614461 Intrauterine growth retardation, neonatal lactic acidosis and renal tubular dysfunction	[267,303]
UQCC3	Cytochrome <i>b</i> assembly factor	Lactic acidosis, hypoglycemia, hypotonia, and delayed development	[268]
VPS53	Heme lyase (Cytochrome c1)	Complicated hereditary spastic paraparesis	[304]
BCS1L	AAA-ATPase involved in Rieske protein incorporation. Stabilization, incorporation, and metabolism of UQCRCF1	GRACILE Syndrome, Bjornstad Syndrome, myopathy, encephalopathy, proximal tubulopathy, and liver failure	[26,275,294,306-308,310]

MZM1L/LYRM-7	Matrix protein involved in Rieske protein incorporation. Stabilization, incorporation, and metabolism of UQCRFS1	Neurological decompensation and regression, leukoencephalopathy and liver failure, infantile CIII deficiency associated with cavitating leukoencephalopathy metabolic decompensation	[293,311-313]
TTC19	Rieske protein metabolism. Stabilization, incorporation, and metabolism of UQCRFS1	Progressive encephalopathy, ataxia, spastic paraparesis, and psychiatric phenotype	[292,314- 317]

4.4 Cytochrome *c* oxidase - Complex IV

Cytochrome *c* oxidase (COX, complex IV, CIV) is the terminal complex of the ETC. The enzyme transfers electrons from cytochrome *c* to molecular oxygen. In humans, it is composed of 14 subunits, with the NDUFA4, the most recently discovered subunit initially attributed to CI [318,319], found to be incorporated in the structure of monomeric human CIV [320]. Only two, MT-CO1 and MT-CO2, are catalytical subunits. MTCO1 contains three prosthetic groups: cytochrome *a*3 and CuB, which form the bi-nuclear center that binds oxygen, and cytochrome *a*. MT-CO2 incorporates the CuA center [321]. MT-CO3 is necessary to provide additional stability to the enzyme while it undergoes turnover [322]. The remaining subunits (COX4, 5A, 5B, 6A, 6B, 6C, 7A, 7B, 7C, 8A) are supposed to have a structural role in the stabilization of the complex. CIV is the only ETC complex that evolved tissue, developmental and species-specific isoforms for COX subunits 4, 6A, 6B, 7A, 7B, and 8A [323,324].

CIV assembly grows with a modular process through the incorporation of modules formed by different subunits and defined by each of the mtDNA-encoded core subunits [124,325-326]. Any subunit of complex IV could carry mutations and rise a mitochondriopathy [324,327-329]. Mutations in the MT-CO1, MT-CO2, and MT-CO3 are causative of COX deficiency and mitochondrial disease with an extreme clinical heterogeneity (Table 5).

Table 5. Complex IV subunits with associated clinical phenotypes and references. Adapted from [264] and [70].

Subunits	Associated clinical phenotypes	References
MTCO1	MELAS syndrome, myopathy, myoglobinuria, motor neuron disease, exercise intolerance, epilepsy, multisystem disorders, deafness, LHON, or mitochondrial sensorineural hearing loss	[330-334]
MTCO2	Encephalomyopathy, LHON, myopathy, hypertrophic cardiomyopathy	[335-338]
MTCO3	MIDD, LHON, myopathy, Leigh disease, myoglobinuria, sporadic bilateral optic neuropathy, rhabdomyolysis, encephalopathy	[339-344]
COX4I1	Short stature, poor weight gain, mild dysmorphic features, Fanconi anemia, Leigh-like syndrome	[345,346]
COX4I2	Exocrine pancreatic insufficiency, dyserythropoietic anemia, calvarial hyperostosis	[347]
COX5A	Early-onset pulmonary arterial hypertension, lactic acidemia, failure to thrive	[348]
COX6A1	Charcot-Marie-Tooth disease	[349]
COX6A2	Muscle weakness and hypotonia, cardiomyopathy	[350]
COX6B1	Severe infantile encephalomyopathy	[328,329]
COX7A1	Failure to thrive, encephalopathy, hypotonia	[351]
COX7B	Microphthalmia with linear skin lesions	[352]
COX8A	Leigh-like syndrome presenting with leukodystrophy and severe epilepsy	[353]

NDUFA4	Leigh syndrome	[318]
--------	----------------	-------

Pathological variants in ‘supernumerary’ COX subunits have been reported in tissue and development-specific isoforms [323]. Among the assembly factors, the most representative is *SURF1*, the functional absence of which causes LS [237,238,261] or even Charcot–Marie–Tooth disease [354]. The elucidation of the pathogenetic mechanism has received an impulse recently [68]. Mutations in *COX10*, which catalyzes the farnesylation of a vinyl group of heme *b*, cause LS and other forms of the fatal early-onset neurological syndrome [355-357]. Mutations in *COX15*, which catalyzes the subsequent step of heme synthesis, cause variable clinical presentations [358-360]. Copper delivery to the active sites of *MT-CO1* and *MT-CO2* involves factors essential for COX activity [234,361]. *SCO1*, *SCO2*, and *COA6* have been found mutated in patients showing CIV deficiency and fatal outcomes [325,355,362-373]. Among complex IV proteins, *COX6B1* assists CIV assembly, working as a linking subunit at the dimeric interface of CIV [374].

The specific functions of the remaining proteins (all associated with human diseases, see **Table 6**) are known only in part and require additional studies.

Table 6. Complex IV assembly factors with function, associated clinical phenotypes, and references. Adapted from [263] and [70].

Assembly Factors	Function	Associated clinical phenotypes	References
RNA stability and translation			
TACO1	Translational activator of mitochondria encoded MTCO1.	Leigh syndrome	[375,376]
LRPPRC	Mitochondrial mRNA stability.	French Canadian type of Leigh syndrome	[377]
FASTKD2	Involved in post-transcriptional RNA maturation, ribosome biogenesis and translation.	Brain atrophy, epilepsy, delayed psychomotor development, bilateral optic atrophy, spastic hemiparesis, cardiomyopathy	[378-380]
Heme a biosynthesis and insertion			
COX10	Heme a synthesis (conversion of heme b into heme o).	Leigh syndrome, encephalopathy, cardiomyopathy, sensorineural deafness, and metabolic acidosis	[356,357,381,382]
COX15	Heme a synthesis (conversion of heme o into heme a).	Leigh syndrome, encephalopathy, cardiomyopathy, sensorineural deafness, and metabolic acidosis	[356,360,383,384]
SURF1	Involved in the insertion or stabilization of heme a3.	Leigh syndrome, Charcot–Marie–Tooth disease	[237,238,261,354,385]
Copper metabolism and insertion			
COA5/C2ORF64	Involved in the unknown step of CIV biogenesis	Fatal infantile cardioencephalomyopathy	[386]
COA6/C1ORF31??	Copper homeostasis and transport to CIV.	Fatal infantile cardioencephalopathy	[372,373,387]
SCO1	Incorporation of copper atoms. (biogenesis of CuA center	Cardioencephalomyopathy, Leigh syndrome-like symptoms, spinal muscular atrophy-like presentations, Charcot–Marie–Tooth disease type 4. CIV deficiency, neonatal hepatopathy, encephalopathy with hepatopathy and cardiomyopathy, pure encephalopathy, metabolic syndrome with exclusively fatal lactic acidosis	[362,368,370,382,388,389]
SCO2	Incorporation of copper atoms. (biogenesis of CuA center	Cardioencephalomyopathy, Leigh syndrome-like symptoms, spinal muscular	[237,238,354,364- 368]

		atrophy-like presentations, Charcot–Marie–Tooth disease type 4. CIV deficiency, cardiac hypertrophy	
COX11	Copper chaperone.	Coloboma, Ocular, With Or Without Hearing Impairment, Cleft Lip/Palate, And/Or Mental Retardation and Spinal Muscular Atrophy, Distal, X-Linked 3	[390]
COX16	MTCO2 maturation.		[391,392]
COX17	Copper transfer.	-	[393]
COX19	Stabilization of COX11.		[394,395]
COX20	Stabilization of MT-CO2.	Cerebellar ataxia	[396-398]
Assembly			
COA3/MITRAC12	Required for MTCO1 stability and assembly. Involved in translational regulation of MTCO1 and prevention of MTCO1 aggregation before assembly.	Mild phenotype, exercise intolerance, peripheral neuropathy, obesity, and short stature	[399-402]
COA7	Unknown.	Ataxia and peripheral neuropathy, cognitive impairments, leukodystrophy	[161,403]
COX14/c12orf62	MTCO1 stability and assembly; avoid MTCO1 aggregation.	Severe lactic acidosis and dysmorphic features	[397,404]
CMC1	Stabilizes the interaction between MTCO1, COX14, and COA3.		[405]
COX20/FAM36A	MTCO2 chaperone for copper metalation.	Growth delay, hypotonia, cerebellar ataxia	[397,398,406]
PET100	Stabilizes MT-CO2 module	Early-onset psychomotor delay, seizures, hypotonia, Leigh syndrome, CIV deficiency, and fatal infantile lactic acidosis	[309,407-409]
PET117	Assembly factor: possible role in Cox15 oligomerization and function, stabilizes MT-CO2 module	Neurodevelopmental regression. And bulbar lesions	[410-412]
MR-1S	Interacts with PET117 and PET100.		[326]
APOPT1/COA8	Intermediate assembly steps. Putative role in CIV protection from ROS damage, enhance CIV biogenesis	Leukodystrophy, neurological signs	[413-415]
COX18	Promotes the translocation of MTCO2 globular domain through the IMM.	Isolated COX deficiency in infancy	[416-418]
COX19	Stabilization of COX11.	Isolated COX deficiency in infancy	[394,395,418]
COA-X	Putative assembly factor		[419]
HIGD2A	Promotes incorporation of MT-CO3 module		

4.5 ATP synthase - Complex V

ATP synthase (Complex V, CV) is the enzyme that catalyzes the synthesis of ATP required as an energy source for various cellular processes from ADP and phosphate utilizing the proton-motive force generated through electron transfer. The F₁F₀ ATP synthase has two functional domains: the hydrophilic matrix-facing F₁ domain for ATP generation and the membrane-facing F₀ domain for proton translocation [420,421]. The proton translocation leads to rotary motion of the c-ring in the F₀ domain that is connected to the F₁ catalytic subunit by the peripheral stalk (PS). The human CV is composed of 29 proteins of 18 kinds, including the Inhibitory factor 1, IF₁, in which only F₀-ATP₆ and ATP₈ are mtDNA encoded [422]. The complete structure of the dimeric and monomeric mammalian mitochondrial F₁F₀-ATP synthase has been just recently resolved by Cryo-EM [423,424].

The assembly pathway of human CV is also modular [420,422,425,426] since three subcomplexes, F₁ module, c-ring, and PS are formed individually and then associate together. The assembly starts with the three alpha and three beta subunits of the F₁ domain to which the rest of the subunits bind subsequently. The eight units composing the c-ring are assembled within the IMM. When the two subcomplexes get together, the subunits of the PS join followed by the remaining subunits of the membrane domain, including MT-ATP₆ and MT-ATP₈ [124]. Only three assembly factors have been described among which ATPAF1 and ATPAF2, that binds and stabilizes subunit beta [427] and subunit alpha [428], respectively.

Pathogenic mutations have been reported both in mtDNA and nDNA encoded ATP synthase subunits. The coding sequences of two F₀ subunits are overlapping in the human mtDNA and pathological variants in both are the cause of sporadic and maternally inherited mitochondrial disease (Table 7).

Table 7. Complex V subunits and assembly factors with function, associated clinical phenotypes, and references. Adapted from [263], integrated with [70].

Subunits	Location	Associated clinical phenotypes	References
MT-ATP6	Fo domain	Mitochondrial CV deficiency Neuropathy, Ataxia and Retinitis Pigmentosa (NARP) syndrome Leigh syndrome Adult-onset ataxia and polyneuropathy Bilateral striatal necrosis Motor neuron syndrome Mitochondrial myopathy, lactic acidosis, and sideroblastic anemia	[90,91,429-444]
MT-ATP8	Fo domain	Mitochondrial CV deficiency Valproate-induced reversible brain atrophy Hypertrophic cardiomyopathy	[445,446]
MT-ATP6/8 overlap region	Fo domain	Mitochondrial CV deficiency Infantile hypertrophic cardiomyopathy	[444]
ATP5F1A (GENE ATP5A1)	F1 domain	Mitochondrial CV deficiency Combined OXPHOS deficiency Fatal infantile encephalopathy	[447,448]
ATP5F1D	F1 domain	Mitochondrial CV deficiency Metabolic decompensation with lactic acidosis, hypoglycemia, hyperammonemia, and 3-methylglucaconic aciduria, encephalopathy	[449]
ATP5F1E	F1 domain	Mitochondrial Cdeficiency	[134,450]

		Neonatal-onset lactic acidosis, 3-methylglutaconic aciduria, mild mental retardation, hypertrophic cardiomyopathy, and peripheral neuropathy	
Assembly factors			
ATPAF1	Binds and stabilizes subunit beta of F1 Domain	Asthma in children	[451]
ATPAF2	Binds and stabilizes subunit alpha of F1 domain	Degenerative encephalopathy, elevated lactate levels, developmental delay	[452]
TMEM70	Unknown	Neonatal mitochondrial encephalocardiomyopathy. ATP synthase deficiency	[134]

Mutations in *MT-ATP6* have been found in Neuropathy, Ataxia, and Retinitis Pigmentosa (NARP) syndrome [90] and Maternally Inherited Leigh Syndrome (MILS) [429,453]. NARP is an adult-onset slowly progressive form, whereas MILS is an early onset, highly disabling, often fatal condition. In most cases, NARP and MILS are associated with mutations in position m.8993 changing from T to C or G [430,454,455]. The T>G transition is usually more severe, and the severity of the disease depends rather tightly on the degree of heteroplasmy found in post-mitotic tissues [91,455]. Until now, only three out of the sixteen nuclear-encoded CV subunits and three assembly factors (e.g., ATPAF2; ATP12 and TMTM70) have been associated with mitochondrial disease (see **Table 7**).

4.6 Respiratory supercomplexes

The OxPhos complexes interact with each other forming higher-order structures of the respiratory chain enzymes which have been called supercomplexes (SCs). Complexes IV and V can form dimers and oligomers [456-458]. According to their molecular size and subunit composition, the main SCs have been assigned the following stoichiometries: III₂IV₁, I₁III₂, I₁III₂IV₁, and I₂III₂IV₁₋₂. In particular, the association of complexes I, III₂ and IV, SC I₁III₂IV₁, considered as a functional unit capable of transferring electrons from NADH to O₂, is defined as the ‘respirasome’ [459] whereas the supercomplex I₂III₂IV₂ has been named as ‘respiratory megacomplex’ [123]. High-resolution Cryo-EM structures of the respirasome of several mammalian species, including humans, have been recently resolved [123,460-463]. The respirasome organization was supposed to be functionally advantageous making electron transfer from CI to CIV through CIII₂ more efficient and decreasing the formation of deleterious ROS [464-467]. The fact that the biogenesis of CIII₂ and CIV occurs independently but the CI assembly is not, can be an explanation of the reason why mutations in CIII₂ and CIV may result in secondary effects also on CI assembly but only rarely the *vice versa* [194,468]. High-throughput proteomics techniques have recently been applied in human cybrids holo-CIII₂-deficient, demonstrating the loss of SCs containing CIII₂ and CI when the CIII₂ is not fully assembled. In this model, the combination of null CIII and markedly reduced CI enzymatic activity, confirmed the well-established connection between CIII₂ deficiency and hampered assembly process in CI [469].

In astrocytes, most of CI is free, resulting in poor mitochondrial respiration but high ROS production; while, in neurons, CI is mostly embedded into supercomplexes, thus resulting in high mitochondrial respiration and low ROS production. Interestingly, the proteins that shape the *cristae*, the mitochondrial contact sites, and the *cristae*-organizing system (*MICOS*) complex were found to be essential for the assembly and functionality of the OxPhos system [470]. Understanding the structure and assembly of the SCs is very relevant to explain the cases of combined respiratory chain deficiency due to complexes’ functional and structural interdependency.

For expert reviews on the relationship between *cristae* dynamics and mitochondrial OxPhos bioenergetics, please refer to references [471,472]. For a detailed review on the formation and function of supercomplexes, see Reference [473].

5. Conclusions

An integrative approach that combines multi-omics data could represent a strategic way to solve, at least in part, the complexity of mitochondrial diseases and mitochondrial medicine highlighting the interrelationships of the involved OxPhos complexes and their functions, and the knowledge about genotype-phenotype correlation. However, the science behind combined omic approaches, will need the integration of data from genomics, transcriptomics, proteomics, and metabolomics, to include also the novel approaches looking to the post-translational modifications that provide the link between metabolism, mitochondrial proteome, and the two cellular genomes. The recent application of CRISPR/Cas9 technology to patient-specific iPSCs, to generate neural cultures and cerebral organoids is providing patient-specific cellular and tissue models that allow the investigation of the defects of neuronal morphogenesis caused by specific mutations (**Figure 1**). Thus, in the multi-omic era, the opportunity to understand the cause of each mitochondrial disease becomes ever more tangible.

Author Contributions: Conceptualization, P.Z., and V.P.; writing—original draft preparation, P.Z.; S.D. and V.P.; writing, review, supervision and editing, P.Z. and V.P.; supervision, V.P. and F.M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by grants from REGIONE PUGLIA-MALATTIE RARE-Petruzzella Del. N.246 10.09.2019_UPB-SMBNOS and by donations of Parents' Associations (to VP). This study received partial financial support from the Italian Ministry of Health-Ricerca Corrente, MITO-NEXT, Mit-OMICS (to FMS), Fondazione Telethon (Grant numbers GUP09004). The authors alone are responsible for the content and writing.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We thank the patients' association MITOCON and UILDM (Unione Italiana Lotta alla Distrofia Muscolare) and CollaGe-associazione-Genitori-Manzoni-Poli-Molfetta.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Chandel, N.S. Mitochondria as Signaling Organelles. *BMC Biol* **2014**, *12*, 34, doi:10.1186/1741-7007-12-34.
2. Mitchell, P. Coupling of Phosphorylation to Electron and Hydrogen Transfer by a Chemi-Osmotic Type of Mechanism. *Nature* **1961**, *191*, 144–148, doi:10.1038/191144a0.
3. Spinelli, J.B.; Haigis, M.C. The Multifaceted Contributions of Mitochondria to Cellular Metabolism. *Nat Cell Biol* **2018**, *20*, 745–754, doi:10.1038/s41556-018-0124-1.
4. De Stefani, D.; Raffaello, A.; Teardo, E.; Szabò, I.; Rizzuto, R. A Forty-Kilodalton Protein of the Inner Membrane Is the Mitochondrial Calcium Uniporter. *Nature* **2011**, *476*, 336–340, doi:10.1038/nature10230.
5. Green, D.R.; Galluzzi, L.; Kroemer, G. Metabolic Control of Cell Death. *Science* **2014**, *345*, 1250256–1250256, doi:10.1126/science.1250256.
6. Gorman, G.S.; Chinnery, P.F.; DiMauro, S.; Hirano, M.; Koga, Y.; McFarland, R.; Suomalainen, A.; Thorburn, D.R.; Zeviani, M.; Turnbull, D.M. Mitochondrial Diseases. *Nat Rev Dis Primers* **2016**, *2*, 16080, doi:10.1038/nrdp.2016.80.
7. Muraresku, C.C.; McCormick, E.M.; Falk, M.J. Mitochondrial Disease: Advances in Clinical Diagnosis, Management, Therapeutic Development, and Preventative Strategies. *Curr Genet Med Rep* **2018**, *6*, 62–72, doi:10.1007/s40142-018-0138-9.
8. Piel, R.B.; Dailey, H.A.; Medlock, A.E. The Mitochondrial Heme Metabolon: Insights into the Complex(Ity) of Heme Synthesis and Distribution. *Molecular Genetics and Metabolism* **2019**, *128*, 198–203, doi:10.1016/j.ymgme.2019.01.006.
9. Miller, W.L. Steroid Hormone Synthesis in Mitochondria. *Molecular and Cellular Endocrinology* **2013**, *379*, 62–73, doi:10.1016/j.mce.2013.04.014.
10. Anderson, A.J.; Jackson, T.D.; Stroud, D.A.; Stojanovski, D. Mitochondria—Hubs for Regulating Cellular Biochemistry: Emerging Concepts and Networks. *Open Biol.* **2019**, *9*, 190126, doi:10.1098/rsob.190126.

11. Chan, D.C. Fusion and Fission: Interlinked Processes Critical for Mitochondrial Health. *Annu. Rev. Genet.* **2012**, *46*, 265–287, doi:10.1146/annurev-genet-110410-132529.
12. Pickles, S.; Vigie, P.; Youle, R.J. Mitophagy and Quality Control Mechanisms in Mitochondrial Maintenance. *Current Biology* **2018**, *28*, R170–R185, doi:10.1016/j.cub.2018.01.004.
13. Mills, E.L.; Kelly, B.; O'Neill, L.A.J. Mitochondria Are the Powerhouses of Immunity. *Nat Immunol* **2017**, *18*, 488–498, doi:10.1038/ni.3704.
14. Tikou, V.; Tan, M.-W.; Dikic, I. Mitochondrial Functions in Infection and Immunity. *Trends in Cell Biology* **2020**, *30*, 263–275, doi:10.1016/j.tcb.2020.01.006.
15. Schapira, A.H. Mitochondrial Disease. *The Lancet* **2006**, *368*, 70–82, doi:10.1016/S0140-6736(06)68970-8.
16. DiMauro, S.; Schon, E.A. Mitochondrial Respiratory-Chain Diseases. *N Engl J Med* **2003**, *348*, 2656–2668, doi:10.1056/NEJMra022567.
17. Luft, R. The Development of Mitochondrial Medicine. *Proceedings of the National Academy of Sciences* **1994**, *91*, 8731–8738, doi:10.1073/pnas.91.19.8731.
18. La Morgia, C.; Maresca, A.; Caporali, L.; Valentino, M.L.; Carelli, V. Mitochondrial Diseases in Adults. *J Intern Med* **2020**, *287*, 592–608, doi:10.1111/joim.13064.
19. Petruzzella, V.; Tiranti, V.; Fernandez, P.; Ianna, P.; Carrozzo, R.; Zeviani, M. Identification and Characterization of Human CDNAs Specific to BCS1, PET112, SCO1, COX15, and COX11, Five Genes Involved in the Formation and Function of the Mitochondrial Respiratory Chain. *Genomics* **1998**, *54*, 494–504, doi:10.1006/geno.1998.5580.
20. Stenton, S.L.; Prokisch, H. Advancing Genomic Approaches to the Molecular Diagnosis of Mitochondrial Disease. *Essays in Biochemistry* **2018**, *62*, 399–408, doi:10.1042/EBC20170110.
21. Stenton, S.L.; Prokisch, H. Genetics of Mitochondrial Diseases: Identifying Mutations to Help Diagnosis. *EBioMedicine* **2020**, *56*, 102784, doi:10.1016/j.ebiom.2020.102784.
22. Calvo, S.E.; Compton, A.G.; Hershman, S.G.; Lim, S.C.; Lieber, D.S.; Tucker, E.J.; Laskowski, A.; Garone, C.; Liu, S.; Jaffe, D.B.; et al. Molecular Diagnosis of Infantile Mitochondrial Disease with Targeted Next-Generation Sequencing. *Science Translational Medicine* **2012**, *4*, 118ra10–118ra10, doi:10.1126/scitranslmed.3003310.
23. Plutino, M.; Chaussonot, A.; Rouzier, C.; Ait-El-Mkadem, S.; Fragaki, K.; Paquis-Flucklinger, V.; Bannwarth, S. Targeted next Generation Sequencing with an Extended Gene Panel Does Not Impact Variant Detection in Mitochondrial Diseases. *BMC Med Genet* **2018**, *19*, 57, doi:10.1186/s12881-018-0568-y.
24. Anderson, S.; Bankier, A.T.; Barrell, B.G.; de Bruijn, M.H.L.; Coulson, A.R.; Drouin, J.; Eperon, I.C.; Nierlich, D.P.; Roe, B.A.; Sanger, F.; et al. Sequence and Organization of the Human Mitochondrial Genome. *Nature* **1981**, *290*, 457–465, doi:10.1038/290457a0.
25. Garone, C.; Donati, M.A.; Sacchini, M.; Garcia-Diaz, B.; Bruno, C.; Calvo, S.; Mootha, V.K.; DiMauro, S. Mitochondrial Encephalomyopathy Due to a Novel Mutation in *ACAD9*. *JAMA Neurol* **2013**, *70*, 1177, doi:10.1001/jamaneurol.2013.3197.
26. Oláhová, M.; Berti, C.C.; Collier, J.J.; Alston, C.L.; Jameson, E.; Jones, S.A.; Edwards, N.; He, L.; Chinnery, P.F.; Horvath, R.; et al. Molecular Genetic Investigations Identify New Clinical Phenotypes Associated with BCS1L-Related Mitochondrial Disease. *Human Molecular Genetics* **2019**, *28*, 3766–3776, doi:10.1093/hmg/ddz202.
27. Stenton, S.L.; Kremer, L.S.; Kopajtich, R.; Ludwig, C.; Prokisch, H. The Diagnosis of Inborn Errors of Metabolism by an Integrative “Multi-omics” Approach: A Perspective Encompassing Genomics, Transcriptomics, and Proteomics. *Jrnl of Inher Metab Disea* **2020**, *43*, 25–35, doi:10.1002/jimd.12130.
28. Mootha, V.K.; Bunkenborg, J.; Olsen, J.V.; Hjerrild, M.; Wisniewski, J.R.; Stahl, E.; Bolouri, M.S.; Ray, H.N.; Sihag, S.; Kamal, M.; et al. Integrated Analysis of Protein Composition, Tissue Diversity, and Gene Regulation in Mouse Mitochondria. *Cell* **2003**, *115*, 629–640, doi:10.1016/S0092-8674(03)00926-7.
29. Pagliarini, D.J.; Calvo, S.E.; Chang, B.; Sheth, S.A.; Vafai, S.B.; Ong, S.-E.; Walford, G.A.; Sugiana, C.; Boneh, A.; Chen, W.K.; et al. A Mitochondrial Protein Compendium Elucidates Complex I Disease Biology. *Cell* **2008**, *134*, 112–123, doi:10.1016/j.cell.2008.06.016.
30. Gonczarowska-Jorge, H.; Zahedi, R.P.; Sickmann, A. The Proteome of Baker's Yeast Mitochondria. *Mitochondrion* **2017**, *33*, 15–21, doi:10.1016/j.mito.2016.08.007.
31. Clamp, M.; Fry, B.; Kamal, M.; Xie, X.; Cuff, J.; Lin, M.F.; Kellis, M.; Lindblad-Toh, K.; Lander, E.S. Distinguishing Protein-Coding and Noncoding Genes in the Human Genome. *Proceedings of the National Academy of Sciences* **2007**, *104*, 19428–19433, doi:10.1073/pnas.0709013104.
32. Ponomarenko, E.A.; Poverennaya, E.V.; Ilgisonis, E.V.; Pyatnitskiy, M.A.; Kopylov, A.T.; Zgoda, V.G.; Lisitsa, A.V.; Archakov, A.I. The Size of the Human Proteome: The Width and Depth. *International Journal of Analytical Chemistry* **2016**, *2016*, 1–6, doi:10.1155/2016/7436849.
33. Lopez, M.F.; Kristal, B.S.; Chernokalskaya, E.; Lazarev, A.; Shestopalov, A.I.; Bogdanova, A.; Robinson, M. High-Throughput Profiling of the Mitochondrial Proteome Using Affinity Fractionation and Automation. *Electrophoresis* **2000**, *21*, 3427–3440, doi:10.1002/1522-2683(20001001)21:16<3427::AID-ELPS3427>3.0.CO;2-L.
34. Karlberg, O.; Canbäck, B.; Kurland, C.G.; Andersson, S.G. The Dual Origin of the Yeast Mitochondrial Proteome. *Yeast* **2000**, *17*, 170–187, doi:10.1002/1097-0061(20000930)17:3<170::AID-YEA25>3.0.CO;2-V.
35. Cotter, D. MitoProteome: Mitochondrial Protein Sequence Database and Annotation System. *Nucleic Acids Research* **2004**, *32*, 463D – 467, doi:10.1093/nar/gkh048.

36. Guda, P.; Subramaniam, S.; Guda, C. Mitoproteome: Human Heart Mitochondrial Protein Sequence Database. In *Cardiovascular Proteomics*; Humana Press: New Jersey, 2006; Vol. 357, pp. 375–384 ISBN 978-1-59745-214-4.
37. Calvo, S.E.; Clauser, K.R.; Mootha, V.K. MitoCarta2.0: An Updated Inventory of Mammalian Mitochondrial Proteins. *Nucleic Acids Res* **2016**, *44*, D1251–D1257, doi:10.1093/nar/gkv1003.
38. Smith, A.C.; Robinson, A.J. MitoMiner v3.1, an Update on the Mitochondrial Proteomics Database. *Nucleic Acids Res* **2016**, *44*, D1258–D1261, doi:10.1093/nar/gkv1001.
39. Doccini, S.; Morani, F.; Nesti, C.; Pezzini, F.; Calza, G.; Soliymani, R.; Signore, G.; Rocchiccioli, S.; Kanninen, K.M.; Huuskonen, M.T.; et al. Proteomic and Functional Analyses in Disease Models Reveal CLN5 Protein Involvement in Mitochondrial Dysfunction. *Cell Death Discov* **2020**, *6*, 18, doi:10.1038/s41420-020-0250-y.
40. Hung, V.; Lam, S.S.; Udeshi, N.D.; Svinkina, T.; Guzman, G.; Mootha, V.K.; Carr, S.A.; Ting, A.Y. Correction: Proteomic Mapping of Cytosol-Facing Outer Mitochondrial and ER Membranes in Living Human Cells by Proximity Biotinylation. *eLife* **2019**, *8*, e50707, doi:10.7554/eLife.50707.
41. Geladaki, A.; Kočevár Britovšek, N.; Breckels, L.M.; Smith, T.S.; Vennard, O.L.; Mulvey, C.M.; Crook, O.M.; Gatto, L.; Lilley, K.S. Combining LOPIT with Differential Ultracentrifugation for High-Resolution Spatial Proteomics. *Nat Commun* **2019**, *10*, 331, doi:10.1038/s41467-018-08191-w.
42. Sung, A.Y.; Floyd, B.J.; Pagliarini, D.J. Systems Biochemistry Approaches to Defining Mitochondrial Protein Function. *Cell Metabolism* **2020**, *31*, 669–678, doi:10.1016/j.cmet.2020.03.011.
43. Ohue, M.; Matsuzaki, Y.; Uchikoga, N.; Ishida, T.; Akiyama, Y. MEGADOCK: An All-to-All Protein-Protein Interaction Prediction System Using Tertiary Structure Data. *PPL* **2013**, *21*, 766–778, doi:10.2174/09298665113209990050.
44. Ohue, M.; Shimoda, T.; Suzuki, S.; Matsuzaki, Y.; Ishida, T.; Akiyama, Y. MEGADOCK 4.0: An Ultra-High-Performance Protein-Protein Docking Software for Heterogeneous Supercomputers. *Bioinformatics* **2014**, *30*, 3281–3283, doi:10.1093/bioinformatics/btu532.
45. Hayashi, T.; Matsuzaki, Y.; Yanagisawa, K.; Ohue, M.; Akiyama, Y. MEGADOCK-Web: An Integrated Database of High-Throughput Structure-Based Protein-Protein Interaction Predictions. *BMC Bioinformatics* **2018**, *19*, 62, doi:10.1186/s12859-018-2073-x.
46. Floyd, B.J.; Wilkerson, E.M.; Veling, M.T.; Minogue, C.E.; Xia, C.; Beebe, E.T.; Wrobel, R.L.; Cho, H.; Kremer, L.S.; Alston, C.L.; et al. Mitochondrial Protein Interaction Mapping Identifies Regulators of Respiratory Chain Function. *Molecular Cell* **2016**, *63*, 621–632, doi:10.1016/j.molcel.2016.06.033.
47. Formosa, L.E.; Dibley, M.G.; Stroud, D.A.; Ryan, M.T. Building a Complex Complex: Assembly of Mitochondrial Respiratory Chain Complex I. *Seminars in Cell & Developmental Biology* **2018**, *76*, 154–162, doi:10.1016/j.semcdb.2017.08.011.
48. Yu, Y.; Herman, P.; Rothman, D.L.; Agarwal, D.; Hyder, F. Evaluating the Gray and White Matter Energy Budgets of Human Brain Function. *J Cereb Blood Flow Metab* **2018**, *38*, 1339–1353, doi:10.1177/0271678X17708691.
49. Sokoloff, L.; Reivich, M.; Kennedy, C.; Rosiers, M.H.D.; Patlak, C.S.; Pettigrew, K.D.; Sakurada, O.; Shinohara, M. THE [14 C] DEOXYGLUCOSE METHOD FOR THE MEASUREMENT OF LOCAL CEREBRAL GLUCOSE UTILIZATION: THEORY, PROCEDURE, AND NORMAL VALUES IN THE CONSCIOUS AND ANESTHETIZED ALBINO RAT. *J Neurochem* **1977**, *28*, 897–916, doi:10.1111/j.1471-4159.1977.tb10649.x.
50. Bélanger, M.; Allaman, I.; Magistretti, P.J. Brain Energy Metabolism: Focus on Astrocyte-Neuron Metabolic Cooperation. *Cell Metabolism* **2011**, *14*, 724–738, doi:10.1016/j.cmet.2011.08.016.
51. Barros, L.F.; Courjaret, R.; Jakoby, P.; Loaiza, A.; Lohr, C.; Deitmer, J.W. Preferential Transport and Metabolism of Glucose in Bergmann Glia over Purkinje Cells: A Multiphoton Study of Cerebellar Slices. *Glia* **2009**, *57*, 962–970, doi:10.1002/glia.20820.
52. Jha, M.K.; Morrison, B.M. Glia-Neuron Energy Metabolism in Health and Diseases: New Insights into the Role of Nervous System Metabolic Transporters. *Experimental Neurology* **2018**, *309*, 23–31, doi:10.1016/j.expneurol.2018.07.009.
53. Parpura, V.; Basarsky, T.A.; Liu, F.; Jeftinija, K.; Jeftinija, S.; Haydon, P.G. Glutamate-Mediated Astrocyte-Neuron Signalling. *Nature* **1994**, *369*, 744–747, doi:10.1038/369744a0.
54. Xin, W.; Bonci, A. Functional Astrocyte Heterogeneity and Implications for Their Role in Shaping Neurotransmission. *Front. Cell. Neurosci.* **2018**, *12*, 141, doi:10.3389/fncel.2018.00141.
55. Oliveira, J.F.; Sardinha, V.M.; Guerra-Gomes, S.; Araque, A.; Sousa, N. Do Stars Govern Our Actions? Astrocyte Involvement in Rodent Behavior. *Trends in Neurosciences* **2015**, *38*, 535–549, doi:10.1016/j.tins.2015.07.006.
56. Allen, N.J. Star Power: Astrocytes Regulate Behavior. *Cell* **2019**, *177*, 1091–1093, doi:10.1016/j.cell.2019.04.042.
57. Herrero-Mendez, A.; Almeida, A.; Fernández, E.; Maestre, C.; Moncada, S.; Bolaños, J.P. The Bioenergetic and Antioxidant Status of Neurons Is Controlled by Continuous Degradation of a Key Glycolytic Enzyme by APC/C-Cdh1. *Nat Cell Biol* **2009**, *11*, 747–752, doi:10.1038/ncb1881.
58. Bittner, C.X.; Valdebenito, R.; Ruminot, I.; Loaiza, A.; Larenas, V.; Sotelo-Hitschfeld, T.; Moldenhauer, H.; San Martín, A.; Gutierrez, R.; Zambrano, M.; et al. Fast and Reversible Stimulation of Astrocytic Glycolysis by K⁺ and a Delayed and Persistent Effect of Glutamate. *Journal of Neuroscience* **2011**, *31*, 4709–4713, doi:10.1523/JNEUROSCI.5311-10.2011.
59. Bouzier-Sore, A.-K.; Voisin, P.; Canioni, P.; Magistretti, P.J.; Pellerin, L. Lactate Is a Preferential Oxidative Energy Substrate over Glucose for Neurons in Culture. *J Cereb Blood Flow Metab* **2003**, *23*, 1298–1306, doi:10.1097/01.WCB.0000091761.61714.25.
60. Allaman, I.; Bélanger, M.; Magistretti, P.J. Astrocyte-Neuron Metabolic Relationships: For Better and for Worse. *Trends in Neurosciences* **2011**, *34*, 76–87, doi:10.1016/j.tins.2010.12.001.

61. Fecher, C.; Trovò, L.; Müller, S.A.; Snaidero, N.; Wettmarshausen, J.; Heink, S.; Ortiz, O.; Wagner, I.; Kühn, R.; Hartmann, J.; et al. Cell-Type-Specific Profiling of Brain Mitochondria Reveals Functional and Molecular Diversity. *Nat Neurosci* **2019**, *22*, 1731–1742, doi:10.1038/s41593-019-0479-z.
62. Petrova, V.Y.; Drescher, D.; Kujumdzieva, A.V.; Schmitt, M.J. Dual Targeting of Yeast Catalase A to Peroxisomes and Mitochondria. *Biochemical Journal* **2004**, *380*, 393–400, doi:10.1042/bj20040042
63. Baughman, J.M.; Perocchi, F.; Girgis, H.S.; Plovanich, M.; Belcher-Timme, C.A.; Sancak, Y.; Bao, X.R.; Strittmatter, L.; Goldberger, O.; Bogorad, R.L.; et al. Integrative Genomics Identifies MCU as an Essential Component of the Mitochondrial Calcium Uniporter. *Nature* **2011**, *476*, 341–345, doi:10.1038/nature10234.
64. Lopez-Fabuel, I.; Le Douce, J.; Logan, A.; James, A.M.; Bonvento, G.; Murphy, M.P.; Almeida, A.; Bolaños, J.P. Complex I Assembly into Supercomplexes Determines Differential Mitochondrial ROS Production in Neurons and Astrocytes. *Proc Natl Acad Sci USA* **2016**, *113*, 13063–13068, doi:10.1073/pnas.1613701113.
65. Graham, L.C.; Eaton, S.L.; Brunton, P.J.; Atri, A.; Smith, C.; Lamont, D.J.; Gillingwater, T.H.; Pennetta, G.; Skehel, P.; Wishart, T.M. Proteomic Profiling of Neuronal Mitochondria Reveals Modulators of Synaptic Architecture. *Mol Neurodegeneration* **2017**, *12*, 77, doi:10.1186/s13024-017-0221-9.
66. Stauch, K.L.; Purnell, P.R.; Fox, H.S. Quantitative Proteomics of Synaptic and Nonsynaptic Mitochondria: Insights for Synaptic Mitochondrial Vulnerability. *J. Proteome Res.* **2014**, *13*, 2620–2636, doi:10.1021/pr500295n.
67. Yousefi, R.; Fornasiero, E.F.; Cyganek, L.; Montoya, J.; Jakobs, S.; Rizzoli, S.O.; Rehling, P.; Pacheu-Grau, D. Monitoring Mitochondrial Translation in Living Cells. *EMBO Rep* **2021**, *22*, doi:10.15252/embr.202051635.
68. Inak, G.; Rybak-Wolf, A.; Lisowski, P.; Pentimalli, T.M.; Jüttner, R.; Glazar, P.; Uppal, K.; Bottani, E.; Brunetti, D.; Secker, C.; et al. Defective Metabolic Programming Impairs Early Neuronal Morphogenesis in Neural Cultures and an Organoid Model of Leigh Syndrome. *Nat Commun* **2021**, *12*, 1929, doi:10.1038/s41467-021-22117-z.
69. Papa, S.; Martino, P.L.; Capitanio, G.; Gaballo, A.; De Rasmio, D.; Signorile, A.; Petruzzella, V. The Oxidative Phosphorylation System in Mammalian Mitochondria. In *Advances in Mitochondrial Medicine*; Scatena, R., Bottoni, P., Giardina, B., Eds.; Advances in Experimental Medicine and Biology; Springer Netherlands: Dordrecht, **2012**; Vol. 942, pp. 3–37 ISBN 978-94-007-2868-4.
70. Fernandez-Vizcarra, E.; Zeviani, M. Mitochondrial Disorders of the OXPHOS System. *FEBS Lett* **2021**, *595*, 1062–1106, doi:10.1002/1873-3468.13995.
71. Tang, J.X.; Thompson, K.; Taylor, R.W.; Oláhová, M. Mitochondrial OXPHOS Biogenesis: Co-Regulation of Protein Synthesis, Import, and Assembly Pathways. *IJMS* **2020**, *21*, 3820, doi:10.3390/ijms21113820.
72. Bergman, O.; Ben-Shachar, D. Mitochondrial Oxidative Phosphorylation System (OXPHOS) Deficits in Schizophrenia: Possible Interactions with Cellular Processes. *Can J Psychiatry* **2016**, *61*, 457–469, doi:10.1177/0706743716648290.
73. Chacinska, A.; Koehler, C.M.; Milenkovic, D.; Lithgow, T.; Pfanner, N. Importing Mitochondrial Proteins: Machineries and Mechanisms. *Cell* **2009**, *138*, 628–644, doi:10.1016/j.cell.2009.08.005.
74. Carapito, C.; Kuhn, L.; Karim, L.; Rompais, M.; Rabilloud, T.; Schwenzer, H.; Sissler, M. Two Proteomic Methodologies for Defining N-Termini of Mature Human Mitochondrial Aminoacyl-TRNA Synthetases. *Methods* **2017**, *113*, 111–119, doi:10.1016/j.ymeth.2016.10.012.
75. Russell, O.M.; Gorman, G.S.; Lightowlers, R.N.; Turnbull, D.M. Mitochondrial Diseases: Hope for the Future. *Cell* **2020**, *181*, 168–188, doi:10.1016/j.cell.2020.02.051.
76. Rusecka, J.; Kaliszewska, M.; Bartnik, E.; Tońska, K. Nuclear Genes Involved in Mitochondrial Diseases Caused by Instability of Mitochondrial DNA. *J Appl Genetics* **2018**, *59*, 43–57, doi:10.1007/s13353-017-0424-3.
77. Yasukawa, T.; Kang, D. An Overview of Mammalian Mitochondrial DNA Replication Mechanisms. *The Journal of Biochemistry* **2018**, *164*, 183–193, doi:10.1093/jb/mvy058.
78. Barshad, G.; Marom, S.; Cohen, T.; Mishmar, D. Mitochondrial DNA Transcription and Its Regulation: An Evolutionary Perspective. *Trends in Genetics* **2018**, *34*, 682–692, doi:10.1016/j.tig.2018.05.009.
79. Boczonadi, V.; Ricci, G.; Horvath, R. Mitochondrial DNA Transcription and Translation: Clinical Syndromes. *Essays in Biochemistry* **2018**, *62*, 321–340, doi:10.1042/EBC20170103.
80. Kummer, E.; Ban, N. Mechanisms and Regulation of Protein Synthesis in Mitochondria. *Nat Rev Mol Cell Biol* **2021**, *22*, 307–325, doi:10.1038/s41580-021-00332-2.
81. Wallace, D.C. Bioenergetics in Human Evolution and Disease: Implications for the Origins of Biological Complexity and the Missing Genetic Variation of Common Diseases. *Phil. Trans. R. Soc. B* **2013**, *368*, 20120267, doi:10.1098/rstb.2012.0267.
82. DiMauro, S.; Schon, E.A.; Carelli, V.; Hirano, M. The Clinical Maze of Mitochondrial Neurology. *Nat Rev Neurol* **2013**, *9*, 429–444, doi:10.1038/nrneurol.2013.126.
83. Wallace, D.C. Mitochondrial Genetic Medicine. *Nat Genet* **2018**, *50*, 1642–1649, doi:10.1038/s41588-018-0264-z.
84. Ferreira, C.R.; Rahman, S.; Keller, M.; Zschocke, J.; ICIMD Advisory Group; Abdenur, J.; Ali, H.; Artuch, R.; Ballabio, A.; Barshop, B.; et al. An International Classification of Inherited Metabolic Disorders (ICIMD). *Jrnl of Inher Metab Disea* **2021**, *44*, 164–177, doi:10.1002/jimd.12348.
85. Holt, I.J.; Harding, A.E.; Morgan-Hughes, J.A. Deletions of Muscle Mitochondrial DNA in Patients with Mitochondrial Myopathies. *Nature* **1988**, *331*, 717–719, doi:10.1038/331717a0.
86. Wallace, D.; Singh, G.; Lott, M.; Hodge, J.; Schurr, T.; Lezza, A.; Elsas, L.; Nikoskelainen, E. Mitochondrial DNA Mutation Associated with Leber's Hereditary Optic Neuropathy. *Science* **1988**, *242*, 1427–1430, doi:10.1126/science.3201231.

87. Wallace, D.C.; Zheng, X.; Lott, M.T.; Shoffner, J.M.; Hodge, J.A.; Kelley, R.I.; Epstein, C.M.; Hopkins, L.C. Familial Mitochondrial Encephalomyopathy (MERRF): Genetic, Pathophysiological, and Biochemical Characterization of a Mitochondrial DNA Disease. *Cell* **1988**, *55*, 601–610, doi:10.1016/0092-8674(88)90218-8.
88. Carelli, V.; La Morgia, C. Clinical Syndromes Associated with MtDNA Mutations: Where We Stand after 30 Years. *Essays in Biochemistry* **2018**, *62*, 235–254, doi:10.1042/EBC20170097.
89. Leber, Th. Ueber hereditäre und congenital-angelegte Sehnervenleiden. *Graefe's Archiv für Ophthalmologie* **1871**, *17*, 249–291, doi:10.1007/BF01694557.
90. Holt, I.J.; Harding, A.E.; Petty, R.K.; Morgan-Hughes, J.A. A New Mitochondrial Disease Associated with Mitochondrial DNA Heteroplasmy. *Am J Hum Genet* **1990**, *46*, 428–433.
91. Tatuch, Y.; Christodoulou, J.; Feigenbaum, A.; Clarke, J.T.; Wherret, J.; Smith, C.; Rudd, N.; Petrova-Benedict, R.; Robinson, B.H. Heteroplasmic MtDNA Mutation (T---G) at 8993 Can Cause Leigh Disease When the Percentage of Abnormal MtDNA Is High. *Am J Hum Genet* **1992**, *50*, 852–858.
92. Prezant, T.R.; Agapian, J.V.; Bohlman, M.C.; Bu, X.; Öztas, S.; Qiu, W.-Q.; Arnos, K.S.; Cortopassi, G.A.; Jaber, L.; Rotter, J.I.; et al. Mitochondrial Ribosomal RNA Mutation Associated with Both Antibiotic-Induced and Non-Syndromic Deafness. *Nat Genet* **1993**, *4*, 289–294, doi:10.1038/ng0793-289.
93. Fukuhara, N.; Tokiguchi, S.; Shirakawa, K.; Tsubaki, T. Myoclonus Epilepsy Associated with Ragged-Red Fibres (Mitochondrial Abnormalities): Disease Entity or a Syndrome? *Journal of the Neurological Sciences* **1980**, *47*, 117–133, doi:10.1016/0022-510X(80)90031-3.
94. Shoffner, J.M.; Lott, M.T.; Lezza, A.M.S.; Seibel, P.; Ballinger, S.W.; Wallace, D.C. Myoclonic Epilepsy and Ragged-Red Fiber Disease (MERRF) Is Associated with a Mitochondrial DNA TRNAlys Mutation. *Cell* **1990**, *61*, 931–937, doi:10.1016/0092-8674(90)90059-N.
95. Pavlakis, S.G.; Phillips, P.C.; DiMauro, S.; De Vivo, D.C.; Rowland, L.P. Mitochondrial Myopathy, Encephalopathy, Lactic Acidosis, and Strokelike Episodes: A Distinctive Clinical Syndrome. *Ann Neurol* **1984**, *16*, 481–488, doi:10.1002/ana.410160409.
96. Goto, Y.; Nonaka, I.; Horai, S. A Mutation in the TRNAlau(UUR) Gene Associated with the MELAS Subgroup of Mitochondrial Encephalomyopathies. *Nature* **1990**, *348*, 651–653, doi:10.1038/348651a0.
97. Kearns, T.P.; Sayre, G.P. Retinitis Pigmentosa, External Ophthalmoplegia, and Complete Heart Block: Unusual Syndrome with Histologic Study in One of Two Cases. *AMA Arch Ophthalmol* **1958**, *60*, 280–289.
98. Zeviani, M.; Moraes, C.T.; DiMauro, S.; Nakase, H.; Bonilla, E.; Schon, E.A.; Rowland, L.P. Deletions of Mitochondrial DNA in Kearns-Sayre Syndrome. *Neurology* **1988**, *38*, 1339–1339, doi:10.1212/WNL.38.9.1339.
99. Pearson, H.A.; Lobel, J.S.; Kocoshis, S.A.; Naiman, J.L.; Windmiller, J.; Lammi, A.T.; Hoffman, R.; Marsh, J.C. A New Syndrome of Refractory Sideroblastic Anemia with Vacuolization of Marrow Precursors and Exocrine Pancreatic Dysfunction. *The Journal of Pediatrics* **1979**, *95*, 976–984, doi:10.1016/S0022-3476(79)80286-3.
100. Rötig, A.; Cormier, V.; Blanche, S.; Bonnefont, J.P.; Ledest, F.; Romero, N.; Schmitz, J.; Rustin, P.; Fischer, A.; Saudubray, J.M. Pearson's Marrow-Pancreas Syndrome. A Multisystem Mitochondrial Disorder in Infancy. *J. Clin. Invest.* **1990**, *86*, 1601–1608, doi:10.1172/JCI114881.
101. Carelli, V.; Ghelli, A.; Ratta, M.; Bacchilega, E.; Sangiorgi, S.; Mancini, R.; Leuzzi, V.; Cortelli, P.; Montagna, P.; Lugaresi, E.; et al. Leber's Hereditary Optic Neuropathy: Biochemical Effect of 11778/ND4 and 3460/ND1 Mutations and Correlation with the Mitochondrial Genotype. *Neurology* **1997**, *48*, 1623–1632, doi:10.1212/WNL.48.6.1623.
102. Tatuch, Y.; Robinson, B.H. The Mitochondrial DNA Mutation at 8993 Associated with NARP Slows the Rate of ATP Synthesis in Isolated Lymphoblast Mitochondria. *Biochemical and Biophysical Research Communications* **1993**, *192*, 124–128, doi:10.1006/bbrc.1993.1390.
103. Bernes, S.M.; Bacino, C.; Prezant, T.R.; Pearson, M.A.; Wood, T.S.; Fournier, P.; Fischel-Ghodsian, N. Identical Mitochondrial DNA Deletion in Mother with Progressive External Ophthalmoplegia and Son with Pearson Marrow-Pancreas Syndrome. *The Journal of Pediatrics* **1993**, *123*, 598–602, doi:10.1016/S0022-3476(05)80962-X.
104. Shanske, S.; Tang, Y.; Hirano, M.; Nishigaki, Y.; Tanji, K.; Bonilla, E.; Sue, C.; Krishna, S.; Carlo, J.R.; Willner, J.; et al. Identical Mitochondrial DNA Deletion in a Woman with Ocular Myopathy and in Her Son with Pearson Syndrome. *The American Journal of Human Genetics* **2002**, *71*, 679–683, doi:10.1086/342482.
105. Hirst, J. Mitochondrial Complex I. *Annu. Rev. Biochem.* **2013**, *82*, 551–575, doi:10.1146/annurev-biochem-070511-103700.
106. Ripple, M.O.; Kim, N.; Springett, R. Mammalian Complex I Pumps 4 Protons per 2 Electrons at High and Physiological Proton Motive Force in Living Cells*. *Journal of Biological Chemistry* **2013**, *288*, 5374–5380, doi:10.1074/jbc.M112.438945.
107. Efremov, R.G.; Baradaran, R.; Sazanov, L.A. The Architecture of Respiratory Complex I. *Nature* **2010**, *465*, 441–445, doi:10.1038/nature09066.
108. Zhu, J.; Vinothkumar, K.R.; Hirst, J. Structure of Mammalian Respiratory Complex I. *Nature* **2016**, *536*, 354–358, doi:10.1038/nature19095.
109. Vinothkumar, K.R.; Zhu, J.; Hirst, J. Architecture of Mammalian Respiratory Complex I. *Nature* **2014**, *515*, 80–84, doi:10.1038/nature13686.
110. Clason, T.; Ruiz, T.; Schagger, H.; Peng, G.; Zickermann, V.; Brandt, U.; Michel, H.; Radermacher, M. The Structure of Eukaryotic and Prokaryotic Complex I. *Journal of Structural Biology* **2010**, *169*, 81–88, doi:10.1016/j.jsb.2009.08.017.
111. Baradaran, R.; Berrisford, J.M.; Minhas, G.S.; Sazanov, L.A. Crystal Structure of the Entire Respiratory Complex I. *Nature* **2013**, *494*, 443–448, doi:10.1038/nature11871.

112. Agip, A.-N.A.; Blaza, J.N.; Bridges, H.R.; Viscomi, C.; Rawson, S.; Muench, S.P.; Hirst, J. Cryo-EM Structures of Complex I from Mouse Heart Mitochondria in Two Biochemically Defined States. *Nat Struct Mol Biol* **2018**, *25*, 548–556, doi:10.1038/s41594-018-0073-1.
113. Hirst, J.; Roessler, M.M. Energy Conversion, Redox Catalysis and Generation of Reactive Oxygen Species by Respiratory Complex I. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2016**, *1857*, 872–883, doi:10.1016/j.bbabi.2015.12.009.
114. Stroud, D.A.; Surgenor, E.E.; Formosa, L.E.; Reljic, B.; Frazier, A.E.; Dibley, M.G.; Osellame, L.D.; Stait, T.; Beilharz, T.H.; Thorburn, D.R.; et al. Accessory Subunits Are Integral for Assembly and Function of Human Mitochondrial Complex I. *Nature* **2016**, *538*, 123–126, doi:10.1038/nature19754.
115. Yip, C.; Harbour, M.E.; Jayawardena, K.; Fearnley, I.M.; Sazanov, L.A. Evolution of Respiratory Complex I. *Journal of Biological Chemistry* **2011**, *286*, 5023–5033, doi:10.1074/jbc.M110.194993.
116. Sazanov, L.A. A Giant Molecular Proton Pump: Structure and Mechanism of Respiratory Complex I. *Nat Rev Mol Cell Biol* **2015**, *16*, 375–388, doi:10.1038/nrm3997.
117. Zickermann, V.; Wirth, C.; Nasiri, H.; Siegmund, K.; Schwalbe, H.; Hunte, C.; Brandt, U. Mechanistic Insight from the Crystal Structure of Mitochondrial Complex I. *Science* **2015**, *347*, 44–49, doi:10.1126/science.1259859.
118. Parey, K.; Wirth, C.; Vonck, J.; Zickermann, V. Respiratory Complex I — Structure, Mechanism and Evolution. *Current Opinion in Structural Biology* **2020**, *63*, 1–9, doi:10.1016/j.sbi.2020.01.004.
119. Kampjut, D.; Sazanov, L.A. The Coupling Mechanism of Mammalian Respiratory Complex I. *Science* **2020**, *370*, eabc4209, doi:10.1126/science.abc4209.
120. Grba, D.N.; Hirst, J. Mitochondrial Complex I Structure Reveals Ordered Water Molecules for Catalysis and Proton Translocation. *Nat Struct Mol Biol* **2020**, *27*, 892–900, doi:10.1038/s41594-020-0473-x.
121. Klusch, N.; Senkler, J.; Yildiz, Ö.; Kühlbrandt, W.; Braun, H.-P. A Ferredoxin Bridge Connects the Two Arms of Plant Mitochondrial Complex I; *Plant Biology*, 2020;
122. Soufari, H.; Parrot, C.; Kuhn, L.; Waltz, F.; Hashem, Y. Specific Features and Assembly of the Plant Mitochondrial Complex I Revealed by Cryo-EM. *Nat Commun* **2020**, *11*, 5195, doi:10.1038/s41467-020-18814-w.
123. Guo, R.; Zong, S.; Wu, M.; Gu, J.; Yang, M. Architecture of Human Mitochondrial Respiratory Megacomplex I₂III₂IV₂. *Cell* **2017**, *170*, 1247–1257.e12, doi:10.1016/j.cell.2017.07.050.
124. Signes, A.; Fernandez-Vizarra, E. Assembly of Mammalian Oxidative Phosphorylation Complexes I–V and Supercomplexes. *Essays in Biochemistry* **2018**, *62*, 255–270, doi:10.1042/EBC20170098.
125. Zurita Rendón, O.; Antonicka, H.; Horvath, R.; Shoubridge, E.A. A Mutation in the Flavin Adenine Dinucleotide-Dependent Oxidoreductase FOXRED1 Results in Cell-Type-Specific Assembly Defects in Oxidative Phosphorylation Complexes I and II. *Mol. Cell. Biol.* **2016**, *36*, 2132–2140, doi:10.1128/MCB.00066-16.
126. Vartak, R.S.; Semwal, M.K.; Bai, Y. An Update on Complex I Assembly: The Assembly of Players. *J Bioenerg Biomembr* **2014**, *46*, 323–328, doi:10.1007/s10863-014-9564-x.
127. Carroll, J.; Ding, S.; Fearnley, I.M.; Walker, J.E. Post-Translational Modifications near the Quinone Binding Site of Mammalian Complex I*. *Journal of Biological Chemistry* **2013**, *288*, 24799–24808, doi:10.1074/jbc.M113.488106.
128. Nouws, J.; Nijtmans, L.; Houten, S.M.; van den Brand, M.; Huynen, M.; Venselaar, H.; Hoefs, S.; Gloerich, J.; Kronick, J.; Hutchin, T.; et al. Acyl-CoA Dehydrogenase 9 Is Required for the Biogenesis of Oxidative Phosphorylation Complex I. *Cell Metabolism* **2010**, *12*, 283–294, doi:10.1016/j.cmet.2010.08.002.
129. Haack, T.B.; Danhauser, K.; Haberberger, B.; Hoser, J.; Strecker, V.; Boehm, D.; Uziel, G.; Lamantea, E.; Invernizzi, F.; Poulton, J.; et al. Exome Sequencing Identifies ACAD9 Mutations as a Cause of Complex I Deficiency. *Nat Genet* **2010**, *42*, 1131–1134, doi:10.1038/ng.706.
130. Vogel, R.O.; Janssen, R.J.R.J.; van den Brand, M.A.M.; Dieteren, C.E.J.; Verkaart, S.; Koopman, W.J.H.; Willems, P.H.G.M.; Pluk, W.; van den Heuvel, L.P.W.J.; Smeitink, J.A.M.; et al. Cytosolic Signaling Protein Ecsit Also Localizes to Mitochondria Where It Interacts with Chaperone NDUFAF1 and Functions in Complex I Assembly. *Genes & Development* **2007**, *21*, 615–624, doi:10.1101/gad.408407.
131. Formosa, L.E.; Mimaki, M.; Frazier, A.E.; McKenzie, M.; Stait, T.L.; Thorburn, D.R.; Stroud, D.A.; Ryan, M.T. Characterization of Mitochondrial FOXRED1 in the Assembly of Respiratory Chain Complex I. *Human Molecular Genetics* **2015**, *24*, 2952–2965, doi:10.1093/hmg/ddv058.
132. Calvo, S.E.; Tucker, E.J.; Compton, A.G.; Kirby, D.M.; Crawford, G.; Burt, N.P.; Rivas, M.; Guiducci, C.; Bruno, D.L.; Goldberger, O.A.; et al. High-Throughput, Pooled Sequencing Identifies Mutations in NUBPL and FOXRED1 in Human Complex I Deficiency. *Nat Genet* **2010**, *42*, 851–858, doi:10.1038/ng.659.
133. Andrews, B.; Carroll, J.; Ding, S.; Fearnley, I.M.; Walker, J.E. Assembly Factors for the Membrane Arm of Human Complex I. *Proceedings of the National Academy of Sciences* **2013**, *110*, 18934–18939, doi:10.1073/pnas.1319247110.
134. Čížková, A.; Stránecký, V.; Mayr, J.A.; Tesařová, M.; Havlíčková, V.; Paul, J.; Ivánek, R.; Kuss, A.W.; Hansíková, H.; Kaplanová, V.; et al. TMEM70 Mutations Cause Isolated ATP Synthase Deficiency and Neonatal Mitochondrial Encephalomyopathy. *Nat Genet* **2008**, *40*, 1288–1290, doi:10.1038/ng.246.
135. Sánchez-Caballero, L.; Elurbe, D.M.; Baertling, F.; Guerrero-Castillo, S.; van den Brand, M.; van Strien, J.; van Dam, T.J.P.; Rodenburg, R.; Brandt, U.; Huynen, M.A.; et al. TMEM70 Functions in the Assembly of Complexes I and V. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2020**, *1861*, 148202, doi:10.1016/j.bbabi.2020.148202.

136. Vogel, R.O.; Janssen, R.J.R.J.; Ugalde, C.; Grovenstein, M.; Huijbens, R.J.; Visch, H.-J.; van den Heuvel, L.P.; Willems, P.H.; Zeviani, M.; Smeitink, J.A.M.; et al. Human Mitochondrial Complex I Assembly Is Mediated by NDUFAF1. *FEBS Journal* **2005**, *272*, 5317–5326, doi:10.1111/j.1742-4658.2005.04928.x.
137. Dunning, C.J.R.; McKenzie, M.; Sugiana, C.; Lazarou, M.; Silke, J.; Connelly, A.; Fletcher, J.M.; Kirby, D.M.; Thorburn, D.R.; Ryan, M.T. Human CIA30 Is Involved in the Early Assembly of Mitochondrial Complex I and Mutations in Its Gene Cause Disease. *EMBO J* **2007**, *26*, 3227–3237, doi:10.1038/sj.emboj.7601748.
138. Ogilvie, I. A Molecular Chaperone for Mitochondrial Complex I Assembly Is Mutated in a Progressive Encephalopathy. *Journal of Clinical Investigation* **2005**, *115*, 2784–2792, doi:10.1172/JCI26020.
139. Saada, A.; Edvardson, S.; Rapoport, M.; Shaag, A.; Amry, K.; Miller, C.; Lorberboum-Galski, H.; Elpeleg, O. C6ORF66 Is an Assembly Factor of Mitochondrial Complex I. *The American Journal of Human Genetics* **2008**, *82*, 32–38, doi:10.1016/j.ajhg.2007.08.003.
140. Saada, A.; Vogel, R.O.; Hoefs, S.J.; van den Brand, M.A.; Wessels, H.J.; Willems, P.H.; Venselaar, H.; Shaag, A.; Barghuti, F.; Reish, O.; et al. Mutations in NDUFAF3 (C3ORF60), Encoding an NDUFAF4 (C6ORF66)-Interacting Complex I Assembly Protein, Cause Fatal Neonatal Mitochondrial Disease. *The American Journal of Human Genetics* **2009**, *84*, 718–727, doi:10.1016/j.ajhg.2009.04.020.
141. Sugiana, C.; Pagliarini, D.J.; McKenzie, M.; Kirby, D.M.; Salemi, R.; Abu-Amero, K.K.; Dahl, H.-H.M.; Hutchison, W.M.; Vascotto, K.A.; Smith, S.M.; et al. Mutation of C20orf7 Disrupts Complex I Assembly and Causes Lethal Neonatal Mitochondrial Disease. *The American Journal of Human Genetics* **2008**, *83*, 468–478, doi:10.1016/j.ajhg.2008.09.009.
142. Rhein, V.F.; Carroll, J.; Ding, S.; Fearnley, I.M.; Walker, J.E. NDUFAF5 Hydroxylates NDUF57 at an Early Stage in the Assembly of Human Complex I. *Journal of Biological Chemistry* **2016**, *291*, 14851–14860, doi:10.1074/jbc.M116.734970.
143. McKenzie, M.; Tucker, E.J.; Compton, A.G.; Lazarou, M.; George, C.; Thorburn, D.R.; Ryan, M.T. Mutations in the Gene Encoding C8orf38 Block Complex I Assembly by Inhibiting Production of the Mitochondria-Encoded Subunit ND1. *Journal of Molecular Biology* **2011**, *414*, 413–426, doi:10.1016/j.jmb.2011.10.012.
144. Carilla-Latorre, S.; Gallardo, M.E.; Annesley, S.J.; Calvo-Garrido, J.; Graña, O.; Accari, S.L.; Smith, P.K.; Valencia, A.; Garesse, R.; Fisher, P.R.; et al. MidA Is a Putative Methyltransferase That Is Required for Mitochondrial Complex I Function. *Journal of Cell Science* **2010**, *123*, 1674–1683, doi:10.1242/jcs.066076.
145. Rhein, V.F.; Carroll, J.; Ding, S.; Fearnley, I.M.; Walker, J.E. NDUFAF7 Methylates Arginine 85 in the NDUF52 Subunit of Human Complex I. *Journal of Biological Chemistry* **2013**, *288*, 33016–33026, doi:10.1074/jbc.M113.518803.
146. Alston, C.L.; Veling, M.T.; Heidler, J.; Taylor, L.S.; Alaimo, J.T.; Sung, A.Y.; He, L.; Hopton, S.; Broomfield, A.; Pavaine, J.; et al. Pathogenic Bi-Allelic Mutations in NDUFAF8 Cause Leigh Syndrome with an Isolated Complex I Deficiency. *The American Journal of Human Genetics* **2020**, *106*, 92–101, doi:10.1016/j.ajhg.2019.12.001.
147. Sheftel, A.D.; Stehling, O.; Pierik, A.J.; Netz, D.J.A.; Kerscher, S.; Elsässer, H.-P.; Wittig, I.; Balk, J.; Brandt, U.; Lill, R. Human Ind1, an Iron-Sulfur Cluster Assembly Factor for Respiratory Complex I. *MCB* **2009**, *29*, 6059–6073, doi:10.1128/MCB.00817-09.
148. Bych, K.; Kerscher, S.; Netz, D.J.A.; Pierik, A.J.; Zwicker, K.; Huynen, M.A.; Lill, R.; Brandt, U.; Balk, J. The Iron-Sulphur Protein Ind1 Is Required for Effective Complex I Assembly. *EMBO J* **2008**, *27*, 1736–1746, doi:10.1038/emboj.2008.98.
149. Protasoni, M.; Bruno, C.; Donati, M.A.; Mohamoud, K.; Severino, M.; Allegri, A.; Robinson, A.J.; Reyes, A.; Zeviani, M.; Garone, C. Novel Compound Heterozygous Pathogenic Variants in Nucleotide-Binding Protein like Protein (NUBPL) Cause Leukoencephalopathy with Multi-Systemic Involvement. *Molecular Genetics and Metabolism* **2020**, *129*, 26–34, doi:10.1016/j.ymgme.2019.11.003.
150. Guarani, V.; Paulo, J.; Zhai, B.; Huttlin, E.L.; Gygi, S.P.; Harper, J.W. TIMMDC1/C3orf1 Functions as a Membrane-Embedded Mitochondrial Complex I Assembly Factor through Association with the MCIA Complex. *Mol. Cell. Biol.* **2014**, *34*, 847–861, doi:10.1128/MCB.01551-13.
151. Kremer, L.S.; Bader, D.M.; Mertes, C.; Kopajtich, R.; Pichler, G.; Iuso, A.; Haack, T.B.; Graf, E.; Schwarzmayr, T.; Terrile, C.; et al. Genetic Diagnosis of Mendelian Disorders via RNA Sequencing. *Nat Commun* **2017**, *8*, 15824, doi:10.1038/ncomms15824.
152. Désir, J.; Coppeters, F.; Van Regemorter, N.; De Baere, E.; Abramowicz, M.; Cordonnier, M. TMEM126A Mutation in a Moroccan Family with Autosomal Recessive Optic Atrophy. *Mol Vis* **2012**, *18*, 1849–1857.
153. Hanein, S.; Perrault, I.; Roche, O.; Gerber, S.; Khadom, N.; Rio, M.; Boddaert, N.; Jean-Pierre, M.; Brahimi, N.; Serre, V.; et al. TMEM126A, Encoding a Mitochondrial Protein, Is Mutated in Autosomal-Recessive Nonsyndromic Optic Atrophy. *The American Journal of Human Genetics* **2009**, *84*, 493–498, doi:10.1016/j.ajhg.2009.03.003.
154. Kloth, K.; Synofzik, M.; Kernstock, C.; Schimpf-Linzenbold, S.; Schuettauf, F.; Neu, A.; Wissinger, B.; Weisschuh, N. Novel Likely Pathogenic Variants in TMEM126A Identified in Non-Syndromic Autosomal Recessive Optic Atrophy: Two Case Reports. *BMC Med Genet* **2019**, *20*, 62, doi:10.1186/s12881-019-0795-x.
155. La Morgia, C.; Caporali, L.; Tagliavini, F.; Palombo, F.; Carbonelli, M.; Liguori, R.; Barboni, P.; Carelli, V. First TMEM126A Missense Mutation in an Italian Proband with Optic Atrophy and Deafness. *Neurol Genet* **2019**, *5*, e329, doi:10.1212/NXG.0000000000000329.
156. Meyer, E.; Michaelides, M.; Tee, L.J.; Robson, A.G.; Rahman, F.; Pasha, S.; Luxon, L.M.; Moore, A.T.; Maher, E.R. Nonsense Mutation in TMEM126A Causing Autosomal Recessive Optic Atrophy and Auditory Neuropathy. *Mol Vis* **2010**, *16*, 650–664.
157. Heide, H.; Bleier, L.; Steger, M.; Ackermann, J.; Dröse, S.; Schwamb, B.; Zörnig, M.; Reichert, A.S.; Koch, I.; Wittig, I.; et al. Complexome Profiling Identifies TMEM126B as a Component of the Mitochondrial Complex I Assembly Complex. *Cell Metabolism* **2012**, *16*, 538–549, doi:10.1016/j.cmet.2012.08.009.

158. Sánchez-Caballero, L.; Ruzzenente, B.; Bianchi, L.; Assouline, Z.; Barcia, G.; Metodiev, M.D.; Rio, M.; Funalot, B.; van den Brand, M.A.M.; Guerrero-Castillo, S.; et al. Mutations in Complex I Assembly Factor TMEM126B Result in Muscle Weakness and Isolated Complex I Deficiency. *The American Journal of Human Genetics* **2016**, *99*, 208–216, doi:10.1016/j.ajhg.2016.05.022.
159. Alston, C.L.; Compton, A.G.; Formosa, L.E.; Strecker, V.; Oláhová, M.; Haack, T.B.; Smet, J.; Stouffs, K.; Diakumis, P.; Ciara, E.; et al. Biallelic Mutations in TMEM126B Cause Severe Complex I Deficiency with a Variable Clinical Phenotype. *The American Journal of Human Genetics* **2016**, *99*, 217–227, doi:10.1016/j.ajhg.2016.05.021.
160. Guerrero-Castillo, S.; Baertling, F.; Kownatzki, D.; Wessels, H.J.; Arnold, S.; Brandt, U.; Nijtmans, L. The Assembly Pathway of Mitochondrial Respiratory Chain Complex I. *Cell Metabolism* **2017**, *25*, 128–139, doi:10.1016/j.cmet.2016.09.002.
161. Martinez Lyons, A.; Ardisson, A.; Reyes, A.; Robinson, A.J.; Moroni, I.; Ghezzi, D.; Fernandez-Vizarra, E.; Zeviani, M. COA7 (C1orf163 /RESA1) Mutations Associated with Mitochondrial Leukoencephalopathy and Cytochrome c Oxidase Deficiency. *J Med Genet* **2016**, *53*, 846–849, doi:10.1136/jmedgenet-2016-104194.
162. Dang, Q.-C.L.; Phan, D.H.; Johnson, A.N.; Pasapuleti, M.; Alkhalidi, H.A.; Zhang, F.; Vik, S.B. Analysis of Human Mutations in the Supernumerary Subunits of Complex I. *Life (Basel)* **2020**, *10*, doi:10.3390/life10110296.
163. Bugiani, M.; Invernizzi, F.; Alberio, S.; Briem, E.; Lamantea, E.; Carrara, F.; Moroni, I.; Farina, L.; Spada, M.; Donati, M.A.; et al. Clinical and Molecular Findings in Children with Complex I Deficiency. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2004**, *1659*, 136–147, doi:10.1016/j.bbabo.2004.09.006.
164. Malfatti, E.; Bugiani, M.; Invernizzi, F.; de Souza, C.F.-M.; Farina, L.; Carrara, F.; Lamantea, E.; Antozzi, C.; Confalonieri, P.; Sanseverino, M.T.; et al. Novel Mutations of ND Genes in Complex I Deficiency Associated with Mitochondrial Encephalopathy. *Brain* **2007**, *130*, 1894–1904, doi:10.1093/brain/awm114.
165. Fassone, E.; Rahman, S. Complex I Deficiency: Clinical Features, Biochemistry and Molecular Genetics. *J Med Genet* **2012**, *49*, 578–590, doi:10.1136/jmedgenet-2012-101159.
166. Rodenburg, R.J. Mitochondrial Complex I-Linked Disease. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2016**, *1857*, 938–945, doi:10.1016/j.bbabo.2016.02.012.
167. Yu-Wai-Man, P.; Griffiths, P.G.; Brown, D.T.; Howell, N.; Turnbull, D.M.; Chinnery, P.F. The Epidemiology of Leber Hereditary Optic Neuropathy in the North East of England. *Am J Hum Genet* **2003**, *72*, 333–339, doi:10.1086/346066.
168. Carelli, V.; Rugolo, M.; Sgarbi, G.; Ghelli, A.; Zanna, C.; Baracca, A.; Lenaz, G.; Napoli, E.; Martinuzzi, A.; Solaini, G. Bioenergetics Shapes Cellular Death Pathways in Leber's Hereditary Optic Neuropathy: A Model of Mitochondrial Neurodegeneration. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2004**, *1658*, 172–179, doi:10.1016/j.bbabo.2004.05.009.
169. Yu-Wai-Man, P.; Griffiths, P.G.; Chinnery, P.F. Mitochondrial Optic Neuropathies – Disease Mechanisms and Therapeutic Strategies. *Progress in Retinal and Eye Research* **2011**, *30*, 81–114, doi:10.1016/j.preteyeres.2010.11.002.
170. Giordano, C.; Iommarini, L.; Giordano, A.; Pisano, A.; Valentino, M.L.; Caporali, L.; Liguori, R.; Deceglie, S.; Roberti, M.; et al. Efficient Mitochondrial Biogenesis Drives Incomplete Penetrance in Leber's Hereditary Optic Neuropathy. *Brain* **2014**, *137*, 335–353, doi:10.1093/brain/awt343.
171. Bianco, A.; Martínez-Romero, I.; Bisceglia, L.; D'Agruma, L.; Favia, P.; Ruiz-Pesini, E.; Guerriero, S.; Montoya, J.; Petruzzella, V. Mitochondrial DNA Copy Number Differentiates the Leber's Hereditary Optic Neuropathy Affected Individuals from the Unaffected Mutation Carriers. *Brain* **2016**, *139*, e1–e1, doi:10.1093/brain/awv216.
172. Bianco, A.; Bisceglia, L.; Russo, L.; Palese, L.L.; D'Agruma, L.; Emperador, S.; Montoya, J.; Guerriero, S.; Petruzzella, V. High Mitochondrial DNA Copy Number Is a Protective Factor From Vision Loss in Heteroplasmic Leber's Hereditary Optic Neuropathy (LHON). *Invest. Ophthalmol. Vis. Sci.* **2017**, *58*, 2193, doi:10.1167/iovs.16-20389.
173. Bianco, A.; Valletti, A.; Longo, G.; Bisceglia, L.; Montoya, J.; Emperador, S.; Guerriero, S.; Petruzzella, V. Mitochondrial DNA Copy Number in Affected and Unaffected LHON Mutation Carriers. *BMC Res Notes* **2018**, *11*, 911, doi:10.1186/s13104-018-4025-y.
174. Tun, A.W.; Chaiyari, S.; Kaewsutthi, S.; Katanyoo, W.; Chuenkongkaew, W.; Kuwano, M.; Tomonaga, T.; Peerapittayamongkol, C.; Thongboonkerd, V.; Lertrit, P. Profiling the Mitochondrial Proteome of Leber's Hereditary Optic Neuropathy (LHON) in Thailand: Down-Regulation of Bioenergetics and Mitochondrial Protein Quality Control Pathways in Fibroblasts with the 11778G>A Mutation. *PLoS ONE* **2014**, *9*, e106779, doi:10.1371/journal.pone.0106779.
175. Lenaz, G.; Baracca, A.; Carelli, V.; D'Aurelio, M.; Sgarbi, G.; Solaini, G. Bioenergetics of Mitochondrial Diseases Associated with MtDNA Mutations. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2004**, *1658*, 89–94, doi:10.1016/j.bbabo.2004.03.013.
176. Brown, M.D.; Trounce, I.A.; Jun, A.S.; Allen, J.C.; Wallace, D.C. Functional Analysis of Lymphoblast and Cybrid Mitochondria Containing the 3460, 11778, or 14484 Leber's Hereditary Optic Neuropathy Mitochondrial DNA Mutation. *Journal of Biological Chemistry* **2000**, *275*, 39831–39836, doi:10.1074/jbc.M006476200.
177. Floreani, M.; Napoli, E.; Martinuzzi, A.; Pantano, G.; De Riva, V.; Trevisan, R.; Bisetto, E.; Valente, L.; Carelli, V.; Dabbeni-Sala, F. Antioxidant Defences in Cybrids Harboring MtDNA Mutations Associated with Leber's Hereditary Optic Neuropathy: Antioxidant Defences in LHON Cybrids. *FEBS Journal* **2005**, *272*, 1124–1135, doi:10.1111/j.1742-4658.2004.04542.x.
178. Simon, DavidK.; Friedman, J.; Breakefield, XandraO.; Jankovic, J.; Brin, MitchellF.; Provias, J.; Bressman, SusanB.; Charness, MichaelE.; Tarsy, D.; Johns, DonaldR.; et al. A Heteroplasmic Mitochondrial Complex I Gene Mutation in Adult-Onset Dystonia. *neurogenetics* **2003**, *4*, 199–205, doi:10.1007/s10048-003-0150-3.
179. Kirby, D.M. Mutations of the Mitochondrial ND1 Gene as a Cause of MELAS. *Journal of Medical Genetics* **2004**, *41*, 784–789, doi:10.1136/jmg.2004.020537.

180. Howell, N.; Bindoff, L.A.; McCullough, D.A.; Kubacka, I.; Poulton, J.; Mackey, D.; Taylor, L.; Turnbull, D.M. Leber Hereditary Optic Neuropathy: Identification of the Same Mitochondrial ND1 Mutation in Six Pedigrees. *Am J Hum Genet* **1991**, *49*, 939–950.
181. Johns, D.R.; Berman, J. Alternative, Simultaneous Complex I Mitochondrial DNA Mutations in Leber's Hereditary Optic Neuropathy. *Biochemical and Biophysical Research Communications* **1991**, *174*, 1324–1330, doi:10.1016/0006-291X(91)91567-V.
182. McFarland, R.; Kirby, D.M.; Fowler, K.J.; Ohtake, A.; Ryan, M.T.; Amor, D.J.; Fletcher, J.M.; Dixon, J.W.; Collins, F.A.; Turnbull, D.M.; et al. De Novo Mutations in the Mitochondrial ND3 Gene as a Cause of Infantile Mitochondrial Encephalopathy and Complex I Deficiency. *Ann Neurol* **2004**, *55*, 58–64, doi:10.1002/ana.10787.
183. Torroni, A.; Petrozzi, M.; D'Urbano, L.; Sellitto, D.; Zeviani, M.; Carrara, F.; Carducci, C.; Leuzzi, V.; Carelli, V.; Barboni, P.; et al. Haplotype and Phylogenetic Analyses Suggest That One European-Specific MtDNA Background Plays a Role in the Expression of Leber Hereditary Optic Neuropathy by Increasing the Penetrance of the Primary Mutations 11778 and 14484. *Am J Hum Genet* **1997**, *60*, 1107–1121.
184. Lertrit, P.; Noer, A.S.; Jean-Francois, M.J.; Kapsa, R.; Dennett, X.; Thyagarajan, D.; Lethlean, K.; Byrne, E.; Marzuki, S. A New Disease-Related Mutation for Mitochondrial Encephalopathy Lactic Acidosis and Strokelike Episodes (MELAS) Syndrome Affects the ND4 Subunit of the Respiratory Complex I. *Am J Hum Genet* **1992**, *51*, 457–468.
185. Brown, M.D.; Starikovskaya, E.; Derbeneva, O.; Hosseini, S.; Allen, J.C.; Mikhailovskaya, I.E.; Sukernik, R.I.; Wallace, D.C. The Role of MtDNA Background in Disease Expression: A New Primary LHON Mutation Associated with Western Eurasian Haplogroup J. *Hum Genet* **2002**, *110*, 130–138, doi:10.1007/s00439-001-0660-8.
186. Brown, M.D.; Voljavec, A.S.; Lott, M.T.; Macdonald, I.; Wallace, D.C. Leber's Hereditary Optic Neuropathy: A Model for Mitochondrial Neurodegenerative Diseases. *FASEB j.* **1992**, *6*, 2791–2799, doi:10.1096/fasebj.6.10.1634041.
187. Liolitsa, D.; Rahman, S.; Benton, S.; Carr, L.J.; Hanna, M.G. Is the Mitochondrial Complex I ND5 Gene a Hot-Spot for MELAS Causing Mutations? *Ann Neurol* **2003**, *53*, 128–132, doi:10.1002/ana.10435.
188. Ravn, K.; Wibbrand, F.; Hansen, F.J.; Horn, N.; Rosenberg, T.; Schwartz, M. An MtDNA Mutation, 14453G→A, in the NADH Dehydrogenase Subunit 6 Associated with Severe MELAS Syndrome. *Eur J Hum Genet* **2001**, *9*, 805–809, doi:10.1038/sj.ejhg.5200712.
189. Schuelke, M.; Smeitink, J.; Mariman, E.; Loeffen, J.; Plecko, B.; Trijbels, F.; Stöckler-Ipsiroglu, S.; Heuvel, L. van den Mutant NDUFV1 Subunit of Mitochondrial Complex I Causes Leukodystrophy and Myoclonic Epilepsy. *Nat Genet* **1999**, *21*, 260–261, doi:10.1038/6772.
190. Bénit, P.; Chretien, D.; Kadhon, N.; de Lonlay-Debeney, P.; Cormier-Daire, V.; Cabral, A.; Peudener, S.; Rustin, P.; Munnich, A.; Rötig, A. Large-Scale Deletion and Point Mutations of the Nuclear NDUFV1 and NDUF51 Genes in Mitochondrial Complex I Deficiency. *The American Journal of Human Genetics* **2001**, *68*, 1344–1352, doi:10.1086/320603.
191. Bénit, P.; Beugnot, R.; Chretien, D.; Giurgea, I.; De Lonlay-Debeney, P.; Issartel, J.-P.; Corral-Debrinski, M.; Kersch, S.; Rustin, P.; Rötig, A.; et al. Mutant NDUFV2 Subunit of Mitochondrial Complex I Causes Early Onset Hypertrophic Cardiomyopathy and Encephalopathy: NDUFV2 AND CARDIOMYOPATHY/ENCEPHALOPATHY. *Hum. Mutat.* **2003**, *21*, 582–586, doi:10.1002/humu.10225.
192. Loeffen, J.; Elpeleg, O.; Smeitink, J.; Smeets, R.; Stöckler-Ipsiroglu, S.; Mandel, H.; Sengers, R.; Trijbels, F.; van den Heuvel, L. Mutations in the Complex I NDUF52 Gene of Patients with Cardiomyopathy and Encephalomyopathy. *Ann Neurol* **2001**, *49*, 195–201, doi:10.1002/1531-8249(20010201)49:2<195::aid-ana39>3.0.co;2-m.
193. Benit, P. Mutant NDUF53 Subunit of Mitochondrial Complex I Causes Leigh Syndrome. *Journal of Medical Genetics* **2004**, *41*, 14–17, doi:10.1136/jmg.2003.014316.
194. Budde, S.M.S.; van den Heuvel, L.P.W.J.; Janssen, A.J.; Smeets, R.J.P.; Buskens, C.A.F.; DeMeirleir, L.; Van Coster, R.; Baethmann, M.; Voit, T.; Trijbels, J.M.F.; et al. Combined Enzymatic Complex I and III Deficiency Associated with Mutations in the Nuclear Encoded NDUF54 Gene. *Biochemical and Biophysical Research Communications* **2000**, *275*, 63–68, doi:10.1006/bbrc.2000.3257.
195. Spiegel, R.; Shaag, A.; Mandel, H.; Reich, D.; Penyakov, M.; Hujeirat, Y.; Saada, A.; Elpeleg, O.; Shalev, S.A. Mutated NDUF56 Is the Cause of Fatal Neonatal Lactic Acidemia in Caucasus Jews. *Eur J Hum Genet* **2009**, *17*, 1200–1203, doi:10.1038/ejhg.2009.24.
196. Taylor, R.W.; Pyle, A.; Griffin, H.; Blakely, E.L.; Duff, J.; He, L.; Smertenko, T.; Alston, C.L.; Neeve, V.C.; Best, A.; et al. Use of Whole-Exome Sequencing to Determine the Genetic Basis of Multiple Mitochondrial Respiratory Chain Complex Deficiencies. *JAMA* **2014**, *312*, 68, doi:10.1001/jama.2014.7184.
197. Smeitink, J.; van den Heuvel, L. Human Mitochondrial Complex I in Health and Disease. *The American Journal of Human Genetics* **1999**, *64*, 1505–1510, doi:10.1086/302432.
198. Loeffen, J.; Smeitink, J.; Triepels, R.; Smeets, R.; Schuelke, M.; Sengers, R.; Trijbels, F.; Hamel, B.; Mullaart, R.; van den Heuvel, L. The First Nuclear-Encoded Complex I Mutation in a Patient with Leigh Syndrome. *The American Journal of Human Genetics* **1998**, *63*, 1598–1608, doi:10.1086/302154.
199. Berger, I.; Hershkovitz, E.; Shaag, A.; Edvardson, S.; Saada, A.; Elpeleg, O. Mitochondrial Complex I Deficiency Caused by a Deleterious NDUF51 Mutation. *Ann Neurol* **2008**, *63*, 405–408, doi:10.1002/ana.21332.
200. Fernandez-Moreira, D.; Ugalde, C.; Smeets, R.; Rodenburg, R.J.T.; Lopez-Laso, E.; Ruiz-Falco, M.L.; Briones, P.; Martin, M.A.; Smeitink, J.A.M.; Arenas, J. X-Linked NDUF51 Gene Mutations Associated with Mitochondrial Encephalomyopathy. *Ann Neurol* **2007**, *61*, 73–83, doi:10.1002/ana.21036.

201. Hoefs, S.J.G.; Dieteren, C.E.J.; Distelmaier, F.; Janssen, R.J.R.J.; Epplen, A.; Swarts, H.G.P.; Forkink, M.; Rodenburg, R.J.; Nijtmans, L.G.; Willems, P.H.; et al. NDUFA2 Complex I Mutation Leads to Leigh Disease. *The American Journal of Human Genetics* **2008**, *82*, 1306–1315, doi:10.1016/j.ajhg.2008.05.007.
202. Alston, C.L.; Heidler, J.; Dibley, M.G.; Kremer, L.S.; Taylor, L.S.; Fratter, C.; French, C.E.; Glasgow, R.I.C.; Feichtinger, R.G.; Delon, I.; et al. Bi-Allelic Mutations in NDUFA6 Establish Its Role in Early-Onset Isolated Mitochondrial Complex I Deficiency. *The American Journal of Human Genetics* **2018**, *103*, 592–601, doi:10.1016/j.ajhg.2018.08.013.
203. van den Bosch, B.J.C.; Gerards, M.; Sluiter, W.; Stegmann, A.P.A.; Jongen, E.L.C.; Hellebrekers, D.M.E.I.; Oegema, R.; Lambrichs, E.H.; Prokisch, H.; Danhauser, K.; et al. Defective NDUFA9 as a Novel Cause of Neonatally Fatal Complex I Disease. *J Med Genet* **2012**, *49*, 10–15, doi:10.1136/jmedgenet-2011-100466.
204. Ostergaard, E.; Rodenburg, R.J.; van den Brand, M.; Thomsen, L.L.; Duno, M.; Batbayli, M.; Wibrand, F.; Nijtmans, L. Respiratory Chain Complex I Deficiency Due to NDUFA12 Mutations as a New Cause of Leigh Syndrome. *Journal of Medical Genetics* **2011**, *48*, 737–740, doi:10.1136/jmg.2011.088856.
205. Angebault, C.; Charif, M.; Guegen, N.; Piro-Megy, C.; Mousson de Camaret, B.; Procaccio, V.; Guichet, P.-O.; Hebrard, M.; Manes, G.; Leboucq, N.; et al. Mutation in NDUFA13/GRIM19 Leads to Early Onset Hypotonia, Dyskinesia and Sensorial Deficiencies, and Mitochondrial Complex I Instability. *Human Molecular Genetics* **2015**, *24*, 3948–3955, doi:10.1093/hmg/ddv133.
206. Haack, T.B.; Madignier, F.; Herzer, M.; Lamantea, E.; Danhauser, K.; Invernizzi, F.; Koch, J.; Freitag, M.; Drost, R.; Hillier, I.; et al. Mutation Screening of 75 Candidate Genes in 152 Complex I Deficiency Cases Identifies Pathogenic Variants in 16 Genes Including NDUFB9. *J Med Genet* **2012**, *49*, 83–89, doi:10.1136/jmedgenet-2011-100577.
207. Piekutowska-Abramczuk, D.; Assouline, Z.; Mataković, L.; Feichtinger, R.G.; Koňáriková, E.; Jurkiewicz, E.; Stawiński, P.; Gusic, M.; Koller, A.; Pollak, A.; et al. NDUFB8 Mutations Cause Mitochondrial Complex I Deficiency in Individuals with Leigh-like Encephalomyopathy. *The American Journal of Human Genetics* **2018**, *102*, 460–467, doi:10.1016/j.ajhg.2018.01.008.
208. Friederich, M.W.; Erdogan, A.J.; Coughlin, C.R.; Elos, M.T.; Jiang, H.; O'Rourke, C.P.; Lovell, M.A.; Wartchow, E.; Gowan, K.; Chatfield, K.C.; et al. Mutations in the Accessory Subunit NDUFB10 Result in Isolated Complex I Deficiency and Illustrate the Critical Role of Intermembrane Space Import for Complex I Holoenzyme Assembly. *Hum. Mol. Genet.* **2016**, dddw431, doi:10.1093/hmg/ddw431.
209. van Rahden, V.A.; Fernandez-Vizarra, E.; Alawi, M.; Brand, K.; Fellmann, F.; Horn, D.; Zeviani, M.; Kutsche, K. Mutations in NDUFB11, Encoding a Complex I Component of the Mitochondrial Respiratory Chain, Cause Microphthalmia with Linear Skin Defects Syndrome. *The American Journal of Human Genetics* **2015**, *96*, 640–650, doi:10.1016/j.ajhg.2015.02.002.
210. Reinson, K.; Kovacs-Nagy, R.; Öglane-Shlik, E.; Pajusalu, S.; Nöukas, M.; Wintjes, L.T.; van den Brandt, F.C.A.; Brink, M.; Acker, T.; Ahting, U.; et al. Diverse Phenotype in Patients with Complex I Deficiency Due to Mutations in NDUFB11. *European Journal of Medical Genetics* **2019**, *62*, 103572, doi:10.1016/j.ejmg.2018.11.006.
211. Kohda, M.; Tokuzawa, Y.; Kishita, Y.; Nyuzuki, H.; Moriyama, Y.; Mizuno, Y.; Hirata, T.; Yatsuka, Y.; Yamashita-Sugahara, Y.; Nakachi, Y.; et al. A Comprehensive Genomic Analysis Reveals the Genetic Landscape of Mitochondrial Respiratory Chain Complex Deficiencies. *PLoS Genet* **2016**, *12*, e1005679, doi:10.1371/journal.pgen.1005679.
212. Alahmad, A.; Nasca, A.; Heidler, J.; Thompson, K.; Oláhová, M.; Legati, A.; Lamantea, E.; Meisterknecht, J.; Spagnolo, M.; He, L.; et al. Bi-allelic Pathogenic Variants in NDUFC2 Cause Early-onset Leigh Syndrome and Stalled Biogenesis of Complex I. *EMBO Mol Med* **2020**, *12*, doi:10.15252/emmm.202012619.
213. Baertling, F.; Sánchez-Caballero, L.; Timal, S.; van den Brand, M.A.; Ngu, L.H.; Distelmaier, F.; Rodenburg, R.J.; Nijtmans, L.G. Mutations in Mitochondrial Complex I Assembly Factor NDUFAF3 Cause Leigh Syndrome. *Molecular Genetics and Metabolism* **2017**, *120*, 243–246, doi:10.1016/j.ymgme.2016.12.005.
214. Baertling, F.; Sánchez-Caballero, L.; van den Brand, M.A.M.; Wintjes, L.T.; Brink, M.; van den Brandt, F.A.; Wilson, C.; Rodenburg, R.J.T.; Nijtmans, L.G.J. NDUFAF4 Variants Are Associated with Leigh Syndrome and Cause a Specific Mitochondrial Complex I Assembly Defect. *Eur J Hum Genet* **2017**, *25*, 1273–1277, doi:10.1038/ejhg.2017.133.
215. Ishiyama, A.; Muramatsu, K.; Uchino, S.; Sakai, C.; Matsushima, Y.; Makioka, N.; Ogata, T.; Suzuki, E.; Komaki, H.; Sasaki, M.; et al. NDUFAF3 Variants That Disrupt Mitochondrial Complex I Assembly May Associate with Cavitating Leukoencephalopathy. *Clin Genet* **2018**, *93*, 1103–1106, doi:10.1111/cge.13215.
216. Ugarteburu, O.; Teresa Garcia-Silva, M.; Aldamiz-Echevarria, L.; Gort, L.; Garcia-Villoria, J.; Tort, F.; Ribes, A. Complex I Deficiency, Due to NDUFAF4 Mutations, Causes Severe Mitochondrial Dysfunction and Is Associated to Early Death and Dysmorphia. *Mitochondrion* **2020**, *55*, 78–84, doi:10.1016/j.mito.2020.09.003.
217. Petruzzella, V. A Nonsense Mutation in the NDUF54 Gene Encoding the 18 KDa (AQDQ) Subunit of Complex I Abolishes Assembly and Activity of the Complex in a Patient with Leigh-like Syndrome. *Human Molecular Genetics* **2001**, *10*, 529–535, doi:10.1093/hmg/10.5.529.
218. Petruzzella, V.; Papa, S. Mutations in Human Nuclear Genes Encoding for Subunits of Mitochondrial Respiratory Complex I: The NDUF54 Gene. *Gene* **2002**, *286*, 149–154, doi:10.1016/S0378-1119(01)00810-1.
219. Petruzzella, V.; Panelli, D.; Torraco, A.; Stella, A.; Papa, S. Mutations in the NDUF54 Gene of Mitochondrial Complex I Alter Stability of the Splice Variants. *FEBS Letters* **2005**, *579*, 3770–3776, doi:10.1016/j.febslet.2005.05.035.
220. Scacco, S.; Petruzzella, V.; Budde, S.; Vergari, R.; Tamborra, R.; Panelli, D.; van den Heuvel, L.P.; Smeitink, J.A.; Papa, S. Pathological Mutations of the Human NDUF54 Gene of the 18-KDa (AQDQ) Subunit of Complex I Affect the Expression of the Protein and the Assembly and Function of the Complex. *Journal of Biological Chemistry* **2003**, *278*, 44161–44167, doi:10.1074/jbc.M307615200.

221. Lamont, R.E.; Beaulieu, C.L.; Bernier, F.P.; Sparkes, R.; Innes, A.M.; Jackel-Cram, C.; Ober, C.; Parboosingh, J.S.; Lemire, E.G. A Novel NDUF54 Frameshift Mutation Causes Leigh Disease in the Hutterite Population. *Am. J. Med. Genet.* **2017**, *173*, 596–600, doi:10.1002/ajmg.a.37983.
222. Budde, S.M.S.; van den Heuvel, L.P.W.J.; Smeets, R.J.P.; Skladal, D.; Mayr, J.A.; Boelen, C.; Petruzzella, V.; Papa, S.; Smeitink, J.A.M. Clinical Heterogeneity in Patients with Mutations in the NDUF54 Gene of Mitochondrial Complex I. *J Inherit Metab Dis* **2003**, *26*, 813–815, doi:10.1023/B:BOLI.0000010003.14113.af.
223. Leshinsky-Silver, E.; Lebre, A.-S.; Minai, L.; Saada, A.; Steffann, J.; Cohen, S.; Rötig, A.; Munnich, A.; Lev, D.; Lerman-Sagie, T. NDUF54 Mutations Cause Leigh Syndrome with Predominant Brainstem Involvement. *Molecular Genetics and Metabolism* **2009**, *97*, 185–189, doi:10.1016/j.ymgme.2009.03.002.
224. Ortigoza-Escobar, J.D.; Oyarzabal, A.; Montero, R.; Artuch, R.; Jou, C.; Jiménez, C.; Gort, L.; Briones, P.; Muchart, J.; López-Gallardo, E.; et al. Ndufs4 Related Leigh Syndrome: A Case Report and Review of the Literature. *Mitochondrion* **2016**, *28*, 73–78, doi:10.1016/j.mito.2016.04.001.
225. Finsterer, J.; Zarrouk-Mahjoub, S. NDUF54-Related Leigh Syndrome in Hutterites. *Am. J. Med. Genet.* **2017**, *173*, 1450–1451, doi:10.1002/ajmg.a.38225.
226. Rahman, S.; Blok, R.B.; Dahl, H.-H.M.; Danks, D.M.; Kirby, D.M.; Chow, C.W.; Christodoulou, J.; Thorburn, D.R. Leigh Syndrome: Clinical Features and Biochemical and DNA Abnormalities. *Ann Neurol.* **1996**, *39*, 343–351, doi:10.1002/ana.410390311.
227. Leigh, D. SUBACUTE NECROTIZING ENCEPHALOMYELOPATHY IN AN INFANT. *Journal of Neurology, Neurosurgery & Psychiatry* **1951**, *14*, 216–221, doi:10.1136/jnnp.14.3.216.
228. Assouline, Z.; Jambou, M.; Rio, M.; Bole-Feysot, C.; de Lonlay, P.; Barnerias, C.; Desguerre, I.; Bonnemains, C.; Guillermet, C.; Steffann, J.; et al. A Constant and Similar Assembly Defect of Mitochondrial Respiratory Chain Complex I Allows Rapid Identification of NDUF54 Mutations in Patients with Leigh Syndrome. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **2012**, *1822*, 1062–1069, doi:10.1016/j.bbdis.2012.01.013.
229. Lazarou, M.; McKenzie, M.; Ohtake, A.; Thorburn, D.R.; Ryan, M.T. Analysis of the Assembly Profiles for Mitochondrial- and Nuclear-DNA-Encoded Subunits into Complex I. *MCB* **2007**, *27*, 4228–4237, doi:10.1128/MCB.00074-07.
230. Breuer, M.E.; Willems, P.H.G.M.; Smeitink, J.A.M.; Koopman, W.J.H.; Nooteboom, M. Cellular and Animal Models for Mitochondrial Complex I Deficiency: A Focus on the NDUF54 Subunit. *IUBMB Life* **2013**, *65*, 202–208, doi:10.1002/iub.1127.
231. Ingraham, C.A.; Burwell, L.S.; Skalska, J.; Brookes, P.S.; Howell, R.L.; Sheu, S.-S.; Pinkert, C.A. NDUF54: Creation of a Mouse Model Mimicking a Complex I Disorder. *Mitochondrion* **2009**, *9*, 204–210, doi:10.1016/j.mito.2009.02.001.
232. Ma, H.; Folmes, C.D.L.; Wu, J.; Morey, R.; Mora-Castilla, S.; Ocampo, A.; Ma, L.; Poulton, J.; Wang, X.; Ahmed, R.; et al. Metabolic Rescue in Pluripotent Cells from Patients with MtDNA Disease. *Nature* **2015**, *524*, 234–238, doi:10.1038/nature14546.
233. Galera-Monge, T.; Zurita-Díaz, F.; Canals, I.; Grønning Hansen, M.; Rufián-Vázquez, L.; Ehinger, J.K.; Elmer, E.; Martin, M.A.; Garesse, R.; Ahlenius, H.; et al. Mitochondrial Dysfunction and Calcium Dysregulation in Leigh Syndrome Induced Pluripotent Stem Cell Derived Neurons. *IJMS* **2020**, *21*, 3191, doi:10.3390/ijms21093191.
234. Zheng, X.; Boyer, L.; Jin, M.; Kim, Y.; Fan, W.; Bardy, C.; Berggren, T.; Evans, R.M.; Gage, F.H.; Hunter, T. Alleviation of Neuronal Energy Deficiency by MTOR Inhibition as a Treatment for Mitochondria-Related Neurodegeneration. *eLife* **2016**, *5*, e13378, doi:10.7554/eLife.13378.
235. Lorenz, C.; Lesimple, P.; Bukowiecki, R.; Zink, A.; Inak, G.; Mlody, B.; Singh, M.; Semtner, M.; Mah, N.; Auré, K.; et al. Human iPSC-Derived Neural Progenitors Are an Effective Drug Discovery Model for Neurological MtDNA Disorders. *Cell Stem Cell* **2017**, *20*, 659–674.e9, doi:10.1016/j.stem.2016.12.013.
236. Romero-Morales, A.I.; Rastogi, A.; Robertson, G.L.; Rasmussen, M.L.; Temuri, H.; Chakrabarty, R.P.; McElroy, G.S.; Hsu, L.; Almonacid, P.M.; Millis, B.A.; et al. Human iPSC-Derived Cerebral Organoids Model Features of Leigh Syndrome and Reveal Abnormal Corticogenesis. *Cell Biology*, 2020;
237. Zhu, Z.; Yao, J.; Johns, T.; Fu, K.; Bie, I.D.; Macmillan, C.; Cuthbert, A.P.; Newbold, R.F.; Wang, J.; Chevette, M.; et al. SURF1, Encoding a Factor Involved in the Biogenesis of Cytochrome c Oxidase, Is Mutated in Leigh Syndrome. *Nat Genet* **1998**, *20*, 337–343, doi:10.1038/3804.
238. Tiranti, V.; Hoertnagel, K.; Carrozzo, R.; Galimberti, C.; Munaro, M.; Granatiero, M.; Zelante, L.; Gasparini, P.; Marzella, R.; Rocchi, M.; et al. Mutations of SURF-1 in Leigh Disease Associated with Cytochrome c Oxidase Deficiency. *The American Journal of Human Genetics* **1998**, *63*, 1609–1621, doi:10.1086/302150.
239. Saneto, R.; Ruhoy, I. The Genetics of Leigh Syndrome and Its Implications for Clinical Practice and Risk Management. *TACG* **2014**, *221*, doi:10.2147/TACG.S46176.
240. Sun, F.; Huo, X.; Zhai, Y.; Wang, A.; Xu, J.; Su, D.; Bartlam, M.; Rao, Z. Crystal Structure of Mitochondrial Respiratory Membrane Protein Complex II. *Cell* **2005**, *121*, 1043–1057, doi:10.1016/j.cell.2005.05.025.
241. Van Vranken, J.G.; Na, U.; Winge, D.R.; Rutter, J. Protein-Mediated Assembly of Succinate Dehydrogenase and Its Cofactors. *Critical Reviews in Biochemistry and Molecular Biology* **2015**, *50*, 168–180, doi:10.3109/10409238.2014.990556.
242. Ghezzi, D.; Goffrini, P.; Uziel, G.; Horvath, R.; Klopstock, T.; Lochmüller, H.; D'Adamo, P.; Gasparini, P.; Strom, T.M.; Prokisch, H.; et al. SDHAF1, Encoding a LYR Complex-II Specific Assembly Factor, Is Mutated in SDH-Defective Infantile Leukoencephalopathy. *Nat Genet* **2009**, *41*, 654–656, doi:10.1038/ng.378.
243. Munnich, A.; Rustin, P. Clinical Spectrum and Diagnosis of Mitochondrial Disorders. *Am. J. Med. Genet.* **2001**, *106*, 4–17, doi:10.1002/ajmg.1391.

244. Bourgeron, T.; Rustin, P.; Chretien, D.; Birch-Machin, M.; Bourgeois, M.; Viegas-Péquignot, E.; Munnich, A.; Rötig, A. Mutation of a Nuclear Succinate Dehydrogenase Gene Results in Mitochondrial Respiratory Chain Deficiency. *Nat Genet* **1995**, *11*, 144–149, doi:10.1038/ng1095-144.
245. Parfait, B.; Chretien, D.; Rötig, A.; Marsac, C.; Munnich, A.; Rustin, P. Compound Heterozygous Mutations in the Flavoprotein Gene of the Respiratory Chain Complex II in a Patient with Leigh Syndrome. *Human Genetics* **2000**, *106*, 236–243, doi:10.1007/s004390051033.
246. Van Coster, R.; Seneca, S.; Smet, J.; Van Hecke, R.; Gerlo, E.; Devreese, B.; Van Beeumen, J.; Leroy, J.G.; De Meirleir, L.; Lissens, W. Homozygous Gly555Glu Mutation in the Nuclear-Encoded 70 kDa Flavoprotein Gene Causes Instability of the Respiratory Chain Complex II. *Am. J. Med. Genet.* **2003**, *120A*, 13–18, doi:10.1002/ajmg.a.10202.
247. Pagnamenta, A.T.; Hargreaves, I.P.; Duncan, A.J.; Taanman, J.-W.; Heales, S.J.; Land, J.M.; Bitner-Glindzicz, M.; Leonard, J.V.; Rahman, S. Phenotypic Variability of Mitochondrial Disease Caused by a Nuclear Mutation in Complex II. *Molecular Genetics and Metabolism* **2006**, *89*, 214–221, doi:10.1016/j.ymgme.2006.05.003.
248. Horvath, R. Leigh Syndrome Caused by Mutations in the Flavoprotein (Fp) Subunit of Succinate Dehydrogenase (SDHA). *Journal of Neurology, Neurosurgery & Psychiatry* **2006**, *77*, 74–76, doi:10.1136/jnnp.2005.067041.
249. Jain-Ghai, S.; Cameron, J.M.; Al Maawali, A.; Blaser, S.; MacKay, N.; Robinson, B.; Raiman, J. Complex II Deficiency-A Case Report and Review of the Literature. *Am. J. Med. Genet.* **2013**, *161*, 285–294, doi:10.1002/ajmg.a.35714.
250. Nesti, C.; Meschini, M.C.; Meunier, B.; Sacchini, M.; Doccini, S.; Romano, A.; Petrillo, S.; Pezzini, I.; Seddiki, N.; Rubegni, A.; et al. Additive Effect of Nuclear and Mitochondrial Mutations in a Patient with Mitochondrial Encephalomyopathy. *Human Molecular Genetics* **2015**, *24*, 3248–3256, doi:10.1093/hmg/ddv078.
251. Levitas, A.; Muhammad, E.; Harel, G.; Saada, A.; Caspi, V.C.; Manor, E.; Beck, J.C.; Sheffield, V.; Parvari, R. Familial Neonatal Isolated Cardiomyopathy Caused by a Mutation in the Flavoprotein Subunit of Succinate Dehydrogenase. *Eur J Hum Genet* **2010**, *18*, 1160–1165, doi:10.1038/ejhg.2010.83.
252. Burnichon, N.; Brière, J.-J.; Libé, R.; Vescovo, L.; Rivière, J.; Tissier, F.; Jouanno, E.; Jeunemaitre, X.; Bénit, P.; Tzagoloff, A.; et al. SDHA Is a Tumor Suppressor Gene Causing Paranglioma. *Human Molecular Genetics* **2010**, *19*, 3011–3020, doi:10.1093/hmg/ddq206.
253. Astuti, D.; Latif, F.; Dallol, A.; Dahia, P.L.M.; Douglas, F.; George, E.; Sköldbberg, F.; Husebye, E.S.; Eng, C.; Maher, E.R. Gene Mutations in the Succinate Dehydrogenase Subunit SDHB Cause Susceptibility to Familial Pheochromocytoma and to Familial Paranglioma. *The American Journal of Human Genetics* **2001**, *69*, 49–54, doi:10.1086/321282.
254. Janeway, K.A.; Kim, S.Y.; Lodish, M.; Nosé, V.; Rustin, P.; Gaal, J.; Dahia, P.L.M.; Liegl, B.; Ball, E.R.; Raygada, M.; et al. Defects in Succinate Dehydrogenase in Gastrointestinal Stromal Tumors Lacking KIT and PDGFRA Mutations. *Proc Natl Acad Sci USA* **2011**, *108*, 314–318, doi:10.1073/pnas.1009199108.
255. Baysal, B.E. An Alu-Mediated Partial SDHC Deletion Causes Familial and Sporadic Paranglioma. *Journal of Medical Genetics* **2004**, *41*, 703–709, doi:10.1136/jmg.2004.019224.
256. McWhinney, S.R.; Pasini, B.; Stratakis, C.A. Familial Gastrointestinal Stromal Tumors and Germ-Line Mutations. *N Engl J Med* **2007**, *357*, 1054–1056, doi:10.1056/NEJMc071191.
257. Baysal, B.E. Mutations in SDHD, a Mitochondrial Complex II Gene, in Hereditary Paranglioma. *Science* **2000**, *287*, 848–851, doi:10.1126/science.287.5454.848.
258. Hao, H.-X.; Khalimonchuk, O.; Schraders, M.; Dephoure, N.; Bayley, J.-P.; Kunst, H.; Devilee, P.; Cremers, C.W.R.J.; Schiffman, J.D.; Bentz, B.G.; et al. SDH5, a Gene Required for Flavination of Succinate Dehydrogenase, Is Mutated in Paranglioma. *Science* **2009**, *325*, 1139–1142, doi:10.1126/science.1175689.
259. Sköldbberg; Grimelius; Woodward; Rorsman; Van Schothorst; Winqvist; Karlsson; Åkerström; Kämpe; Husebye A Family with Hereditary Extra-Adrenal Parangliomas without Evidence for Mutations in the von Hippel-Lindau Disease or Ret Genes: Hereditary Extra-Adrenal Paranglioma. *Clinical Endocrinology* **1998**, *48*, 11–16, doi:10.1046/j.1365-2265.1998.00320.x.
260. Lussey-Lepoutre, C.; Buffet, A.; Gimenez-Roqueplo, A.-P.; Favier, J. Mitochondrial Deficiencies in the Predisposition to Paranglioma. *Metabolites* **2017**, *7*, 17, doi:10.3390/metabo7020017.
261. Ghezzi, D.; Zeviani, M. Human Diseases Associated with Defects in Assembly of OXPHOS Complexes. *Essays in Biochemistry* **2018**, *62*, 271–286, doi:10.1042/EBC20170099.
262. Dwight, T.; Na, U.; Kim, E.; Zhu, Y.; Richardson, A.L.; Robinson, B.G.; Tucker, K.M.; Gill, A.J.; Benn, D.E.; Clifton-Bligh, R.J.; et al. Analysis of SDHAF3 in Familial and Sporadic Pheochromocytoma and Paranglioma. *BMC Cancer* **2017**, *17*, 497, doi:10.1186/s12885-017-3486-z.
263. Kudryavtseva, A.V.; Kalinin, D.V.; Pavlov, V.S.; Savvateeva, M.V.; Fedorova, M.S.; Pudova, E.A.; Kobelyatskaya, A.A.; Golovyuk, A.L.; Guvatova, Z.G.; Razmakhaev, G.S.; et al. Mutation Profiling in Eight Cases of Vagal Parangliomas. *BMC Med Genomics* **2020**, *13*, 115, doi:10.1186/s12920-020-00763-4.
264. Protasoni, M.; Zeviani, M. Mitochondrial Structure and Bioenergetics in Normal and Disease Conditions. *IJMS* **2021**, *22*, 586, doi:10.3390/ijms22020586.
265. Iwata, S. Complete Structure of the 11-Subunit Bovine Mitochondrial Cytochrome bc₁ Complex. *Science* **1998**, *281*, 64–71, doi:10.1126/science.281.5373.64.
266. Hildenbeutel, M.; Hegg, E.L.; Stephan, K.; Gruschke, S.; Meunier, B.; Ott, M. Assembly Factors Monitor Sequential Hemylation of Cytochrome b to Regulate Mitochondrial Translation. *Journal of Cell Biology* **2014**, *205*, 511–524, doi:10.1083/jcb.201401009.

267. Tucker, E.J.; Wanschers, B.F.J.; Szklarczyk, R.; Mountford, H.S.; Wijeyeratne, X.W.; van den Brand, M.A.M.; Leenders, A.M.; Rodenburg, R.J.; Reljić, B.; Compton, A.G.; et al. Mutations in the UQCC1-Interacting Protein, UQCC2, Cause Human Complex III Deficiency Associated with Perturbed Cytochrome b Protein Expression. *PLoS Genet* **2013**, *9*, e1004034, doi:10.1371/journal.pgen.1004034.
268. Wanschers, B.F.J.; Szklarczyk, R.; van den Brand, M.A.M.; Jonckheere, A.; Suijskens, J.; Smeets, R.; Rodenburg, R.J.; Stephan, K.; Helland, I.B.; Elkamil, A.; et al. A Mutation in the Human CBP4 Ortholog UQCC3 Impairs Complex III Assembly, Activity and Cytochrome b Stability. *Human Molecular Genetics* **2014**, *23*, 6356–6365, doi:10.1093/hmg/ddu357.
269. Stephan, K.; Ott, M. Timing of Dimerization of the Bc Complex during Mitochondrial Respiratory Chain Assembly. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2020**, *1861*, 148177, doi:10.1016/j.bbabi.2020.148177.
270. Bottani, E.; Cerutti, R.; Harbour, M.E.; Ravaglia, S.; Dogan, S.A.; Giordano, C.; Fearnley, I.M.; D'Amati, G.; Viscomi, C.; Fernandez-Vizarra, E.; et al. TTC19 Plays a Husbandry Role on UQCRFS1 Turnover in the Biogenesis of Mitochondrial Respiratory Complex III. *Molecular Cell* **2017**, *67*, 96–105.e4, doi:10.1016/j.molcel.2017.06.001.
271. Atkinson, A.; Smith, P.; Fox, J.L.; Cui, T.-Z.; Khalimonchuk, O.; Winge, D.R. The LYR Protein Mzm1 Functions in the Insertion of the Rieske Fe/S Protein in Yeast Mitochondria. *Molecular and Cellular Biology* **2011**, *31*, 3988–3996, doi:10.1128/MCB.05673-11.
272. Cui, T.-Z.; Smith, P.M.; Fox, J.L.; Khalimonchuk, O.; Winge, D.R. Late-Stage Maturation of the Rieske Fe/S Protein: Mzm1 Stabilizes Rip1 but Does Not Facilitate Its Translocation by the AAA ATPase Bcs1. *Molecular and Cellular Biology* **2012**, *32*, 4400–4409, doi:10.1128/MCB.00441-12.
273. Sánchez, E.; Lobo, T.; Fox, J.L.; Zeviani, M.; Winge, D.R.; Fernández-Vizarra, E. LYRM7/MZM1L Is a UQCRFS1 Chaperone Involved in the Last Steps of Mitochondrial Complex III Assembly in Human Cells. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2013**, *1827*, 285–293, doi:10.1016/j.bbabi.2012.11.003.
274. Cruciat, C.-M. Bcs1p, an AAA-Family Member, Is a Chaperone for the Assembly of the Cytochrome Bc1 Complex. *The EMBO Journal* **1999**, *18*, 5226–5233, doi:10.1093/emboj/18.19.5226.
275. Fernandez-Vizarra, E.; Bugiani, M.; Goffrini, P.; Carrara, F.; Farina, L.; Procopio, E.; Donati, A.; Uziel, G.; Ferrero, I.; Zeviani, M. Impaired Complex III Assembly Associated with BCS1L Gene Mutations in Isolated Mitochondrial Encephalopathy. *Human Molecular Genetics* **2007**, *16*, 1241–1252, doi:10.1093/hmg/ddm072.
276. Tang, W.K.; Borgnia, M.J.; Hsu, A.L.; Esser, L.; Fox, T.; de Val, N.; Xia, D. Structures of AAA Protein Translocase Bcs1 Suggest Translocation Mechanism of a Folded Protein. *Nat Struct Mol Biol* **2020**, *27*, 202–209, doi:10.1038/s41594-020-0373-0.
277. Wagener, N.; Ackermann, M.; Funes, S.; Neupert, W. A Pathway of Protein Translocation in Mitochondria Mediated by the AAA-ATPase Bcs1. *Molecular Cell* **2011**, *44*, 191–202, doi:10.1016/j.molcel.2011.07.036.
278. Peruzzo, R.; Corrà, S.; Costa, R.; Brischigliaro, M.; Varanita, T.; Biasutto, L.; Rampazzo, C.; Ghezzi, D.; Leanza, L.; Zoratti, M.; et al. Exploiting Pyocyanin to Treat Mitochondrial Disease Due to Respiratory Complex III Dysfunction. *Nat Commun* **2021**, *12*, 2103, doi:10.1038/s41467-021-22062-x.
279. Andreu, A.L.; Bruno, C.; Dunne, T.C.; Tanji, K.; Shanske, S.; Sue, C.M.; Krishna, S.; Hadjigeorgiou, G.M.; Shtilbans, A.; Bonilla, E.; et al. A Nonsense Mutation (G15059A) in the Cytochrome b Gene in a Patient with Exercise Intolerance and Myoglobinuria. *Ann Neurol* **1999**, *45*, 127–130, doi:10.1002/1531-8249(199901)45:1<127::aid-art20>3.0.co;2-y.
280. Andreu, A.L.; Hanna, M.G.; Reichmann, H.; Bruno, C.; Penn, A.S.; Tanji, K.; Pallotti, F.; Iwata, S.; Bonilla, E.; Lach, B.; et al. Exercise Intolerance Due to Mutations in the Cytochrome b Gene of Mitochondrial DNA. *N Engl J Med* **1999**, *341*, 1037–1044, doi:10.1056/NEJM199909303411404.
281. De Co, I.F.; Renier, W.O.; Ruitenbeek, W.; Ter Laak, H.J.; Bakker, M.; Schägger, H.; Van Oost, B.A.; Smeets, H.J. A 4-Base Pair Deletion in the Mitochondrial Cytochrome b Gene Associated with Parkinsonism/MELAS Overlap Syndrome. *Ann Neurol* **1999**, *45*, 130–133, doi:10.1002/1531-8249(199901)45:1<130::aid-art21>3.3.co;2-q.
282. Keightley, J.A.; Anitori, R.; Burton, M.D.; Quan, F.; Buist, N.R.M.; Kennaway, N.G. Mitochondrial Encephalomyopathy and Complex III Deficiency Associated with a Stop-Codon Mutation in the Cytochrome b Gene. *The American Journal of Human Genetics* **2000**, *67*, 1400–1410, doi:10.1086/316900.
283. Andreu, A.L.; Bruno, C.; Shanske, S.; Shtilbans, A.; Hirano, M.; Krishna, S.; Hayward, L.; Systrom, D.S.; Brown, R.H.; DiMauro, S. Missense Mutation in the MtDNA Cytochrome b Gene in a Patient with Myopathy. *Neurology* **1998**, *51*, 1444–1447, doi:10.1212/WNL.51.5.1444.
284. Lamantea, E.; Carrara, F.; Mariotti, C.; Morandi, L.; Tiranti, V.; Zeviani, M. A Novel Nonsense Mutation (Q352X) in the Mitochondrial Cytochrome b Gene Associated with a Combined Deficiency of Complexes I and III. *Neuromuscular Disorders* **2002**, *12*, 49–52, doi:10.1016/S0960-8966(01)00244-9.
285. Mancuso, M.; Filosto, M.; Stevens, J.C.; Patterson, M.; Shanske, S.; Krishna, S.; DiMauro, S. Mitochondrial Myopathy and Complex III Deficiency in a Patient with a New Stop-Codon Mutation (G339X) in the Cytochrome b Gene. *Journal of the Neurological Sciences* **2003**, *209*, 61–63, doi:10.1016/S0022-510X(02)00462-8.
286. Andreu, A.L.; Checcarelli, N.; Iwata, S.; Shanske, S.; DiMauro, S. A Missense Mutation in the Mitochondrial Cytochrome b Gene in a Revisited Case with Histiocytoid Cardiomyopathy. *Pediatr Res* **2000**, *48*, 311–314, doi:10.1203/00006450-200009000-00008.
287. Wibrand, F.; Ravn, K.; Schwartz, M.; Rosenberg, T.; Horn, N.; Vissing, J. Multisystem Disorder Associated with a Missense Mutation in the Mitochondrial Cytochrome b Gene. *Ann Neurol* **2001**, *50*, 540–543, doi:10.1002/ana.1224.
288. Schuelke, M.; Krude, H.; Finckh, B.; Mayatepek, E.; Janssen, A.; Schmelz, M.; Trefz, F.; Trijbels, F.; Smeitink, J. Septo-Optic Dysplasia Associated with a New Mitochondrial cytochrome b Mutation. *Ann Neurol* **2002**, *51*, 388–392, doi:10.1002/ana.10151.

289. Ghelli, A.; Tropeano, C.V.; Calvaruso, M.A.; Marchesini, A.; Iommarini, L.; Porcelli, A.M.; Zanna, C.; De Nardo, V.; Martinuzzi, A.; Wibrand, F.; et al. The Cytochrome b p.278Y>C Mutation Causative of a Multisystem Disorder Enhances Superoxide Production and Alters Supramolecular Interactions of Respiratory Chain Complexes. *Human Molecular Genetics* **2013**, *22*, 2141–2151, doi:10.1093/hmg/ddt067.
290. Carossa, V.; Ghelli, A.; Tropeano, C.V.; Valentino, M.L.; Iommarini, L.; Maresca, A.; Caporali, L.; La Morgia, C.; Liguori, R.; Barboni, P.; et al. A Novel In-Frame 18-Bp Microdeletion in *MT-CYB* Causes a Multisystem Disorder with Prominent Exercise Intolerance. *Human Mutation* **2014**, *35*, 954–958, doi:10.1002/humu.22596.
291. Fernández-Vizarra, E.; Zeviani, M. Nuclear Gene Mutations as the Cause of Mitochondrial Complex III Deficiency. *Front. Genet.* **2015**, *6*, doi:10.3389/fgene.2015.00134.
292. Ghezzi, D.; Arzuffi, P.; Zordan, M.; Da Re, C.; Lamperti, C.; Benna, C.; D'Adamo, P.; Diodato, D.; Costa, R.; Mariotti, C.; et al. Mutations in *TTC19* Cause Mitochondrial Complex III Deficiency and Neurological Impairment in Humans and Flies. *Nat Genet* **2011**, *43*, 259–263, doi:10.1038/ng.761.
293. Invernizzi, F.; Tigano, M.; Dallabona, C.; Donnini, C.; Ferrero, I.; Cremonte, M.; Ghezzi, D.; Lamperti, C.; Zeviani, M. A Homozygous Mutation in *LYRM7/MZM1L* Associated with Early Onset Encephalopathy, Lactic Acidosis, and Severe Reduction of Mitochondrial Complex III Activity. *Human Mutation* **2013**, *34*, 1619–1622, doi:10.1002/humu.22441.
294. de Lonlay, P.; Valnot, I.; Barrientos, A.; Gorbatyuk, M.; Tzagoloff, A.; Taanman, J.-W.; Benayoun, E.; Chrétien, D.; Kadhom, N.; Lombès, A.; et al. A Mutant Mitochondrial Respiratory Chain Assembly Protein Causes Complex III Deficiency in Patients with Tubulopathy, Encephalopathy and Liver Failure. *Nat Genet* **2001**, *29*, 57–60, doi:10.1038/ng706.
295. Lin, C.-H.; Tsai, P.-I.; Lin, H.-Y.; Hattori, N.; Funayama, M.; Jeon, B.; Sato, K.; Abe, K.; Mukai, Y.; Takahashi, Y.; et al. Mitochondrial UQCRC1 Mutations Cause Autosomal Dominant Parkinsonism with Polyneuropathy. *Brain* **2020**, *143*, 3352–3373, doi:10.1093/brain/awaa279.
296. Miyake, N.; Yano, S.; Sakai, C.; Hatakeyama, H.; Matsushima, Y.; Shiina, M.; Watanabe, Y.; Bartley, J.; Abdenur, J.E.; Wang, R.Y.; et al. Mitochondrial Complex III Deficiency Caused by a Homozygous *UQCRC2* Mutation Presenting with Neonatal-Onset Recurrent Metabolic Decompensation. *Human Mutation* **2013**, *34*, 446–452, doi:10.1002/humu.22257.
297. Brown, M.D.; Voljavec, A.S.; Lott, M.T.; Torroni, A.; Yang, C.C.; Wallace, D.C. Mitochondrial DNA Complex I and III Mutations Associated with Leber's Hereditary Optic Neuropathy. *Genetics* **1992**, *130*, 163–173, doi:10.1093/genetics/130.1.163.
298. Fouad Bouzidi, M.; Schägger, H.; Collombet, J.-M.; Carrier, H.; Flocard, F.; Quard, S.; Mousson, B.; Godinot, C. Decreased Expression of Ubiquinol-Cytochrome c Reductase Subunits in Patients Exhibiting Mitochondrial Myopathy with Progressive Exercise Intolerance. *Neuromuscular Disorders* **1993**, *3*, 599–604, doi:10.1016/0960-8966(93)90123-2.
299. Gagnard, P.; Menezes, M.; Schiff, M.; Bayot, A.; Rak, M.; Ogier de Baulny, H.; Su, C.-H.; Gilleron, M.; Lombes, A.; Abida, H.; et al. Mutations in *CYC1*, Encoding Cytochrome C1 Subunit of Respiratory Chain Complex III, Cause Insulin-Responsive Hypoglycemia. *The American Journal of Human Genetics* **2013**, *93*, 384–389, doi:10.1016/j.ajhg.2013.06.015.
300. Gusic, M.; Schottmann, G.; Feichtinger, R.G.; Du, C.; Scholz, C.; Wagner, M.; Mayr, J.A.; Lee, C.-Y.; Yépez, V.A.; Lorenz, N.; et al. Bi-Allelic *UQCRC1* Variants Are Associated with Mitochondrial Complex III Deficiency, Cardiomyopathy, and Alopecia Totalis. *The American Journal of Human Genetics* **2020**, *106*, 102–111, doi:10.1016/j.ajhg.2019.12.005.
301. Haut, S.; Brivet, M.; Touati, G.; Rustin, P.; Lebon, S.; Garcia-Cazorla, A.; Saudubray, J.M.; Boutron, A.; Legrand, A.; Slama, A. A Deletion in the Human *QP-C* Gene Causes a Complex III Deficiency Resulting in Hypoglycaemia and Lactic Acidosis. *Hum Genet* **2003**, *113*, 118–122, doi:10.1007/s00439-003-0946-0.
302. Barel, O.; Shorer, Z.; Flusser, H.; Ofir, R.; Narkis, G.; Finer, G.; Shalev, H.; Nasasra, A.; Saada, A.; Birk, O.S. Mitochondrial Complex III Deficiency Associated with a Homozygous Mutation in *UQCRC1*. *The American Journal of Human Genetics* **2008**, *82*, 1211–1216, doi:10.1016/j.ajhg.2008.03.020.
303. Feichtinger, R.G.; Brunner-Krainz, M.; Alhaddad, B.; Wortmann, S.B.; Kovacs-Nagy, R.; Stojakovic, T.; Erwa, W.; Resch, B.; Windischhofer, W.; Verheyen, S.; et al. Combined Respiratory Chain Deficiency and *UQCRC2* Mutations in Neonatal Encephalomyopathy: Defective Supercomplex Assembly in Complex III Deficiencies. *Oxidative Medicine and Cellular Longevity* **2017**, *2017*, 1–11, doi:10.1155/2017/7202589.
304. Hausman-Kedem, M.; Ben-Shachar, S.; Menascu, S.; Geva, K.; Sagie, L.; Fattal-Valevski, A. *VPS53* Gene Is Associated with a New Phenotype of Complicated Hereditary Spastic Paraparesis. *Neurogenetics* **2019**, *20*, 187–195, doi:10.1007/s10048-019-00586-1.
305. Baker, R.A.; Priestley, J.R.C.; Wilstermann, A.M.; Reese, K.J.; Mark, P.R. Clinical Spectrum of *BCS1L* Mitopathies and Their Underlying Structural Relationships. *Am J Med Genet* **2019**, *179*, 373–380, doi:10.1002/ajmg.a.61019.
306. Visapää, I.; Fellman, V.; Varilo, T.; Palotie, A.; Raivio, K.O.; Peltonen, L. Assignment of the Locus for a New Lethal Neonatal Metabolic Syndrome to 2q33-37. *The American Journal of Human Genetics* **1998**, *63*, 1396–1403, doi:10.1086/302123.
307. Siddiqi, S.; Siddiqi, S.; Mansoor, A.; Oostrik, J.; Ahmad, N.; Kazmi, S.A.R.; Kremer, H.; Qamar, R.; Schraders, M. Novel Mutation in AAA Domain of *BCS1L* Causing Björnstad Syndrome. *J Hum Genet* **2013**, *58*, 819–821, doi:10.1038/jhg.2013.101.
308. Hinson, J.T.; Fantin, V.R.; Schönberger, J.; Breivik, N.; Siem, G.; McDonough, B.; Sharma, P.; Keogh, I.; Godinho, R.; Santos, F.; et al. Missense Mutations in the *BCS1L* Gene as a Cause of the Björnstad Syndrome. *N Engl J Med* **2007**, *356*, 809–819, doi:10.1056/NEJMoa055262.
309. Gil-Borlado, M.C.; González-Hoyuela, M.; Blázquez, A.; García-Silva, M.T.; Gabaldón, T.; Manzanares, J.; Vara, J.; Martín, M.A.; Seneca, S.; Arenas, J.; et al. Pathogenic Mutations in the 5' Untranslated Region of *BCS1L* mRNA in Mitochondrial Complex III Deficiency. *Mitochondrion* **2009**, *9*, 299–305, doi:10.1016/j.mito.2009.04.001.

310. Blázquez, A.; Gil-Borlado, M.C.; Morán, M.; Verdú, A.; Cazorla-Calleja, M.R.; Martín, M.A.; Arenas, J.; Ugalde, C. Infantile Mitochondrial Encephalomyopathy with Unusual Phenotype Caused by a Novel BCS1L Mutation in an Isolated Complex III-Deficient Patient. *Neuromuscular Disorders* **2009**, *19*, 143–146, doi:10.1016/j.nmd.2008.11.016.
311. Dallabona, C.; Abbink, T.E.M.; Carrozzo, R.; Torracco, A.; Legati, A.; van Berkel, C.G.M.; Niceta, M.; Langella, T.; Verrigni, D.; Rizza, T.; et al. LYRM7 Mutations Cause a Multifocal Cavitating Leukoencephalopathy with Distinct MRI Appearance. *Brain* **2016**, *139*, 782–794, doi:10.1093/brain/awv392.
312. Hempel, M.; Kremer, L.S.; Tsiakas, K.; Alhaddad, B.; Haack, T.B.; Löbel, U.; Feichtinger, R.G.; Sperl, W.; Prokisch, H.; Mayr, J.A.; et al. LYRM7 - Associated Complex III Deficiency: A Clinical, Molecular Genetic, MR Tomographic, and Biochemical Study. *Mitochondrion* **2017**, *37*, 55–61, doi:10.1016/j.mito.2017.07.001.
313. Kremer, L.S.; L'hermitte-Stead, C.; Lesimple, P.; Gilleron, M.; Filaut, S.; Jardel, C.; Haack, T.B.; Strom, T.M.; Meitinger, T.; Azouz, H.; et al. Severe Respiratory Complex III Defect Prevents Liver Adaptation to Prolonged Fasting. *Journal of Hepatology* **2016**, *65*, 377–385, doi:10.1016/j.jhep.2016.04.017.
314. Morino, H.; Miyamoto, R.; Ohnishi, S.; Maruyama, H.; Kawakami, H. Exome Sequencing Reveals a Novel TTC19 Mutation in an Autosomal Recessive Spinocerebellar Ataxia Patient. *BMC Neurol* **2014**, *14*, 5, doi:10.1186/1471-2377-14-5.
315. Nogueira, C.; Barros, J.; Sá, M.J.; Azevedo, L.; Taipa, R.; Torracco, A.; Meschini, M.C.; Verrigni, D.; Nesti, C.; Rizza, T.; et al. Novel TTC19 Mutation in a Family with Severe Psychiatric Manifestations and Complex III Deficiency. *Neurogenetics* **2013**, *14*, 153–160, doi:10.1007/s10048-013-0361-1.
316. Habibzadeh, P.; Inaloo, S.; Silawi, M.; Dastsooz, H.; Farazi Fard, M.A.; Sadeghipour, F.; Faghihi, Z.; Rezaeian, M.; Yavarian, M.; Böhm, J.; et al. A Novel TTC19 Mutation in a Patient With Neurological, Psychological, and Gastrointestinal Impairment. *Front. Neurol.* **2019**, *10*, 944, doi:10.3389/fneur.2019.00944.
317. Mordaunt, D.A.; Jolley, A.; Balasubramaniam, S.; Thorburn, D.R.; Mountford, H.S.; Compton, A.G.; Nicholl, J.; Manton, N.; Clark, D.; Bratkovic, D.; et al. Phenotypic Variation of TTC19 -Deficient Mitochondrial Complex III Deficiency: A Case Report and Literature Review. *Am. J. Med. Genet.* **2015**, *167*, 1330–1336, doi:10.1002/ajmg.a.36968.
318. Balsa, E.; Marco, R.; Perales-Clemente, E.; Szklarczyk, R.; Calvo, E.; Landázuri, M.O.; Enríquez, J.A. NDUFA4 Is a Subunit of Complex IV of the Mammalian Electron Transport Chain. *Cell Metabolism* **2012**, *16*, 378–386, doi:10.1016/j.cmet.2012.07.015.
319. Pitceathly, R.D.S.; Rahman, S.; Wedatilake, Y.; Polke, J.M.; Cirak, S.; Foley, A.R.; Sailer, A.; Hurler, M.E.; Stalker, J.; Hargreaves, I.; et al. NDUFA4 Mutations Underlie Dysfunction of a Cytochrome c Oxidase Subunit Linked to Human Neurological Disease. *Cell Reports* **2013**, *3*, 1795–1805, doi:10.1016/j.celrep.2013.05.005.
320. Zong, S.; Wu, M.; Gu, J.; Liu, T.; Guo, R.; Yang, M. Structure of the Intact 14-Subunit Human Cytochrome c Oxidase. *Cell Res* **2018**, *28*, 1026–1034, doi:10.1038/s41422-018-0071-1.
321. Hill, B.C. The Sequence of Electron Carriers in the Reaction of Cytochrome c Oxidase with Oxygen. *J Bioenerg Biomembr* **1993**, *25*, 115–120, doi:10.1007/BF00762853.
322. Sharma, V.; Ala-Vannessluoma, P.; Vattulainen, I.; Wikström, M.; Róg, T. Role of Subunit III and Its Lipids in the Molecular Mechanism of Cytochrome c Oxidase. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2015**, *1847*, 690–697, doi:10.1016/j.bbabi.2015.04.007.
323. Sinkler, C.A.; Kalpage, H.; Shay, J.; Lee, I.; Malek, M.H.; Grossman, L.I.; Hüttemann, M. Tissue- and Condition-Specific Isoforms of Mammalian Cytochrome c Oxidase Subunits: From Function to Human Disease. *Oxidative Medicine and Cellular Longevity* **2017**, *2017*, 1–19, doi:10.1155/2017/1534056.
324. Kadenbach, B.; Hüttemann, M. The Subunit Composition and Function of Mammalian Cytochrome c Oxidase. *Mitochondrion* **2015**, *24*, 64–76, doi:10.1016/j.mito.2015.07.002.
325. Nijtmans, L.G.J.; Taanman, J.-W.; Muijsers, A.O.; Speijer, D.; Van den Bogert, C. Assembly of Cytochrome-c Oxidase in Cultured Human Cells. *Eur J Biochem* **1998**, *254*, 389–394, doi:10.1046/j.1432-1327.1998.2540389.x.
326. Vidoni, S.; Harbour, M.E.; Guerrero-Castillo, S.; Signes, A.; Ding, S.; Fearnley, I.M.; Taylor, R.W.; Tiranti, V.; Arnold, S.; Fernandez-Vizarra, E.; et al. MR-1S Interacts with PET100 and PET117 in Module-Based Assembly of Human Cytochrome c Oxidase. *Cell Reports* **2017**, *18*, 1727–1738, doi:10.1016/j.celrep.2017.01.044.
327. Rak, M.; Bénit, P.; Chrétien, D.; Bouchereau, J.; Schiff, M.; El-Khoury, R.; Tzagoloff, A.; Rustin, P. Mitochondrial Cytochrome c Oxidase Deficiency. *Clinical Science* **2016**, *130*, 393–407, doi:10.1042/CS20150707.
328. Massa, V.; Fernandez-Vizarra, E.; Alshahwan, S.; Bakhsh, E.; Goffrini, P.; Ferrero, I.; Mereghetti, P.; D'Adamo, P.; Gasparini, P.; Zeviani, M. Severe Infantile Encephalomyopathy Caused by a Mutation in COX6B1, a Nucleus-Encoded Subunit of Cytochrome C Oxidase. *The American Journal of Human Genetics* **2008**, *82*, 1281–1289, doi:10.1016/j.ajhg.2008.05.002.
329. Abdulhag, U.N.; Soiferman, D.; Schueler-Furman, O.; Miller, C.; Shaag, A.; Elpeleg, O.; Edvardson, S.; Saada, A. Mitochondrial Complex IV Deficiency, Caused by Mutated COX6B1, Is Associated with Encephalomyopathy, Hydrocephalus and Cardiomyopathy. *Eur J Hum Genet* **2015**, *23*, 159–164, doi:10.1038/ejhg.2014.85.
330. Lamperti, C.; Diodato, D.; Lamantea, E.; Carrara, F.; Ghezzi, D.; Mereghetti, P.; Rizzi, R.; Zeviani, M. MELAS-like Encephalomyopathy Caused by a New Pathogenic Mutation in the Mitochondrial DNA Encoded Cytochrome c Oxidase Subunit I. *Neuromuscular Disorders* **2012**, *22*, 990–994, doi:10.1016/j.nmd.2012.06.003.
331. Valente, L.; Piga, D.; Lamantea, E.; Carrara, F.; Uziel, G.; Cudia, P.; Zani, A.; Farina, L.; Morandi, L.; Mora, M.; et al. Identification of Novel Mutations in Five Patients with Mitochondrial Encephalomyopathy. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2009**, *1787*, 491–501, doi:10.1016/j.bbabi.2008.10.001.

332. Comi, G.P.; Bordoni, A.; Salani, S.; Franceschina, L.; Sciacco, M.; Prella, A.; Fortunato, F.; Zeviani, M.; Napoli, L.; Bresolin, N.; et al. Cytochrome c Oxidase Subunit I Microdeletion in a Patient with Motor Neuron Disease. *Ann Neurol* **1998**, *43*, 110–116, doi:10.1002/ana.410430119.
333. D'Aurelio, M.; Pallotti, F.; Barrientos, A.; Gajewski, C.D.; Kwong, J.Q.; Bruno, C.; Beal, M.F.; Manfredi, G. In Vivo Regulation of Oxidative Phosphorylation in Cells Harboring a Stop-Codon Mutation in Mitochondrial DNA-Encoded Cytochrome c Oxidase Subunit I. *Journal of Biological Chemistry* **2001**, *276*, 46925–46932, doi:10.1074/jbc.M106429200.
334. Nishigaki, Y.; Ueno, H.; Coku, J.; Koga, Y.; Fujii, T.; Sahashi, K.; Nakano, K.; Yoneda, M.; Nonaka, M.; Tang, L.; et al. Extensive Screening System Using Suspension Array Technology to Detect Mitochondrial DNA Point Mutations. *Mitochondrion* **2010**, *10*, 300–308, doi:10.1016/j.mito.2010.01.003.
335. Clark, K.M.; Taylor, R.W.; Johnson, M.A.; Chinnery, P.F.; Chrzanowska-Lightowlers, Z.M.A.; Andrews, R.M.; Nelson, I.P.; Wood, N.W.; Lamont, P.J.; Hanna, M.G.; et al. An MtDNA Mutation in the Initiation Codon of the Cytochrome C Oxidase Subunit II Gene Results in Lower Levels of the Protein and a Mitochondrial Encephalomyopathy. *The American Journal of Human Genetics* **1999**, *64*, 1330–1339, doi:10.1086/302361.
336. Abu-Amro, K.K.; Bosley, T.M. Mitochondrial Abnormalities in Patients with LHON-like Optic Neuropathies. *Invest. Ophthalmol. Vis. Sci.* **2006**, *47*, 4211, doi:10.1167/iovs.06-0295.
337. Rahman, S.; Taanman, J.-W.; Cooper, J.M.; Nelson, I.; Hargreaves, I.; Meunier, B.; Hanna, M.G.; García, J.J.; Capaldi, R.A.; Lake, B.D.; et al. A Missense Mutation of Cytochrome Oxidase Subunit II Causes Defective Assembly and Myopathy. *The American Journal of Human Genetics* **1999**, *65*, 1030–1039, doi:10.1086/302590.
338. Wei, Y.-L.; Yu, C.-A.; Yang, P.; Li, A.-L.; Wen, J.-Y.; Zhao, S.-M.; Liu, H.-X.; Ke, Y.-N.; Campbell, W.; Zhang, Y.-G.; et al. NOVEL MITOCHONDRIAL DNA MUTATIONS ASSOCIATED WITH CHINESE FAMILIAL HYPERTROPHIC CARDIOMYOPATHY. *Clinical and Experimental Pharmacology and Physiology* **2009**, *36*, 933–939, doi:10.1111/j.1440-1681.2009.05183.x.
339. Tabeji, M.; Mkaouer-Rebai, E.; Mnif, M.; Kallabi, F.; Ben Mahmoud, A.; Ben Saad, W.; Charfi, N.; Keskes-Ammar, L.; Kamoun, H.; Abid, M.; et al. A Novel Mutation MT-COIII m.9267G>C and MT-COI m.5913G>A Mutation in Mitochondrial Genes in a Tunisian Family with Maternally Inherited Diabetes and Deafness (MIDD) Associated with Sever Nephropathy. *Biochemical and Biophysical Research Communications* **2015**, *459*, 353–360, doi:10.1016/j.bbrc.2015.01.151.
340. Horvath, R. Childhood Onset Mitochondrial Myopathy and Lactic Acidosis Caused by a Stop Mutation in the Mitochondrial Cytochrome c Oxidase III Gene. *Journal of Medical Genetics* **2002**, *39*, 812–816, doi:10.1136/jmg.39.11.812.
341. Mkaouer-Rebai, E.; Ellouze, E.; Chamkha, I.; Kammoun, F.; Triki, C.; Fakhfakh, F. Molecular-Clinical Correlation in a Family With a Novel Heteroplasmic Leigh Syndrome Missense Mutation in the Mitochondrial Cytochrome c Oxidase III Gene. *J Child Neurol* **2011**, *26*, 12–20, doi:10.1177/0883073810371227.
342. Bosley, T.M.; Brodsky, M.C.; Glasier, C.M.; Abu-Amro, K.K. Sporadic Bilateral Optic Neuropathy in Children: The Role of Mitochondrial Abnormalities. *Invest. Ophthalmol. Vis. Sci.* **2008**, *49*, 5250, doi:10.1167/iovs.08-2193.
343. Marotta, R.; Chin, J.; Kirby, D.M.; Chiotis, M.; Cook, M.; Collins, S.J. Novel Single Base Pair COX III Subunit Deletion of Mitochondrial DNA Associated with Rhabdomyolysis. *Journal of Clinical Neuroscience* **2011**, *18*, 290–292, doi:10.1016/j.jocn.2010.06.001.
344. Hanna, M.G.; Nelson, I.P.; Rahman, S.; Lane, R.J.M.; Land, J.; Heales, S.; Cooper, M.J.; Schapira, A.H.V.; Morgan-Hughes, J.A.; Wood, N.W. Cytochrome c Oxidase Deficiency Associated with the First Stop-Codon Point Mutation in Human MtDNA. *The American Journal of Human Genetics* **1998**, *63*, 29–36, doi:10.1086/301910.
345. Abu-Libdeh, B.; Douiev, L.; Amro, S.; Shahrour, M.; Ta-Shma, A.; Miller, C.; Elpeleg, O.; Saada, A. Mutation in the COX4I1 Gene Is Associated with Short Stature, Poor Weight Gain and Increased Chromosomal Breaks, Simulating Fanconi Anemia. *Eur J Hum Genet* **2017**, *25*, 1142–1146, doi:10.1038/ejhg.2017.112.
346. Pillai, N.R.; AlDhaheri, N.S.; Ghosh, R.; Lim, J.; Streff, H.; Nayak, A.; Graham, B.H.; Hanchard, N.A.; Elsea, S.H.; Scaglia, F. Biallelic Variants in COX4I1 Associated with a Novel Phenotype Resembling Leigh Syndrome with Developmental Regression, Intellectual Disability, and Seizures. *Am J Med Genet* **2019**, *179*, 2138–2143, doi:10.1002/ajmg.a.61288.
347. Shteyer, E.; Saada, A.; Shaag, A.; Al-Hijawi, F.A.; Kidess, R.; Revel-Vilk, S.; Elpeleg, O. Exocrine Pancreatic Insufficiency, Dyserythropoietic Anemia, and Calvarial Hyperostosis Are Caused by a Mutation in the COX4I2 Gene. *The American Journal of Human Genetics* **2009**, *84*, 412–417, doi:10.1016/j.ajhg.2009.02.006.
348. Baertling, F.; Al-Murshedi, F.; Sánchez-Caballero, L.; Al-Senaidi, K.; Joshi, N.P.; Venselaar, H.; van den Brand, M.A.; Nijtmans, L.G.; Rodenburg, R.J. Mutation in Mitochondrial Complex IV Subunit COX5A Causes Pulmonary Arterial Hypertension, Lactic Acidemia, and Failure to Thrive: BAERTLING et AL. *Human Mutation* **2017**, *38*, 692–703, doi:10.1002/humu.23210.
349. Tamiya, G.; Makino, S.; Hayashi, M.; Abe, A.; Numakura, C.; Ueki, M.; Tanaka, A.; Ito, C.; Toshimori, K.; Ogawa, N.; et al. A Mutation of COX6A1 Causes a Recessive Axonal or Mixed Form of Charcot-Marie-Tooth Disease. *The American Journal of Human Genetics* **2014**, *95*, 294–300, doi:10.1016/j.ajhg.2014.07.013.
350. Inoue, M.; Uchino, S.; Iida, A.; Noguchi, S.; Hayashi, S.; Takahashi, T.; Fujii, K.; Komaki, H.; Takeshita, E.; Nonaka, I.; et al. COX6A2 Variants Cause a Muscle-specific Cytochrome c Oxidase Deficiency. *Ann Neurol* **2019**, *86*, 193–202, doi:10.1002/ana.25517.
351. Vondrackova, A.; Vesela, K.; Hansikova, H.; Zajicova Docekalova, D.; Rozsypalova, E.; Zeman, J.; Tesarova, M. High-Resolution Melting Analysis of 15 Genes in 60 Patients with Cytochrome-c Oxidase Deficiency. *J Hum Genet* **2012**, *57*, 442–448, doi:10.1038/jhg.2012.49.

352. Indrieri, A.; van Rahden, V.A.; Tiranti, V.; Morleo, M.; Iaconis, D.; Tammaro, R.; D'Amato, I.; Conte, I.; Maystadt, I.; Demuth, S.; et al. Mutations in COX7B Cause Microphthalmia with Linear Skin Lesions, an Unconventional Mitochondrial Disease. *The American Journal of Human Genetics* **2012**, *91*, 942–949, doi:10.1016/j.ajhg.2012.09.016.
353. Hallmann, K.; Kudin, A.P.; Zsurka, G.; Kornblum, C.; Reimann, J.; Stüve, B.; Waltz, S.; Hattingen, E.; Thiele, H.; Nürnberg, P.; et al. Loss of the Smallest Subunit of Cytochrome c Oxidase, COX8A, Causes Leigh-like Syndrome and Epilepsy. *Brain* **2016**, *139*, 338–345, doi:10.1093/brain/awv357.
354. Echaniz-Laguna, A.; Ghezzi, D.; Chassagne, M.; Mayencon, M.; Padet, S.; Melchionda, L.; Rouvet, I.; Lannes, B.; Bozon, D.; Latour, P.; et al. SURF1 Deficiency Causes Demyelinating Charcot-Marie-Tooth Disease. *Neurology* **2013**, *81*, 1523–1530, doi:10.1212/WNL.0b013e3182a4a518.
355. Valnot, I. A Mutation in the Human Heme A:Farnesyltransferase Gene (COX10) Causes Cytochrome c Oxidase Deficiency. *Human Molecular Genetics* **2000**, *9*, 1245–1249, doi:10.1093/hmg/9.8.1245.
356. Antonicka, H. Mutations in COX10 Result in a Defect in Mitochondrial Heme A Biosynthesis and Account for Multiple, Early-Onset Clinical Phenotypes Associated with Isolated COX Deficiency. *Human Molecular Genetics* **2003**, *12*, 2693–2702, doi:10.1093/hmg/ddg284.
357. Coenen, M.J.H.; van den Heuvel, L.P.; Ugalde, C.; ten Brinke, M.; Nijtmans, L.G.J.; Trijbels, F.J.M.; Beblo, S.; Maier, E.M.; Munttau, A.C.; Smeitink, J.A.M. Cytochromec Oxidase Biogenesis in a Patient with a Mutation in COX10 Gene. *Ann Neurol*. **2004**, *56*, 560–564, doi:10.1002/ana.20229.
358. Antonicka, H.; Mattman, A.; Carlson, C.G.; Glerum, D.M.; Hoffbuhr, K.C.; Leary, S.C.; Kennaway, N.G.; Shoubbridge, E.A. Mutations in COX15 Produce a Defect in the Mitochondrial Heme Biosynthetic Pathway, Causing Early-Onset Fatal Hypertrophic Cardiomyopathy. *The American Journal of Human Genetics* **2003**, *72*, 101–114, doi:10.1086/345489.
359. Alfadhel, M.; Lillquist, Y.P.; Waters, P.J.; Sinclair, G.; Struys, E.; McFadden, D.; Henderson, G.; Hyams, L.; Shoffner, J.; Vallance, H.D. Infantile Cardioencephalopathy Due to a COX15 Gene Defect: Report and Review. *Am. J. Med. Genet.* **2011**, *155*, 840–844, doi:10.1002/ajmg.a.33881.
360. Bugiani, M. Novel Mutations in COX15 in a Long Surviving Leigh Syndrome Patient with Cytochrome c Oxidase Deficiency. *Journal of Medical Genetics* **2005**, *42*, e28–e28, doi:10.1136/jmg.2004.029926.
361. Fernández-Vizarra, E.; Tiranti, V.; Zeviani, M. Assembly of the Oxidative Phosphorylation System in Humans: What We Have Learned by Studying Its Defects. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **2009**, *1793*, 200–211, doi:10.1016/j.bbamcr.2008.05.028.
362. Leary, S.C.; Antonicka, H.; Sasarman, F.; Weraarpachai, W.; Cobine, P.A.; Pan, M.; Brown, G.K.; Brown, R.; Majewski, J.; Ha, K.C.H.; et al. Novel Mutations in SCO1 as a Cause of Fatal Infantile Encephalopathy and Lactic Acidosis. *Human Mutation* **2013**, *34*, 1366–1370, doi:10.1002/humu.22385.
363. Brix, N.; Jensen, J.M.; Pedersen, I.S.; Ernst, A.; Frost, S.; Bogaard, P.; Petersen, M.B.; Bender, L. Mitochondrial Disease Caused by a Novel Homozygous Mutation (Gly106del) in the SCO1 Gene. *Neonatology* **2019**, *116*, 290–294, doi:10.1159/000499488.
364. Papadopoulou, L.C.; Sue, C.M.; Davidson, M.M.; Tanji, K.; Nishino, I.; Sadlock, J.E.; Krishna, S.; Walker, W.; Selby, J.; Glerum, D.M.; et al. Fatal Infantile Cardioencephalomyopathy with COX Deficiency and Mutations in SCO2, a COX Assembly Gene. *Nat Genet* **1999**, *23*, 333–337, doi:10.1038/15513.
365. Jaksch, M. Mutations in SCO2 Are Associated with a Distinct Form of Hypertrophic Cardiomyopathy and Cytochrome c Oxidase Deficiency. *Human Molecular Genetics* **2000**, *9*, 795–801, doi:10.1093/hmg/9.5.795.
366. Sue, C.M.; Karadimas, C.; Checcarelli, N.; Tanji, K.; Papadopoulou, L.C.; Pallotti, F.; Guo, F.L.; Shanske, S.; Hirano, M.; De Vivo, D.C.; et al. Differential Features of Patients with Mutations in Two COX Assembly Genes, SURF-1 and SCO2. *Ann Neurol* **2000**, *47*, 589–595.
367. Jaksch, M.; Horvath, R.; Horn, N.; Auer, D.P.; Macmillan, C.; Peters, J.; Gerbitz, K.-D.; Kraegeloh-Mann, I.; Munttau, A.; Karcagi, V.; et al. Homozygosity (E140K) in SCO2 Causes Delayed Infantile Onset of Cardiomyopathy and Neuropathy. *Neurology* **2001**, *57*, 1440–1446, doi:10.1212/WNL.57.8.1440.
368. Pronicki, M.; Kowalski, P.; Piekutowska-Abramczuk, D.; Taybert, J.; Karkucinska-Wieckowska, A.; Szymanska-Debinska, T.; Karczmarewicz, E.; Pajdowska, M.; Migdal, M.; Milewska-Bobula, B.; et al. A Homozygous Mutation in the SCO2 Gene Causes a Spinal Muscular Atrophy like Presentation with Stridor and Respiratory Insufficiency. *European Journal of Paediatric Neurology* **2010**, *14*, 253–260, doi:10.1016/j.ejpn.2009.09.008.
369. Pronicka, E.; Piekutowska-Abramczuk, D.; Szymańska-Dębińska, T.; Bielecka, L.; Kowalski, P.; Łuczak, S.; Karkucińska-Wieckowska, A.; Migdał, M.; Kubalska, J.; Zimowski, J.; et al. The Natural History of SCO2 Deficiency in 36 Polish Children Confirmed the Genotype–Phenotype Correlation. *Mitochondrion* **2013**, *13*, 810–816, doi:10.1016/j.mito.2013.05.007.
370. Rebelo, A.P.; Saade, D.; Pereira, C.V.; Farooq, A.; Huff, T.C.; Abreu, L.; Moraes, C.T.; Mnatsakanova, D.; Mathews, K.; Yang, H.; et al. SCO2 Mutations Cause Early-Onset Axonal Charcot-Marie-Tooth Disease Associated with Cellular Copper Deficiency. *Brain* **2018**, *141*, 662–672, doi:10.1093/brain/awx369.
371. Barcia, G.; Assouline, Z.; Pennisi, A.; Gitiaux, C.; Schiff, M.; Boddaert, N.; Munnich, A.; Bonnefont, J.-P.; Rötig, A. Cytochrome c Oxidase Deficiency Caused by Biallelic SCO2 Mutations in Two Sibs with Cerebellar Ataxia and Progressive Peripheral Axonal Neuropathy. *Molecular Genetics and Metabolism Reports* **2019**, *21*, 100528, doi:10.1016/j.ymgmr.2019.100528.
372. Ghosh, A.; Trivedi, P.P.; Timbalia, S.A.; Griffin, A.T.; Rahn, J.J.; Chan, S.S.L.; Gohil, V.M. Copper Supplementation Restores Cytochrome c Oxidase Assembly Defect in a Mitochondrial Disease Model of COA6 Deficiency. *Human Molecular Genetics* **2014**, *23*, 3596–3606, doi:10.1093/hmg/ddu069.

373. Baertling, F.; A.M. van den Brand, M.; Hertecant, J.L.; Al-Shamsi, A.; P. van den Heuvel, L.; Distelmaier, F.; Mayatepek, E.; Smeitink, J.A.; Nijtmans, L.G.J.; Rodenburg, R.J.T. Mutations in COA6 Cause Cytochrome c Oxidase Deficiency and Neonatal Hypertrophic Cardiomyopathy. *Human Mutation* **2015**, *36*, 34–38, doi:10.1002/humu.22715.
374. Salvador-Severo, K.; Gómez-Caudillo, L.; Quezada, H.; García-Trejo, J. de J.; Cárdenas-Conejo, A.; Vázquez-Memije, M.E.; Minauro-Sanmiguel, F. Mitochondrial Proteomic Profile of Complex IV Deficiency Fibroblasts: Rearrangement of Oxidative Phosphorylation Complex/Supercomplex and Other Metabolic Pathways. *Boletín Médico del Hospital Infantil de México* **2017**, *74*, 175–180, doi:10.1016/j.bmhmx.2017.05.001.
375. Weraarpachai, W.; Antonicka, H.; Sasarman, F.; Seeger, J.; Schrank, B.; Kolesar, J.E.; Lochmüller, H.; Chevrette, M.; Kaufman, B.A.; Horvath, R.; et al. Mutation in TACO1, Encoding a Translational Activator of COX I, Results in Cytochrome c Oxidase Deficiency and Late-Onset Leigh Syndrome. *Nat Genet* **2009**, *41*, 833–837, doi:10.1038/ng.390.
376. Oktay, Y.; Güngör, S.; Zeltner, L.; Wiethoff, S.; Schöls, L.; Sonmezler, E.; Yilmaz, E.; Munro, B.; Bender, B.; Kernstock, C.; et al. Confirmation of TACO1 as a Leigh Syndrome Disease Gene in Two Additional Families. *JND* **2020**, *7*, 301–308, doi:10.3233/JND-200510.
377. Xu, F.; Morin, C.; Mitchell, G.; Ackerley, C.; Robinson, B.H. The Role of the LRPPRC (Leucine-Rich Pentatricopeptide Repeat Cassette) Gene in Cytochrome Oxidase Assembly: Mutation Causes Lowered Levels of COX (Cytochrome c Oxidase) I and COX III MRNA. *Biochemical Journal* **2004**, *382*, 331–336, doi:10.1042/BJ20040469.
378. Antonicka, H.; Shoubridge, E.A. Mitochondrial RNA Granules Are Centers for Posttranscriptional RNA Processing and Ribosome Biogenesis. *Cell Reports* **2015**, *10*, 920–932, doi:10.1016/j.celrep.2015.01.030.
379. Ghezzi, D.; Saada, A.; D'Adamo, P.; Fernandez-Vizarra, E.; Gasparini, P.; Tiranti, V.; Elpeleg, O.; Zeviani, M. FASTKD2 Non-sense Mutation in an Infantile Mitochondrial Encephalomyopathy Associated with Cytochrome C Oxidase Deficiency. *The American Journal of Human Genetics* **2008**, *83*, 415–423, doi:10.1016/j.ajhg.2008.08.009.
380. Wei, X.; Du, M.; Li, D.; Wen, S.; Xie, J.; Li, Y.; Chen, A.; Zhang, K.; Xu, P.; Jia, M.; et al. Mutations in FASTKD2 Are Associated with Mitochondrial Disease with Multi-OXPHOS Deficiency. *Human Mutation* **2020**, *41*, 961–972, doi:10.1002/humu.23985.
381. Nobrega, M.P.; Nobrega, F.G.; Tzagoloff, A. COX10 Codes for a Protein Homologous to the ORF1 Product of Paracoccus Denitrificans and Is Required for the Synthesis of Yeast Cytochrome Oxidase. *J Biol Chem* **1990**, *265*, 14220–14226.
382. Valnot, I.; Osmond, S.; Gigarel, N.; Mehaye, B.; Amiel, J.; Cormier-Daire, V.; Munnich, A.; Bonnefont, J.-P.; Rustin, P.; Rötig, A. Mutations of the SCO1 Gene in Mitochondrial Cytochrome c Oxidase Deficiency with Neonatal-Onset Hepatic Failure and Encephalopathy. *The American Journal of Human Genetics* **2000**, *67*, 1104–1109, doi:10.1016/S0002-9297(07)62940-1.
383. Glerum, D.M.; Muroff, I.; Jin, C.; Tzagoloff, A. COX15 Codes for a Mitochondrial Protein Essential for the Assembly of Yeast Cytochrome Oxidase. *Journal of Biological Chemistry* **1997**, *272*, 19088–19094, doi:10.1074/jbc.272.30.19088.
384. Oquendo, C.E. Functional and Genetic Studies Demonstrate That Mutation in the COX15 Gene Can Cause Leigh Syndrome. *Journal of Medical Genetics* **2004**, *41*, 540–544, doi:10.1136/jmg.2003.017426.
385. Smith, D.; Gray, J.; Mitchell, L.; Antholine, W.E.; Hosler, J.P. Assembly of Cytochrome-c Oxidase in the Absence of Assembly Protein Surf1p Leads to Loss of the Active Site Heme. *Journal of Biological Chemistry* **2005**, *280*, 17652–17656, doi:10.1074/jbc.C500061200.
386. Huigslot, M.; Nijtmans, L.G.; Szklarczyk, R.; Baars, M.J.H.; van den Brand, M.A.M.; HendriksFranssen, M.G.M.; van den Heuvel, L.P.; Smeitink, J.A.M.; Huynen, M.A.; Rodenburg, R.J.T. A Mutation in C2orf64 Causes Impaired Cytochrome c Oxidase Assembly and Mitochondrial Cardiomyopathy. *The American Journal of Human Genetics* **2011**, *88*, 488–493, doi:10.1016/j.ajhg.2011.03.002.
387. Stroud, D.A.; Maher, M.J.; Lindau, C.; Vögtle, F.-N.; Frazier, A.E.; Surgenor, E.; Mountford, H.; Singh, A.P.; Bonas, M.; Oeljeklaus, S.; et al. COA6 Is a Mitochondrial Complex IV Assembly Factor Critical for Biogenesis of MtDNA-Encoded COX2. *Hum. Mol. Genet.* **2015**, *24*, 5404–5415, doi:10.1093/hmg/ddv265.
388. Leary, S.C. Human SCO1 and SCO2 Have Independent, Cooperative Functions in Copper Delivery to Cytochrome c Oxidase. *Human Molecular Genetics* **2004**, *13*, 1839–1848, doi:10.1093/hmg/ddh197.
389. Stiburek, L.; Vesela, K.; Hansikova, H.; Hulkova, H.; Zeman, J. Loss of Function of Sco1 and Its Interaction with Cytochrome c Oxidase. *American Journal of Physiology-Cell Physiology* **2009**, *296*, C1218–C1226, doi:10.1152/ajpcell.00564.2008.
390. Hiser, L.; Di Valentin, M.; Hamer, A.G.; Hosler, J.P. Cox11p Is Required for Stable Formation of the CuBand Magnesium Centers of Cytochrome c Oxidase. *Journal of Biological Chemistry* **2000**, *275*, 619–623, doi:10.1074/jbc.275.1.619.
391. Cerqua, C.; Morbidoni, V.; Desbats, M.A.; Doimo, M.; Frasson, C.; Sacconi, S.; Baldoin, M.C.; Sartori, G.; Basso, G.; Salvati, L.; et al. COX16 Is Required for Assembly of Cytochrome c Oxidase in Human Cells and Is Involved in Copper Delivery to COX2. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2018**, *1859*, 244–252, doi:10.1016/j.bbabo.2018.01.004.
392. Carlson, C.G.; Barrientos, A.; Tzagoloff, A.; Glerum, D.M. COX16 Encodes a Novel Protein Required for the Assembly of Cytochrome Oxidase in Saccharomyces Cerevisiae. *Journal of Biological Chemistry* **2003**, *278*, 3770–3775, doi:10.1074/jbc.M209893200.
393. Glerum, D.M.; Shtanko, A.; Tzagoloff, A. Characterization of , a Yeast Gene Involved in Copper Metabolism and Assembly of Cytochrome Oxidase. *Journal of Biological Chemistry* **1996**, *271*, 14504–14509, doi:10.1074/jbc.271.24.14504.
394. Bode, M.; Woellhaf, M.W.; Bohnert, M.; Laan, M. van der; Sommer, F.; Jung, M.; Zimmermann, R.; Schroda, M.; Herrmann, J.M. Redox-Regulated Dynamic Interplay between Cox19 and the Copper-Binding Protein Cox11 in the Intermembrane Space of Mitochondria Facilitates Biogenesis of Cytochrome c Oxidase. *MBoC* **2015**, *26*, 2385–2401, doi:10.1091/mbc.E14-11-1526.

395. Nobrega, M.P.; Bandeira, S.C.B.; Beers, J.; Tzagoloff, A. Characterization of COX19, a Widely Distributed Gene Required for Expression of Mitochondrial Cytochrome Oxidase. *Journal of Biological Chemistry* **2002**, *277*, 40206–40211, doi:10.1074/jbc.M207348200.
396. Naess, K.; Bruhn, H.; Stranneheim, H.; Freyer, C.; Wibom, R.; Mourier, A.; Engvall, M.; Nennesmo, I.; Lesko, N.; Wredenberg, A.; et al. Clinical Presentation, Genetic Etiology, and Coenzyme Q10 Levels in 55 Children with Combined Enzyme Deficiencies of the Mitochondrial Respiratory Chain. *The Journal of Pediatrics* **2021**, *228*, 240–251.e2, doi:10.1016/j.jpeds.2020.08.025.
397. Szklarczyk, R.; Wanschers, B.F.J.; Nijtmans, L.G.; Rodenburg, R.J.; Zschocke, J.; Dikow, N.; van den Brand, M.A.M.; Hendriks-Franssen, M.G.M.; Gilissen, C.; Veltman, J.A.; et al. A Mutation in the FAM36A Gene, the Human Ortholog of COX20, Impairs Cytochrome c Oxidase Assembly and Is Associated with Ataxia and Muscle Hypotonia. *Human Molecular Genetics* **2013**, *22*, 656–667, doi:10.1093/hmg/ddt473.
398. Doss, S.; Lohmann, K.; Seibler, P.; Arns, B.; Klopstock, T.; Zühlke, C.; Freimann, K.; Winkler, S.; Lohnau, T.; Drungowski, M.; et al. Recessive Dystonia-Ataxia Syndrome in a Turkish Family Caused by a COX20 (FAM36A) Mutation. *J Neurol* **2014**, *261*, 207–212, doi:10.1007/s00415-013-7177-7.
399. Clemente, P.; Peralta, S.; Cruz-Bermudez, A.; Echevarría, L.; Fontanesi, F.; Barrientos, A.; Fernandez-Moreno, M.A.; Garesse, R. HCOA3 Stabilizes Cytochrome c Oxidase 1 (COX1) and Promotes Cytochrome c Oxidase Assembly in Human Mitochondria*. *Journal of Biological Chemistry* **2013**, *288*, 8321–8331, doi:10.1074/jbc.M112.422220.
400. Mick, D.U.; Vukotic, M.; Piechura, H.; Meyer, H.E.; Warscheid, B.; Deckers, M.; Rehling, P. Coa3 and Cox14 Are Essential for Negative Feedback Regulation of COX1 Translation in Mitochondria. *Journal of Cell Biology* **2010**, *191*, 141–154, doi:10.1083/jcb.201007026.
401. Mick, D.U.; Dennerlein, S.; Wiese, H.; Reinhold, R.; Pacheu-Grau, D.; Lorenzi, I.; Sasarman, F.; Weraarpachai, W.; Shoubridge, E.A.; Warscheid, B.; et al. MITRAC Links Mitochondrial Protein Translocation to Respiratory-Chain Assembly and Translational Regulation. *Cell* **2012**, *151*, 1528–1541, doi:10.1016/j.cell.2012.11.053.
402. Ostergaard, E.; Weraarpachai, W.; Ravn, K.; Born, A.P.; Jønson, L.; Duno, M.; Wibrand, F.; Shoubridge, E.A.; Vissing, J. Mutations in COA3 Cause Isolated Complex IV Deficiency Associated with Neuropathy, Exercise Intolerance, Obesity, and Short Stature. *J Med Genet* **2015**, *52*, 203–207, doi:10.1136/jmedgenet-2014-102914.
403. Higuchi, Y.; Okunushi, R.; Hara, T.; Hashiguchi, A.; Yuan, J.; Yoshimura, A.; Murayama, K.; Ohtake, A.; Ando, M.; Hiramatsu, Y.; et al. Mutations in COA7 Cause Spinocerebellar Ataxia with Axonal Neuropathy. *Brain* **2018**, *141*, 1622–1636, doi:10.1093/brain/awy104.
404. Weraarpachai, W.; Sasarman, F.; Nishimura, T.; Antonicka, H.; Auré, K.; Rötig, A.; Lombès, A.; Shoubridge, E.A. Mutations in C12orf62, a Factor That Couples COX I Synthesis with Cytochrome c Oxidase Assembly, Cause Fatal Neonatal Lactic Acidosis. *The American Journal of Human Genetics* **2012**, *90*, 142–151, doi:10.1016/j.ajhg.2011.11.027.
405. Bourens, M.; Barrientos, A. A CMC1 -knockout Reveals Translation-independent Control of Human Mitochondrial Complex IV Biogenesis. *EMBO Rep* **2017**, *18*, 477–494, doi:10.15252/embr.201643103.
406. Hell, K.; Tzagoloff, A.; Neupert, W.; Stuart, R.A. Identification of Cox20p, a Novel Protein Involved in the Maturation and Assembly of Cytochrome Oxidase Subunit 2. *Journal of Biological Chemistry* **2000**, *275*, 4571–4578, doi:10.1074/jbc.275.7.4571.
407. Church, C.; Chapon, C.; Poyton, R.O. Cloning and Characterization of PET100, a Gene Required for the Assembly of Yeast Cytochrome c Oxidase. *Journal of Biological Chemistry* **1996**, *271*, 18499–18507, doi:10.1074/jbc.271.31.18499.
408. Lim, S.C.; Smith, K.R.; Stroud, D.A.; Compton, A.G.; Tucker, E.J.; Dasvarma, A.; Gandolfo, L.C.; Marum, J.E.; McKenzie, M.; Peters, H.L.; et al. A Founder Mutation in PET100 Causes Isolated Complex IV Deficiency in Lebanese Individuals with Leigh Syndrome. *The American Journal of Human Genetics* **2014**, *94*, 209–222, doi:10.1016/j.ajhg.2013.12.015.
409. Oláhová, M.; Haack, T.B.; Alston, C.L.; Houghton, J.A.; He, L.; Morris, A.A.; Brown, G.K.; McFarland, R.; Chrzanowska-Lightowlers, Z.M.; Lightowlers, R.N.; et al. A Truncating PET100 Variant Causing Fatal Infantile Lactic Acidosis and Isolated Cytochrome c Oxidase Deficiency. *Eur J Hum Genet* **2015**, *23*, 935–939, doi:10.1038/ejhg.2014.214.
410. Taylor, N.G.; Swenson, S.; Harris, N.J.; Germany, E.M.; Fox, J.L.; Khalimonchuk, O. The Assembly Factor Pet117 Couples Heme a Synthase Activity to Cytochrome Oxidase Assembly. *Journal of Biological Chemistry* **2017**, *292*, 1815–1825, doi:10.1074/jbc.M116.766980.
411. McEwen, J.E.; Hong, K.H.; Park, S.; Preciado, G.T. Sequence and Chromosomal Localization of Two PET Genes Required for Cytochrome c Oxidase Assembly in *Saccharomyces Cerevisiae*. *Curr Genet* **1993**, *23*, 9–14, doi:10.1007/BF00336742.
412. Renkema, G.H.; Visser, G.; Baertling, F.; Wintjes, L.T.; Wolters, V.M.; van Montfrans, J.; de Kort, G.A.P.; Nikkels, P.G.J.; van Hasselt, P.M.; van der Crabben, S.N.; et al. Mutated PET117 Causes Complex IV Deficiency and Is Associated with Neurodevelopmental Regression and Medulla Oblongata Lesions. *Hum Genet* **2017**, *136*, 759–769, doi:10.1007/s00439-017-1794-7.
413. Signes, A.; Cerutti, R.; Dickson, A.S.; Benincá, C.; Hinchey, E.C.; Ghezzi, D.; Carrozzo, R.; Bertini, E.; Murphy, M.P.; Nathan, J.A.; et al. APOPT1/COA8 Assists COX Assembly and Is Oppositely Regulated by UPS and ROS. *EMBO Mol Med* **2019**, *11*, doi:10.15252/emmm.201809582.
414. Melchionda, L.; Damseh, N.S.; Abu Libdeh, B.Y.; Nasca, A.; Elpeleg, O.; Zanolini, A.; Ghezzi, D. A Novel Mutation in TTC19 Associated with Isolated Complex III Deficiency, Cerebellar Hypoplasia, and Bilateral Basal Ganglia Lesions. *Front. Genet.* **2014**, *5*, doi:10.3389/fgene.2014.00397.
415. Hedberg-Oldfors, C.; Darin, N.; Thomsen, C.; Lindberg, C.; Oldfors, A. COX Deficiency and Leukoencephalopathy Due to a Novel Homozygous APOPT1/COA8 Mutation. *Neurol Genet* **2020**, *6*, e464, doi:10.1212/NXG.0000000000000464.

416. Souza, R.L.; Green-Willms, N.S.; Fox, T.D.; Tzagoloff, A.; Nobrega, F.G. Cloning and Characterization of COX18, ASaccharomyces Cerevisiae PET Gene Required for the Assembly of Cytochrome Oxidase. *Journal of Biological Chemistry* **2000**, *275*, 14898–14902, doi:10.1074/jbc.275.20.14898.
417. Bourens, M.; Barrientos, A. Human Mitochondrial Cytochrome c Oxidase Assembly Factor COX18 Acts Transiently as a Membrane Insertase within the Subunit 2 Maturation Module. *Journal of Biological Chemistry* **2017**, *292*, 7774–7783, doi:10.1074/jbc.M117.778514.
418. Sacconi, S.; Trevisson, E.; Pistollato, F.; Baldoin, M.C.; Rezzonico, R.; Bourget, I.; Desnuelle, C.; Tenconi, R.; Basso, G.; DiMauro, S.; et al. HCOX18 and HCOX19: Two Human Genes Involved in Cytochrome c Oxidase Assembly. *Biochemical and Biophysical Research Communications* **2005**, *337*, 832–839, doi:10.1016/j.bbrc.2005.09.127.
419. Legati, A.; Reyes, A.; Nasca, A.; Invernizzi, F.; Lamantea, E.; Tiranti, V.; Garavaglia, B.; Lamperti, C.; Ardisson, A.; Moroni, I.; et al. New Genes and Pathomechanisms in Mitochondrial Disorders Unraveled by NGS Technologies. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2016**, *1857*, 1326–1335, doi:10.1016/j.bbabo.2016.02.022.
420. Jonckheere, A.I.; Smeitink, J.A.M.; Rodenburg, R.J.T. Mitochondrial ATP Synthase: Architecture, Function and Pathology. *J Inherit Metab Dis* **2012**, *35*, 211–225, doi:10.1007/s10545-011-9382-9.
421. Abrahams, J.P.; Leslie, A.G.W.; Lutter, R.; Walker, J.E. Structure at 2.8 Å Resolution of F1-ATPase from Bovine Heart Mitochondria. *Nature* **1994**, *370*, 621–628, doi:10.1038/370621a0.
422. He, J.; Ford, H.C.; Carroll, J.; Douglas, C.; Gonzales, E.; Ding, S.; Fearnley, I.M.; Walker, J.E. Assembly of the Membrane Domain of ATP Synthase in Human Mitochondria. *Proc Natl Acad Sci USA* **2018**, *115*, 2988–2993, doi:10.1073/pnas.1722086115.
423. Spikes, T.E.; Montgomery, M.G.; Walker, J.E. Structure of the Dimeric ATP Synthase from Bovine Mitochondria. *Proc Natl Acad Sci USA* **2020**, *117*, 23519–23526, doi:10.1073/pnas.2013998117.
424. Pinke, G.; Zhou, L.; Sazanov, L.A. Cryo-EM Structure of the Entire Mammalian F-Type ATP Synthase. *Nat Struct Mol Biol* **2020**, *27*, 1077–1085, doi:10.1038/s41594-020-0503-8.
425. Nijtmans, L.G.J.; Klement, P.; Houštěk, J.; van den Bogert, C. Assembly of Mitochondrial ATP Synthase in Cultured Human Cells: Implications for Mitochondrial Diseases. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1995**, *1272*, 190–198, doi:10.1016/0925-4439(95)00087-9.
426. Wittig, I.; Meyer, B.; Heide, H.; Steger, M.; Bleier, L.; Wumaier, Z.; Karas, M.; Schägger, H. Assembly and Oligomerization of Human ATP Synthase Lacking Mitochondrial Subunits a and A6L. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2010**, *1797*, 1004–1011, doi:10.1016/j.bbabo.2010.02.021.
427. Wang, Z.G.; Ackerman, S.H. The Assembly Factor Atp11p Binds to the Beta-Subunit of the Mitochondrial F(1)-ATPase. *J Biol Chem* **2000**, *275*, 5767–5772, doi:10.1074/jbc.275.8.5767.
428. Wang, Z.-G. The Alpha -Subunit of the Mitochondrial F1 ATPase Interacts Directly with the Assembly Factor Atp12p. *The EMBO Journal* **2000**, *19*, 1486–1493, doi:10.1093/emboj/19.7.1486.
429. Shoffner, J.M.; Fernhoff, P.M.; Krawiecki, N.S.; Caplan, D.B.; Holt, P.J.; Koontz, D.A.; Takei, Y.; Newman, N.J.; Ortiz, R.G.; Polak, M.; et al. Subacute Necrotizing Encephalopathy: Oxidative Phosphorylation Defects and the ATPase 6 Point Mutation. *Neurology* **1992**, *42*, 2168–2168, doi:10.1212/WNL.42.11.2168.
430. White, S.L.; Shanske, S.; Biros, I.; Warwick, L.; Dahl, H.M.; Thorburn, D.R.; Di Mauro, S. Two Cases of Prenatal Analysis for the Pathogenic T to G Substitution at Nucleotide 8993 in Mitochondrial DNA. *Prenat Diagn* **1999**, *19*, 1165–1168.
431. Burrage, L.C.; Tang, S.; Wang, J.; Donti, T.R.; Walkiewicz, M.; Luchak, J.M.; Chen, L.-C.; Schmitt, E.S.; Niu, Z.; Erana, R.; et al. Mitochondrial Myopathy, Lactic Acidosis, and Sideroblastic Anemia (MLASA) plus Associated with a Novel de Novo Mutation (m.8969G>A) in the Mitochondrial Encoded ATP6 Gene. *Molecular Genetics and Metabolism* **2014**, *113*, 207–212, doi:10.1016/j.ymgme.2014.06.004.
432. Rantamäki, M.T.; Soini, H.K.; Finnilä, S.M.; Majamaa, K.; Udd, B. Adult-Onset Ataxia and Polyneuropathy Caused by Mitochondrial 8993T→C Mutation. *Ann Neurol* **2005**, *58*, 337–340, doi:10.1002/ana.20555.
433. Craig, K.; Elliott, H.R.; Keers, S.M.; Lambert, C.; Pyle, A.; Graves, T.D.; Woodward, C.; Sweeney, M.G.; Davis, M.B.; Hanna, M.G.; et al. Episodic Ataxia and Hemiplegia Caused by the 8993T→C Mitochondrial DNA Mutation. *Journal of Medical Genetics* **2007**, *44*, 797–799, doi:10.1136/jmg.2007.052902.
434. Pfeffer, G.; Blakely, E.L.; Alston, C.L.; Hassani, A.; Boggild, M.; Horvath, R.; Samuels, D.C.; Taylor, R.W.; Chinnery, P.F. Adult-Onset Spinocerebellar Ataxia Syndromes Due to MTATP6 Mutations. *J Neurol Neurosurg Psychiatry* **2012**, *83*, 883–886, doi:10.1136/jnnp-2012-302568.
435. De Meirleir, L.; Seneca, S.; Lissens, W.; Schoentjes, E.; Desprechins, B. Bilateral Striatal Necrosis with a Novel Point Mutation in the Mitochondrial ATPase 6 Gene. *Pediatric Neurology* **1995**, *13*, 242–246, doi:10.1016/0887-8994(95)00184-H.
436. Thyagarajan, D.; Shanske, S.; Vazquez -Memije, M.; Devivo, D.; Dimauro, S. A Novel Mitochondrial ATPase 6 Point Mutation in Familial Bilateral Striatal Necrosis. *Ann Neurol* **1995**, *38*, 468–472, doi:10.1002/ana.410380321.
437. Brum, M.; Semedo, C.; Guerreiro, R.; Pinto Marques, J. Motor Neuron Syndrome as a New Phenotypic Manifestation of Mutation 9185T>C in Gene MTATP6. *Case Reports in Neurological Medicine* **2014**, *2014*, 1–4, doi:10.1155/2014/701761.
438. Takahashi, S.; Makita, Y.; Oki, J.; Miyamoto, A.; Yanagawa, J.; Naito, E.; Goto, Y.; Okuno, A. De Novo MtDNA Nt 8993 (T→G) Mutation Resulting in Leigh Syndrome. *The American Journal of Human Genetics* **1998**, *62*, 717–719, doi:10.1086/301751.
439. Vilarinho, L.; Barbot, C.; Carrozzo, R.; Calado, E.; Tessa, A.; Dionisi-Vici, C.; Guimarães, A.; Santorelli, F.M. Clinical and Molecular Findings in Four New Patients Harboring the MtDNA 8993T>C Mutation. *J Inherit Metab Dis* **2001**, *24*, 883–884, doi:10.1023/A:1013908728445.

440. Carrozzo, R.; Tessa, A.; Vazquez-Memije, M.E.; Piemonte, F.; Patrono, C.; Malandrini, A.; Dionisi-Vici, C.; Vilarinho, L.; Villanova, M.; Schagger, H.; et al. The T9176G MtDNA Mutation Severely Affects ATP Production and Results in Leigh Syndrome. *Neurology* **2001**, *56*, 687–690, doi:10.1212/WNL.56.5.687.
441. Lopez-Gallardo, E.; Solano, A.; Herrero-Martin, M.D.; Martinez-Romero, I.; Castano-Perez, M.D.; Andreu, A.L.; Herrera, A.; Lopez-Perez, M.J.; Ruiz-Pesini, E.; Montoya, J. NARP Syndrome in a Patient Harboring an Insertion in the MT-ATP6 Gene That Results in a Truncated Protein. *Journal of Medical Genetics* **2008**, *46*, 64–67, doi:10.1136/jmg.2008.060616.
442. D'Aurelio, M.; Vives-Bauza, C.; Davidson, M.M.; Manfredi, G. Mitochondrial DNA Background Modifies the Bioenergetics of NARP/MILS ATP6 Mutant Cells. *Human Molecular Genetics* **2010**, *19*, 374–386, doi:10.1093/hmg/ddp503.
443. de Coo, I.F.; Smeets, H.J.; Gabreëls, F.J.; Arts, N.; van Oost, B.A. Isolated Case of Mental Retardation and Ataxia Due to a de Novo Mitochondrial T8993G Mutation. *Am J Hum Genet* **1996**, *58*, 636–638.
444. Ware, S.M.; El-Hassan, N.; Kahler, S.G.; Zhang, Q.; Y-W; Miller, E.; Wong, B.; Spicer, R.L.; Craigen, W.J.; Kozel, B.A.; et al. Infantile Cardiomyopathy Caused by a Mutation in the Overlapping Region of Mitochondrial ATPase 6 and 8 Genes. *Journal of Medical Genetics* **2009**, *46*, 308–314, doi:10.1136/jmg.2008.063149.
445. Galimberti, C.A.; Diegoli, M.; Sartori, I.; Uggetti, C.; Brega, A.; Tartara, A.; Arbustini, E. Brain Pseudoatrophy and Mental Regression on Valproate and a Mitochondrial DNA Mutation. *Neurology* **2006**, *67*, 1715–1717, doi:10.1212/01.wnl.0000242882.58086.9a.
446. Jonckheere, A.I.; Hogeveen, M.; Nijtmans, L.; van den Brand, M.; Janssen, A.; Diepstra, H.; van den Brandt, F.; van den Heuvel, B.; Hol, F.; Hofste, T.; et al. A Novel Mitochondrial ATP8 Gene Mutation in a Patient with Apical Hypertrophic Cardiomyopathy and Neuropathy. *Case Reports* **2009**, *2009*, bcr0720080504–bcr0720080504, doi:10.1136/bcr.07.2008.0504.
447. Jonckheere, A.I.; Renkema, G.H.; Bras, M.; van den Heuvel, L.P.; Hoischen, A.; Gilissen, C.; Nabuurs, S.B.; Huynen, M.A.; de Vries, M.C.; Smeitink, J.A.M.; et al. A Complex V ATP5A1 Defect Causes Fatal Neonatal Mitochondrial Encephalopathy. *Brain* **2013**, *136*, 1544–1554, doi:10.1093/brain/awt086.
448. Lieber, D.S.; Calvo, S.E.; Shanahan, K.; Slate, N.G.; Liu, S.; Hershman, S.G.; Gold, N.B.; Chapman, B.A.; Thorburn, D.R.; Berry, G.T.; et al. Targeted Exome Sequencing of Suspected Mitochondrial Disorders. *Neurology* **2013**, *80*, 1762–1770, doi:10.1212/WNL.0b013e3182918c40.
449. Oláhová, M.; Yoon, W.H.; Thompson, K.; Jangam, S.; Fernandez, L.; Davidson, J.M.; Kyle, J.E.; Grove, M.E.; Fisk, D.G.; Kohler, J.N.; et al. Biallelic Mutations in ATP5F1D, Which Encodes a Subunit of ATP Synthase, Cause a Metabolic Disorder. *The American Journal of Human Genetics* **2018**, *102*, 494–504, doi:10.1016/j.ajhg.2018.01.020.
450. Mayr, J.A.; Havlickova, V.; Zimmermann, F.; Magler, I.; Kaplanova, V.; Jesina, P.; Pecinova, A.; Nuskova, H.; Koch, J.; Sperl, W.; et al. Mitochondrial ATP Synthase Deficiency Due to a Mutation in the ATP5E Gene for the F1 Subunit. *Human Molecular Genetics* **2010**, *19*, 3430–3439, doi:10.1093/hmg/ddq254.
451. Schaubberger, E.M.; Ewart, S.L.; Arshad, S.H.; Huebner, M.; Karmaus, W.; Holloway, J.W.; Friderici, K.H.; Ziegler, J.T.; Zhang, H.; Rose-Zerilli, M.J.; et al. Identification of ATPAF1 as a Novel Candidate Gene for Asthma in Children. *Journal of Allergy and Clinical Immunology* **2011**, *128*, 753–760.e11, doi:10.1016/j.jaci.2011.04.058.
452. De Meirleir, L. Respiratory Chain Complex V Deficiency Due to a Mutation in the Assembly Gene ATP12. *Journal of Medical Genetics* **2004**, *41*, 120–124, doi:10.1136/jmg.2003.012047.
453. de Vries, D.D.; van Engelen, B.G.M.; Gabreëls, F.J.M.; Ruitenbeek, W.; van Oost, B.A. A Second Missense Mutation in the Mitochondrial ATPase 6 Gene in Leigh's Syndrome: MtDNA Mutation in Leigh's Syndrome. *Ann Neurol* **1993**, *34*, 410–412, doi:10.1002/ana.410340319.
454. White, S.L.; Shanske, S.; McGill, J.J.; Mountain, H.; Geraghty, M.T.; DiMauro, S.; Dahl, H.-H.M.; Thorburn, D.R. Mitochondrial DNA Mutations at Nucleotide 8993 Show a Lack of Tissue- or Age-Related Variation. *J Inherit Metab Dis* **1999**, *22*, 899–914, doi:10.1023/A:1005639407166.
455. Houštěk, J.; Pícková, A.; Vojtíšková, A.; Mráček, T.; Pecina, P.; Ješina, P. Mitochondrial Diseases and Genetic Defects of ATP Synthase. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2006**, *1757*, 1400–1405, doi:10.1016/j.bbabo.2006.04.006.
456. Schagger, H. Respiratory Chain Supercomplexes of Mitochondria and Bacteria. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2002**, *1555*, 154–159, doi:10.1016/S0005-2728(02)00271-2.
457. Wittig, I.; Schagger, H. Structural Organization of Mitochondrial ATP Synthase. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2008**, *1777*, 592–598, doi:10.1016/j.bbabo.2008.04.027.
458. Mourier, A.; Matic, S.; Ruzzenente, B.; Larsson, N.-G.; Milenkovic, D. The Respiratory Chain Supercomplex Organization Is Independent of COX7a2l Isoforms. *Cell Metabolism* **2014**, *20*, 1069–1075, doi:10.1016/j.cmet.2014.11.005.
459. Acín-Pérez, R.; Fernández-Silva, P.; Peleato, M.L.; Pérez-Martos, A.; Enriquez, J.A. Respiratory Active Mitochondrial Supercomplexes. *Molecular Cell* **2008**, *32*, 529–539, doi:10.1016/j.molcel.2008.10.021.
460. Gu, J.; Wu, M.; Guo, R.; Yan, K.; Lei, J.; Gao, N.; Yang, M. The Architecture of the Mammalian Respirasome. *Nature* **2016**, *537*, 639–643, doi:10.1038/nature19359.
461. Wu, M.; Gu, J.; Guo, R.; Huang, Y.; Yang, M. Structure of Mammalian Respiratory Supercomplex I 1 III 2 IV 1. *Cell* **2016**, *167*, 1598–1609.e10, doi:10.1016/j.cell.2016.11.012.
462. Letts, J.A.; Fiedorczuk, K.; Sazanov, L.A. The Architecture of Respiratory Supercomplexes. *Nature* **2016**, *537*, 644–648, doi:10.1038/nature19774.
463. Sousa, J.S.; Mills, D.J.; Vonck, J.; Kühlbrandt, W. Functional Asymmetry and Electron Flow in the Bovine Respirasome. *eLife* **2016**, *5*, e21290, doi:10.7554/eLife.21290.

464. Lapuente-Brun, E.; Moreno-Loshuertos, R.; Acín-Pérez, R.; Latorre-Pellicer, A.; Colás, C.; Balsa, E.; Perales-Clemente, E.; Quirós, P.M.; Calvo, E.; Rodríguez-Hernández, M.A.; et al. Supercomplex Assembly Determines Electron Flux in the Mitochondrial Electron Transport Chain. *Science* **2013**, *340*, 1567–1570, doi:10.1126/science.1230381.
465. Genova, M.L.; Lenaz, G. Functional Role of Mitochondrial Respiratory Supercomplexes. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2014**, *1837*, 427–443, doi:10.1016/j.bbabi.2013.11.002.
466. Lenaz, G.; Tioli, G.; Falasca, A.I.; Genova, M.L. Complex I Function in Mitochondrial Supercomplexes. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2016**, *1857*, 991–1000, doi:10.1016/j.bbabi.2016.01.013.
467. Calvo, E.; Cogliati, S.; Hernansanz-Agustín, P.; Loureiro-López, M.; Guarás, A.; Casuso, R.A.; García-Marqués, F.; Acín-Pérez, R.; Martí-Mateos, Y.; Silla-Castro, Jc.; et al. Functional Role of Respiratory Supercomplexes in Mice: SCAF1 Relevance and Segmentation of the Q_{pool}. *Sci. Adv.* **2020**, *6*, eaba7509, doi:10.1126/sciadv.aba7509.
468. Loeffen, J.L.C.M.; Smeitink, J.A.M.; Trijbels, J.M.F.; Janssen, A.J.M.; Triepels, R.H.; Sengers, R.C.A.; van den Heuvel, L.P. Isolated complex I deficiency in children: Clinical, biochemical and genetic aspects. *Human Mutation* **2000**, *15.2*, 123–134.
469. Protasoni, M.; Pérez-Pérez, R.; Lobo-Jarne, T.; Harbour, M.E.; Ding, S.; Peñas, A.; Diaz, F.; Moraes, C.T.; Fearnley, I.M.; Zeviani, M.; et al. Respiratory Supercomplexes Act as a Platform for Complex III -mediated Maturation of Human Mitochondrial Complexes I and IV. *EMBO J* **2020**, *39*, doi:10.15252/embj.2019102817
470. Pernas, L.; Scorrano, L. Mito-Morphosis: Mitochondrial Fusion, Fission, and Cristae Remodeling as Key Mediators of Cellular Function. *Annu. Rev. Physiol.* **2016**, *78*, 505–531, doi:10.1146/annurev-physiol-021115-105011.
471. Cogliati, S.; Enriquez, J.A.; Scorrano, L. Mitochondrial Cristae: Where Beauty Meets Functionality. *Trends in Biochemical Sciences* **2016**, *41*, 261–273, doi:10.1016/j.tibs.2016.01.001.
472. Silva Ramos, E.; Larsson, N.-G.; Mourier, A. Bioenergetic Roles of Mitochondrial Fusion. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **2016**, *1857*, 1277–1283, doi:10.1016/j.bbabi.2016.04.002.
473. Letts, J.A.; Sazanov, L.A. Clarifying the Supercomplex: The Higher-Order Organization of the Mitochondrial Electron Transport Chain. *Nat Struct Mol Biol* **2017**, *24*, 800–808, doi:10.1038/nsmb.3460.